1 **Unlocking prophage potential: *In silico* and experimental analysis of a**
2 **novel *Mycobacterium fortuitum* LysinB containing a peptidoglycan-binding**
3 **domain**
4

Ritam Das[1,2], Kanika Nadar[3], Ritu Arora[3] and Urmi Bajpai[3]*

[1]Faculty of Biological Sciences, Friedrich Schiller University, Jena-07737, Germany

[2]Department of Life Science, Acharya Narendra Dev College, University of Delhi, Govindpuri,

Kalkaji-110019, New Delhi, India

[3]*Department of Biomedical Science, Acharya Narendra Dev College, University of Delhi,

Govindpuri, Kalkaji-110019, New Delhi, India, urmibajpai@andc.du.ac.in

Address for correspondence:
Urmi Bajpai, PhD (Professor)
Department of Biomedical Science,
Acharya Narendra Dev College (University of Delhi),
Govindpuri, Kalkaji, New-Delhi-110019, India.
Email id: urmibajpai@andc.du.ac.in

Abstract

Endolysins are highly evolved bacteriophage-encoded lytic enzymes produced to damage the bacterial cell wall for phage progeny release. They offer promising potential as highly specific lytic proteins with a low chance of bacterial resistance. The diversity in lysin sequences and domain organization can be staggering. *In silico* analysis of bacteriophage and prophage genomes can help identify endolysins exhibiting unique features and high antibacterial activity, hence feeding the pipeline of narrow-spectrum protein antibiotics. Mycobacteriophage lysis cassettes mostly have two lytic enzymes, LysinA and LysinB. The enzyme LysinA targets peptidoglycan in the cell wall and possesses a modular architecture. LysinB typically contains a single domain and acts upon the mycolyl ester linkages in mycolyl-arabinogalactan-peptidoglycan (Payne *et al.*, 2010). This study aimed to find novel LysinBs against *Mycobacterium fortuitum*. After a detailed *in silico* characterization of lysis cassettes from three *M. fortuitum* prophages, we chose to work on a LysinB (hereafter described as LysinB_MF) found in an incomplete prophage (phiE1336, 9.4 kb in strain E1336).

LysinB_MF showed low sequence similarity with any other endolysins in the database and formed a separate clade on phylogenetic analysis. LysinB_MF's structure, extracted from the AlphaFold Protein Structure Database, demonstrated a modular architecture with two structurally distinct domains: a peptidoglycan-binding domain (PGBD) at the N-terminal and the characteristic alpha/beta hydrolase domain connected via a linker peptide. We found the alpha/beta hydrolase domain, which is the enzyme-active domain (EAD), contains the conserved Ser-Asp-His catalytic triad with a tunnel-like topology and forms intermolecular hydrogen bonds. The PGBD shows structural similarity to the cell-wall binding domain of an amidase from *Clostridium acetobutylicum,* hinting at its acquisition due to domain mobility. Our *in silico* electrostatic potential analysis suggested that PGBD might be essential to the enzyme activity. This was experimentally validated by generating a truncated version of the enzyme, which demonstrated about six-fold decreased activity compared to its native form. The antimycobacterial activity of this enzyme was also compromised in its absence. Based on our analysis, PGBD emerged as an integral constituent of enzymes with diverse functional properties and is predicted to be a conserved cross-kingdom. Overall, this study highlights the importance of mining mycobacterial prophages as a novel endolysin source. It also provides unique insights into the diverse architecture of mycobacteriophage-encoded endolysins and the importance of functional domains for their catalytic activities.

Keywords: Prophage; Endolysins; LysinB; Structure; Domains; Esterase activity

The modular LysinB_MF

1. Introduction

Mycobacterium infections are among the most debilitating, if not fatal (Gordon & Parish, 2018; Das *et al.*, 2021). *Mycobacterium tuberculosis*, the causative agent of Tuberculosis, alone was the cause of 1.3 million deaths in the year 2022, the second-highest infectious killer after COVID-19 (Bagcchi, 2023). Infections due to nontuberculous *Mycobacterium* (NTM) are also becoming an emerging issue of concern, and their prevalence is increasing worldwide. While *Mycobacterium fortuitum*-associated pulmonary diseases are the most prevalent (Khosravi *et al.*, 2018), the bacterium also causes extra-pulmonary infections in humans, such as cutaneous and bone lesions, which are difficult to treat, chronic, occult, resistant, and recurrent (Sethi *et al.*, 2014; Fraga *et al.*, 2022).

Mycobacteriophages are viruses that specifically infect *Mycobacterium spp*. Most mycobacteriophages code for two endolysins: LysinA and LysinB, which target the bacterial cell wall and cause lysis of their host cells at the end of their life cycle (Payne *et al.*, 2010; Payne & Hatfull, 2012). LysinA is a peptidoglycan hydrolase, and LysinB is a mycolylarabinogalactan esterase, which hydrolyzes the mycolyl ester linkages in mycolyl-arabinogalactan-peptidoglycan

layer present in mycobacteria. Since the exogenous application of endolysins has been

demonstrated to show an antibacterial effect, lysins have been explored as an alternative/

combination therapy with antibiotics in several studies (Fraga *et al.*, 2019; Abouhmad *et al.*, 2020;

Hurst-Hess *et al.*, 2023; Singh *et al.*, 2023). While the reports on structural and functional

characterization of mycobacteriophage-encoded lysins (Payne *et al.*, 2010; Eniyan *et al.*, 2020;

Gigante *et al.*, 2021) have provided insights into their structural diversity, these lysins are relatively

less reported when compared to phage lysins against Gram-negative and Gram-positive bacteria.

Besides the isolated mycobacteriophages, the genomes of nontuberculous *Mycobacterium* (NTM)

that are enriched with prophages (Abad *et al.*, 2023), could also serve as a promising source for

diverse antibacterial proteins (Schmitz *et al.*, 2011; Shah *et al.*, 2023). However, these prophages

are largely unannotated. In this study, we report a novel LysinB encoded by one such unique

prophage in a strain of *M. fortuitum,* with three characteristic features: i) unlike the usual single-

domain architecture of LysinB enzymes, LysinB_MF has a structurally distinct peptidoglycan

binding domain (PGBD), ii) its enzyme active domain shows structural similarity to a hydrolase

(amidase) of *Clostridium acetobutylicum,* and iii) PGBD has an important role in enzyme activity.

As a first step towards understanding the molecular properties, we have characterized the lytic

cassette of three *M. fortuitum* prophages, and carried out a detailed examination of LysinB_MF,

showing the highest similarity with D29 LysinB, which we used as a bait sequence for searching

LysinB sequences in *M.fortuitum* prophages. We have carried out an extensive sequence and

structural analysis of lysis cassettes and LysinB_MF and hypothesized and biochemically

characterized the importance of the PGBD in its activity.

2. Results and Discussion

2.1 *Mycobacterium* prophages as a source of novel lysin sequences

D29 and Ms6 LysB enzymes are the extensively studied mycolylarabinogalactan esterases with pre-clinical characterization (Payne & Hatfull, 2012; Gigante *et al.*, 2017; Fraga *et al.*, 2019; Hurst-Hess *et al.*, 2023). However, to facilitate the growing interest in mycobacteriophage lysins to treat drug-resistant *Mycobacterium spp.,* more lysins need to be discovered and investigated to understand their biological features and to develop them as novel therapeutics. Non-tubercular mycobacteria (NTM) are reported to harbour a large proportion of prophages (Abad *et al.*, 2023). We used D29 LysB (O64205) as a bait sequence and performed BLAST to find similar but not identical LysinB sequences against all the *M. fortuitum* genomes in the database. The workflow for the *in silico* identification of LysinB enzymes is represented in Figure 1A. The top three hits for LysinB were found to be encoded by prophages of *M. fortuitum* strains (Table 1). In this study, we have carried out the structural and functional characterization of a unique LysinB enzyme (mentioned as LysinB_MF) encoded by a prophage, called phiE1336, present in the sequenced strain E1336 of *Mycobacterium fortuitum* (tax id: 1766). A detailed characterization of this prophage has been done in the later sections.

We found the prophage phiE1336 genome to be incomplete (9.4 kb), however, its lysis cassette was intact. It constituted LysinA, three small hypothetical proteins, LysinB and Holin (Figure 1B). LysinB_MF, with an overall identity of 49% with D29 LysinB, is a peptidoglycan-binding domain-containing protein (WP_064866887.1). The second hit, with an overall identity of 48%, was a hypothetical protein (HP) (WP_278289780.1) from *M. fortuitum* strain DVD-1301 (NZ_JARUNL010000131.1). Reannotation of the genome and PHASTER prediction indicated the protein to be a LysinB, encoded by an intact prophage (hereby called phiDVD-1301). This

prophage genome size was 39.6 kb with a GC content of 62.86%, similar to what is observed for

other mycobacteriophages. Nucleotide BLAST analysis of the genome indicated phiDVD-1301 to

have approximately 80% sequence identity with cluster P phages Tortellini (KX648391),

Phayonce (KR080195) and Purky (MN096355). Annotation demonstrated the presence of a lytic

cassette (Figure 1B) with LysinA (385 amino acids) as the largest protein-coding gene. LysinB

(326 amino acids) from phiDVD-1301 showed 48% identity with D29 LysB, however, domain

predictions for this enzyme remained unsuccessful. The third hit (WP_269975983.1) showing 36%

sequence identity with D29 LysB was a hypothetical protein (HP), encoded by an *M. fortuitum*

strain MF GZ001 (NZ_CP107719.1) prophage. This prophage (hereby called phiGZ001) has a

size of 38.9 kb with distinct attL (CACACTGTTCTTC) and attR (CACACTGTTCTTC) sites

recognized by serine integrases (Suzuki *et al.*, 2020). Nucleotide BLAST analysis of the genome

indicated phiGZ001 to have approximately 86% sequence identity with cluster F1 phages XFactor

(KT281795), Tootsieroll (MZ747519) and Blexus (MN586012). Sequence reannotation and

domain analysis of the identified HP (WP_269975983.1) predicted an alpha/beta hydrolase

domain. In phiGZ001, we did not find a holin gene in the lysis cassette (Figure 1B). We speculate

this prophage either lyses the host independent of holin (Eniyan *et al.*, 2020) or the lytic cassette

is incomplete.

Figure 1: Analysis of prophages in *M. fortuitum* strains. A) The workflow of prophage detection and characterization of the lytic regions. **B)** Lytic cassette of prophages phiE1336, phiDVD-1301 and phiGZ001.

Table 1: Information on *M. fortuitum* prophages identified and characterized in the study

S. no.	Name	Size (kb)	Completeness	GC%	Cluster similarity	*M. fortuitum* lysogen
1	phiE1336	9.4	Incomplete	62	-	NZ_LZSJ01000108.1
2	phiDVD-1301	39.6	Intact	62.8	P	NZ_JARUNL010000131.1
3	phiGZ001	38.9	Intact	63.4	F1	NZ_CP107719.1

2.2 The incomplete prophage in the *M. fortuitum* strain E1336

PHAST analysis predicted the presence of an incomplete genome of prophage phiE1336. Its size

is only 9.4 kb (Figure 2A) but the GC content (62%) is comparable to the host (Ho *et al.*, 2012).

Mycobacteriophages are grouped into clusters based on the similarities and differences in their

nucleotide sequences (Hatfull, 2022). Currently, 32 clusters of mycobacteriophages have been

defined, and the unclassified phages are denoted as singletons (https://phagesdb.org/). We

performed nucleotide BLAST analysis to identify the possible cluster of phiE1336 and to predict

phages identical to it. None of the whole genome sequences in the database showed more than

10% query coverage, a trend similar to singletons. However, considering the short genomic

fragment of phiE1336, it is difficult to state conclusively whether it belongs among the singleton

phages or to a cluster.

2.3 The lytic cassette in prophage phiE1336

We extracted the 9.4 kb fragment and performed annotation using RAST, which predicted it to

have 13 ORFs (Figure 2A). Among the predicted ORFs, six could be assigned putative functions,

which included lytic and structural proteins (Table S1). The lytic cassette begins with LysinA (44.3

kDa), a peptidoglycan hydrolase consisting of an amidase domain spanning from 171–317 and an

overlapping peptidoglycan recognition protein (PGRP) region from 183–317 amino acids (Figure

2B). Co-occurrence of domains is common in phage lysins (Payne & Hatfull, 2012; Hatfull, 2018).

The amidase domain targets the N-acetylmuramoyl-L-alanine linkages, which connect the sugar

backbone to the peptide crosslink (Payne & Hatfull, 2012). BLAST analysis showed phiE1336

LysinA to share 71.9% identity with LysinA (gp24) from F1 sub-cluster mycobacteriophage

Koella (MH316564) and 70.5% identity with LysinA (gp69) from O cluster phage Smooch

(MN428052). Domain construction of lysins from these phages is similar and constitutes the active

site of the amidase with an overlaying peptidoglycan recognition protein (PGRP) region.

The two small HPs between LysinA and LysinB of phiE1336 (Figure 1B) are of size 10.6 kDa and

3.9 kDa and do not possess a recognizable domain. Neither of the HPs shows any significant

similarity to other sequences in the database. This arrangement of the lytic cassette, with HPs

between Lysin A and LysinB, appears unique for phiE1336 among the other prophages

characterized in this study (Figure 1B). The other reported mycobacterium prophages, such as

phiMAV1 and MabR, constitute lytic regions with LysinA and LysinB present together (Payne *et*

al., 2010; Cote *et al.*, 2022).

Upstream to the two HPs in the cassette are the genes encoding for mycolylarabinogalactan

esterase (LysinB_MF), another small HP (3.1 kDa), and holin (Figure 2B). LysinB_MF is a

multidomain enzyme with a molecular weight of 35.3 kDa. While we could not precisely predict

the function of the 3.1 kDa protein, BLAST analysis showed similar proteins to have an ATP

binding activity. The holin gene encodes for a protein of size 7.4 kDa, which shares 61.1% identity

with the holin from cluster I2 mycobacteriophage Che9c. Comparable to other phage-encoded

holin, this, too, consists of two transmembrane helices, which could destabilize the membrane

proton motive force of the host in a time-dependent manner (Pollenz *et al.*, 2022).

Figure 2: Characterization of phiE1366 and the encoded LysinB_MF. A) phiE1336 has a
genome size of 9.4 kb with a GC content of 62%. The genome consists of 13 ORFs, which encode for a few
structural proteins and proteins of the lytic cassette **B)** Domain architecture of LysinB_MF from

phiE1336 demonstrates distinct PGBD and EAD. **C)** Phylogenetic analysis of LysinB_MF to understand
its evolutionary relationship. Representative sequences of LysinBs from all sub-clusters and singletons were
taken for analysis. Phylogeny analysis displayed LysinB_MF forming its separate clade (in red arrow), which
branched from the neighbouring Cluster V LysinB at some point. Green arrows show the clades disparate with
the evolutionary relationships of mycobacteriophage clusters.
200

2.4 LysinB_MF

Although the information on mycobacteriophage-encoded mycolylarabinogalactan esterase

(LysinB) has been discussed earlier in several studies (Payne *et al.*, 2010; Korany *et al.*, 2019;

Abouhmad *et al.*, 2020), reports on the characterization of a multidomain LysinB is limited

(Gigante *et al.*, 2021). We found LysinB_MF to be a modular enzyme composed of two

structurally distinct domains connected by a linker peptide (Figure 2B). The enzyme (319 amino

acids) is comprised of two functionally defined regions: PGBD and the alpha/beta hydrolase

domain (Figure S1). The PGBD is present at the N-terminal region of LysinB_MF and spans from

amino acids 39–70, and a linker of 14 amino acids connects it to the catalytic enzyme-active

domain (EAD) (Figure 2B). The EAD, like most mycolylarabinogalactan esterases, is comprised

of an alpha/beta hydrolase domain that spans right after the linker until the C-terminal end,

covering a substantial part of the enzyme. Through BLAST analysis, we found LysinB from a V

cluster mycobacteriophage Cosmo (KP027195) to be the closest predicted homologue of

LysinB_MF. The sequences showed a high query coverage of 98% and 56.43% identity (e-value

3e-116). Besides this, we could not find duplicate or identical proteins to LysinB_MF in the

database. Therefore, to examine the evolutionary relationship of LysinB_MF, we performed the

phylogenetic analysis using LysinB sequences from singletons and mycobacteriophages belonging

to all the sub-clusters (Figure 2C). There were 187 sequences in the repository file prepared with

at least one sequence and a maximum of three. The multi-FASTA file was subjected to sequence

alignment and then processed using the iTOL to generate a rooted phylogenetic tree based on the

maximum likelihood method.

Interestingly, we found a separate clade for LysinB_MF, as observed for LysinBs from singleton

phages (Figure 2C). The LysinB_MF sequence appeared to have branched from the cluster V

endolysins and had a branch length of 0.79. A PGBD in these LysinBs remains unannotated or

lacking, although conserved residues in the N-terminal region were observed in the multiple

sequence file. It appears Cluster V LysinBs and LysinB_MF may have had a common ancestor at

some point; however, they diverged into forming their clade, and LysinB_MF retained the PGBD

function.

Though the phylogenetic analysis gave us some clue about the evolutionary relationship of

LysinB_MF, what was interesting to note was the unconventional grouping of endolysins from

different sub-clusters in the same clade (shown in green arrows, Figure 2C). Phylogenetic analysis

of tape-measure proteins in mycobacteriophage genomes distinctly shows separate clades, called

clusters or even specific sub-clusters (Jacobs-Sera *et al.*, 2012). Phages belonging to the same

cluster are grouped into the same clade. However, while studying LysinB, we observed sequences

belonging to entirely different clusters, if not sub-clusters, forming a common sub-clade and

demonstrating the possibility of a common ancestor at some point in the evolutionary history.

LysinB from clusters A7 and K6; F1, F4 and F5; B6 and B12; P1, P3 and I1, respectively, were

present in the same clade (Figure 2C). In some instances, the evolution of LysinB appeared to be

independent of the evolution of mycobacteriophages. Mycobacteriophage-encoded LysinBs,

therefore, emerge as enzymes with a complex evolutionary pattern, and their lineage requires a

greater analysis.

Figure 3: Structural analysis of LysinB_MF. A) Three-dimensional (3D) model of LysinB_MF demonstrates a multi-domain structure comprising two structurally distinct domains: the PGBD (39-70), a short linker region followed by the EAD, an α/β hydrolase (85-317). **B)** The surface model of LysinB_MF demonstrates a bulky C-terminal end with a distinct active-site groove formed with residues Ser-Asp-His. The groove of the EAD, indicated with a yellow arrow, displays a tunnel-like topology. **C)** LysinB_MF can be superimposed on D29 LysB (in pink) with an overall RMSD of 0.56. Structural alignment of the two endolysins shows similar active-site arrangement, although PGBD in D29 Lysin B is absent. **D)** Active-site conformation in LysinB_MF and D29 LysinB. LysinB_MF shows a narrow but deeper topology (shown with a yellow arrow). D29 LysB displays a larger edge, having a funnel-like conformation (shown with a purple arrow). **E)** Active site residues in LysinB_MF (Ser156-Asp236-His305) and D29 LysB (S82-Asp166-His240).

2.5 LysinB_MF has a complex enzyme active domain (EAD)

To analyze the structural information of LysinB_MF, we attempted to characterize the 3D model

of the enzyme available in the AlphaFold Protein Structure Database (AFDB: A0A1A0NU50).

The structure of this enzyme was available as a peptidoglycan-binding protein derived from the

M. fortuitum strain described earlier. The protein structure showed a high model confidence

(pLDDT > 90) and was taken forward for further analysis (Figure 3).

The EAD of LysinB_MF comprises the α/β sandwich belonging to the α/β hydrolase family and

contains the catalytic triad Ser-Asp-His. This domain spanned from amino acid 85–317, covering

a major portion of the enzyme and demonstrated a highly twisted structure, consisting of a C-

terminal linker region from 246-310. The active site of LysinB_MF displayed four β-strands (β2–

β5), surrounded from all sides by α-helices (α4-α6). A surface model of the enzyme demonstrated

the active-site groove with a tunnel-like topology (Figure 3B). A similar topology is observed in

D29 LysB, which serves as a binding pocket for substrates such as several *p*-NP derivatives,

trehalose monomycolate and mAGP from mycobacterial cell wall (Payne *et al.*, 2010). A DALI

search with high confidence (DALI Z-score 29.7) identified the structural relationship of

LysinB_MF with the tertiary structure of D29 LysB (PDB accession 3HC7). The two esterases

could be superimposed with an overall RMSD of 0.56 (153 C^α atoms) (Figure 3C). Both the

structures displayed similar β-sheet conformation around the active site surrounded by the helices.

Given the high structural similarity observed between LysinB_MF and D29 LysinB, the topology

of D29 LysB's binding pocket, unlike LysinB_MF, showed an edge with a wider circumference

and a funnel-like appearance (Figure 3D). Furthermore, a bird's eye view and analysis of the

conserved active-site residues (Ser82, Asp166 & His240 in D29 LysB and Ser156, Asp236 &

His305 in LysinB_MF) (Figure 4A and 4D) of these enzymes displayed notable differences.

Unlike their counterparts, the residues Asp166 in D29 LysB and His305 in LysinB_MF were part

of alpha-helices in these enzymes, which added to the differences in their groove topology. Also,

unlike D29 LysB, the active-site residue His305 in LysinB_MF was found to form polar contacts

with residues Ser156 and Asp236 (Figure 3E). These bonds are known to provide stability by

acting as electrostatic attractive forces which influence the charge redistribution in the enzyme and

are responsible for the uptake of electrons (pull) from the active site towards the amino acid

residues (Qiu *et al.*, 2018). Also, in the case of serine proteases with similar Ser-Asp-His catalytic

triads, the removal of hydrogen bonds resulted in the partial release of the active-site residues in

molecular dynamic simulations and a decrease in the overall stability of the transition states, which

affects the enzyme's turnover number (k_{cat}) (Zheng *et al.*, 2006).

Figure 4: Conserved residues in LysinB_MF. A) Multiple sequence alignment of LysinB enzymes from
various clusters demonstrate key residues in PGBD and EAD. The EAD displays the conserved active-site
residues Ser-Asp-His and the GXP motif. **B)** Analysis using POLYVIEW-MM identified and mapped conserved
residues in the tertiary structure of LysinB_MF. A scale of 0-9 is used to quantify the conservation index, where
0 indicates the least conserved and nine is the most. Exposed residues in the structure of LysinB_MF are found
to be least conserved among all others, while most of the highly conserved residues are found buried. **C)** Bird's
eye view of PGBD displays conserved amino acids. Ser41, Tyr43 Phe44, Asp45, Val52, Glu54, Gln56, Gly66
and Ile67 are a few highly conserved residues in this domain found both in sequence alignment and in structural

analysis. **D)** Bird's eye view of EAD displays conserved amino acids. The active site Ser156, Asp236 & His305
and Gly191 & Pro193 from the GXP motif were highly conserved in LysinBs.
300

2.6 The PGBD of LysinB_MF is structurally similar to an amidase

In most of the mycolylarabinogalactan esterases reported earlier (Fraga *et al.*, 2019; Abouhmad *et*

al., 2020; Eniyan *et al.*, 2020; Singh *et al.*, 2023), an extended N-terminal region consisting of a

PGBD hasn't been reported. It is a rare feature that has been observed in Ms6 LysinB (Gigante *et*

al., 2021) and our analysis has predicted only seven out of sixty LysinB sequences to contain

PGBD (unpublished data). Region 7–71 in LysinB_MF was interpreted as a PGBD-like

superfamily, with region 39–70 predicted to be the functional PGBD (highlighted in light blue in

Figure 2A). Only a single β-sheet in this region is present in the enzyme's N-terminal end, and the

rest of the structure appeared to be highly twisted, dominated by three α-helices. The annotated

PGBD is comprised of two α helices (α2 and α3), and it tapers to form the linker region connecting

to the EAD. The multiple sequence alignment and structural analysis of LysinB_MF demonstrated

several conserved residues in this region (Figure 4), which are also found in LysinBs without the

N-terminal PGBD. The amino acids with the highest score in the conservation index are mostly

buried in the enzyme and could be involved in the functional properties of the PGBD (Figures 4A-

4C).

Considering the unavailability of structural information about the PGBD in a mycobacteriophage-

encoded LysinB and failure to acquire the same through structural similarity analysis based on the

native structure of LysinB_MF, we separately built the PGBD and analyzed it further. The tertiary

structure of PGBD was prepared using LysinB_MF as a template, and it had a QMEANDisCo

Global score of 0.64 ± 0.11, which ensured high quality (Figure S2A). We then performed

structural similarity prediction using DALI, which predicted 20 possible neighbours, and these

were visually sorted based on their Z-score. Interestingly, the top scoring structure (Z-score from

DALI 8.9) found similar to the PGBD of LysinB_MF was an enzyme with amidase activity (PDB

accession code 4XXT). This zinc-dependent amidase is a PGBD-containing protein from

Clostridium acetobutylicum (ATCC 824). To analyze the structural similarity to PGBD of

LysinB_MF, we downloaded the protein 3D file from the PDB database and carried out structural

alignment using PyMOL. The PGBDs from the amidase and LysinB_MF could be superpositioned

with an overall RMSD as low as 0.46, even though the sequences do not show identical residues

(Figure S2B). Also, amidases (or lysozyme-like domains) possessing a peptidoglycan-binding and

hydrolyzing function are usually the features of LysinA enzymes. Due to the unavailability of the

3D structure of a LysinA in the PDB database, we could not examine the structural similarity of

PGBD specifically to a mycobacteriophage-encoded N-acetylmuramoyl-L-alanine amidase.

However, it is noteworthy that structural similarity of PGBD of LysinB_MF with high confidence

was achieved with an amidase from a bacterial species. We speculate that domain mobility could

be a reason behind the generation of modular proteins, and they play a crucial role in their

functioning (Basu *et al.*, 2009).

2.7 PGBD is essential for the esterase activity of LysinB_MF

To understand the role of PGBD in the activity of LysinB_MF, we first computationally generated

its truncated version (LysinB_MF**Δ**PGBD), lacking the N-terminal domain (Figures 5A and 5B).

Using eF-surf, a comparative analysis of the native and the truncated structures indicated a change

in the electrostatic potential of the active site. The catalytic tunnel appeared more electronegative

in LysinB_MF**Δ**PGBD than its native structure (Figures 5C and 5D). A change in electrostatic

potential could impact the interactions between an enzyme and its substrate and, hence, the

enzyme's activity. The importance of electrostatics in enzyme-mediated catalysis has been

demonstrated for amylases where a change in the charge near the active site residues resulted in

an enzyme with drastically reduced activity (Nielsen *et al.*, 1999). To find out if the deletion of

PGBD also hindered the *in vitro* activity of LysinB_MF, we purified the complete (LysinB_MF)

and the truncated enzyme (LysinB_MF**Δ**PGBD) as recombinant proteins (Figures 5E-5F).

The lipolytic assay of the truncated protein yielded low Ca^{2+} precipitation (Figure 5H). The *in vitro*

esterase assay also revealed the truncated LysinB_MF**Δ**PGBD (Figures 5A, 5B and 5E) to show

about a six-fold decrease in activity when compared with the activity of the whole enzyme (Figure

5G), indicating PGBD's role in enzyme activity. A similar effect has also been observed for an N-

acetylmuramoyl-L-alanine amidase from bacteriophage PSA, where truncation of the cell wall

binding domain led to a several-fold decrease in enzyme activity (Korndorfer *et al.*, 2006). The

stabilizing effect of PGBD in both mycolylarabinogalactan esterase (LysinB) and N-

acetylmuramoyl-L-alanine amidase (LysinA) is significant and requires more investigation.

Figure 5: PGBD is essential for the optimum activity of LysinB_MF. A) Domain architecture and comparison of native LysinB_MF and truncated LysinB_MFΔPGBD (EAD). **B)** The structure of LysinB_MF without the PGBD (LysinB_MFΔPGBD) was built using the native structure as a template. LysinB_MFΔPGBD structure comprises the linker region (in grey) and the EAD (in green). **C-D)** Comparison of the intact LysinB_MF and LysinB_MFΔPGBD demonstrates a change in electrostatic potential in the active site (indicated with a yellow arrow). **E)** SDS-PAGE (12%) analysis of purified LysinB_MF and LysinB_MFΔPGBD. Lane 1: protein molecular mass standards. Lane 2: purified recombinant protein LysinB_MF (36.3kDa). Lane 3: LysinB_MFΔPGBD (26 kDa). **F)** Immunoblot of N-terminal His-tagged LysinB_MF. Lane 1: protein molecular mass standards. Lane 2: N-terminal His-tagged LysinB_MF. **G)** A white zone indicates Ca^{2+}-precipitation by LysinB_MF and LysinB_MFΔPGBD (5 μg each). **H)** Esterase

372 activity of whole LysinB_MF and LysinB, determined by the PNP release assay (n=6). **I)** Spot test
373 for intrinsic activity of LysinB_MF and LysinB_MF**ΔPGBD** against *M. smegmatis* and *M.*
374 *fortuitum*. All the assays were performed using 5 µg of each enzyme preparation.
375

2.8 Antimycobacterial activity of LysinB_MF and LysinB_MFΔPGBD

Given the complex, hydrophobic cell envelope of mycobacterial spp., the ability of an endolysin

to cause cell lysis on external application is highly desired since it obviates the need for an outer

membrane permeabilizer (OMP) or a membrane-destabilizer. This property also makes lysins

more efficacious candidates as antibiotics (enzybiotics) or as an adjunct to antibiotics. To check

whether LysinB_MF possessed intrinsic cell lysis activity, we performed a lysin susceptibility test

on *M. fortuitum* and *M. smegmatis*. A zone of clearance on both the mycobacterium species was

observed, indicating its ability to lyse mycobacterial cells unaided. Considering the significant

decrease in LysinB_MF**ΔPGBD** activity observed in the esterase assay (Figure 5H), we tested its

cell lysis activity against *M. fortuitum* (and *M. smegmatis*) and, unlike whole LysinB_MF, did not

find clearance zone on the bacterial lawn (Figures 5I). It appears that the PGBD not only supports

the enzymatic activity of LysinB_MF but might also have a role in destabilizing the cell wall of

M.fortuitum and *M.smegmatis*. The cross-species activity of the PGBD has not been reported for

a mycobacteriophage lysin. It will be exciting to identify the PGBD-interacting region in the

mycobacterial cell wall.

2.9 Conservation of PGBD across different kingdoms

Through protein truncation and enzymatic assays, we demonstrated the integral role of PGBD in

LysinB_MF activity (Figures 5H- 5I). However, this feature is not constant in all other endolysins.

While in the case of bacteriophage PSA, removing the cell-wall binding domain of endolysin

demonstrated a similar decrease in enzymatic activity (Korndorfer *et al.*, 2006), the lytic activity

of a *Bacillus anthracis* prophage endolysin was enhanced when the C-terminal cell-wall binding

domain was removed (Low *et al.*, 2005). PGBD appears to have diverse functionality; however, a

proper understanding of the conservation and importance of PGBD is limited. To comprehend the

importance of this functional domain in other organisms, we analyzed all the sequences present in

the UniProt database against the Pfam entry for the PGBD (PF01471). There are approximately

122,000 protein sequences consisting of this domain. From our curated analysis of these sequences

based on TaxId from different organisms, we found this domain is conserved across all kingdoms

(Figure 6). Among the prokaryotes, PF01471 was predicted to be the highest in *E. coli* (Figure 6),

where it is present in LD-transpeptidases, which are involved in cell wall adaptation to stress and

for providing outer membrane stability (Aliashkevich & Cava, 2022). In *M. fortuitum*, PF01471

was found chiefly in enzymes with a PG hydrolase activity. In the evolutionary Tree of Life in

higher kingdoms, with *Arabidopsis thaliana* as an example, our curated search predicted 27

UniProt protein matches against PF01471 (Figure 6). Most of the matched proteins in this species

were found to be zinc-dependent endo-peptidases or matrix metalloproteinases, which are involved

in root development and water uptake (Mishra *et al.*, 2021). In higher animals, including humans,

we received 54 UniProt protein matches against PF01471, which, notably, was higher than most

prokaryotes analyzed in this study (Figure 6). In humans, a diverse group of essential enzymes

possess a PGBD, including matrix metalloproteinases (also observed in *A. thaliana*), enzymes

involved in the degradation of extracellular matrix and development of human bones such as

Stromelysin-2 (Bord *et al.*, 1998) and collagenases, which are implicated in the turnover of

connective tissue matrix constituents (Knauper *et al.*, 1996). The cross-kingdom conservation of

the short PGBD observed here would require a deeper analysis and is beyond the scope of our

study.

Figure 6: Cross-kingdom conservation of a PGBD. The UniProt database was screened against the Pfam ID of PGBD (PF01471), which resulted in approximately 122,000 protein hits with this domain. The curated database of hits was then individually screened against TaxIDs from various organisms across all kingdoms.

3. Conclusion

We have carried out an *in silico* analysis of three prophages in *M. fortuitum*, a medically important

NTM. The analysis includes extraction, annotation and sequence characterization of prophages,

followed by the structural analysis of the lysis cassettes. In this study, a detailed analysis of one of

the three prophages, 'phiE1336' and its encoded LysinB (LysinB_MF) was done. Besides the

catalytic alpha/beta domain and a C-terminal linker region, LysinB_MF has an extended N-

terminal, containing a peptidoglycan-binding domain. A conservation analysis predicted this

domain in diverse enzymes encoded by several organisms. Though LysinB_MF is encoded by an

incomplete prophage, which should be incapable of forming phage progeny on induction, its *in*

vitro esterase activity is greater than the reported LysinB enzymes. It will be worthwhile to

investigate whether this highly active, prophage-derived LysinB enzyme has a role in the host

M.fortuitum life cycle. Based on our *in silico* analysis, we predicted PGBD's role in the enzyme's

activity and validated it experimentally. In addition, we also observed a loss of the intrinsic

antibacterial activity in the truncated enzyme.

In summary, the diversity in domain organization of lysins in the prophage-encoded lysis cassettes

and the efficient *in vitro* and antimycobacterial activity of LysinB_MF highlights the importance

of prophages as a source of novel efficacious endolysins, which could be developed as promising

anti-mycobacterial protein antibiotics.

4. Materials and Methods

4.1 *In silico* identification of LysinB_MF and characterization

Lysin B from D29 (GenBank: AAC18452.1) was blasted (Altschul *et al.*, 1990) against all the

genomes available in the NCBI database for *M. fortuitum* (tax id: 1766), possibly harbouring

prophages. The *M. fortuitum* genome with the top lysin hits (LysinB_MF) was subjected to

PHAST (Zhou *et al.*, 2011) and to annotate and fix frameshifts in these genomes, the RAST server

was used. The CGview tool (Grant & Stothard, 2008) was used to build the genome map of the

first hit, phiE1336. Cluster-wise sequences of LysinB were downloaded from the

Actinobacteriophage Database (https://phagesdb.org/), and domains of those lysins were predicted

using InterProScan 5 (Quevillon *et al.*, 2005). ClustalW (Thompson *et al.*, 1994) and ESPript

(https://espript.ibcp.fr) (Robert & Gouet, 2014) were used to generate alignment files which were

then used to create phylogenetic trees using Interactive Tree Of Life (iTOL) v5 (Letunic & Bork,

2021). The tertiary structure of LysinB_MF was downloaded from the AlphaFold Protein Structure

Database (Varadi *et al.*, 2023) and visualized using PyMOL (https://pymol.org/2/). The

POLYVIEW-MM tool was used to predict conserved residues in the tertiary structure of

LysinB_MF (Porollo & Meller, 2010). The DALI server was used to find structures similar to the

LysinB_MF (Holm, 2022) and eF-surf was used to predict the electrostatic potential of the enzyme

(Nielsen *et al.*, 1999). Conservation analysis was done by analyzing the Pfam entry of PGBD

(PF01471) against all the proteins available in the UniProt database (https://www.uniprot.org/).

TaxIDs were used to detect the presence of this domain across several species using the InterPro

classification of protein family website (https://www.ebi.ac.uk/interpro/).

4.2 Cloning, overexpression and purification of recombinant LysinB_MF and

LysinB_MFΔPGBD

The LysinB_MF gene was synthesized and cloned in the pET28a vector by Biomatik Corporation,

Kitchener, Ontario. To clone the truncated version of LysinB_MF lacking the PGBD

(LysinB_MFΔPGBD), internal primers were designed to PCR amplify the EAD and the amplicon

was ligated to the pET28a vector. The recombinant constructs pET28a-LysinB_MF and pET28a-

LysinB_MF**Δ**PGBD were overexpressed in BL21(DE3) and C41(DE3) strains of *E. coli,*

respectively. For soluble expression, induction with 0.1 mM IPTG and overnight incubation at

16ºC was done. The induced culture pellet was collected by centrifugation at 8,000 rpm for ten

minutes at 4ºC and sonicated at 90% amplitude with 30 seconds each on or off pulse for ten cycles

(Eniyan *et al.*, 2020; Das & Bajpai, 2023). The cell lysate was centrifuged at 13,000 rpm for 45

minutes at 4ºC, and the supernatant was considered the soluble fraction.

Recombinant LysinB_MF and LysinB_MF**Δ**PGBD were purified by Ni-NTA affinity

chromatography. Briefly, the soluble fraction containing the His$_6$-tagged protein was kept for

binding with the Ni-NTA matrix for 2 hours at 4 ºC, and the bound resin was packed in a column.

After packing and discarding the flow-through, the column was gradient-washed with buffer

containing 20-40 mM imidazole and the bound protein was eluted at 200-250 mM imidazole

concentrations. The eluted fractions were analyzed using 12% SDS-PAGE (Laemmli, 1970) and

the purified proteins were dialyzed with a buffer containing 50 mM Tris (pH 7.9) and 150 mM

NaCl. The concentration of each protein was estimated by Bradford assay (Bradford, 1976). For

immunoblotting, the protein(s) in the gel were transferred to a PVDF membrane using a Mini-

PROTEAN Tetra Cell (Bio-Rad, United States). After blocking the membrane with 3% BSA at

4°C overnight, the membrane was incubated with mouse anti-His6 antiserum (Santacruz Biotech,

USA) for 4 hours at a dilution of 1:2000. The membrane was washed once with PBS and twice

with PBS-Tween, followed by incubation with IgG HRP-conjugated antibody at a dilution of

1:10000 for 45 minutes at 4 °C. The blot was developed by using 3, 3'-Diaminobenzidine (DAB-

H$_2$O$_2$) as the substrate.

4.3 Assay for the esterase activity of LysinB

The micro-titre plate-based p-Nitrophenol butyrate (pNPB) esterase assay (Fraga *et al.*, 2019) was

performed to estimate the activity of LysinB_MF and LysinB_MFΔPGBD. The reaction

conditions for both the recombinant proteins were kept constant: 5 µg of enzyme 10 mM pNPB in

25 mM Tris buffer (pH 7.2) in a reaction volume of 200 µl at 37ºC for 30 minutes. Absorbance at

410 nm was measured to determine the release of p-nitrophenol (PNP). In the control reaction,

enzyme buffer was added in place of the enzymes. The experiment was done twice in three sets,

and the absorbance value (A_{410}) was plotted as mean ± SD.

4.4 Lipolytic activity of LysinB

510   The enzyme activity was also observed by $CaCl_2$ precipitation. Five µg each of LysinB_MF and

511   LysinB_MFΔPGBD was spot-tested on Middlebrook 7H10 plates containing 1mM $CaCl_2$ and 1%

512   (v/v) Tween-20. Precipitation, confirming lipolytic activity, was observed with 10 min incubation

513   at room temperature.

4.5 Spot Test of LysinB

For the antimycobacterial activity, the spot test on bacterial lawns was performed. Briefly,

Mycobacterium smegmatis strain mc²155 and *M. fortuitum* (NIHJ 1615) were grown for 48 hours

at 37°C using Middlebrook 7H9 broth (HiMedia, India), supplemented with 0.4% glycerol.

Bacterial lawn in 7H10 Middlebrook agar plates containing 0.05% Tween 80 plates were prepared

by spreading 0.5 mL of *M. smegmatis* and *M. fortuitum* using 7H10 soft agar. Five µg each of

LysinB_MF and LysinB_MFΔPGBD were added to the prepared lawn as spots and the plates were

incubated at 37°C.

Acknowledgement

We thank the Science and Engineering Research Board (SERB), Department of Science and Technology (DST), India, for supporting the project (SERB PROJECT EMR/2017/004051). Acharya Narendra Dev College (ANDC), University of Delhi, is acknowledged for providing the infrastructural facilities and ELITE fellowship to RD. KN received the Innovation in Science Pursuit for Inspired Research (INSPIRE) fellowship from the Department of Science and Technology (DST), India; RA received the Senior Research Fellowship (SRF) from the University Grants Commission (UGC), New Delhi, India. The graphical abstract was created using BioRender.com (https://biorender.com).

ORCID

RD: 0000-0003-3171-2011

RA: 0000-0001-7666-7790

KN: 0009-0007-8807-730X

UB: 0000-0003-2441-6089

Author Contribution

RD: Data curation, Software, Visualization, Investigation, Validation, Writing- original draft. KN: Data curation, Experimental Investigation, Visualization, Validation. RA: Primer-designing, Data curation, Visualization.   UB: Conceptualization, Methodology, Experimental Investigation Supervision, Validation, Writing– review & editing.

Funding

This study was supported by the funding provided to the project investigator UB by the Science and Engineering Research Board (SERB), Department of Science and Technology (DST), India

(SERB PROJECT EMR/2017/004051).

Availability of data and materials

This published article and its supplementary information files include all data generated or

analyzed during this study.

Ethics approval and consent to participate.

Not applicable.

Competing interests

The authors declare that they have no competing interests.

555
556

557 **Supplementary Information**
558
559 **Table S1:** Genome annotation of phiE1336.
560

Gene	Start	Stop	Strand	Annotation	Domain
gp1	1	2469	+	Phage tail protein	-
gp2	2595	3632	+	Phage tail protein	-
gp3	3654	3821	+	HP	-
gp4	3825	4187	+	HP	-
gp5	4376	5575	+	LysinA	Amidase/PGRP
gp6	6065	5772	-	HP	-
gp7	6081	6191	+	HP	-
gp8	6188	7147	+	LysinB_MF	PG binding and AB_hydrolase
gp9	7147	7236	+	HP	-
gp10	7242	7457	+	Holin	-
gp11	7454	7900	+	HP	-
gp12	8509	8093	-	HP	-
gp13	8899	8513	-	HP	-

562
563
564 **Figure S1: Comparison of domain organization of LysinBs.** Domain prediction and architecture
565 of well-studied LysinBs indicate the modular nature of LysinB_MF. Unlike other LysinBs, this
566 lysin has a PGBD in the N-terminal end of the enzyme.
567

A.

B.

Figure S2: Structural prediction of PGBD from LysinB_MF and alignment with an amidase. A) To
facilitate our understanding of the structural similarity of PGBD from LysinB_MF, we separately built its 3D
model using the structure of the lysin as a template. The structure retains its original α-helices, as observed in
the native LysinB_MF. However, the single β-strand is missing. The structure shows very high confidence and
was taken forward for further analysis. B) A zinc-dependent amidase (PDB accession 4XXT), a peptidoglycan-
binding domain-containing protein from *Clostridium acetobutylicum* (ATCC 824), displays high structural
similarity with the PGBD from LysinB_MF. The two structures can be superimposed and superpositioned with
an overall RMSD as low as 0.46.
578

References

Abad L, Gauthier CH, Florian I, Jacobs-Sera D, Hatfull GF. 2023. The heterogeneous and diverse population of prophages in Mycobacterium genomes. *mSystems* **8**(5): e0044623.

Abouhmad A, Korany AH, Grey C, Dishisha T, Hatti-Kaul R. 2020. Exploring the Enzymatic and Antibacterial Activities of Novel Mycobacteriophage Lysin B Enzymes. *Int J Mol Sci* **21**(9).

Aliashkevich A, Cava F. 2022. LD-transpeptidases: the great unknown among the peptidoglycan cross-linkers. *Febs Journal* **289**(16): 4718-4730.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. 1990. Basic local alignment search tool. *J Mol Biol* **215**(3): 403-410.

Bagcchi S. 2023. WHO's Global Tuberculosis Report 2022. *Lancet Microbe* **4**(1): e20.

Basu MK, Poliakov E, Rogozin IB. 2009. Domain mobility in proteins: functional and evolutionary implications. *Brief Bioinform* **10**(3): 205-216.

Bord S, Horner A, Hembry RM, Compston JE. 1998. Stromelysin-1 (MMP-3) and stromelysin-2 (MMP-10) expression in developing human bone: potential roles in skeletal development. *Bone* **23**(1): 7-12.

Bradford MM. 1976. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem* **72**: 248-254.

Cote J, Welch C, Kimble M, Archambault D, Ross JC, Orellana H, Amero K, Bourett C, Daigle A, Hutchison KW, et al. 2022. Characterization of the cluster MabR prophages of Mycobacterium abscessus and Mycobacterium chelonae. *G3 (Bethesda)* **12**(9).

Das R, Bajpai U. 2023. Functional characterization of a DNA-dependent AAA ATPase in a F-cluster mycobacteriophage. *Virus Res* **323**: 198957.

Das R, Eniyan K, Bajpai U. 2021. Computational identification and characterization of antigenic properties of Rv3899c of Mycobacterium tuberculosis and its interaction with human leukocyte antigen (HLA). *Immunogenetics* **73**(5): 357-368.

Eniyan K, Sinha A, Ahmad S, Bajpai U. 2020. Functional characterization of the endolysins derived from mycobacteriophage PDRPxv. *World J Microbiol Biotechnol* **36**(6): 83.

Fraga AG, Trigo G, Murthy RK, Akhtar S, Hebbur M, Pacheco AR, Dominguez J, Silva-Gomes R, Goncalves CM, Oliveira H, et al. 2019. Antimicrobial activity of Mycobacteriophage D29 Lysin B during Mycobacterium ulcerans infection. *PLoS Negl Trop Dis* **13**(8): e0007113.

Fraga K, Maireles M, Jordan M, Soldevila L, Murillo O. 2022. Mycobacterium fortuitum osteomyelitis of the cuboid bone treated with CERAMENT G and V: a case report. *J Bone Jt Infect* **7**(4): 163-167.

Gigante AM, Hampton CM, Dillard RS, Gil F, Catalao MJ, Moniz-Pereira J, Wright ER, Pimentel M. 2017. The Ms6 Mycolyl-Arabinogalactan Esterase LysB is Essential for an Efficient Mycobacteriophage-Induced Lysis. *Viruses* **9**(11).

Gigante AM, Olivenca F, Catalao MJ, Leandro P, Moniz-Pereira J, Filipe SR, Pimentel M. 2021. The Mycobacteriophage Ms6 LysB N-Terminus Displays Peptidoglycan Binding Affinity. *Viruses* **13**(7).

Gordon SV, Parish T. 2018. Microbe Profile: Mycobacterium tuberculosis: Humanity's deadly microbial foe. *Microbiology (Reading)* **164**(4): 437-439.

Grant JR, Stothard P. 2008. The CGView Server: a comparative genomics tool for circular genomes. *Nucleic Acids Res* **36**(Web Server issue): W181-184.

Hatfull GF. 2018. Mycobacteriophages. *Microbiol Spectr* **6**(5).

Hatfull GF. 2022. Mycobacteriophages: From Petri dish to patient. *PLoS Pathog* **18**(7): e1010602.

Ho YS, Adroub SA, Aleisa F, Mahmood H, Othoum G, Rashid F, Zaher M, Ali S, Bitter W, Pain A, et al. 2012. Complete genome sequence of Mycobacterium fortuitum subsp. fortuitum type strain DSM46621. *J Bacteriol* **194**(22): 6337-6338.

Holm L. 2022. Dali server: structural unification of protein families. *Nucleic Acids Res* **50**(W1): W210-W215.

Hurst-Hess K, Walz A, Yang Y, McGuirk H, Gonzalez-Juarrero M, Hatfull GF, Ghosh P, Ojha AK. 2023. Intrapulmonary Treatment with Mycobacteriophage LysB Rapidly Reduces Mycobacterium abscessus Burden. *Antimicrob Agents Chemother* **67**(6): e0016223.

Jacobs-Sera D, Marinelli LJ, Bowman C, Broussard GW, Guerrero Bustamante C, Boyle MM, Petrova ZO, Dedrick RM, Pope WH, Science Education Alliance Phage Hunters Advancing G, et al. 2012. On the nature of mycobacteriophage diversity and host preference. *Virology* **434**(2): 187-201.

Khosravi AD, Mirsaeidi M, Farahani A, Tabandeh MR, Mohajeri P, Shoja S, Hoseini Lar KhosroShahi SR. 2018. Prevalence of nontuberculous mycobacteria and high efficacy of d-cycloserine and its synergistic effect with clarithromycin against Mycobacterium fortuitum and Mycobacterium abscessus. *Infect Drug Resist* **11**: 2521-2532.

Knauper V, Lopez-Otin C, Smith B, Knight G, Murphy G. 1996. Biochemical characterization of human collagenase-3. *J Biol Chem* **271**(3): 1544-1550.

Korany AH, Abouhmad A, Bakeer W, Essam T, Amin MA, Hatti-Kaul R, Dishisha T. 2019. Comparative Structural Analysis of Different Mycobacteriophage-Derived Mycolylarabinogalactan Esterases (Lysin B). *Biomolecules* **10**(1).

Korndorfer IP, Danzer J, Schmelcher M, Zimmer M, Skerra A, Loessner MJ. 2006. The crystal structure of the bacteriophage PSA endolysin reveals a unique fold responsible for specific recognition of Listeria cell walls. *J Mol Biol* **364**(4): 678-689.

Laemmli UK. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**(5259): 680-685.

Letunic I, Bork P. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation. *Nucleic Acids Res* **49**(W1): W293-W296.

Low LY, Yang C, Perego M, Osterman A, Liddington RC. 2005. Structure and lytic activity of a Bacillus anthracis prophage endolysin. *J Biol Chem* **280**(42): 35433-35439.

Mishra LS, Kim SY, Caddell DF, Coleman-Derr D, Funk C. 2021. Loss of Arabidopsis matrix metalloproteinase-5 affects root development and root bacterial communities during drought stress. *Physiol Plant* **172**(2): 1045-1058.

Nielsen JE, Beier L, Otzen D, Borchert TV, Frantzen HB, Andersen KV, Svendsen A. 1999. Electrostatics in the active site of an alpha-amylase. *Eur J Biochem* **264**(3): 816-824.

Payne K, Sun Q, Sacchettini J, Hatfull GF. 2010. Mycobacteriophage Lysin B is a novel mycolylarabinogalactan esterase. *Molecular Microbiology*.

Payne KM, Hatfull GF. 2012. Mycobacteriophage endolysins: diverse and modular enzymes with multiple catalytic activities. *PLoS One* **7**(3): e34052.

Pollenz RS, Bland J, Pope WH. 2022. Bioinformatic characterization of endolysins and holin-like membrane proteins in the lysis cassette of phages that infect Gordonia rubripertincta. *PLoS One* **17**(11): e0276603.

Porollo A, Meller J. 2010. POLYVIEW-MM: web-based platform for animation and analysis of
molecular simulations. *Nucleic Acids Res* **38**(Web Server issue): W662-666.
Qiu S, Azofra LM, MacFarlane DR, Sun C. 2018. Hydrogen bonding effect between active site
and protein environment on catalysis performance in H2-producing [NiFe] hydrogenases.
Physical Chemistry Chemical Physics.
Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, Lopez R. 2005.
InterProScan: protein domains identifier. *Nucleic Acids Res* **33**(Web Server issue): W116-
120.
Robert X, Gouet P. 2014. Deciphering key features in protein structures with the new ENDscript
server. *Nucleic Acids Res* **42**(Web Server issue): W320-324.
Schmitz JE, Ossiprandi MC, Rumah KR, Fischetti VA. 2011. Lytic enzyme discovery through
multigenomic sequence analysis in Clostridium perfringens. *Appl Microbiol Biotechnol*
89(6): 1783-1795.
Sethi S, Arora S, Gupta V, Kumar S. 2014. Cutaneous Mycobacterium fortuitum Infection:
Successfully Treated with Amikacin and Ofloxacin Combination. *Indian J Dermatol* **59**(4):
383-384.
Shah S, Das R, Chavan B, Bajpai U, Hanif S, Ahmed S. 2023. Beyond antibiotics: phage-
encoded lysins against Gram-negative pathogens. *Front Microbiol* **14**: 1170418.
Singh AK, Gangakhedkar R, Thakur HS, Raman SK, Patil SA, Jain V. 2023.
Mycobacteriophage D29 Lysin B exhibits promising anti-mycobacterial activity against
drug-resistant Mycobacterium tuberculosis. *Microbiol Spectr* **11**(6): e0459722.
Suzuki S, Yoshikawa M, Imamura D, Abe K, Eichenberger P, Sato T. 2020. Compatibility of
Site-Specific Recombination Units between Mobile Genetic Elements. *iScience* **23**(1):
100805.
Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of
progressive multiple sequence alignment through sequence weighting, position-specific
gap penalties and weight matrix choice. *Nucleic Acids Res* **22**(22): 4673-4680.
Varadi M, Bertoni D, Magana P, Paramval U, Pidruchna I, Radhakrishnan M, Tsenkov M,
Nair S, Mirdita M, Yeo J, et al. 2023. AlphaFold Protein Structure Database in 2024:
providing structure coverage for over 214 million protein sequences. *Nucleic Acids Res*.
Zheng ZL, Ye MQ, Zuo ZY, Liu ZG, Tai KC, Zou GL. 2006. Probing the importance of
hydrogen bonds in the active site of the subtilisin nattokinase by site-directed mutagenesis
and molecular dynamics simulation. *Biochem J* **395**(3): 509-515.
Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. 2011. PHAST: a fast phage search tool.
Nucleic Acids Res **39**(Web Server issue): W347-352.
705