Multiomics uncovers the epigenomic and transcriptomic response to viral and bacterial stimulation in turbot

Oscar Aramburu [1]*, Belén Gómez-Pardo[1], Paula Rodríguez-Villamayor[1], Andrés Blanco-Hortas[1], Jesús Lamas[1], Pooran Dewari[2], Diego Perojil-Morata[2], Pierre Boudinot[3], Daniel J. Macqueen[2], Carmen Bouza[1], Paulino Martínez[1]*

[1] University of Santiago de Compostela, Spain

[2] The Roslin Institute and Royal (Dick) School of Veterinary Studies, University of Edinburgh, Easter Bush Campus, UK

[3]Université Paris-Saclay, INRAE, UVSQ, VIM, Jouy-en-Josas, 78350, France.

* Corresponding authors

Abstract

Uncovering the epigenomic regulation of immune responses is essential for a comprehensive understanding of host defence mechanisms but remains poorly described in farmed fish. Here, we report the first annotation of the innate immune regulatory response in the genome of turbot (*Scophthalmus maximus*), a farmed flatfish. We integrated RNA-Seq with ATAC-Seq and ChIP-Seq (histone marks H3K4me3, H3K27ac and H3K27me3) using samples from head kidney. Sampling was performed 24 hours post-stimulation with viral (poly I:C) and bacterial (inactivate *Vibrio anguillarum*) mimics *in vivo* and *in vitro* (primary leukocyte cultures). Among the 8,797 differentially expressed genes (DEGs), we observed enrichment of transcriptional activation pathways in response to *Vibrio* and immune response pathways - including interferon stimulated genes - for poly I:C. Meanwhile, metabolic and cell cycle were downregulated by both mimics. We identified notable differences in chromatin accessibility (20,617 *in vitro*, 59,892 *in vivo*) and H3K4me3 bound regions (11,454 *in vitro*, 10,275 *in vivo*) - i.e. marking active promoters - between stimulations and controls. Overlaps of DEGs with promoters showing differential accessibility or histone mark binding revealed a significant coupling of the transcriptome and chromatin state. DEGs with activation marks in their promoters were enriched for similar functions to the global DEG set, but not in all cases, suggesting key regulatory genes were in poised or bivalent states. Active promoters and putative enhancers were differentially enriched in transcription factor binding motifs, many of them common to viral and bacterial responses. Finally, an in-depth analysis of immune response changes in chromatin state surrounding key DEGs encoding transcription factors was performed. This comprehensive multi-omics investigation provides an improved understanding of the epigenomic basis for the turbot immune responses and provides novel functional genomic information that can be leveraged in selective breeding towards enhanced disease resistance.

Keywords: turbot, immune response, epigenomics, chromatin state, transcription factor

INTRODUCTION

The functional annotation of farm animal genomes is important for understanding traits with complex genetic architecture, such as disease resistance, growth, feed efficiency or reproduction (Andersson et al., 2015; Tuggle et al., 2016; Giuffra et al., 2019; Clark et al., 2020). Until recently, functional annotation mainly focused on protein-coding genes using transcriptomics. Transcriptome annotation is now consolidated with robust pipelines (Raghavan et al., 2022) and, throughout the years, transcriptome annotations for human (Frankish et al., 2023), model species (He et al., 2020; Lawson et al., 2020), terrestrial livestock (Summers et al., 2020; Halstead et al., 2021; Overbey et al., 2021) and some aquaculture species (Ramberg et al., 2021; Johnston et al., 2024) have been published.

Non-coding regulatory elements, including promoters, enhancers, silencers and insulators, have been studied in several livestock species but are mostly unexplored in aquaculture species. These elements play essential roles in regulating gene expression and their state can change depending on tissue, cell type, sex, age, and health status (Kellis et al., 2014). Thus, annotation of regulatory elements in different contexts not only aids to address basic questions related to morphology and physiology (Halstead et al., 2020; Pan et al., 2023), but also functional genomic responses to environmental variation (Feinberg, 2007; Ecker et al., 2018; Hu and Barrett, 2017). Genetic variation at non-coding elements also underpins phenotypic variation (Villar et al., 2020; Boltsis et al., 2021), and can thus be leveraged to improve our ability to predict polygenic traits using genomic data (Zhu et al., 2023). In this respect, more than 90% of phenotype-associated single nucleotide polymorphisms (SNPs) identified in human genome-wide association studies (GWAS) are located in non-coding regions (Giral et al., 2018), with similar results reported for livestock (Prowse-Wilkins et al., 2021).

In the past decade, human and livestock functional annotation initiatives have investigated epigenetic mechanisms involved in gene regulation through the study of chromatin state modifications across the genome (Yan et al., 2020a). Chromatin can switch dynamically between active and inactive states in minutes to hours, leaving epigenetic footprints that can be transmitted vertically following DNA replication (Moazed, 2011; Cuvier and Fierz, 2017). Many different sequencing assays have been developed to infer chromatin epigenetic status, including chromatin accessibility (ATAC-Seq; Buenrostro et al., 2015), protein-DNA interactions (ChIP-Seq; Park, 2009) and long-range chromatin interactions (Hi-C; van Berkum et al., 2010). These and other assays are being applied by the FAANG Consortium (Functional Annotation of Animal Genomes; Fang et al., 2019; Foissac et al., 2019; Liu et al., 2020; Clark et al., 2020). Current annotations of chromatin state and regulatory elements remain limited to a few terrestrial farm animal species (Summers et al., 2020, Pan et al., 2021; 2023; Xiang et al., 2021). However, a catalogue of regulatory elements is being generated for several important fish species used in global aquaculture (Johnston et al., 2024), currently the fastest growing animal production sector (Subasinghe et al., 2009; Troell et al., 2023).

Turbot (*Scophthalmus maximus*) is a valuable farmed fish in Europe and Asia (more than 100,000 tons), with the highest production in China (Gao et al., 2023) followed by Spain (APROMAR 2022). Turbot is in its 6[th] generation of selective breeding and infectious disease outbreaks constitute one of the main challenges this young industry faces (Martínez et al., 2021; Gao et al., 2023). This is a broader trend shared by global aquaculture, where infectious diseases cause losses of totalling more than 5,000 M€ per year (Mishra et al., 2018). Functional annotation of

the turbot transcriptome has been performed against high-quality reference genomes (Figueras
et al., 2016; Maroso et al., 2018; Xu et al., 2020; Martinez et al., 2021), including for immune-
organs stimulated with viruses (Diaz-Rosales et al., 2012), bacteria (Millán et al., 2011; Libran-
Pérez et al., 2022) and parasites (Pardo et al., 2012; Robledo et al., 2014; Ronza et al., 2016; Valle
et al., 2020). Candidate genes for disease resistance have been further explored by mapping
differentially expressed genes (DEGs) within QTL regions (Martínez, 2016; Saura et al., 2019;
Aramburu et al., 2023). However, limited attention has been given to non-coding regulatory
elements, beyond a recent analysis of chromatin accessibility focussed on early development
(Guerrero-Peña et al., 2023). How chromatin state and non-coding regulatory elements are
regulated during immune responses remains undefined in turbot and scarcely explored in other
farmed finfish.

The head kidney has been targeted in all previous functional genomics studies in turbot
investigating pathogen responses (Millán et al., 2011; Díaz-Rosales et al., 2012; Pardo et al.,
2012; Robledo et al., 2014; Ronza et al., 2016; Librán-Pérez et al., 2022; Aramburu et al., 2023),
due to its central role in fish immunity (Mokhtar et al., 2023). Head kidney is a key lymphoid
organ in most marine fishes and, analogous to the mammalian bone marrow, responsible for the
production of high leukocyte diversity, including B-lymphocytes, early-stage T-lymphocytes, as
well myeloid cells such as granulocytes and monocytes/macrophages (Klosterhoff et al., 2015;
Geven et al., 2017; Chen et al., 2022). Innate immunity provides the first line of defence against
pathogens and acts following the binding of pathogen-associated molecular patterns (PAMPs) to
germline pattern recognition receptors (PRRs), leading to various effector cellular functions
targeting pathogen destruction and clearance.

The aim of this study was to generate the first comprehensive functional annotation of the innate
immune response of turbot using a chromosome-level reference genome sequence (Martínez et
al., 2021). Live fish and primary immune cell cultures were stimulated using mimics of viral and
bacterial infections and compared to controls to capture changes in the transcriptome alongside
chromatin accessibility and epigenetic state by integrating RNA-Seq, ATAC-Seq and ChIP-Seq
data. Our results provide a deeper understanding of the epigenomic basis for innate immunity
in turbot and a novel resource to prioritize genetic variation associated with non-coding
elements regulating immune responses.

MATERIALS AND METHODS

Animals

Thirty 8-month-old immature turbot specimens provided by Stolt Sea Farm SA (Ribeira, Spain)
were housed in indoor tanks with recirculating seawater at the facilities of the Aquarium of the
University of Santiago de Compostela (Spain) for a period of acclimation of 15 d at 16 °C
(Supplementary table 1). All fish were fasted for 24 h before stimulations were performed.
Eighteen fish were stimulated *in vivo* by intraperitoneal injection, while the other 12 were used
for leukocyte isolation for *in vitro* stimulation (see following sections). Fish were anesthetized by
bath (MS-222; 100mg/L) and then euthanized by anaesthetic overdose (MS-222; 150 mg/L)
before tissue sampling. All animal procedures were approved by the Bioethics Committee of the
University of Santiago de Compostela (body authorized according to R.D. 53/2013) and with the
authorization of the Xunta de Galicia Regional Government.

Protocols

Detailed protocols for the *in vivo* and *in vitro* stimulations, RNA isolation, ATAC-Seq and ChIP-Seq
(including library preparation) are available in the FAANG repository (data.faang.org; URLs for
protocols in Supplementary tables 1 and 2).

***In vivo* immunostimulation**

Six fish were used per experimental condition for the *in vivo* stimulations: i) poly I:C for viral
mimic immunostimulation; ii) killed *Vibrio anguillarum* for bacterial immunostimulation; and iii)
phosphate buffered saline (PBS) for control. For poly I:C (Sigma P1530), we prepared a working
stock at 5 mg/ml in PBS, preheated to 55 °C (15 min) and cooled at room temperature (20 min)
before use. Fish were then injected intraperitoneally with 5 µg per g fish weight. For bacterial
immunostimulation, an extract of *V. anguillarum* (strain P0382; INRA, France) was used. Bacteria
were cultured in a tryptic soy broth medium to an OD600 (optical density at 600 nm) of 1.5. The
bacterial pellet (derived from 100 ml of full-grown culture) was washed in an isotonic solution
of NaCl (9 g/L) four times and resuspended in 1 ml of the same solution. Bacteria were killed by
incubation for 30 sec at 100 °C, allowed to cool at room temperature and stored at -80 °C. The
bacteria extract was inoculated in each specimen, diluted in PBS (1:10) for a final volume of 100
µl. Control fish were injected with 100 µl of PBS. After 24 h, head kidney samples were extracted,
washed with PBS and cut into at least three pieces (> 20 mg each); two were flash frozen on dry
ice for ATAC-Seq and ChIP-Seq and the other was immersed in RNAlater (Thermofisher Scientific)
for RNA extraction and RNA-Seq. All samples were then stored at -80°C (Figure 1A;
Supplementary table 2).

***In vitro* stimulation: leukocyte isolation and culture**

Leukocytes were isolated from 12 fish. The entire head kidney was aseptically isolated and placed
in a Petri dish with 40 ml of cell isolation media (500 ml of Leibovitz L-15 medium (L-15), 10 ml
FBS (2 %), 0.02 % EDTA). Samples were then cut into small pieces and passed through a 100 µm
nylon mesh with constant flow of cell isolation media. Leukocytes were separated by
centrifugation of 40 ml of the cell suspension gently layered in a 50 ml tube containing 51 %
Percoll (400 x g, 30 min, 4 °C, no brakes). The interface layer was collected by centrifugation (400
x g, 10 min, 4 °C) and washed three times with L-15 medium containing 0.1 FBS, keeping the
pelleted leukocytes (Figure 1B).

To ensure enough cells were available, samples were pooled by pairs after cell counting and cell
viability was evaluated by the trypan blue exclusion test, totalling 6 pools (each 18×10^6 cells;
Supplementary table 1). Then, each pool was divided into nine aliquots of 2×10^6 cells (in 2 ml)
that were dispensed into wells, for a total of 54 wells (6 pools x 9 aliquots): 18 stimulated with
20 µl of poly I:C solution; 18 with 20 µl of inactivated *V. anguillarum*; and the remaining 18 wells
used as controls (Figure 1B). All leukocyte cultures were incubated for 24 h at 16 °C. Cells were
then collected in 2 ml Eppendorf tubes and pelleted at 500 x g for 5 mins at room temperature.
Eighteen pellets were resuspended in RNAlater and stored at -80 °C for RNA-Seq, while the other
36 were flash frozen in dry ice and stored at -80 °C for ATAC-Seq and µChIPmentation,
respectively (Supplementary table 1).
162

164 **Figure 1.** Experimental design followed for turbot immunostimulations: **A)** *in vivo*
165 immunostimulation of immature fish with subsequent sampling of head kidney; **B)** *in vitro*
166 stimulation of head kidney leukocytes of immature fish; 18 samples were used for RNA-Seq, 18
167 for ATAC-Seq and 18 for ChIP-Seq/µChIPmentation with antibodies to H3K4me3, H3K27ac and
168 H3K27me3 histone marks.

RNA isolation and sequencing

After sample thawing, total RNA was extracted and purified using the miRNeasy Kit (QIAGEN)
with specific modifications for: i) frozen head kidney (> 20 mg), following a protocol for "Total
RNA extraction for tissues", and ii) frozen leukocytes (2×10^6 cells) following a protocol for "Total
RNA extraction for frozen cells" (Supplementary table 2). RNA integrity and quantity were
evaluated in a Bioanalyzer (Bonsai Technologies, Madrid, Spain) and in a NanoDrop® ND-1000
spectrophotometer (NanoDrop® Technologies Inc., Wilmington, DE, USA). RNA integrity number
(RIN) averaged 8.3 across all samples, always above 7.3. RNA samples were delivered to
Novogene (UK) for library preparation using NEBNext Ultra Directional RNA Library Prep Kits for
Illumina and sequenced using an Illumina NovaSeq S4 platform to generate 150 bp paired end
reads.

RNA-Seq data processing

RNA-Seq data was processed using nf-core/rnaseq 3.10.1 (Patel et al., 2023) run with default
parameters, using the turbot Ensembl genome ASM1334776v1 (Martínez et al., 2021) as
reference. In brief, the pipeline evaluated quality of raw reads using FASTQC (Andrews, 2010)
and trimmed adapters and low-quality bases using Trim Galore! (Martin et al., 2011). Reads were
then mapped using STAR (Dobin et al., 2013). Normalized transcript read counts were obtained
using RSEM (Li and Dewey, 2011). After nf-core processing, the resulting count tables were
filtered to remove genes with expression below 5 transcripts per million (TPM < 5) and
represented in only one sample across all conditions.

Differential gene expression and gene ontology analysis

Differentially expressed genes (DEGs) between stimulated and control samples were identified
using the R/Bioconductor package DESeq2 v1.38.1 (Love et al., 2014). Genes with false discovery
rate (FDR) adjusted p < 0.05 were considered DEGs. Functional enrichment of the DEG lists was
performed using ShinyGO v0.77 (Ge et al., 2020). Gene ontology (GO) terms for Biological

Process of each list were ranked by statistical significance (FDR-adjusted p < 0.05). All expressed
genes across conditions were used as the background for GO analyses.

ATAC-Seq: Library preparation and sequencing

Following a standard protocol for "Nuclei isolation for ATAC-Seq procedures" (Supplementary
table 2), the frozen head kidney fragments (> 20 mg; *in vivo* assay) and cell pellets ($2x10^6$ cells;
in vitro assay) were thawed and resuspended in 1 ml of TST buffer. Each tissue fragment was cut
into smaller pieces with a scalpel, mashed with the rubber back of a syringe, and filtered through
a 40 μm cell strainer, while cell pellets were resuspended by gentle pipetting. The number and
integrity of nuclei was assessed with a haemocytometer (minimum of ~50,000 nuclei in a 16.5
μl suspension; ~3,000 nuclei/μl) before carrying out the Tn5 transposase reaction with Illumina
Tagment DNA TDE1 enzyme (37°C, 30 min, 1000 RPM, Illumina) following the standard
"OmniATAC protocol" (Corces et al., 2017; Supplementary table 2). The resulting DNA was
purified with a MinElute PCR purification kit (Qiagen), and DNA concentration assessed with a
Qubit using the dsDNA HS kit (ThermoFisher Scientific). Library amplification (10-12 PCR cycles)
was carried out using the NEBNext Ultra II DNA Library Prep Kit (New England Biolabs), with IDT
for Illumina UD Indexes (96x, Plate A, Set 1, Illumina). Library size selection was performed to
remove fragments below 180 bp and above 700 bp using AMPure XP beads (Beckman coulter).
Finally, DNA fragment size distribution was assessed with the Bioanalyzer High Sensitivity DNA
Assay kit (Agilent Technologies). ATAC-Seq libraries were delivered to Novogene (UK) to be
sequenced on an Illumina NovaSeq S4 platform generating 150 bp paired end reads.

ChIP-Seq and μChIPmentation: Library preparation and sequencing

The frozen head kidney fragments (> 20 mg) and leukocyte pellets ($2x10^6$ cells) were thawed on
ice. Following a standard "ChIP-seq" protocol (Supplementary table 2), the tissue fragments were
transferred into a Douncer homogenizer containing a protease inhibitor cocktail (PIC, Roche, 1
tablet in 50 ml of PBS) solution immersed in ice and homogenized using pestles A and B (from
less to more plunger adjustment). The leukocyte pellets were resuspended in the PBS and PIC
solution. The amount and quality of nuclei was assessed with a haemocytometer and trypan
blue staining (> 10 million cells for head kidney; > 100,000 for leukocyte cultures). Due to the
low number of nuclei recovered from the leukocyte cultures, a μChIPmentation protocol
(Diagenode) was used, following a modified protocol (Supplementary table 2).

In both cases, chromatin crosslinking was done using a 1 % formaldehyde solution followed by
quenching with glycine (0.125 M). Nuclei were pelleted and resuspended in complete sonication
buffer, while leukocyte nuclei were resuspended in Hanks' Balanced Salt Solution (HBSS, Thermo
Fisher Scientific) - tL1 buffer (Supplementary table 2). The chromatin was sheared using a Covaris
S2 focused ultrasonicator with the following parameters: 2 % duty cycle, intensity 3, with 200
cycles per burst, at 4 °C for 8 min and 6 min for head kidney tissue and leukocytes, respectively.

The immunoprecipitation was performed using Diagenode antibodies for three marks: H3K4me3
(marking active promoter regions; cat. No. C15410003; 1.3 μg/μl), H3K27ac (marking active
enhancer and promoter regions; cat. No. C15410196; 2.8 μg/μl) and H3K27me3 (marking
Polycomb repressed regions; cat. No. C15410195; 1.1 μg/μl). For head kidney samples,
antibodies were coupled with pre-washed protein A and protein G beads, and the tubes left
under rotation overnight (~16 h) at 4 °C, following the standard ChIP-Seq protocol. After washing
the beads and decrosslinking, the samples were purified using the MinElute PCR purification kit
(Qiagen) and later quantified with DNA HS Qubit (ThermoFisher Scientific). Immunoprecipitated
chromatin was stored at -20 °C until library preparation, using a Microplex v3 kit (Diagenode).

For the leukocyte samples, the µChipmentation kit for histones (Diagenode) was used for
chromatin immunoprecipitation and ChIP-Seq library preparation (Supplementary table 2).

Before sequencing, the quantity and quality of purified libraries was assessed using the Qubit
DNA HS kit (ThermoFisher Scientific) and the High Sensitivity DNA Assay kit (Agilent
Technologies), respectively. A minimum of 60 % of the chromatin was required to have a size
distribution between 200-700 bp (centered around 350-400 bp). ChIP-Seq libraries were
delivered to Novogene (UK) for sequencing on an Illumina NovaSeq S4 platform generating 150
bp paired-end reads.

ATAC-Seq and ChIP-Seq data processing

ATAC-Seq and ChIP-Seq data were processed using the nf-core/atacseq v1.2.2. and nf-
core/chipseq v1.2.2 pipelines (Ewels et al., 2023), respectively, run with the narrow_peak option
for ATAC-Seq and H3K4me3 and H3K27ac ChIP-Seq datasets, and with the broad_peak option for
H3K27me3. The other parameters were kept by default. Quality assessment of the reads was
carried out with FASTQC (Andrews, 2010), and adapters and low-quality bases trimmed with Trim
Galore! (Martin et al., 2011). Reads were mapped to the turbot genome using BWA (Li and
Durbin, 2009). Further filtering was done with SAMtools (Danecek et al., 2021), BAMtools
(Barnett et al., 2011) and Pysam (Danecek et al., 2021). Genome-wide immunoprecipitation (IP)
enrichment relative to controls was done with deepTools (Ramírez et al., 2014) and
broad/narrow peaks were called using MACS2 (Zhang et al., 2008). Once nf-core was finished,
suboptimal replicates with very low peak numbers were excluded after visualizing their bigwig
files on the Integrative Genomics Viewer (IGV; Robinson et al., 2023).

ChIP-Seq and µChIPmentation blacklist

To improve the signal-to-noise ratio of the ChIP-Seq and µChIPmentation data, a blacklist
consisting of high signal and low mappability regions was constructed using ChIP-Seq and
µChIPmentation inputs, including 21 control ChIPseq turbot samples (ENA accession
PRJEB57784). The mappability of the turbot genome for read lengths of 100 bp and 150 bp (k-
mers 100 and 150) was quantified using the umap software package (Karimzadeh et al., 2018).
The generated mappability files were fed into the ENCODE blacklist software (Amemiya et al.,
2019) to generate the blacklist. Alignment and peak files derived from the nf-core pipeline were
filtered with BAMtools to remove reads and peaks located in the blacklist regions.

Differential histone modification regions and differentially accessible regions

271 Differential histone modification regions (DHMRs) and differentially accessible regions (DARs;
272 adjusted p < 0.05) between stimulated samples and controls were identified using DiffBind (Stark
273 and Brown, 2011) with default settings.

Integration of ChIP-Seq and ATAC-Seq data with RNA-Seq data

275 For each condition tested, the promoters of DEGs overlapping with regions tagged as DARs
276 and/or DHMRs were identified. When applicable, a hypergeometric test was performed to check
277 the significance of overlapping between each pair (p < 0.05).

Chromatin state inferences

Genome-wide chromatin states for each condition were predicted using ChromHMM (Ernst and
Kellis, 2012) integrating the ChIP-Seq data (µChIPmentation for *in vitro* samples) for the three
histone marks (H3K4me3, H3K27ac and H3K27me3) and ATAC-Seq data. Chromatin state

prediction was performed by testing ChromHMM models including from 8 to 15 states, keeping
the one that returned the most biologically relevant chromatin states, for head kidney and
leukocyte data separately (Pan et al., 2021; Baranasic et al., 2022; Vu and Enst, 2022). The
genome-wide distribution of resulting chromatin states was visualized on IGV. Regions annotated
as enhancer-related states by ChromHMM were retrieved, and each stimulation dataset was
compared against its respective control to identify potential enhancer-related regions. Then,
each list of regions including potential enhancers annotated as "intergenic" or "intron" were kept
as differential enhancer-state regions for further analysis.

Transcription factor motif analysis

Enriched transcription factor (TF) binding motifs (TFBM) included in the HOMER software (Heinz
et al., 2010) were identified using the *findMotifsGenome.pl* function (settings: -size given -mask
-mset vertebrates) in the different lists of promoter-associated DHMRs/DARs and putative
enhancers for each condition. Random genomic regions with GC-content matching each input
genomic list were used as the background for automatic motif analysis by HOMER. Only TFBMs
with $p < 0.01$ and percentage of target sequences > 5 % were considered enriched.

GO analysis of the TFs predicted to bind enriched TFBMs was performed with Metascape (Zhou
et al., 2019) using zebrafish *Danio rerio* as the reference. Finally, a selection of TF genes that were
DEGs and contained promoter regions overlapping DARs or DHMRs were explored with IGV
(Robinson et al., 2023).

RESULTS

Raw sequencing data and sample metadata

A total of 186 multiomic datasets were produced in this study, including RNA-Seq (36), ATAC-Seq
(36) and ChIP-Seq (108; 36 per histone mark, plus 6 ChIP-Seq input controls). Full information on
samples and metadata is shared in Supplementary tables 1 and 2.

RNA-Seq

On average, 69,580,246 raw reads per library were produced across the 36 RNA-Seq samples,
with 97.2 % mapping to the turbot genome (Supplementary table 3A). Principal Component
Analysis (PCA) showed that 77 % of the transcriptome variance was explained by PC1, separating
the *in vivo* and *in vitro* stimulations (Figure 2A). Despite all three *in vivo* conditions grouping
together, a suggestive spatial segregation was observed mainly across PC1, with the control at
one end and the *Vibrio* stimulation at the other. For the *in vitro* stimulations, PC2 clearly
separated the *Vibrio* stimulation from poly I:C and control samples, the last two showing some
overlap (Figure 2B).

Table 1. Number of differentially expressed genes (DEGs) for the *in vitro* and *in vivo* stimulations with
Vibrio and poly I:C

Stimulant	Stimulation	Upregulated DEGs	Downregulated DEGs	Total DEGs
Vibrio	*in vitro*	3,321	3,217	6,538
	in vivo	910	929	1,839
poly I:C	*in vitro*	858	544	1,402
	in vivo	2,001	1,918	3,919

Differential expression analysis was performed by comparing each stimulated condition to the
respective controls, both for the *in vitro* and *in vivo* stimulations. In total, 8,797 DEGs were
identified across all comparisons (Table 1, Supplementary table 4). For the *in vitro* stimulations,
a stronger response was observed for *Vibrio* than poly I:C stimulation, both for up and
downregulated genes. Meanwhile, poly I:C showed a higher number of DEGs than *Vibrio* for the
in vivo stimulations both for up and downregulated genes (Table 1, Supplementary table 4).
Overall, more DEGs were detected in the *in vitro* than in the *in vivo* stimulations (7,940 vs 5,758,
respectively).

Functional enrichment among DEGs

Gene Ontology (GO) analysis identified enriched biological processes for up and downregulated
DEGs in all conditions (Supplementary table 5). Among the downregulated DEGs, metabolism,
cell cycle and cytoskeleton organization terms were enriched for the *in vitro* stimulations, while
a limited number of terms were detected for *in vivo* stimulations (Supplementary table 5).

Figure 2. PCA of the *in vivo* and *in vitro* transcriptomic response to stimulation with poly I:C and *Vibrio*.
A) PCA of the whole transcriptomic dataset; **B)** Independent PCAs for the *in vitro* and *in vivo* transcriptomic
assays.

More abundant and specific enriched GO terms were detected among the upregulated DEGs. Although RNA metabolism was enriched for most stimulations, poly I:C stimulation was linked to a more specific activation of key immune functions (particularly *in vitro*) such as interferon-stimulated genes and cytokine pathways, and regulation of toll-like receptors, besides more general immune terms. DEGs for the *in vitro Vibrio* stimulation were enriched for immune-related terms associated with cytokine and several transport pathways, whereas *in vivo* upregulated genes displayed strong enrichment of terms related to cytoskeleton organization, tissue development and syncytium formation (Supplementary table 5).

Comparing DEGs between conditions, important immune-related GO terms were commonly enriched between the *in vitro* poly I:C and *Vibrio* stimulations (Figure 3; Supplementary table 6), including interferon type I stimulated genes (*socs1a, socs1b, nod2, nmi*), cytokine signalling (the same genes plus *traf2*, *stat4* and *il15ra*) and major histocompatibility complex type I (MHC-I) pathways (*erap1b*, *tapbpl*, *tapbp*.2). Activation of transcription, protein localization, and a small number of terms associated with the immune-related "peptidyl-arginine modification" (*prmt1, prmt3, prmt5, prmt7*) were overrepresented *in vivo*.

We detected several DEGs regulated in opposite directions for poly I:C and Vibrio stimulations (Table 2; Figure 3; Supplementary table 6), suggestive of divergent immune responses for bacteria and virus. Among the *in vivo* upregulated DEGs for *Vibrio* and downregulated for poly I:C, we observed enrichment of general immune functions, including genes such as *il-1b*, *traf4a* and *f2r*. Conversely, immune-related genes including *nod2*, *sting1*, *irf1b* and *apaf1*, were downregulated for *Vibrio* and upregulated for poly I:C *in vivo*. Interestingly, most genes involved in the type-I IFN response orthologs of human and mouse (interferome database; Rusinova et al., 2023; Clark et al., 2023) were upregulated in the poly I:C stimulations, while downregulated with *Vibrio* extracts, especially in the *in vivo* stimulation (Table 2).

Figure 3. Venn diagrams illustrating overlap of DEGs for: **A)** *in vitro* poly I:C vs *Vibrio*, and **B)** *in vivo* poly I:C vs *Vibrio*. The white boxes show enriched GO terms related to the immune system found in specific subsets. Asterisks (*) highlight cases where suggestive immune-related GO terms were identified, despite not being significantly enriched. Double asterisks (**) highlight specific genes (in parentheses) associated with relevant immune functions. GO terms for the genes in the list were extracted from Ensembl-BioMart.

Table 2. Immune-related DEGs showing opposite responses following *Vibrio* and poly I:C stimulations.

Comparison	Differentially expressed genes (DEGs)
In vitro - upregulated by poly I:C, downregulated by *Vibrio*	**aldh2-like**, **arhgap22**, <u>arrdc2</u>, <u>asb9-like</u> **ascc3**, **atxn7l1**, bpifcl, c1qa, **cep170b**, chaf1b, cnnm4b, **dync1h1**, **ece2b**, **ermp1-like**, **gna14**, **gpr155a**, **has1**, **ip6k2-like**, **mapk8a**, **mycbp2**, nfkb1, **notch2**, pld4-like, psda, rilp, serinc1-like, sgk2b, sp100.1, **spsb4a**, **sptbn1**, stim2b, synpr, **tank**, tent4b, **tlr8**, tmem269-like, tnk2b, tpm4b, **trim25-like**, **trim35-14**, **usp1**, vcanb
In vitro - upregulated by *Vibrio*, downregulated by poly I:C	bcat-like, <u>ctns</u>, gid8b-like, pcxb, **psat1**, rab11b-like, rtf2, sfxn2, **slc25a43**, **stat3**, thoc7, tollip, vhl, vps26c, <u>vps41</u>
In vivo - upregulated by poly I:C, downregulated by *Vibrio*	adora4a, **ankrd10b**, **apaf1**, <u>arhgap12b</u>, atg9a, **atxn7l1**, bpifcl, camkk1b, **casp1a-like**, cylda, emilin1, **fem1c**, gfral, **gna14**, gnptab, <u>il22ra2</u>, **irf1b**, <u>nlrc3-like</u>, **nod2**, <u>nsmce4a</u>, otulina, **parp15-like**, **pstpip1b**, **pstpip2**, **rasgef1b**, rnf146, **rnf170**, **sting1**, **themis2**, **txk**, ubp15, <u>ucp2-like</u>, **xcr1-like**, **xkr8-like**
In vivo - upregulated by *Vibrio*, downregulated by poly I:C	aebp1, alpl, anxa13l, <u>anxa1a</u>, <u>cald1a</u>, capn2b, chchd6a, col17a1-like, <u>f2r</u>, <u>ggh-like</u>, icn-like, il-1b, <u>il8-like</u>, <u>il11-like</u>, kdm2aa, <u>lims2</u>, map1lc3b, mfsd10, msto1, olfm5-like, osbpl10, plpp1like, plxnb2b, <u>slc4a4-like</u>, tlr13-like, <u>traf4a</u>, <u>vat1</u>, <u>zan-like</u>

Turbot orthologs of human/mouse type I IFN stimulated genes (ISG) are highlighted (in bold upregulated and underlined downregulated), based on the database Interferome (Rusinova et al., 2013) and Clark et al. (2023). Note that most of these genes are downregulated in response to Vibrio anguillarum.

Considering condition-specific DEGs (Figure 3; Supplementary table 6), upregulated by poly I:C
in vitro DEGs were enriched for terms associated with toll-like receptor signalling pathway (*usp4,*
tasl, irak3), while transcriptional activation terms were found among DEGs in *Vibrio in vitro.* For
the *in vivo* stimulations, transcriptional activation GO terms were enriched for upregulated DEGs
after poly I:C stimulation, whereas response to stimulus (explained by *tlr3, hamp, tnip1, fxc1a*
and *ccr12a*, among other genes) and syncytium formation (explained by *kirrel3l, jam2a,*
plekho1b) were enriched for *Vibrio.*

ATAC-Seq and ChIP-Seq

Excluding some ATAC-Seq *in vitro* libraries not associated with the stimulant, most libraries
showed > 95% read mapping to the turbot genome (Supplementary table 3B). On average, we
identified 24,251 (*in vitro*) and 62,013 (*in vivo*) peaks per sample for ATAC-Seq; 26,199
(H3K4me3), 13,461 (H3K27ac) and 27,765 (H3K27me3) peaks for the *in vivo* ChIP-Seq data; and
15,870 (H3K4me3), 5,363 (H3K27ac) and 38,784 (H3K27me3) peaks for the *in vitro* ChIP-Seq
data. Samples in the PCA plot were mostly segregated across PC1, although PC2 explained most
of the variance between histone marks, particularly differentiating repressive (H3K27me3) from
active (H3K4me3 and H3K27ac) marks (Figure 4). *Vibrio* showed the greatest differentiation

among the stimulations, while poly I:C and controls were mostly intermingled, particularly for
ATAC-Seq (*in vitro* and *in vivo*) and H3K4me3-ChIP-Seq (*in vitro*).

Figure 4. PCA of the *in vivo and in vitro* epigenomic response to stimulation with poly I:C and *Vibrio*. **A)** PCA of the whole dataset; **B)** Separated PCAs for each assay and condition.

ChIP-Seq blacklist

Certain genomic regions obscure epigenetic analyses because of anomalous, unstructured, or
high signal due to particular genomic features (Amemiya et al., 2019). Using the 21 ChIP-Seq and
µChIPmentation input controls (ENA project PRJEB57784) we constructed a blacklist of low
confident genomic regions for turbot ChIP assays (Supplementary table 7). On average, 6.98 %
(~39Mb) of the turbot genome was included in the blacklist, consisted of high input signal (5.58
%) and low mappability (1.40 %) regions (Supplementary figure 1).

Chromatin state annotation

Genome-wide chromatin state predictions were produced using ChromHMM employing the
ChIP-Seq and ATAC-Seq data from stimulated and control samples from head kidney and
leukocytes (Figure 5, Supplementary table 8). For the *in vivo* head kidney samples (Figure 5A), a
10-state model was chosen including promoters/transcription start sites (TSS; States 1, 2, 3 and
4), potential enhancer regions (States 5 and 6), ATAC islands (State 7, i.e. ATAC-peaks lacking
histone marks), repressed regions (States 8 and 9) and low signal regions (State 10). For the *in*
vivo leukocytes (Figure 5B), an 8-state chromatin model was chosen including promoters/TSS
(States 1, 2 and 3), potential enhancer regions (States 4 and 5), ATAC islands (State 6), repressed
regions (State 7) and low signal regions (State 8). For each chromatin state map, ±2kb regions
around the TSS showed signal specifically for chromatin states 1, 2, 3 and 4 for head kidney (*in*
vivo) and states 1, 2 and 3 for leukocytes (*in vitro*), mostly corresponding to promoter regions
and/or transcriptionally active regions.

Figure 5. Chromatin state models of the turbot genome: **A)** 10-state model based on *in vivo* samples; **B)**
8-state model based on *in vitro* samples. Heatmaps of the emission parameters for each chromatin state
are shown for ChIP-Seq data with three histone marks and ATAC-Seq (left), the neighbourhood around the
TSS (middle) and the emission parameters of each chromatin state considered alongside features within
the turbot genome (right). TSS: Transcription start site; TES: Transcription end site.

Differential chromatin accessibility regions (DAR) and differential histone modification regions (DHMR) following stimulation

We next aimed to identify regulatory regions in the turbot genome affected by the immune stimulation. Significant DARs and DHMRs (FDR-adjusted p < 0.05) contrasting *Vibrio* and poly I:C stimulations to control samples were identified (Table 3, Supplementary table 9). Globally, a higher number of DARs or DHMRs were detected for up- than for down-regulated regions, more *in vivo* than *in vitro*, and more for *Vibrio* than poly I:C comparisons. Additionally, histone marks H3K27ac and H3K27me3 showed much lower DHMRs than H3K3me3 or ATAC-Seq DARs.

Table 3. Differential Accessibility Regions (DARs) and Differential Histone Modification Regions (DHMRs) for the *in vitro* and *in vivo* immune stimulations with viral and bacterial PAMPs

PAMP	Stimulation	Assay	Upregulated DAR/DHMR	Downregulated DAR/DHMR	All DAR/DHMR
Vibrio	*in vitro*	ATAC	17,878	2,739	20,617
		H3K4me3	10,755	699	11,454
		H3K27ac	623	2	625
		H3K27me3	1,211	6	1,217
	in vivo	ATAC	16,733	43,159	59,892
		H3K4me3	9,988	287	10,275
		H3K27ac	3	0	3
		H3K27me3	0	4	4
poly I:C	*in vitro*	ATAC	0	0	0
		H3K4me3	0	0	0
		H3K27ac	20	0	20
		H3K27me3	0	0	0
	in vivo	ATAC	15	25	40
		H3K4me3	1,036	0	1,036
		H3K27ac	38	0	38
		H3K27me3	0	126	126

Association of ATAC-Seq and ChIP-Seq data with RNA-Seq data

We tested if DARs and DHMRs for H3K4me3 and H3K27ac marks annotated as promoter/TSS regions (up to -1kb upstream of TSSs) corresponded to DEGs under the same experimental conditions (Hypergeometric distribution test, p < 0.05). DARs and DHMRs were much more overrepresented at the promoter regions of up- rather than down-regulated DEGs (Table 4, Supplementary table 10), suggesting changes in chromatin state associated with activation of genes. We performed GO enrichment analyses of DEGs overlapping with DARs and DHMRs for activation marks (H3K4me3 and H3K27ac) at promoter regions within each experimental condition through Hypergeometric distribution tests (P < 0.05, Table 4). Significant enrichment was detected for upregulated genes (p < 0.05), including several metabolic activities and particular immune functions, including antigen processing/presentation and apoptotic/cell death pathways (Supplementary table 11). Specifically, enriched terms were mostly associated with RNA processing (in particular tRNA, rRNA, ncRNA), ribosome biogenesis, and translation for the *Vibrio in vitro* stimulation, whereas for protein localization to nucleus and organelle, DNA replication and carbohydrate metabolism for the *Vibrio in vivo* stimulation. The term 'peptidyl-arginine modification' was enriched for the intersection of DAR/DHMRs and DEGs (*prmt1, 3, 5*

442 and *7*), as outlined before for the whole DEG analysis (RNA-Seq). The poly I:C *in vivo* stimulation
443 showed enrichment in immune-related processes, including 'antigen processing and
444 presentation via MHC-I' (explained by *erap1b, tapbpl*) and 'regulation of programmed cell death'
445 (explained by *apaf1, bida, casp8 and 10, socs3a, grinab, bcl2l10*; Supplementary table 11).
446

447 **Table 4.** Overlap of promoter DARs and DHMRs with DEG promoters (Hypergeometric test, p < 0.05).
448 Significant results are underlined. Refer to Supplementary table 10 for further details.

| Experimental condition | DARs/DHMRs at promoter regions | | Integration of DARs/DHMRs at promoter regions of DEGs | | | |
| | | | Down | | Up | |
	Down	Up	DEG	DAR/DHMR + DEG	DEG	DAR/DHMR + DEG
ATAC poly I:C *in vitro*	-	-	544	-	858	-
ATAC *Vibrio in vitro*	55	4922	3217	1	3321	1259
ATAC poly I:C *in vivo*	4	1	1918	0	2001	0
ATAC *Vibrio in vivo*	1350	6115	929	34	910	395
H3K4me3 poly I:C *in vitro*	-	-	544	0	858	0
H3K4me3 *Vibrio in vitro*	53	4289	3217	14	3321	1164
H3K4me3 poly I:C *in vivo*	-	368	1918	-	2001	151
H3K4me3 *Vibrio in vivo*	22	4024	929	3	910	242
H3K27ac poly I:C *in vitro*	-	0	544	-	858	0
H3K27ac *Vibrio in vitro*	0	72	3217	0	3321	49
H3K27ac poly I:C *in vivo*	-	4	1918	-	2001	3
H3K27ac *Vibrio in vivo*	-	1	929	-	910	1
Active prom poly I:C *in vitro*	-	-	544	-	858	-
Active prom *Vibrio in vitro*	106	6744	3217	15	3321	1782
Active prom poly I:C *in vivo*	4	370	1918	0	2001	151
Active prom *Vibrio in vivo*	1263	7082	929	37	910	463

prom: promoter; DAR: differential accessibility regions; DHMR: differential histone modification regions

Transcription factor binding motif (TFBM) analysis

To establish TFs potentially associated with chromatin state regulation following immune
stimulation, the enrichment of TFBMs within DAR/DHMRs annotated as promoter or enhancer
regions were examined. Significant enrichment was detected for poly I:C *in vivo* and the *Vibrio*
in vitro and *in* vivo stimulations. On average across simulations, 4 (range: 0-11) and 71 (range: 0-
132) enriched TFBMs were detected for down- and upregulated promoters, respectively, while
14 (range: 6-27) and 49 (range: 1-80) respective TFBMs were enriched for downregulated and
upregulated enhancers (Supplementary table 12).

Most TFBMs and associated TFs predicted for active promoters and enhancers were enriched for
several stimulations (Figure 6; Supplementary table 12): 68 TFs for both *Vibrio* stimulations and
in vivo poly I:C; 59 TFs for both *Vibrio* stimulations; and 20 TF for both *in vivo* stimulations.

To further investigate the functions of TFs with enriched motifs, we used the Molecular Complex
Detection algorithm (MCODE) and protein-protein interaction data (PPI) to identify connected
functions among differentially expressed TFs and among TFs with DARs or DHMRs on their
promoters (Table 5, Figure 7, Supplementary Figure 2).

Figure 6. Venn diagram illustrating overlap of enriched transcription factor binding motifs (TFBMs) in
upregulated promoter and enhancer regions in response to the different stimulants. Each coloured box
shows the TFBM found.

Table 5. Upregulated TFs with binding motif enrichment predicted on DARs and DHMRs (H3K4me3 and
H3K27ac) of active promoters and enhancer regions. The second column shows TF with promoters
overlapping with DAR/DHMR, while the third column shows TF that are differentially expressed genes
(DEGs). TF that are in both columns are underlined.
474

Stimulation		Transcription factor DAR/DHMR	Transcription factor DEG
poly I:C	*in vitro*	-	-
	in vivo	BATF, BHLHE40, PU.1, RUNX1, ZEB2	ELK4, ETV4, FRA1, IRF1, IRF8, MEF2C, PU.1, STAT3
Vibrio	*in vitro*	AP1, ATF1, BATF, BCL11A, <u>BHLHE40</u>, BHLHE41, BMYB, JUN, <u>MYCA</u>, EGR1, ELK1, EN1, FOS, FOSL2, FOXK1, FOXK2, <u>FOXO1</u>, FOXO3, FRA2, HIF1A, <u>HIF1B</u>, HINFP, <u>IRF3</u>, IRF4, <u>IRF8</u>, JUNB, JUND, KLF6, KLF9, MEIS1, <u>MITF</u>, MNT, MYB, NFkB-p65, PU.1 RUNX1, SMAD3, SP1, SP2, <u>STAT4</u>, USF1, USF2, ZEB2, ZNF341	ATF3, <u>BHLHE40</u>, BMYB, <u>MYCA</u>, ELF3, <u>FOXO1</u>, FOSL1A, FRA1, <u>HIF1B</u>, <u>IRF3</u>, <u>IRF8</u>, <u>MITF</u>, p63, <u>STAT4</u>
	in vivo	AP1, ATF1, ATF2, ATF4, ATF7, BATF, <u>BCL11A</u>, BHLHE49, BMYB, JUN, MYCA, CRE, <u>EGR1</u>, ELK1, EN1, ETV4, FOS, FOSL2, FOXK2, FOXO1, FOXO3, FRA2, HIF1B, HINFP, IRF3, IRF4, IRF8, JUNB, JUND, KLF6, KLF9, MAFB, MEF2A, MEF2C, <u>MEIS1</u>, MITF, MNT, MYB, PRDM1, PU.1, RUNX1, SMAD3, SP1, SP2, STAT3, STAT4, TCF12, USF1, USF2, <u>ZEB2</u>, ZNF341, ZNF711	BCL11A, CLOCK, <u>EGR1</u>, FRA1, MEF2C, <u>MEIS1</u>, <u>ZEB2</u>

DAR: differential accessibility regions; DHMR: differential histone modification regions

For the *Vibrio in vitro* stimulation, interconnected clusters of TFs associated with hemopoiesis
and immune functions, particularly myeloid leukocyte differentiation, FGF signalling pathway,
regulation of neutrophil differentiation, lymphocyte activation, apoptosis and the ERK1/2
immune signalling, were identified. For the *Vibrio in vivo* stimulation, we detected clusters of TFs
associated with similar functions (e.g. hemopoiesis, the ERK1/2 signalling pathway, myeloid cell
development and cell death), but also other functions such as insulin signalling, regeneration,
and oestrogen-dependent expression (Figure 7). Finally for the poly I:C *in vivo* stimulation,
myeloid leukocyte differentiation, regulation of cell differentiation, cell morphogenesis and
biosynthesis and growth showed connected TF subsets.

Using the Integrative Genomics Viewer (IGV), we visualized the variation in chromatin states
around the promoters of 12 TF-DEGs selected from Table 5 to show the variance in chromatin
state distribution across TFs and their regulatory elements following activation by the different
stimulants (Figure 8; Supplementary Figure 3).

489
490
Figure 7. Functional interconnection of active TFs detected in response to the different stimulants: **A)**
Ontology enrichment clusters of upregulated TFs showing promoter DARs/DHMRs with TFBMs enriched in
promoter or enhancer DARs/DHMRs of DEGs detected in *Vibrio in vivo* stimulation. **B)** Protein-protein
interaction MCODE network of upregulated TFs induced by *Vibrio in vivo* stimulation. Each cluster is
coloured with the most statistically significant term among terms that cluster together. The size of each
term is given by -log10 p, and the stronger the similarity among nearby terms, the thicker the edges
between them.

Figure 8. Chromatin state predictions and RNA/ATAC/ChIP-Seq tracks in the transcription factor coding gene *foxo1a*, which was upregulated and showed DAR around the promoter for *Vibrio in vitro* response. Transcription factor binding motifs for FOXO1A were also enriched in the promoter DARs of DEGs upregulated in response to *Vibrio in vitro*.

DISCUSSION

This study represents the first epigenomic analysis of the turbot head kidney, the primary hematopoietic and lymphoid organ of teleosts, in response to two mimics of viral (poly I:C) and bacterial (inactivated *V. anguillarum*) stimulation. While head kidney transcriptomic responses to bacteria, virus and parasites have been extensively investigated in turbot (Pardo et al., 2008; Millán et al., 2010; Domínguez et al., 2013; Díaz-Rosales et al., 2012; Robledo et al., 2014; Ronza et al., 2020; Valle et al., 2020; Libran-Pérez et al., 2022), the epigenetic regulation of chromatin states following immune stimulation has not been explored before. The head kidney response was explored through intraperitoneal injection, reflecting the response in the whole-body, including interactions among immune organs, and through *in vitro* stimulation of primary leukocyte cultures, reflecting their direct interaction with stimulants.

Transcriptomic response to immune stimulation

Upregulated genes with key immune roles were found in both *in vitro* stimulations, but also in the poly I:C *in vivo* stimulation. Poly I:C interacts with TLR3 (Toll-like receptor 3) and induces IFN production (Kumar et al., 2006). *tlr3* was upregulated in the *Vibrio in vivo* stimulation and not differentially expressed in poly I:C *in vivo*, but it was downregulated in both *in vitro* stimulations. A possible explanation is that *in vitro* downregulation could reflect signal exhaustion, suggesting that 24 hours postinjection is too long a period to capture the direct interaction between leukocytes and stimulants, and in fact, a shorter period was applied in other species with a consistent response (Herrera-Uribe et al., 2020). However, we cannot discard alternative hypotheses related to changes or diversification of the TLR3 signalling pathway in flatfish or specifically in turbot, as suggested by its activation in response to *Vibrio in vivo* stimulation and to parasites, such as *P. dicentrarchi* (Figueras et al., 2016). In fact, we identified other interferon-stimulated DEGs in poly I:C both *in vitro* and *in vivo*, including *socs1a* and b (suppressor of cytokine signalling 1), *nod2* (nucleotide binding oligomerization domain containing 2) and *nmi* (N-Myc interactor), typically participating in type I IFN responses (Blumer et al., 2017; Ahn et al., 2021). *socs1* is an inducible negative regulator of IFN (Blumer et al., 2017), a mechanism to maintain tissue homeostasis following the IFN response; *nod2* encodes for an intracellular receptor triggering innate antibacterial and antiviral signalling pathways, including IFN (Nie et

al., 2017); and *nmi* increases STAT-mediated transcription in response to IFN-gamma (Zhu et al.,
1999). The activation of genes involved in positive and negative regulation of IFN- signalling
suggests a fine adjustment to an intense IFN response to recover tissue homeostasis (Ivashkiv
and Donlin, 2014; Murira and Lamarre, 2016). Also, the upregulation of these four key genes
after *Vibrio in vitro* stimulation is consistent with past reports that place IFN regulation as a key
point of the immune response (Kopitar-Jerala, 2017).

Upregulated DEGs shared by both *in vivo* stimulations were involved in the activation of
transcription and translation and protein modification/localization. This included four members
of the immune-related protein arginine methyltransferase family (*prmt1*, *prmt3*, *prmt5* and
prmt7), enriched under the term "peptidyl-arginine modification". *Prmt* family members
regulate transcription and translation and are also involved in signal transduction during
inflammation and response against poly I:C and bacterial lipopolysaccharide (LPS; Srour, 2022).
These four genes were also shared between both *Vibrio* comparisons, suggesting an important
role in the early response to *Vibrio*.

We also checked if immune genes could be regulated in opposite directions after stimulation
with viral and bacterial mimics, which could reflect antagonistic immune responses. Increased
susceptibility to bacterial superinfections induced by innate antiviral responses has been
reported in several models in mammals (Barman and Metzger, 2021) in type I and type II IFN
responses (Navarini et al., 2006; Sun and Metzger, 2008). Such mechanisms are likely present in
teleosts due to the high conservation of these pathways, and in fact, different patterns of
resistance to viral and bacterial infections were observed among isogenic rainbow trout lines
(Fraslin et al., 2020). However, no negative correlations for resistance to bacteria and viruses has
been observed in the few studies addressing this issue in relation to selective breeding (Ødegård
et al., 2007; 2011; Bangera et al., 2011). Indeed, understanding the cellular basis of antagonistic
responses to viruses and bacteria in fish will require more research, but here we identified
examples of opposite gene expression responses, including genes from the core type I IFN
response, conserved between teleosts and humans (Levraud et al., 2019; Clark et al., 2023), such
as *sting1* (stimulator of interferon response CGAMP interactor 1, Zhang et al., 2021*), irf1b*
(interferon regulatory factor 1; Sullivan et al., 2021) and *nod2* (mentioned above),
downregulated by *Vibrio*s and upregulated by poly I:C *in vivo*. Reciprocally, the critical pro-
inflammatory cytokine *il1b* (interleukin 1 beta, Sahoo et al., 2011) and the regulator of
inflammation *traf4a* (TNF receptor associated factor 4, Marinis et al., 2012) were upregulated by
Vibrio but downregulated by poly I: C. Interestingly, a majority of genes induced by poly I:C and
repressed by *Vibrio* extracts had human/mouse ISG orthologs (22/42 *in vitro*, 18/34 *in vivo*); in
contrast, many genes induced by *Vibrio* extracts and repressed by poly I:C, especially *in vivo,* had
human/mouse orthologs downregulated by type I IFN. Overall, these observations indicate that
a significant part of these contrasted responses is mediated by genes functionally conserved
between fish and mammals.

Epigenetic assays and their association with transcriptomic response

Activation and recruitment of transcription factor (TF) subsets is at the top of the specific
transcriptome cascade response (Alvarez et al., 2020; Weidemüller et al., 2021). In our study, the
exploration of promoters of differentially expressed TFs, also involving differential accessibility
regions (DARs) and differential histone modification regions (DHMRs), barely showed changes in
the chromatin state distribution between treatments (Supplementary Figure 3). This is the case
for instance of egr*1* (early growth response 1), which plays a key role in cell survival, proliferation
and cell death, and regulates the expression of *il1b* and *cxcl2* (CXC motif chemokine ligand 2).

The same was observed for *meis1* (myeloid ecotropic viral integration site homeobox 1), related
to bone marrow hematopoiesis in mammals, and *zeb2* (zinc finger e-box binding homeobox 2),
associated with macrophage infiltration (Taylor et al., 2022). In all these examples, the TF genes
showed bivalent/poised promoters in the three *in vivo* samples (control, *Vibrio* and poly I:C). A
similar situation was found *in vitro*, where promoters of genes, such as *irf3* (interferon regulatory
factor 3) and *mitf* (melanocyte inducing transcription factor), showed activation signals
regardless of the experimental condition. In previous studies, these bivalent/poised states have
been interpreted as a mechanism allowing rapid responses, e.g. for genes induced during the
pro-inflammatory response (Herrera-Uribe et al., 2020; Martínez de Paz and Josefowicz, 2021)
or during mammal embryonic development and in germ cells (Lesch and Page, 2014; Barbieri et
al., 2017). Thus, the expression of primary response genes, such as the TF genes exemplified
above, may be allowed by permissive chromatin states.

Regardless of predicted chromatin states around gene promoters, we observed an increased
enrichment of these immune-related genes following poly I:C and *Vibrio* stimulation that would
suggest further chromatin unpacking (Table 3), even if the chromatin state is not strongly
affected. In fact, many differential regions were detected between stimulations, particularly
DARs (i.e. ATAC-Seq) and DHMRs (for H3K4me3) for *Vibrio*. In comparison, fewer DARs and
DHMRs were detected for the poly I:C stimulations. We interpret this absence of differential
regions as a return, both *in vivo* and *in vitro*, to the native chromatin state after 24 hpi, which
may suggest the initial response has already exhausted. Poly I:C is a synthetic double-stranded
RNA that acts as a PAMP binding to a specific PRR (TLR3). Meanwhile, inactive Gram-negative
bacteria such as *Vibrio* sp. harbour a variety of PAMPs, such as LPS, porins, peptidoglycan,
mannose-rich sugars and flagellin, detected by a plethora of PRRs, such as TLRs, NOD-like and C-
type lectin receptors, and PGRPs (Boltaña et al., 2011; Li and Wu, 2021). Thus, we explain our
findings as the effect of a shorter but likely intense innate response against poly I:C and a more
sustained response to *Vibrio*. In a previous study on pig alveolar macrophages stimulated with
poly I:C and LPS, Herrera-Uribe et al. (2020) noted a decrease in the number of H3K27ac-DHMRs
from 2 to 6 hpi, which could explain our observation in turbot (24 hpi).

The presence of differential activation signals at promoter regions, either in the form of open
chromatin regions or histone marks such as H3K4me3 and H3K27ac, are typically associated with
differential gene expression (Stępniak et al., 2021). H3K27ac has been widely catalogued as one
of the most dynamic activation marks in eukaryotes, (Saeed et al., 2014). Surprisingly, we did not
find important changes between experimental conditions in our study, in which H3K4me3 was
the most dynamic among the ChIP-Seq marks. We suspect that the decrease in H3K27ac and
increase in H3K4me3 could also be an effect of our choice to sample 24 hours post-injection,
especially for the *in vivo* stimulations. Previous studies have reported that H3K4me3 may not
have an active role in activating transcription, but instead reflects transcriptional changes itself,
suggesting that H3K4me3 modification may have role on transcriptional consistency or memory
of previous states (Howe et al., 2017).

DEG promoters overlapped with DAR/DHMRs for upregulated genes in most conditions (Table
4). GO enrichment of DEGs with activation signals at promoters (Supplementary table 11) was
similar to that observed with the whole DEG dataset (Supplementary table 5), with both *Vibrio*
stimulations enriched in terms associated with transcription activation, while poly I:C *in vivo*
showed a more specific response, enriched in immune-related processes (Supplementary table
6). However, DEGs of some key immune functions, such as response to cytokine stimulus, did
not show DAR/DHMRs, suggesting that these genes could be already accessible before

stimulation. Cytokines are one of the core initiators of the inflammatory response (Secombes et
al., 2001), thus a bivalent state, ready for activation or inactivation of expression, might explain
this observation (Figure 5).

Predicted TFBMs enriched within DARs/DHMRs in the promoters and enhancer-state regions
predicted by ChromHMM of upregulated DEGs, provided a more detailed picture of regulatory
elements changed by the immune stimulations (Figure 6, Supplementary table 12). Most of the
enriched TFBMs were shared between at least two treatments, which supports their general
regulatory function. Among those shared between most stimulations (excluding poly I:C *in vitro*),
we found several TFBMs of the ETS-family, which are involved in cell proliferation, apoptosis and
lymphocyte development (Watson et al., 2002). We also found TFBMs enrichment of PU.1, a
master TF for the myeloid lineage, that promotes chromatin accessibility, also identified in a
similar study in pig (Turkistany and DeKoter, 2011; Chen et al., 2020; Herrera-Uribe et al., 2020).
PU.1 is thought to promote binding of other TFs that were enriched in our study, including AP1.
This protein is activated by TLRs during the immune response (Shan et al., 2018) and regulates
gene expression in response to cytokines, stress, and bacterial and viral infections (Kim et al.,
2014), and notably the IFN regulatory factors (IRFs, particularly IRF3, IRF4 and IRF8). The IRFs are
key regulators of innate antiviral and antibacterial responses in vertebrates including fish (Clark
et al., 2021; Han et al., 2023). The observed ontology enrichment clustering of these TFs
reinforces the similarities between the conditions in our study (Figure 7, Supplementary Figure
2).

To illustrate the functionality of our turbot epigenomic atlas, we explored the chromatin state
changes in a selection of 12 TF genes (Figure 8, Supplementary Figure 4). These genes were
differentially expressed and showed DARs/DHMRs in their promoters. Immune-responsive
chromatin state remodelling was found, for instance, for *irf8* (interferon regulatory factor 8) in
the *Vibrio in vitro* stimulation. Here, an extension of the "strongly active promoter" state around
the TSS region was visible compared to the non-stimulated and poly I:C stimulated samples,
alongside an overall expansion of the "strongly active transcript" and "ATAC island" states
downstream. IRF8 is a key regulator of the NF-κB signalling pathway during inflammation, along
with IRF3 (Yan et al., 2020b). Both *irf* genes were differentially expressed in poly I:C and *Vibrio in*
vitro stimulations (*irf8* also in poly I:C *in vivo*) and showed promoter DARs in *Vibrio* stimulations.
Interestingly, chromatin state changes were barely observed in the vicinity of the *irf3* promoter
(Supplementary Figure 4).

Another example was found in *Vibrio* and poly I:C *in vivo* stimulations for *bcl11a* (BCL11
transcription factor A), a key regulator of dendritic cell differentiation (Ippolito et al., 2014) and
negative regulator of p53 (Yu et al., 2012). This gene was differentially expressed in both
stimulations, but only showed a DAR promoter in response to *Vibrio* stimulation. The
promoter/TSS region was annotated by ChromHMM as "weak repressed Polycomb" in the
control, which changed in both the *Vibrio* and poly I:C stimulations, extending to
"bivalent/poised" state, and in the case of *Vibrio* showing an extension to "Strongly active
promoter/transcript" state. Weak and strong signals of active enhancers were also detected in
the poly I:C and *Vibrio* stimulations within the second intron of the *bcl11a*, that were not present
in the controls (Supplementary Figure 4). A similar situation was found *in vitro* for *foxo1a*
(Forkhead box O1), a TF-coding gene that participates in mucosal (innate) immune response
regulating the expression of antimicrobial peptides and promoting phagocytosis during bacterial
and parasitic infections (Cabrera-Ortega et al., 2017; Graves and Milovanova, 2019). This gene
was differentially expressed in *Vibrio in vitro* and showed an extension of the "Strongly active

promoter" state compared to the control and specially the poly I:C stimulation, while conserving
the "bivalent/poised TSS" stretches downstream of the promoter. This was followed by many
"ATAC islands" along the gene that were also present, although more scarcely, in the control, and
almost absent in poly I:C. Finally, a "medium enhancer" state was detected within the first intron,
present only in the *Vibrio* condition.

Conclusions and perspectives

In summary, our study provides the first atlas of regulatory elements in turbot head kidney and
leukocytes during the early response to viral and bacterial stimulation, as a contribution to the
AQUA-FAANG project within the umbrella of the FAANG initiative. The integration of ATAC-Seq
and ChIP-Seq data suggests that changes in chromatin state distribution were not as frequent
between stimulations and controls as expected. However, the presence of DARs and DHMRs
between the stimulations and the controls, which broadly overlapped with DEGs, provides clear
evidence for gene regulation at the epigenetic level that underpins changes in gene expression
driving immune functions. Overall, this epigenomic atlas will help to decode the molecular
mechanisms underlying turbot immune responses to viral and bacterial stimuli and offers a novel
resource for developing selective breeding strategies for controlling diseases, one of the main
concerns of turbot industry. Future work will benefit from linking the regulatory annotations
generated in this study with genetic variants defined by whole genome re-sequencing (Johnston
et al., 2024), to help prioritize causal genetic variants for disease resistance traits underpinned
by gene expression responses to pathogens.

DATA AVAILABILITY STATEMENT

All raw RNA-Seq, ATAC-Seq and ChIP-Seq datasets can be accessed through the ENA repository
under accession numbers PRJEB47933, PRJEB47934 and PRJEB57784, respectively.

AUTHOR CONTRIBUTIONS

OA: Methodology, Software, Formal analysis, Investigation, Data Curation, Visualization, Writing
– original draft, Writing – review & editing; **BGP:** Methodology, Investigation, Resources, Writing
– review & editing; **PRV:** Investigation, Resources, Writing – review & editing; **ABH:** Software,
Formal Analysis, Data Curation, Visualization, Writing – review & editing; **JL**: Investigation,
Resources, Writing – review & editing; **PD:** Software, Formal Analysis, Writing – review &
editing; **DPM:** Software, Formal Analysis, Writing – review & editing; **PB**: Resources, Writing –
review & editing; **DM**: Funding Acquisition, Methodology, Supervision, Resources, Writing –
review & editing; **CB**: Conceptualization, Formal Analysis, Resources, Writing – review & editing,
Project Administration, Supervision; **PM**: Conceptualization, Methodology, Formal Analysis,
Resources, Writing – review & editing, Project Administration, Supervision.

FUNDING

This study was funded by the AQUA-FAANG project, which received funding from the European
Union's Horizon 2020 research and innovation programme under grant agreement No 817923.
Additional funding was provided by Xunta de Galicia local government (Spain) (ED431C 2022/33),
which also supported the research fellowships of OA and PRV (refs. ED481A-2020/119 and
ED481A-2020/491430, respectively). Contributions from the Roslin Institute were further
supported by the BBSRC Institutional Strategic Programme grants BBS/E/D/10002070,
BBS/E/D/20002174, BBS/E/RL/230001B and BBS/E/RL/230002B.

ACKNOWLEDGMENTS

We acknowledge the technical support and informatic resources provided by the Centro de Supercomputación de Galicia (CESGA). We acknowledge Professor Carolina Tafalla (INIA-CSIC) for the support given in the revision of this article.

REFERENCES

Ahn JH, Park JY, Kim DY, Lee TS, Jung DH, Kim YJ, Lee YJ, Lee YJ, Seo IS, Song EJ, Jang AR, Yang SJ, Shin SJ, Park JH (2021). **Type I Interferons Are Involved in the Intracellular Growth Control of** *Mycobacterium abscessus* **by Mediating NOD2-Induced Production of Nitric Oxide in Macrophages**. Frontiers in Immunology, 12. doi: 10.3389/fimmu.2021.738070.

Alvarez JM, Schinke AL, Brooks MD, Pasquino A, Leonelli L, Varala K, Safi A, Krouk G, Krapp A, Coruzzi GM (2020). **Transient genome-wide interactions of the master transcription factor NLP7 initiate a rapid nitrogen-response cascade**. *Nature Communications*, *11*(1), 1157. doi: 10.1038/s41467-020-14979-6.

Amemiya HM, Kundaje A, Boyle AP (2019). **The ENCODE Blacklist: Identification of Problematic Regions of the Genome**. *Sci Rep* 9, 9354. doi: 10.1038/s41598-019-45839-z.

Andersson L, Archibald AL, Bottema CD, Brauning R, Burgess SC, Burt DW, Casas E, Cheng HH, Clarke L, Couldrey C, Dalrymple BP, Elsik CG, Foissac S, Giuffra E, Groenen MA, Hayes BJ, Huang LS, Khatib H, Kijas JW, … Zhou H (2015). **Coordinated international action to accelerate genome-to-phenome with FAANG, the Functional Annotation of Animal Genomes project**. *Genome Biology*, *16*(1), 57. doi: 10.1186/s13059-015-0622-4.

Andrews S (2010). **FastQC: A Quality Control Tool for High Throughput Sequence Data.** Available online at: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

APROMAR (2022). **La acuicultura en España 2022**. *Asociación Empresarial de Acuicultura de España, Cádiz* (Online, accessed October 17, 2023 at www.apromar.es)

Aramburu O, Blanco A, Bouza C, Martínez P (2023). **Integration of host-pathogen functional genomics data into the chromosome-level genome assembly of turbot (***Scophthalmus maximus***)**. *Aquaculture*, 564, 739067. doi: 10.1016/j.aquaculture.2022.739067.

Bangera R, Ødegård J, Præbel AK, Mortensen A, Nielsen HM (2011). **Genetic correlations between growth rate and resistance to vibriosis and viral nervous necrosis in Atlantic cod (***Gadus morhua* **L.)**. *Aquaculture*, 317(1–4), 67–73. doi: 10.1016/j.aquaculture.2011.04.018.

Baranasic D, Hörtenhuber M, Balwierz PJ, Zehnder T, Mukarram AK, Nepal C, Várnai C, Hadzhiev Y, Jimenez-Gonzalez A, Li N, Wragg J, D'Orazio FM, Relic D, Pachkov M, Díaz N, Hernández-Rodríguez B, Chen Z, Stoiber M, Dong M, Stevens I, Ross SE, Eagle A, Martin R, Obasaju O, … Müller F (2022). **Multiomic atlas with functional stratification and developmental dynamics of zebrafish cis-regulatory elements**. *Nat Genet* 54, 1037–1050 (2022). doi: 10.1038/s41588-022-01089-w.

Barbieri M, Xie SQ, Torlai-Triglia E, Chiariello AM, Bianco S, de Santiago I, Branco MR, Rueda D, Nicodemi M, Pombo A (2017). **Active and poised promoter states drive folding of the extended HoxB locus in mouse embryonic stem cells**. *Nature Structural & Molecular Biology*, *24*(6), 515–524. doi: 10.1038/nsmb.3402

Barman TK, Metzger DW (2021). **Disease Tolerance during Viral-Bacterial Co-Infections.** *Viruses*, *13*(12), 2362. doi: 10.3390/v13122362.

Barnett DW, Garrison EK, Quinlan AR, Strömberg MP, Marth GT (2011). **BamTools: a C++ API**
and toolkit for analyzing and managing BAM files. *Bioinformatics*, *27*(12), 1691–1692. doi:
10.1093/bioinformatics/btr174.

Blumer T, Coto-Llerena M, Duong FHT, Heim MH (2017). **SOCS1 is an inducible negative**
regulator of interferon λ (IFN-λ)–induced gene expression in vivo. *Journal of Biological*
Chemistry, *292*(43), 17928–17938. doi: 10.1074/jbc.M117.788877.

Boltaña S, Roher N, Goetz FW, MacKenzie SA (2011). **PAMPs, PRRs and the genomics of gram-**
negative bacterial recognition in fish. *Developmental & Comparative Immunology*, *35*(12),
1195–1203. doi: 10.1016/j.dci.2011.02.010.

Boltsis I, Grosveld F, Giraud G, Kolovos P (2021). **Chromatin Conformation in Development and**
Disease. *Frontiers in Cell and Developmental Biology*, 9. doi: 10.3389/fcell.2021.723859.

Buenrostro JD, Wu B, Chang HY, Greenleaf WJ (2015). **ATAC-seq: A Method for Assaying**
Chromatin Accessibility Genome-Wide. *Current Protocols in Molecular Biology*, 109(1). doi:
10.1002/0471142727.mb2129s109.

Cabrera-Ortega AA, Feinberg D, Liang Y, Rossa C, Graves DT (2017). **The Role of Forkhead Box 1**
(FOXO1) in the Immune System: Dendritic Cells, T Cells, B Cells, and Hematopoietic Stem Cells.
Critical Reviews in Immunology, *37*(1), 1–13. doi: 10.1615/CritRevImmunol.2017019636.

Chen S, Yang J, Wei Y, Wei X (2020). **Epigenetic regulation of macrophages: from homeostasis**
maintenance to host defense. *Cellular & Molecular Immunology*, *17*(1), 36–49. doi:
10.1038/s41423-019-0315-0.

Chen W, Huang J, Wang W, Wang Y, Chen H, Wang Q, Zhang Y, Liu Q, Yang D (2022). **Multi-tissue**
scRNA-seq reveals immune cell landscape of turbot (*Scophthalmus maximus*). *Fundamental*
Research, *2*(4), 550–561. doi: 10.1016/j.fmre.2021.12.015.

Clark EL, Archibald AL, Daetwyler HD, Groenen MAM, Harrison PW, Houston RD, Kühn C, Lien S,
Macqueen DJ, Reecy JM, Robledo D, Watson M, Tuggle CK, Giuffra E (2020). **From FAANG to**
fork: application of highly annotated genomes to improve farmed animal production. *Genome*
Biology, *21*(1), 285. doi: 10.1186/s13059-020-02197-8.

Clark TC, Boudinot P, Collet B (2021). **Evolution of the IRF Family in Salmonids**. *Genes*, *12*(2),
238. doi: 10.3390/genes12020238.

Clark TC, Naseer S, Gundappa MK, Laurent A, Perquis A, Collet B, Macqueen DJ, Martin SAM,
Boudinot P (2023). **Conserved and divergent arms of the antiviral response in the duplicated**
genomes of salmonid fishes. *Genomics*, *115*(4), 110663. doi: 10.1016/j.ygeno.2023.110663.

Corces MR, Trevino AE, Hamilton EG, Greenside PG, Sinnott-Armstrong NA, Vesuna S, Satpathy
AT, Rubin AJ, Montine KS, Wu B, Kathiria A, Cho SW, Mumbach MR, Carter AC, Kasowski M,
Orloff LA, Risca VI, Kundaje A, Khavari PA, … Chang HY (2017). **An improved ATAC-seq protocol**
reduces background and enables interrogation of frozen tissues. *Nature Methods*, *14*(10), 959–
962. doi: 10.1038/nmeth.4396.

Cuvier O, Fierz B (2017). **Dynamic chromatin technologies: from individual molecules to**
epigenomic regulation in cells. *Nature Reviews Genetics*, 18(8), 457–472. doi:
10.1038/nrg.2017.28.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,
McCarthy SA, Davies RM, Li H (2021). **Twelve years of SAMtools and BCFtools**. *GigaScience*,
10(2). doi: 10.1093/gigascience/giab008

Díaz-Rosales P, Romero A, Balseiro P, Dios S, Novoa B, Figueras A (2012). **Microarray-Based**
Identification of Differentially Expressed Genes in Families of Turbot (*Scophthalmus maximus*)
After Infection with Viral Haemorrhagic Septicaemia Virus (VHSV). *Marine Biotechnology*,
14(5), 515–529. doi:10.1007/s10126-012-9465-0.

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR
(2013). **STAR: ultrafast universal RNA-seq aligner**. *Bioinformatics*, *29*(1), 15–21. doi:
10.1093/bioinformatics/bts635.

Domínguez B, Pardo BG, Noia M, Millán A, Gómez-Tato A, Martínez P, Leiro J, Lamas J (2013).
Microarray analysis of the inflammatory and immune responses in head kidney turbot
leucocytes treated with resveratrol. *International Immunopharmacology*, *15*(3), 588–596. doi:
10.1016/j.intimp.2013.01.024.

Ecker S, Pancaldi V, Valencia A, Beck S, Paul DS (2018). **Epigenetic and Transcriptional Variability**
Shape Phenotypic Plasticity. *BioEssays*, *40*(2), 1700148. doi: 10.1002/bies.201700148.

Ernst J, Kellis M (2012). **ChromHMM: automating chromatin-state discovery and**
characterization. *Nature Methods*, *9*(3), 215–216. doi: 10.1038/nmeth.1906.

Ewels P, Peltzer A, Fillinger S, Patel H, Alneberg J, Wilm A, Ulysse-Garcia M, Di Tommaso P,
Nahnsen S (2023). **The nf-core framework for community-curated bioinformatics pipelines**.
Zenodo. doi: 10.5281/zenodo.3240506.

Fang L, Liu S, Liu M, Kang X, Lin S, Li B, Connor EE, Baldwin RL, Tenesa A, Ma L, Liu GE, Li C (2019).
Functional annotation of the cattle genome through systematic discovery and characterization
of chromatin states and butyrate-induced variations. *BMC Biology*, 17(1), 68. doi:
10.1186/s12915-019-0687-8.

Feinberg AP (2007). **Phenotypic plasticity and the epigenetics of human disease**. *Nature*,
447(7143), 433–440. doi: 10.1038/nature05919.

Figueras A, Robledo D, Corvelo A, Hermida M, Pereiro P, Rubiolo JA, Gómez-Garrido J, Carreté L,
Bello X, Gut M, Gut IG, Marcet-Houben M, Forn-Cuní G, Galán B, García JL, Abal-Fabeiro JL, Pardo
BG, Taboada X, Fernández C, … Martínez P (2016). **Whole genome sequencing of turbot**
(*Scophthalmus maximus*; *Pleuronectiformes*): a fish adapted to demersal life. *DNA Research*,
23(3), 181–192. doi: 10.1093/dnares/dsw007.

Foissac S, Djebali S, Munyard K, Vialaneix N, Rau A, Muret K, Esquerré D, Zytnicki M, Derrien T,
Bardou P, Blanc F, Cabau C, Crisci,E, Dhorne-Pollet S, Drouet F, Faraut T, Gonzalez I, Goubil A,
Lacroix-Lamandé S, … Giuffra E (2019). **Multi-species annotation of transcriptome and**
chromatin structure in domesticated animals. *BMC Biology*, 17(1), 108. doi: 10.1186/s12915-
019-0726-5.

Frankish A, Carbonell-Sala S, Diekhans M, Jungreis I, Loveland JE, Mudge JM, Sisu C, Wright JC,
Arnan C, Barnes I, Banerjee A, Bennett R, Berry A, Bignell A, Boix C, Calvet F, Cerdán-Vélez D,
Cunningham F, Davidson C, … Flicek P (2023). **GENCODE: reference annotation for the human**

and mouse genomes in 2023. *Nucleic Acids Research*, *51*(D1), D942–D949. doi:
10.1093/nar/gkac1071

Fraslin C, Quillet E, Rochat T, Dechamp N, Bernardet JF, Collet B, Lallias D, Boudinot P (2020).
Combining Multiple Approaches and Models to Dissect the Genetic Architecture of Resistance
to Infections in Fish. *Frontiers in Genetics*, 11. doi: 10.3389/fgene.2020.00677.

Gao Y, Wang Q, Liu Y, Ma Y, Jin H, Liu J, Wang H, Yan Y, Li J (2023). **Epidemiology of turbot**
bacterial diseases in China between October 2016 and December 2019. *Frontiers in Marine*
Science, *10*. doi: 10.3389/fmars.2023.1145083.

Ge SX, Jung D, Yao R (2020). **ShinyGO: a graphical gene-set enrichment tool for animals and**
plants. *Bioinformatics*, *36*(8), 2628–2629. doi: 10.1093/bioinformatics/btz931.

Graves DT, Milovanova TN (2019). **Mucosal Immunity and the FOXO1 Transcription Factors**.
Frontiers in Immunology, *10*. doi: 10.3389/fimmu.2019.02530.

Guerrero-Peña L, Suarez-Bregua P, Gil-Gálvez A, Naranjo S, Méndez-Martínez L, Tur R, García-
Fernández P, Tena JJ, Rotllant J (2023). **Genome-wide chromatin accessibility and gene**
expression profiling during flatfish metamorphosis. *Scientific Data*, *10*(1), 196. doi:
10.1038/s41597-023-02111-4.

Geven EJW, Klaren PHM (2017). **The teleost head kidney: Integrating thyroid and immune**
signalling. *Developmental & Comparative Immunology*, 66, 73–83. doi:
10.1016/j.dci.2016.06.025.

Giral H, Landmesser U, Kratzer A (2018). **Into the Wild: GWAS Exploration of Non-coding RNAs**.
Frontiers in Cardiovascular Medicine, 5. doi: 10.3389/fcvm.2018.00181.

Giuffra E, Tuggle CK (2019). **Functional Annotation of Animal Genomes (FAANG): Current**
Achievements and Roadmap. *Annual Review of Animal Biosciences*, *7*(1), 65–88. doi:
10.1146/annurev-animal-020518-114913.

Halstead MM, Ma X, Zhou C, Schultz RM, Ross PJ (2020). **Chromatin remodelling in bovine**
embryos indicates species-specific regulation of genome activation. *Nature Communications*,
11(1), 4654. doi: 10.1038/s41467-020-18508-3.

Halstead MM, Islas-Trejo A, Goszczynski DE, Medrano JF, Zhou H, Ross PJ (2021). **Large-Scale**
Multiplexing Permits Full-Length Transcriptome Annotation of 32 Bovine Tissues from a Single
Nanopore Flow Cell. *Frontiers in Genetics*, 12. doi: 10.3389/fgene.2021.664260.

Han C, Huang W, Peng S, Zhou J, Zhan H, Li W, Gong J, Li Q (2023). **Characterization and**
expression analysis of the interferon regulatory factor (IRF) gene family in zig-zag eel
(*Mastacembelus armatus*) against *Aeromonas veronii* infection. *Developmental & Comparative*
Immunology, *140*, 104622. doi: 10.1016/j.dci.2022.104622.

He P, Williams BA, Trout D, Marinov GK, Amrhein H, Berghella L, Goh ST, Plajzer-Frick I, Afzal V,
Pennacchio LA, Dickel DE, Visel A, Ren B, Hardison RC, Zhang Y, Wold BJ (2020). **The changing**
mouse embryo transcriptome at whole tissue and single-cell resolution. *Nature*, *583*(7818),
760–767. doi: 10.1038/s41586-020-2536-x.

Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, Cheng JX, Murre C, Singh H, Glass CK
(2010). **Simple Combinations of Lineage-Determining Transcription Factors Prime cis-**

Regulatory Elements Required for Macrophage and B Cell Identities. *Molecular Cell*, *38*(4), 576–589. doi: 10.1016/j.molcel.2010.05.004.

Herrera-Uribe J, Liu H, Byrne KA, Bond ZF, Loving CL, Tuggle CK (2020). **Changes in H3K27ac at Gene Regulatory Regions in Porcine Alveolar Macrophages Following LPS or PolyIC Exposure**. Frontiers in Genetics, 11. doi: 10.3389/fgene.2020.00817.

Howe FS, Fischl H, Murray SC, Mellor J (2017). **Is H3K4me3 instructive for transcription activation?** *BioEssays*, *39*(1), e201600095. doi: 10.1002/bies.201600095.

Hu J, Barrett RDH (2017). **Epigenetics in natural animal populations**. *Journal of Evolutionary Biology*, *30*(9), 1612–1632. doi: 10.1111/jeb.13130.

Ippolito GC, Dekker JD, Wang YH, Lee BK, Shaffer AL, Lin J, Wall JK, Lee BS, Staudt LM, Liu YJ, Iyer VR, Tucker HO (2014). **Dendritic cell fate is determined by BCL11A**. *Proceedings of the National Academy of Sciences*, *111*(11). doi: 10.1073/pnas.1319228111.

Ivashkiv LB, Donlin LT (2014). **Regulation of type I interferon responses**. *Nature Reviews Immunology*, 14(1), 36–49. doi:10.1038/nri3581.

Johnston IA, Kent MP, Boudinot P, Looseley M, Bargelloni L, Faggion S, Merino GA, Ilsley GR, Bobe J, Tsigenopoulos CS, Robertson J, Harrison PW, Martinez P, Robledo D, Macqueen DJ, Lien S (2024). **Advancing fish breeding in aquaculture through genome functional annotation**. *Aquaculture*, *583*, 740589. doi: 10.1016/j.aquaculture.2024.740589

Karimzadeh M, Ernst C, Kundaje A, Hoffman MM (2018). **Umap and Bismap: quantifying genome and methylome mappability**. *Nucleic Acids Research*. doi: 10.1093/nar/gky677.

Kellis M, Wold B, Snyder MP, Bernstein BE, Kundaje A, Marinov GK, Ward LD, Birney E, Crawford GE, Dekker J, Dunham I, Elnitski LL, Farnham PJ, Feingold EA, Gerstein M, Giddings MC, Gilbert D. M, Gingeras TR, Green ED, … Hardison RC (2014). **Defining functional DNA elements in the human genome**. *Proceedings of the National Academy of Sciences*, 111(17), 6131–6138. doi: 10.1073/pnas.1318948111.

Kim J, Lee KW, Lee HJ (2014). **Polyphenols Suppress and Modulate Inflammation**. In *Polyphenols in Human Health and Disease* (pp. 393–408). Elsevier. doi 10.1016/B978-0-12-398456-2.00029-3.

Klosterhoff MC, Pereira-Junio J, Rodrigues RV, Gusmão EP, Sampaio LA, Tesser MB, Romano L A (2015). **Ontogenic development of kidney, thymus and spleen and phenotypic expression of CD3 and CD4 receptors on the lymphocytes of cobia (*Rachycentroncanadum*)**. *Anais Da Academia Brasileira de Ciências*, 87(4), 2111–2121. doi: 10.1590/0001-3765201520140623.

Kopitar-Jerala N (2017). **The Role of Interferons in Inflammation and Inflammasome Activation**. *Frontiers in Immunology*, *8*. doi: 10.3389/fimmu.2017.00873.

Kumar A, Zhang J, Yu FSX (2006). **Toll-like receptor 3 agonist poly(I:C)-induced antiviral response in human corneal epithelial cells**. *Immunology*, *117*(1), 11–21. doi: 10.1111/j.1365-2567.2005.02258.x.

Lawson ND, Li R, Shin M, Grosse A, Yukselen O, Stone OA, Kucukural A, Zhu L (2020). **An improved**
zebrafish transcriptome annotation for sensitive and comprehensive detection of cell type-
specific genes. *ELife*, 9. doi: 10.7554/eLife.55792.

Levraud JP, Jouneau L, Briolat V, Laghi V, Boudinot P (2019). **IFN-Stimulated Genes in Zebrafish**
and Humans Define an Ancient Arsenal of Antiviral Immunity. *The Journal of Immunology*,
203(12), 3361–3373. doi: 10.4049/jimmunol.1900804.
920

Lesch BJ, Page DC (2014). **Poised chromatin in the mammalian germ line**. *Development*,
141(19), 3619–3626. doi: 10.1242/dev.113027.

Li H, Durbin R (2009). **Fast and accurate short read alignment with Burrows–Wheeler**
transform. *Bioinformatics*, *25*(14), 1754–1760. doi: 10.1093/bioinformatics/btp324.

Li B, Dewey CN (2011). **RSEM: accurate transcript quantification from RNA-Seq data with or**
without a reference genome. *BMC Bioinformatics*, *12*(1), 323. doi: 10.1186/1471-2105-12-323.

Li D, Wu M (2021). **Pattern recognition receptors in health and diseases**. *Signal Transduction*
and Targeted Therapy, *6*(1), 291. doi: 10.1038/s41392-021-00687-0.

Librán-Pérez M, Pereiro P, Figueras A, Novoa B (2022). **Transcriptome Analysis of Turbot**
(*Scophthalmus maximus*) Infected With *Aeromonas salmonicida* Reveals a Direct Effect on
Leptin Synthesis as a Neuroendocrine Mediator of Inflammation and Metabolism Regulation.
Frontiers in Marine Science, *9*. doi: 10.3389/fmars.2022.888115.

Liu S, Yu Y, Zhang S, Cole JB, Tenesa A, Wang T, McDaneld TG, Ma L, Liu GE, Fang L (2020).
Epigenomics and genotype-phenotype association analyses reveal conserved genetic
architecture of complex traits in cattle and human. *BMC Biology*, 18(1), 80. doi:
10.1186/s12915-020-00792-6.

Love MI, Huber W, Anders S (2014). **Moderated estimation of fold change and dispersion for**
RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550. doi: 10.1186/s13059-014-0550-8.

Marinis JM, Hutti JE, Homer CR, Cobb BA, Cantley LC, McDonald C, Abbott DW (2012). **IκB Kinase**
***α* Phosphorylation of TRAF4 Downregulates Innate Immune Signaling**. *Molecular and Cellular*
Biology, *32*(13), 2479–2489. doi: 10.1128/MCB.00106-12.

Maroso F, Hermida M, Millán A, Blanco A, Saura M, Fernández A, Dalla-Rovere G, Bargelloni L,
Cabaleiro S, Villanueva B, Bouza C, Martínez P (2018). **Highly dense linkage maps from 31 full-**
sibling families of turbot (*Scophthalmus maximus*) provide insights into recombination
patterns and chromosome rearrangements throughout a newly refined genome assembly.
DNA Research, 25(4), 439–450. doi: 10.1093/dnares/dsy015.

Martin M (2011). **Cutadapt removes adapter sequences from high-throughput sequencing**
reads. *EMBnet.Journal*, *17*(1), 10. doi: 10.14806/ej.17.1.200

Martínez P (2016). **Genomics advances for boosting aquaculture breeding programs in Spain**.
Aquaculture, *464*, 117–120. doi: 10.1016/j.aquaculture.2016.06.021.

Martínez P, Robledo D, Taboada X, Blanco A, Moser M, Maroso F, Hermida M, Gómez-Tato A,
Álvarez-Blázquez B, Cabaleiro S, Piferrer F, Bouza C, Lien S, Viñas AM (2021). **A genome-wide**
association study, supported by a new chromosome-level genome assembly, suggests sox2 as

**a main driver of the undifferentiated ZZ/ZW sex determination of turbot (*Scophthalmus*
maximus)**. *Genomics*, 113(4), 1705–1718. doi: 10.1016/j.ygeno.2021.04.007.

Martínez de Paz A, Josefowicz S Z (2021). **Signaling-to-chromatin pathways in the immune**
system. *Immunological Reviews*, *300*(1), 37–53. doi: 10.1111/imr.12955.

Millán A, Gómez-Tato A, Fernández C, Pardo BG, Álvarez-Dios JA, Calaza M, Bouza C, Vázquez M,
Cabaleiro S, Martínez P (2010). **Design and Performance of a Turbot (*Scophthalmus maximus*)**
Oligo-microarray Based on ESTs from Immune Tissues. *Marine Biotechnology*, *12*(4), 452–465.
doi: 10.1007/s10126-009-9231-0.

Millán A, Gómez-Tato A, Pardo BG, Fernández C, Bouza C, Vera M, Alvarez-Dios JA, Cabaleiro S,
Lamas J, Lemos ML, & Martínez P (2011). **Gene Expression Profiles of the Spleen, Liver, and**
Head Kidney in Turbot (*Scophthalmus maximus*) Along the Infection Process with *Aeromonas*
***salmonicida* Using an Immune-Enriched Oligo-microarray**. *Marine Biotechnology*, 13(6), 1099–
1114. doi: 10.1007/s10126-011-9374-7.

Mishra S, Das R, Swain P (2018). **Status of Fish Diseases in Aquaculture and Assessment of**
Economic Loss Due to Disease; Today and Tomorrow's Printers and Publishers: New Delhi, India,
Volume 1.

Moazed D (2011). **Mechanisms for the Inheritance of Chromatin States**. *Cell*, 146(4), 510–518.
doi: 10.1016/j.cell.2011.07.013.

Mokhtar DM, Zaccone G, Alesci A, Kuciel M, Hussein MT, Sayed RKA (2023). **Main Components**
of Fish Immunity: An Overview of the Fish Immune System. *Fishes*, *8*(2), 93. doi:
10.3390/fishes8020093.

Murira A, Lamarre A (2016). **Type-I Interferon Responses: From Friend to Foe in the Battle**
against Chronic Viral Infection. *Frontiers in Immunology*, 7. doi: 10.3389/fimmu.2016.00609.

Navarini AA, Recher M, Lang KS, Georgiev P, Meury S, Bergthaler A, Flatz L, Bille J, Landmann R,
Odermatt B, Hengartner H, Zinkernagel RM (2006). **Increased susceptibility to bacterial**
superinfection as a consequence of innate antiviral responses. *Proceedings of the National*
Academy of Sciences, 103(42), 15535–15539. doi: 10.1073/pnas.0607325103.

Nie L, Xu XX, Xiang LX, Shao JZ, Chen J (2017). **Mutual Regulation of NOD2 and RIG-I in Zebrafish**
Provides Insights into the Coordination between Innate Antibacterial and Antiviral Signaling
Pathways. *International Journal of Molecular Sciences*, *18*(6), 1147. doi: 10.3390/ijms18061147.

Ødegård J, Olesen I, Gjerde B, Klemetsdal G (2007). **Positive genetic correlation between**
resistance to bacterial (furunculosis) and viral (infectious salmon anaemia) diseases in farmed
Atlantic salmon (*Salmo salar*). *Aquaculture, 271*(1–4), 173–177. doi:
10.1016/j.aquaculture.2007.06.006.

Ødegård J, Baranski M, Gjerde B, Gjedrem T (2011**). Methodology for genetic evaluation of**
disease resistance in aquaculture species: challenges and future prospects. *Aquaculture*
Research, 42, 103–114. doi: 10.1111/j.1365-2109.2010.02669.x.

Overbey EG, Ng TT, Catini P, Griggs LM, StewartP, Tkalcic S, Hawkins RD, Drechsler Y (2021).
Transcriptomes of an Array of Chbiicken Ovary, Intestinal, and Immune Cells and Tissues.
Frontiers in Genetics, 12. doi: 10.3389/fgene.2021.664424.

Pan Z, Yao Y, Yin H, Cai Z, Wang Y, Bai L, Kern C, Halstead M, Chanthavixay G, Trakooljul N,
Wimmers K, Sahana G, Su G, Lund MS, Fredholm M, Karlskov-Mortensen P, Ernst CW, Ross P,
Tuggle CK, … Zhou H (2021). **Pig genome functional annotation enhances the biological
interpretation of complex traits and human disease**. *Nature Communications*, 12(1), 5848. doi:
10.1038/s41467-021-26153-7.

Pan Z, Wang Y, Wang M, Wang Y, Zhu X, Gu S, Zhong C, An L, Shan M, Damas J, Halstead MM,
Guan D, Trakooljul N, Wimmers K, Bi Y, Wu S, Delany ME, Bai X, Cheng HH, … Zhou H (2023). **An
atlas of regulatory elements in chicken: A resource for chicken genetics and genomics**. *Science
Advances*, 9(18). doi: 10.1126/sciadv.ade1204.

Pardo BG, Fernández C, Millán A, Bouza C, Vázquez-López A, Vera M, Alvarez-Dios JA, Calaza M,
Gómez-Tato A, Vázquez M, Cabaleiro S, Magariños B, Lemos ML, Leiro JM, Martínez P (2008).
**Expressed sequence tags (ESTs) from immune tissues of turbot (*Scophthalmus maximus*)
challenged with pathogens**. *BMC Veterinary Research*, 4(1), 37. doi: 10.1186/1746-6148-4-37.

Pardo BG, Millán A, Gómez-Tato A, Fernández C, Bouza C, Alvarez-Dios JA, Cabaleiro S, Lamas J,
Leiro JM, Martínez P (2012). **Gene Expression Profiles of Spleen, Liver, and Head Kidney in
Turbot (*Scophthalmus maximus*) Along the Infection Process with *Philasterides dicentrarchi*
Using an Immune-Enriched Oligo-Microarray**. *Marine Biotechnology*, 14(5), 570–582. doi:
10.1007/s10126-012-9440-9.

Park PJ (2009). **ChIP–seq: advantages and challenges of a maturing technology**. *Nature Reviews
Genetics*, 10(10), 669–680. doi: 10.1038/nrg2641.

Patel H, Ewels P, Peltzer A, Botvinnik O, Sturm G, Moreno D, Vemuri P, Garcia MU, Morins S,
Pantano L, Binzer-Panchal M, Syme R, Zepper M, Kelly G, Hanssen F, Fellows Yates JA, Cheshire
C, Espinosa-Carrasco J, Miller E, Talbot A, Zhou P, Guinchard S, Hörtenhuber M, Gabernet G,
Mertes C, Straub D, Di Tommaso P (2023). **nf-core/rnaseq: nf-core/rnaseq v3.10.1 – Plastered
Rhodium Rudolph**. *Zenodo*. doi: 10.5281/zenodo.7505987.

Prowse-Wilkins CP, Wang J, Xiang R, Garner JB, Goddard ME, Chamberlain AJ (2021). **Putative
Causal Variants Are Enriched in Annotated Functional Regions From Six Bovine Tissues**.
Frontiers in Genetics, *12*. doi: 10.3389/fgene.2021.664379.

Raghavan V, Kraft L, Mesny F, Rigerte L (2022). **A simple guide to *de novo* transcriptome
assembly and annotation**. *Briefings in Bioinformatics*, 23(2). doi: 10.1093/bib/bbab563.

Ramberg S, Høyheim B, Østbye TKK, Andreassen R (2021). **A *de novo* Full-Length mRNA
Transcriptome Generated from Hybrid-Corrected PacBio Long-Reads Improves the Transcript
Annotation and Identifies Thousands of Novel Splice Variants in Atlantic Salmon**. *Frontiers in
Genetics*, 12. doi: 10.3389/fgene.2021.656334.

Ramírez F, Dündar F, Diehl S, Grüning BA, Manke T (2014). **deepTools: a flexible platform for
exploring deep-sequencing data**. *Nucleic Acids Research*, *42*(W1), W187–W191. doi:
10.1093/nar/gku365.

Robinson JT, Thorvaldsdottir H, Turner D, Mesirov JP (2023). **igv.js: an embeddable JavaScript**
implementation of the Integrative Genomics Viewer (IGV). *Bioinformatics*, *39*(1). doi:
10.1093/bioinformatics/btac830.

Robledo D, Ronza P, Harrison PW, Losada AP, Bermúdez R, Pardo BG, José-Redondo M, Sitjà-
Bobadilla A, Quiroga MI, Martínez P (2014). **RNA-seq analysis reveals significant transcriptome**
changes in turbot (*Scophthalmus maximus*) suffering severe enteromyxosis. *BMC Genomics*,
15(1), 1149. doi: 10.1186/1471-2164-15-1149.

Robledo D, Martin AP, Álvarez-Dios JA, Bouza C, Pardo BG, Martínez P (2017). **First**
characterization and validation of turbot microRNAs. *Aquaculture*, *472*, 76–83. do:
10.1016/j.aquaculture.2016.05.002.

Ronza P, Robledo D, Bermúdez R, Losada AP, Pardo BG, Sitjà-Bobadilla A, Quiroga MI, Martínez P
(2016). **RNA-seq analysis of early enteromyxosis in turbot (*Scophthalmus maximus*): new**
insights into parasite invasion and immune evasion strategies. *International Journal for*
Parasitology, 46(8), 507–517. doi: 10.1016/j.ijpara.2016.03.007.

Ronza P, Robledo D, Losada AP, Bermúdez R, Pardo BG, Martínez P, Quiroga MI (2020). **The**
Teleost Thymus in Health and Disease: New Insights from Transcriptomic and
Histopathological Analyses of Turbot, *Scophthalmus maximus*. *Biology*, *9*(8), 221. doi:
10.3390/biology9080221.

Saeed S, Quintin J, Kerstens HHD, Rao NA, Aghajanirefah A, Matarese F, Cheng SC, Ratter J,
Berentsen K, van der Ent MA, Sharifi N, Janssen-Megens EM, ter Huurne M, Mandoli A, van

Sahoo M, Ceballos-Olvera I, del Barrio L, Re F (2011). **Role of the Inflammasome, IL-1β, and IL-**
18 in Bacterial Infections. *The Scientific World JOURNAL*, *11*, 2037–2050. doi:
10.1100/2011/212680.

Saura M, Carabaño MJ, Fernández A, Cabaleiro S, Doeschl-Wilson AB, Anacleto O, Maroso F,
Millán A, Hermida M, Fernández C, Martínez P, Villanueva B (2019). **Disentangling Genetic**
Variation for Resistance and Endurance to Scuticociliatosis in Turbot Using Pedigree and
Genomic Information. *Frontiers in Genetics*, *10*. doi: 10.3389/fgene.2019.00539.

Secombes CJ, Wang T, Hong S, Peddie S, Crampe M, Laing KJ, Cunningham C, Zou J (2001).
Cytokines and innate immunity of fish. *Developmental & Comparative Immunology*, *25*(8–9),
713–723. doi: 10.1016/S0145-305X(01)00032-5.

Shan S, Liu R, Jiang L, Zhu Y, Li H, Xing W, Yang G (2018). **Carp Toll-like receptor 8 (Tlr8): An**
intracellular Tlr that recruits TIRAP as adaptor and activates AP-1 pathway in immune
response. *Fish & Shellfish Immunology*, *82*, 41–49. doi: 10.1016/j.fsi.2018.08.001.

Stark R, Brown G (2011). **DiffBind: differential binding analysis of ChIP-Seq peak data**.
Bioconductor. doi: 10.18129/B9.bioc.DiffBind.

Stępniak K, Machnicka MA, Mieczkowski J, Macioszek A, Wojtaś B, Gielniewski B, Poleszak K,
Perycz M, Król SK, Guzik R, Dąbrowski MJ, Dramiński M, Jardanowska M, Grabowicz I, Dziedzic
A, Kranas H, Sienkiewicz K, Diamanti K, Kotulska K, … Wilczyński B (2021). **Mapping chromatin**
accessibility and active regulatory elements reveals pathological mechanisms in human
gliomas. *Nature Communications*, *12*(1), 3621. doi: 10.1038/s41467-021-23922-2.

Srour N, Khan S, Richard S (2022**). The Influence of Arginine Methylation in Immunity and Inflammation**. *Journal of Inflammation Research*, *Volume 15*, 2939–2958. doi: 10.2147/JIR.S364190.

Subasinghe R, Soto D, Jia J (2009). **Global aquaculture and its role in sustainable development**. *Reviews in Aquaculture*, *1*(1), 2–9. doi: 10.1111/j.1753-5131.2008.01002.x.

Sullivan C, Soos BL, Millard PJ, Kim CH, King BL (2021). **Modeling Virus-Induced Inflammation in Zebrafish: A Balance Between Infection Control and Excessive Inflammation**. *Frontiers in Immunology*, *12*. doi: 10.3389/fimmu.2021.636623.

Summers KM, Bush SJ, Wu C, Su AI, Muriuki C, Clark EL, Finlayson HA, Eory L, Waddell LA, Talbot R, Archibald AL, Hume DA (2020). **Functional Annotation of the Transcriptome of the Pig, *Sus scrofa*, Based Upon Network Analysis of an RNAseq Transcriptional Atlas**. *Frontiers in Genetics*, 10. doi: 10.3389/fgene.2019.01355.

Sun K, Metzger DW (2008). **Inhibition of pulmonary antibacterial defense by interferon-γ during recovery from influenza infection**. *Nature Medicine*, 14(5), 558–564. doi: 10.1038/nm1765.

Taylor RS, Ruiz Daniels R, Dobie R, Naseer S, Clark TC, Henderson NC, Boudinot P, Martin SAM, Macqueen DJ (2022). **Single cell transcriptomics of Atlantic salmon (*Salmo salar* L.) liver reveals cellular heterogeneity and immunological responses to challenge by** Aeromonas salmonicida. *Frontiers in Immunology*, *13*. doi: 10.3389/fimmu.2022.984799.

Troell M, Costa-Pierce B, Stead S, Cottrell RS, Brugere C, Farmery AK, Little DC, Strand Å, Pullin R, Soto D, Beveridge M, Salie K, Dresdner J, Moraes-Valenti P, Blanchard J, James P, Yossa R, Allison E, Devaney C, Barg U (2023). **Perspectives on aquaculture's contribution to the Sustainable Development Goals for improved human and planetary health**. *Journal of the World Aquaculture Society*, 54(2), 251–342. doi: 10.1111/jwas.12946.

Tuggle CK, Giuffra E, White SN, Clarke L, Zhou H, Ross PJ, Acloque H, Reecy JM, Archibald A, Bellone RR, Boichard M, Chamberlain A, Cheng H, Crooijmans RPMA, Delany ME, Finno CJ, Groenen MAM, Hayes B, Lunney JK, … Watson M (2016). **GO-FAANG meeting: a Gathering On Functional Annotation of Animal Genomes**. *Animal Genetics*, *47*(5), 528–533. doi: 10.1111/age.12466.

Turkistany SA, DeKoter RP (2011). **The Transcription Factor PU.1 is a Critical Regulator of Cellular Communication in the Immune System**. *Archivum Immunologiae et Therapiae Experimentalis*, *59*(6), 431–440. doi: 10.1007/s00005-011-0147-9.

Valle A, Leiro JM, Pereiro P, Figueras A, Novoa B, Dirks RPH, Lamas J (2020). **Interactions between the Parasite Philasterides dicentrarchi and the Immune System of the Turbot *Scophthalmus maximus*. A Transcriptomic Analysis**. *Biology*, 9(10), 337. doi: 10.3390/biology9100337

van Berkum NL, Lieberman-Aiden E, Williams L, Imakaev M, Gnirke A, Mirny LA, Dekker J, Lander ES (2010). **Hi-C: A Method to Study the Three-dimensional Architecture of Genomes**. *Journal of Visualized Experiments*, 39. doi: 10.3791/1869.

Villar D, Frost S, Deloukas P, Tinker A (2020). **The contribution of non-coding regulatory elements to cardiovascular disease**. *Open Biology*, *10*(7). doi: 10.1098/rsob.200088.

Vu H, Ernst J (2022). **Universal annotation of the human genome through integration of over a thousand epigenomic datasets**. *Genome Biology*, *23*(1), 9. doi: 10.1186/s13059-021-02572-z.

Watson DK, Li R, Sementchenko VI, Mavrothalassitis G, Seth A (2002). **ETS Family of Transcription Factors**. In *Encyclopedia of Cancer* (pp. 189–196). Elsevier. doi: 10.1016/B0-12-227555-1/00078-2.

Weidemüller P, Kholmatov M, Petsalaki E, Zaugg JB (2021). **Transcription factors: Bridge between cell signaling and gene regulation**. *PROTEOMICS*, *21*(23–24). doi: 10.1002/pmic.202000034.

Xiang R, MacLeod IM, Daetwyler HD, de Jong G, O'Connor E, Schrooten C, Chamberlain AJ, Goddard ME (2021). **Genome-wide fine-mapping identifies pleiotropic and functional variants that predict many traits across global cattle populations**. *Nature Communications*, *12*(1), 860. doi: 10.1038/s41467-021-21001-0.

Xu X, Shao C, Xu H, Zhou Q, You F, Wang N, Li W, Li M, Chen S (2020). **Draft genomes of female and male turbot *Scophthalmus maximus***. *Scientific Data*, *7*(1), 90. doi: 10.1038/s41597-020-0426-6.

Yan F, Powell DR, Curtis DJ, Wong NC (2020a). **From reads to insight: a hitchhiker's guide to ATAC-seq data analysis**. *Genome Biology*, 21(1), 22. doi: 10.1186/s13059-020-1929-3.

Yan X, Zhao X, Huo R, Xu T (2020b). **IRF3 and IRF8 Regulate NF-κB Signaling by Targeting MyD88 in Teleost Fish**. *Frontiers in Immunology*, *11*. doi: 10.3389/fimmu.2020.00606.

Yu Y, Wang J, Khaled W, Burke S, Li P, Chen X, Yang W, Jenkins NA, Copeland NG, Zhang S, Liu P (2012). **Bcl11a is essential for lymphoid development and negatively regulates p53**. *Journal of Experimental Medicine*, *209*(13), 2467–2483. doi: 10.1084/jem.20121846.

Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, Nusbaum C, Myers RM, Brown M, Li W, Liu XS (2008). **Model-based Analysis of ChIP-Seq (MACS)**. *Genome Biology*, *9*(9), R137. doi: 10.1186/gb-2008-9-9-r137.

Zhang R, Kang R, Tang D (2021). **The STING1 network regulates autophagy and cell death**. *Signal Transduction and Targeted Therapy*, *6*(1), 208. doi: 10.1038/s41392-021-00613-4.

Zhou Y, Zhou B, Pache L, Chang M, Khodabakhshi AH, Tanaseichuk O, Benner C, Chanda SK (2019). **Metascape provides a biologist-oriented resource for the analysis of systems-level datasets**. *Nature Communications*, *10*(1), 1523. doi: 10.1038/s41467-019-09234-6.

Zhu M, John S, Berg M, Leonard WJ (1999). **Functional Association of Nmi with Stat5 and Stat1 in IL-2- and IFN γ-Mediated Signaling**. *Cell*, *96*(1), 121–130. doi: 10.1016/S0092-8674(00)80965-4.

Zhu XN, Wang YZ, Li C, Wu HY, Zhang R, Hu XX. (2023). **Chicken chromatin accessibility atlas accelerates epigenetic annotation of birds and gene fine-mapping associated with growth traits**. *Zoological Research*, 44(1), 53–62. doi: 10.24272/j.issn.2095-8137.2022.228.