Spatiotemporal Changes in Netrin/Dscam1 Signaling Dictate Axonal Projection Direction in *Drosophila* Small Ventral Lateral Clock Neurons

Jingjing Liu[1], Yuedong Wang[1], Junhai Han[1,2, *], Yao Tian[1, *]

[1] School of Life Science and Technology, the Key Laboratory of Developmental Genes and Human Disease, Southeast University, Nanjing 210096, China

[2] Co-innovation Center of Neuroregeneration, Nantong University, Nantong, China

* Correspondence: junhaihan@seu.edu.cn, yaotian@seu.edu.cn,

Abstract

Axon projection is a spatial and temporal-specific process in which the growth cone receives environmental signals guiding axons to their final destination. However, the mechanisms underlying changes in axonal projection direction without well-defined landmarks remain elusive. Here, we present evidence showcasing the dynamic nature of axonal projections in *Drosophila*'s small ventral lateral clock neurons (s-LNvs). Our findings reveal that these axons undergo an initial vertical projection in the early larval stage, followed by a subsequent transition to a horizontal projection in the early-to-mid third instar larvae. The vertical projection of s-LNv axons correlates with mushroom body calyx expansion, while the s-LNv-expressed Down syndrome cell adhesion molecule (Dscam1) interacts with Netrins to regulate the horizontal projection. During a specific temporal window, locally newborn dorsal clock neurons (DNs) secrete Netrins, facilitating the transition of axonal projection direction in s-LNvs. Our study establishes a compelling *in vivo* model to probe the mechanisms of axonal projection direction switching in the absence of clear landmarks. These findings underscore the significance of dynamic local microenvironments in the synergetic regulation of axonal projection direction transitions.

Keywords: Axon projection; projection direction switch; Dscam1; Netrin; s-LNvs; *Drosophila*;

Introduction

During nervous system development, neurons extend axons to reach their targets in order to build functional neural circuits (Cang & Feldheim, 2013; Kaas, 1997). The growth cone, which is specialized at the tip of the extending axon, is critical for receiving multiple guidance signals from the external environment to guide axon projection (Lowery & Vactor, 2009; Mortimer et al., 2008; Vitriol & Zheng, 2012). Guidance cues regulate cytoskeletal dynamics through growth cone-specific receptors, steering axons via attractive or repulsive signals (Araújo & Tear, 2003; Koch et al., 2012; Kolodkin & Tessier-Lavigne, 2011; Esther T Stoeckli, 2018; Zang et al., 2021). Over the last 30 years, several guidance cues have been identified and divided into four classical families: Semaphorins, Ephrins, Netrins, and Slits (Derijck et al., 2010; Dickson, 2002; Goodhill, 2016; X.-T. Li et al., 2014; Orioli & Klein, 1997). The emergence of other kinds of guidance cues, including morphogens, growth factors, and cell adhesion molecules, increases the complexity of axon pathfinding (Chan & Odde, 2008; Chédotal et al., 2012; Kahn & Baas, 2016; Vitriol & Zheng, 2012).

When going through a long-distance pathfinding process in the complex and highly dynamic *in vivo* environment, axons frequently undergo multiple changes in their projection directions (Squarzoni et al., 2015). Crucially, intermediate targets play a pivotal role by providing vital guiding information that enables axons to transition their projection directions and embark upon the subsequent stages of their journey, ultimately leading them to their final destination (de Ramon Francàs et al., 2017; Garel & Rubenstein, 2004; Squarzoni et al., 2015). Certain intermediate targets can respond to the growth cone's signaling to maintain a dynamic balance between attractive and repulsive forces(Bron et al., 2007; Martins et al., 2022). One of the well-studied intermediate target projection types is the midline cross model (Derijck et al., 2010; Yang et al., 2009). As a clear landmark, axons need to project to the midline first and then leave it immediately (Gorla & Bashaw, 2020; Neuhaus-Follini & Bashaw, 2015). The midline floor plate cells secreted long-range attractive signals to attract commissural axons towards them (Kennedy et al., 1994; Serafini et al., 1996). Once there, these cells emit signals of rejection to prevent excessive closeness of axons, thus aiding the axon in departing the floor plate to reach its intended destination (Kidd et al., 1998; Long et al., 2004; E T Stoeckli et al., 1997). Comparatively, the

typical axonal projection process within the *in vivo* milieu undergoes multiple transitions in

projection direction without a distinct landmark, such as the midline (Garel & López-Bendito,

2014; Gezelius & López-Bendito, 2017). However, how the axons change their projection

direction without well-defined landmarks is still unclear.

Drosophila small ventral lateral clock neurons (s-LNvs) exhibit the typical tangential projection

pattern in the central brain (Agrawal & Hardin, 2016; Gummadova et al., 2009; Hardin, 2017;

Helfrich-Förster, 1997). The s-LNv axons originate from the ventrolateral soma and project to the

dorsolateral area of the brain, and subsequently undergo a axon projection direction shift,

projecting horizontally toward the midline with a short dorsal extension (Helfrich-Förster et al.,

2007). Several studies have reported the impact of various factors, including the Slit-Robo signal

(Oliva et al., 2016), Lar (leukocyte-antigen-related) (Agrawal & Hardin, 2016), dfmr1 (Okray et

al., 2015), dTip60 (Pirooznia et al., 2012), and Dscam1 long 3' UTR (Zhang et al., 2019), on the

axon outgrowth of s-LNvs in the adult flies' dorsal projection. However, the mechanism

underlying the projection direction switch of s-LNv axons remains largely unknown. Here, we

show that s-LNvs and newborn dorsal clock neurons (DNs) generate spatiotemporal-specific

guidance cues to precisely regulate the transition of projection direction in s-LNv axons. This

regulation uncovers unexpected interdependencies between axons and local microenvironment

dynamics during the navigation process, enabling accurate control over axon projection.

Results

s-LNvs axons change their projection direction in the early-to-mid third instar larvae

The stereotypical trajectory of the s-LNvs axon projection can be succinctly characterized as an

initial vertical extension originating from the ventrolateral brain, followed by a directional pivot at

the dorsolateral protocerebrum, ultimately leading to a horizontal projection towards the midline.

Immunoreactivity for pigment-dispersing factor (PDF) (Cyran et al., 2005), representing the

pattern of LNv neurons, is initially detected in the brains of 1st-instar larvae 4-5 hours after larval

hatching (ALH) (Helfrich-Förster, 1997). To elucidate the intricate axon pathfinding process

during development, we employed a dual-copy *Pdf-GAL4* to drive the expression of double-copy

UAS-mCD8::GFP, visualizing the s-LNvs at the larval stage (Figure 1A). At 8 hours ALH, the

GFP signal strongly co-localized with the PDF signal within the axons. This remarkable

co-localization continued throughout the larval stage, particularly concentrated at the axon's

terminal end (Figure 1A).

We then quantified the process of s-LNvs axonal projection. The vertical projection distance of the

s-LNvs axon was ascertained by measuring the vertical span between the lowest point of the

optical lobe and the highest point of the axon's projection terminal. Statistical analyses revealed a

steadfast linear increase in the vertical projection distance of the s-LNvs axon as larval

development progressed (Figure 1A-B). To determine the horizontal projection distance of the

s-LNvs axon, we applied the methodology employed by Olive *et al.* for measuring adult flies

(Oliva et al., 2016). A tangent line is drawn precisely at the pivotal juncture where the s-LNvs

axon undergoes a shift in its projection direction. We define the Arbitrary Unit (A.U.) as the ratio

between the horizontal distances from this point to both the end of the axon projection and the

midline (Figure 1A and 1C). Remarkably, at the time points of 8, 24, 48, and 72 hours ALH, the

A.U. value for s-LNvs axon projection in the horizontal direction remained relatively stable at

approximately 0.13. However, at 96 and 120 hours ALH, a significant and striking increase in the

A.U. value ensued, eventually reaching approximately 0.25 (Figure 1C). These findings suggest

that the precise transition of s-LNv axon projection from vertical towards the horizontal direction

occurs in the early-to-mid third instar larvae, specifically during 72 to 96 hours ALH (Figure 1D).

Vertical projection of s-LNvs axons correlate with mushroom body calyx expansion

The axons of s-LNv project dorsally, coming in close proximity to the dendritic tree of the

mushroom body (MB), specifically the calyx (Figures 2A and 2C). This spatial arrangement

facilitates potential interactions between the s-LNvs axon and the MB. MB predominantly

comprises 2500 intrinsic neurons, known as Kenyon Cells (KCs) (Puñal et al., 2021). The

development of the MB is a dynamic process characterized by three primary types of KCs: α/β,

α'/β', and γ, each following distinct temporal schedules of birth. The γ cells emerge from the

embryonic stage until early 3rd-instar larval phase, while the α'/β' cells are born during the latter

half of larval life, and the α/β cells appear in the post-larval stage (Lee et al., 1999; Puñal et al.,

2021). We noticed that s-LNv axons exhibit only vertical growth before early 3rd-instar larval

stage, which coincides with the emergence of γ KCs. Consequently, we have contemplated the

potential correlation between the vertical projection of s-LNv axons and the growth of the

mushroom body.

Next, we closely monitored the spatial relationship between the calyx and the axon projections of

the s-LNvs throughout different larval developmental stages (Figure 2A). We used *OK107-GAL4*

and *Tab2-201Y-GAL4* to drive the expression of mCD8::GFP in all KC subtypes or specifically in

γ KCs (Pauls et al., 2010), respectively, to visualize MB KCs. As development progressed, the

calyx area defined by both GAL4 drivers exhibited a consistent expansion. Statistical analyses

further confirmed that the calyx area increased proportionally with the advancement of

developmental time, demonstrating a linear relationship (Figure 2B-C). Additionally, Pearson

correlation analysis unveiled a robust positive correlation between the calyx area and the vertical

projection distance of s-LNvs axon ($r = 0.9987$, $P < 0.0001$ for *OK107-GAL4*; $r = 0.9963$, $P =$

0.0003 for *Tab2-201Y-GAL4*) (Figure 2B-C). These compelling observations suggest that the MB

calyx, and potentially specifically the calyx of γ KCs, exert significant influence on the vertical

projection of the s-LNvs axon.

To validate our idea, we employed a straightforward approach to evaluate any changes in s-LNvs

axon projection after ablating γ KCs using the apoptosis-promoting factors RPRC and HID driven

by the *Tab2-201Y GAL4*. Initially, we confirmed the effectiveness of our approach. In the control

group, the GFP signal in γ KCs remained unchanged, while the ablation group showed a

significant reduction or complete absence of the signal (Figure 2-figure supplement 1).

Subsequently, we examined the axon projection of s-LNvs under these conditions. In the

non-ablated group, the vertical distance increased linearly during development, following the

expected pattern (Figure 2D-E). However, in the ablation group, although the initial vertical

projection of s-LNvs appeared normal, there was a noticeable decrease in vertical projection

distance at 48 hours ALH. This vertical development trend ceased thereafter, with the vertical

projection distance measuring only 20 to 30 µm at 72 hours ALH (Figure 2D-E). Importantly, in

later larval stages, we observed that s-LNvs in certain larvae became undetectable (data not

shown), suggesting the potential occurrence of apoptosis in these neurons following MB ablation.

These results underscore a significant association between MB development, specifically the γ

KCs, and the vertical projection dynamics of s-LNv axons.

s-LNv-expressed Dscam1 mediate s-LNv axons horizontal projection

To decipher the signaling pathways governing s-LNvs axon horizontal projection, we performed a

screening with a total of 285 targets by using the RNA-interfering (RNAi) approach

(Supplementary File 3). To optimize efficiency, we chose to use *Clk856-GAL4*, which exhibits

high expression in s-LNv during early larval stages, despite also being expressed in subsets of

other clock neurons (Gummadova et al., 2009). Significantly, the knockdown of Down syndrome

cell adhesion molecule (Dscam1) exhibited a marked reduction in the horizontal projection

distance of s-LNv axons (Figure 3-figure supplement 1A-B). To validate the role of

s-LNv-expressed Dscam1 in governing axon horizontal projection, we employed the *Pdf-GAL4*,

which specifically targeted s-LNvs during the larval stage. At 96 hours ALH, vertical projection

showed no significant changes in *Pdf-GAL4;UAS-Dscam1-RNAi* flies, but a noticeable deficit was

observed in the horizontal projection of the s-LNvs axon (Figure 3A-C). These observations

strongly indicate that Dscam1 specifically regulates the horizontal projection of s-LNvs axon.

We further checked s-LNvs axon projection in several well-defined *Dscam1* mutants, including

Dscam1²¹, Dscam1¹, and Dscam1⁰⁵⁵¹⁸ (Hummel et al., 2003; Schmucker et al., 2000). While all

heterozygous mutants displayed normal horizontal axon projection in s-LNvs, trans-heterozygous

mutants and homozygotes exhibited shortened horizontal axon projection (Figure 3D-E).

Additionally, our immunostaining results indicated concentrated Dscam1 expression at the growth

cone of s-LNv axons in 3rd-instar larvae (Figure 3F). Taken together, these data demonstrate that

Dscam1 controls the horizontal axon projection of s-LNvs in a cell-autonomous manner.

Dscam1 is a well-recognized cell adhesion molecule that plays a crucial role in axon guidance

(Chen et al., 2006; Hummel et al., 2003; Zhan et al., 2004; Zhang et al., 2019). The receptors

present on growth cones sensor the extracellular axon guidance molecules to initiate the

reorganization of the cellular cytoskeleton, leading to the facilitation of axonal projection.

Convincingly, knockdown of the cytoskeletal molecules *tsr* (the *Drosophila* homolog of *cofilin*)

(Sudarsanam et al., 2020) and *chic* (the *Drosophila* homolog of *profilin*) (Shields et al., 2014) in

s-LNvs recaptured the horizontal axon projection deficits in *Dscam1* knockdown flies or *Dscam1*

mutants (Figure 3-figure supplement 1C-D). Consistently, knockdown of *Dock* and *Pak*, the

guidance receptor partners of Dscam1 (Schmucker et al., 2000), or *SH3PX1*, the critical linker

between Dscam1 and the cytoskeleton (Worby et al., 2001), phenocopied *Dscam1* knockdown

flies or *Dscam1* mutants (Figure 3-figure supplement 1C-D). Taken together, these findings

provide solid evidence to support that Dscam1 signaling as a crucial regulator of s-LNvs axon

horizontal projection (Figure 3-figure supplement 1E).

Neuronal-derived Netrins act upstream of Dscam1 to govern the horizontal projection of

s-LNvs axon

Two classical guidance molecules, Slit and Netrin, have been shown to specifically bind to the

extracellular domain of Dscam1 (Alavi et al., 2016; Andrews et al., 2008). The *Drosophila*

genome contains the sole *slit* gene and two *Netrin* genes: *Netrin-A* (*NetA*) and *Netrin-B* (*NetB*)

(Harris et al., 1996; Mitchell et al., 1996). Interestingly, knockdown of both *NetA* and *NetB*

(hereafter referred to as *Netrins*), rather than *slit*, resulted in the defective horizontal axon

projection of s-LNvs. Convincingly, *NetA,NetB* double mutant (Brankatschk & Dickson, 2006),

but not *NetA* or *NetB* single mutant, exhibited the defective horizontal axon projection of s-LNvs

(Figure 4A-B and Figure 4-figure supplement 1). These results imply that two Netrin molecules

function redundantly in regulating horizontal axon projection of s-LNvs.

To determine the source of Netrins, which are known to be secreted axon guidance molecules, we

conducted experiments using pan-neuronal (*nsyb-GAL4*) and pan-glial (*repo-GAL4*) drivers to

selectively knock down *Netrins* (Figure 4C-D). Strikingly, when *Netrins* were knocked down in

neurons but not in glia, we observed severe defects in the horizontal axon projection of s-LNvs.

These findings reveal that neuron-secreted Netrins serve as ligands for Dscam1, controlling the

horizontal axon projection of s-LNvs.

Dorsal neuron-secreted Netrins mediate the horizontal axon projection of s-LNvs

To further identify the source of Netrins, we focused on the neurons located in the dorsolateral

protocerebrum, where the s-LNvs axons change their projection direction. We first excluded the

possibility that MB-secreted Netrins mediate the horizontal axon projection of s-LNvs, as

knockdown of *Netrins* with *OK107-GAL4* showed normal horizontal axon projection of s-LNvs

(Figure 5A-B).

The Dorsal neurons (DNs), a subset of clock neurons, also situate in the dorsolateral

protocerebrum. DNs are categorized into three types, DN1, DN2, and DN3 (Reinhard et al., 2022),

and previous studies have shown that both DN2 and DN1 form synaptic connections with s-LNvs

axon terminals at the adult stage (Schlichting et al., 2022). Interestingly, knockdown of *Netrins* in

a substantial portion of DN2, DN3, and DN1 (Kaneko et al., 1997) (*Per-GAL4,*

Pdf-GAL80;UAS-Netrins-RNAi) resulted in the defective horizontal axon projection of s-LNvs

(Figure 5A-B). To further validate the involvement of DNs in mediating the horizontal axon

projection of s-LNvs, we conducted cell ablation experiments. Notably, the impairments in the

horizontal axonal projection of s-LNvs due to DNs ablation were exclusively observed at 96 hours

ALH, which is after the occurrence of horizontal projection (Figure 5C-D). In contrast, ablating

crz$^+$ neurons, which occupy a similar location to DNs, had no significant effects on the horizontal

axon projection of the s-LNvs (Figure 5-figure supplement 1). Moreover, by expressing NetB in

DNs in *NetA,NetB* double mutants, we successfully restored the defective horizontal axon

projection of s-LNvs (*NetAB$^\Delta$;Per-GAL4, Pdf-GAL80;UAS-NetB*) (Figure 5E-F). These results

demonstrate that DNs secrete Netrins to guide the horizontal axon projection of s-LNvs.

Newborn DNs secret Netrins to regulate the horizontal axon projection of s-LNvs

Finally, we wonder which population of DNs secret Netrins to regulate the horizontal axon

projection of s-LNvs. Therefore, we monitored the number and location of DNs during the s-LNvs

axon projection and found that the number of DNs significantly increased during this process.

Per-GAL4, Pdf-GAL80; UAS-mCD8::GFP only labeled 4-5 cells at 48 hours ALH, and raise to

10-15 by 72 hours, and subsequently increased to approximately 25 by 96 hours ALH (Figure

6A-B). It is worth noting that the location of these newly formed DNs resides lateral to the

transition point of axon projection direction of the s-LNvs, while maintaining a basic parallelism

with the horizontal axon projection of the s-LNvs (Figure 6A). The sharp increase in the number

of DNs coincides remarkably with the timing of the switch in axon projection direction of the

s-LNvs.

Next, we asked whether these newly generated DNs expressed Netrins. To visualize the expression

of endogenous NetB, we applied the fly strains that insert either GFP or myc tag in *NetB* gene. At

72 hours ALH, we were unable to detect any NetB signals within the DNs. In contrast, at 96 hours

ALH, we easily detected prominent NetB signals in approximately 6-8 newborn DNs (Figure

6C-D, Figure 6-figure supplement 1A).

Previous studies have shown that DSCAM is involved in Netrin-1-mediated axonal attraction

(Andrews et al., 2008; G. Liu et al., 2009; Ly et al., 2008). In addition, DSCAM also functions as

a repulsive receptor, associating with Uncoordinated-5C (UNC5C) to mediate Netrin-1-induced

axon growth cone collapse (Purohit et al., 2012). To dissect the role of Netrin signaling, we

ectopically expressed Netrins in neurons marked by *R78G02-GAL4* (Jenett et al., 2012; Suzuki et

al., 2022), located in front of the horizontal projection of s-LNv axons, closer to the midline

(Figure 6-figure supplement 1B). Expression of either NetA or NetB in *R78G02-GAL4*-labeled

neurons significantly suppressed the horizontal axon projection of s-LNvs (Figure 6-figure

supplement 1B-C). Taken together, these findings reveal that newborn DNs secrete Netrins to

orchestrate the transition of axon projection in s-LNvs from a vertical to a horizontal direction

(Figure 7).

Discussion

The mechanisms underlying axonal responses to intricate microenvironments and the precise

development of axons within the brain have long remained enigmatic. In the past, researchers have

identified a large number of axon guidance cues and receptors using different models *in vivo* and

in vitro (Esther T Stoeckli, 2018). In addition, transient cell-cell interactions through intermediate

targets play a crucial role in guiding axonal projection step by step towards its final destination

(Chao et al., 2009; Garel & Rubenstein, 2004). The landmark midline, a crucial intermediate target,

serves as a model in the majority of early studies exploring the directional transitions of axonal

projections (Evans & Bashaw, 2010). However, so many neural projections do not cross the

midline that it is challenging to understand how these axon projections are guided (Goodhill,

2016). In our study, we established an excellent model to investigate the mechanism of axonal

projection direction switch by the *Drosophila* s-LNvs. We discovered a coordinated growth pattern

between the vertical projection of s-LNvs and the MB calyx. Furthermore, the synergistic

interaction between Dscam1 expressed in s-LNvs and the emerging DN-secreted Netrins precisely

modulates the transition of s-LNv axons from a vertical to a horizontal projection within a specific

time window (Figure 7). These findings reveal the mechanism behind the transition of axonal

projection direction, emphasizing the significance of developmental microenvironments in

ensuring precise axon projection.

The dependence of s-LNv vertical projection on mushroom body calyx expansion

During our monitoring of the axonal projection of s-LNvs, we observed that the axon terminals

reached the dorsolateral brain area at an early stage but continued their vertical growth. This

phenomenon raises the question of what drives this growth. The mushroom body, a sophisticated

central hub within the fruit fly's brain, exhibits proximity to the projection of s-LNv axons in

spatial arrangement. Upon careful examination, we found a positive correlation between the

vertical length increase of s-LNv axons and the growth of the mushroom body calyx (Figure 2).

Remarkably, when we selectively removed KC cells, we observed a striking effect on the axonal

projection of s-LNvs. The s-LNv axonal projections either stalled at the initial stage or even

completely disappeared (Figure 2 and data not shown). Hence, the vertical projection of s-LNv

axons is dependent on mushroom body calyx expansion. Our hypothesis is supported by the

findings of Helfrich-Förster *et al.*, who reported locomotor activity and circadian rhythm defects

in some MB mutants (HELFRICH-FÖRSTER et al., 2002), potentially attributed to the

abnormality of s-LNvs axon. Moreover, it was observed that the MB mutants had minimal impact

on the l-LNvs. This can be attributed to the fact that l-LNvs fibers emerge at a later stage during

pupal development and are situated at a spatial distance from the mushroom body.

Unfortunately, we did not identify any molecules that have a discernible impact on the vertical

projection of s-LNv axons through screening. This indicates that the successful completion of the

vertical projection process may play a pivotal role in determining the overall existence of s-LNv

axonal projections. Furthermore, the dorsal projection of s-LNv axons intersects with the ventral

projection of neurons such as DN1 in the dorsolateral region adjacent to the calyx (Keene et al.,

2011), resembling the corpus callosum in mammals (Fothergill et al., 2014; Hutchins et al., 2011;

Keeble et al., 2006; Piper et al., 2009; Unni et al., 2012). Thus, during early developmental stages,

multiple guiding cues may redundantly function between the mushroom body calyx and

neighboring neural processes to ensure the smooth progression of neural development and

establish a stable brain structure. However, to gain a more comprehensive understanding, it is

worthwhile to explore whether the development of the mushroom body calyx influences the

projection of other neurons traversing the same territory and the guidance cues involved. Insights

from existing single-cell sequencing data obtained from multiple stages of larval life (Avalos et al.,

2019; Corrales et al., 2022), might offer some clues and indications.

Netrin/Dscam signaling specifically controls the horizontal projection of s-LNv axons

In this paper, when we specifically knockdown *Dscam1* in s-LNvs, we observed a significant

defect in the axonal horizontal projection. This suggests that Dscam1 autonomously regulates the

horizontal axonal projection of s-LNvs (Figure 3). Our findings align with a recent study that

reported abnormal horizontal axonal projection of s-LNv neurons in adult flies when Dscam1

mRNAs lack the long 3' UTR (Zhang et al., 2019).

The extracellular domain of Dscam1 has been verified to be capable of binding to Netrin or Slit

(Alavi et al., 2016). In *Drosophila*, at the midline of the embryonic central nervous system,

Dscam1 forms a complex with Robo1 to receive the Slit signal and promote the growth of

longitudinal axons (Alavi et al., 2016). In adult flies, Slit/Robo signaling restricts the medial

growth of s-LNv axons (Oliva et al., 2016). However, we found that reducing *Netrins* levels,

rather than *slit*, led to a decrease in the horizontal projection distance of s-LNv axons. Moreover,

in the *NetAB*[A] double mutant, the extent of defects in the horizontal axonal projection of s-LNv

neurons is similar to that observed when knocking down *Dscam1* in s-LNvs. Indeed, previous

studies have provided evidence showcasing the widespread involvement of Dscam1 as a receptor

for Netrin in mediating axon growth and pathfinding (G. Liu et al., 2009; Ly et al., 2008;

Matthews & Grueber, 2011). When we disturbed the expression of Netrin signals in the axon

targeting microenvironment, it was also sufficient to cause abnormal s-LNv projection (Figure

6-figure supplement 1B-C). Therefore, our findings suggest that Netrin, as a ligand for Dscam1,

regulates the process of switching axon projection direction (Figure 4).

The newborn DNs-secreted Netrin coordinates with s-LNv-expressed Dscam1 to switch the

projection direction of s-LNv axons

While the midline serves as a prominent landmark, it presents a daunting challenge to comprehend

the guidance and directional transitions of axons in many neural projections that do not actually

cross this central axis (Evans & Bashaw, 2010). Following vertical projection, the s-LNvs growth

cones remain within the dorsolateral area for at least 48 hours before initiating horizontal

projection (Figure 1). This phenomenon aligns harmoniously with the outcomes observed in

earlier studies that the axonal growth of other types of neurons slows down at the specific choice

point, the midline (Bak & Fraser, 2003; Godement et al., 1994; T. Li et al., 2021), indicating that

the dorsolateral area serves as an intermediate targets to facilitate the subsequent phase of s-LNvs

projection journey.

Extracellular cues released by the final or intermediate targets play a crucial role in guiding axonal

projection. The axon terminals of s-LNvs showed close spatial associations with the somas and

processes of DNs. In this study, the knockdown of *Netrins* specifically in per[+],Pdf[-] neurons, but

not in the mushroom body, resulted in axon projection defects in s-LNvs (Figure 5). Furthermore,

the ablation of DNs resulted in the inhibition of horizontal growth of s-LNvs axons (Figure 5).

These findings suggest that the Netrin signaling microenvironment secreted by DNs is involved in

regulating the horizontal projection of s-LNv axons.

Ideas and Speculation

The emergence of NetB-positive DNs and the aberrant axonal projection of s-LNv neurons caused

by DN ablation occur concurrently during development (Figure 6). In *Drosophila*, three types of

circadian oscillator neurons, including two DN1s, two DN2s, and 4-5 s-LNvs, can be identified as

early as embryonic stage 16 (Houl et al., 2008). The DN3 group, which represents the largest

contingent within the central circadian neuron network with over 35 neurons, emerges during the

larval stage (T. Liu et al., 2015). Little is known about the function of DN3 neurons, particularly

during the larval stage. Due to the lack of cell-specific labeling tools, we can only speculate about

the identity of these newly generated Netrin-secreting neurons as DN3s based on their spatial

distribution. Taken together, these results unveil a novel regulatory mechanism where axons,

during the process of pathfinding, await guidance cues from newly born guidepost cells. This

enables a switch in the direction of axon projection, facilitating the axon's subsequent journey.

Acknowledgements

We thank Dr. Haihuai He for providing *Dscam1* mutant flies; Dr. Yufeng Pan for the fly line used

in cell ablation; Dr. Renjun Tu for the *NetB-GFP* fly; Dr. Xuan Guo for collecting the tool flies;

the Bloomington stock center and Tsinghua fly center for providing flies; Dr. Tzumin Lee for

Dscam1 antibodies; Dr. Ranhui Duan HA-NetA and HA-NetB plasmids. This work was supported

by the National Natural Science Foundation of China (32170970 to Y.T., 32230039 to J.H.), STI

2030- Major Projects-2021ZD0202500 to J.H., and the Guangdong Key Project-2018B030335001

to J.H.

Author Contributions

J. L. designed and performed experiments, interpreted data and wrote the manuscript; Y. W.

performed experiments, Y.T. designed experiments, interpreted data and wrote the manuscript; J.

H. designed experiments, interpreted data, wrote the manuscript, and supervised the project.

Declaration of Interests

The authors declare no competing financial interests.

Corresponding authors

Correspondence to: Yao Tian and Junhai Han, School of Life Science and Technology, Southeast

University, 2 Sipailou Road, Nanjing 210096, China. Tel: +86-25-83790962; Fax:

+86-25-83790962; E-mail: yaotian@seu.edu.cn, junhaihan@seu.edu.cn.

Figure legends

Figure 1. s-LNvs axon projection dynamics during the larval stage.

(A) Images showing the growth process of s-LNvs during larval development. Larval brains were

stained with anti-PDF (white) and HRP (magenta) antibodies. Different time points indicate the

hours after larval hatching (ALH).

(B) Left panel: schematic diagram illustrating the method used to measure the degree of vertical

projection. One hemisphere of the larvae brain is depicted. larval neuropil (gray), s-LNvs (green),

optic lobe (blue). Right panel: graphs showing the average vertical projection length at different

developmental stages, presented as mean ± SD.

(C) Left panel: schematic diagram illustrating the method used to measure the degree of horizontal

projection. Right panel: graphs showing the average horizontal projection (A.U.: arbitrary unit) at

different developmental stages, presented as mean ± SD. The red line segment indicates the stage

at which the axonal projection undergoes a directional transition.

(D) Schematic representation of s-LNvs vertical to horizontal projection directional shift during 72

-96 hours ALH. One hemisphere of the larvae brain (gray), larval neuropil (white), s-LNvs

(green).

For (B), (C), 8 h (n = 7), 24 h (n = 10), 48 h (n = 14), 72 h (n = 16), 96 h (n = 13), 120 h (n = 16).

Figure 2. Development of vertical projection process alongside Mushroom Body growth.

(A) Top: Spatial position relationship between calyx and s-LNvs labeled by *OK107-GAL4*, which

labels MB γ and α'β' neurons, at different developmental time points. Bottom: Spatial position

relationship between calyx and s-LNvs labeled by *Tab2-201Y-GAL4*, which labels MB γ neurons,

Larvae brains were stained with anti-PDF (green) and GFP (magenta) antibodies. Different time

indicated hours ALH. The white dotted line indicates the calyx region.

(B) Graphs showing the mushroom body calyx's area and vertical projection length averaged over

different development stage are presented as mean ± SD. *OK107*: 24 h (n = 10), 48 h (n = 8), 72 h

(n = 12), 96 h (n = 12), 120 h (n = 9). Pearson's r = 0.9987, P < 0.0001. *Tab2-201Y*: 24 h (n = 6),

48 h (n = 12), 72 h (n = 13), 96 h (n = 13), 120 h (n = 11). Pearson's r = 0.9963, P = 0.0003.

(C) Schematic representation of synergetic development of s-LNvs vertical projection and

mushroom body calyx. Mushroom body (purple), s-LNvs (Green).

(D) Images of mushroom body ablation in the developing larval stages. Larvae brains were stained

with anti-PDF (white) and HRP (magenta) antibodies. Different time indicated hours ALH.

(E) Quantification of vertical projection length in Control *(Tab2-201Y >GFP)* and Ablation

(Tab2-201Y >GFP,rpr,hid) flies. Data are presented as mean ± SD. Control: 24 h (n = 6), 48 h (n =

14), 72 h (n = 13), Ablation: 24 h (n = 6), 48 h (n = 12), 72 h (n = 12). Two-tailed Student's t tests

were used. ns, $p > 0.05$; ****$p < 0.0001$.

Figure 2- figure supplement 1. Validation of MB ablation efficiency.

(A) Images of mushroom body ablation in the developing larval stages. Larvae brains were stained

with anti-GFP (green) and HRP (magenta) antibodies. Different time indicated after larval

hatching (ALH).

(B) Quantification of MB calyx area in Control *(201Y >GFP)* and Ablation *(201Y >GFP,rpr,hid)*

flies. Data are presented as mean ± SD. Control: 24 h (n = 6), 48 h (n = 12), 72 h (n = 13),

Ablation: 24 h (n = 6), 48 h (n = 12), 72 h (n = 12). Two-tailed Student's t tests, ****$p < 0.0001$.

Figure 3. Critical role of Dscam1 in horizontal projection.

(A) Images of s-LNvs in *Pdf >GFP* and *Pdf >GFP, Dscam1^{RNAi}* fly. White line segment represents

the horizontal projection distance of s-LNvs. Larvae brains were stained with anti-PDF (white)

and HRP (magenta) antibodies at 96 hours ALH. *Pdf >GFP*: (n = 13), *Pdf >GFP, Dscam1^{RNAi}*: (n

= 15).

(B) Quantification of vertical projection length in *Pdf >GFP* and *Pdf >GFP, Dscam1^{RNAi}* flies.

Data are presented as mean ± SD. Two-tailed Student's t tests, ns, $p > 0.05$.

(C) Quantification of horizontal A.U. in *Pdf >GFP* and *Pdf >GFP, Dscam1^{RNAi}* flies. Data are

presented as mean ± SD. Two-tailed Student's t tests, ****$p < 0.0001$.

(D) Images of *Dscam1* mutant s-LNvs projection phenotype. Larvae brains were collected at late

3rd larvae. Heads were stained with anti-PDF (white) and HRP (magenta) antibodies.

(E) Quantification of horizontal A.U. in *Dscam1* mutant flies. Data are presented as mean ± SD.

w^{1118} (n = 22) *Dscam1^{21}/+* (n = 5), *Dscam1^{1}/+* (n = 8), *Dscam1^{05518}/+* (n = 10), *Dscam1^{21}/*

Dscam1^{05518} (n = 6), *Dscam1^{21}/Dscam1^{1}* (n = 10), *Dscam1^{05518}* (n = 14). One-way ANOVA with

Tukey's post hoc, ns, $p > 0.05$, ****$p < 0.0001$.

(F) Endogenous Dscam1 co-localizes with the s-LNvs axon terminal. *Pdf >GFP* fly heads were

collected at 120 h ALH and stained with anti-Dscam1 (red). The white dotted line indicates the

s-LNvs axon.

Figure 3- figure supplement 1. Dscam1 and its downstream signaling pathway in s-LNvs

horizontal projection.

(A) Images of immunostained in *Clk856-GAL4* knockdown Dscam1. Larvae brains were collected

at late 3rd larvae. Heads were stained with anti-PDF (white) and HRP (magenta) antibodies.

(B) Quantification of horizontal A.U. in *Clk856-GAL4* knockdown Dscam1. Data are presented as

mean ± SD. *Clk856-GAL4/+* (n = 6), *Clk856-GAL4 >Dscam1^{RNAi}* (n = 10). Two-tailed Student's t

tests, ****p < 0.0001.

(C) Images of immunostained in *Pdf-GAL4* knockdown cytoskeleton-associated regulatory

proteins and Dscam1 interaction molecule. Larvae brains were collected at late 3rd larvae. Heads

were stained with anti-PDF (white) and HRP (magenta) antibodies.

(D) Quantification of horizontal A.U. in *Pdf-GAL4* knockdown *Dscam1* interaction molecule and

cytoskeleton-associated regulatory proteins. Data are presented as mean ± SD. *Pdf-GAL4/+* (n =

14), *Pdf-GAL4 >tsr^{RNAi}* (n = 6), *Pdf-GAL4 >chic^{RNAi}* (n = 8), *Pdf-GAL4 >SH3PX1^{RNAi}* (n = 6),

Pdf-GAL4 >Dock^{RNAi} (n = 7), *Pdf-GAL4 >pak^{RNAi}* (n = 6). One-way ANOVA with Dunnett's post

hoc, ***p < 0.001 , ****p < 0.0001.

(E) Schematic representation of Dscam1 mediates s-LNvs horizontal projection downstream

signaling pathways. Dscam1 (brown) activates the downstream signaling cascade involving Dock

(cyan), SH3PX1 (yellow), Pak (blue), and cofilin/profilin (depicted in purple and blue balls,

respectively). These signaling events induce alterations in cell cytoskeleton proteins (gray balls),

facilitating the guidance of horizontal projection

Figure 4. Neuron-derived Netrin guides s-LNvs horizontal projection.

(A) Images of immunostained in *w^{1118}*, *NetA^{Δ}*, *NetB^{Δ}* and *NetAB^{Δ}* fly. Larvae brains were collected

at late 3rd larvae. Heads were stained with anti-PDF (white) and HRP (magenta) antibodies.

(B) Quantification of horizontal A.U. *w^{1118}*, *NetA^{Δ}*, *NetB^{Δ}* and *NetAB^{Δ}* fly. Data are presented as

mean ± SD, *w^{1118}* (n = 17), *NetA^{Δ}* (n = 15), *NetB^{Δ}* (n = 24), *NetAB^{Δ}* (n = 23). One-way ANOVA

with Dunnett's post hoc, ns, $p > 0.05$, ***$p < 0.001$.

(C) Images of immunostained in *nSyb-GAL4* and *repo-GAL4* knockdown *Netrins*. Larvae brains

were collected at late 3rd larvae heads were stained with anti-PDF (white) and HRP (magenta)

antibodies.

(D) Quantification of horizontal A.U. in *nSyb-GAL4* and *repo-GAL4* knockdown *Netrins* fly. Data

are presented as mean ± SD. *nSyb-GAL4/+* (n = 13), *nSyb-GAL4 >Netrins^{RNAi}* (n = 16),

repo-GAL4/+ (n = 34), *repo-GAL4 >Netrins^{RNAi}* (n = 20). Two-tailed Student's t tests, ns, $p > 0.05$,

***$p < 0.001$.

Figure 4- figure supplement 1. Identifying the upstream signal of Dscam1.

(A) Images of immunostained in *tubulin-GAL4/+*, *tubulin-GAL4 >Netrins^{RNAi}*,

tubulin-GAL4 >slit^{RNAi} fly. Larvae brains were heads were collected at late 3rd larvae. Head

stained with anti-PDF (white) and HRP (magenta) antibodies.

(B) Quantification of horizontal A.U. in *tubulin-GAL4/+,* *tubulin-GAL4 >Netrins^{RNAi,}*

tubulin-GAL4 >slit^{RNAi} fly. Data are presented as mean ± SD. *tubulin-GAL4/+* (n = 13),

tubulin-GAL4 >Netrins^{RNAi} (n = 20), *tubulin-GAL4 >slit^{RNAi}* (n = 9). One-way ANOVA with

Dunnett's post hoc, ns, $p > 0.05$, **$p < 0.01$.

Figure 5. Netrin secreted by DN neurons guides time-specific horizontal projection of

s-LNvs.

(A) Images of immunostained in *OK107-GAL4* and *per-GAL4, Pdf-GAL80* knockdown *Netrins.*

Larvae brains were collected at late 3rd larvae. Heads were stained with anti-PDF (white) and

HRP (magenta) antibodies.

(B) Quantification of horizontal A.U. in *OK107-GAL4* and *per-GAL4, Pdf-GAL80* knockdown

Netrins fly. Data are presented as mean ± SD. *OK107-GAL4/+* (n = 20),

OK107-GAL4 >Netrins^{RNAi} (n = 13). *per-GAL4, Pdf-GAL80/+* (n = 18), *per-GAL4,*

Pdf-GAL80 >Netrins^{RNAi} (n = 16), Two-tailed Student's t tests for *OK107-GAL4* knockdown

Netrins, ns, $p > 0.05$. Mann-Whitney test for *per-GAL4, Pdf-GAL80* knockdown *Netrins*, ****$p <$

0.0001.

(C) Images of DN ablation in the developing larval stages. Larvae brains were stained with

anti-PDF (white) and HRP (magenta) antibodies. Different time indicated hours ALH.

(D) Quantification of horizontal A.U. in Control (*per-GAL4, Pdf-GAL80 >GFP*) and Ablation

(*per-GAL4, Pdf-GAL80 >GFP,rpr,hid*) flies. Data are presented as mean ± SD. Control: 72 h (n =

5), 96 h (n = 12), Ablation: 72 h (n = 5), 96 h (n = 10). Two-tailed Student's t tests were used to

compare conditions. ns, p > 0.05, ****p < 0.0001.

(E) Images of immunostained in *per-GAL4, Pdf-GAL80* overexpress *NetB* in *NetAB^4*. Larvae

brains were collected at late 3rd larvae. Heads were stained with anti-PDF (white) and HRP

(magenta) antibodies.

(F) Quantification of horizontal A.U. in *per-GAL4, Pdf-GAL80* overexpress *NetB* in *NetAB^4*. Data

are presented as mean ± SD. *NetAB^4, UAS-NetB* (n = 12), *NetAB^4, per-GAL4,Pdf-GAL80 /+* (n =

5), *NetAB^4,per-GAL4,Pdf-GAL80 >UAS-NetB* (n = 9). One-way ANOVA with Bonferroni post

hoc, ns, p > 0.05, ***p < 0.001.

Figure 5- figure supplement 1. Ablation of DN neurons leads to reduced s-LNvs horizontal

projection.

(A) Images of per-GAL4, pdf-GAL80 and crz-GAL4 ablation. Larvae brains were stained with

anti-PDF(white) and HRP(magenta) antibodies at 120 h ALH.

(B) Quantification of horizontal A.U. in *per-GAL4, Pdf-GAL80* and *crz-GAL4* ablation fly. Data

are presented as mean ± SD. *per-GAL4, Pdf-GAL80 >GFP* (n = 14), *per-GAL4,*

Pdf-GAL80 >GFP,rpr,hid (n = 20), *crz-GAL4 >GFP* (n = 4), *crz-GAL4 >GFP,rpr,hid* (n = 3).

Two-tailed Student's t tests, ns, p > 0.05; ****p < 0.0001.

Figure 6. Dynamic changes in DN neurons coordinated with s-LNvs axonal targeting.

(A) Top: Images of s-LNvs axon and DNs growth process in the developing larval stages. Larvae

brains were stained with anti-PDF (white) and GFP (green) antibodies. Different time indicated

hours ALH are shown. Bottom: schematic diagram of s-LNvs horizontal projection and the

corresponding increase in the number of DN neurons. s-LNvs axon (gray), DN neurons (green).

(B) Quantification of the number of DN neurons labeled by *per-GAL4, Pdf-GAL80* at different

developmental stages. Data are presented as mean ± SD.

(C) Images of newborn DN neurons were colocalized with *Netrin-B* at different developmental

times. Larvae brains were stained with anti-GFP (green), mcherry (red) and PDF (white)

antibodies. Different time indicated hours ALH. The white arrow demarcates the co-localization of

red and green signals.

(D) Schematic representation s-LNvs projection directional transition and the corresponding

increase in the number of DN neurons. s-LNvs axon (gray), DN neurons (green), newborn DN

neurons (light green at 72 h, orange at 96 h).

Figure 6- figure supplement 1. Anterior Netrin ectopic expression reduces horizontal

projection length.

(A) Left: Schematic representation s-LNvs projection directional transition and the corresponding

increase in the number of DN neurons. s-LNvs axon (gray), DN neurons(green), newborn DN

neurons (light green at 72 h, orange at 96h). Right: images of DN neurons were colocalized with

Netrin-B at different developmental times. Larvae brains were stained with anti-Myc (red) and

GFP (green) antibodies. Different time indicated after larval hatching (ALH). The white arrow

demarcates the co-localization of red and green signals.

(B) Images of Netrin ectopic expressed in *R78G02-GAL4*. Larvae brains were collected at late 3rd

larvae. Heads were stained with anti-PDF (white) and HRP (magenta) antibodies.

(C) Quantification of horizontal A.U. in Netrin ectopic expressed in *R78G02-GAL4* fly. Data are

presented as mean ± SD. *R78G0-GAL4/+* (n = 11), *R78G02-GAL4 >UAS-NetA^{OE}* (n = 6),

R78G02-GAL4 >UAS-NetB^{OE} (n = 11). One-way ANOVA with Dunnett's post hoc, ***$p < 0.001$,

****$p < 0.0001$.

Figure 7. Dynamic cellular molecular environment during the s-LNv projection directional

shift.

Cartoon depicting the dynamic cellular molecular microenvironment during the axonal projection

directional shift from vertical to horizontal projection in s-LNvs. 72 h and 96 h represent 72 h and

96 h after larval hatching, respectively. s-LNvs (blue), neuropile (pink), brain lobe (gray circule),

DNs (orange and light orange circles), optic lobe (gray dotted line), Netrin (Orange combination

molding), Dscam1 (pruple).

Materials and methods

Fly Genetics

The flies were maintained on standard medium at 25°C with 60–80% relative humidity. The wild-type flies used in this study were w^{1118}. The *Dscam1 (CG17800)* null mutant allele, *Dscam1^{21}*, was obtained from Haihuai He's laboratory. *UAS-rprC;;UAS-hid* fly were kindly provided by Yufeng Pan's laboratory. The RNAi lines used in these studies were purchased from Tsinghua University; other flies were obtained from Bloomington Stock Center. Full genotypes of the flies shown in the main figures and supplemental figures are listed in Supplementary File 2.

Generation of Transgenic Flies

To generate the NetA and NetB transgenes, the full-length NetA cDNAs and the the full-length NetB cDNAs were sub-cloned into the pUAST vectors and injected into w^{1118} flies.

Antibodies

Antibodies were obtained from Developmental Studies Hybridoma Bank (anti-PDF, anti-FasII) Jackson Immuno Research (anti-HRP), Invitrogen (anti-GFP Alexa-488 goat anti-chicken IgY), Rockland (anti-RFP), and Abcam (Alexa-555 goat anti-rabbit IgG, Alexa-647 goat anti-mouse IgG, Alexa 488 goat anti-rabbit IgG and Alexa 488 goat anti-mouse IgG), gifted from Tzumin Lee's Lab (anti-Dscam1).

Collection larvae

~100 females lay eggs on the collecting medium (1% agar and 30% juice) in Petri dishes for a 4-hour period. To obtain larvae of defined ages, freshly emerged larvae were selected at 2 hours' intervals. They were then allowed to grow on the cornmeal medium until they reached the desired age, and they were dissected at 0.5 hour intervals.

Immunostaining

Larvae brain were dissected in phosphate-buffered saline (PBS) and fixed with 4% formaldehyde (PFA) for 25 min at room temperature. After fixation and three washes with 0.3% Triton X-100 in PBS. Brains were blocked with 5% goat serum in PBS with 0.3% Triton X-100 at 4°C 1 hour and

were incubated with primary antibodies in PBS with 0.3% Triton X-100 at 4°C overnight. After

four washes, brains were incubated with secondary antibodies in PBS with 0.3% Triton X-100 at

room temperature for 2-3 h. After four washes, brains were mounted for microscopy in vectashield

without DAPI (Vector Laboratories). Primary antibodies were mouse anti-PDF (1:300), mouse

anti-FasII (1:50), Rabbit anti- HRP (1:500), mouse anti-Dscam1- 18mAb (1:20), rabbit anti-GFP

(1:200), rabbit anti-RFP (1:500), chicken anti-GFP (1:2000), rabbit anti-Myc (1:200), Secondary

antibodies were Alexa Fluor 555 goat anti-Rabbit IgG (1:200), Alexa Fluor 647 goat anti-mouse

IgG (1:200), Alexa Fluor 488 goat anti-rabbit IgG (1:200) and Alexa Fluor 488 goat anti-chicken

IgY (1:200). Samples were imaged on an LSM 700 confocal microscope (Zeiss).

ImageJ software (National Institutes of Health) was used for the quantification vertical and

horizontal axonal length of PDF immunostaining in s-LNvs. For s-LNvs horizontal measurements,

results are presented as A.U. representing the fraction between the lengths of the horizontal

projections divided by the distance between cell bodies and midline. Data are presented as means

± standard deviation from examined brains.

RNAi Screening

The UAS-RNAi line were obtained from Tsinghua Fly Center, and Bloomington Drosophila Stock

Center. The RNAi screen fly was generated as follows: recombine *Pdf-GAL4* with

UAS-mCD8-GFP or use *Clk856-GAL4* directly, which has a broader expression pattern.

UAS-RNAi males were crossed to this two screening lines, and the resulting flies were kept at

25 °C, and dissected at desired age.

Neuron ablation

Tab2-201Y-GAL4 which is expressed in larval mushroom body γ neuron and

per-GAL4,Pdf-GAL80 which is expressed in DN neurons was used to drive the expression of

reaper (rpr) and hid .The GAL4 flies integrated UAS-GFP, and the efficiency of ablation was

confirmed by GFP signals in mushroom body and DN neurons. Animals were raised at 25°C to

desired time.

<u>Quantification and Statistical Analysis</u>

Statistical analysis was performed with GraphPad Prism V8.0.2 software. Data are presented as

means ± standard deviation (SD). Shapiro-Wilk test was used to verify whether the data

conformed to the normal distribution. F test was used to verify homogenous variances between

two groups. Bartlett test was used to verify homogenous variances three or more conditions and

genotypes. Statistical significance was set as: * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$; **** $p <$

0.0001, ns, no significance, $p > 0.05$. Statistical details of the experiments are found in the figure

legends.

Supplemental Information

Supplementary File 1. Key resource table

Supplementary File 2. Full genotypes of the flies that are shown in the main figures, supplemental

figures.

Supplementary File 3. List of genes used for RNAi screen

Reference

Agrawal, P., & Hardin, P. E. (2016). The drosophila receptor protein tyrosine phosphatase LAR is

required for development of circadian pacemaker neuron processes that support rhythmic activity

in constant darkness but not during light/dark cycles. *Journal of Neuroscience*, *36*(13), 3860–

3870. https://doi.org/10.1523/JNEUROSCI.4523-15.2016

Alavi, M., Song, M., King, G. L. A., Gillis, T., Propst, R., Lamanuzzi, M., Bousum, A., Miller, A.,

Allen, R., & Kidd, T. (2016). Dscam1 Forms a Complex with Robo1 and the N-Terminal

Fragment of Slit to Promote the Growth of Longitudinal Axons. *PLoS Biology*, *14*(9), 1–31.

https://doi.org/10.1371/journal.pbio.1002560

Andrews, G. L., Tanglao, S., Farmer, W. T., Morin, S., Brotman, S., Berberoglu, M. A., Price, H.,

Fernandez, G. C., Mastick, G. S., Charron, F., & Kidd, T. (2008). Dscam guides embryonic

axons by Netrin-dependent and -independent functions. *Development*, *135*(23), 3839–3848.

https://doi.org/10.1242/dev.023739

Araújo, S. J., & Tear, G. (2003). Axon guidance mechanisms and molecules: Lessons from

invertebrates. *Nature Reviews Neuroscience*, *4*(11), 910–922. https://doi.org/10.1038/nrn1243

Avalos, C. B., Brugmann, R., & Sprecher, S. G. (2019). Single cell transcriptome atlas of the

drosophila larval brain. *ELife*, *8*, 1–25. https://doi.org/10.7554/eLife.50354

Bak, M., & Fraser, S. E. (2003). Axon fasciculation and differences in midline kinetics between

pioneer and follower axons within commissural fascicles. *Development*, *130*(20), 4999–5008.

https://doi.org/10.1242/dev.00713

Brankatschk, M., & Dickson, B. J. (2006). Netrins guide Drosophila commissural axons at short range.

Nature Neuroscience, *9*(2), 188–194. https://doi.org/10.1038/nn1625

Bron, R., Vermeren, M., Kokot, N., Andrews, W., Little, G. E., Mitchell, K. J., & Cohen, J. (2007).

Boundary cap cells constrain spinal motor neuron somal migration at motor exit   points by a

semaphorin-plexin mechanism. *Neural Development*, *2*, 21.

https://doi.org/10.1186/1749-8104-2-21

Cang, J., & Feldheim, D. A. (2013). Developmental mechanisms of topographic map formation and

alignment. *Annual Review of Neuroscience*, *36*, 51–77.

https://doi.org/10.1146/annurev-neuro-062012-170341

Chan, C. E., & Odde, D. J. (2008). Traction dynamics of filopodia on compliant substrates. *Science*

(New York, N.Y.), *322*(5908), 1687–1691. https://doi.org/10.1126/science.1163595

Chao, D. L., Ma, L., & Shen, K. (2009). Transient cell-cell interactions in neural circuit formation.

Nature Reviews Neuroscience, *10*(4), 262–271. https://doi.org/10.1038/nrn2594

Chédotal, A., Richards, L. J., Dent, E. W., Gupton, S. L., Frank, B., Adams, R. H., Eichmann, A.,

Engle, E. C., Raper, J., & Mason, C. (2012). *Navigating Intermediate Targets : The Nervous*

System Midline Navigating Intermediate Targets : The Nervous System Midline.

https://doi.org/10.1101/cshperspect.a002055

Chen, B. E., Kondo, M., Garnier, A., Watson, F. L., Püettmann-Holgado, R., Lamar, D. R., &

Schmucker, D. (2006). The Molecular Diversity of Dscam Is Functionally Required for Neuronal

Wiring Specificity in Drosophila. *Cell*, *125*(3), 607–620.

https://doi.org/10.1016/j.cell.2006.03.034

Corrales, M., Cocanougher, B. T., Kohn, A. B., Wittenbach, J. D., Long, X. S., Lemire, A., Cardona,

A., Singer, R. H., Moroz, L. L., & Zlatic, M. (2022). A single-cell transcriptomic atlas of

complete insect nervous systems across multiple life stages. *Neural Development*, *17*(1), 8.

https://doi.org/10.1186/s13064-022-00164-6

Cyran, S. A., Yiannoulos, G., Buchsbaum, A. M., Saez, L., Young, M. W., & Blau, J. (2005). The

double-time protein kinase regulates the subcellular localization of the   Drosophila clock protein

period. *The Journal of Neuroscience : The Official Journal of the Society for   Neuroscience*,

25(22), 5430–5437. https://doi.org/10.1523/JNEUROSCI.0263-05.2005

de Ramon Francàs, G., Zuñiga, N. R., & Stoeckli, E. T. (2017). The spinal cord shows the way – How

axons navigate intermediate targets. *Developmental Biology*, *432*(1), 43–52.

https://doi.org/10.1016/j.ydbio.2016.12.002

Derijck, A. A. H. A., Van Erp, S., & Pasterkamp, R. J. (2010). Semaphorin signaling: molecular

switches at the midline. *Trends in Cell Biology*, *20*(9), 568–576.

https://doi.org/10.1016/j.tcb.2010.06.007

Dickson, B. J. (2002). Molecular mechanisms of axon guidance. *Science (New York, N.Y.)*, *298*(5600),

1959–1964. https://doi.org/10.1126/science.1072165

Evans, T. A., & Bashaw, G. J. (2010). *Axon guidance at the midline : of mice and flies*. 79–85.

https://doi.org/10.1016/j.conb.2009.12.006

Fothergill, T., Donahoo, A.-L. S., Douglass, A., Zalucki, O., Yuan, J., Shu, T., Goodhill, G. J., &

Richards, L. J. (2014). Netrin-DCC Signaling Regulates Corpus Callosum Formation Through

Attraction of Pioneering Axons and by Modulating Slit2-Mediated Repulsion. *Cerebral Cortex*,

24(5), 1138–1151. https://doi.org/10.1093/cercor/bhs395

Garel, S., & López-Bendito, G. (2014). Inputs from the thalamocortical system on axon pathfinding

mechanisms. *Current Opinion in Neurobiology*, *27*, 143–150.

https://doi.org/https://doi.org/10.1016/j.conb.2014.03.013

Garel, S., & Rubenstein, J. L. R. (2004). Intermediate targets in formation of topographic projections:

Inputs from the thalamocortical system. *Trends in Neurosciences*, *27*(9), 533–539.

https://doi.org/10.1016/j.tins.2004.06.014

Gezelius, H., & López-Bendito, G. (2017). Thalamic neuronal specification and early circuit formation.

Developmental Neurobiology, *77*(7), 830–843. https://doi.org/https://doi.org/10.1002/dneu.22460

Godement, P., Wang, L. C., & Mason, C. A. (1994). Retinal axon divergence in the optic chiasm:

dynamics of growth cone behavior at the midline [published erratum appears in J Neurosci 1995

Mar;15(3):following table of contents]. *The Journal of Neuroscience*, *14*(11), 7024 LP – 7039.

https://doi.org/10.1523/JNEUROSCI.14-11-07024.1994

Goodhill, G. J. (2016). Can Molecular Gradients Wire the Brain? *Trends in Neurosciences*, *39*(4), 202–

211. https://doi.org/10.1016/j.tins.2016.01.009

Gorla, M., & Bashaw, G. J. (2020). Molecular mechanisms regulating axon responsiveness at the

midline. *Developmental Biology*, *466*(1), 12–21.

https://doi.org/https://doi.org/10.1016/j.ydbio.2020.08.006

Gummadova, J. O., Coutts, G. A., & Glossop, N. R. J. (2009). Analysis of the drosophila clock

promoter reveals heterogeneity in expression between subgroups of central oscillator cells and

identifies a novel enhancer region. *Journal of Biological Rhythms*, *24*(5), 353–367.

https://doi.org/10.1177/0748730409343890

Hardin, P. E. (2017). VRILLE Controls PDF Neuropeptide Accumulation and Arborization Rhythms in

Small Ventrolateral Neurons to Drive Rhythmic Behavior in Drosophila VRILLE Controls PDF

Neuropeptide Accumulation and Arborization Rhythms in Small Ventrolateral Neurons to Drive

R. *Current Biology*, *27*(22), 3442-3453.e4. https://doi.org/10.1016/j.cub.2017.10.010

Harris, R., Sabatelli, L. M., & Seeger, M. A. (1996). Guidance cues at the Drosophila CNS midline:

identification and characterization   of two Drosophila Netrin/UNC-6 homologs. *Neuron*, *17*(2),

217–228. https://doi.org/10.1016/s0896-6273(00)80154-3

Helfrich-Förster, C. (1997). Development of pigment-dispersing hormone-immunoreactive neurons in

the nervous system of Drosophila melanogaster. *Journal of Comparative Neurology*, *380*(3),

335–354.

https://doi.org/10.1002/(SICI)1096-9861(19970414)380:3<335::AID-CNE4>3.3.CO;2-M

Helfrich-Förster, C., Shafer, O. T., Wülbeck, C., Grieshaber, E., Rieger, D., & Taghert, P. (2007).

Development and morphology of the clock-gene-expressing lateral neurons of   Drosophila

melanogaster. *The Journal of Comparative Neurology*, *500*(1), 47–70.

https://doi.org/10.1002/cne.21146

HELFRICH-FÖRSTER, C., WULF, J., & DE BELLE, J. S. (2002). MUSHROOM BODY

INFLUENCE ON LOCOMOTOR ACTIVITY AND CIRCADIAN RHYTHMS IN

DROSOPHILA MELANOGASTER. *Journal of Neurogenetics*, *16*(2), 73–109.

https://doi.org/10.1080/01677060213158

Houl, J. H., Ng, F., Taylor, P., & Hardin, P. E. (2008). CLOCK expression identifies developing

circadian oscillator neurons in the brains   of Drosophila embryos. *BMC Neuroscience*, *9*, 119.

https://doi.org/10.1186/1471-2202-9-119

Hummel, T., Vasconcelos, M. L., Clemens, J. C., Fishilevich, Y., Vosshall, L. B., & Zipursky, S. L.

(2003). Axonal targeting of olfactory receptor neurons in Drosophila is controlled by Dscam.

Neuron, *37*(2), 221–231. https://doi.org/10.1016/S0896-6273(02)01183-2

Hutchins, B. I., Li, L., & Kalil, K. (2011). Wnt/calcium signaling mediates axon growth and guidance

in the developing corpus callosum. *Developmental Neurobiology*, *71*(4), 269–283.

https://doi.org/https://doi.org/10.1002/dneu.20846

Jenett, A., Rubin, G. M., Ngo, T.-T. B., Shepherd, D., Murphy, C., Dionne, H., Pfeiffer, B. D.,

Cavallaro, A., Hall, D., Jeter, J., Iyer, N., Fetter, D., Hausenfluck, J. H., Peng, H., Trautman, E.

T., Svirskas, R. R., Myers, E. W., Iwinski, Z. R., Aso, Y., … Zugates, C. T. (2012). A

GAL4-driver line resource for Drosophila neurobiology. *Cell Reports*, *2*(4), 991–1001.

https://doi.org/10.1016/j.celrep.2012.09.011

Kaas, J. H. (1997). Topographic maps are fundamental to sensory processing. *Brain Research Bulletin*,

44(2), 107–112. https://doi.org/10.1016/s0361-9230(97)00094-4

Kahn, O. I., & Baas, P. W. (2016). Microtubules and Growth Cones: Motors Drive the Turn. *Trends in*

Neurosciences, *39*(7), 433–440. https://doi.org/https://doi.org/10.1016/j.tins.2016.04.009

Kaneko, M., Helfrich-Förster, C., & Hall, J. C. (1997). Spatial and temporal expression of the period

and timeless genes in the developing nervous system of drosophila: Newly identified pacemaker

candidates and novel features of clock gene product cycling. *Journal of Neuroscience*, *17*(17),

6745–6760. https://doi.org/10.1523/jneurosci.17-17-06745.1997

Keeble, T. R., Halford, M. M., Seaman, C., Kee, N., Macheda, M., Anderson, R. B., Stacker, S. A., &

Cooper, H. M. (2006). The Wnt Receptor Ryk Is Required for Wnt5a-Mediated Axon Guidance

on the Contralateral Side of the Corpus Callosum. *The Journal of Neuroscience*, *26*(21), 5840 LP

– 5848. https://doi.org/10.1523/JNEUROSCI.1175-06.2006

Keene, A. C., Mazzoni, E. O., Zhen, J., Younger, M. A., Yamaguchi, S., Blau, J., Desplan, C., &

Sprecher, S. G. (2011). Distinct visual pathways mediate Drosophila larval light avoidance and

circadian   clock entrainment. *The Journal of Neuroscience : The Official Journal of the Society*

for   Neuroscience, *31*(17), 6527–6534. https://doi.org/10.1523/JNEUROSCI.6165-10.2011

Kennedy, T. E., Serafini, T., de la Torre, J. R., & Tessier-Lavigne, M. (1994). Netrins are diffusible

chemotropic factors for commissural axons in the embryonic   spinal cord. *Cell*, *78*(3), 425–435.

https://doi.org/10.1016/0092-8674(94)90421-9

Kidd, T., Brose, K., Mitchell, K. J., Fetter, R. D., Tessier-Lavigne, M., Goodman, C. S., & Tear, G.

(1998). Roundabout controls axon crossing of the CNS midline and defines a novel   subfamily

of evolutionarily conserved guidance receptors. *Cell*, *92*(2), 205–215.

https://doi.org/10.1016/s0092-8674(00)80915-0

Koch, D., Rosoff, W. J., Jiang, J., Geller, H. M., & Urbach, J. S. (2012). Strength in the periphery:

growth cone biomechanics and substrate rigidity   response in peripheral and central nervous

system neurons. *Biophysical Journal*, *102*(3), 452–460. https://doi.org/10.1016/j.bpj.2011.12.025

Kolodkin, A. L., & Tessier-Lavigne, M. (2011). Mechanisms and molecules of neuronal wiring: A

primer. *Cold Spring Harbor Perspectives in Biology*, *3*(6), 1–14.

https://doi.org/10.1101/cshperspect.a001727

Lee, T., Lee, A., & Luo, L. (1999). Development of the Drosophila mushroom bodies: sequential

generation of three   distinct types of neurons from a neuroblast. *Development (Cambridge,*

England), *126*(18), 4065–4076. https://doi.org/10.1242/dev.126.18.4065

Li, T., Fu, T.-M., Wong, K. K. L., Li, H., Xie, Q., Luginbuhl, D. J., Wagner, M. J., Betzig, E., & Luo,

794   L. (2021). Cellular bases of olfactory circuit assembly revealed by systematic time-lapse imaging.

795   *Cell*, *184*(20), 5107-5121.e14. https://doi.org/https://doi.org/10.1016/j.cell.2021.08.030

796 Li, X.-T., Zhou, Q.-S., Yu, Q., Zhao, X., & Liu, Q.-X. (2014). [Current progress in functions of axon

797   guidance molecule Robo and underlying molecular mechanism]. *Sheng li xue bao : [Acta*

798   *physiologica Sinica]*, *66*(3), 373–385.

799 Liu, G., Li, W., Wang, L., Kar, A., Guan, K. L., Rao, Y., & Wu, J. Y. (2009). DSCAM functions as a

800   netrin receptor in commissural axon pathfinding. *Proceedings of the National Academy of*

801   *Sciences of the United States of America*, *106*(8), 2951–2956.

802   https://doi.org/10.1073/pnas.0811083106

803 Liu, T., Mahesh, G., Houl, J. H., & Hardin, P. E. (2015). Circadian activators are expressed days before

804   they initiate clock function in late pacemaker neurons from Drosophila. *Journal of Neuroscience*,

805   *35*(22), 8662–8671. https://doi.org/10.1523/JNEUROSCI.0250-15.2015

806 Long, H., Sabatier, C., Ma, L., Plump, A., Yuan, W., Ornitz, D. M., Tamada, A., Murakami, F.,

807   Goodman, C. S., & Tessier-Lavigne, M. (2004). Conserved roles for Slit and Robo proteins in

808   midline commissural axon guidance. *Neuron*, *42*(2), 213–223.

809   https://doi.org/10.1016/s0896-6273(04)00179-5

810 Lowery, L. A., & Vactor, D. Van. (2009). The trip of the tip: understanding the growth cone machinery.

811   *Nature Reviews Molecular Cell Biology*, *10*(5), 332–343. https://doi.org/10.1038/nrm2679

812 Ly, A., Nikolaev, A., Suresh, G., Zheng, Y., Tessier-Lavigne, M., & Stein, E. (2008). DSCAM Is a

813   Netrin Receptor that Collaborates with DCC in Mediating Turning Responses to Netrin-1. *Cell*,

814   *133*(7), 1241–1254. https://doi.org/10.1016/j.cell.2008.05.030

815 Martins, L. F., Brambilla, I., Motta, A., de Pretis, S., Bhat, G. P., Badaloni, A., Malpighi, C., Amin, N.

816   D., Imai, F., Almeida, R. D., Yoshida, Y., Pfaff, S. L., & Bonanomi, D. (2022). Motor neurons

817   use push-pull signals to direct vascular remodeling critical for their connectivity. *Neuron*,

818   *110*(24), 4090-4107.e11. https://doi.org/10.1016/j.neuron.2022.09.021

819 Matthews, B. J., & Grueber, W. B. (2011). Dscam1-mediated self-avoidance counters netrin-dependent

820   targeting of dendrites in drosophila. *Current Biology*, *21*(17), 1480–1487.

821   https://doi.org/10.1016/j.cub.2011.07.040

822 Mitchell, K. J., Doyle L., J. L., Serafini, T., Kennedy, T. E., Tessier-Lavigne, M., Goodman, C. S., &

823   Dickson, B. J. (1996). Genetic analysis of Netrin genes in Drosophila: Netrins guide CNS

commissural axons and peripheral motor axons. *Neuron*, *17*(2), 203–215.

https://doi.org/10.1016/S0896-6273(00)80153-1

Mortimer, D., Fothergill, T., Pujic, Z., Richards, L. J., & Goodhill, G. J. (2008). Growth cone

chemotaxis. *Trends in Neurosciences*, *31*(2), 90–98.

https://doi.org/https://doi.org/10.1016/j.tins.2007.11.008

Neuhaus-Follini, A., & Bashaw, G. J. (2015). Crossing the embryonic midline: molecular mechanisms

regulating axon responsiveness at an intermediate target. *WIREs Developmental Biology*, *4*(4),

377–389. https://doi.org/https://doi.org/10.1002/wdev.185

Okray, Z., de Esch, C. E. F., Van Esch, H., Devriendt, K., Claeys, A., Yan, J., Verbeeck, J., Froyen, G.,

Willemsen, R., de Vrij, F. M. S., & Hassan, B. A. (2015). A novel fragile X syndrome mutation

reveals a conserved role for the   carboxy-terminus in FMRP localization and function. *EMBO*

Molecular Medicine, *7*(4), 423–437. https://doi.org/10.15252/emmm.201404576

Oliva, C., Soldano, A., Mora, N., De Geest, N., Claeys, A., Erfurth, M. L., Sierralta, J., Ramaekers, A.,

Dascenco, D., Ejsmont, R. K., Schmucker, D., Sanchez-Soriano, N., & Hassan, B. A. (2016).

Regulation of Drosophila Brain Wiring by Neuropil Interactions via a Slit-Robo-RPTP Signaling

Complex. *Developmental Cell*, *39*(2), 267–278. https://doi.org/10.1016/j.devcel.2016.09.028

Orioli, D., & Klein, R. (1997). The Eph receptor family: axonal guidance by contact repulsion. *Trends*

in Genetics : TIG, *13*(9), 354–359. https://doi.org/10.1016/s0168-9525(97)01220-1

Pauls, D., Selcho, M., Gendre, N., Stocker, R. F., & Thum, A. S. (2010). Drosophila larvae establish

appetitive olfactory memories via mushroom body neurons of embryonic origin. *Journal of*

Neuroscience, *30*(32), 10655–10666. https://doi.org/10.1523/JNEUROSCI.1281-10.2010

Piper, M., Plachez, C., Zalucki, O., Fothergill, T., Goudreau, G., Erzurumlu, R., Gu, C., & Richards, L.

J. (2009). Neuropilin 1-Sema Signaling Regulates Crossing of Cingulate Pioneering Axons

during Development of the Corpus Callosum. *Cerebral Cortex*, *19*(suppl_1), i11–i21.

https://doi.org/10.1093/cercor/bhp027

Pirooznia, S. K., Chiu, K., Chan, M. T., Zimmerman, J. E., & Elefant, F. (2012). Epigenetic Regulation

of Axonal Growth of Drosophila Pacemaker Cells by Histone Acetyltransferase Tip60 Controls

Sleep. *Genetics*, *192*(4), 1327–1345. https://doi.org/10.1534/genetics.112.144667

Puñal, V. M., Ahmed, M., Thornton-Kolbe, E. M., & Clowney, E. J. (2021). Untangling the wires:

development of sparse, distributed connectivity in the mushroom body calyx. *Cell and Tissue*

Research, *383*(1), 91–112. https://doi.org/10.1007/s00441-020-03386-4

Purohit, A. A., Li, W., Qu, C., Dwyer, T., Shao, Q., Guan, K. L., & Liu, G. (2012). Down syndrome

cell adhesion molecule (DSCAM) associates with uncoordinated-5C (UNC5C) in

netrin-1-mediated growth cone collapse. *Journal of Biological Chemistry*, *287*(32), 27126–27138.

https://doi.org/10.1074/jbc.M112.340174

Reinhard, N., Schubert, F. K., Bertolini, E., Hagedorn, N., Manoli, G., Sekiguchi, M., Yoshii, T.,

Rieger, D., & Helfrich-Förster, C. (2022). The Neuronal Circuit of the Dorsal Circadian Clock

Neurons in Drosophila melanogaster. *Frontiers in Physiology*, *13*, 886432.

https://doi.org/10.3389/fphys.2022.886432

Schlichting, M., Richhariya, S., Herndon, N., Ma, D., Xin, J., Lenh, W., Abruzzi, K., & Rosbash, M.

(2022). Dopamine and GPCR-mediated modulation of DN1 clock neurons gates the circadian

timing of sleep. *Proceedings of the National Academy of Sciences of the United States of*

America, *119*(34), e2206066119. https://doi.org/10.1073/pnas.2206066119

Schmucker, D., Clemens, J. C., Shu, H., Worby, C. A., Xiao, J., Muda, M., Dixon, J. E., & Zipursky, S.

L. (2000). Drosophila Dscam is an axon guidance receptor exhibiting extraordinary molecular

diversity. *Cell*, *101*(6), 671–684. https://doi.org/10.1016/s0092-8674(00)80878-8

Serafini, T., Colamarino, S. A., Leonardo, E. D., Wang, H., Beddington, R., Skarnes, W. C., &

Tessier-Lavigne, M. (1996). Netrin-1 is required for commissural axon guidance in the

developing vertebrate nervous system. *Cell*, *87*(6), 1001–1014.

https://doi.org/10.1016/s0092-8674(00)81795-x

Shields, A. R., Spence, A. C., Yamashita, Y. M., Davies, E. L., & Fuller, M. T. (2014). The

actin-binding protein profilin is required for germline stem cell maintenance and germ cell

enclosure by somatic cyst cells. *Development*, *141*(1), 73–82. https://doi.org/10.1242/dev.101931

Squarzoni, P., Thion, M. S., & Garel, S. (2015). Neuronal and microglial regulators of cortical wiring:

usual and novel guideposts. *Frontiers in Neuroscience*, *9*, 248.

https://doi.org/10.3389/fnins.2015.00248

Stoeckli, E T, Sonderegger, P., Pollerberg, G. E., & Landmesser, L. T. (1997). Interference with

axonin-1 and NrCAM interactions unmasks a floor-plate activity inhibitory for commissural

axons. *Neuron*, *18*(2), 209–221. https://doi.org/10.1016/s0896-6273(00)80262-7

Stoeckli, Esther T. (2018). Understanding axon guidance: are we nearly there yet? *Development*,

145(10), dev151415. https://doi.org/10.1242/dev.151415

Sudarsanam, S., Yaniv, S., Meltzer, H., & Schuldiner, O. (2020). Cofilin regulates axon growth and

branching of Drosophila γ-neurons. *Journal of Cell Science*, *133*(8).

https://doi.org/10.1242/jcs.232595

Suzuki, Y., Kurata, Y., & Sakai, T. (2022). Dorsal–lateral clock neurons modulate consolidation and

maintenance of long-term memory in Drosophila. *Genes to Cells*, *27*(4), 266–279.

https://doi.org/https://doi.org/10.1111/gtc.12923

Unni, D. K., Piper, M., Moldrich, R. X., Gobius, I., Liu, S., Fothergill, T., Donahoo, A.-L. S., Baisden,

J. M., Cooper, H. M., & Richards, L. J. (2012). Multiple Slits regulate the development of

midline glial populations and the corpus callosum. *Developmental Biology*, *365*(1), 36–49.

https://doi.org/https://doi.org/10.1016/j.ydbio.2012.02.004

Vitriol, E. A., & Zheng, J. Q. (2012). Growth cone travel in space and time: the cellular ensemble of

cytoskeleton,   adhesion, and membrane. *Neuron*, *73*(6), 1068–1081.

https://doi.org/10.1016/j.neuron.2012.03.005

Worby, C. A., Simonson-Leff, N., Clemens, J. C., Kruger, R. P., Muda, M., & Dixon, J. E. (2001). The

sorting nexin, DSH3PX1, connects the axonal guidance receptor, Dscam, to the   actin

cytoskeleton. *The Journal of Biological Chemistry*, *276*(45), 41782–41789.

https://doi.org/10.1074/jbc.M107080200

Yang, L., Garbe, D. S., & Bashaw, G. J. (2009). A frazzled/DCC-dependent transcriptional switch

regulates midline axon guidance. *Science (New York, N.Y.)*, *324*(5929), 944–947.

https://doi.org/10.1126/science.1171320

Zang, Y., Chaudhari, K., & Bashaw, G. J. (2021). New insights into the molecular mechanisms of axon

guidance receptor regulation   and signaling. *Current Topics in Developmental Biology*, *142*,

147–196. https://doi.org/10.1016/bs.ctdb.2020.11.008

Zhan, X. L., Clemens, J. C., Neves, G., Hattori, D., Flanagan, J. J., Hummel, T., Vasconcelos, M. L.,

Chess, A., & Zipursky, S. L. (2004). Analysis of Dscam diversity in regulating axon guidance in

Drosophila mushroom bodies. *Neuron*, *43*(5), 673–686.

https://doi.org/10.1016/j.neuron.2004.07.020

Zhang, Z., So, K., Peterson, R., Bauer, M., Ng, H., Zhang, Y., Kim, J. H., Kidd, T., & Miura, P. (2019).

Elav-Mediated Exon Skipping and Alternative Polyadenylation of the Dscam1 Gene Are

Required for Axon Outgrowth. *Cell Reports*, *27*(13), 3808-3817.e7.

https://doi.org/10.1016/j.celrep.2019.05.083

Figure 1

Figure 2

A

	24 h	48 h	72 h	96 h	120 h

OK107-GAL4 >GFP
PDF
GFP
20 µm

201Y-GAL4>GFP
PDF
GFP
20 µm

B

OK107
γ,α',β' neuron

201Y
γ neuron

r = 0.9987 P < 0.001
r = 0.9963 P = 0.0003

OK107 · 201Y · Vertical

MB calyx area (µm²) / Vertical Distance (µm)

C

Vertical projection
MB calyx
s-LNvs
24 h · 48 h · 72 h · 96 h · 120 h
D M L V

D

	24 h	48 h	72 h

201Y >GFP
PDF
HRP
20 µm

201Y >GFP ,rpr,hid

E

201Y >GFP
201Y >GFP,rpr,hid

Vertical Distance (µm)

ns **** ****

24 48 72 (h)

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7