FLOWERING LOCUS C integrates carbon and nitrogen signaling for the proper timing of flowering in Arabidopsis

Vladislav Gramma[1‡#], Justyna Jadwiga Olas[1‡#], Vasiliki Zacharaki[2#], Jathish Ponnu[3], Urszula Marta Luzarowska[1], Magdalena Musialak-Lange[1‡], and Vanessa Wahl[1,4*]

[1]Max Planck Institute of Molecular Plant Physiology, Department Metabolic Networks, Potsdam, Germany.

[2]Umeå Plant Science Centre, Department of Plant Physiology, Umeå University, Umeå, Sweden.

[3]Joseph Gottlieb Kölreuter Institute for Plant Sciences (JKIP), Karlsruhe Institute of Technology (KIT), Germany.

[4]The James Hutton Institute, Department of Cell and Molecular Sciences, Dundee, UK.

*Corresponding author/author responsible for distribution of materials

#These three authors (in alphabetical order) contributed equally.

‡**Current affiliations:** University of Applied Sciences Berlin (VG); Leibniz Institute of Vegetable and Ornamental Crop, Großbeeren, Germany (JJO); Metasysx GmbH, Am Mühlenberg 11, Potsdam, Germany (MML)

Author emails and ORCID:

Vladislav.Gramma@HTW-Berlin.de (0000-0002-9890-3042)

Olas@igzev.de (0000-0002-4311-6738)

Vasiliki.Zacharaki@umu.se (0000-0002-5543-2332)

Urszula.Luzarowska@gmail.com (0000-0002-3350-5297)

Jathish.Ponnu@kit.edu (0000-0002-3276-7068)

musialak-lange@metasysx.com (0000-0002-0388-8960)

Vanessa.Wahl@hutton.ac.uk (0000-0001-7421-8801)

Short title: C and N signaling regulate FLC.

Abstract

The timing of flowering in plants is modulated by both carbon (C) and nitrogen

(N) signaling pathways. In a previous study, we established a pivotal role of the

sucrose-signaling trehalose 6-phosphate pathway in regulating flowering under

N-limited short-day conditions. In this work, we expand on our finding that wild-

type plants grown under N-limited short days require an active trehalose 6-

phosphate pathway to be able to flower. Both wild-type plants grown under N-

limited conditions and knock-down plants of *TREHALOSE PHOSPHATE*

SYNTHASE1 induce *FLOWERING LOCUS C* expression, a well-known floral

repressor associated with the vernalization response. When exposed to an

extended period of cold, a mutant of *FLOWERING LOCUS C* fails to respond

to N availability, and flowers at the same time under N-limited and full-nutrition

conditions. Our data suggest that SUCROSE NON-FERMENTING 1 RELATED

KINASE 1-dependent trehalose 6-phosphate-mediated C signaling and a novel

mechanism downstream of N signaling likely involving NIN-LIKE PROTEIN 7

impact the expression of *FLOWERING LOCUS C.* Collectively, our data

underscore the existence of a multi-factor regulatory system in which both C

and N signaling pathways jointly govern the regulation of flowering in plants.

Keywords: TREHALOSE PHOSPHATE SYNTHASE (TPS1), Trehalose 6-

phosphate (T6P), FLOWERING LOCUS C (FLC), nitrogen, N signaling,

flowering time

Introduction

Owing to their sessile nature, plants adapt to environmental changes by modifying their development and growth. These processes require significant amounts of energy. Plants are in constant feedback with the environment and their nutrient status, especially carbon (C) and nitrogen (N), that serve as crucial bases for energy production and biomass generation. Low levels of C or N in the cells suppresses development and growth in plants and triggers the onset of senescence. To balance energy-intensive developmental processes with endogenous nutrient availability, plants have evolved intricate signaling networks (Fernie et al., 2020).

Flowering is an important developmental process in the life cycle of plants with correct timing being essential for reproductive success. It is regulated by a sophisticated genetic network that integrates various environmental and endogenous signals to regulate the expression of the floral integrator genes such as the florigen, *FLOWERING LOCUS T* (*FT*), and *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1* (*SOC1*) (Srikanth and Schmid, 2011; Romera-Branchat et al., 2014; Song et al., 2015). FT integrates signals perceived in the leaves and conveys this information to the shoot apical meristem (SAM) to induce flowering (Corbesier et al., 2007; Jaeger and Wigge, 2007; Mathieu et al., 2007). At the SAM, FT interacts with the bZIP transcription factor FLOWERING LOCUS D (FD) to form a complex that directly activates *SOC1* along with floral meristem identity genes such as *APETALA1* (*AP1*) (Abe et al., 2005; Wigge et al., 2005).

In addition to other stimuli, temperature impacts greatly the time of flowering. Increased ambient temperature results in earlier flowering due to decreased SVP protein stability (Lee et al., 2013; Lee et al., 2014). SVP forms a temperature-dependent flowering repressor complex with partners such as FLOWERING LOCUS M (FLM)/MADS AFFECTING FLOWERING1 (MAF1), an orthologue of FLOWERING LOCUS C (FLC) (Pose et al., 2013; Sureshkumar et al., 2016), resulting in earlier flowering when plants are exposed to warmer conditions (Pose et al., 2013). SVP was also shown to interact with FLC in a flowering repressor complex (Fujiwara et al., 2008; Li et al., 2008). This delays floral transition by directly reducing the expression of *FT*,

FD, and *SOC1* (Hepworth et al., 2002; Helliwell et al., 2006; Searle et al., 2006;

Lee et al., 2007; Li et al., 2008). In winter-annual accessions of *Arabidopsis*

thaliana (Arabidopsis) flowering is suppressed due to active *FRIGIDA* (*FRI*)

resulting in promoted expression of *FLC*, unless the plants are exposed to a

long period of cold (vernalization process) (Sheldon et al., 2000). This

regulation involves a plethora of proteins and complexes acting in many layers

of gene regulation, ranging from RNA structures, epigenetic modification to

transcriptional and mRNA processing control (reviewed in Whittaker and Dean,

2017; and Sharma et al., 2020; Xu et al., 2021; Xu et al., 2021; Yang et al.,

2022).

Organic C and N supply is essential in particular for vegetative growth and plant

development (Sulpice et al., 2013). It is known that nutrients are essential for

developmental transitions (Fernie et al., 2020), but the underlying mechanisms

continue to be subject to active investigation. Interestingly, *FLC* expression was

observed to increase significantly in NITRATE TRANSPORTER 1.1 (NRT1.1)

and NRT1.13 defective mutant plants (Teng et al., 2019; Chen et al., 2021).

While NRT1.13 is suggested to be a nitrate transporter, NRT1.1 is a key

component of nitrate signaling functioning as both a transporter and a sensor

in roots (Li et al., 2021). This suggests a nitrate signaling-dependent control of

FLC as proposed by Kant and colleagues (Kant et al., 2011). This is supported

by the introduction of an *flc-3* mutation into the late-flowering NRT1.1 deficient

plant background which restored wild-type flowering (Teng et al., 2019).

Previous studies have identified multiple factors that influence N-regulated

flowering, which often vary and depend on the cultivation systems used (Lin

and Tsay, 2017). We are using a soil-based N-limited system developed by

(Tschoep et al., 2009), which allows plant adaptation and the investigation of

flowering time without stress-related symptoms (Olas et al., 2019; Olas et al.,

2021). With this system we previously reported that nitrate-regulated flowering

depends on SAM factors. Notably, in N-limiting conditions, nitrate-responsive

gene expression is affected and nitrate assimilation is reduced in the SAM (Olas

et al., 2019). The early nitrate response involves the NIN-LIKE PROTEIN (NLP)

transcription factors NLP6 and NLP7. They accumulate in the nucleus in the

presence of nitrate, regulating gene expression through nitrate responsive cis-

elements (NRE) (Konishi and Yanagisawa, 2013; Marchive et al., 2013).
Limited nitrate availability delays flowering due to decreased expression of
SOC1, likely through NLP6/NLP7-regulated expression of the SQUAMOSA
PROMOTER-BINDING PROTEIN-LIKE transcription factors encoding genes
SPL3 and *SPL5* (Olas et al., 2019).

The sucrose signal trehalose 6-phosphate (T6P) regulates a plethora of
developmental and physiological responses (reviewed in Fichtner and Lunn,
2021). In Arabidopsis*,* T6P is synthesized by TREHALOSE PHOSPHATE
SYNTHASE1 (TPS1) (Vandesteene et al., 2010; Yang et al., 2012) and it acts
mainly by modulating the SUCROSE NON-FERMENTING 1 RELATED
KINASE 1 (SnRK1) activity. Moreover, T6P was suggested to be able to bind
directly to the SnRK1 upstream activating kinases and inhibit their activity (Zhai
et al., 2018). SnRK1 is a key sensor of energy status and it is required for both
normal growth and plant responses to stresses that impact plant fitness and
survival (Polge and Thomas, 2007; Baena-Gonzalez and Sheen, 2008).
Although single mutants of SnRK1 catalytic subunits resemble wild-type plants
(Baena-Gonzalez et al., 2007; Jeong et al., 2015), *tps1* mutants (*tps1-2*) are
embryo-lethal (Eastmond et al., 2002). This can be bypassed by ectopically
expressing dexamethasone-inducible *TPS1* (*GVG::TPS1*) during seed set (van
Dijken et al., 2004). However, plants grown from these seeds remain in the
vegetative phase for a highly extended period or fail to flower entirely (van
Dijken et al., 2004; Wahl et al., 2013). T6P signaling induces flowering in leaves
via *FT* and also acts at the SAM through microRNA156 (miR156) and its target
transcripts, *SPL3-5* (Wahl et al., 2013), at least partially via the modulation of
the SUCROSE NON-FERMENTING 1 RELATED KINASE 1 (SnRK1) complex
activity (Zacharaki et al., 2022). This was supported by the observation that loss
of SnRK1 activity in the *tps1-2,GVG::TPS1* plants led to early induction of *FT*
in the leaves, reduced *miR156* levels and strong induction of *SPL3* in the SAM
during bolting (Zacharaki et al., 2022). Taken together these findings indicate
that both C and N signaling can target the same components of the flowering
network at the SAM (Wahl et al., 2013; Olas et al., 2019; Zacharaki et al., 2022),
underscoring their joint importance for the proper timing of flowering.

Even though the current understanding implies a straightforward output
downstream of nutrient signaling, our data now indicate a more complex

relationship between nutrient signaling and developmental programs. Here, we

demonstrate that the T6P pathway, which controls flowering under N limitation

in short days (Olas et al., 2019), impacts on the expression of *FLC* in addition

to *FLC* being differentially expressed upon exposure to contrasting N levels.

Our findings suggest that both C- and N-dependent pathways regulate

Arabidopsis flowering time by modulating *FLC* expression, implying a role in

the composition and timing of the FLC-SVP repressor complex within a

developmental context.

Results

Sucrose signaling represses *FLC*

We have previously reported that plants grown under N-limited conditions and short days (SD) accumulate both sucrose and T6P towards the end of the vegetative growth phase. Importantly, *TPS1* knock-down plants (*35S::amiRTPS1*) did not flower under these conditions (Olas et al., 2019). To understand this phenomenon, we analyzed a developmental series of rosette samples from both Col-0 and *35S::amiRTPS1* plants, focusing on candidate genes, which specifically change their expression before the floral transition. This analysis included multiple flowering time genes assessed by RT-qPCR. Notably, we found a strong up-regulation of *FLC* expression in 4- to 6-days-old and *MAF5* (*MADS AFFECTING FLOWERING 5*) expression in 12-days-old *35S::amiRTPS1* plants (Fig. 1A, Fig. S1), similar to Zeng et al. (2024). Considering that *FLC* expression has previously been suggested to be modulated in response to N availability (Kant et al., 2011), it is an interesting candidate for further investigation. We observed that *FLC* expression declines before the floral transition, which occurs at 10 days after germination (DAG) in Col-0 wild-type plants and 19 DAG in *35S::amiRTPS1* grown in full nutrition soil (Wahl et al., 2013). This suggests that the T6P pathway fundamentally contributes to the full repression of *FLC* in young seedlings. While we initially did not anticipate that the T6P pathway could affect *FLC* expression at later stages, we found that when the *flc-3* mutation is introduced into the *tps1-2 GVG::TPS1* background, it partially rescues the late flowering and the delayed vegetative phase transition observed in this *tps1-2 GVG::TPS1* (Fig. 1B,C; Fig. S2; Table S1,2). Our data, therefore, suggest that the T6P pathway is involved in *FLC* regulation to promote flowering and facilitate the vegetative phase change.

FLC integrates N-signaling into the flowering network

It has been previously shown that flowering is delayed in wild-type plants grown in the limited N (LN) soil (Olas et al., 2019). Furthermore, some data suggest that *FLC* expression may be influenced by N availability (Kant et al., 2011). Thus, we conducted experiments to investigate the potential regulation of *FLC* expression by N status. We grew wild-type Col-0 plants in a soil-based growth

system (Tschoep et al., 2009), consisting of a soil with optimal N (ON) and one

with LN source. We observed elevated *FLC* expression levels in LN in both

rosettes and apices of Col-0 plants grown continuously in SD conditions (Fig.

2A, B), and in apices of plants that were initially grown in SD conditions and

subsequently transferred to LD conditions (Fig. 2C). Similarly, we found

upregulation of *MAF5* levels in rosettes of LN grown plants (Fig. S3).

Considering that the *MAF5* was found to act downstream of the T6P pathway,

this finding suggests a link between N and T6P signaling pathways. (Fig. S1),

MAF5 was also found to be upregulated in response to N limitation (Fig. S3).

Next, to obtain information on the expression pattern at higher spatial

resolution, we used *FLC* as a probe and performed RNA *in situ* hybridization

(Fig. 2D). *FLC* transcript was detectable at the SAM and in young leaves of LN

grown plants, confirming our previous observations that limited N availability

enhances *FLC* expression in plants. This finding suggests that FLC plays a role

in the regulation of flowering time in response to N availability.

It is well established that exposure to low temperatures decreases *FLC*

expression in plants (Searle et al., 2006). For this reason, we grew wild-type

plants at 4°C in SD for 8 weeks, followed by a transfer to 22°C until flowering.

This treatment resulted in wild-type plants flowering at the same time in both N

regimes, suggesting that FLC contributes to the delayed flowering time

observed in plants grown in the LN soil (Fig. 3A; Table S1).

FLC is known to form a flowering repressor complex with SVP to suppress

SOC1 at the SAM (Li et al., 2008). Unlike *FLC*, *SVP* was not differentially

expressed in either LN-grown plants or *TPS1* knock-down plants (Fig. S4; Fig.

S5). Importantly, neither *flc-3* nor *svp-32* mutant plants responded to the

reduced N content in the LN soil (Fig. 3B; Table S1), flowering at the same time

in ON and LN conditions. This indicates that both FLC and SVP play a role in

the N-dependent regulation of flowering time.

N-signaling affects *FLC* via NLP7

NLPs are key regulators of nitrate sensing and signaling, with NLP6 and NLP7

being two of the most well-characterized members of this family in *Arabidopsis*

(Fredes et al., 2019). In the presence of nitrate, NLP7 is retained in the nucleus

through phosphorylation, where it binds to NREs present in N-responsive
genes to promote their expression (Konishi and Yanagisawa, 2013).
Interestingly, we observed a significant reduction of *FLC* expression in the *nlp7-*
1 mutant, indicating that an active NLP7 modulates *FLC* expression when N is
not limited (Fig. 4).

Since the *FLC* gene does not carry an NRE in its promoter, genomic or
downstream sequences, we expanded our analysis to include other flowering
time genes that regulate *FLC* (Table S3). Notably, *FRI*, a key regulator
upstream of *FLC,* has four putative NREs (Table S3). However, in the Col-0
background, the *FRI* locus encodes an inactive protein and therefore does not
influence *FLC*. In addition to *FRI*, other genomic loci encoding *FLC* regulators
were also found with putative NREs (Table S3), but their expression was
unaffected under N limited conditions (Fig. S6). This was also the case in
35S::amiRTPS1 plants (Fig. S7). Taken together, this suggests that *FLC*
suppression involves an as yet unknown transcription factor(s), whose activity
is regulated by NLP7.

Sucrose and N-signals interconnect at the level of *FLC* for coordinated
flowering time regulation
We have previously demonstrated that the T6P pathway and sufficient nitrate
levels are necessary for floral induction in SD (Olas et al., 2019). The fact that
35S::amiRTPS1 plants fail to flower when N is limited and that *FLC* expression
is modulated by N availability, prompted us to test whether *FLC* is a target of
both N signaling and the T6P pathway.

We observed that *FLC* transcription was elevated in rosettes of wild-type plants
grown under SD with limited N which was even more pronounced in
35S::amiRTPS1 plants (Fig. 5A). This suggests an additive effect between N
signaling and the T6P pathway, both converging on the *SPL3-5* node at the
SAM (Wahl et al., 2013; Olas et al., 2019). To test whether *FLC* could be
regulated through SPL3-5, we measured its expression in *spl345* mutants (Xie
et al., 2020). However, *FLC* expression in rosette leaves of *spl345* mutants was
comparable to that of wild-type plants (Fig. S8A), indicating that both pathways
regulate *FLC* expression via another mechanism. Similarly to *FLC*, we did not

observe any difference in *SVP* expression in *spl345* compared to Col-0 plants
(Fig. S8B).

The T6P pathway is known to function by directly modulating SnRK1 activity
(Zhang et al., 2009). Loss of SnRK1 activity restores flowering of *tps1* mutants
in LD by initial induction of *FT* in the leaves and subsequent suppression of
miR156 followed by SPLs induction in the SAM (Zacharaki et al., 2022). Thus,
we tested whether *FLC* regulation in *tps1* mutants is also mediated by SnRK1.
We found that indeed *FLC* expression was increased in the *tps1 GVG::TPS1*
mutant where SnRK1 is fully active. Interestingly, introducing non-catalytically
active mutations in SnRK1 within the *tps1 GVG::TPS1* background restores
FLC expression to wild-type levels in both rosette leaves and apex tissue (Fig.
5B,C). The suppression of *FLC* in the double mutant is more pronounced in the
apex than the rosette leaves, underscoring the critical role of the T6P pathway
in controlling developmental transitions. Our data suggest that *FLC* expression
is regulated by both nitrate and sugar availability via NPLs and the T6P pathway
through the SnRK1 complex, respectively (Fig. 6).

Discussion
C and N are essential for plant growth and development and the ability of plants
to properly sense their availability is crucial due to their sessile nature. C in the
form of sucrose is produced via photosynthesis in the leaves while N can be
taken up in both inorganic forms, as nitrate and ammonia, or organic forms, as
amino acids.

In *Arabidopsis*, a key sugar sensor is the T6P pathway which functions via
SnRK1 activity. The T6P pathway has a key role in plants' developmental
transitions, such as flowering. So far, it has been shown that both *miR156* and
FT regulation in the SAM and leaves, respectively, are required for *tps1* plants
to complete their transition to flowering (Wahl et al., 2013; Ponnu et al., 2020).
Here we found that *FLC,* a repressor of flowering, is also regulated by the T6P
pathway (Fig. 1A) and that loss of functional *FLC* partially restores flowering in
tps1 (Fig. 1B,C). Although, we do not expect that *FLC* regulation is the prime
target of the T6P pathway under normal growth conditions, it could represent
an additional mechanism to prevent flowering under non-optimal growth
conditions.

Plants experiencing a sudden shift to colder temperatures have increased
amounts sucrose previously proposed to serve as a freezing protectant and
concomitant rising T6P levels (reviewed in Stitt and Hurry, 2002; Carillo et al.,
2013). During long cold exposure, *FLC* is suppressed and this regulation
involves several mechanisms, ranging from RNA structures to epigenetic
control. *FLC* suppression allows induction of *FT* and *SOC1* and flowering to
commence (Whittaker and Dean, 2017). In this scenario, when plants
experience cooler temperatures, nutrients that provide plants with C and N, are
transported and stored to serve as a basis for rapid growth for when conditions
are optimal again or used and metabolically transformed into cryoprotectants
to protect the cells from freezing damage (Kaplan et al., 2007). Thus, in sub-
optimal growth conditions, the T6P pathway might contribute to the suppression
of *FLC* in response to the C status.
N availability is a key factor in the regulation of plants' developmental processes
and phase transitions including the timing of flowering (Klebs, 1913; Dickens
and Van Staden, 1988; Bernier et al., 1993; Olas et al., 2019). Arabidopsis
cultivated on synthetic substrates exhibit early flowering in response to reduced
N levels (Castro Marin et al., 2011; Kant et al., 2011; Liu et al., 2013).
Conversely, soil-grown plants subjected to N limitation flower later than those
cultivated in soil without N limitation, which we previously linked to the induction
of *SPL3* and *SPL5* by NLP6 and NLP7 (Olas et al., 2019). In this study we
discovered that this phenotype can additionally be explained by significantly
elevated levels of *FLC* under N limitation (Fig. 2A-D). Furthermore, we found
that flowering time in plants with suppressed *FLC* due to the vernalization
response or with a non-functional *flc-3* allele is independent of N availability
(Fig. 3A,B). These results demonstrate that despite the general belief that *FLC*
does not play a major role in the regulation of flowering time in rapid-cycling
accessions, such as Col-0, *FLC* is required for fine-tuning the timing of floral
transition downstream of N signaling. Similar to *flc-3*, *svp-32* mutants flower at
the same time in the ON and LN soils (Fig. 3B), suggesting a role of *SVP* in N-
dependent flowering time regulation. However, in contrast to *FLC*, *SVP* is not
differentially expressed in plants grown in ON and LN soil (Fig. S5). FLC and
SVP proteins form a flowering repressor complex that delays floral transition by
directly reducing the expression of *FT* and *SOC1* (Hepworth et al., 2002;

Helliwell et al., 2006; Lee et al., 2007; Li et al., 2008). Given that both functional

FLC and *SVP* loci are required for the adjustment of flowering time in response

to N availability, it is likely that the N signal is integrated at the level of the FLC-

SVP complex. In this scenario, the formation of the repressor complex would

be tuned by the adjustment of *FLC* expression downstream of N-signaling.

Several transcription factors that are transcriptionally responsive to the N status

have been identified as prime responsive genes to N availability (Vidal et al.,

2015). NLPs are transcription factors facilitating nitrate signaling in plants, with

NLP6 and NLP7 representing the master regulators and the two most studied

(Fredes et al., 2019). In the absence of nitrate, NLP7 localizes strictly to the

cytosol, while exposure to nitrate triggers its localization into the nucleus where

it binds directly to NREs of nitrate-regulated genes (Konishi and Yanagisawa,

2013; Marchive et al., 2013). Since NREs are not present in the *FLC* locus

(Table S3), it is unlikely to be directly controlled by NLPs. Other examples of

FLC regulation related to N availability, are the *nrt1.1* and *nrt1.13*, mutants of

the nitrate sensor and transporter NRT1.1 and transporter NRT1.13 (Teng et

al., 2019; Chen et al., 2021). Similar to our findings (Fig. S7), expression of

known upstream regulators of *FLC* was not changed in *nrt1.13*, suggesting that

NRT1.13 regulates *FLC* expression and flowering time independently of these

known pathways (Chen et al., 2021).

Interestingly, we found that *FLC* was significantly downregulated in the late

flowering *nlp7-1* and *nlp6-2 nlp7-1* mutants grown on standard soil (Fig. 4A,

Fig. S9) indicating that NLP7 plays a role in the modulation of *FLC* expression.

Given the fact that NLP7 was found to control most of the nitrate-related gene

response (Marchive et al., 2013; Alvarez et al., 2020), the *nlp7-1* mutant is

thought to mimic a low nitrate state. Hence, this result appears to contradict our

observation of *FLC* accumulation in LN-grown plants (Fig. 2A,B). This could be

explained by the presence of an unknown NLP-independent mechanism

responsible for *FLC* upregulation in LN conditions. However, it should be noted

that in contrast to the mutant background, functional NLP7 is still present in

wild-type plants exposed to limited N. Thus, *nlp7-1* might not entirely mimic the

low-nitrate state after all and the absence of a functional NLP7 likely leads to

compensation by other NLPs. Furthermore, NLP proteins contain a PB1

domain, which mediates protein-protein interactions influencing NLP activity

(Konishi and Yanagisawa, 2019). Given this, NLP7 might form a complex with
an unknown *FLC* repressor, thereby preventing its nuclear localization under
low-nitrate conditions. In the absence of NLP7 or when plants are grown under
optimal N conditions, this potential repressor would localize into the nucleus,
leading to a repression of *FLC* expression. It will be interesting to further dissect
the mechanisms of *FLC* regulation downstream of N-signaling in the future.

Our data demonstrate that both the T6P and N-signaling pathways possibly
affect *FLC* expression via different mechanisms. Previous studies have
demonstrated that both pathways act via the miR156/SPLs node (Wahl et al.,
2013; Olas et al., 2019; Ponnu et al., 2020; Zacharaki et al., 2022). In particular,
the expression of *SPL3* and *SPL5* is reduced in plants grown in N-limited
environment (Bi et al., 2007; Pant et al., 2009; Krapp et al., 2011; Liang et al.,
2012; Fischer et al., 2013), suggesting a role for the miR156/SPL3/5 module in
the regulation of flowering time when N is limited. Similarly, the T6P pathway
acts via miR156 downregulation and SPL3-5 upregulation to induce flowering
and the vegetative phase change (Wahl et al., 2013; Ponnu et al., 2020;
Zacharaki et al., 2022). Although both pathways converge on the miR156/SPLs
module, *FLC* regulation seems to be independent (Fig. S8).

T6P has a key role in promoting growth and development by suppressing
SnRK1 complex activity, via direct binding to the SnRK1 upstream kinases
(Zhai et al., 2018). In a previous study, it was shown that *FT* was induced in the
double *tps1-2 GVG::TPS1 kin10-5* and *tps1-2 GVG::TPS1 snf4* mutant as early
as in wild-type plants (Zacharaki et al., 2022). Although this early *FT* induction
promoted the floral transition in wild-type plants within a few days, this was not
the case in both double mutants. The elevated expression of *FLC* in rosette
leaves of these mutants (Fig. 5C) could thus at least partially explain this
phenomenon. *FLC* downregulation is directly correlated with early *FT*
upregulation previously observed in the double mutants (Zacharaki et al.,
2022). Interestingly, we observed that *FLC* was also downregulated in the
double mutants in the apex (Fig. 5B) with more striking differences later on,
coinciding with the timing of floral transition (Zacharaki et al., 2022). In addition,
ectopic *FLC* expression in the SAM has been associated with delayed flowering
and reduced *SOC1* and *FD* expression (Sheldon et al., 2002; Noh and
Amasino, 2003; Searle et al., 2006). This is also the case in *tps1-2 GVG::TPS1*,

while gene expression is restored in the double mutants (Fig. S10) (Zacharaki

et al., 2022). Our data combined with the findings of Zeng et al. (2024) suggest

that the regulation of SnRK1 activity is essential for T6P-dependent floral

induction, which has several modes of action throughout the floral network to

ensure that sufficient energy is available for this demanding developmental

transition. Finally, our findings shed further light on the multifactorial aspects of

C- and N-dependent regulation of flowering time.

Material and Methods

<u>Plant material and growth conditions</u>

Arabidopsis thaliana plants used for this study are of the Columbia (Col-0) ecotype. Mutant and transgenic lines such as *flc-3, svp-32, 35S::amiRTPS1, tps1-2,GVG::TPS1, tps1-2,GVG::TPS1,kin10-5, tps1-2,GVG::TPS1,snf4-1, nlp6-2, nlp7-1, nlp6-2,nlp7-1* and *spl345* were previously described (Michaels and Amasino, 1999; Lee et al., 2007; Wahl et al., 2013; Olas et al., 2019; Xie et al., 2020; Zacharaki et al., 2022). The *flc-3,tps1-2,GVG::TPS1* double mutant lines were generated by crossing. Genotypes were confirmed by a genotyping PCR using the oligonucleotides listed in Table S4.

Arabidopsis plants were grown in controlled growth chambers (Model E-36L, Percival Scientific Inc., Perry, IA, USA) at 22°C in long-day (LD, 16h light/8h dark) or short-day (SD, 8h dark/16h light) conditions. Light intensity was approximately 160 µmol/m²s. Controlled induction of flowering was performed by transferring the plants from non-inductive (SD) to inductive conditions (LD) as described (Schmid et al., 2003).

A previously established, almost natural, soil-based N-limited growth system consisting of ON and LN soil was used to grow plants (Tschoep et al., 2009). Briefly, the growth system consists of two types of peat-based soil mixtures with either an optimal level of N (ON, ~850 mg (N)/kg) or a limited level of N (LN, ~40 mg (N)/kg). Soil mixtures were prepared as described (Olas et al., 2019).

<u>Phenotypic analyses</u>

Flowering time was defined as days to flowering (DTF), which describes the days after germination to the day of bolting (inflorescence length 0.5cm), and by the total number of leaves (TLN). At least 16 plants were used to determine flowering time of each genotype. For vegetative phase change, juvenile leaf numbers were recorded and the leaf shape was digitally documented as described (Ponnu et al., 2020). A student's *t*-test was used to test the significance of the phenotypic differences.

<u>Reverse transcription quantitative PCR (RT-qPCR)</u>

Sampling, RNA extraction and RT-qPCR analysis of *FLC* in the *tps1-2,GVG::TPS1*, *tps1-2,GVG::TPS1,kin10-5* and *tps1-2,GVG::TPS1,snf4-1* were

performed as described (Zacharaki et al., 2022). RNA extraction and RT-qPCR
analyses of all the other genes were performed according to Wahl et al. (2013).
Relative expression values were calculated with the 2^DDCt method using Ct
values of a housekeeping gene index of *TUB2* (At5g62690), *SAND*
(At2g28390), *UBQ10* (At4g05320), and *PDF2* (At1g13320). RT-qPCR
analyses were performed in three or four biological replicates (n=3 or 4). A
Student's *t*-test was used to test for statistical significance.

RNA *in situ* hybridization
Wax embedding, sectioning, RNA *in situ* hybridization, and imaging were
performed as described (Wahl et al., 2013; Gramma and Wahl, 2023). Probes
were synthesized using the DIG RNA Labeling Kit (Roche, Mannheim,
Germany) for CDS of the *FLC* gene cloned into the pGEM®-T Easy vector
(Promega, Madison, Wisconsin, US). Oligonucleotides and construct IDs are
listed in Table S4.
451

Accession numbers
TPS1 (At1g78580), FLC (At5g10140), SVP (At2g22540), MAF5 (At5g65080),
FCA (At2g19520), EMF1 (At5g11530), PIE1/SNF2 (At3g12810), NLP6
(At1g64530), NLP7 (At4g24020), FRI (At4g00650), SUF4 (At1g30970), ELF7
(At1g79730), SEF (At5g37055), VRN1 (At3g18990), VRN2 (At4g16845),
EMF2/CYR1 (At5g51230), TFL2 (At5g17690), FVE (At2g19520), HUA2
(At2g19520), SNF4 (At1g09020), KIN10 (At3g01090), SPL3 (At2g33810),
SPL4 (At1g53160), SPL5 (At3g15270).
460

461 **Data availability**
462 The data supporting the findings of this study are included in this manuscript or
463 the supplemental information and material can be obtained from the
464 corresponding author upon reasonable request.

Acknowledgments and Funding

We thank Rob Hancock, Markus Schmid and the Wahl group for discussions and comments on the manuscript, Armin Schlereth, Christin Abel, Kerstin Zander, and Sebastian Kindermann for technical support. Work in the Wahl group was supported by the BMBF (031B0191), the DFG (SPP1530: WA3639/1-2, 2-1), the Max Planck Society and the James Hutton Institute.

Author's contribution

VW conceived and designed the experiments and prepared the figures. All authors performed essential experiments and analyzed data: JJO and VW performed the RNA *in situ* hybridizations; VG, JJO, VZ, and MML the RT-qPCRs; JJO, UML and JP performed phenotypic analyses. VG, VZ and VW wrote the manuscript with contributions from the other authors. All authors have read and commented on the text and figures within this manuscript.

References

Abe M, Kobayashi Y, Yamamoto S, Daimon Y, Yamaguchi A, Ikeda Y, Ichinoki H, Notaguchi M, Goto K, Araki T (2005) FD, a bZIP protein mediating signals from the floral pathway integrator FT at the shoot apex. Science **309:** 1052-1056

Alvarez JM, Schinke AL, Brooks MD, Pasquino A, Leonelli L, Varala K, Safi A, Krouk G, Krapp A, Coruzzi GM (2020) Transient genome-wide interactions of the master transcription factor NLP7 initiate a rapid nitrogen-response cascade. Nat Commun **11:** 1157

Baena-Gonzalez E, Rolland F, Thevelein JM, Sheen J (2007) A central integrator of transcription networks in plant stress and energy signalling. Nature **448:** 938-942

Baena-Gonzalez E, Sheen J (2008) Convergent energy and stress signaling. Trends Plant Sci **13:** 474-482

Bernier G, Havelange A, Houssa C, Petitjean A, Lejeune P (1993) Physiological Signals That Induce Flowering. Plant Cell **5:** 1147-1155

Bi YM, Wang RL, Zhu T, Rothstein SJ (2007) Global transcription profiling reveals differential responses to chronic nitrogen stress and putative nitrogen regulatory components in Arabidopsis. BMC Genomics **8:** 281

Carillo P, Feil R, Gibon Y, Satoh-Nagasawa N, Jackson D, Blasing OE, Stitt M, Lunn JE (2013) A fluorometric assay for trehalose in the picomole range. Plant Methods **9:** 21

Castro Marin I, Loef I, Bartetzko L, Searle I, Coupland G, Stitt M, Osuna D (2011) Nitrate regulates floral induction in Arabidopsis, acting independently of light, gibberellin and autonomous pathways. Planta **233:** 539-552

Chen HY, Lin SH, Cheng LH, Wu JJ, Lin YC, Tsay YF (2021) Potential transceptor AtNRT1.13 modulates shoot architecture and flowering time in a nitrate-dependent manner. Plant Cell **33:** 1492-1505

Corbesier L, Vincent C, Jang S, Fornara F, Fan Q, Searle I, Giakountis A, Farrona S, Gissot L, Turnbull C, Coupland G (2007) FT protein movement contributes to long-distance signaling in floral induction of Arabidopsis. Science **316:** 1030-1033

Dickens CWS, Van Staden J (1988) The In Vitro Flowering of Kalanchöe blossfeldiana Poellniz: I. ROLE OF CULTURE CONDITIONS AND NUTRIENTS. Journal of Experimental Botany **39:** 461-471

Eastmond PJ, van Dijken AJ, Spielman M, Kerr A, Tissier AF, Dickinson HG, Jones JD, Smeekens SC, Graham IA (2002) Trehalose-6-phosphate synthase 1, which catalyses the first step in trehalose synthesis, is essential for Arabidopsis embryo maturation. Plant J **29:** 225-235

Fernie AR, Bachem CWB, Helariutta Y, Neuhaus HE, Prat S, Ruan YL, Stitt M, Sweetlove LJ, Tegeder M, Wahl V, Sonnewald S, Sonnewald U (2020) Synchronization of developmental, molecular and metabolic aspects of source-sink interactions. Nat Plants **6:** 55-66

Fichtner F, Lunn JE (2021) The Role of Trehalose 6-Phosphate (Tre6P) in Plant Metabolism and Development. Annu Rev Plant Biol **72:** 737-760

Fischer JJ, Beatty PH, Good AG, Muench DG (2013) Manipulation of microRNA expression to improve nitrogen use efficiency. Plant Sci **210:** 70-81

Fredes I, Moreno S, Diaz FP, Gutierrez RA (2019) Nitrate signaling and the control of Arabidopsis growth and development. Curr Opin Plant Biol **47:** 112-118

Fujiwara S, Oda A, Yoshida R, Niinuma K, Miyata K, Tomozoe Y, Tajima T, Nakagawa M, Hayashi K, Coupland G, Mizoguchi T (2008) Circadian clock proteins LHY and CCA1 regulate SVP protein accumulation to control flowering in Arabidopsis. Plant Cell **20:** 2960-2971

Gramma V, Wahl V (2023) RNA In Situ Hybridization on Plant Tissue Sections: Expression Analysis at Cellular Resolution. Methods Mol Biol **2686:** 331-350

Helliwell CA, Wood CC, Robertson M, Peacock WJ, Dennis ES (2006) The Arabidopsis FLC protein interacts directly in vivo with SOC1 and FT chromatin and is part of a high-molecular-weight protein complex. Plant Journal **46:** 183-192

Hepworth SR, Valverde F, Ravenscroft D, Mouradov A, Coupland G (2002) Antagonistic regulation of flowering-time gene SOC1 by CONSTANS and FLC via separate promoter motifs. EMBO J **21:** 4327-4337

Jaeger KE, Wigge PA (2007) FT protein acts as a long-range signal in Arabidopsis. Curr Biol **17:** 1050-1054

Jeong EY, Seo PJ, Woo JC, Park CM (2015) AKIN10 delays flowering by inactivating IDD8 transcription factor through protein phosphorylation in Arabidopsis. BMC Plant Biol **15:** 110

Kant S, Peng M, Rothstein SJ (2011) Genetic regulation by NLA and microRNA827 for maintaining nitrate-dependent phosphate homeostasis in arabidopsis. PLoS Genet **7:** e1002021

Kaplan F, Kopka J, Sung DY, Zhao W, Popp M, Porat R, Guy CL (2007) Transcript and metabolite profiling during cold acclimation of Arabidopsis reveals an intricate relationship of cold-regulated gene expression with modifications in metabolite content. Plant J **50:** 967-981

Klebs G (1913) Uber des Verhaltnis der Aubenwelt zur Entwicklung der Pflanze. Sitz-Ber. Akad. Wiss. Heidelberg Ser. B **5:** 3-47

Konishi M, Yanagisawa S (2013) Arabidopsis NIN-like transcription factors have a central role in nitrate signalling. Nat Commun **4:** 1617

Konishi M, Yanagisawa S (2019) The role of protein-protein interactions mediated by the PB1 domain of NLP transcription factors in nitrate-inducible gene expression. BMC Plant Biol **19:** 90

Krapp A, Berthome R, Orsel M, Mercey-Boutet S, Yu A, Castaings L, Elftieh S, Major H, Renou JP, Daniel-Vedele F (2011) Arabidopsis roots and shoots show distinct temporal adaptation patterns toward nitrogen starvation. Plant Physiol **157:** 1255-1282

Lee JH, Park SH, Lee JS, Ahn JH (2007) A conserved role of SHORT VEGETATIVE PHASE (SVP) in controlling flowering time of Brassica plants. Biochimica Et Biophysica Acta-Gene Structure and Expression **1769:** 455-461

Lee JH, Ryu HS, Chung KS, Pose D, Kim S, Schmid M, Ahn JH (2013) Regulation of temperature-responsive flowering by MADS-box transcription factor repressors. Science **342:** 628-632

Lee JH, Sook Chung K, Kim SK, Ahn JH (2014) Post-translational regulation of SHORT VEGETATIVE PHASE as a major mechanism for thermoregulation of flowering. Plant Signal Behav **9:** e28193

Lee JH, Yoo SJ, Park SH, Hwang I, Lee JS, Ahn JH (2007) Role of SVP in the control of flowering time by ambient temperature in Arabidopsis. Genes Dev **21:** 397-402

Li D, Liu C, Shen L, Wu Y, Chen H, Robertson M, Helliwell CA, Ito T, Meyerowitz E, Yu H (2008) A repressor complex governs the integration of flowering signals in Arabidopsis. Developmental Cell **15:** 110-120

Li L, Liu KH, Sheen J (2021) Dynamic Nutrient Signaling Networks in Plants. Annu Rev Cell Dev Biol **37:** 341-367

Liang G, He H, Yu D (2012) Identification of nitrogen starvation-responsive microRNAs in Arabidopsis thaliana. PLoS One **7:** e48951

Lin YL, Tsay YF (2017) Influence of differing nitrate and nitrogen availability on flowering control in Arabidopsis. J Exp Bot **68:** 2603-2609

Liu T, Li Y, Ren J, Qian Y, Yang X, Duan W, Hou X (2013) Nitrate or NaCl regulates floral induction in Arabidopsis thaliana. Biologia **68:** 215-222

Marchive C, Roudier F, Castaings L, Brehaut V, Blondet E, Colot V, Meyer C, Krapp A (2013) Nuclear retention of the transcription factor NLP7 orchestrates the early response to nitrate in plants. Nat Commun **4:** 1713

Mathieu J, Warthmann N, Kuttner F, Schmid M (2007) Export of FT protein from phloem companion cells is sufficient for floral induction in Arabidopsis. Curr Biol **17:** 1055-1060

Michaels SD, Amasino RM (1999) FLOWERING LOCUS C encodes a novel MADS domain protein that acts as a repressor of flowering. Plant Cell **11:** 949-956

Noh YS, Amasino RM (2003) PIE1, an ISWI family gene, is required for FLC activation and floral repression in Arabidopsis. Plant Cell **15:** 1671-1682

Olas JJ, Apelt F, Watanabe M, Hoefgen R, Wahl V (2021) Developmental stage-specific metabolite signatures in Arabidopsis thaliana under optimal and mild nitrogen limitation. Plant Sci **303:** 110746

Olas JJ, Van Dingenen J, Abel C, Dzialo MA, Feil R, Krapp A, Schlereth A, Wahl V (2019) Nitrate acts at the Arabidopsis thaliana shoot apical meristem to regulate flowering time. New Phytol

Pant BD, Musialak-Lange M, Nuc P, May P, Buhtz A, Kehr J, Walther D, Scheible WR (2009) Identification of nutrient-responsive Arabidopsis and rapeseed microRNAs by comprehensive real-time polymerase chain reaction profiling and small RNA sequencing. Plant Physiol **150:** 1541-1555

Polge C, Thomas M (2007) SNF1/AMPK/SnRK1 kinases, global regulators at the heart of energy control? Trends Plant Sci **12:** 20-28

Ponnu J, Schlereth A, Zacharaki V, Dzialo MA, Abel C, Feil R, Schmid M, Wahl V (2020) The trehalose 6-phosphate pathway impacts vegetative phase change in Arabidopsis thaliana. Plant J **104:** 768-780

Pose D, Verhage L, Ott F, Yant L, Mathieu J, Angenent GC, Immink RG, Schmid M (2013) Temperature-dependent regulation of flowering by antagonistic FLM variants. Nature **503:** 414-417

Romera-Branchat M, Andres F, Coupland G (2014) Flowering responses to seasonal cues: what's new? Curr Opin Plant Biol **21:** 120-127

Schmid M, Uhlenhaut NH, Godard F, Demar M, Bressan R, Weigel D, Lohmann JU (2003) Dissection of floral induction pathways using global expression analysis. Development **130:** 6001-6012

Searle I, He Y, Turck F, Vincent C, Fornara F, Krober S, Amasino RA, Coupland G (2006) The transcription factor FLC confers a flowering response

to vernalization by repressing meristem competence and systemic signaling in Arabidopsis. Genes Dev **20:** 898-912

Sharma N, Geuten K, Giri BS, Varma A (2020) The molecular mechanism of vernalization in Arabidopsis and cereals: role of Flowering Locus C and its homologs. Physiol Plant **170:** 373-383

Sheldon CC, Conn AB, Dennis ES, Peacock WJ (2002) Different regulatory regions are required for the vernalization-induced repression of FLOWERING LOCUS C and for the epigenetic maintenance of repression. Plant Cell **14:** 2527-2537

Sheldon CC, Rouse DT, Finnegan EJ, Peacock WJ, Dennis ES (2000) The molecular basis of vernalization: the central role of FLOWERING LOCUS C (FLC). Proc Natl Acad Sci U S A **97:** 3753-3758

Song YH, Shim JS, Kinmonth-Schultz HA, Imaizumi T (2015) Photoperiodic flowering: time measurement mechanisms in leaves. Annu Rev Plant Biol **66:** 441-464

Srikanth A, Schmid M (2011) Regulation of flowering time: all roads lead to Rome. Cell Mol Life Sci **68:** 2013-2037

Stitt M, Hurry V (2002) A plant for all seasons: alterations in photosynthetic carbon metabolism during cold acclimation in Arabidopsis. Curr Opin Plant Biol **5:** 199-206

Sulpice R, Nikoloski Z, Tschoep H, Antonio C, Kleessen S, Larhlimi A, Selbig J, Ishihara H, Gibon Y, Fernie AR, Stitt M (2013) Impact of the carbon and nitrogen supply on relationships and connectivity between metabolism and biomass in a broad panel of Arabidopsis accessions. Plant Physiol **162:** 347-363

Sureshkumar S, Dent C, Seleznev A, Tasset C, Balasubramanian S (2016) Nonsense-mediated mRNA decay modulates FLM-dependent thermosensory flowering response in Arabidopsis. Nat Plants **2:** 16055

Teng Y, Liang Y, Wang M, Mai H, Ke L (2019) Nitrate Transporter 1.1 is involved in regulating flowering time via transcriptional regulation of FLOWERING LOCUS C in Arabidopsis thaliana. Plant Sci **284:** 30-36

Tschoep H, Gibon Y, Carillo P, Armengaud P, Szecowka M, Nunes-Nesi A, Fernie AR, Koehl K, Stitt M (2009) Adjustment of growth and central metabolism to a mild but sustained nitrogen-limitation in Arabidopsis. Plant Cell Environ **32:** 300-318

van Dijken AJ, Schluepmann H, Smeekens SC (2004) Arabidopsis trehalose-6-phosphate synthase 1 is essential for normal vegetative growth and transition to flowering. Plant Physiol **135:** 969-977

Vandesteene L, Ramon M, Le Roy K, Van Dijck P, Rolland F (2010) A single active trehalose-6-P synthase (TPS) and a family of putative regulatory TPS-like proteins in Arabidopsis. Mol Plant **3:** 406-419

Vidal EA, Alvarez JM, Moyano TC, Gutierrez RA (2015) Transcriptional networks in the nitrate response of Arabidopsis thaliana. Curr Opin Plant Biol **27:** 125-132

Wahl V, Ponnu J, Schlereth A, Arrivault S, Langenecker T, Franke A, Feil R, Lunn JE, Stitt M, Schmid M (2013) Regulation of flowering by trehalose-6-phosphate signaling in Arabidopsis thaliana. Science **339:** 704-707

Whittaker C, Dean C (2017) The FLC Locus: A Platform for Discoveries in Epigenetics and Adaptation. Annu Rev Cell Dev Biol **33:** 555-575

Wigge PA, Kim MC, Jaeger KE, Busch W, Schmid M, Lohmann JU, Weigel
D (2005) Integration of spatial and temporal information during floral induction
in Arabidopsis. Science **309:** 1056-1059
Xie Y, Zhou Q, Zhao Y, Li Q, Liu Y, Ma M, Wang B, Shen R, Zheng Z, Wang
H (2020) FHY3 and FAR1 Integrate Light Signals with the miR156-SPL Module-
Mediated Aging Pathway to Regulate Arabidopsis Flowering. Mol Plant **13:**
483-498
Xu C, Fang X, Lu T, Dean C (2021) Antagonistic cotranscriptional regulation
through ARGONAUTE1 and the THO/TREX complex orchestrates FLC
transcriptional output. Proc Natl Acad Sci U S A **118**
Xu C, Wu Z, Duan HC, Fang X, Jia G, Dean C (2021) R-loop resolution
promotes co-transcriptional chromatin silencing. Nat Commun **12:** 1790
Yang HL, Liu YJ, Wang CL, Zeng QY (2012) Molecular evolution of trehalose-
6-phosphate synthase (TPS) gene family in Populus, Arabidopsis and rice.
PLoS One **7:** e42438
Yang M, Zhu P, Cheema J, Bloomer R, Mikulski P, Liu Q, Zhang Y, Dean
C, Ding Y (2022) In vivo single-molecule analysis reveals COOLAIR RNA
structural diversity. Nature **609:** 394-399
Zacharaki V, Ponnu J, Crepin N, Langenecker T, Hagmann J, Skorzinski
N, Musialak-Lange M, Wahl V, Rolland F, Schmid M (2022) Impaired KIN10
function restores developmental defects in the Arabidopsis trehalose 6-
phosphate synthase1 (tps1) mutant. New Phytol **235:** 220-233
Zeng L, Zacharaki V, Wang Y, Schmid M (2024) Mutations in HUA2 restore
flowering in the Arabidopsis trehalose 6-phosphate synthase1 (tps1) mutant.
bioRxiv**:** 2024.2001. 2031.578264
Zhai Z, Keereetaweep J, Liu H, Feil R, Lunn JE, Shanklin J (2018) Trehalose
6-Phosphate Positively Regulates Fatty Acid Synthesis by Stabilizing
WRINKLED1. Plant Cell **30:** 2616-2627
Zhang Y, Primavesi LF, Jhurreea D, Andralojc PJ, Mitchell RA, Powers SJ,
Schluepmann H, Delatte T, Wingler A, Paul MJ (2009) Inhibition of SNF1-
related protein kinase1 activity and regulation of metabolic pathways by
trehalose-6-phosphate. Plant Physiol **149:** 1860-1871
708

Figure legends

Figure 1. **The Trehalose 6-phosphate pathway impacts on *FLOWERING LOCUS C*.** (**A**) Expression of *FLOWERING LOCUS C* measured by RT-qPCR in rosettes of Col-0 and *35S::amiRTPS1* plants grown under long days (16h light/ 8h darkness). *n* = 4. (**B**) Flowering time measured as leaf numbers (rosette leaves in gray; cauline leaves in white). n ≥ 15 individual plants per genotype. (**C**) Representative photographs of the plants analyzed in (B). Abbreviations: days after germination (DAG). Data represents mean, error bars are standard deviations (s.d.), statistically significant difference compared to Col-0 wild-type (Student *t*-test, *P<0.05, **P<0.01, and ***P<0.001).

Figure 2. ***FLOWERING LOCUS C* in response to nitrogen limitation.** (**A, B**) Expression of *FLOWERING LOCUS C* measured by RT-qPCR in rosettes (A) and apices (B) of Col-0 plants grown in optimal nitrogen (ON) and limited-nitrogen (LN) conditions under short days (8h light/ 16h dark). (**C**) *FLOWERING LOCUS C* expression measured by RT-qPCR in apices of plants initially grown under short days (30 days) and then transferred to long days to initiate the floral transition for 3, 5, and 7 days. (**D**) RNA *in situ* hybridization using *FLOWERING LOCUS C* specific probe on longitudinal sections through vegetative apices of Col-0 plants grown in ON and LN soils. Abbreviations: days after germination (DAG); days after shift (DAS). Data represents mean, error bars are standard deviations (s.d.), n=3, statistically significant difference between ON and LN (Student's *t*-test, *P<0.05, **P<0.01, ***P<0.001). Star indicates apex summit.

Figure 3. **FLOWERING LOCUS C and SHORT VEGETATIVE PHASE are required for the limited nitrogen-dependent flowering response.** (**A**) Flowering time of Col-0 wild-type plants treated with an 8-week period of cold. Note that afterwards plants were transferred to 22°C until flowering. (**B**) Flowering time of Col-0, *flc-3,* and *svp-32* mutant plants grown under short-day (8h light/16h darkness) conditions. Data represents mean, error bars are standard deviations (s.d.), n ≥ 15 individual plants per genotype, statistically significant difference between ON and LN (Student's *t*-test, ***P<0.001).

Figure 4. *FLOWERING LOCUS C* **expression downstream of NIN-LIKE PROTEIN 6 (NLP6) and NIN-LIKE PROTEIN 7 (NLP7).** Expression of *FLOWERING LOCUS C* measured by RT-qPCR at 10 days after germination (DAG) in rosettes of Col-0, *nlp6-2,nlp7-1*, and *nlp6-2,nlp7-1* plants. Data represents mean, error bars are standard deviations (s.d.), n=3, statistically significant difference compared to Col-0 wild-type (Student's *t*-test, *$P<0.05$, **$P<0.01$).

Figure 5. **Trehalose 6-phosphate pathway and nitrogen-signaling converge at *FLOWERING LOCUS C*.** (**A**, **B**, **C**) Expression of *FLOWERING LOCUS C* measured by RT-qPCR in (A) rosettes of wild-type Col-0 and *35S::amiRTPS1* plants grown in optimal nitrogen (ON) and limited-nitrogen (LN) conditions under short days (8h light/ 16h dark) at 60 days after germination (DAG), in (B) apices and (C) rosettes of wild-type Col-0, *tps1-2,GVG:TPS1*, *snf4,tps1-2,GVG:TPS1* and *kin10,tps1-2,GVG:TPS1* plants grown in standard soil under long days (16h light/ 8h dark). Data represents mean, error bars are standard deviations (s.d.), statistically significant difference compared to Col-0 wild-type (Student *t*-test, *$P<0.05$, **$P<0.01$, and ***$P<0.001$).

Figure 6. **Carbon and nitrogen signaling target similar components of the flowering network in the shoot apical meristem for the proper timing of flowering**. *FLOWERING LOCUS C,* a key repressor of flowering, is not only regulated by cold temperature as part of the vernalization process, but is also affected by nutrient availability. The Trehalose 6-phosphate pathway negatively impacts *FLOWERING LOCUS C* via SUCROSE NON-FERMENTING 1 RELATED KINASE 1. Nitrogen signaling controls *FLOWERING LOCUS C* via a yet to identify mechanism involving NIN-LIKE PROTEIN 7. The repressor complex composed of FLOWERING LOCUS C and SVP is eventually tuned by the adjustment of *FLOWERING LOCUS C* expression downstream of both carbon and nitrogen signaling to control *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1* in the shoot apical meristem. Independently, both nutrient pathways work via the age pathway (SQUAMOSA PROMOTER-BINDING PROTEIN-LIKE 3-5) to induce flowering.

773 **Supplemental Material**

Supplemental Figure S1. *MADS AFFECTING FLOWERING 5* in *35S::amiRTPS1* plants.

Supplemental Figure S2. *flc-3* partially suppresses the delayed vegetative phase change phenotype of *tps1-2,GVG::TPS1* plants.

Supplemental Figure S3. *MADS AFFECTING FLOWERING 5* in response to nitrogen limitation.

Supplemental Figure S4. *SHORT VEGETATIVE PHASE* in *35S::amiRTPS1* plants.

Supplemental Figure S5. *SHORT VEGETATIVE PHASE* in response to nitrogen limitation.

Supplemental Figure S6. Regulators upstream *FLOWERING LOCUS C* in response to N limitation.

Supplemental Figure S7. Regulators upstream *FLOWERING LOCUS C* in *35S::amiRTPS1* plants.

Supplemental Figure S8. *FLOWERING LOCUS C* and *SHORT VEGETATIVE PHASE* in *spl345* mutant plants.

Supplemental Figure S9. Plant phenotype of *nlp6* and *nlp7* mutant plants.

Supplemental Figure S10. *FLOWERING LOCUS D* in *snrk1,tps1-2,GVG::TPS1* mutants.

Supplemental Table S1. Flowering time data of experiments described in this study (Figure S2).

Supplemental Table S2. Vegetative phase change data of experiment described in this study

Supplemental Table S3. Putative nitrate responsive *cis*-elements (NREs) in regulators upstream *FLOWERING LOCUS C.*

Supplemental Table S4. Oligonucleotides used in this study.

Figure 1. **The Trehalose 6-phosphate pathway impacts on *FLOWERING LOCUS C*.** (**A**) Expression of *FLOWERING LOCUS C* measured by RT-qPCR in rosettes of Col-0 and *35S::amiRTPS1* plants grown under long days (16h light/ 8h darkness). *n* = 4. (**B**) Flowering time measured as leaf numbers (rosette leaves in gray; cauline leaves in white). n ≥ 15 individual plants per genotype. (**C**) Representative photographs of the plants analyzed in (B). Abbreviations: days after germination (DAG). Data represents mean, error bars are standard deviations (s.d.), statistically significant difference compared to Col-0 wild-type (Student *t*-test, *$P<0.05$, **$P<0.01$, and ***$P<0.001$).

Figure 2. *FLOWERING LOCUS C* **in response to nitrogen limitation.** (**A, B**) Expression of *FLOWERING LOCUS C* measured by RT-qPCR in rosettes (A) and apices (B) of Col-0 plants grown in optimal nitrogen (ON) and limited-nitrogen (LN) conditions under short days (16h light/ 8h dark). (**C**) *FLC* expression measured by RT-qPCR in apices of plants initially grown under short days (30 days) and then transferred to long days to initiate the floral transition for 3, 5, and 7 days. (**D**) RNA *in situ* hybridization using *FLOWERING LOCUS C* specific probe on longitudinal sections through vegetative apices of Col-0 plants grown in ON and LN soils. Abbreviations: days after germination (DAG); days after shift (DAS). Data represents mean, error bars are standard deviations (s.d.), n=3, statistically significant difference between ON and LN (Student's *t*-test, *$P<0.05$, **$P<0.01$, ***$P<0.001$). Star indicates apex summit.

Figure 3. **FLOWERING LOCUS C and SHORT VEGETATIVE PHASE are required for the limited nitrogen-dependent flowering response.** (**A**) Flowering time of Col-0 wild-type plants treated with an 8-week period of cold. Note that afterwards plants were transferred to 22°C until flowering. (**B**) Flowering time of Col-0, *flc-3,* and *svp-32* mutant plants grown under short-day (8h light/16h darkness) conditions. Data represents mean, error bars are standard deviations (s.d.), n ≥ 15 individual plants per genotype, statistically significant difference between ON and LN (Student's *t*-test, ***P<0.001).

Figure 4. *FLOWERING LOCUS C expression downstream of NIN-LIKE PROTEIN 6 (NLP6) and NIN-LIKE PROTEIN 7 (NLP7).* Expression of *FLOWERING LOCUS C* measured by RT-qPCR at 10 days after germination (DAG) in rosettes of Col-0, *nlp6-2*, *nlp7-1*, and *nlp6-2,nlp7-1* plants. Data represents mean, error bars are standard deviations (s.d.), n=3, statistically significant difference compared to Col-0 wild-type (Student's *t*-test, *$P<0.05$, **$P<0.01$).

Figure 5. **Trehalose 6-phosphate pathway and nitrogen-signaling converge at *FLOWERING LOCUS C*.** (**A**, **B**, **C**) Expression of *FLOWERING LOCUS C* measured by RT-qPCR in (A) rosettes of wild-type Col-0 and *35S::amiRTPS1* plants grown in optimal nitrogen (ON) and limited-nitrogen (LN) conditions under short days (8h light/ 16h dark) at 60 days after germination (DAG), in (B) apices and (C) rosettes of wild-type Col-0, *tps1-2,GVG:TPS1*, *snf4,tps1-2,GVG:TPS1* and *kin10,tps1-2,GVG:TPS1* plants grown in standard soil under long days (16h light/ 8h dark). Data represents mean, error bars are standard deviations (s.d.), significant difference compared to Col-0 wild-type (Student *t*-test, *$P<0.05$, **$P<0.01$, and ***$P<0.001$).

Figure 6. **Carbon and nitrogen signaling target similar components of the flowering network in the shoot apical meristem for the proper timing of flowering**. *FLOWERING LOCUS C,* a key repressor of flowering, is not only regulated by cold temperature as part of the vernalization process, but is also affected by nutrient availability. The Trehalose 6-phosphate pathway negatively impacts *FLOWERING LOCUS C* via SUCROSE NON-FERMENTING 1 RELATED KINASE 1. Nitrogen signaling controls *FLOWERING LOCUS C* via a yet to identify mechanism involving NIN-LIKE PROTEIN 7. The repressor complex composed of FLOWERING LOCUS C and SVP is eventually tuned by the adjustment of *FLOWERING LOCUS C* expression downstream of both carbon and nitrogen signaling to control *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1* in the shoot apical meristem. Independently, both C and N pathways work via the age pathway (SQUAMOSA PROMOTER-BINDING PROTEIN-LIKE 3-5) to induce flowering.