Comparing arm to whole-body motor control disambiguates age-related deterioration from compensation

Robin Mathieu*[1], Florian Chambellant[1], Elizabeth Thomas[1], Charalambos Papaxanthis[1,2], Pauline Hilt[1], Patrick Manckoundia[1,3], France Mourey[1], Jérémie Gaveau[1]

[1] Inserm UMR 1093-CAPS, Université de Bourgogne, UFR des Sciences du Sport, F-21000, Dijon. Institut National de la Santé et de la Recherche Médicale : UMR 1093, Université de Bourgogne, France.
[2] Pôle Recherche et Santé Publique, CHU Dijon Bourgogne, F-21000 Dijon, France
[3] Département de Gérontologie, Hôpital de Champmaillot, Centre Hospitalier Universitaire de Dijon, Dijon, France

*Corresponding author
Correspondence: robin.mathieu@u-bourgogne.fr

ABSTRACT

As the global population ages, it is crucial to understand sensorimotor compensation mechanisms that allow older adults to remain in good physical health. Although age-related compensation has long been conceptualized and despite important research efforts, compensatory processes remain essentially chimeras. A major problem with their identification is the ambiguous interpretation of age-related alterations. The question of whether an alteration reflects deterioration or compensation is difficult to answer. Here we compared the electromyographic and kinematic patterns of different motor tasks in younger (n = 20; mean age = 23.6 years) and older adults (n = 24; mean age = 72 years). Building on the knowledge that humans take advantage of gravity effects to minimize their muscle effort, we probed the ability of younger and older adults to plan energetically efficient movement during arm-only and whole-body movements. In line with previous studies and compared to younger adults, muscle activation patterns revealed that older adults used a less efficient movement strategy during whole-body movement tasks. Does this reflect deterioration or compensation? We found that this age-related alteration was task-specific. It did not affect arm movements, thereby supporting the hypothesis that healthy older adults maintain the ability to plan energetically efficient movements. More importantly, we found that the decreased whole-body movement efficiency was negatively correlated with kinematic measures of balance control (i.e., the center-of-mass movement amplitude and speed). The more efficient the movement strategy, the more challenging the balance. Overall, these results suggest that decreased movement efficiency in older adults does not reflect a deterioration but rather a compensation process that adapts movement strategy to the task specificities. When balance is at stake, healthy older adults prefer stability to energy efficiency.

Keywords: Aging, motor control, electromyography, gravity, sensorimotor compensation, equilibrium, posture

Introduction

Living old and healthy, also known as successful aging, is a blessing but is nonetheless associated with deterioration in various organs and functions. In terms of motor deterioration, aging is associated with loss of muscle mass (Larsson et al., 2019), sensory receptor degradation (Goble et al., 2009; Zalewski, 2015; Saftari & Kwon, 2018), and cortical atrophy (Hoffstaedter et al., 2015; Salat, 2004). Functionally, this translates into a decline in muscle strength and power (Larsson et al., 2019; Pousson et al., 2001) and movements that tend to become slower and more variable (Buckles, 1993; Darling et al., 1989). If the deteriorations are too great, they lead to reductions in quality of life and, ultimately, to dependency. Importantly, successful aging is thought to depend on compensatory processes that offset deteriorations (Baltes & Baltes, 1990; Martin et al., 2015; Zhang & Radhakrishnan, 2018). Even the most elementary concept of health includes compensatory processes at its core. The World Health Organization defined health as "a state of complete physical, mental, and social well-being and not merely the absence of disease or infirmity" (1948). Scientists and clinicians later redefined it even more generally as "the ability to adapt and to self-manage" (Huber et al., 2011; The Lancet, 2009). So, despite the normal deterioration associated with age, compensatory processes enable older adults to adapt and remain in good health (i.e., aging successfully), and thus continue to live comfortably.

In a world with a rapidly aging population (Rudnicka et al., 2020), it is essential to understand the compensatory processes that enable older people to remain healthy. This represents a critical step toward implementing interventions aimed at detecting, preventing, or reducing frailty and later dependency (for reviews, see Barulli & Stern, 2013; Ouwehand et al., 2007; Poirier et al., 2021; Zhang & Radhakrishnan, 2018). Compensation has long been theorized and could be defined as "a response to loss in means (resources) used to maintain success or desired levels of functioning (outcomes)" (Baltes, 1997). In contexts of severe deterioration, the most basic form of compensation is the use of external aids (e.g., a crutch for walking). Such compensations are observed in frail or dependent older adults, i.e. when deterioration is severe. When considering more subtle deterioration levels, identifying compensation becomes challenging. In these cases, compensatory processes enable older adults to maintain behavioral performances similar to those of younger adults, at least for the less demanding tasks of daily life (Barulli & Stern, 2013). These compensatory processes are the result of neurophysiological and behavioral adaptations that are more difficult to observe with the naked eye. Almost thirty years ago, in his famous theory of selection, optimization, and compensation, Paul Baltes and his colleagues already noted this difficulty (Baltes, 1997; Baltes & Baltes, 1990).

Since then, countless studies have explored compensatory processes using powerful tools and analyses (for recent reviews, see Bunzeck et al., 2024; Fettrow et al., 2021; Poirier et al., 2021). These studies have considerably advanced the description and understanding of age-related neural alterations. Nevertheless, behavioral compensatory processes and their underlying neural mechanisms remain essentially chimeras. Building on the theoretical work of Krakauer et al. (2017), we recently proposed that an important reason for this failure may be that studies focusing on age-related neural alterations have used overly crude behavioral paradigms (Poirier et al., 2021). Typically, these studies have used broad measures such as muscle strength, reaction time, or movement time. Although these measures and paradigms tested important functional motor performances, they measured the combination of several behavioral strategies and subtending neural mechanisms. Since these strategies and mechanisms likely showed different levels of age-related deterioration, previous studies have likely mixed deterioration and compensation processes (Poirier et al., 2021). Identifying neural compensation requires linking the brain to behavior and, to establish a precise link, we need fine behavioral measures and experimental paradigms that allow approaching the constituent processes of a behavior (Krakauer et al., 2017; Pereira et al., 2020; Urai et al., 2022). It is therefore essential to first develop detailed knowledge of age-related compensation at the behavioral level.

We sought to fill this gap by building upon the results of two different bodies of literature. On one hand, several studies have demonstrated that the brain plans efficient arm movements that take advantage of the mechanical effects of gravity to minimize muscle effort (Berret et al., 2008; Crevecoeur et al., 2009; Gaveau et al., 2014, 2016, 2021; Gaveau & Papaxanthis, 2011; Gueugneau et al., 2023; for a review, see White et al., 2020). Importantly, recent work demonstrates that this ability is maintained and maybe even upregulated in older adults (Healy et al., 2023; Huang & Ahmed, 2014; Poirier et al., 2020; Summerside et al., 2024) On the other hand, studies probing the control of movements performed with the entire body have reported a different conclusion. Even-though such movements involve increased effort in older compared to younger adults (Hortobagyi et al., 2003, 2011; Julius et al., 2012; VanSwearingen & Studenski, 2014), results from

kinematic studies suggest that older adults plan movements that are less-efficient (Casteran et al., 2018; Paizis
et al., 2008). Since the ability to plan efficient movements is maintained in older adults, as testified by arm
movements studies, one may speculate that this decreased efficiency reflects an age-related compensation
that changes movement strategy (i.e. an age-related motor adaptation process). However, because these
literatures used very different experimental paradigms and measurements, this conclusion is highly
speculative. More importantly, the results of numerous other studies could also interpret the decreased
efficiency observed in whole-body movements as a deterioration of the ability to produce efficient motor
patterns (Goodpaster et al., 2006; Henry & Baudry, 2019; Quinlan et al., 2018; Vernazza-Martin et al., 2008).
Here we test the hypothesis that age-related alterations in movement efficiency correspond to an adaptation
process, i.e. a change in movement strategy that compensates for other deteriorated sensorimotor
components. To overcome the aforementioned limitations, using a specific muscle activation pattern analysis
that has proven relevant to focusing on this precise process of effort efficiency (Chambellant et al., 2023;
Gaveau et al., 2021; Poirier et al., 2022, 2024; Thomas et al., 2023), we compare older to younger adults on
tasks involving either arm or whole-body movements.

Methods

Participants

Because we had no a-priori data to calculate the ideal sample size, we included as many participants as
possible over a fixed recruitment period. Twenty younger adults (23.6 ± 2.1 y.o.) and twenty-four older adults
(72 ± 5.3 y.o.) were included in the study after giving their oral informed consent. Participants had normal or
corrected-to-normal vision and did not present any neurological or muscular disorders. The laterality index of
each participant was superior to 60 (Edinburgh Handedness Inventory, Oldfield 1971), indicating that all
participants were right-handed. The study was carried out following legal requirements and international
norms (Declaration of Helsinki, 1964) and approved by the French National Ethics Committee (2019-A01558-
49).

Experimental Protocol

All participants performed four tasks in a randomized order. These tasks either required moving the arm
only (Figure 1A), or the whole-body (Figure 1B-D). Whole-body movements consisted in seat-to-stand/back-
to-sit (STS/BTS, Figure 1B), whole-body reaching toward a near target (WBR D1, Figure 1C), and whole-body
reaching toward a far target (WBR D2, Figure 1D). The arm task was selected because it is the reference task
that has been studied to demonstrate how muscle patterns take advantage of gravity effects to minimize
muscle effort. The whole-ody tasks were selected because they include an equilibrium constraint, are

Figure 1. Illustration of the four tasks. Each panel illustrates the extreme body positions between which participants performed their movements. Each position was alternatively the starting or the ending target of a movement, depending on movement direction. **A**: Single degree of freedom arm movements flexion/extension around the shoulder joint (flexion/extension). Participants performed upward and downward arm movements. **B**: Seat-to-stand/Back-to-sit movements. Participants performed vertical multi-articular whole-body movements to either stand up from the stool (upward movement) or sit on it (downward movement). **C**: Whole-body reaching task toward a near target. Participants performed vertical multi-articular whole-body movements to either reach towards targets that were located nearby the floor (downward movement) or to bounce back from this position toward a rest vertical standing position (upward movement). **D**: same as C but with targets that were placed farther away on the antero-posterior axis.

ARM task

 This task was similar to a task used in several previous studies probing human movement adaptation to the gravity environment (Gaveau et al., 2014, 2016, 2021; Gaveau & Papaxanthis, 2011; Gentili et al., 2007; Hondzinski et al., 2016; Le Seac'h & McIntyre, 2007; Poirier et al., 2020, 2022; Yamamoto & Kushiro, 2014a). Using their right arm, participants carried out single-degree-of-freedom vertical arm flexion/extension movements around the shoulder joint. Two blocks of arm movements were performed in a randomized order. One block consisted of six slow movements, and one block consisted of twelve fast movements. Two targets (diameter of 3 cm) were placed in front of the participant's right shoulder

(in a parasagittal plane) at a distance corresponding to the length of their fully extended arm plus two centimeters. The prescribed movement amplitude between the two targets was 45°, corresponding to an 112.5° (upward target, 22.5° above horizontal) and 67.5° (downward target, 22.5° below horizontal) shoulder flexion/extension.

STS/BTS task
    This task was similar to those of previous studies (Jeon et al., 2021; Manckoundia et al., 2006; Millington et al., 1992; Mourey et al., 1998). Participants were seated on an armless stool whose height was adjusted to correspond to 30% of the participant's height. The hands were positioned on the hips, and the back was instructed to be maintained about vertical. Participants were asked to stand up from the stool, make a short pause (about 2s), and then sit back on the stool. Similarly to arm movements, participants executed two blocks of movements in a randomized order. One block consisted of six slow movements, and the other consisted of 12 fast movements.

WBR task
    This task was similar to those of Casteran et al. (2018) and Paizis et al. (2008). Starting from an upright position, we asked participants to perform whole body reaching movements (WBR) toward two targets nearby the floor with their two index fingers (10% of their heights above the floor). The two targets (4 × 2 cm) were spaced by 0.5m on a medio-lateral axis and centered on the participant's sagittal axis. They were placed in front of the participant at two different distances, corresponding to 15% (D1) or 30% (D2) of their height on the antero-posterior axis. Distances were measured from the participant's big toe. Similarly to the previous two tasks, for each distance and in a randomized order, participants executed two blocks of trials performed at two different speeds. One block consisted of six slow movements and the other twelve fast movements (total of four blocks: two speeds × two distances).

Trial organization
    The organization of a trial was similar for all tasks. It was carried out as follows: i) the experimenter indicated to get ready; ii) the participant adopted the requested initial position; iii) after a brief delay (~1 second), the experimenter verbally informed the participant that she/he was free to reach the requested final position whenever she or he wanted. Note that reaction time was not emphasized in our experiment; iv) the participant was requested to maintain the final position for a brief period (about 1 second); v) the experimenter instructed to move back to the starting position (reversed movement) whenever desired; vi) lastly, the participant was asked to relax. A short rest period (~20 s) separated trials to prevent muscle fatigue. Additionally, participants were free to rest as long as they wanted between blocks. Participants were allowed to perform a few practice trials (~3 trials) before each block. Low-speed and high-speed blocks were similar except that the instructions were to perform the movements in roughly 5 seconds or as fast as possible, respectively.

Data Collection

Kinematics
    We used the Plug-In Gait full body model (Vicon, Oxford Metrics, UK) following their recommendations to place the 39 reflective markers on the participant's head (temples and backs of the head to form a rigid plan with the head), back (C7, T10 and on the right scapula), torso (jugular notch where the clavicles meet the sternum and on the xiphoid of the sternum), shoulders (acromion), arms (upper lateral 1/3 for the left arm, and 2/3 for the right arm), elbows (lateral epicondyle), forearms (lower lateral 1/3 for the left forearm, and 2/3 for the right forearm), wrists (both cubitus styloid processes), hands (middle of the proximal knuckle of the index), pelvis (anterior and posterior superior iliac spine), thighs (upper lateral 1/3 for the left leg, and 2/3 for the right leg), knees (lateral side of the flexion-extension axis), calves (upper lateral 1/3 for the left calf, and 2/3 for the right calf), ankles (lateral malleolus), and feet (second metatarsal head and heel). The markers on the scapula, on the arms, on the forearms, on the thighs, and on the calves have been deliberately placed asymmetrically so that the model can best dissociate the right and left sides, these markers are not used for the analyses presented in this manuscript.

We recorded the position of all markers with an optoelectronic motion capture system (Vicon system,
Oxford Metrics, UK; 18 cameras) at a sampling frequency of 200 Hz. The spatial variable error of the system
was less than 0.5mm.

EMG
We placed sixteen bipolar surface electrodes (Cosmed, pico EMG, sampling frequency: 1000Hz) on the
anterior (AD) and posterior (PD) heads of the deltoid, vastus lateralis (VL), biceps femoris (BF), spinal erectors
on L1 (ESL1) and on T7 (EST7), the soleus (SOL), and on the tibialis anterior (TA) to record EMG activity.
Electrodes were placed bilaterally. The location of each electrode was determined following the
recommendations from Barbero et al. (2012).
The Giganet unit (Vicon, Oxford Metrics, UK) synchronously recorded kinematic and EMG data.

Data Analysis
We processed kinematic and EMG data using custom programs written in Matlab (Mathworks, Natick, MA).
Data processing was inspired by previous studies (Gaveau et al., 2021; Poirier et al., 2022) and was similar for
all tasks.

Kinematics analysis
First, we filtered position using a third-order low-pass Butterworth filter (5 Hz cut-off, zerophase distortion,
"butter" and "filtfilt" functions). We then computed the amplitude of the movement using steady phases
(200ms for fast movements and 500ms for slow movements) before and after the movement, using the marker
of the right shoulder (Figure 2). The amplitude was computed on the Z axis for fast movements and on X, Y and
Z axes for slow movements. For slow movements, we used 3D position to minimize detection error on signals
that were more variable than those obtained during fast movements. Last, we automatically defined
movement onset and offset as the moments when the displacement rose above or felt below a threshold
corresponding to 5% and 95% of the total movement amplitude, respectively.

Figure 2. Detection of movement start and end. This panel displays the recording of two successively opposite fast Sit to stand / Back to sit movements. The black trace represents the position of a shoulder marker through time. Rest position is collected during steady phases, before and after each movement (ascending or descending). Based on the data obtained in steady phases, a recursive algorithm automatically defined movement onset and offset as the moments when the displacement rose above or felt below a threshold corresponding to 5% and 95% of the total movement amplitude, respectively.

213   On behalf of using the kinematics to define the start and end of movement, we analyzed the displacement
214   of the Center of Mass (CoM) in three dimensions to understand how equilibrium was maintained during the
215   whole-body tasks. This was done to reproduce the work of Casteran et al., (2018) and Paizis et al., (2008), but

also, and more importantly, to perform a simple analyse to test whether our main criterion, quantified via electromyographic activity, is linked to a simple, interpretable kinematic change. Our analysis utilized a seven-segment mathematical model incorporating rigid segments such as the Trunk, Thigh, Shank, Foot, Upper arm, Forearm, and Hand. We used anthropometric data from (Winter, 2009), as performed by previous studies (Berret et al., 2009; Stapley et al., 1999). Our choice of movement segmentation for this specific kinematic analysis has been guided by the works of Casteran et al., (2018) and Paizis et al., (2008). We determined movement onset and ofsset on velocity profiles, using a threshold of 5% of the peak velocity. We further explored the kinematics of the whole-body tasks using two simple parameters: i) the total displacement of the center of mass, calculated as the distance between the start and end positions and normalized by the subject's height ; and ii) the peak velocity of the center of mass. We focused on downard movements as these are the ones that have been studied and present the greatest challenge to balance. The specific process to compute criteria used by previous studies (Casteran et al., 2018 and Paizis et al., 2008) are detailed and available in Supplementary Figure 1.

EMG analysis

   Rational about effort. A body of computational studies has demonstrated that human arm movements take advantage of gravity effects to minimie muscle effort (Berret et al., 2008; Crevecoeur et al., 2009; Gaveau et al., 2011, 2014, 2016, 2021). Most of theses studies used mathematical models that minimized the absolute work of muscle force to produce the arm displacement (Berret et al., 2008; Gaveau et al., 2011, 2014, 2016, 2021). More sepcifically, the study from Berret et al. 2008 formally demonstrated that this effort-cost, and associated behavior, corresponded to an energetic-like optimum. Thus, building upon this literature, we interchangebly use "energetic efficiency" and "effort minimization" in the present work. Because previous studies have shown that the amplitude of kinematic and electromygographic markers directly relates to how much effort is minimized (Gaveau et al., 2016; Poirier et al., 2022), here we compare the amplitude of an EMG marker between younger and older adults. If the EMG marker increases, this means that muscle effort minimization increases – i.e., the minimization process is upregulated – thus that muscle effort decreases. If the EMG marker decreases, this means that muscle effort minimization decreases – i.e., the minimization process is downregulated – thus that muscle effort increases. Below, following methodologies developed by several previous studies, we detail how we obtain this EMG marker.

   Pre-processing. EMG signals were first rectified and filtered using a bandpass third-order Butterworth filter (bandpass 30-300 Hz, zero-phase distortion, "butter" and "filtfilt" functions) followed by a low-pass third-order Butterworth filter (low-pass frequency: 5 Hz) to highlight important features of muscular activities. Signals were integrated using a 100ms sliding window (trapezoidal numerical integration from Matlab [Mat, 2021]), and cut off. For fast movements, EMG signals were cut off from 240ms before movement onset to 90ms after movement offset. For slow movements, EMG signals cut off from 75ms before movement onset to 75ms after movement offset. These timing values were obtained from preliminary analyses detecting EMG activity start and stop before and after all movements. The result is the average of all participants. Importantly, those values were kept constant for all participants and, thus, should not bias group comparisons.

   Phasic/tonic separation. We then computed the phasic component of each EMG signal using a well-known subtraction procedure that has mostly been used to study arm movements (Buneo et al., 1994; d'Avella et al., 2006, 2008; Flanders et al., 1994; Flanders & Herrmann, 1992; Gaveau et al., 2021). This processing allows quantifying how much the central nervous system takes advantage of the gravity torque when moving the body in the gravity environment (Gaveau et al., 2021; Poirier et al., 2020, 2022, 2024). Here, we customized this procedure to investigate whole body movements since movements are not one-degree-of-freedom movements. First, the tonic signal was obtained from the six slow movements. For that purpose, the cut movements (as described earlier with delays) were normalized in duration to be finally averaged together in one tonic signal. Second, to improve signal to noise ratio, EMG traces of fast movements were ordered according to movement mean velocity and averaged across two trials (from the two slowest to the two fastest movements). This resulted in six EMG traces to be analyzed for each block. Each set of two traces was normalized in duration (corresponding to the mean duration of the two traces) before averaging. Third, the phasic component was obtained by subtracting the tonic EMG from the EMG trace of each pair of fast

movements. Finally, to set the data of all participants on a common scale, phasic activity was normalized by the maximal EMG value recorded in each task for each participant.

Figure 3: Illustration of the calculation method to obtain phasic EMG components. Electromyographic signals (mV) are presented as a function of time. Pattern duration and amplitude are normalized (see methods). **A**: Six integrated Vastus Lateralis EMG signals during fast BTS movements of a typical participant (BTS: Back-to-seat) ; **B**: Six integrated Vastus Lateralis EMG signals recorded during slow BTS movements of a typical participant . These signals represent the the tonic component. ; **C**: Integrated phasic EMG component computed using the six fast (panel **A**) and slow movements (panel **B**). The phasic is calculated by subtracting the mean of the slow acquisitions from the fast acquisitions (Phasic = Fast – Tonic).

Muscles selection. It was recently shown that the phasic EMG activity of antigravity muscles, those that pull against the gravity vector, consistently exhibits negative epochs (Chambellant et al., 2024; Gaveau et al., 2021; Poirier et al., 2022; Thomas et al., 2023) when the arm acceleration sign is coherent with the gravity acceleration sign (i.e., in the acceleration phase of downward movement and in the deceleration phase of upward movements). This observation likely reflects an optimal predictive motor strategy where muscle activity is decreased when gravity assists arm movements, thereby discounting muscle effort (Gaveau et al., 2021). In the present study, the antigravity muscles are: i) the Anterior Deltoïd (DA), flexing the shoulder joint; ii) the Vastus Lateralis (VL), extending the knee joint; iii) the Erector Spinae L1 (ESL1), extending the rachis; iv) the Erector Spinae D7 (ESD7), extending the rachis; v) the Soleus (SOL), flexing the ankle in the plantar direction. Because the Erector Spinae D7 and the Soleus muscles did not play a strong focal role but a rather postural one in the present tasks, we focused our analyses on the remaining three muscles (DA, VL and ESL1). Probing the activation of a postural muscle does not seem appropriate to test whether the nervous system takes advantage of gravity to produce the movement. We focused on DA during arm movements and on VL and ESL1 during movement of the entire body.

Quantifying negativity. We defined negative epochs as an interval where the phasic EMG signal was inferior to zero minus three times the standard deviation of the stable phase preceding the movement, and this for at least 40ms. This duration has been chosen after preliminary tests to avoid detecting false-positive. We kept it constant for all analyses. We used this value as a threshold to automatically detect negativity onset and offset. On each negativity phase, we computed: i) a negativity index, defined as T x NA / TA, with NA the Negative Area integrated on the phasic signal between negativity onset and offset, TA the Tonic Area

integrated on the tonic signal between the negativity onset and offset, and T the duration of the negative epoch normalized by movement duration. This value is always negative or null. The lower the value, the greater the efficiency; ii) negativity occurrence, defined as the number of trials where a negative epoch was automatically detected, divided by the total number of trials in the condition; iii) negativity duration, defined as the duration of the negative epoch normalized by movement duration; iv) negativity amplitude, defined as the minimal $Phasic\ value\ /\ Tonic\ Value \times 100$ during the negative period. A value of -100 indicates that the muscle is completely relaxed and a value of 0 indicates that the muscle exactly compensated the gravity torque.

As is often the case with EMG recordings, some of the EMG signals exhibited aberrant values. Those signals are usually due to poor contact between the electrodes and the skin. Supplementary Table 1 summarizes the issues encountered with all electrodes and participants.

Machine Learning

Here we were interested in comparing arm movement and whole-body movement control because the scientific literature has reported that whole-body movements control changes with age, whilst arm movement control does not (Casteran et al., 2018; Paizis et al., 2008; Poirier et al., 2020, 2024; Vernazza-Martin et al., 2008). Focusing on a limited number of muscles is problematic, as we risk probing muscles whose activation patterns are not affected by age. To ensure that our restrictive theory-driven analysis provides meaningful results, we verified that our cherry-picked muscles truly conveyed information about age-related modifications of whole-body movements control. To this aim, we employed machine learning analyses that quantified how much each muscle activation was altered by age. This allowed controling that we were actually focusing on muscles that discriminated movement control between younger and older adults.

Our rationale was the following: If the algorithm can successfully separate the data of younger and older adults, using antigravity muscle activation patterns, this would demonstrate that important information is contained in those muscles regarding age-related modifications of movement control. For more details on similar use and operation of machine learning algorithms on EMG signals, please see (Chambellant et al., 2024; Thomas et al., 2023; Tolambiya et al., 2011). Here we present the results of a Linear Discriminant Analysis (LDA, Johnson & Wichern, 1988) but we verified that we obtained similar conclusions with two other algorithms, namely the Quadratic Discriminant Analysis (QDA, Cover, 1965) and the Support Vector Machine (SVM, Vapnik and Lerner, 1965). We used custom Matlab scripts [Mat, 2021] to perform all machine learning analyses. The ESL1G was not considered for these analyses because the electrode was defective for several younger participants (see Supplementary Table 1).

The input data was the phasic EMG signals of the 15 muscles taken individually or the whole set at once. These vectors were fed to the algorithms using binary classification setups, where the algorithm learned to distinguish between the EMGs of the two groups. To ensure robustness of the results, we employed a five-fold cross-validation method. This involved splitting the whole dataset into five sets while ensuring equal representation of both directions in each set. The algorithm was trained on four of those sets before being evaluated on the fifth set (containing data unknown to the trained algorithm). This operation was repeated five times so each set was tested once. Cross-validation allowed computing the average accuracy and its variance across the testing sets, thereby providing a reliable estimate of the accuracy obtained by the algorithm. Finally, we could compare the accuracy of the algorithm for each muscle.

Univariate Statistics

After an initial kinematic analysis (detailed in the results section), we observed a difference in movement duration between younger and older adults (conducting a repeated measure analyses of variance with a between factor *Age* with two levels: Young/Older and a within factor *Task-type* with two levels: Arm/Whole-body movements). Because movement duration is known to influence phasic EMG negativity (Poirier et al., 2023), we added movement duration as a covariate. We performed repeated measure analyses of covariance (ANCOVA) using JASP software. Two ANCOVA analyses were carried out. We first used a mixed ANCOVA with a between factor *Age* (two levels: Young/Older) and a within factor *Task-type* (two levels: Arm/Whole-body movements) to test whether age effects on movement control depended on the type of task being performed. Second, to detail the age differences observed during movements of the entire body, we used a mixed ANCOVA with a between factor *Age* (two levels: Young/Older) and a within factor *Whole-Body-Tasks* (three levels: STS_BTS/WBR D1/WBR D2). In all cases, the significance level was set to 0.05.

341   To test for a possible beneficial effect (i.e., a compensation) of the EMG alterations that we observed with
342   aging, we performed a kinematic analysis of the center of mass. We then used independent Student-tests and
343   Pearson correlation coefficients to study potential differences between groups and associations between
344   variables.

Results

Movement duration of fast movements varied between tasks and was slightly reduced in younger
compared to older participants (see Figure 4 and Supplementary Table 2 for detailed results). Overall older
adults were 3.5% slower than younger adults. A repeated measures ANOVA revealed that this age-difference
was significant ($F_{(1,42)}$= 14.5, P=4.58E-05, η^2=0.256). For this reason, we used movement duration as a
covariate in the following statistical analyses. Nevertheless, as revealed by Figure 4, it is important to note that
an important number of older adults moved with durations that were similar to those of younger adults.

Figure 4. Mean ± SD movement durations (s) for fast movements performed in all tasks and both groups (STS: Seat-to-stand, BTS: Back-to-seat, WBR: Whole-body-reaching, D1: Short distance=15% of the heigh of the subject and D2: Long distance=30% of the heigh of the subject). Each point correponds to the average duration of the trials of one participant. The blue points represent the young participants and the green points correpond to the older participants.

Figure 5 displays average phasic EMG profiles for each muscle, direction, and task. As recently reported,
phasic EMG signals of arm movements show negative phases during the deceleration of upward and the
acceleration of downward arm movements, i.e., where gravity torque helps generate the arm's motion
(Gaveau et al., 2021; Poirier et al., 2022, 2024). Previous works demonstrated that this negativity is not erratic
but systematic and indicate that muscles contract less than necessary to compensate for gravity effects. It is
therefore especially prominent on antigravity muscles and reveals that the central nervous system (CNS)
exploits gravity effects to produce efficient movements, i.e., motor patterns that save unnecessary muscle
effort. Here, we extend this result to movements performed with the entire body. Indeed, for STS/BTS and
WBR movements, Figure 2B-D unveils phasic EMG negativity during the deceleration of upward movements
and the acceleration of downward movements, i.e. when gravity can help produce the motion. This first
qualitative result demonstrates that movements that are performed with the entire body, similarly to more
focal arm movements, exploit gravity effects to save unnecessary muscle effort (Gaveau et al., 2021). More
importantly, the present results qualitatively reveal that older adults also use such an efficient strategy, both
when moving their arm and their entire body.

The present study aimed to test for possible age-differences in the extent to which older and younger adults use gravity effects to minimize muscle effort. To ensure that restricting the analysis to antigravity muscles would nonetheless provide meaningful information on how age alters muscle activation patterns, we used a machine learning approach. This analysis indeed revealed that antigravity muscles contained important information, allowing separating age-groups with some of the best success-rates (see Figure 6 for results regarding LDA accuracy). The vastus lateralis (VL) and the spinal erectors on L1 (ESL1) achieve classification accuracies of 57.72% and 59.51% respectively (considering that these classifications are significantly better than chance if they are above 52.5% according to a fairness test).Thus, building on the complementary results of the theory-driven approach (Chambellant et al., 2024; Gaveau et al., 2021; Poirier et al., 2022, 2024; Thomas et al., 2023) and the present data-driven control analysis (Figure 6), in the following we focus the analysis on

Figure 5: Mean (±SE) integrated phasic EMGs recorded for both groups (n = 20 for younger and n = 24 for older) during arm (**panel A**), Seat-to-stand/Back-to-seat (**panel B**), and whole body reaching movements (**panel C** for short distance D1 and **panel D** for long distance D2). Blue traces present EMGs recorded for younger participants, while green traces present EMGs recorded for older participants. The dotted line divides the movement in two: the first half is acceleration and the second is deceleration. (DA: Anterior deltoid, DP: Posterior deltoid, VL: Vastus Lateralis, BF: Biceps Femoris, ESL1: Erector Spinae in L1, EST7: Erector Spinae in T7, TA: Tibialis Anterior, SOL: Soleus).

Figure 6. A. Heatmap representing the accuracy of the LDA algorithm to discriminate young from older adults using phasic EMGs recorded during the tasks mobilizing the entire body. The height first lines correspond to the accuracy for each individual muscle. The last line corresponds to the accuracy taking all muscles simultaneously (DA: Anterior deltoid, DP: Posterior deltoid, VL: Vastus Lateralis, BF: Biceps Femoris, ESL1: Erector Spinae in L1, EST7: Erector Spinae in T7, TA: Tibialis Anterior, SOL: Soleus). The first six columns correspond to the six whole-body tasks of the experiment (STS = Sit To Stand; BTS = Back To Sit; WBR D1 = Whole Body Reaching near target; WBR D2 = Whole body reaching far target), the last column corresponds to the average accuracy of the six tasks for each muscle. This analysis has been conducted to showcase which muscles are important for discrimination. We can see those antigravity muscles (muscles that act against gravity, in our case, the VL and the ESL1) contain relevant information as they reach the highest scores. **B**. Mean ± SD distance between groups representations by the LDA algorithm for each individual muscle (DA: Anterior deltoid; DP: Posterior deltoid; VL: Vastus Lateralis; BF: Biceps Femoris; ESL1: Erector Spinae in L1; EST7: Erector Spinae in T7; TA: Tibialis Anterior; SOL: Soleus). The higher the distance, the more differentiable the groups are. Values corresponds to the average distance obtain for all six tasks. We can see those antigravity muscles (muscles that act against gravity, in our case, the VL and the ESL1) contain relevant information as they reach some of the highest values. Error bars correspond to standard error across a five-fold cross-validation.

Main analysis

Following our primary hypothesis, we first analyzed a single metric quantifying phasic EMG negativity on
an average muscle activation pattern (vastus lateralis and erector spinae in L1 were averaged for whole-body
tasks and deltoid anterior was used for arm tasks), namely the negative area of phasic EMG patterns (see
methods and (Poirier et al., 2022, 2024). The bigger the negativity index, the more efficient the muscle
contractions, in the sense that gravity effects were maximally exploited to minimize muscle effort (Gaveau et
al., 2021). Figure 7 displays the results of this ANCOVA analysis (Age × Task-Type), revealing a significant
interaction between age and task factors ($F_{(1,42)}$= 5.48, P=2.44E-02, η^2=0.120) but no Age or Tasks effect (for
detailed statistical results, please see Supplementary Table 3). This result demonstrates that age differently
alters motor strategies for arm movements vs whole-body movements. Older adults used gravity effects to a
similar extent as younger ones when performing arm movements (older adults, mean ± SD: -10.7 ± 5.6, 95%
CI: [-8.4;-13.0]; younger adults, -11.4 ± 3.6, [-9.8;-13.0]), but to a lesser extent when performing whole-body
movements (older adults, -9.7 ± 3.2, [-8.0;-11.5]; younger adults, -15.6 ± 3.3, [-14.1;-17.0]). As recently
reported by Poirier et al. (2024), similar arm results in younger and older adults suggest that the ability to plan
movements that optimally use gravity effects to minimize muscle effort remains functional in older adults. The
results obtained in whole-body movement tasks (STS/BTS and WBR) could thus suggest that the difference
observed between older and younger adults does not reflect a deterioration of the ability to plan movements
that are optimally adapted to the gravity environment. Instead, it would suggest a change in movement

strategy that compensates for other deteriorated control processes (for example, the loss of muscle mass &
force).
We performed a complementary analysis to determine whether every whole-body tasks showed the same
age-effect (ANCOVA Age x Whole-Body-Tasks). This test did not reveal any interaction effect ($F_{(2,42)}$= 0.77,
P=4.67E-01, η^2=0.019), further supporting the interpretation that this is the "whole body" aspect of the task
that impacts the motor strategy in older adults (please see Supplementary Table 3 for full analysis).
Previous studies have proposed that the change in kinematic strategies observed between older and
younger adults during whole-body movements could be explained as a strategy maximizing equilibrium
maintenance rather than muscular effort (Casteran et al., 2018; Paizis et al., 2008). Following this hypothesis,
one would predict increasing differences between younger and older adults when the equilibrium constraint
increases. In the present experiment, increased equilibrium constraint was produced by increasing the target
distance during whole body reaching movements (WBR D1 vs WBR D2; alike Casteran et al., 2018). The Age x
Whole-Body-Tasks ANCOVA however did not reveal such a difference.

Figure 7. Negativity index computed for **A**. arm and whole-body movements in both groups (WB: Whole Body combines seat-to-stand/back-to-seat, whole-body reaching from D1 and whole-body reaching from D2) and **B**. each whole-body tasks (STS/BTS: seat-to-stand/back-to-sit, WBR D1: whole-body reaching from D1 and WBR D2: whole-body reaching from D2). The negativity index, defined as T x NA / TA, with NA the Negative Area integrated on the phasic signal between negativity onset and offset, TA the Tonic Area integrated on the tonic signal between the negativity onset and offset, and T the duration of the negative epoch normalized by movement duration. The blue points correspond to the younger participants and the green points correpond to the older participants. Each point corresponds to the mean value of one participant (mean accross trials and antigravity muscles and/or tasks).

Last, we analysed kinematic patterns in order to investigate whether the decreased effort minimization
observed during whole body tasks in older adults could actually be interpreted as a compensation. We tested
whether the negativity of phasic EMGs correlated with kinematic parameters that are related to balance
control (the COM displacement, and COM peak velocity, see Figure 8 and see Supplementary Figure 1 for
detailed results of the reproduction of the tests conducted by Casteran et al., (2018) Paizis et al., (2008). The
EMG criterion during the Back to Seat task was found significantly correlated with the COM displacement
(Pearson correlation, P=2.2E-2, Pearson's r=-0.343) and the COM peak velocity (Pearson correlation, P=1.9E-
3, Pearson's r=-0.476). This same EMG criterion also turned out to be significantly correlated for the Whole-
Body Bending task with the COM displacement (Pearson correlation, P=3.2E-3, Pearson's r=-0.435) and with
the COM peak velocity (Pearson correlation, P=1.2E-7, Pearson's r=-0.700). The linear regressions revealed
that the more a participant used the effects of gravity, the more and the quicker she/he displaced his COM.
One could interpret this result as demonstrating that older adults lose the ability to plan energetically efficient
movement and, thus, move their whole-body less and more slowly. However, the null age effect on arm
movement control supports the hypothesis that planning efficient movements remains functional in older
adults, as also supported by recent other results (Healy et al., 2023; Huang & Ahmed, 2014; Poirier et al., 2020;
Summerside et al., 2024). Overall, during movements performed with the entire body, i.e., when equilibrium
maintenance is challenged, the present results support an age-related adaptation process that selects a less
energetically efficient but more stable movement strategy in healthy older adults.

Exploratory analyses

To provide a fine-grained analysis of the age effect on phasic EMG negativity during whole-body motion, we probed negativity duration, negativity amplitude, and negativity occurrence across tasks and age-groups (see Figure 9). Here also, the bigger the values, the bigger the use of gravity effects to produce body motion. A repeated measures ANCOVA Age x Tasks (Young/Older x STS_BTS/WBRD1/WBRD2) revealed a significant age effect where negativity duration was larger in younger compared to older participants ($F_{(1,36)}$= 21.49, P=4.54E-05, η^2=0.374). The age effect did not reach significance for negativity occurrence ($F_{(1,36)}$= 3.62, P=0.065, η^2=0.091) nor for negativity amplitude ($F_{(1,36)}$= 1.16, P=0.28, η^2=0.031). No interaction between Age and the other factors reached significance (see Supplementary Table 3 for detailed results). Overall, all variables showed qualitatively smaller negativity on phasic EMGs, thus reduced use of gravity effects, in older compared to younger adult. As already observed for arm movements (Poirier et al. 2024), it is mainly the duration of inactivation that is modulated.

Figure 8. Center of mass analyses. Linear relationship between EMG negativity (Vastus Lateralis and Spinal Erector L1) and total displacement of the center of mass during **A**. back-to-sit movements and **B**. bending movements from the whole-body reaching tasks (averaged between distances D1 and D2).

Figure 9. Mean ± SD negativity quantification for all tasks and groups (STS: Seat-to-stand, BTS: Back-to-seat, WBR: Whole-body-reaching, D1: Short distance=15% of the heigh of the subject and D2: Long distance=30% of the heigh of the subject). Quantification was carried out using three criteria: negativity duration (panel **A**), negativity amplitude (panel **B**) an negativity occurrence (panel **C**) on the antigravity muscles (vastus lateralis and the erector spinae). The blue points correspond to the younger participants and the green points correspond to the older participants. Each point corresponds to the mean value accross trials and antigravity muscles.

Discussion

In younger and older adults, we investigated the muscle activation patterns responsible for arm and
whole-body movements. The results revealed an age-related alteration of muscle activation that differ
between the type of tasks. Comparing older adults to younger ones, we found that a muscle marker of
effort minimization was reduced during whole-body movements but the results show no evidence that this
marker was reduced during arm movements. Previous works have demonstrated that this marker allows
quantifying the output of a sensorimotor control process that adapts human movements to gravity
(Gaveau et al., 2021; Poirier et al., 2022, 2024). More precisely, this marker allows for quantifying how
much one harvests gravity effects to minimize muscle effort. Here, arm movement results reveal that this
effort-discounting process remains functional in older adults. During whole-body movements, however,
the present results reveal that a criteria linked to effort-minimization was downregulated in older adults
compared to younger adults. Overall, the present results suggest a compensation process that modulates
planning strategies to maximize equilibrium in older adults.

Age-related compensatory processes in sensorimotor control. A number of studies have proposed
that the differences observed between younger and older adults can be interpreted as compensations for
age-related deteriorations. Of particular interest are studies from the last decade that have sought to
investigate specific motor control processes rather than broad motor performance. For example, some of
these studies indirectly suggested that older adults favor feedforward rather than feedback control (Moran
et al., 2014; Wolpe et al., 2016) to compensate for the attenuation of sensory processing with increasing
age (Moran et al., 2014; Parthasharathy et al., 2022; Saenen et al., 2023). Others indirectly suggested that
older adults favor movement efficiency over precision (Healy et al., 2023; Poirier et al., 2020) to
compensate for their increased energetic cost (Didier et al., 1993; Hortobagyi et al., 2011; John et al., 2009).
Yet, because the focus of these studies was not on compensatory processes, they did not include dedicated
experimental conditions. The aim of the present study was to fill this gap.

Maintained efficiency of arm movements in older adults. The metabolic rate is known to influence
resource use, body size, rate of senescence, and survival probability (Brown et al., 2004; DeLong et al.,
2010; Strotz et al., 2018; Van Voorhies & Ward, 1999). The nervous system has therefore developed the
ability to design movement strategies that minimize our every-day efforts (Cheval et al., 2018; Gaveau et
al., 2016; Huang et al., 2012; Morel et al., 2017; Selinger et al., 2015; Shadmehr et al., 2016). The present
findings confirm the results of previous arm movement studies that proposed a theory according to which
motor control takes advantage of gravity effects to minimize muscle effort (Berret et al., 2008; Crevecoeur
et al., 2009; Gaveau et al., 2014, 2016, 2021; Gaveau & Papaxanthis, 2011). Here, we focused on the muscle
activation marker of gravity-related effort minimization, i.e. the negativity of phasic EMG. Previous
modeling and experimental work demonstrated that this phasic EMG negativity results from an optimal
control process that plans efficient arm movements in the gravity field (Gaveau et al., 2021). As reported
by (Poirier et al., 2024), we found similar phasic EMG negativity during arm movements in older and
younger adults. Thus, arm movements equally optimized gravity effects in younger and older adults. These
results align with those of studies that probed progressive motor adaptation to a new environment in older
adults. Using locally induced force fields in a robotic environment, these studies revealed that older adults
decreased their metabolic costs similarly to younger adults while adapting to new environmental dynamics
(Healy et al., 2023; Huang & Ahmed, 2014). Overall, results from arm movement studies advocate for the
maintenance of the ability to optimally integrate environmental dynamics and plan arm movements that
minimize muscle effort in older adults.

Whole-body movements also harvests gravity effects to minimize muscle effort. Current results also
extend the current knowledge on the planning of energetically efficient movements to more global
movements, both in younger and older adults. They unravel that deactivating muscles, below the tonic
level that would be necessary to compensate for external dynamics are not only relevant to control focal
arm movement but also for whole-body movements. Using a combination of modeling and experimental
work, previous studies demonstrated that healthy participants move their arms following trajectories and
using muscular patterns that minimize muscle effort in the gravity environment (Berret et al., 2008;
Crevecoeur et al., 2009; Gaveau et al., 2014, 2016, 2021; Gaveau & Papaxanthis, 2011). To isolate gravity
effects, most studies focused on one-degree-of-freedom arm movements. Although those studies allowed

us to clearly demonstrate how motor planning integrates gravity effects into motor planning, one-degree-of-freedom movements are hardly representative of the rich and complex human movement repertoire. The present study, using more ecological movements, basically extends the optimal integration of gravity effects theory to every-day movements.

Decreased efficiency of whole-body movements in older adults. Contrary to focal arm movements, we observed a strong age difference during global movements that engaged the entire body, here sit to stand / back to sit and whole-body reaching movements. Specifically, the negativity of phasic EMG was significantly reduced in older compared to younger adults. This suggests that whole-body movements are less energetically efficient in older adults than in younger ones, adding to the general result that global movements are more energy-demanding for older adults compared to younger adults (Didier et al., 1993; Hortobagyi et al., 2003, 2011; John et al., 2009). Previous kinematic studies suggested that older adults favor movement strategies that maximize balance maintenance rather than effort minimization (Casteran et al., 2018; Paizis et al., 2008). However, age differences observed during whole-body movements may also be interpreted as an inability to minimize muscle effort when coordinating complex movements (Goodpaster et al., 2006; Henry & Baudry, 2019; Quinlan et al., 2018; Vernazza-Martin et al., 2008). Here, contrasting results from arm and whole-body movements in the same participants, we provide support for a compensation process that adapts movement strategy in older adults, rather than a deterioration of the ability to optimally coordinate whole-body movements. Since arm movements revealed that older participants maintained the ability to plan energetically efficient movements, altered whole-body movement may be explained as an adaptation of movement strategy rather than a deteriorated motor planning. More, we found that decreased efficiency was associated to decreased center-of-mass displacement and speed, i.e. less instability. This further suggests that increased effort in older adults is a compensation process that trades movement efficiency with equilibrium maintenance. This could be explained as an optimal motor planning process that minimizes a composite cost function; i.e. effort and unstability. It has been proposed that the central nervous system combines different costs – related to energy, precision, or duration for example – when planning a movement (Berret et al., 2011; Gielen, 2009; Healy et al., 2023; Liu & Todorov, 2007; Mombaur et al., 2010; Poirier et al., 2024; Tanis et al., 2023; Vu et al., 2016). In older adults, this combination would increase the relative weighting of the unstability (equilibrium) cost and decrease the relative weighting of the effort cost. Future work may use this framework to probe age-related motor adaptation.

Effect of target distance. During the whole-body reaching task, reusing the protocol (Casteran et al., 2018), we varied the antero-posterior distance of the target to be reached. Casteran et al. (2018) found larger differences between younger and older participants when the target was further. Consequently, we hypothesized that the further away the target, the greater the age differences in the negativity epochs of phasic EMGs. The present results do not validate this hypothesis (see Supplementary Table 3).

Age-related compensation in the brain. In the sensorimotor field, following the consensus that aging is associated with increased activation and increased spatial recruitment, numerous studies have attempted to establish a correlation between brain activation and behavioral performance in older adults (for reviews, see Fettrow et al., 2021; Poirier et al., 2021; Seidler et al., 2010; Ward, 2006). This literature has not reached a consensus on the neural changes underlying compensatory mechanisms in the aging brain. Several studies reported a positive correlation (Cassady, Gagnon, et al., 2020; Clark et al., 2014; Harada et al., 2009; Heuninckx et al., 2008; Holtzer et al., 2015; Jor'dan et al., 2017; Larivière et al., 2019; Mattay et al., 2002; Spedden et al., 2019), and as many reported no correlation or even a negative correlation (Bernard & Seidler, 2012; Cassady et al., 2019; Cassady, Ruitenberg, et al., 2020; Fernandez et al., 2019; Hawkins et al., 2018; Holtzer et al., 2016; Loibl et al., 2011; Riecker et al., 2006; Ward et al., 2008). Building on the theoretical work of Krakauer et al. (2017), we recently proposed that an important reason for this lack of consensus may be that previous studies, while focusing on brain activations, used crude behavioral paradigms that likely mixed deteriorated and compensatory processes (Poirier et al., 2021). Using behavioral paradigms that focus on specific motor control processes, as performed here, could help differentiating compensatory mechanisms from deteriorative ones.

Role of physical and cogntive fitness in age-related compensation. Physical and cognitive fitness may influence how much older adults favor stability over energetic efficiency. It is well-known that physical and cognitive fitness significantly impact functional mobility in older adults (Marusic et al., 2018; Wickramarachchi et al., 2023; Zhao et al., 2014). One could speculate that physical and cognitive fitness is

inversely related to the level of physical and cognitive deterioration. For example, muscle force and sensory integration are crucial for controlling balance. The more deteriorated they are, the more the need for compensatory process to adapt movement control to the participant capacities. Future research should account for variations in physical and cognitive fitness to better understand their role in the development of compensatory mechanisms.

Simple mono-articular vs complex multi-articular arm movements. Another aspect that needs to be highlighted here is the choice of the arm task, which is not representative of all existing tasks for studying upper limb motor skills. Using this very same task, the results from two previous studies also support the preservation of arm movement efficiency in older adults (Poirier et al., 2020, 2024). One may wonder whether the present conclusions would hold for more complex arm movements. Using multi-degree of freedom arm movements to study motor adaptation to an externally imposed force-field, other studies also reported results showing that, alike younger adults, older adults maintain the ability to produce movements that are energetically efficient (Healy et al., 2023; Summerside et al., 2024). The present mono-articular results therefore likely generalize to other types of arm movements. In young adults, the efficient integration of gravity effects to minimize muscle effort has been demonstrated with varied arm movements, such as single or muti-degree of freedom pointing movements, drawing movements, reach to grasp movements, or arm movements that transport a hand-grasped object (Berret et al., 2008; Crevecoeur et al., 2009; Gaveau et al., 2011; Gaveau & Papaxanthis, 2011; Le Seac'h & McIntyre, 2007; Papaxanthis et al., 1998, 2005; Yamamoto & Kushiro, 2014b). Futurre work may test whether the present conclusions extend to more complex and functional arm movements.

In conclusion, probing a specific motor control process, the present study provides a set of behavioral results that support the interpretation of a compensatory process that counterbalances other deteriorated processes in older adults. Probing age effects on specific sensorimotor control processes may help disentangle compensation from deterioration processes that occur through healthy aging (Poirier et al., 2021). We believe that understanding compensation at a behavioral level is an important step toward pinpointing its neural underpinning (Krakauer et al., 2017) and, later, preventing unhealthy aging (Baltes & Baltes, 1990; Martin et al., 2015; Zhang & Radhakrishnan, 2018).

Acknowledgements

We thank Yves Ballay, Denis Barbusse, and Gabriel Poirier for their support during the pilot study. We also thank all the participants who took part in the experiment.

Data, scripts, code, and supplementary information availability

Data are available online: 10.5281/zenodo.10619701, webpage hosting the data: https://doi.org/10.5281/zenodo.10619701 (*citation of the data eg* Mathieu et al, 2024);
Scripts and code are available online: 10.5281/zenodo.10634004, webpage hosting the scripts: https://doi.org/10.5281/zenodo.10634004 (*citation of the scripts eg* Mathieu et al, 2024);
Supplementary information is available online: 10.5281/zenodo.10671496, webpage hosting the file: https://doi.org/10.5281/zenodo.12671953 (*citation of the supplementary file eg* Mathieu et al, 2024);

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.
Jérémie Gaveau is a member of the managing board of the PCI Health & Movement Sciences.

Funding

This entire study is part of a thesis funded by the National Research Agency (ANR I-SITE BFC).

References

Baltes, P. B. (1997). On the incomplete architecture of human ontogeny : Selection, optimization, and compensation as foundation of developmental theory. *American Psychologist*, *52*(4), 366-380. https://doi.org/10.1037/0003-066X.52.4.366

Baltes, P. B., & Baltes, M. M. (1990). Psychological perspectives on successful aging : The model of selective optimization with compensation. In P. B. Baltes & M. M. Baltes (Éds.), *Successful Aging* (1^re éd., p. 1-34). Cambridge University Press. https://doi.org/10.1017/CBO9780511665684.003

Barbero, M., Merletti, R., & Rainoldi, A. (2012). *Atlas of Muscle Innervation Zones*. Springer Milan. https://doi.org/10.1007/978-88-470-2463-2

Barulli, D., & Stern, Y. (2013). Efficiency, capacity, compensation, maintenance, plasticity : Emerging concepts in cognitive reserve. *Trends in Cognitive Sciences*, *17*(10), 502-509. https://doi.org/10.1016/j.tics.2013.08.012

Bernard, J. A., & Seidler, R. D. (2012). Evidence for motor cortex dedifferentiation in older adults. *Neurobiology of Aging*, *33*(9), 1890-1899. https://doi.org/10.1016/j.neurobiolaging.2011.06.021

Berret, B., Bonnetblanc, F., Papaxanthis, C., & Pozzo, T. (2009). Modular Control of Pointing beyond Arm's Length. *Journal of Neuroscience*, *29*(1), 191-205. https://doi.org/10.1523/JNEUROSCI.3426-08.2009

Berret, B., Chiovetto, E., Nori, F., & Pozzo, T. (2011). Evidence for Composite Cost Functions in Arm Movement Planning : An Inverse Optimal Control Approach. *PLoS Computational Biology*, *7*(10), e1002183. https://doi.org/10.1371/journal.pcbi.1002183

Berret, B., Darlot, C., Jean, F., Pozzo, T., Papaxanthis, C., & Gauthier, J. P. (2008). The Inactivation Principle : Mathematical Solutions Minimizing the Absolute Work and Biological Implications for the Planning of Arm Movements. *PLoS Computational Biology*, *4*(10), e1000194. https://doi.org/10.1371/journal.pcbi.1000194

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., & West, G. B. (2004). TOWARD A METABOLIC THEORY OF ECOLOGY. *Ecology*, *85*(7), 1771-1789. https://doi.org/10.1890/03-9000

Buckles, V. D. (1993). Age-Related Slowing. In G. E. Stelmach & V. Hömberg (Éds.), *Sensorimotor Impairment in the Elderly* (p. 73-87). Springer Netherlands. https://doi.org/10.1007/978-94-011-1976-4_6

Buneo, C. A., Soechting, J. F., & Flanders, M. (1994). Muscle activation patterns for reaching : The representation of distance and time. *Journal of Neurophysiology*, *71*(4), 1546-1558. https://doi.org/10.1152/jn.1994.71.4.1546

Bunzeck, N., Steiger, T. K., Krämer, U. M., Luedtke, K., Marshall, L., Obleser, J., & Tune, S. (2024). Trajectories and contributing factors of neural compensation in healthy and pathological aging. *Neuroscience & Biobehavioral Reviews*, *156*, 105489. https://doi.org/10.1016/j.neubiorev.2023.105489

Cassady, K., Gagnon, H., Freiburger, E., Lalwani, P., Simmonite, M., Park, D. C., Peltier, S. J., Taylor, S. F., Weissman, D. H., Seidler, R. D., & Polk, T. A. (2020). Network segregation varies with neural distinctiveness in sensorimotor cortex. *NeuroImage*, *212*, 116663. https://doi.org/10.1016/j.neuroimage.2020.116663

Cassady, K., Gagnon, H., Lalwani, P., Simmonite, M., Foerster, B., Park, D., Peltier, S. J., Petrou, M., Taylor, S. F., Weissman, D. H., Seidler, R. D., & Polk, T. A. (2019). Sensorimotor network segregation declines with age and is linked to GABA and to sensorimotor performance. *NeuroImage*, *186*, 234-244. https://doi.org/10.1016/j.neuroimage.2018.11.008

Cassady, K., Ruitenberg, M. F. L., Reuter-Lorenz, P. A., Tommerdahl, M., & Seidler, R. D. (2020). Neural Dedifferentiation across the Lifespan in the Motor and Somatosensory Systems. *Cerebral Cortex*, *30*(6), 3704-3716. https://doi.org/10.1093/cercor/bhz336

Casteran, M., Hilt, P. M., Mourey, F., Manckoundia, P., French, R., & Thomas, E. (2018). Shifts in Key Time Points and Strategies for a Multisegment Motor Task in Healthy Aging Subjects. *The Journals of Gerontology: Series A*, *73*(12), 1609-1617. https://doi.org/10.1093/gerona/gly066

Chambellant, F., Gaveau, J., Papaxanthis, C., & Thomas, E. (2023). *Deactivation and Collective Phasic Muscular Tuning for Pointing Direction : Insights from Machine Learning* [Preprint]. Neuroscience. https://doi.org/10.1101/2023.03.15.532800

Chambellant, F., Gaveau, J., Papaxanthis, C., & Thomas, E. (2024). Deactivation and Collective Phasic Muscular Tuning for Pointing Direction : Insights from Machine Learning. *Heliyon*, e33461. https://doi.org/10.1016/j.heliyon.2024.e33461

Cheval, B., Tipura, E., Burra, N., Frossard, J., Chanal, J., Orsholits, D., Radel, R., & Boisgontier, M. P. (2018). Avoiding sedentary behaviors requires more cortical resources than avoiding physical activity : An EEG study. *Neuropsychologia*, *119*, 68-80. https://doi.org/10.1016/j.neuropsychologia.2018.07.029

Clark, D. J., Christou, E. A., Ring, S. A., Williamson, J. B., & Doty, L. (2014). Enhanced Somatosensory Feedback Reduces Prefrontal Cortical Activity During Walking in Older Adults. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, *69*(11), 1422-1428. https://doi.org/10.1093/gerona/glu125

Cover, T. M. (1965). Geometrical and Statistical Properties of Systems of Linear Inequalities with Applications in Pattern Recognition. *IEEE Transactions on Electronic Computers*, *EC-14*(3), 326-334. https://doi.org/10.1109/PGEC.1965.264137

Crevecoeur, F., Thonnard, J.-L., & Lefèvre, P. (2009). Optimal Integration of Gravity in Trajectory Planning of Vertical Pointing Movements. *Journal of Neurophysiology*, *102*(2), 786-796. https://doi.org/10.1152/jn.00113.2009

Darling, W. G., Cooke, J. D., & Brown, S. H. (1989). Control of simple arm movements in elderly humans. *Neurobiology of Aging*, *10*(2), 149-157. https://doi.org/10.1016/0197-4580(89)90024-9

d'Avella, A., Fernandez, L., Portone, A., & Lacquaniti, F. (2008). Modulation of Phasic and Tonic Muscle Synergies With Reaching Direction and Speed. *Journal of Neurophysiology*, *100*(3), 1433-1454. https://doi.org/10.1152/jn.01377.2007

d'Avella, A., Portone, A., Fernandez, L., & Lacquaniti, F. (2006). Control of Fast-Reaching Movements by Muscle Synergy Combinations. *Journal of Neuroscience*, *26*(30), 7791-7810. https://doi.org/10.1523/JNEUROSCI.0830-06.2006

DeLong, J. P., Okie, J. G., Moses, M. E., Sibly, R. M., & Brown, J. H. (2010). Shifts in metabolic scaling, production, and efficiency across major evolutionary transitions of life. *Proceedings of the National Academy of Sciences*, *107*(29), 12941-12945. https://doi.org/10.1073/pnas.1007783107

Didier, J. P., Mourey, F., Brondel, L., Marcer, I., Milan, C., Casillas, J. M., Verges, B., & Winsland, J. K. D. (1993). The Energetic Cost of Some Daily Activities : A Comparison in a Young and Old Population. *Age and Ageing*, *22*(2), 90-96. https://doi.org/10.1093/ageing/22.2.90

Fernandez, N. B., Hars, M., Trombetti, A., & Vuilleumier, P. (2019). Age-related changes in attention control and their relationship with gait performance in older adults with high risk of falls. *NeuroImage*, *189*, 551-559. https://doi.org/10.1016/j.neuroimage.2019.01.030

Fettrow, T., Hupfeld, K., Reimann, H., Choi, J., Hass, C., & Seidler, R. (2021). Age differences in adaptation of medial-lateral gait parameters during split-belt treadmill walking. *Scientific Reports*, *11*(1), 21148. https://doi.org/10.1038/s41598-021-00515-z

Flanders, M., & Herrmann, U. (1992). Two components of muscle activation : Scaling with the speed of arm movement. *Journal of Neurophysiology*, *67*(4), 931-943. https://doi.org/10.1152/jn.1992.67.4.931

Flanders, M., Pellegrini, J. J., & Soechting, J. F. (1994). Spatial/temporal characteristics of a motor pattern for reaching. *Journal of Neurophysiology*, *71*(2), 811-813. https://doi.org/10.1152/jn.1994.71.2.811

Gaveau, J., Berret, B., Angelaki, D. E., & Papaxanthis, C. (2016). Direction-dependent arm kinematics reveal optimal integration of gravity cues. *eLife*, *5*, e16394. https://doi.org/10.7554/eLife.16394

Gaveau, J., Berret, B., Demougeot, L., Fadiga, L., Pozzo, T., & Papaxanthis, C. (2014). Energy-related optimal control accounts for gravitational load : Comparing shoulder, elbow, and wrist rotations. *Journal of Neurophysiology*, *111*(1), 4-16. https://doi.org/10.1152/jn.01029.2012

Gaveau, J., Grospretre, S., Berret, B., Angelaki, D. E., & Papaxanthis, C. (2021). A cross-species neural integration of gravity for motor optimization. *Science Advances*, *7*(15), eabf7800. https://doi.org/10.1126/sciadv.abf7800

Gaveau, J., Paizis, C., Berret, B., Pozzo, T., & Papaxanthis, C. (2011). Sensorimotor adaptation of point-to-point arm movements after spaceflight : The role of internal representation of gravity force in trajectory planning. *Journal of Neurophysiology*, *106*(2), 620-629. https://doi.org/10.1152/jn.00081.2011

Gaveau, J., & Papaxanthis, C. (2011). The Temporal Structure of Vertical Arm Movements. *PLoS ONE*, *6*(7), e22045. https://doi.org/10.1371/journal.pone.0022045

Gentili, R., Cahouet, V., & Papaxanthis, C. (2007). Motor planning of arm movements is direction-dependent in the gravity field. *Neuroscience*, *145*(1), 20-32. https://doi.org/10.1016/j.neuroscience.2006.11.035

Gielen, S. (2009). Review of Models for the Generation of Multi-Joint Movements in 3-D. In D. Sternad (Éd.), *Progress in Motor Control* (Vol. 629, p. 523-550). Springer US. https://doi.org/10.1007/978-0-387-77064-2_28

Goble, D. J., Coxon, J. P., Wenderoth, N., Van Impe, A., & Swinnen, S. P. (2009). Proprioceptive sensibility in the elderly : Degeneration, functional consequences and plastic-adaptive processes. *Neuroscience & Biobehavioral Reviews*, *33*(3), 271-278. https://doi.org/10.1016/j.neubiorev.2008.08.012

Goodpaster, B. H., Park, S. W., Harris, T. B., Kritchevsky, S. B., Nevitt, M., Schwartz, A. V., Simonsick, E. M., Tylavsky, F. A., Visser, M., Newman, A. B., & for the Health ABC Study. (2006). The Loss of Skeletal Muscle Strength, Mass, and Quality in Older Adults : The Health, Aging and Body Composition Study. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, *61*(10), 1059-1064. https://doi.org/10.1093/gerona/61.10.1059

Gueugneau, N., Martin, A., Gaveau, J., & Papaxanthis, C. (2023). Gravity-efficient motor control is associated with contraction-dependent intracortical inhibition. *iScience*, *26*(7), 107150. https://doi.org/10.1016/j.isci.2023.107150

Harada, T., Miyai, I., Suzuki, M., & Kubota, K. (2009). Gait capacity affects cortical activation patterns related to speed control in the elderly. *Experimental Brain Research*, *193*(3), 445-454. https://doi.org/10.1007/s00221-008-1643-y

Hawkins, K. A., Fox, E. J., Daly, J. J., Rose, D. K., Christou, E. A., McGuirk, T. E., Otzel, D. M., Butera, K. A., Chatterjee, S. A., & Clark, D. J. (2018). Prefrontal over-activation during walking in people with mobility deficits : Interpretation and functional implications. *Human Movement Science*, *59*, 46-55. https://doi.org/10.1016/j.humov.2018.03.010

Healy, C. M., Berniker, M., & Ahmed, A. A. (2023). Learning vs. minding : How subjective costs can mask motor learning. *PLOS ONE*, *18*(3), e0282693. https://doi.org/10.1371/journal.pone.0282693

Henry, M., & Baudry, S. (2019). Age-related changes in leg proprioception : Implications for postural control. *Journal of Neurophysiology*, *122*(2), 525-538. https://doi.org/10.1152/jn.00067.2019

Heuninckx, S., Wenderoth, N., & Swinnen, S. P. (2008). Systems Neuroplasticity in the Aging Brain : Recruiting Additional Neural Resources for Successful Motor Performance in Elderly Persons. *The Journal of Neuroscience*, *28*(1), 91-99. https://doi.org/10.1523/JNEUROSCI.3300-07.2008

Hoffstaedter, F., Grefkes, C., Roski, C., Caspers, S., Zilles, K., & Eickhoff, S. B. (2015). Age-related decrease of functional connectivity additional to gray matter atrophy in a network for movement initiation. *Brain Structure and Function*, *220*(2), 999-1012. https://doi.org/10.1007/s00429-013-0696-2

Holtzer, R., Mahoney, J. R., Izzetoglu, M., Wang, C., England, S., & Verghese, J. (2015). Online fronto-cortical control of simple and attention-demanding locomotion in humans. *NeuroImage*, *112*, 152-159. https://doi.org/10.1016/j.neuroimage.2015.03.002

Holtzer, R., Verghese, J., Allali, G., Izzetoglu, M., Wang, C., & Mahoney, J. R. (2016). Neurological Gait Abnormalities Moderate the Functional Brain Signature of the Posture First Hypothesis. *Brain Topography*, *29*(2), 334-343. https://doi.org/10.1007/s10548-015-0465-z

Hondzinski, J. M., Soebbing, C. M., French, A. E., & Winges, S. A. (2016). Different damping responses explain vertical endpoint error differences between visual conditions. *Experimental Brain Research*, *234*(6), 1575-1587. https://doi.org/10.1007/s00221-015-4546-8

Hortobagyi, T., Finch, A., Solnik, S., Rider, P., & DeVita, P. (2011). Association Between Muscle Activation and Metabolic Cost of Walking in Young and Old Adults. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, *66A*(5), 541-547. https://doi.org/10.1093/gerona/glr008

Hortobagyi, T., Mizelle, C., Beam, S., & DeVita, P. (2003). Old Adults Perform Activities of Daily Living Near Their Maximal Capabilities. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, *58*(5), M453-M460. https://doi.org/10.1093/gerona/58.5.M453

Huang, H. J., & Ahmed, A. A. (2014). Older adults learn less, but still reduce metabolic cost, during motor adaptation. *Journal of Neurophysiology*, *111*(1), 135-144. https://doi.org/10.1152/jn.00401.2013

Huang, H. J., Kram, R., & Ahmed, A. A. (2012). Reduction of Metabolic Cost during Motor Learning of Arm Reaching Dynamics. *Journal of Neuroscience*, *32*(6), 2182-2190. https://doi.org/10.1523/JNEUROSCI.4003-11.2012

Huber, M., Knottnerus, J. A., Green, L., Horst, H. V. D., Jadad, A. R., Kromhout, D., Leonard, B., Lorig, K., Loureiro, M. I., Meer, J. W. M. V. D., Schnabel, P., Smith, R., Weel, C. V., & Smid, H. (2011). How should we define health? *BMJ*, *343*(jul26 2), d4163-d4163. https://doi.org/10.1136/bmj.d4163

Jeon, W., Hsiao, H.-Y., & Griffin, L. (2021). Effects of different initial foot positions on kinematics, muscle activation patterns, and postural control during a sit-to-stand in younger and older adults. *Journal of Biomechanics*, *117*, 110251. https://doi.org/10.1016/j.jbiomech.2021.110251

John, E. B., Liu, W., & Gregory, R. W. (2009). Biomechanics of Muscular Effort : Age-Related Changes. *Medicine & Science in Sports & Exercise*, *41*(2), 418-425. https://doi.org/10.1249/MSS.0b013e3181884480

Johnson, R. A., & Wichern, D. W. (1988). *Applied multivariate statistical analysis* (2. ed). Prentice Hall.

Jor'dan, A. J., Poole, V. N., Iloputaife, I., Milberg, W., Manor, B., Esterman, M., & Lipsitz, L. A. (2017). Executive Network Activation is Linked to Walking Speed in Older Adults : Functional MRI and TCD Ultrasound Evidence From the MOBILIZE Boston Study. *The Journals of Gerontology: Series A*, *72*(12), 1669-1675. https://doi.org/10.1093/gerona/glx063

Julius, L. M., Brach, J. S., Wert, D. M., & VanSwearingen, J. M. (2012). Perceived Effort of Walking : Relationship With Gait, Physical Function and Activity, Fear of Falling, and Confidence in Walking in Older Adults With Mobility Limitations. *Physical Therapy*, *92*(10), 1268-1277. https://doi.org/10.2522/ptj.20110326

Krakauer, J. W., Ghazanfar, A. A., Gomez-Marin, A., MacIver, M. A., & Poeppel, D. (2017). Neuroscience Needs Behavior : Correcting a Reductionist Bias. *Neuron*, *93*(3), 480-490. https://doi.org/10.1016/j.neuron.2016.12.041

Larivière, S., Xifra-Porxas, A., Kassinopoulos, M., Niso, G., Baillet, S., Mitsis, G. D., & Boudrias, M. (2019). Functional and effective reorganization of the aging brain during unimanual and bimanual hand movements. *Human Brain Mapping*, *40*(10), 3027-3040. https://doi.org/10.1002/hbm.24578

Larsson, L., Degens, H., Li, M., Salviati, L., Lee, Y. I., Thompson, W., Kirkland, J. L., & Sandri, M. (2019). Sarcopenia : Aging-Related Loss of Muscle Mass and Function. *Physiological Reviews*, *99*(1), 427-511. https://doi.org/10.1152/physrev.00061.2017

Le Seac'h, A. B., & McIntyre, J. (2007). Multimodal reference frame for the planning of vertical arms movements. *Neuroscience Letters*, *423*(3), 211-215. https://doi.org/10.1016/j.neulet.2007.07.034

Liu, D., & Todorov, E. (2007). Evidence for the Flexible Sensorimotor Strategies Predicted by Optimal Feedback Control. *The Journal of Neuroscience*, *27*(35), 9354-9368. https://doi.org/10.1523/JNEUROSCI.1110-06.2007

Loibl, M., Beutling, W., Kaza, E., & Lotze, M. (2011). Non-effective increase of fMRI-activation for motor performance in elder individuals. *Behavioural Brain Research*, *223*(2), 280-286. https://doi.org/10.1016/j.bbr.2011.04.040

Manckoundia, P., Mourey, F., Pfitzenmeyer, P., & Papaxanthis, C. (2006). Comparison of motor
strategies in sit-to-stand and back-to-sit motions between healthy and Alzheimer's disease
elderly subjects. *Neuroscience*, *137*(2), 385-392.
https://doi.org/10.1016/j.neuroscience.2005.08.079

Martin, P., Kelly, N., Kahana, B., Kahana, E., Willcox, B. J., Willcox, D. C., & Poon, L. W. (2015).
Defining Successful Aging : A Tangible or Elusive Concept? *The Gerontologist*, *55*(1), 14-25.
https://doi.org/10.1093/geront/gnu044

Marusic, U., Verghese, J., & Mahoney, J. R. (2018). Cognitive-Based Interventions to Improve
Mobility : A Systematic Review and Meta-analysis. *Journal of the American Medical Directors*
Association, *19*(6), 484-491.e3. https://doi.org/10.1016/j.jamda.2018.02.002

Mattay, V. S., Fera, F., Tessitore, A., Hariri, A. R., Das, S., Callicott, J. H., & Weinberger, D. R. (2002).
Neurophysiological correlates of age-related changes in human motor function. *Neurology*,
58(4), 630-635. https://doi.org/10.1212/WNL.58.4.630

Millington, P. J., Myklebust, B. M., & Shambes, G. M. (1992). Biomechanical analysis of the sit-to-
stand motion in elderly persons. *Archives of Physical Medicine and Rehabilitation*, *73*(7),
609-617.

Mombaur, K., Truong, A., & Laumond, J.-P. (2010). From human to humanoid locomotion—An
inverse optimal control approach. *Autonomous Robots*, *28*(3), 369-383.
https://doi.org/10.1007/s10514-009-9170-7

Moran, R. J., Symmonds, M., Dolan, R. J., & Friston, K. J. (2014). The Brain Ages Optimally to Model
Its Environment : Evidence from Sensory Learning over the Adult Lifespan. *PLoS*
Computational Biology, *10*(1), e1003422. https://doi.org/10.1371/journal.pcbi.1003422

Morel, L., Chiang, M. S. R., Higashimori, H., Shoneye, T., Iyer, L. K., Yelick, J., Tai, A., & Yang, Y.
(2017). Molecular and Functional Properties of Regional Astrocytes in the Adult Brain. *The*
Journal of Neuroscience, *37*(36), 8706-8717. https://doi.org/10.1523/JNEUROSCI.3956-
16.2017

Mourey, F., Pozzo, T., Rouhier-Marcer, I., & Didier, J.-P. (1998). A kinematic comparison between
elderly and young subjects standing up from and sitting down in a chair. *Age and Ageing*, *27*(2),
137-146. https://doi.org/10.1093/ageing/27.2.137

Ouwehand, C., De Ridder, D. T. D., & Bensing, J. M. (2007). A review of successful aging models :
Proposing proactive coping as an important additional strategy. *Clinical Psychology Review*,
27(8), 873-884. https://doi.org/10.1016/j.cpr.2006.11.003

Paizis, C., Papaxanthis, C., Berret, B., & Pozzo, T. (2008). Reaching beyond arm length in normal
aging : Adaptation of hand trajectory and dynamic equilibrium. *Behavioral Neuroscience*,
122(6), 1361-1370. https://doi.org/10.1037/a0013280

Papaxanthis, C., Pozzo, T., & McIntyre, J. (2005). Kinematic and dynamic processes for the control of
pointing movements in humans revealed by short-term exposure to microgravity.
Neuroscience, *135*(2), 371-383. https://doi.org/10.1016/j.neuroscience.2005.06.063

Papaxanthis, C., Pozzo, T., Vinter, A., & Grishin, A. (1998). The representation of gravitational force
during drawing movements of the arm. *Experimental Brain Research*, *120*(2), 233-242.
https://doi.org/10.1007/s002210050397

Parthasharathy, M., Mantini, D., & Orban De Xivry, J.-J. (2022). Increased upper-limb sensory
attenuation with age. *Journal of Neurophysiology*, *127*(2), 474-492.
https://doi.org/10.1152/jn.00558.2020

Pereira, T. D., Shaevitz, J. W., & Murthy, M. (2020). Quantifying behavior to understand the brain.
Nature Neuroscience, *23*(12), 1537-1549. https://doi.org/10.1038/s41593-020-00734-z

Poirier, G., Mourey, F., Sirandre, C., Papaxanthis, C., & Gaveau, J. (2023). *Speed-dependent*
optimization of gravity effects for motor control. https://doi.org/10.1101/2023.03.14.532654

Poirier, G., Ohayon, A., Juranville, A., Mourey, F., & Gaveau, J. (2021). Deterioration, Compensation
and Motor Control Processes in Healthy Aging, Mild Cognitive Impairment and Alzheimer's
Disease. *Geriatrics*, *6*(1), 33. https://doi.org/10.3390/geriatrics6010033

Poirier, G., Papaxanthis, C., Lebigre, M., Juranville, A., Mathieu, R., Savoye-Laurens, T.,
Manckoundia, P., Mourey, F., & Gaveau, J. (2024). *Aging decreases the lateralization of*
gravity-related effort minimization during vertical arm movements.
https://doi.org/10.1101/2021.10.26.465988

Poirier, G., Papaxanthis, C., Mourey, F., & Gaveau, J. (2020). Motor Planning of Vertical Arm
Movements in Healthy Older Adults : Does Effort Minimization Persist With Aging? *Frontiers*
in Aging Neuroscience, *12*, 37. https://doi.org/10.3389/fnagi.2020.00037
Poirier, G., Papaxanthis, C., Mourey, F., Lebigre, M., & Gaveau, J. (2022). Muscle effort is best
minimized by the right-dominant arm in the gravity field. *Journal of Neurophysiology*,
jn.00324.2021. https://doi.org/10.1152/jn.00324.2021
Pousson, M., Lepers, R., & Van Hoecke, J. (2001). Changes in isokinetic torque and muscular activity
of elbow flexors muscles with age. *Experimental Gerontology*, *36*(10), 1687-1698.
https://doi.org/10.1016/S0531-5565(01)00143-7
Quinlan, J. I., Maganaris, C. N., Franchi, M. V., Smith, K., Atherton, P. J., Szewczyk, N. J., Greenhaff,
P. L., Phillips, B. E., Blackwell, J. I., Boereboom, C., Williams, J. P., Lund, J., & Narici, M. V.
(2018). Muscle and Tendon Contributions to Reduced Rate of Torque Development in Healthy
Older Males. *The Journals of Gerontology: Series A*, *73*(4), 539-545.
https://doi.org/10.1093/gerona/glx149
Riecker, A., Gröschel, K., Ackermann, H., Steinbrink, C., Witte, O., & Kastrup, A. (2006). Functional
significance of age-related differences in motor activation patterns. *NeuroImage*, *32*(3),
1345-1354. https://doi.org/10.1016/j.neuroimage.2006.05.021
Rudnicka, E., Napierała, P., Podfigurna, A., Męczekalski, B., Smolarczyk, R., & Grymowicz, M.
(2020). The World Health Organization (WHO) approach to healthy ageing. *Maturitas*, *139*,
6-11. https://doi.org/10.1016/j.maturitas.2020.05.018
Saenen, L., Verheyden, G., & Orban De Xivry, J.-J. (2023). The differential effect of age on upper limb
sensory processing, proprioception, and motor function. *Journal of Neurophysiology*, *130*(5),
1183-1193. https://doi.org/10.1152/jn.00364.2022
Saftari, L. N., & Kwon, O.-S. (2018). Ageing vision and falls : A review. *Journal of Physiological*
Anthropology, *37*(1), 11. https://doi.org/10.1186/s40101-018-0170-1
Salat, D. H. (2004). Thinning of the Cerebral Cortex in Aging. *Cerebral Cortex*, *14*(7), 721-730.
https://doi.org/10.1093/cercor/bhh032
Seidler, R. D., Bernard, J. A., Burutolu, T. B., Fling, B. W., Gordon, M. T., Gwin, J. T., Kwak, Y., &
Lipps, D. B. (2010). Motor control and aging : Links to age-related brain structural, functional,
and biochemical effects. *Neuroscience & Biobehavioral Reviews*, *34*(5), 721-733.
https://doi.org/10.1016/j.neubiorev.2009.10.005
Selinger, J. C., O'Connor, S. M., Wong, J. D., & Donelan, J. M. (2015). Humans Can Continuously
Optimize Energetic Cost during Walking. *Current Biology*, *25*(18), 2452-2456.
https://doi.org/10.1016/j.cub.2015.08.016
Shadmehr, R., Huang, H. J., & Ahmed, A. A. (2016). A Representation of Effort in Decision-Making
and Motor Control. *Current Biology*, *26*(14), 1929-1934.
https://doi.org/10.1016/j.cub.2016.05.065
Spedden, M. E., Choi, J. T., Nielsen, J. B., & Geertsen, S. S. (2019). Corticospinal control of normal
and visually guided gait in healthy older and younger adults. *Neurobiology of Aging*, *78*, 29-41.
https://doi.org/10.1016/j.neurobiolaging.2019.02.005
Stapley, P. J., Pozzo, T., Cheron, G., & Grishin, A. (1999). Does the coordination between posture and
movement during human whole-body reaching ensure center of mass stabilization?
Experimental Brain Research, *129*(1), 134-146. https://doi.org/10.1007/s002210050944
Strotz, L. C., Saupe, E. E., Kimmig, J., & Lieberman, B. S. (2018). Metabolic rates, climate and
macroevolution : A case study using Neogene molluscs. *Proceedings of the Royal Society B:*
Biological Sciences, *285*(1885), 20181292. https://doi.org/10.1098/rspb.2018.1292
Summerside, E. M., Courter, R. J., Shadmehr, R., & Ahmed, A. A. (2024). Slowing of Movements in
Healthy Aging as a Rational Economic Response to an Elevated Effort Landscape. *The Journal*
of Neuroscience, *44*(15), e1596232024. https://doi.org/10.1523/JNEUROSCI.1596-23.2024
Tanis, D., Calalo, J. A., Cashaback, J. G. A., & Kurtzer, I. L. (2023). Accuracy and effort costs together
lead to temporal asynchrony of multiple motor commands. *Journal of Neurophysiology*, *129*(1),
1-6. https://doi.org/10.1152/jn.00435.2022
The Lancet. (2009). What is health? The ability to adapt. *The Lancet*, *373*(9666), 781.
https://doi.org/10.1016/S0140-6736(09)60456-6

Thomas, E., Ali, F. B., Tolambiya, A., Chambellant, F., & Gaveau, J. (2023). Too much information is no information : How machine learning and feature selection could help in understanding the motor control of pointing. *Frontiers in Big Data*, *6*, 921355. https://doi.org/10.3389/fdata.2023.921355

Tolambiya, A., Thomas, E., Chiovetto, E., Berret, B., & Pozzo, T. (2011). An Ensemble Analysis of Electromyographic Activity during Whole Body Pointing with the Use of Support Vector Machines. *PLoS ONE*, *6*(7), e20732. https://doi.org/10.1371/journal.pone.0020732

Urai, A. E., Doiron, B., Leifer, A. M., & Churchland, A. K. (2022). Large-scale neural recordings call for new insights to link brain and behavior. *Nature Neuroscience*, *25*(1), 11-19. https://doi.org/10.1038/s41593-021-00980-9

Van Voorhies, W. A., & Ward, S. (1999). Genetic and environmental conditions that increase longevity in *Caenorhabditis elegans* decrease metabolic rate. *Proceedings of the National Academy of Sciences*, *96*(20), 11399-11403. https://doi.org/10.1073/pnas.96.20.11399

VanSwearingen, J. M., & Studenski, S. A. (2014). Aging, Motor Skill, and the Energy Cost of Walking : Implications for the Prevention and Treatment of Mobility Decline in Older Persons. *The Journals of Gerontology Series A: Biological Sciences and Medical Sciences*, *69*(11), 1429-1436. https://doi.org/10.1093/gerona/glu153

Vernazza-Martin, S., Tricon, V., Martin, N., Mesure, S., Azulay, J. P., & Le Pellec-Muller, A. (2008). Effect of aging on the coordination between equilibrium and movement : What changes? *Experimental Brain Research*, *187*(2), 255-265. https://doi.org/10.1007/s00221-008-1301-4

Vu, V. H., Isableu, B., & Berret, B. (2016). On the nature of motor planning variables during arm pointing movement : Compositeness and speed dependence. *Neuroscience*, *328*, 127-146. https://doi.org/10.1016/j.neuroscience.2016.04.027

Ward, N. S. (2006). Compensatory mechanisms in the aging motor system. *Ageing Research Reviews*, *5*(3), 239-254. https://doi.org/10.1016/j.arr.2006.04.003

Ward, N. S., Swayne, O. B. C., & Newton, J. M. (2008). Age-dependent changes in the neural correlates of force modulation : An fMRI study. *Neurobiology of Aging*, *29*(9), 1434-1446. https://doi.org/10.1016/j.neurobiolaging.2007.04.017

White, O., Gaveau, J., Bringoux, L., & Crevecoeur, F. (2020). The gravitational imprint on sensorimotor planning and control. *Journal of Neurophysiology*, *124*(1), 4-19. https://doi.org/10.1152/jn.00381.2019

Wickramarachchi, B., Torabi, M. R., & Perera, B. (2023). Effects of Physical Activity on Physical Fitness and Functional Ability in Older Adults. *Gerontology & Geriatric Medicine*, *9*, 23337214231158476. https://doi.org/10.1177/23337214231158476

Winter, D. A. (2009). *Biomechanics and Motor Control of Human Movement*. John Wiley & Sons, Inc. https://doi.org/10.1002/9780470549148

Wolpe, N., Ingram, J. N., Tsvetanov, K. A., Geerligs, L., Kievit, R. A., Henson, R. N., Wolpert, D. M., Cam-CAN, Tyler, L. K., Brayne, C., Bullmore, E., Calder, A., Cusack, R., Dalgleish, T., Duncan, J., Matthews, F. E., Marslen-Wilson, W., Shafto, M. A., Campbell, K., … Rowe, J. B. (2016). Ageing increases reliance on sensorimotor prediction through structural and functional differences in frontostriatal circuits. *Nature Communications*, *7*(1), 13034. https://doi.org/10.1038/ncomms13034

Yamamoto, S., & Kushiro, K. (2014). Direction-dependent differences in temporal kinematics for vertical prehension movements. *Experimental Brain Research*, *232*(2), 703-711. https://doi.org/10.1007/s00221-013-3783-y

Zalewski, C. (2015). Aging of the Human Vestibular System. *Seminars in Hearing*, *36*(03), 175-196. https://doi.org/10.1055/s-0035-1555120

Zhang, W., & Radhakrishnan, K. (2018). Evidence on selection, optimization, and compensation strategies to optimize aging with multiple chronic conditions : A literature review. *Geriatric Nursing*, *39*(5), 534-542. https://doi.org/10.1016/j.gerinurse.2018.02.013

Zhao, E., Tranovich, M. J., & Wright, V. J. (2014). The role of mobility as a protective factor of cognitive functioning in aging adults : A review. *Sports Health*, *6*(1), 63-69. https://doi.org/10.1177/1941738113477832