**Transinfection of *Wolbachia w*AlbB into *Culex quinquefasciatus* mosquitoes does not**

**alter vector competence for Hawaiian avian malaria (*Plasmodium relictum* GRW4)**

A. Marm Kilpatrick[1,*], Christa M. Seidl[1], Isaiah J. Ipsaro[1], Chris E. Garrison[1], Giulia Fabbri[2], Paul

I. Howell[2], Austin G. McGowan[2], Bradley J. White[2], Sara N. Mitchell[2]

[1]Dept. of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA 95064

[2]Verily Life Sciences, South San Francisco, CA, 94080, USA

*corresponding author: akilpatr@ucsc.edu

**Short Title:** *Wolbachia w*AlbB transinfection and vector competence for avian malaria

**Keywords:** *Wolbachia pipientis*; *Culex quinquefasciatus*, *Plasmodium relictum*; vector

competence; disease control; cytoplasmic incompatibility; incompatible insect technique

**Competing Interests:** GF, PIH, AGM, BJW, and SNM report employment and equity ownership

at Verily Life Sciences, a for-profit company developing new technologies for mosquito control.

**Abstract**

Avian malaria is expanding upslope with warmer temperatures and driving multiple species of

Hawaiian birds towards extinction. Methods to reduce malaria transmission are urgently needed

to prevent further declines. Releasing *Wolbachia*-infected incompatible male mosquitoes

suppress mosquito populations and releasing *Wolbachia*-infected female mosquitoes could

reduce pathogen transmission if the *Wolbachia* strain reduced vector competence. We cleared

*Culex quinquefasciatus* of their natural *Wolbachia pipientis* *w*Pip infection and transinfected

them with *Wolbachia* *w*AlbB isolated from *Aedes albopictus*. We show that *w*AlbB infection was

transmitted transovarially, and demonstrate cytoplasmic incompatibility with wild-type

mosquitoes infected with *w*Pip from Oahu and Maui, Hawaii. We measured vector competence

for avian malaria, *Plasmodium relictum*, lineage GRW4, of seven mosquito lines (two with

*w*AlbB; three with natural *w*Pip infection, and two cleared of *Wolbachia* infection) by allowing

them to feed on canaries infected with recently collected field isolates of Hawaiian *P. relictum*.

We tested 73 groups ($N_{total}$ = 1176) of mosquitoes for *P. relictum* infection in abdomens and

disseminated (thorax) infections 6-14 days after feeding across a range of parasitemias from

0.028% to 2.49%, and a smaller subset of salivary glands. We found no measurable effect of

*Wolbachia* on any endpoint, but strong effects of parasitemia, days post feeding, and mosquito

strain on both abdomen infection prevalence and disseminated infection prevalence. These

results suggest that releasing male *w*AlbB-infected *C. quinquefasciatus* mosquitoes could

suppress *w*Pip-infected mosquito populations, but would have little positive or negative impact

on mosquito vector competence for *P. relictum* if *w*AlbB became established in local mosquito

populations. More broadly, the lack of *Wolbachia* effects on vector competence we observed

highlights the variable impacts of both native and transfected *Wolbachia* infections in

mosquitoes.

**Introduction**

Hawaiian birds are experiencing an extinction crisis. Of the 47 species of Hawaiian

honeycreepers known to science, forty-one are extinct or federally endangered (Paxton *et al.*

2022). One of the key threats to the remaining species is avian malaria, *Plasmodium relictum*

(lineage GRW4), transmitted by *Culex quinquefasciatus* mosquitoes (Beadell *et al.* 2006;

Paxton *et al.* 2022). Rising global temperatures due to climate change have increased the

distribution of mosquito populations and the transmission of malaria in higher elevations on

multiple islands, as predicted two decades ago (Benning *et al.* 2002). This has led to further

population declines in many species, with two additional species now nearly extinct in the wild

(Atkinson *et al.* 2014; Paxton *et al.* 2022). New tools are urgently needed to reduce transmission

of avian malaria.

One recent breakthrough for reducing transmission of mosquito-borne diseases is the

use of *Wolbachia* bacteria either to suppress mosquito populations or to reduce mosquito vector

competence (Flores & O'Neill 2018; Ross *et al.* 2019). *Wolbachia* bacteria are intracellular

parasites that naturally infect many mosquitoes and insect species (Hilgenboecker *et al.* 2008),

and they can be transinfected into new species or populations (Hughes & Rasgon 2014).

Different strains of *Wolbachia* can have different effects on mosquitoes, including reducing adult

lifespan, reduced vector competence for some pathogens, and a reduction in mosquito

populations through cytoplasmic incompatibility (CI) (Iturbe-Ormaetxe *et al.* 2011). CI caused by

*Wolbachia* results in inviable embryos unless both male and female mosquitoes have

compatible *Wolbachia* strains. Control strategies involving the release of *Wolbachia*-infected

incompatible males could be further improved if the *Wolbachia* strain also reduces vector

competence in accidentally released female mosquitoes. Similarly, if a *Wolbachia* strain reduces

vector competence, then population replacement via large scale releases of female mosquitoes

infected with this *Wolbachia* strain could reduce vector competence and transmission, as has

been recently demonstrated for dengue virus and *Aedes aegypti* mosquitoes (Utarini *et al.* 2021;

Velez *et al.* 2023). The EPA recently approved an emergency exemption to release large

numbers of *w*AlbB transinfected male mosquitoes to reduce populations of *C. quinquefasciatus*

in Hawaii (US EPA, 2023), making this a viable strategy to reduce transmission of avian malaria

in Hawaii.

Our goals were threefold: first, to transinfect *Wolbachia w*AlbB into a *C. quinquefasciatus*

line to use for incompatible male releases for population suppression in Hawaii; second, to

confirm complete maternal transmission of *Wolbachia* infection and CI of transinfected males

with wild-type females from Hawaii infected with *Wolbachia w*Pip; and third, to determine if the

transfinfected *Wolbachia w*AlbB strain alters the vector competence of transinfected female *C.*

*quinquefasciatus* mosquitoes for the avian malaria lineage found in Hawaii, *P. relictum* GRW4.

We did not have an *a priori* hypothesis about whether the *w*AlbB strain of *Wolbachia* would

increase or reduce vector competence of *C. quinquefasciatus* for *P. relictum* because effects of

transinfected *Wolbachia* in other mosquito species have been highly variable (Hughes *et al.*

2014; Ross *et al.* 2019) and no studies of the effects of transinfected *Wolbachia* on malaria

competence have been done in *C. quinquefasciatus*. Previous studies of *C. quinquefasciatus*

naturally infected with *Wolbachia w*Pip found higher susceptibility for *P. relictum* lineage SGS1

than mosquitoes cleared of *w*Pip with antibiotics (Zélé *et al.* 2014a). However, there was no

effect of *Wolbachia w*Pip on infection prevalence in the field (Zélé *et al.* 2014b). Finally, a

previous study that created a *w*AlbB transinfected *C. quinquefasciatus* line didn't quantify the

effects on vector competence for any pathogen (Ant *et al.* 2020), and this line has been lost in

an insectary malfunction (Steven Sinkins, personal communication).

**Methods**

***Culex quinquefasciatus* Mosquito Strains**

We studied eight *C. quinquefasciatus* mosquito lines (Table S1) which were

combinations of four mosquito strains (Palmyra Atoll (abbreviated Palmyra or Palm in figures),

Oahu, Maui, Field (Captain Cook, Hawaii)) with one of two *Wolbachia* strains (native *w*Pip or

transinfected *w*AlbB) or cleared of *Wolbachia* infection via antibiotic treatment ("None"). We

refer to lines using the strain and *Wolbachia* type (e.g. Palmyra-*w*Pip, or Oahu-None). We used

subsets of these eight lines for three types of experiments: maternal inheritance of transinfected

*w*AlbB *Wolbachia*, cytoplasmic incompatibility, and vector competence for avian malaria, *P.*

*relictum* GRW4 (Table S1; Supplemental Material, Methods: Mosquito strains).

**Clearing *Wolbachia* *w*Pip and transinfecting *Wolbachia* *w*Alb**

We created two mosquito lines without *Wolbachia*, Palmyra-None and Oahu-None, by clearing

native *Wolbachia* wPip with antibiotics and then used these to create two additional mosquito

lines transinfected with *Wolbachia* wAlbB (Supplementary material: Methods: Transinfection of

*Culex quinquefasciatus* with *Wolbachia* *w*AlbB). We examined maternal transmission of *w*AlbB

in Palmyra-*w*AlbB ("DQB", *D*ebug *q*uinquefasciatus wAlb*B*."), and bi-directional cytoplasmic

incompatibility between Palmyra-*w*AlbB and both Oahu-*w*Pip and Maui-*w*Pip (Supplementary

material, Methods: Maternal transmission determination and Cytoplasmic incompatibility (CI)

testing).

***Plasmodium relictum* isolates**

We collected two isolates of *Plasmodium relictum* (lineage GRW4) from wild birds at two

sites on Hawai'i Island, one from an 'Apapane (*Himatione sanguinea*) from Pu'u Wa'awa'a

Forest Reserve (19.738154°N, 155.875234°W, 1,230 m above sea level) in February 2020 and

another from a Warbling White-eye (*Zosterops japonicus*) from the same Captain Cook, HI site

where Field mosquitoes were collected (Supplemental Material: Methods: *Plasmodium relictum*

isolates).

**Experimental *Plasmodium relictum* infections and mosquito feeding**

We inoculated canaries intramuscularly with 50-200 μL of whole blood (parasitemia: 0.1–

2.85%) containing an avian malaria isolate that had been passaged 1–7 times and was either a

thawed deglycerolized sample, or was fresh blood from another infected canary. Starting on day

5 post-infection afterward we took 5-10 μL of blood by brachial venipuncture and screened thin

blood smears by microscopy (Supplemental Material: Methods: *Plasmodium relictum* isolates)

and by qPCR (Neddermeyer *et al.* 2023; Paxton *et al.* 2023) to detect infection and estimate

parasitemia. Once infection was detected, we allowed approximately 50 mosquitoes from each

of three mosquito lines to feed on each infected bird simultaneously. We differentiated the three

mosquito lines during feedings by spraying each line with a green or red fluorescent marker or

leaving it unmarked (Figure S1; Supplementary Material, Methods: Mosquito marking); the line

receiving each spray color (or none) was randomly selected. We fed most mosquitoes by

restraining an infected canary on top of a container containing the mosquitoes that had holes in

the lid that allowed the bird's legs and feet to be inside the container (Figure S2a). Field

mosquitoes were more hesitant to feed and were placed in a Bugdorm with an unrestrained bird

in a PVC cylinder with a perch and allowed to feed for ~ 8 hrs overnight (Figure S2b).

We collected engorged mosquitoes with an aspirator and transferred them into a cage in

an incubator set to 26 °C. We provided mosquitoes with cottons soaked in a 10% sucrose

solution and held them until dissection. We collected unfed females, knocked them down on ice,

and counted the number of each spray color under a UV light to quantify feeding success for

each group.

We killed engorged mosquitoes 6-14 days after feeding and dissected them by cleanly

separating the thorax from the abdomen at the scutellum with sterile dissection needles. We

placed abdomens in 96-well DNA extraction plates containing Chemagic DNA lysis buffer

(PerkinElmer, Waltham, MA, USA) and placed the head, thorax, and legs of each individual in a

second plate to test for a disseminated infection. When DNA extraction plates were not

available, abdomens and thoraxes (including the head and legs) were placed in separate 1 mL

vials containing 0.5 mL of 70% ethanol. All samples were stored at -20 °C until DNA extraction

and processing by qPCR (Supplemental Information, Methods: *Plasmodium relictum* qPCR and

ddPCR). We also examined *P. relictum* infection in the salivary glands for a subset of 106 Oahu

mosquitoes of all three Wolbachia types (*w*AlbB, *w*Pip, and None) (Supplemental Information,

Methods: Salivary gland infection).

**Statistical analyses**

We analyzed the fraction of mosquitoes testing positive for *P. relictum* by qPCR with

generalized linear models with a binomial distribution and a log link. We analyzed abdomen and

thorax infections separately. We included log(Parasitemia), days post feeding, mosquito strain

(Oahu, Palmyra, or Field), and *Wolbachia* strain (*w*AlbB, *w*Pip, or None) as predictors. We

excluded small batches of mosquitoes with N<4 (a batch is a unique mosquito strain+*Wolbachia*

strain+day post feeding+parasitemia), but results were qualitatively identical if we included all

mosquitoes. We examined two and three-way interactions among the predictors

$\log_{10}$(Parasitemia), days post feeding, and mosquito strain and used AIC to determine which

interactions were best supported by the data. We did not include spray color or *P. relictum*

isolate in the final model because neither had support when added to the best fitting model (see

Table S4; Spray color: $\square^2$ = 1.47, df = 2, P = 0.48; PUWA Isolate coef.: -0.225, SE = 0.18, Z = -

1.23, P =  0.22).

To examine salivary gland infection, we log transformed the ratio of *P. relictum* DNA to

mosquito DNA in the salivary glands and analyzed it with a linear model with *Wolbachia* type as

a predictor. There was heteroscedasticy in the residuals so we performed a robust comparison

using the *coeftest* function in the *lmtest* package with a variance-covariance matrix estimated

using the *sandwich* package. All analyses were performed in R, v.4.3.1.

**Results**

**Transinfected line maternal transmission and Cytoplasmic Incompatibility (CI)**

We cleared *w*Pip *Wolbachia* infection from the Palmyra-*w*Pip strain of *C.*

*quinquefasciatus* with antibiotics and successfully transinfected a single female with the *w*AlbB

strain of *Wolbachia*. We then examined maternal transmission in 991 adults from generations 7

thru 13 of the mosquito line from this female ($N_{ave}$ = 188.8 (range 187-192)/generation,

excluding generation 9 which had 32 mosquitoes). Every sample from every generation tested

positive for *w*AlbB by ddPCR, indicating complete maternal transmission of *Wolbachia* across

generations (mean ratio of wAlbB gene copies to *C. quinquefasciatus* gene copies: 2.95; SE =

0.077).

Bidirectional cytoplasmic incompatibility between mosquitoes infected with *w*AlbB and

*w*Pip was nearly complete in both directions. In crosses between Palmyra-*w*AlbB males ("DQB")

with Oahu-*w*Pip females, only 0.93% of 18,457 eggs hatched (99.07% were inviable), and with

Maui-*w*Pip females, only 1.54% of 13,898 eggs hatched (98.46% were inviable) (Figure 1; Table

S2). Similarly, when Palmyra-*w*AlbB females were mated with Oahu-*w*Pip males only 0.03% of

10,496 eggs hatched (99.97% inviable), and when crossed with Maui-wPip males only 0.05% of

14,706 eggs hatched (99.95% inviable) (Figure 1; Table S3).

**Figure 1. Bi-directional cytoplasmic incompatibility results between Palmyra mosquitoes**

**infected with *Wolbachia wAlbB* (green ovals) and two mosquito strains (Oahu and Maui)**

**infected with native *Wolbachia wPip* (yellow ovals). Three replicates are shown for each**

**of four crosses that include both males and females of each mosquito strain: A1-A3:**

**Palmyra-*w*AlbB males X Oahu-*w*Pip females; B1-B3: Oahu-*w*Pip males X Palmyra-*w*AlbB**

**females; C1-C3: Palmyra-*w*AlbB males X Maui-*w*Pip females; D1-D3: Maui-*w*Pip males X**

**Palmyra-*w*AlbB females. Each of the 12 replicates had $N_{mean}$ = 4796 (range 786-7131); the**

**total number of eggs for each cross is shown above the bars.**

**Vector competence measurements**

We infected fourteen canaries with *P. relictum* GRW4, which had peak parasitemias

averaging 1.22% (range 0.04% - 4.7%; note that we didn't sample birds every day and could

have missed individual peak parasitemias). Peak parasitemia increased with passage number

(Log(peak parasitemia) = -6.81+ 0.465 (SE 0.19) * Passage number; P = 0.032; N= 13; $R^2$ =

35.3%), but was unrelated to the dose injected (# of parasites), or isolate (dose coeff.: -0.39, SE

4.02, N = 13, P = 0.925; isolate coef., PUWA vs CACO: -0.042, SE 0.68, N = 13, P = 0.952, in

separate models with passage number). Over the course of infection, birds had daily

parasitemias of 0.016% to 4.7% and we used parasitemias between 0.028% and 2.49% to

measure vector competence of *C. quinquefasciatus* mosquitoes (Figure S3).

We fed 68 batches (mean N = 57; SD = 17.9; range 20-130) of *C. quinquefasciatus*

mosquitoes on these infected canaries (Figure S4; Table S4; see Supplementary Material:

Feeding success). We tested the 1176 fed mosquitoes from these 68 batches in 73 groups for

abdomen and thorax infection, by qPCR (mean N = 16/group; SD = 7.5; range 4-46). Despite

this enormous number of groups, there was no evidence for differences in disseminated (thorax)

infection prevalence among *Wolbachia* strains (*w*AlbB, native *w*Pip, or None) (compare the

height of points with the same color in each panel of Figure 2 and height of different colored

lines in Figure 3; Table S5).

However, the fraction of mosquitoes with disseminated infections increased sharply with

parasitemia and days since feeding, and the slope of parasitemia increased over time (Figures

2,3, Table S5). Disseminated infection prevalence was slightly higher in the Oahu strain than in

the Palmyra strain (Figures S5; Table S5), and highest in the Field strain (Figure S5; Table S5).

The Results were qualitatively identical when we analyzed each *Wolbachia*-mosquito strain pair

separately (Figure S6; Table S6).

**Figure 2. Fraction of thoraxes infected (and binomial 95% CI) plotted against *Wolbachia***

**type in the mosquitoes (*w*AlbB, *w*Pip, or None).** The color shows the parasitemia (percent of

red blood cells infected) of the bird the mosquitoes fed upon (on a $\log_2$ scale), the panel rows

show the mosquito strain (Oahu or Palmyra), the panel columns show the days post-feeding

when the mosquitoes were dissected (6-14 days), and the size of the points shows the sample

size for each point (range 4-39). The fitted model (Table S5) indicates there are no consistent

differences among *Wolbachia* types (points of the same color in each panel are, on average, at

the same height). Points have been slightly jittered along the x-axis to aid in visualization. Note

that in a small number of experiments one of the three *Wolbachia* groups had insufficient

mosquitoes that successfully fed and is not shown (e.g. bottom middle panel, Palm-None on

day 10).

**Figure 3. Fraction of thoraxes infected (and binomial 95% CI) plotted against the parasitemia (percent of red blood cells infected) of the bird the mosquitoes fed upon (on a log$_2$ scale).** This is the same data as in Figure 2, plotted with a different x-axis. Color shows the *Wolbachia* type in the mosquitoes (*w*AlbB, *w*Pip, or None), the panel rows show the mosquito strain (Oahu or Palmyra), the panel columns show the days post-feeding when the mosquitoes were dissected (6-14 days), and the size of the points shows the sample size (range 4-39). The lines show the fitted model for each of the three *Wolbachia* types, which are on top of each other and difficult to distinguish because there was no measurable differences among *Wolbachia* types (Table S5). Points have been slightly jittered along the x-axis to aid in visualization.

Abdomen infection prevalence was higher than disseminated infection prevalence, but patterns were similar, with there being no detectable effect of *Wolbachia* type, and infection prevalence increased with parasitemia, and differed between the mosquito strains (Figures 4, S7; Table S7). Abdomen infection prevalence increased faster with days since infection for the

Palmyra and Field strains than for the Oahu strain, which resulted in differences between the

mosquito strains changing with days since feeding (prevalence was higher in Oahu mosquitoes

6-10 days after infection but was higher in the Palmyra and Field strains 12-14 days after

infection; Figures S7; Table S7).

**Figure 4. Fraction of abdomens infected (and 95% CI) plotted against *Wolbachia* type in**

**the mosquitoes (*w*AlbB, *w*Pip, or None).** The color shows the parasitemia (percent of red

blood cells infected) of the bird the mosquitoes fed upon (on a $\log_2$ scale), the panel rows show

the mosquito strain (Oahu or Palmyra), the panel columns show the days post-feeding when the

mosquitoes were dissected (6-14 days), and the size of the points shows the sample size

(range 4-39). Points have been slightly jittered along the x-axis to aid in visualization.

We tested salivary glands from 106 Oahu mosquitoes for *P. relictum* DNA by ddPCR

that had fed on a canary with a parasitemia of 0.12% 14 days earlier. Salivary glands from all

106 mosquitoes tested positive for *P. relictum* DNA, and there were no differences in the

amount of *P. relictum* DNA among the three *Wolbachia* types (Figure 5).

**Figure 5. *Wolbachia* strain and salivary gland infection.** *P. relictum* DNA measured by

ddPCR, on a log scale, in salivary glands from 106 Oahu mosquitoes infected *w*AlbB, *w*Pip, or

no *Wolbachia* infection*.* Red points show mean and 95% CI. There was no difference among

*Wolbachia* types (robust F = 1.16, df = 2, P = 0.32).

**Discussion**

We successfully transinfected the *w*AlbB *Wolbachia* strain into *C. quinquefasciatus*

mosquitoes originating from Palmyra Atoll after clearing them of their natural *w*Pip *Wolbachia*

infection. The resultant Palmyra-*w*AlbB line of mosquitoes exhibited complete maternal

transmission of *w*AlbB across generations, and almost 100% bidirectional CI with mosquitoes

from Hawaii (Maui and Oahu) infected with *w*Pip. We then created another *w*AlbB-infected line

by backcrossing the Palmyra-*w*AlbB into an Oahu strain of *C. quinquefasciatus* that we cleared

of natural *w*Pip infection. The creation of these two lines of *w*AlbB-infected *C. quinquefasciatus*

mosquitoes makes population suppression of *C. quinquefasciatus* via widespread release of

*w*AlbB-infected males possible both in Hawaii and elsewhere. The bidirectional CI we observed

between *w*AlbB and *w*Pip mosquitoes would result in any accidentally released *w*AlbB-infected

females having inviable offspring if they mate with native *w*Pip males.

We then examined the effect of *w*AlbB, *w*Pip or no *Wolbachia* on vector competence in

two mosquito strains, Palmyra and Oahu, for *P. relictum* GRW4, the only lineage of avian

malaria in Hawaii (Beadell *et al.* 2006). We found infection with *Wolbachia* strains *w*AlbB and

*w*Pip had no detectable effect on vector competence of *C. quinquefasciatus* mosquitoes for

avian malaria *P. relictum* GRW4. There were no detectable differences among *Wolbachia*

groups in thorax infection or abdomen infection in either the Oahu or Palmyra mosquito strains.

We also found no difference among *Wolbachia* groups in the amount of *P. relictum* DNA in

salivary glands in a subset of mosquitoes from the Oahu strain. This is the first study, to our

knowledge, to assess the impact of both natural and stable transinfections of *Wolbachia* on

malaria vector competence in a natural mosquito-pathogen system. Taken together these

results indicate that the purposeful or accidental introduction of the *w*AlbB strain of *Wolbachia* to

mosquito populations in Hawaii would neither help nor hurt conservation efforts to reduce

transmission of avian malaria (Wild 2023).

Our results differ from two previous studies that found that *Wolbachia* altered

*Plasmodium* prevalence in malaria vectors. In *C. quinquefasciatus*, natural *w*Pip infections

increased vector competence for *P. relictum* lineage SGS1 compared to those cleared of

*Wolbachia* infection (Zélé *et al.* 2014a). In contrast, in *Anopheles stephensi,* *w*AlbB reduced

midgut infection prevalence of *Plasmodium falciparum* (Bian *et al.* 2013). The conflicting results

between these two studies suggests that the impacts of *Wolbachia* on *Plasmodium* infection are

not uniform, which is consistent with the highly variable effects of transinfected *Wolbachia* on

other pathogens in other vectors (Ross *et al.* 2019). We had much larger sample sizes than all

past studies combined, and examined seven lines of mosquitoes, including two inbred mosquito

strains with two strains of *Wolbachia* or no *Wolbachia* and one Field mosquito strain, suggesting

that a lack of an effect of *Wolbachia* on *Plasmodium* vector competence was not due to a lack of

power.

In contrast to the lack of differences among *Wolbachia* lines, we found large differences

among mosquito strains in vector competence, with Field mosquitoes having the highest vector

competence, followed by the inbred Oahu strain and then the inbred Palmyra strain. The

differences in vector competence among mosquito strains indicates that we had ample power to

detect differences in vector competence. They also underline the extensive variability in vector

competence among populations of the same mosquito species, which is common for many

vector-pathogen pairs (Kilpatrick *et al.* 2010; Reisen *et al.* 2008).

We found a relatively steep relationship between parasitemia and both thorax and

abdomen infection, and the relationship became steeper with time since feeding. Twelve days

after feeding the relationship was very steep, with parasitemias up to 0.1% leading to relatively

few disseminated thorax infections (<15.0%), whereas parasitemias only three-fold higher

(0.3%) led to most (73.0%) mosquitoes having disseminated infections. In contrast, eight days

after feeding, a 10-fold range of parasitemias (0.06% to 0.6%) produced a similar range in

thorax infection. The steep relationship between host parasitemia and disseminated infection

prevalence made it more challenging to determine the appropriate day to feed mosquitoes on a

canary and which day to dissect and test mosquitoes to obtain an intermediate level of infection.

In many of our feedings most or almost none of the mosquitoes in all groups had disseminated

infections (Figures 2, 3, S5). This steep relationship, if it is also present in wild mosquitoes,

would lead to highly heterogeneous infectiousness among birds (and bird species), with birds

infecting nearly all or almost none of the mosquitoes that fed on them depending on whether

their parasitemia was above or below a relatively narrow threshold parasitemia (e.g. ~0.1%-

0.3% on day 12 post feeding). Our data from the Field mosquito strain was limited, with only one

day post feeding (day 12) where we had fed mosquitoes on canaries with a range of

parasitemias (four parasitemias, ranging from 0.028% to 1.225%; Figure S7). Analysis of just

this (small) dataset of wild G0 mosquitoes produced a much less steep relationship (Figure S7),

suggesting the steep relationship we observed for the dataset composed of mosquitoes from

two inbred lines (Oahu and Palmyra), may be a result of limited genetic variation.

In summary, the rapid decline of many species of Hawaiian birds over the last decade,

due to a climate-change driven increase in malaria transmission at higher elevations, requires

urgent action to prevent extinction. We created two lines of *w*AlbB-infected *C. quinquefasciatus*

mosquitoes with high CI with Hawaiian *C. quinquefasciatus* infected with wPip, which could be

used to suppress mosquito populations via widespread release of *w*AlbB-infected males. We

examined whether the transinfected *Wolbachia* strain *w*AlbB increased or decreased vector

competence of these two lines for *P. relictum* GRW4. We found no effect of *Wolbachia* infection

on thorax, abdomen, or salivary gland infection, suggesting that replacement of the current

*Wolbachia* in Hawaiian mosquitoes (*w*Pip) with the *w*AlbB strain would have little impact on their

susceptibility to infection and ability to transmit this parasite. Furthermore, vector competence in

one population of mosquitoes collected directly from the field had much higher infection rates

than the two transinfected lines. This suggests that accidental releases of small numbers of

female mosquitoes with *Wolbachia* strain *w*AlbB with the large numbers of male mosquitoes

with *Wolbachia* strain *w*AlbB that is planned for 2024 (Wild 2023) is unlikely to alter the

transmission dynamics of malaria beyond the effect of the males greatly reducing mosquito

populations.

**Acknowledgements**

We thank Nancy McInerney, Nichelle VanTassel, and Rob Fleischer for genetic analysis of our

thawed *P. relictum* isolate to confirm it was lineage GRW4. Funding was provided by NSF

grants DEB-1717498 and DEB-1911853.

**Literature Cited**

Ant, T.H., Herd, C., Louis, F., Failloux, A.B. & Sinkins, S.P. (2020). Wolbachia transinfections in
Culex quinquefasciatus generate cytoplasmic incompatibility. *Insect Mol. Biol.*, 29, 1–8.
Atkinson, C.T., Utzurrum, R.B., Lapointe, D.A., Camp, R.J., Crampton, L.H., Foster, J.T., *et al.*
(2014). Changing climate and the altitudinal range of avian malaria in the Hawaiian
Islands - an ongoing conservation crisis on the island of Kaua'i. *Glob. Change Biol.*, 20,
2426–2436.
Beadell, J.S., Ishtiaq, F., Covas, R., Melo, M., Warren, B.H., Atkinson, C.T., *et al.* (2006). Global
phylogeographic limits of Hawaii's avian malaria. *Proc. R. Soc. B Biol. Sci.*, 273, 2935–
2944.
Benning, T.L., LaPointe, D., Atkinson, C.T. & Vitousek, P.M. (2002). Interactions of climate
change with biological invasions and land use in the Hawaiian Islands: Modeling the fate
of endemic birds using a geographic information system. *Proc. Natl. Acad. Sci. U. S. A.*,
99, 14246–14249.
Bian, G., Joshi, D., Dong, Y., Lu, P., Zhou, G., Pan, X., *et al.* (2013). Wolbachia Invades
Anopheles stephensi Populations and Induces Refractoriness to Plasmodium Infection.
*Science*, 340, 748–751.
Carlson, J.S., Giannitti, F., Valkiūnas, G., Tell, L.A., Snipes, J., Wright, S., *et al.* (2016). A
method to preserve low parasitaemia Plasmodium-infected avian blood for host and
vector infectivity assays. *Malar. J.*, 15, 154.
Crawford, J.E., Clarke, D.W., Criswell, V., Desnoyer, M., Cornel, D., Deegan, B., *et al.* (2020).
Efficient production of male Wolbachia-infected Aedes aegypti mosquitoes enables
large-scale suppression of wild populations. *Nat. Biotechnol.*, 38, 482–492.
Dobson, S.L. & Rattanadechakul, W. (2001). A Novel Technique for Removing Wolbachia
Infections from Aedes albopictus (Diptera: Culicidae). *J. Med. Entomol.*, 38, 844–849.
Dumas, E., Atyame, C.M., Milesi, P., Fonseca, D.M., Shaikevich, E.V., Unal, S., *et al.* (2013).
Population structure of Wolbachia and cytoplasmic introgression in a complex of
mosquito species. *BMC Evol. Biol.*, 13, 181.
Faiman, R., Krajacich, B.J., Graber, L., Dao, A., Yaro, A.S., Yossi, O., *et al.* (2021). A novel
fluorescence and DNA combination for versatile, long-term marking of mosquitoes.
*Methods Ecol. Evol.*, 12, 1008–1016.
Flores, H.A. & O'Neill, S.L. (2018). Controlling vector-borne diseases by releasing modified
mosquitoes. *Nat. Rev. Microbiol.*, 16, 508–518.
Hilgenboecker, K., Hammerstein, P., Schlattmann, P., Telschow, A. & Werren, J.H. (2008). How
many species are infected with Wolbachia? – a statistical analysis of current data. *FEMS
Microbiol. Lett.*, 281, 215–220.
Hughes, G.L. & Rasgon, J.L. (2014). Transinfection: a method to investigate Wolbachia–host
interactions and control arthropod-borne disease. *Insect Mol. Biol.*, 23, 141–151.
Hughes, G.L., Rivero, A. & Rasgon, J.L. (2014). Wolbachia Can Enhance Plasmodium Infection
in Mosquitoes: Implications for Malaria Control? *PLoS Pathog.*, 10, e1004182.
Iturbe-Ormaetxe, I., Walker, T. & O' Neill, S.L. (2011). Wolbachia and the biological control of
mosquito-borne disease. *EMBO Rep.*, 12, 508–518.
Kilpatrick, A.M., Fonseca, D.M., Ebel, G.D., Reddy, M.R. & Kramer, L.D. (2010). Spatial and
temporal variation in vector competence of Culex pipiens and Cx. restuans mosquitoes
for West Nile virus. *Am. J. Trop. Med. Hyg.*, 77, 667–671.
Moll, K., Kaneko, Akira, Scherf, Arthur, Wahlgren, Mats, & EVIMalaR (Eds.). (2013). *Methods in
Malaria Research*. 6th edn. Glasgow, UK: MR4/ATCC; Manassas, VA, USA: 2013.
Neddermeyer, J.H., Parise, K.L., Dittmar, E., Kilpatrick, A.M. & Foster, J.T. (2023). Nowhere to
fly: Avian malaria is ubiquitous from ocean to summit on a Hawaiian island. *Biol.
Conserv.*, 279, 109943.
Paxton, E.H., Laut, M., Enomoto, S. & Bogardus, M. (2022). *Hawaiian forest bird conservation
strategies for minimizing the risk of extinction: biological and biocultural considerations*

(Report No. 103). Hawaii Cooperative Studies Unit Technical Report.

Paxton, K.L., Cassin-Sackett, L., Atkinson, C.T., Videvall, E., Campana, M.G. & Fleischer, R.C.
(2023). Gene expression reveals immune response strategies of naïve Hawaiian
honeycreepers experimentally infected with introduced avian malaria. *J. Hered.*,
esad017.

Reisen, W.K., Barker, C.M., Fang, Y. & Martinez, V.M. (2008). Does variation in Culex (Diptera:
Culicidae) vector competence enable outbreaks of West Nile virus in California? *J. Med.*
*Entomol.*, 45, 1126–1138.

Ross, P.A., Turelli, M. & Hoffmann, A.A. (2019). Evolutionary Ecology of Wolbachia Releases
for Disease Control. *Annu. Rev. Genet.*, 53, 93–116.

US EPA,. (2023). *EPA Approves Emergency Exemption for Wolbachia Mosquitoes to Protect*
*Endangered Birds in Hawaii*.

Utarini, A., Indriani, C., Ahmad, R.A., Tantowijoyo, W., Arguni, E., Ansari, M.R., *et al.* (2021).
Efficacy of Wolbachia-Infected Mosquito Deployments for the Control of Dengue. *N.*
*Engl. J. Med.*, 384, 2177–2186.

Valkiunas, G. (2004). *Avian Malaria Parasites and other Haemosporidia*. CRC Press, Boca
Raton.

Velez, I.D., Tanamas, S.K., Arbelaez, M.P., Kutcher, S.C., Duque, S.L., Uribe, A., *et al.* (2023).
Reduced dengue incidence following city-wide wMel Wolbachia mosquito releases
throughout three Colombian cities: Interrupted time series analysis and a prospective
case-control study. *PLoS Negl. Trop. Dis.*, 17, e0011713.

Videvall, E., Paxton, K.L., Campana, M.G., Cassin-Sackett, L., Atkinson, C.T. & Fleischer, R.C.
(2021). Transcriptome assembly and differential gene expression of the invasive avian
malaria parasite Plasmodium relictum in Hawaiʻi. *Ecol. Evol.*, 11, 4935–4944.

Wild, S. (2023). Millions of Mosquitoes Will Rain Down on Hawaii to Save an Iconic Bird. *Sci.*
*Am.*

Zehtindjiev, P., Ilieva, M., Westerdahl, H., Hansson, B., Valkiūnas, G. & Bensch, S. (2008).
Dynamics of parasitemia of malaria parasites in a naturally and experimentally infected
migratory songbird, the great reed warbler Acrocephalus arundinaceus. *Exp. Parasitol.*,
119, 99–110.

Zélé, F., Nicot, A., Berthomieu, A., Weill, M., Duron, O. & Rivero, A. (2014a). *Wolbachia*
increases susceptibility to *Plasmodium* infection in a natural system. *Proc. R. Soc. B*
*Biol. Sci.*, 281, 20132837.

Zélé, F., Vézilier, J., L'Ambert, G., Nicot, A., Gandon, S., Rivero, A., *et al.* (2014b). Dynamics of
prevalence and diversity of avian malaria infections in wild Culex pipiens mosquitoes:
the effects of Wolbachia, filarial nematodes and insecticide resistance. *Parasit. Vectors*,
7, 437.