Impact of rice *GENERAL REGULATORY FACTOR14h* (*GF14h*) on low-

temperature seed germination and its application to breeding

Yusaku Sugimura[1], Kaori Oikawa[1], Yu Sugihara[2,¤a], Hiroe Utsushi[1], Eiko Kanzaki[1],

Kazue Ito[1], Yumiko Ogasawara[1], Tomoaki Fujioka[3], Hiroki Takagi[1,¤b], Motoki Shimizu[1],

Hiroyuki Shimono[4,5], Ryohei Terauchi[1,2], Akira Abe[1*]

[1] Iwate Biotechnology Research Center, Kitakami, Iwate, Japan

[2] Crop Evolution Laboratory, Kyoto University, Muko, Kyoto, Japan

[3] Iwate Agricultural Research Center, Kitakami, Iwate, Japan

[4] Faculty of Agriculture, Iwate University, Morioka, Iwate, Japan

[5] Agri-Innovation Center, Iwate University, Morioka, Iwate, Japan

[¤a] Current address: The Sainsbury Laboratory, University of East Anglia, Norwich,

United Kingdom

[¤b] Current address: Faculty of Bioresources and Environmental Science, Ishikawa

Prefectural University, 1-308, Suematsu, Nonoichi, Ishikawa, 921-8836, Japan

* Corresponding author

E-mail: a-abe@ibrc.or.jp (AA)

ORCIDs

Yusaku Sugimura, 0000-0001-7271-7703

Yu Sugihara, 0000-0001-6042-1091

Hiroki Takagi, 0000-0002-8576-9892

Motoki Shimizu, 0000-0002-5622-5554

Hiroyuki Shimono, 0000-0002-7328-0483

Ryohei Terauchi, 0000-0002-0095-4651

Akira Abe, 0000-0002-0344-2643

Summary

- Direct seeding is employed to circumvent the labor-intensive process of rice (*Oryza sativa*) transplantation, but this approach requires varieties with vigorous low-temperature germination (LTG) when sown in cold climates.

- To investigate the genetic basis of LTG, we identified the quantitative trait locus (QTL) *qLTG11* from high-LTG rice variety Arroz da Terra using QTL-seq.

- We delineated the candidate region to a 52-kb interval containing *GENERAL REGULATORY FACTOR14h* (*GF14h*) gene, which is expressed during seed germination. The Arroz da Terra *GF14h* allele encodes functional GF14h, whereas Japanese rice variety Hitomebore harbors a 4-bp deletion in the coding region. Knocking out functional *GF14h* in a near-isogenic line (NIL) carrying the Arroz da Terra allele decreased LTG, whereas overexpressing functional *GF14h* in Hitomebore increased LTG, indicating that *GF14h* is the causal gene behind *qLTG11*. Analysis of numerous Japanese rice accessions revealed that the functional *GF14h* allele was lost from popular varieties during modern breeding. We generated a NIL in the Hitomebore background carrying a 172-kb genomic fragment from Arroz da Terra including *GF14h*. The NIL showed superior LTG compared to Hitomebore, with otherwise comparable agronomic traits.

- The functional *GF14h* allele from Arroz da Terra represents a valuable resource for direct seeding in cold regions.

Key words

low-temperature seed germination, *GF14h*, *Oryza sativa*, cultivar, genetics, breeding, near-isogenic line, direct seeding

Introduction

Low-temperature seed germination (LTG) is a pivotal agronomic trait in rice (*Oryza sativa*). As rice originated from tropical and subtropical regions, it is highly susceptible to low-temperature conditions compared to other cereal crops such as wheat (*Triticum aestivum*) and barley (*Hordeum vulgare*) (Okuno, 2004). Nevertheless, rice is produced in temperate and high-altitude regions, where it frequently experiences temperatures

below 20°C. In Japan, rice is abundantly cultivated in relatively cold areas such as Tohoku

and Hokkaido. In recent years, there has been an increasing demand to shift from

conventional transplantation-based rice cultivation to direct seeding to reduce labor and

costs. However, direct seeding raises the risk of exposure to low temperatures during seed

germination (Iwata *et al.*, 2010). Therefore, to expand the use of direct seeding, it is

crucial to breed rice cultivars with enhanced LTG.

LTG is a quantitative trait regulated by complex molecular mechanisms. Linkage

mapping and genome-wide association studies (GWAS) have identified over 30 LTG-

related quantitative trait loci (QTLs) or genomic regions associated with this trait, located

on all 12 rice chromosomes (Miura *et al.*, 2001; Fujino *et al.*, 2004; Han *et al.*, 2006;

Jiang *et al.*, 2006; Fujino *et al.*, 2008; Nguyen *et al.*, 2012; Li *et al.*, 2013; Fujino *et al.*,

2015; Pan *et al.*, 2015; Satoh *et al.*, 2015; Jiang *et al.*, 2017; Shakiba *et al.*, 2017; Wang

et al., 2018; Jiang *et al.*, 2020; Shim *et al.*, 2020; Yang *et al.*, 2020; Pan *et al.*, 2021; Mao

et al., 2022). However, only a few genes involved in LTG have been described, such as

qLTG3-1 (Fujino *et al.*, 2008) and *STRESS-ASSOCIATED PROTEIN16* (*OsSAP16*)

(Wang *et al.*, 2018). The *qLTG3-1* gene, encoding a protein of unknown function, has a

substantial influence on LTG (Fujino *et al.*, 2008). During seed germination, *qLTG3-1*

expression is strongly induced in embryos, which leads to the loosening of the tissues

covering the embryo by promoting vacuolation (Fujino *et al.*, 2008). *OsSAP16* encodes a

stress-associated protein with two AN1-C2H2 zinc finger domains (Wang *et al.*, 2018).

OsSAP16 presumably acts as a regulator of LTG.

14-3-3 proteins are regulatory proteins that are widely conserved in eukaryotes.

These proteins bind to phosphorylated serine and tyrosine residues in their target proteins

that participate in signal transduction and the regulation of gene expression (Morrison,

2009; Madeira *et al.*, 2015), thus altering their enzymatic activity, subcellular localization,

stability, or protein–protein interactions (Hermeking, 2003; Wilson *et al.*, 2016; Camoni

et al., 2018). The rice genome encodes eight 14-3-3 proteins, named GF14a–h for

GENERAL REGULATORY FACTOR14 (Chen *et al.*, 2006). *GF14h* is involved in rice

seed germination under optimal temperature conditions (Sun *et al.*, 2022; Yoshida *et al.*,

2022). In addition, GF14h contributes to phytohormone signaling, including abscisic acid

and gibberellin signaling (Sun *et al.*, 2022; Yoshida *et al.*, 2022). However, it remains

unclear whether *GF14h* promotes seed germination under low-temperature conditions

(Yoshida *et al.*, 2022).

QTL pyramiding has been proposed as a breeding concept (Ashikari & Matsuoka,

2006) for bringing together several QTLs (or genes) related to agronomically important

traits in the genetic background of locally adapted elite cultivars. In practice, it is essential

to generate pre-breeding materials for QTL pyramiding, i.e., near-isogenic lines (NILs)

that harbor one or a few genomic segments introgressed from the donor parent into the

genome of the recipient parent through a combination of continuous backcrossing and

selfing via marker-assisted selection (Zhang *et al.*, 2022). In this study, we determined

that *GF14h* is responsible for an LTG-related QTL in Portuguese rice variety Arroz da

Terra. We generated a NIL in the background of rice cultivar Hitomebore, which is

adapted for growth in northern Japan, by replacing its *GF14h* genomic fragment with that

from Arroz da Terra and tested its LTG performance.

Materials and Methods

Plant materials

Rice was cultivated in a paddy field at Iwate Agricultural Research Center (39°35'N,

141°11'E). A recombinant inbred line (RIL) population of 200 F_7 lines was generated

from a cross between Japanese variety Iwatekko and the high-LTG variety Arroz da Terra

(Fig. S1) (Takagi *et al.*, 2013). To develop NILs, rice cultivar Hitomebore was used as

the recipient parent to generate NILs harboring the target genomic region from Arroz da

Terra (Fig. S2a), thereby establishing *qLTG3-2*-NIL, *qLTG11*-NIL, and NIL-*GF14h^{Arroz}*

(Figs. 1c, S2b, S3a).

Evaluation of germination rate

Seeds for each line were harvested 45 days after heading, air-dried at 30°C for two days,

and stored at 4°C until use. The seeds were air-dried at 50°C for seven days in the dark

to break dormancy. For germination tests, 50 seeds per replicate were incubated in a Petri

dish filled with distilled water in the dark at 15°C (low-temperature conditions) or 25°C

(optimal temperature conditions). The germination rate was calculated as the total number

of germinated seeds at each time point divided by the number of seeds tested. Seeds were

considered to have germinated when the white coleoptile was visible.

To evaluate resistance to pre-harvest sprouting, panicles were harvested from

NIL-*GF14h^Arroz* and Hitomebore 30 days after heading. The panicles were incubated in

dark, wet conditions (by covering them with filter paper moistened with water) at 28°C

for 10 days, and seed germination was scored.

QTL-seq analysis

LTG data for the 200 RILs derived from a cross between Iwatekko and Arroz da Terra

(Fig. S1) were analyzed (Takagi *et al.*, 2013). The top 20 RILs showing high-LTG and

the bottom 20 RILs showing low-LTG phenotypes were selected to assemble the two bulk

samples with contrasting LTG phenotypes. All seedlings with high or low LTG were

pooled, and DNA was extracted from each bulk as previously described (Takagi *et al.*,

2013). The genomic DNA of the two bulks was used to generate DNA-seq libraries and

sequenced on a GAIIx sequencer (Illumina, CA, USA). QTL-seq was performed to

identify QTLs related to LTG (Takagi *et al.*, 2013; Sugihara *et al.*, 2022).

Map-based cloning of *qLTG11*

To narrow down the *qLTG11* region, genotyping was performed using a cross population

of *qLTG11*-NIL (BC$_2$F$_3$) backcrossed to Hitomebore. The germination rate under low-

temperature conditions (15°C) was measured to characterize LTG activity. High-

resolution fine mapping with ten markers (markers B−K) between 23.466 Mb and 23.609

Mb on chromosome 11 identified six informative recombinants in the target region.

Primers used for mapping are listed in Table S1.

***De novo* assembly of the Hitomebore and Arroz da Terra genomes**

To reconstruct the *qLTG11* regions in Hitomebore and Arroz da Terra, *de novo* assembly

was performed for each genome using Nanopore long reads and Illumina short reads

according to a published method (Sugihara *et al.*, 2023). To extract high-molecular-

weight genomic DNA from leaf tissue for Nanopore sequencing, a NucleoBond high-

molecular-weight DNA kit (MACHEREY-NAGEL, Germany) was used. Following

DNA extraction, low-molecular-weight DNA was eliminated using a Short Read

Eliminator Kit XL (Circulomics, MD, USA). Library preparation was then performed

using a Ligation Sequencing Kit (SQK-LSK-109; Oxford Nanopore Technologies [ONT],

United Kingdom) according to the manufacturer's instructions, and sequencing was

performed using MinION (ONT, UK) for Arroz da Terra. For Hitomebore, Nanopore long

reads sequenced by Sugihara et al. (2023) were used. Base-calling of the Nanopore long

reads was performed using Guppy 4.4.2 (ONT, UK). Sequences derived from the lambda

phage genome were removed from the raw reads with NanoLyse v1.1.0 (De Coster *et al.*,

2018). The first 50 bp of each read were then removed, as were reads with an average

read quality score below 7 and reads shorter than 3,000 bases, using NanoFilt v2.7.1 (De

Coster *et al.*, 2018). The clean Nanopore long reads were assembled using NECAT v0.0.1

(Chen *et al.*, 2021), setting the genome size to 380 Mb. To improve the accuracy of

assembly, Racon v1.4.20 (Vaser *et al.*, 2017) was used twice for error correction using

Nanopore reads, and Medaka v1.4.1 (https://github.com/nanoporetech/medaka) was

subsequently used to correct mis-assembly. Two rounds of consensus correction were

then performed using bwa-mem v0.7.17 (Li & Durbin, 2009) and HyPo v1.0.3 (Kundu *et*

al., 2019) with the Illumina short reads. Redundant contigs were removed using purge-

haplotigs v1.1.1 (Roach *et al.*, 2018), resulting in a *de novo* assembly of 374.8 Mb

comprising 82 contigs for Hitomebore and 376.3 Mb consisting of 82 contigs for Arroz

da Terra. The resulting genome sequences have been deposited in Zenodo

(https://doi.org/10.5281/zenodo.10460309).

Plant transformation

To generate *GF14h* knockout mutants, two single guide RNAs (sgRNAs) targeting exon

4 or exon 5 of *GF14h* were designed using the web-based service CRISPRdirect

(crispr.dbcls.jp) (Naito *et al.*, 2015) and cloned individually into the

pZH::OsU6gRNA::MMCas9 vector (Mikami *et al.*, 2015). The resulting vectors were

introduced into Agrobacterium (*Agrobacterium tumefaciens*) strain EHA105 for

transformation into *qLTG11*-NIL plants (Toki *et al.*, 2006). The target sites in the positive

transformants were sequenced by Sanger sequencing to detect mutations. To obtain

overexpression constructs, the full-length coding sequence of functional *GF14h* was

amplified from total RNA extracted from *qLTG11*-NIL and cloned into the plant binary

vector pCAMBIA1300 under the control of the cauliflower mosaic virus (CaMV) 35S

promoter. The overexpression plasmid was introduced into Agrobacterium strain

EHA105 for transformation of rice variety Hitomebore (Toki *et al.*, 2006). All primers

used are listed in Table S1.

Expression analysis

Total RNA was extracted from germinating seeds using an RNA-suisui S kit (Rizo). Total

RNA was treated with RNase-free DNase I (Nippon Gene). The resulting samples were

reverse transcribed into first-strand cDNA using a PrimeScript RT Reagent Kit (Takara

Bio). Quantitative PCR (qPCR) was conducted using a QuantStudio 3 system (Thermo

Fisher Scientific) with Luna Universal qPCR Master Mix (New England Biolabs). The

cycling parameters were 1 min at 95°C, followed by 40 cycles of amplification (95°C for

15 sec and 60°C for 30 sec). The *Actin* gene (Os03g0718100) served as an internal control,

and the Delta CT method was used to calculate the relative expression levels. The primer

sets are listed in Table S1.

RNA-seq

Total RNA was extracted from Hitomebore and *qLTG11*-NIL seeds at 0, 1, 2, and 3 days

after the onset of seed hydration under low (15°C) or optimum (25°C) temperature

conditions using an RNA-suisui S kit (Rizo, Ibaraki, Japan). Sequencing libraries were

prepared using an NEBNext Ultra II Directional RNA Library Prep Kit for Illumina (New

England Biolabs Japan, Tokyo, Japan) following the manufacturer's protocol. The

libraries were sequenced in paired-end mode using an Illumina HiSeq X instrument

(Illumina, CA, USA). The raw reads have been deposited in the DNA Databank of Japan

(BioProject accession No. PRJDB17450). For quality control, reads shorter than 50 bases

and those with an average read quality below 20 were discarded using Trimmomatic v0.36

(Bolger *et al.*, 2014), and poly(A) sequences were trimmed using PRINSEQ++ v1.2

(Cantu *et al.*, 2019). The resulting clean reads were aligned to the *de novo* assembled

Hitomebore and Arroz da Terra genomes with HISAT2 v2.1 (Kim *et al.*, 2019). BAM

files were sorted and indexed with SAMtools v1.10 (Li *et al.*, 2009), and aligned reads

were assembled into transcripts with StringTie (Pertea *et al.*, 2015) by combining bam

files for each variety.

Haplotype network analysis

Sequencing datasets were obtained for 503 rice accessions. Of these, 379 were FASTQ

files downloaded from the DNA Data Bank of Japan Sequence Read Archive (DRA)

(Yabe *et al.*, 2018; Tanaka *et al.*, 2020; Tanaka *et al.*, 2021; Shimono *et al.*, 2023) and

124 were sequenced in this study (Table S2). Details about DNA extraction, whole-

genome sequencing techniques, and construction of the genotype datasets in VCF format

are provided in a previous report (Shimono *et al.*, 2023). This study specifically focused

on the coding region of *GF14h*. Genotype information related to the coding region of

GF14h was extracted from the VCF dataset. In addition, the *k*-mer analysis program

(https://github.com/taitoh1970/kmer) (Itoh *et al.*, 2020) was used with Illumina short

reads to detect the 4-bp deletion with high sensitivity. Genotype information for the

presence of the 4-bp deletion was added to the VCF file, and 81 samples with

heterozygous genotypes were discarded. A haplotype network was then constructed using

the median-joining network algorithm (Bandelt *et al.*, 1999) implemented in Popart v1.7

(Leigh *et al.*, 2015).

Evaluation of agronomic traits and yield performance of NIL-$GF14h^{Arroz}$

The grain yields of Hitomebore and NIL-$GF14h^{Arroz}$ were investigated in experimental

paddy fields in 2023. Field experiments were conducted at the Iwate Agricultural

Research Center (39°35'N, 141°11'E) in Kitakami, Iwate, Japan. A fertilization regime of

N:P_2O_5:K_2O = 6:6:6 g m^{-2} was applied as a basal dressing, and N:K_2O = 2:2 g m^{-2} was

applied as a top dressing. Seeds were sown in a seedling nursery box on 21 April, and

seedlings were transplanted to the paddy field on 18 May. To evaluate agronomic traits,

the seedlings were transplanted at a rate of one plant per hill, with a planting density of

22.2 hills m^{-2}. Culm length, panicle length, panicle number, grain number per plant, and

grain weight per plant were measured at maturity. To evaluate yield performance,

seedlings were transplanted with three plants per hill at a planting density of 16.7 hills

m^{-2}. The 0.9 × 5.0 m experimental plots in the paddy fields were arranged in a randomized

complete block design with three replicates. At maturity, 50 hills were harvested from
each plot to measure brown rice yield. The hulls were removed using a rice huller (Model
25MC, Ohya Tanzo Factory Co., Ltd., Japan), and the hulled grains were screened with a
grain sorter (1.9-mm sieve size). Brown rice yields were adjusted to 15% moisture content
and converted to weight per hectare.

Results

Evaluation of QTLs associated with low-temperature germination using the Portuguese rice variety Arroz da Terra

The Portuguese rice variety Arroz da Terra germinates more vigorously under low-
temperature conditions than the Japanese cultivars Iwatekko and Hitomebore (Fig. 1a).
To identify the genes responsible for this difference, we searched for QTLs involved in
the high LTG of Arroz da Terra. We previously generated a set of 200 RILs at the F_7
generation derived from a cross between Arroz da Terra and Iwatekko (Fig. S1) (Takagi
et al., 2013). We phenotyped all RILs for LTG and selected the 20 RILs with the highest
LTG and the 20 RILs with the lowest LTG. We assembled two pools of seedlings with
low or high LTG and extracted their genomic DNA for whole-genome sequencing on the
Illumina platform (Fig. S1) (Takagi *et al.*, 2013). We mapped the resulting sequencing
reads to the Nipponbare rice reference genome (IRGSP-1.0) and performed QTL-seq
analysis using our new high-performance pipeline (Sugihara *et al.*, 2022). Based on the
ΔSNP index, we identified three QTLs related to LTG on chromosome 3 (*qLTG3-1* and
qLTG3-2) and chromosome 11 (*qLTG11*) (Fig. 1b), which is consistent with the results of
a previous study (Takagi *et al.*, 2013).

The *qLTG3-1* region contained the gene Os03g0103300, which was reported to be
involved in LTG in a study using rice cultivar Italica Livorno, which has high LTG, and
Hayamasari, which has low LTG (Fujino *et al.*, 2008). An examination of its coding
sequences in Arroz da Terra, as well as Iwatekko and Hitomebore, revealed that they were
identical to those found in Italica Livorno and Hayamasari, respectively (Fig. S4). While
Italica Livorno harbored a functional haplotype for this gene, Hayamasari carried a loss-
of-function haplotype due to a 71-bp deletion (Fig. S4) (Fujino *et al.*, 2008). Therefore,
we propose that the causal gene for the QTL *LTG3-1* is Os03g0103300.

To evaluate the contribution of the two other QTLs to LTG, we generated NILs

harboring a segment from the Arroz da Terra genome for each QTL (approximately 5 Mb)

in the Hitomebore background (Figs. 1c, S2a, S3a). We detected no clear effect on LTG,

as *qLTG3-2*-NIL and Hitomebore showed similar seed germination rates at 15ºC (Fig.

S3b). By contrast, the germinability of *qLTG11*-NIL was more vigorous than that of

Hitomebore at 15ºC (Fig. 1d), indicating that *qLTG11* enhances LTG. *qLTG11*-NIL seeds

also germinated more rapidly than Hitomebore seeds under normal conditions (25ºC),

although with a smaller difference between the two genotypes than at low temperature

(Fig. S5). We therefore focused our analysis on *qLTG11*.

Identification of *GF14h* as the candidate gene for *qLTG11*

To delineate the *qLTG11* region, we carried out map-based cloning using a segregating

population derived from a cross between BC_2F_3 line *qLTG11*-NIL and Japanese elite

cultivar Hitomebore (Fig. S2a). For mapping, we conducted germination tests at 15°C.

We narrowed down the genomic region containing the QTL to a 52-kb segment (from

23.512 bp to 23.564 Mb) on chromosome 11 based on the Nipponbare reference genome

(IRGSP-1.0) (Fig. 2a). This interval contains two annotated genes based on the

Nipponbare genome sequence (Fig. 2b). We compared the genomic sequence of

Hitomebore and Arroz da Terra across the candidate region using *de novo* genome

assembly obtained from Nanopore long reads. The cultivars Hitomebore and Nipponbare

had an identical genomic sequence over the entire candidate region (Fig. S6a). By contrast,

the genome sequence from Arroz da Terra was substantially different from that of

Nipponbare, with the equivalent candidate region spanning approximately 94 kb (Figs 2b,

S6b).

As the causal gene behind the variation in LTG is likely expressed in seeds, we

performed transcriptome deep sequencing (RNA-seq) during seed germination in

Hitomebore and *qLTG11*-NIL. Within the candidate region, the gene Os11g0609600,

corresponding to the *14-3-3* gene *GF14h*, was expressed in both Hitomebore and

qLTG11-NIL. The *GF14h* gene structure and haplotypes in Arroz da Terra and

Hitomebore are shown in Fig. 2c. We detected a 4-bp deletion in the *GF14h* coding region

in Hitomebore, causing a frameshift mutation predicted to introduce a premature stop

codon (Figs. 2c, S7). These results suggest that Hitomebore carries a loss-of-function

allele of *GF14h*. To assess the role of *GF14h* in LTG, we examined the expression pattern

of the putative functional *GF14h* (*GF14h^Arroz^*) allele during seed germination at low

temperature (15°C) using *qLTG11*-NIL. RT-qPCR analysis of *GF14h* expression levels

showed that they were comparable in the embryo and endosperm at 1 and 3 days after the

onset of seed imbibition (Fig. 2d). At the beginning of germination, when a white

coleoptile was visible (5 and 7 days after seed imbibition), *GF14h* expression levels rose

in the endosperm, but not in the embryo (Fig. 2d). By nine days of imbibition, when most

seeds had germinated, *GF14h* expression in the endosperm returned to basal levels (Fig.

2d). These results support the notion that *GF14h* plays a role in seed germination at low

temperature.

GF14h plays a vital role in LTG

To investigate the contribution of GF14h to LTG, we knocked out the functional *GF14h*

copy present in *qLTG11*-NIL by clustered regularly interspersed short palindromic repeat

(CRISPR)/CRISPR-associated nuclease 9 (Cas9)-mediated gene editing and evaluated

LTG. Specifically, we introduced two single guide RNA (sgRNA) constructs targeting the

exons of *GF14h* individually into *qLTG11*-NIL by Agrobacterium-mediated

transformation. We chose to knock out *GF14h* in the *qLTG11*-NIL background rather than

Arroz da Terra to evaluate the specific contribution of *GF14h* to LTG without the

influence of *qLTG3-1*, which would be present in the Arroz da Terra background. To

accurately evaluate the phenotypes of the edited plants, we selected heterozygous plants

in the T_0 generation and isolated homozygous mutant lines and their unedited

homozygous siblings in the T_1 generation. We obtained four knockout lines (*gf14h-1*,

gf14h-2, *gf14h-3*, and *gf14h-4*) and their wild-type sibling (WT^Arroz^) (Fig. S8). We

detected significant drops in LTG rates in all four knockout lines compared to WT^Arroz^

(Fig. 3a,b). Furthermore, we generated transgenic lines in the Hitomebore background

overexpressing the functional *GF14h* allele from Arroz da Terra under the control of the

CaMV 35S promoter. In these overexpression lines, *GF14h* expression increased

approximately 1,000-fold compared to the wild-type sibling (WT^Hitomebore^) (Fig. S9).

Importantly, the overexpression lines showed higher LTG than WT^Hitomebore^ when tested

at 15ºC (Fig. 3c,d). Taken together, these data indicate that *GF14h* is a key gene involved

in LTG.

Loss-of-function alleles *GF14h* and *qLTG3-1* increased in frequency

during rice breeding in Japan

We reconstructed the *GF14h* haplotype network using genotype data obtained from

whole-genome resequencing of 492 *O. sativa* accessions from various collections,

including the World Rice Core Collection (Tanaka *et al.*, 2020), the Rice Core Collection

of Japanese Landraces (Tanaka *et al.*, 2021), and a set of Japanese landraces and modern

varieties (Shimono *et al.*, 2023), in addition to 11 wild rice (*O. rufipogon*) accessions

(Zhao *et al.*, 2018) (Table S2). We distinguished 10 haplotypes for the *GF14h* coding

region based on 11 polymorphic sites comprising one frameshift mutation caused by a 4-

bp deletion, four nonsynonymous single nucleotide polymorphisms (SNPs), and six

synonymous SNPs (Table S3). The conversion of the functional allele Hap2 to its

nonfunctional allele Hap1 required only a single step: a 4-bp deletion (Fig. S10).

We analyzed the haplotype frequencies of *GF14h* and *qLTG3-1* in Japanese

landraces and cultivars, which we grouped according to their time of release. More than

half of all Japanese landraces carried functional alleles of both *GF14h* and *LTG3-1* (Fig.

4a). The next most common allele combination was a nonfunctional *GF14h* allele with a

functional *LTG3-1* allele (Fig. 4a). The percentage of lines with loss-of-function alleles

at both *GF14h* and *LTG3-1* has increased since the beginning of crossbreeding in Japan

in the early 20th century, with more than 80% of varieties released after 2001 carrying

loss-of-function alleles for both genes (Fig. 4a,b). Looking at each gene separately in

landraces, only a few lines carried a loss-of-function allele for *LTG3-1*, whereas roughly

half of all lines already harbored a loss-of-function allele for *GF14h* (Fig. 4b). Modern

breeding thus appears to have increased the proportion of loss-of-function alleles for these

two genes, with a substantial increase in *LTG3-1*, reaching almost 90% among lines bred

after 2001 (Fig. 4b).

The Arroz da Terra *GF14h* allele could be valuable for rice breeding

To assess how useful the above findings might be to practical breeding programs, we

developed new breeding materials. QTL pyramiding, a strategy for introducing multiple

QTLs for desired traits into a single genetic background, is a key strategy employed in

current breeding. An essential step in QTL pyramiding is the generation of NILs

containing the desired QTLs. Therefore, we developed a NIL, termed NIL-$GF14h^{Arroz}$,

using the elite cultivar Hitomebore as the genetic background into which we introgressed

a 172-kb region from the Arroz da Terra genome containing $GF14h$ (Fig. S2). This NIL

showed a higher seed germination rate under low-temperature conditions compared to

Hitomebore (Fig. 5a,b). In addition, NIL-$GF14h^{Arroz}$ was slightly more susceptible to pre-

harvest sprouting than Hitomebore (Fig. S11). Importantly, we observed no substantial

differences in five agronomic traits (culm length, panicle length, panicle number, grain

number, and grain weight) between NIL-$GF14h^{Arroz}$ and Hitomebore (Fig. 5c-g). Brown

rice yield was slightly lower in NIL-$GF14h^{Arroz}$ compared to Hitomebore, but a sufficient

yield was guaranteed (Fig. 5h,i). These results suggest that NIL-$GF14h^{Arroz}$ could be a

valuable parental line for breeding via QTL pyramiding.

Discussion

Here, we demonstrated that the functional $GF14h$ allele present in Portuguese rice variety

Arroz da Terra plays a pivotal role in supporting seed germinability under low-

temperature conditions. Although the regulation of seed germination by $GF14h$ was

recently documented, its activity under low-temperature conditions remained unclear

(Sun *et al.*, 2022; Yoshida *et al.*, 2022). While many genomic regions associated with

LTG have been detected through QTL mapping and GWAS, only a few studies have

identified the causal genes (Miura *et al.*, 2001; Fujino *et al.*, 2004; Han *et al.*, 2006; Jiang

et al., 2006; Fujino *et al.*, 2008; Nguyen *et al.*, 2012; Li *et al.*, 2013; Fujino *et al.*, 2015;

Pan *et al.*, 2015; Satoh *et al.*, 2015; Jiang *et al.*, 2017; Shakiba *et al.*, 2017; Wang *et al.*,

2018; Jiang *et al.*, 2020; Shim *et al.*, 2020; Yang *et al.*, 2020; Pan *et al.*, 2021; Mao *et al.*,

2022). Indeed, LTG is a quantitative trait involving the cumulative effects of multiple

genes and their epistatic relationships, making it difficult to assess the specific effect of a

single genomic region on LTG. To eliminate the influence of other chromosomal regions

on LTG, we first generated *qLTG11*-NIL containing only one of three QTLs detected in

the Arroz da Terra background for genetic analysis. The analysis of *qLTG11*-NIL revealed

that *GF14h* participates in LTG. Significantly, the NIL harboring the functional *GF14h*

allele from Arroz da Terra in the Hitomebore background provides valuable pre-breeding

materials for QTL pyramiding. These findings provide a genetic understanding of low-

temperature germinability as well as new resources for rice breeding.

Influence of *GF14h* on low-temperature germination

In this study, we identified *GF14h* as being implicated in LTG. In a previous study, a

genetic complementation assay with a functional *GF14h* allele introduced into the rice

cultivar Nipponbare background increased the germination rate at 30°C, but only to a

limited extent at 15°C (Yoshida *et al.*, 2022). This result is not consistent with our finding

that introducing functional *GF14h* into Hitomebore resulted in a significant improvement

in germination at low temperature. Perhaps this discrepancy is due to differences in the

rice varieties used in the germination assays. Notably, the haplotypes of *qLTG3-1*, a major

QTL behind LTG (Fujino *et al.*, 2008), are different between Nipponbare and

Hitomebore: whereas Nipponbare, which was released in 1961, harbors the functional

allele of *qLTG3-1*, Hitomebore carries a loss-of-function allele with a deletion of 71 bp

(Fig. S4) (Hori *et al.*, 2010). Moreover, the cultivars Koshihikari and Hayamasari, which

carry the same loss-of-function *qLTG3-1* allele as the Hitomebore variety, were reported

to exhibit lower germination rates at low temperatures than Nipponbare (Hori *et al.*, 2010).

LTG tests using chromosome segment substitution lines derived from a cross between

Koshihikari and Nipponbare indicated that *qLTG3-1* contributes to the difference in LTG

between the two varieties (Hori *et al.*, 2010). Based on these observations, it is likely that

Nipponbare has a better LTG ability than Hitomebore, which may have masked the effect

of functional *GF14h* on LTG in the previous study (Yoshida *et al.*, 2022). Our study

confirmed the involvement of *GF14h* in LTG through map-based cloning and analysis of

knockout and overexpression lines. It is also worth mentioning that our experiments were

performed in the *qLTG11*-NIL background, which allowed us to isolate the contribution

of GF14h to LTG without any influence from *qLTG3-1* or other genes in the Arroz da

Terra background. In summary, we provided multiple lines of evidence that *GF14h*

contributes to LTG.

The expression pattern of functional *GF14h* during seed germination was

previously unclear. While *GF14h* has been shown to be expressed in the aleurone layer surrounding the embryo (Yoshida *et al.*, 2022), it is also highly expressed in the endosperm (Sun *et al.*, 2022). In the current study, we showed that *GF14h* was expressed throughout the seeds, but with a transient induction in expression in the endosperm at roughly the time of initiation of seed germination. GF14h was reported to regulate seed germination by interacting with the abscisic acid and gibberellin signaling pathways at optimal temperatures (Sun *et al.*, 2022; Yoshida *et al.*, 2022). We therefore suggest that GF14h controls LTG by interacting with various phytohormone signaling pathways.

Low-temperature germinability in rice was lost due to selection in modern Japanese breeding

In this study, we performed haplotype network analysis of *GF14h* using many Japanese rice varieties. We identified ten distinct haplotypes based on 11 polymorphic sites in the *GF14h* coding region. Of these, Hap4, encompassing the aus, indica, tropical japonica, temperate japonica, and *O. rufipogon* accessions, was defined at the center of the haplotype network. Furthermore, we determined that a 4-bp deletion converted the functional haplotype Hap2, which was derived from Hap4, into the nonfunctional haplotype Hap1. This finding is consistent with the relationship between Hap6 and Hap1 (which we defined as Hap2 and Hap1, respectively) observed by Sun *et al.* (2022). These results suggest that the nonfunctional allele represented by Hap1 was introduced into temperate *japonica* varieties from tropical *japonica* varieties carrying Hap2 and then spread to Japanese cultivars.

We studied the haplotype frequencies of *GF14h* and *qLTG3-1* in various Japanese landraces and cultivars, considering their time of release from breeding programs into the field. More than half of the Japanese landraces analyzed carried both functional *GF14h* and *qLTG3-1* alleles. However, the frequency of varieties carrying loss-of-function alleles for both *GF14h* and *qLTG3-1* has increased since crossbreeding began in the early 20th century. This trend has continued to the present, perhaps as a result of artificial selection to improve resistance to pre-harvest sprouting, with more than 80% of all varieties bred since 2001 carrying these loss-of-function alleles.

Our study provides a historical perspective on allelic shifts in Japanese rice breeding

while highlighting the influence of modern breeding on genetic diversity. Further research

is needed to elucidate the potential effects of higher frequencies of loss-of-function alleles

on the overall phenotypic characteristics and ecological adaptability of Japanese rice

varieties. In addition, as mentioned in previous reports (Sun *et al.*, 2022; Yoshida *et al.*,

2022), we believe that the reintroduction of functional alleles should be considered in

order to develop cultivars suitable for labor-saving cultivation techniques such as direct

seeding.

Application to direct seeding for rice cultivation

Cultivation stability under direct seeding conditions is important for managing rice

production costs and reducing labor. However, since rice is sensitive to low temperatures,

improving seed germination and seedling establishment at low temperatures is a desirable

breeding trait in high-latitude rice production areas such as Japan. In the current study,

we developed a potentially useful NIL (NIL-*GF14h^{Arroz}*) by introducing the functional

GF14h allele into the Hitomebore background. Overexpressing this functional *GF14h*

allele was previously shown to improve anaerobic germination and tolerance to seedling

establishment under anaerobic conditions in laboratory experiments (Sun *et al.*, 2022).

However, whether NIL-*GF14h^{Arroz}* exhibits strong seedling vigor at low temperatures in

rice fields remains to be determined. We previously identified a QTL associated with

seedling vigor, *qPHS3-2* (QTL for plant height of seedling 3-2) (Abe *et al.*, 2012).

qPHS3-2 most likely corresponds to the gibberellin biosynthesis gene *GA20 oxidase1*

(*OsGA20ox1*), a paralog of *Semi Dwarf1* (*sd-1*, corresponding to *OsGA20ox2*) (Abe *et*

al., 2012). Therefore, the pyramiding of *qPHS3-2* in NIL-*GF14h^{Arroz}* by marker-assisted

selection represents a promising approach for further improving seedling vigor in NIL-

GF14h^{Arroz}.

In recent years, "early-winter direct seeding" has been experimentally tested as a

new system of direct seeding for rice production in Japan (Shimono, 2020). In this system,

seeds are directly sown in the early winter of the previous year instead of the spring. The

sown seeds thus overwinter in snow-covered soil and germinate the following spring. The

major advantage of this approach is that it can significantly decrease the amount of labor

required by farmers during the busy spring season. However, it is challenging to

overwinter the seeds of modern rice varieties in the soil and achieve good seedling establishment (Shimono *et al.*, 2012; Oikawa *et al.*, 2019; Shimono, 2020). We expect that reintroducing beneficial alleles like *GF14h^{Arroz}* that were lost during modern breeding into future rice varieties will enable the implementation of new cultivation practices and increase productivity.

Acknowledgments

We thank the National Agriculture and Food Research Organization (NARO) gene bank, Japan, for providing rice seeds. This work was performed using the National Institute of Genetics (NIG) supercomputer at the Research Organization of Information and Systems (ROIS) National Institute of Genetics and the Academic Center for Computing and Media Studies (ACCMS) supercomputer at Kyoto University.

Competing interests

None declared.

Author contributions

AA, HT, and RT designed the research. YS, AA, KO, YS, HU, EK, KI, YO, TF, HT, MS, and HS performed experiments and analyzed the data. YS wrote the original manuscript. AA and RT edited the manuscript. All authors contributed to the manuscript and approved the submitted version.

References

Abe A, Takagi H, Fujibe T, Aya K, Kojima M, Sakakibara H, Uemura A, Matsuoka M, Terauchi R. 2012. OsGA20ox1, a candidate gene for a major QTL controlling seedling vigor in rice. *Theor Appl Genet* **125**(4): 647-657.

Ashikari M, Matsuoka M. 2006. Identification, isolation and pyramiding of quantitative trait loci for rice breeding. *Trends Plant Sci* **11**(7): 344-350.

Bandelt HJ, Forster P, Rohl A. 1999. Median-joining networks for inferring

intraspecific phylogenies. *Mol Biol Evol* **16**(1): 37-48.

Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**(15): 2114-2120.

Cabanettes F, Klopp C. 2018. D-GENIES: dot plot large genomes in an interactive,

efficient and simple way. *PeerJ* **6**: e4958.

Camoni L, Visconti S, Aducci P, Marra M. 2018. 14-3-3 Proteins in Plant Hormone

Signaling: Doing Several Things at Once. *Front Plant Sci* **9**: 297.

Cantu VA, Sadural J, Edwards R. 2019. PRINSEQ++, a multi-threaded tool for fast

and efficient quality control and preprocessing of sequencing datasets. *PeerJ*

Preprints **7**: e27553v27551.

Chen F, Li Q, Sun L, He Z. 2006. The rice 14-3-3 gene family and its involvement in

responses to biotic and abiotic stress. *DNA Res* **13**(2): 53-63.

Chen Y, Nie F, Xie SQ, Zheng YF, Dai Q, Bray T, Wang YX, Xing JF, Huang ZJ,

Wang DP, et al. 2021. Efficient assembly of nanopore reads via highly accurate

and intact error correction. *Nat Commun* **12**(1): 60.

De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. 2018. NanoPack:

visualizing and processing long-read sequencing data. *Bioinformatics* **34**(15):

2666-2669.

Fujino K, Obara M, Shimizu T, Koyanagi KO, Ikegaya T. 2015. Genome-wide

association mapping focusing on a rice population derived from rice breeding

programs in a region. *Breed Sci* **65**(5): 403-410.

Fujino K, Sekiguchi H, Matsuda Y, Sugimoto K, Ono K, Yano M. 2008. Molecular

identification of a major quantitative trait locus, qLTG3-1, controlling low-

temperature germinability in rice. *Proc Natl Acad Sci U S A* **105**(34): 12623-

12628.

Fujino K, Sekiguchi H, Sato T, Kiuchi H, Nonoue Y, Takeuchi Y, Ando T, Lin SY,

Yano M. 2004. Mapping of quantitative trait loci controlling low-temperature

germinability in rice (Oryza sativa L.). *Theor Appl Genet* **108**(5): 794-799.

Han LZ, Zhang YY, Qiao YL, Cao GL, Zhang SY, Kim JH, Koh HJ. 2006. Genetic

and QTL analysis for low-temperature vigor of germination in rice. *Yi Chuan Xue*

Bao **33**(11): 998-1006.

Hermeking H. 2003. The 14-3-3 cancer connection. *Nat Rev Cancer* **3**(12): 931-943.

Hori K, Sugimoto K, Nonoue Y, Ono N, Matsubara K, Yamanouchi U, Abe A,

Takeuchi Y, Yano M. 2010. Detection of quantitative trait loci controlling pre-

harvest sprouting resistance by using backcrossed populations of japonica rice

cultivars. *Theor Appl Genet* **120**(8): 1547-1557.

Itoh T, Onuki R, Tsuda M, Oshima M, Endo M, Sakai H, Tanaka T, Ohsawa R, Tabei

Y. 2020. Foreign DNA detection by high-throughput sequencing to regulate

genome-edited agricultural products. *Sci Rep* **10**(1): 4914.

Iwata N, Shinada H, Kiuchi H, Sato T, Fujino K. 2010. Mapping of QTLs controlling

seedling establishment using a direct seeding method in rice. *Breeding Science*

60(4): 353-360.

Jiang L, Liu S, Hou M, Tang J, Chen L, Zhai H, Wan J. 2006. Analysis of QTLs for

seed low temperature germinability and anoxia germinability in rice (Oryza sativa

L.). *Field Crops Research* **98**(1): 68-75.

Jiang N, Shi S, Shi H, Khanzada H, Wassan GM, Zhu C, Peng X, Yu Q, Chen X, He

X, et al. 2017. Mapping QTL for Seed Germinability under Low Temperature

Using a New High-Density Genetic Map of Rice. *Front Plant Sci* **8**: 1223.

Jiang S, Yang C, Xu Q, Wang L, Yang X, Song X, Wang J, Zhang X, Li B, Li H, et

al. 2020. Genetic Dissection of Germinability under Low Temperature by

Building a Resequencing Linkage Map in japonica Rice. *Int J Mol Sci* **21**(4).

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol*

37(8): 907-915.

Kundu R, Casey J, Sung W-K. 2019. HyPo: Super Fast & Accurate Polisher for Long

Read Genome Assemblies. *bioRxiv*: 2019.2012.2019.882506.

Leigh JW, Bryant D, Nakagawa S. 2015. popart: full-feature software for haplotype

network construction. *Methods in Ecology and Evolution* **6**(9): 1110-1116.

Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics* **25**(14): 1754-1760.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G,

Durbin R, Genome Project Data Processing S. 2009. The Sequence

Alignment/Map format and SAMtools. *Bioinformatics* **25**(16): 2078-2079.

Li L, Liu X, Xie K, Wang Y, Liu F, Lin Q, Wang W, Yang C, Lu B, Liu S, et al. 2013.

qLTG-9, a stable quantitative trait locus for low-temperature germination in rice

(Oryza sativa L.). *Theor Appl Genet* **126**(9): 2313-2322.

Madeira F, Tinti M, Murugesan G, Berrett E, Stafford M, Toth R, Cole C,

MacKintosh C, Barton GJ. 2015. 14-3-3-Pred: improved methods to predict 14-

3-3-binding phosphopeptides. *Bioinformatics* **31**(14): 2276-2283.

Mao F, Wu D, Lu F, Yi X, Gu Y, Liu B, Liu F, Tang T, Shi J, Zhao X, et al. 2022. QTL

mapping and candidate gene analysis of low temperature germination in rice

(Oryza sativa L.) using a genome wide association study. *PeerJ* **10**: e13407.

Mikami M, Toki S, Endo M. 2015. Comparison of CRISPR/Cas9 expression constructs

for efficient targeted mutagenesis in rice. *Plant Mol Biol* **88**(6): 561-572.

Miura K, Lin SY, Yano M, Nagamine T. 2001. Mapping Quantitative Trait Loci

Controlling Low Temperature Germinability in Rice (Oryza sativa L.). *Breeding*

Science **51**(4): 293-299.

Morrison DK. 2009. The 14-3-3 proteins: integrators of diverse signaling cues that

impact cell fate and cancer development. *Trends Cell Biol* **19**(1): 16-23.

Naito Y, Hino K, Bono H, Ui-Tei K. 2015. CRISPRdirect: software for designing

CRISPR/Cas guide RNA with reduced off-target sites. *Bioinformatics* **31**(7):

1120-1123.

Nguyen HN, Park I-K, Yeo S-M, Yun Y-T, Ahn S-N. 2012. Mapping quantitative trait

loci controlling low-temperature germinability in rice. *Korean Journal of*

Agricultural Science **39**(4): 477-482.

Oikawa S, Nishi M, Yui S, Kashiwagi J, Nakashima T, Ichikawa S, Kimura T,

Ohdaira Y, Nagasuga K, Kuroda E, et al. 2019. Improvement of Seedling

Establishment by Seed Coating with Iron in Early-winter Direct-sowing Rice.

Japanese Journal of Crop Science **88**(4): 259-267.

Okuno K. 2004. Molecular Mechanisms of Cold Tolerance in Rice and Wheat. *Thermal*

Medicine(Japanese Journal of Hyperthermic Oncology) **20**(2): 51-60.

Pan Y, Zhang H, Zhang D, Li J, Xiong H, Yu J, Li J, Rashid MA, Li G, Ma X, et al.

2015. Genetic analysis of cold tolerance at the germination and booting stages in

rice by association mapping. *PLoS One* **10**(3): e0120590.

Pan Z, Tan B, Cao G, Zheng R, Liu M, Zeng R, Wang S, Zhu H, Ye H, Zhao G, et al.

2021. Integrative QTL Identification, Fine Mapping and Candidate Gene Analysis

of a Major Locus qLTG3a for Seed Low-Temperature Germinability in Rice. *Rice*

(N Y) **14**(1): 103.

Pertea M, Pertea GM, Antonescu CM, Chang TC, Mendell JT, Salzberg SL. 2015.

StringTie enables improved reconstruction of a transcriptome from RNA-seq

reads. *Nat Biotechnol* **33**(3): 290-295.

Roach MJ, Schmidt SA, Borneman AR. 2018. Purge Haplotigs: allelic contig

reassignment for third-gen diploid genome assemblies. *BMC Bioinformatics*

19(1): 460.

Satoh T, Tezuka K, Kawamoto T, Matsumoto S, Satoh-Nagasawa N, Ueda K,

Sakurai K, Watanabe A, Takahashi H, Akagi H. 2015. Identification of QTLs

controlling low-temperature germination of the East European rice (Oryza sativa

L.) variety Maratteli. *Euphytica* **207**(2): 245-254.

Shakiba E, Edwards JD, Jodari F, Duke SE, Baldo AM, Korniliev P, McCouch SR,

Eizenga GC. 2017. Genetic architecture of cold tolerance in rice (Oryza sativa)

determined through high resolution genome-wide analysis. *PLoS One* **12**(3):

e0172133.

Shim KC, Kim SH, Lee HS, Adeva C, Jeon YA, Luong NH, Kim WJ, Akhtamov M,

Park YJ, Ahn SN. 2020. Characterization of a New qLTG3-1 Allele for Low-

temperature Germinability in Rice from the Wild Species Oryza rufipogon. *Rice*

(N Y) **13**(1): 10.

Shimono H. 2020. New technique : Early- winter direct- sowing cultivation in rice under

cool climate. *Regulation of Plant Growth & Development* **55**(1): 63-66.

Shimono H, Abe A, Kim CH, Sato C, Iwata H. 2023. Upcycling rice yield trial data

using a weather-driven crop growth model. *Commun Biol* **6**(1): 764.

Shimono H, Tamai M, Hamasaki T, Sagawa R, Ohtani R. 2012. Effects of Autumn

Direct-seeding on Rice Growth and Yield under Cool Climates. *Japanese Journal*

of Crop Science **81**(1): 93-98.

Sugihara Y, Abe Y, Takagi H, Abe A, Shimizu M, Ito K, Kanzaki E, Oikawa K,

Kourelis J, Langner T, et al. 2023. Disentangling the complex gene interaction

networks between rice and the blast fungus identifies a new pathogen effector.

PLoS Biol **21**(1): e3001945.

Sugihara Y, Young L, Yaegashi H, Natsume S, Shea DJ, Takagi H, Booker H, Innan

H, Terauchi R, Abe A. 2022. High-performance pipeline for MutMap and QTL-

seq. *PeerJ* **10**: e13170.

Sun J, Zhang G, Cui Z, Kong X, Yu X, Gui R, Han Y, Li Z, Lang H, Hua Y, et al.

2022. Regain flood adaptation in rice through a 14-3-3 protein OsGF14h. *Nat*

Commun **13**(1): 5664.

Takagi H, Abe A, Yoshida K, Kosugi S, Natsume S, Mitsuoka C, Uemura A, Utsushi

H, Tamiru M, Takuno S, et al. 2013. QTL-seq: rapid mapping of quantitative

trait loci in rice by whole genome resequencing of DNA from two bulked

populations. *Plant J* **74**(1): 174-183.

Tanaka N, Shenton M, Kawahara Y, Kumagai M, Sakai H, Kanamori H, Yonemaru

J, Fukuoka S, Sugimoto K, Ishimoto M, et al. 2020. Whole-Genome

Sequencing of the NARO World Rice Core Collection (WRC) as the Basis for

Diversity and Association Studies. *Plant Cell Physiol* **61**(5): 922-932.

Tanaka N, Shenton M, Kawahara Y, Kumagai M, Sakai H, Kanamori H, Yonemaru

JI, Fukuoka S, Sugimoto K, Ishimoto M, et al. 2021. Investigation of the

Genetic Diversity of a Rice Core Collection of Japanese Landraces using Whole-

Genome Sequencing. *Plant Cell Physiol* **61**(12): 2087-2096.

Toki S, Hara N, Ono K, Onodera H, Tagiri A, Oka S, Tanaka H. 2006. Early infection

of scutellum tissue with Agrobacterium allows high-speed transformation of rice.

Plant J **47**(6): 969-976.

Vaser R, Sovic I, Nagarajan N, Sikic M. 2017. Fast and accurate de novo genome

assembly from long uncorrected reads. *Genome Res* **27**(5): 737-746.

Wang X, Zou B, Shao Q, Cui Y, Lu S, Zhang Y, Huang Q, Huang J, Hua J. 2018.

Natural variation reveals that OsSAP16 controls low-temperature germination in

rice. *J Exp Bot* **69**(3): 413-421.

Wilson RS, Swatek KN, Thelen JJ. 2016. Regulation of the Regulators: Post-

Translational Modifications, Subcellular, and Spatiotemporal Distribution of

Plant 14-3-3 Proteins. *Front Plant Sci* **7**: 611.

Yabe S, Yoshida H, Kajiya-Kanegae H, Yamasaki M, Iwata H, Ebana K, Hayashi T,

Nakagawa H. 2018. Description of grain weight distribution leading to genomic

selection for grain-filling characteristics in rice. *PLoS One* **13**(11): e0207627.

Yang T, Zhou L, Zhao J, Dong J, Liu Q, Fu H, Mao X, Yang W, Ma Y, Chen L, et al.

2020. The Candidate Genes Underlying a Stably Expressed QTL for Low

Temperature Germinability in Rice (Oryza sativa L.). *Rice (N Y)* **13**(1): 74.

Yoshida H, Hirano K, Yano K, Wang F, Mori M, Kawamura M, Koketsu E, Hattori

M, Ordonio RL, Huang P, et al. 2022. Genome-wide association study identifies

a gene responsible for temperature-dependent rice germination. *Nat Commun*

13(1): 5665.

Zhang B, Ma L, Wu B, Xing Y, Qiu X. 2022. Introgression Lines: Valuable Resources

for Functional Genomics Research and Breeding in Rice (Oryza sativa L.). *Front*

Plant Sci **13**: 863789.

Zhao Q, Feng Q, Lu H, Li Y, Wang A, Tian Q, Zhan Q, Lu Y, Zhang L, Huang T, et

al. 2018. Pan-genome analysis highlights the extent of genomic variation in

cultivated and wild rice. *Nat Genet* **50**(2): 278-284.

Fig 1. Effects of quantitative trait loci (QTLs) on low-temperature seed germinability.

(a) Representative photographs showing the germination of seeds from the Iwatekko, Hitomebore, and Arroz da Terra varieties 10 days after the onset of seed imbibition at $15°$ C. Scale bar, 1 cm. **(b)** Map positions of QTLs for low-temperature germination, as determined by QTL-seq. The Δ (SNP-index) values (red lines) were plotted for chromosomes 3 and 11, with statistical confidence intervals under the null hypothesis of no QTL (green, $P < 0.05$; orange, $P < 0.01$). **(c)** Diagram showing the genotype of *qLTG11*-NIL. *qLTG11*-NIL harbors the Arroz da Terra allele at *qLTG11* on chromosome 11. Light blue indicates genomic fragments from Hitomebore; red indicates genomic fragments from Arroz da Terra; dark blue indicates heterozygous regions. **(d)** Germination time courses of Hitomebore, *qLTG11*-NIL, and Arroz da Terra at $15°$ C. Values are means $\pm$ standard deviation (SD) from biologically independent samples (Hitomebore and NIL, $n = 10$; Arroz da Terra, $n = 5$).

Fig 2. Positional cloning of *qLTG11*.
(a) Fine mapping of *qLTG11* to a 52-kb region between markers E and J. The chromosomal positions are based on the Nipponbare reference genome (Os-Nipponbare-Reference-IRGSP-1.0). Germination percentage was determined at 9 days of incubation at 15° C. Red and blue rectangles indicate chromosomal segments homozygous for Arroz da Terra or Hitomebore, respectively. Different lowercase letters indicate significant differences (n = 3 biologically independent samples, P < 0.001, Tukey's HSD test). **(b)** Genomic structure of the candidate genomic region in Arroz da Terra and Hitomebore. Os11g0609600 (shown in red), encoding GF14h, is expressed in germinating seeds. **(c)** Diagram of the *GF14h* gene structure and sequence polymorphisms between Arroz da Terra and Hitomebore. The chromosomal positions are based on the Nipponbare reference genome. The coding region of *GF14h* in Hitomebore is identical to that in Nipponbare. The 4-bp deletion in Hitomebore causes a frameshift and the introduction of a premature stop codon. **(d)** Relative *GF14h* expression levels in germinating seeds of *qLTG11*-NIL. In the boxplots, the box edges represent the upper and lower quantiles, the horizontal line in the middle of the box represents the median value, whiskers represent the lowest quantile to the top quantile, and the black squares show the mean. Five biological replicates were measured independently. Different lowercase letters indicate significant differences based on Tukey's HSD test (P < 0.05). *OsActin1* (Os03g0718100) was used for normalization.

Fig 3. Effect of *GF14h* mutation and overexpression on low-temperature germination.
(a) Representative photographs showing seed germination in wild-type harboring Arroz-type *GF14h* (WT^Arroz) and CRISPR/Cas9 knockout lines (*gf14-1*) at 8 days after the onset of seed imbibition. Scale bar, 1 cm. **(b)** Seed germination rate of WT^Arroz and its CRISPR/Cas9 knockout lines at 7 days of seed imbibition at 15° C. The two target constructs (Fig. S8) were introduced into the *qLTG11*-NIL line. Data are means $\pm$ standard error (SE, *n* = 3). Different lowercase letters indicate significant differences based on Tukey's HSD test (*P* < 0.01). **(c)** Representative photographs showing seed germination of wild-type (WT^Hitomebore) and *OsGF14h^Arroz* overexpression lines (*OsGF14h^Arroz*-Ox #2) at 7 days of seed imbibition at 15ºC. Scale bar, 1 cm. **(d)** Seed germination rate of WT^Hitomebore and *GF14h^Arroz* overexpression lines in the Hitomebore background at 7 days of seed imbibition at 15° C. Data are means $\pm$ SE (*n* = 3). Different lowercase letters indicate significant differences based on Tukey's HSD test (*P* < 0.05).

Fig 4. Gradual selection of loss-of-function alleles in *GF14h* and *qLTG3-1* during rice breeding in Japan.

A total of 350 Japanese varieties were examined, including the World Rice Core Collection (Tanaka *et al*., 2020), the Rice Core Collection of Japanese Landraces (Tanaka *et al*., 2021), and the collection of Japanese core cultivars (Shimono *et al*., 2023). The allele type at each gene was determined to be functional or nonfunctional by the k-mer method using Illumina short reads for each variety. **(a)** Proportion of allele type combinations at *GF14h* and *qLTG3-1* sorted by breeding year. **(b)** Proportion of nonfunctional allele types at *GF14h* and *qLTG3-1* sorted by breeding year.

Fig 5. Phenotypic analysis of a near-isogenic line homozygous for Arroz-type *GF14h* in the Hitomebore genetic background (NIL-*GF14h^Arroz^*).

(a) Gross morphology of Hitomebore and NIL-*GF14h^Arroz^* at 87 days after transplanting (top row), and seed germination at 9 days after the onset of seed imbibition at 15° C (bottom row). Scale bars, 20 cm (top), 1 cm (bottom). **(b)** Low-temperature germination ability of NIL-*GF14h^Arroz^* at 8 days after the onset of seed imbibition at 15° C. Values are means ± SE of biologically independent samples (*n* = 3). Asterisks indicate significant differences, as determined by two-tailed *t*-test. **(c–g)** Agronomic traits in Hitomebore and NIL-*GF14h^Arroz^*: culm length (**c**), panicle length (**d**), panicle number (**e**), grain number per plant (**f**), and grain weight per plant (**g**). Values are means ± SE of biologically independent plants (*n* = 40). **(h)** Side view of Hitomebore and NIL-*GF14h^Arroz^* growing in the field under typical rice-growing conditions at the maturity stage. **(i)** Brown rice yield per unit area. Values are means ± SE of biologically independent plots (*n* = 3). *P*-values calculated by *t*-tests are listed throughout the figure.