Cortical beta oscillations map to shared brain networks modulated by dopamine

Authors:

Meera Chikermane[1], Liz Weerdmeester[1], Nanditha Rajamani[1], Richard M. Köhler[1], Timon Merk[1], Jonathan Vanhoecke[1], Andreas Horn[1,3,4], Wolf-Julian Neumann*[1,2]

1 Movement Disorders and Neuromodulation Unit, Department of Neurology, Charité – Universitätsmedizin Berlin, Charitéplatz 1, 10117 Berlin, Germany

2 Einstein Center for Neurosciences Berlin, Humboldt Universitat, Berlin, Germany

3 Center for Brain Circuit Therapeutics, Department of Neurology, Brigham and Women's Hospital, MA, USA

4 Departments of Neurology and Neurosurgery, Massachusetts General Hospital, MA, USA

* Corresponding author

 mailto:julian.neumann@charite.de

Abstract

Brain rhythms can facilitate neural communication for the maintenance of brain function. Beta rhythms (13–35 Hz) have been proposed to serve multiple domains of human ability, including motor control, cognition, memory and emotion, but the overarching organisational principles remain unknown. To uncover the circuit architecture of beta oscillations, we leverage normative brain data, analysing over 30 hours of invasive brain signals from 1772 cortical areas in epilepsy patients, to demonstrate that beta is the most distributed cortical brain rhythm. Next, we identify a shared brain network from beta dominant areas with deeper brain structures, like the basal ganglia, by mapping parametrised oscillatory peaks to whole-brain functional and structural MRI connectomes. Finally, we show that these networks share significant overlap with dopamine uptake as indicated by positron emission tomography. Our study suggests that beta oscillations emerge in cortico-subcortical brain networks that are modulated by dopamine. It provides the foundation for a unifying circuit-based conceptualisation of the functional role of beta activity beyond the motor domain and may inspire an extended investigation of beta activity as a feedback signal for closed-loop neurotherapies for dopaminergic disorders.

Keywords: Dopamine, Beta oscillations, Basal ganglia, Movement disorders, Neural circuits, Invasive electrophysiology, MRI connectomics, Cortico-subcortical circuits

Introduction

Oscillatory neural activity is proposed to orchestrate brain function, by providing multilateral rhythmic synchronisation of neural excitability and action potential firing[1]. One prominent example in the human brain is the beta rhythm (13 – 35 Hz), which has traditionally been described as a phenomenon of the motor domain, primarily localised to motor cortex and important for the pathophysiology of movement disorders, such as Parkinson's disease (PD)[2,3]. In PD, beta activity, especially in deeper brain structures, such as the basal ganglia, has been associated with the loss of innervation with dopamine[4], a neurotransmitter critically involved in synaptic plasticity and reinforcement learning at cortico-striatal synapses[5]. This relationship is mirrored in rodent and non-human primate models of PD, corroborating the role of dopamine as a key neurotransmitter governing beta synchrony. Meanwhile, the putative function of beta modulation in both human and animal literature has been extended within and beyond PD, to diverse domains of cognition[3], memory[6] and emotions[7], but only few studies so far have sought to relate these observations to dopamine signalling. For each of these domains, localised synchronisation or desynchronisation phenomena in the beta frequency range have been described in specific brain regions. What the common denominators of these brain regions are, and whether beta activity has a general structural and functional overlap with dopaminergic innervation remains unknown. This reflects a significant roadblock to a fundamental understanding of the circuit architecture of beta activity. To overcome this roadblock, we have developed a unique multimodal research data pipeline combining normative sources of invasive brain signal recordings, MRI connectomics and positron emission tomography (PET). We address the hypothesis that beta activity is a ubiquitous synchronisation phenomenon, that is characterised by whole-brain networks with subcortical nuclei that are under modulatory control of dopamine.

Results

Beta activity is a widely distributed resting brain rhythm in invasive cortical recordings

First, we aimed to identify the spatial distribution of cortical sources of beta activity in the resting human brain. Electrophysiology studies describing brain oscillations often rely on non-invasive electroencephalography (EEG) or magnetoencephalography (MEG) that provide access to brain signal recordings in healthy subjects. While beamforming and other source reconstruction techniques can provide estimates of the spatial peaks of certain brain activity patterns in these recordings, it can be hard to rule out that the derived brain regions are free from influence of volume conduction and current mixing from neighbouring brain areas[8]. Thus, estimates of beta activity in prefrontal areas identified with EEG or MEG could to some extent, be the result of motor cortex activity, contaminating sensors in frontal locations. Invasive recordings directly from brain tissue can circumvent such contamination, but are unavailable in healthy subjects. Here, we analysed over 30 hours of invasive resting state brain signals from an open atlas[9] of the normal electroencephalogram in 1772 channels of either electrocorticography (ECoG; n=258) or stereoelectroencephalography (sEEG; n=1514) covering the human cortex. The atlas is derived from 106 epilepsy patients undergoing monitoring for seizure activity for respective surgery, but the signals were selected based on strict criteria to ensure their origin from healthy brain regions,

in relevant distance to epileptogenic zones or lesions[9]. The localised recordings from bipolar channels of one minute resting periods with eyes closed were taken to the frequency domain using Morlet wavelets and parametrised using the "fitting oscillations and one over f" (FOOOF) toolbox[10], to extract the maximum peak amplitude of the periodic component of beta activity (13 – 30 Hz). For comparison and as a control, we extended this analysis to other canonical brain rhythms, including theta (4 – 8 Hz), alpha (8 – 12 Hz) and gamma (30 – 100 Hz) frequency bands. The frequency band with the highest peak amplitude was identified for each channel and depicted as the dominant rhythm for the respective localisation (Figure 1). This revealed that 15.6% (277) channels were identified as theta dominant, 22.4% (397) as alpha dominant, 56.6% (1005) as beta dominant and 5.2% (93) as gamma dominant channels. The focus of this article is beta oscillations, so all further analyses will refer to beta with alpha for comparison as it was the second most frequently occurring frequency band after beta.

Figure 1: Beta activity is a widely distributed resting brain rhythm in invasive cortical recordings. (A) Exemplar sEEG electrodes with dominant beta (teal; 20 Hz) or dominant alpha (purple; 9 Hz) rhythms, alongside the respective raw signal and parametrised power spectrum. **(B)** 1005/1772 channels showed higher maximum peak power in beta (13 – 30 Hz) than theta (4 – 8 Hz), alpha (8 – 12 Hz) and gamma (30 – 100 Hz). 397/1772 were alpha dominant (theta and gamma not shown). **(C)** Beta dominant channels were distributed across the entire cortex, from occipital to frontal regions, with highest density in sensorimotor areas like precentral gyrus, frontal middle gyrus, frontal inferior gyrus and supplementary motor area. Alpha channels were largely concentrated posteriorly in temporoparietal and occipital areas based on the automatic anatomical labelling atlas[11]. Neither, alpha nor beta activity showed systematic hemispheric differences (p>0.05).

To investigate the regional distributions of frequency bands, we counted the occurrence of each dominant rhythm in gyri and lobes (see Figure 1 and Table 1), by projecting the coordinates in MNI (Montreal Neurological Institute) space to the automated anatomical labelling (AAL) atlas parcellation[11]. Notably, many channels showed more than one oscillatory peak, and all covered lobes had channels with dominant rhythms in each, theta, alpha and beta frequency bands (see Table 1). Beta activity was the most frequent and most widely distributed resting rhythm across all lobes of the human brain including frontal, temporal, cingulate, insula and parietal lobes. Interestingly, the occipital lobe was the only brain region, that showed higher percentage of alpha dominant channels (53/94; 56.4%). The coverage of subdural ECoG grids or tissue penetrating sEEG electrodes could vary, but within overlapping regions, no major differences in the distribution of brain rhythms were observed for electrode type (see supplementary Table 1).

Lobe	Channels/lobe	ECoG	sEEG	Sum theta channels		Sum alpha channels		Sum beta channels	
Frontal	616	122	494	73	11.8%	38	6.2%	458	73.4%
Temporal	519	68	451	106	20.4%	180	35.0%	216	41.6%
Cingulate	116	0	116	12	10.3%	21	18.1%	71	61.2%
Insula	128	2	126	19	14.8%	21	16.4%	86	67.2%
Amygdala	3	0	3	1	33.3%	0	0%	1	33.3%
Parietal	272	52	220	50	18.4%	73	26.8%	139	51.1%
Occipital	107	13	94	15	14.0%	62	57.9%	27	25.2%
Subcortex	11	1	10	1	9.1%	2	18.2%	7	63.6%

Table 1: Spread of invasive electrophysiological channels. Given the low number of channels exhibiting resting state gamma activity, we excluded those from the present table.

MRI connectomics reveal shared cortico-subcortical beta networks

Oscillatory synchronisation across brain regions may shape neural communication through modulation of synaptic sum potentials and neural firing[1]. Oscillatory connectivity can be computed through correlation and modelling of phase and amplitude relationships between brain signals from multielectrode recordings. The spatial coverage of such computations, however, is inherently limited to brain regions that are accessible with the respective recording technique. For non-invasive M/EEG, the analysis of connectivity is limited to superficial cortical brain areas, and can be biased by noise and volume conduction[12]. Invasive recordings in both human and animal studies provide more robust estimates of oscillatory connectivity but are limited to a selected number of channels from defined brain regions[13]. Here, we aimed to investigate the whole brain circuit representation of beta activity, which is impossible with current neurophysiology approaches. Instead, we tested the hypothesis that cortical brain areas that exhibit resting beta share a global brain network by means of MRI connectomics. For this, we derived the methodology from lesion and DBS network mapping[14], which has provided groundbreaking insights into the circuit nature of brain lesion and neurostimulation effects and applied it to reveal the shared network fingerprints of brain regions that exhibit a neurophysiological phenomenon; namely, dominant beta activity. For this we followed a multimodal approach, investigating neurophysiologically identified brain regions by means of MRI connectomics, generating functional and structural connectivity maps for

each brain signal recording location, as previously described[15]. In brief, the MNI coordinates of the invasive electrode locations were used as seeds to compute whole-brain connectivity maps based on MRI data from 1000 healthy subjects as available through the Brain Genomics Superstruct project, and MRI data from 32 subjects as available through the Human Connectome Project (HCP) as part of the Lead Mapper pipeline[16]. While the connectivity maps estimated with functional MRI (fMRI) are based on the covariance blood oxygenation level dependent signals (BOLD) that provide an estimate of functional coactivation, the structural connectivity maps are based on diffusion weighted MRI (dMRI) and provide a relative quantification of major anatomical white matter tracts. To allow for a statistical comparison of connectivity patterns, we compared voxel-wise connectivity estimates from recording locations with dominant beta rhythms, to recording locations with dominant alpha rhythms using mass univariate two-sample t-tests in Statistical Parametric Mapping software, with a family wise error corrected voxel threshold set to $p<0.05$.

Figure 2: MRI connectomics reveal shared cortico-subcortical beta networks. Individual network fingerprints **(A)** seeded from representative beta (top) and alpha (bottom) dominant electrodes highlight the methodological approach and showcase the distinct connectivity patterns that arise in dependence of the connected brain regions (for raw signal traces from these locations see Figure 1). Aggregated functional **(B)** and structural **(C)** connectivity maps across all beta (top) and alpha (bottom) dominant electrodes were subjected to mass univariate t-tests comparing beta vs. alpha networks with Statistical Parametric Mapping software and visualised as T-maps. Channel locations from beta dominant channels were associated with more robust functional and structural connectivity to frontal cortex and the basal ganglia, when compared to alpha channel locations, which were more connected to the occipital cortex.

Our analysis revealed a shared whole brain network for cortical brain regions that exhibit dominant beta oscillations, with marked and highly significant differences (Figure 2) to the control network for alpha oscillations, both in whole-brain functional and structural connectivity patterns (see Supplementary Figure 1 for unthresholded averages). Most prominently, the beta network showed significantly higher connectivity to prefrontal, cingulate and parietal cortical areas and the ventral thalamus and basal ganglia, including substantia nigra, striatum, pallidum and subthalamic nucleus (Supplementary Figures 2 & 3 for brain region averages parcellated with the automatic anatomical labelling atlas). In contrast, whole brain networks seeded from alpha-dominant channels showed a more widespread distribution across the posterior cortical areas, occipital and temporal lobes in comparison to the beta networks.

The connectomic beta network correlates with molecular markers of dopamine signalling
Invasive brain signal recordings in patients undergoing deep brain stimulation (DBS) for Parkinson's disease (PD) have robustly established a strong interaction between dopaminergic tone and amplitude of beta oscillations at rest, e.g. by measuring changes in activity after withdrawal and administration of dopaminergic agents[17]. However, this interaction has been largely neglected outside the field of PD, likely because research methods to establish direct relationships between beta and dopamine in healthy subjects are not readily available. With the present project, we aimed to characterise the circuit architecture of cortical beta oscillations. We found that cortical brain areas, that exhibit resting beta activity share a common brain network. This network, unrelated to pathology, includes strong connectivity with the basal ganglia, the primary target for dopaminergic innervation from the substantia nigra and the ventral tegmental area in the human brain. To further quantify the potential relationship between the beta network and dopamine signalling, we made use of another openly available human neuroscience resource: a database of positron emission tomography (PET) imaging, a molecular imaging technique that relies on intravenous injection of a radioactively labelled molecular tracer, that can be visualised to indicate brain areas with high uptake of neurotransmitters or related molecules in the brain[18]. Specifically, we analysed to what degree the spatial uptake patterns of dopamine, as measurable with fluorodopa (FDOPA; cohort average of 12 healthy subjects) and other dopamine signalling related tracers that bind D1/D2 receptors (average of N=17/44 respectively healthy subjects) or the dopamine transporter (DAT; cohort average of N=180 healthy subjects) were correlated with the beta networks. To reduce the dimensionality and complexity of this analysis, we first applied spatial normalisation and then averaged the abovementioned cohort images to yield a PET based dopamine aggregate map (Figure 3). Next, we resliced the beta network map and the PET images to allow for a meaningful comparison, using a combined parcellation that includes cortex[19], basal ganglia[20], and cerebellum[21]. Finally, we performed non-parametric Spearman's correlations between the structural and functional connectivity maps of beta networks with neurotransmitter uptake. This revealed a significant positive relationship between the beta functional and structural networks and dopamine uptake for cortex (rho=0.22/0.40 for functional/structural networks, P=0.0001) and basal ganglia (rho=0.50/0.33, P=0.0001), but not cerebellum (all P>0.05). These patterns were largely qualitatively reproduced, when the analysis was performed on the individual tracers (see Supplementary Figure 4). Additionally, we repeated the analysis with PET images of GABA uptake, to serve as an unspecific molecular control, which revealed no significant relationship between the beta functional network and GABA uptake cortically (rho=0.04, P=0.22), in the basal

ganglia (rho=0.07, P=0.27) or cerebellum (rho=0.10, P=0.22). The structural beta network revealed low correlation with GABA uptake for cortical areas (rho=0.17, P=0.0003), negative correlations for basal ganglia (rho=-0.55, P=0.0001) and no correlation for cerebellum (P>0.05). In summary, we found significant and specific overlap of the functional and structural connectivity maps seeded from beta-dominant cortical brain areas with the spatial patterns of dopamine uptake in the human brain.

Figure 3: The connectomic beta network correlates with molecular markers of dopamine signalling. Dopamine uptake significantly correlates with beta functional connectivity **(A)** A methods schematic explaining how correlations were calculated in Figure 3. Each point on the scatterplot represents a parcellation from the compound brain atlas. For every parcel X_i, a correlation is calculated for that parcel in the PET scan and in the beta network. **(B)** in cortex (rho=0.22, P=0.0001) and basal ganglia (rho=0.5, P=0.0001) but not cerebellum (P>0.05), and beta structural connectivity **(C)** in cortex (rho=0.4, P=0.0001) and basal ganglia (rho=0.33, P=0.001) but not cerebellum (P>0.05). GABA was used as a control molecule which revealed no significant correlations with beta functional connectivity **(D)** in cortex, basal ganglia or cerebellum (P>0.05). Beta structural connectivity **(E)** revealed weaker positive correlation with GABA in cortex (rho=0.17, P=0.0003), a negative correlation in the basal ganglia (rho=0.55, P=0.0001) and no correlation in cerebellum (P>0.05).

Discussion

The present study combined three openly available datasets of invasive neurophysiology, MRI connectomics and molecular neuroimaging in healthy humans to characterise the spatial distribution of brain regions exhibiting resting beta activity, their shared circuit architecture, and its correlation with molecular markers of dopamine signalling in the human brain. Our results can be broken down into three key insights: First, we demonstrate that the beta rhythm is the most widely distributed dominant neural synchronisation phenomenon, extending far beyond motor areas. Next, we show that these distributed areas share a common functional and structural whole-brain network that connects frontal, cingulate and other large-scale brain areas with the basal ganglia. Finally, we show that this network shares significant spatial overlap with molecular markers of dopamine signalling. We interpret these findings as supportive evidence for the hypothesis that beta activity is generated and sustained by a cortico-subcortical brain network that is modulated by dopamine. This conceptualisation moves away from functionally specific domains, towards a more holistic circuit understanding of beta activity as a brain circuit synchronisation phenomenon.

Brain functions associated with beta oscillations share brain networks modulated by dopamine

Our study investigates the connectivity profile of cortical brain regions that spontaneously exhibit beta oscillations as the dominant brain rhythm. Thus, our approach does not allow for a direct comparison of the role of beta oscillations in human or animal behaviour. Our results motivate to conceptualise the capacity to generate beta oscillations as a feature that arises from a large-scale global brain network that is modulated by dopamine. However, we do not see this concept as contradictory to previous reports of functional correlates of beta activity in the domains of motor control, cognition, memory and emotion[3,6,7,22]. Instead, we believe that the overlap in oscillatory activity across these domains can be conceptualised through a circuit definition of beta activity and its relationship to dopamine signalling. Foremost, in the motor domain there is direct evidence that pathological changes in dopamine causally modulates beta activity, as seen in human patients with Parkinson's disease[17], but also in animals lesioned with neurotoxins targeting dopaminergic neurons, such as 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) in non-human primates[23] or 6-hydroxydopamine (6-OHDA) in rodents[24]. Meanwhile, decades of research have documented the robust relationship of dopaminergic midbrain activity with movement[25] and recent advances in fiberphotometry combined with designed genetically encoded sensors like dLight1[26] now provide direct access to measure temporally specific dopamine release and have sparked a renaissance in the characterisation of dopamine signals associated with motor control in rodents[27]. Similarly, the robust beta modulation that is elicited by voluntary action in sensorimotor cortex and its correlation with motor symptoms of Parkinson's disease is long known. Even in our study, motor and sensorimotor cortex showed higher beta peak amplitudes than any other brain regions. Thus, it is tempting to infer that beta activity and its relation to dopamine is characteristic to the motor domain. On the other hand, numerous studies have reported specific changes in beta activity in relation to many other neuroscientific domains that define human ability. For example, it was shown that beta is implicated in working memory[28], utilisation of salient sensory cues[29], language processing[30],

motivation[31], sleep[32], emotion recognition[33] and may even serve as a biomarker for depressive symptom severity in the anterior cingulate cortex[34]. Our connectomic characterisation of the beta network provides a network basis for these functions and highlights that similar synchronisation phenomena occur in brain areas associated with vastly different functions, such as prefrontal, cingulate and parietal areas, which give rise to parallel striatal and other subcortical projections. In line with this more complex picture, first fast-cyclic voltammetry experiments that allowed for combined measurement of dopamine release with invasive neurophysiology have provided first evidence that beta band oscillations in healthy non-human primates can differentially link dopamine release, beta oscillations and reward and motor control, depending on the contextual information and striatal domain[35]. Thus, an obvious differentiator of these reported functions can be the specific brain region in which beta activity has been recorded from. However, the aim of our study was the opposite, namely to provide evidence for a shared network that unifies these functions. An authoritative perspective article has proposed that the unifying function of beta oscillations is to signal the maintenance of "status quo", e.g. to indicate whether or not a motor or cognitive state should be changed or not[36]. We hypothesise that, this "status quo" hypothesis could be equally or maybe even more adequately posed on the neural level. In fact, one could argue that it can be contextualised in a recently described framework of neural reinforcement, that serves to orchestrate the re-entrance and refinement of neural population dynamics for the production of neural trajectories[37]. On the basis of our observed network architecture described for beta oscillations in this article, we speculate that the "status quo" hypothesis and each of the abovementioned behavioural correlates can be summarized to reflect states with either loss or invigoration of neural reinforcement[38], the brain's intrinsic capacity to modulate the stability of population dynamics via cortico-striatal projections that are orchestrated by dopamine. To simplify this, maintenance of a behavioural, cognitive or emotional "status quo" may be reflected by beta activity on the neural circuit level, which in turn relates to the stability of neural dynamics in frontostriatal circuits that can be dynamically modulated by tonic and phasic changes of dopamine release. If beta is high, the neural dynamics remain stable. If beta is low, new neural trajectories may arise that lead to neural and behavioural state changes and neural learning. Thus, we speculate that the observed beta network and its relation to molecular markers of dopamine in this study may indicate that beta modulation occurs in situations that are associated with neural learning and optimisation of neural trajectories, or lack thereof. Notably, many functional correlates of beta activity could be summarised under the "status quo" hypothesis, but some functions are not easily interpreted in this framework, such as the specific observations in the context of memory processing[28], or regulation of emotions and depression. On the other hand, the role for dopaminergic reinforcement and its implication in these processes is robustly established, both in memory[39] and the pathophysiology of depression[40]. At this point, this discussion remains speculative, but advances in neuroscientific research methods that combine neurophysiology with neurotransmitter release may soon provide new evidence in favour or in opposition to these concepts.

Implications for neurotechnology and clinical brain computer interfaces
Neurotechnology is on the rise, promising to revolutionise the treatment of brain disorders[41]. A key component of clinical brain computer interfaces is the bidirectional communication between brain activity and closed-loop control of neurotherapies. The robust association of the hypodopaminergic

state in Parkinson's disease with excessive beta synchrony has led to the conceptualisation of a closed-loop neurostimulation therapy using beta activity as a feedback signal to adjust administration of deep brain stimulation (DBS) in real-time to therapeutic demand[42]. This approach, also termed adaptive DBS, was first demonstrated based on cortical beta activity that was used to adapt pallidal DBS in the MPTP non-human primate model of PD[43], after which it was quickly translated to first experimental studies using cortical or subcortical beta signals in human patients[44]. Practically, beta activity is used in this novel therapeutic approach to infer the dopaminergic tone and consequential therapeutic demand in PD, for rapid adaptation of therapeutic delivery to the individual state that the patients are facing in real-time. Our study provides support for the validity of this approach, by providing further evidence for a relationship of dopamine and beta activity at the global circuit level. More importantly, our findings may suggest that this relationship could be of greater utility of neurotechnology to treat other dopaminergic disorders. In fact, this notion is in part already backed by empirical data, that shows that beta can be a valuable biomarker for depression severity as identified by machine learning algorithms in invasive neurophysiological recordings from brain implants[7]. Given our observation that beta activity is the most widespread cortical brain rhythm, it is foreseeable that future experimental in-human trials aiming to test the utility of invasive neurophysiology for the adaptation of neurotherapies will be capable to track beta dynamics in higher frequency (sampling rate) and higher precision than any other brain rhythm. Our findings could encourage clinical neuroscience and neural engineering teams to conceptualise the dynamic fluctuations of beta oscillations in context of cortico-striatal processing and dopaminergic modulation, which could lead to new hypothesis driven treatment avenues. Specifically, our demonstration that the distributed occurrence of beta activity shares a brain network that is modulated by dopamine, may motivate the use of beta oscillations to control brain computer interfaces or closed-loop neurostimulation for emerging indications related to dopaminergic dysfunction. In the future, targeted interventions that can increase and decrease beta activity, as recently shown through phase specific modulation[45] could be utilised to mimic phasic dopamine release as a neuroprosthetic approach to alter neural reinforcement[38].

Limitations
While the abovementioned prospects are promising we should cautiously consider the limitations of our study. The most important limitation of this project is aggregated and derived nature of the used brain data. Three independent datasets from different study cohorts, were leveraged to fuse the invasive neurophysiology with MRI and PET based neuroimaging. On the one hand, the independence of the source data may be seen as a form of validation, e.g. for functional and structural connectivity that showed highly overlapping findings while stemming from entirely different cohorts. On the other hand, precisely this aspect severely limits the generalisability of our results to the level of the individual human subject. Similarly, the foundation of the results is built on patients undergoing brain surgery for epilepsy, a severe intractable brain disorder. But at the same time, the reported observations follow a long line of research and corroborate anecdotal or clinical knowledge, that beta activity is a ubiquitous phenomenon. In this light, our study provides an unprecedented and methodologically robust quantitative approach, by using only localised bipolar invasive recordings with oscillatory activity parametrised for optimal comparability. We believe that this rigorous neurophysiological methodology, alongside the fact that the brain signals have been meticulously controlled for pathological activity, by several experienced

neurophysiologists, justifies its use and superiority over aggregates of non-invasive recordings from healthy subjects. Another limitation in our study is the fact that the statistical approach for the comparison of beta and alpha networks follows a winner takes all logic that is, by definition, a simplification, as most areas will contribute to more than one spatio-spectrally distinct oscillatory network. A previous study combining invasive basal ganglia neurophysiology with non-invasive cortical magnetoencephalography (MEG) recordings has identified the strongest oscillatory network of the internal pallidum, to connect to frontal cortex in the beta frequency range, which is in line with our findings[46]. However, additional networks, such as an alpha network with the cerebellum and a theta network with the temporal lobe were also observable in these parallel recordings. Thus, the presented MRI based statistical maps in this study can be interpreted to reflect shared large-scale and robustly estimable whole-brain circuits of dominant beta vs. alpha oscillations but should not be misunderstood to reflect mutual exclusivity for the presence or communication in additional brain rhythms. In fact, when looking at individual parcels of structural connectivity, it is noticeable, that both beta and alpha networks share strong connections to cingulate cortex. This is interesting, because previous invasive neurophysiology studies in patients undergoing deep brain stimulation surgery for depression have reported both excessive alpha and beta activity as potential biomarkers for depressive symptoms in the human cingulum[34], potentially related to differential modulation of neurotransmitters involved in the disease. It should also be noted that in this project, the data are static, but beta oscillations and dopamine are dynamic[47]. Therefore, while we cannot infer the direction of the relationship, it is very likely that the relationship would be negative implying higher beta would mean lower dopamine. Finally, our study is limited to resting state activity and thus does not allow for interpretation of functional changes with relation to behaviour or neural processes, that may individually be associated with other or additional brain rhythms. Most notably, gamma band oscillations are known to be reflective of local processing but notoriously absent in resting state activity and thus are likely equally or even more ubiquitous than beta activity. However, we do not see this as a contradiction, as resting beta activity provides insights into a different state of the cortical brain area, potentially more associated with global circuit synchrony, while gamma band activity may be more reflective of local activity in the cortical microcircuit.

Conclusion

In summary, our study suggests that the beta rhythm is the most widespread cortical oscillatory activity pattern in the resting brain, that shares connectivity with the basal ganglia via a whole brain network that is modulated by dopamine. We interpret our findings to provide a unifying circuit-based integration of previous reports on the functional role of beta oscillations. In the future, our findings may inspire and encourage the investigation of beta oscillations for the use as a feedback signal for neurotechnological interventions in the treatment of dopaminergic disorders.

Methods

Atlas of invasive electrophysiology

The collection of data, cohort and electrophysiology of the MNI Open iEEG Atlas (https://mni-open-ieegatlas.research.mcgill.ca/)is as described in the work of Frauscher and colleagues[9]. Intracranial EEG (iEEG) recordings from the healthy hemispheres of epilepsy patients were collected from three hospitals across Canada. Inclusion criteria for the recordings included that it must 1) contain at least one channel with normal activity, 2) have peri-implantation imaging, 3) be recorded after a minimum of 72 hours post-implantation for stereo-EEG electrodes or 1 week for subdural grids or strips, and 4) have a minimum sampling frequency of 200 Hz. The recordings were analysed to identify frequencies in the classical Berger frequency bands (0.3−70 Hz). Tissue containing channels exhibiting normal activity was classified as normal or healthy tissue. This meant it was outside of the zone of seizure onset, i.e., no interictal epileptic discharge and no discharges or slow wave anomalies were found there. Ethical approval was granted at the MNI (Montreal Neurological Institute) as lead ethics organization (REB vote: MUHC-15-950). Electrode locations were registered to a common stereotaxic space.

Oscillatory analysis and neurophysiological pre-processing

Data from the iEEG atlas were preprocessed using the Statistical Parametric Mapping toolbox (https://www.fil.ion.ucl.ac.uk/spm12/). The original data contained zero padded intervals between signal fragments which were deleted. Next the data were Fourier transformed to obtain a time-frequency signal using Morlet Wavelets with seven cycles. All time-frequency signals were averaged over time to obtain one frequency spectrum of 60 second sections of iEEG measurements. The resulting power spectra were quantified in terms of spectral peak properties and background level activity using FOOOF[10] with the following parameters; limits for possible peak widths were [0.5, 12 Hz], the maximum number of detectable peaks were set to infinite so as to not limit the number of peaks detected, the minimum peak height was 0 while the peak threshold was within 2 standard deviations of modelled power. Extracted spectral properties were z-scored within electrodes, within subjects and within the whole dataset to quantify neural activity from several frequency bands of the spectrum. Frequency-specific activity was quantified using maximum peak amplitude for specific frequency ranges. The maximum peak amplitude was extracted from the z-scored parameters for four frequency bands; theta (4−8 Hz), alpha (8−12 Hz), beta (13−35 Hz), and gamma (30−100 Hz).

Connectivity analysis

Resting state functional connectivity data was collected from 1,000 healthy individuals using a 3T Siemens MRI scanner as part of the Brain Genomics Superstruct Project. Preprocessing included global signal regression and spatial smoothing at 6mm full width at half maximum. The group connectome for structural connectivity was obtained from multishell diffusion-weighted and T2-weighted imaging data subjects from the Human Connectome Project[48]. The imaging data were acquired using a custom MRI scanner with enhanced gradients. Whole brain tractography fibre sets were generated using a generalised q-sampling imaging algorithm implemented in DSI Studio. Connectivity analyses were performed in MATLAB using normative connectomes calculated with

Lead-Mapper[49]. Three-dimensional spherical nifti files representing channel locations were extracted and used as seeds to calculate connectivity maps.

The fMRI and dMRI data underwent processing using the Lead Connectome Mapper. Every contact location from the iEEG atlas was represented by a spherical seed nifti image (r=5mm) To minimise the impact of noise on the results, all maps underwent smoothing with a Gaussian kernel (standard deviation = 8mm). Subsequently, the fMRI maps were masked with a grey matter mask to avoid the inclusion of white matter correlations, which would have introduced artifacts.

To investigate differences between beta dominant and alpha dominant functional connectivity networks, a two sample t-test was calculated for the condition where beta was greater than alpha and vice versa using SPM. Here, the connectivity maps from each dominant channel (1005 beta functional connectivity maps and 397 alpha connectivity maps) underwent preprocessing and were entered into a general linear model (GLM). Estimation of model parameters yielded t-values for each voxel, indicating the strength and direction of differences between the two contrasts (beta > alpha, alpha > beta). This incorporated experimental design conditions and nuisance regressors. The GLM also generated a contrast image to compare activation between the two groups. To address the issue of multiple comparisons, we applied Family-Wise Error (FWE) correction, adjusting significance thresholds such that only voxels with $p < 0.05$ would be included. Each voxel received a t-value, representative of differences between groups relative to variability within group. This t-value was converted into an SPM thresholded map that was subsequently used for further analyses.

A functional connectivity map, i.e., a mapping of the functional connectivity (based on a normative connectome) between a seed region and every other brain region, was derived for each individual channel (N=1772). To do so, each channel was used as a seed region and its respective functional connectivity map was calculated, resulting in 1772 connectivity maps, one for every channel. These maps were subdivided based on the channel's dominant frequency band. This resulted in 277 functional connectivity maps for the theta, 397 for the alpha and 1005 for the beta band. Group level statistics were performed for connectivity maps of each frequency band. All functional connectivity maps were averaged in SPM ImCalc to reveal a group-level functional network for each frequency band.

Like functional connectivity maps, structural connectivity maps (N=1772) were calculated for each individual channel. These were again subdivided based on dominant frequency band per channel and gave a resulting set of structural connectivity maps for each frequency band of 277 (theta), 397 (alpha) and 1005 (beta). Each map contained a set of fibre tracts that were identified as correlated with the specific channel through the structural connectivity analysis. Therefore, to investigate group level statistics, the maps had to be summed and not averaged as in the case of functional connectivity. The resulting structural networks for each frequency band were hence a sum calculated on SPM ImCalc of all identified fibres for all channels for a given frequency band.

Neurotransmitter uptake

A custom master parcellation in MNI space was created in Matlab using SPM functions by combining three existing parcellations to include cortical regions[19], structures of the basal ganglia[20] and cerebellar regions[21]. Regions that were (partially) overlapping between the atlases were only selected once. The final compound parcellation had 476 regions in total.

To investigate dopamine uptake, an open dataset of neurotransmitter uptake was used an open data set of PET data measuring neurotransmitter uptake[18]. Exclusively healthy samples of subjects were selected for various dopamine receptors such as D1[50] (healthy controls (HC)=13), D2[51,52] (HC=34), the dopamine transporter (DAT)[53,54] (HC=180) and fluorodopa (FDOPA)[55](HC=12). To study the broad effects of dopamine, all dopamine receptor PET scans were resliced and averaged using the SPM ImCalc function. A spatial normalisation was then applied to finally create a unified aggregate image of all available dopamine receptors.

The dopamine PET scan and the beta connectivity networks were parcellated based on the master parcellation. Spearman's ranked correlation was used to calculate correlations between the dopamine aggregate image and both beta connectivity networks (Figure 3). The individual dopamine receptors were also similarly parcellated and correlated with the beta networks (S2).

To validate these results, a control analysis was performed using a GABA PET scan from the same open dataset of neurotransmitter uptake. This GABA image was also parcellated and correlated with the two beta networks (Figure 2A, 2B).

Data availability

All data used in the course of this work were publicly available open datasets and can be found online.

1. The iEEG atlas is available on the McGill database and can be accessed via their publication (https://www.mcgill.ca/frauscher-lab/files/frauscher-lab/frauscher_brain2018.pdf).
2. The normative connectomes are available within the Lead-DBS software (www.lead-dbs.org).
3. The PET scans for dopamine uptake as well as their data pipeline are available on the Network Neuroscience Lab GitHub repository(https://github.com/netneurolab/hansen_receptors).

Code availability

All original code used to preprocess and analyze these data, and create the main and supplementary results is openly available on a GitHub repository (https://github.com/meera-cm/iEEG_connectomics).

Additional code from the open datasets is available:

1. Lead-DBS and Lead-Connectome Mapper: (https://github.com/leaddbs/leaddbs)
2. Hansen receptors: (https://github.com/netneurolab/hansen_receptors/tree/main/code)

References

1. Fries, P. Rhythms for Cognition: Communication through Coherence. *Neuron* **88**, 220–235 (2015).

2. Hammond, C., Bergman, H. & Brown, P. Pathological synchronization in Parkinson's disease: networks, models and treatments. *Trends Neurosci* **30**, 357–364 (2007).

3. Paulo, D. L. *et al.* Corticostriatal beta oscillation changes associated with cognitive function in Parkinson's disease. *Brain* **146**, 3662–3675 (2023).

4. Iskhakova, L. *et al.* Modulation of dopamine tone induces frequency shifts in cortico-basal ganglia beta oscillations. *Nat Commun* **12**, 7026 (2021).

5. Berke, J. D. What does dopamine mean? *Nat Neurosci* **21**, 787–793 (2018).

6. Hanslmayr, S., Staresina, B. P. & Bowman, H. Oscillations and Episodic Memory: Addressing the Synchronization/Desynchronization Conundrum. *Trends in Neurosciences* **39**, 16–25 (2016).

7. Alagapan, S. *et al.* Cingulate dynamics track depression recovery with deep brain stimulation. *Nature* **622**, 130–138 (2023).

8. Hedrich, T., Pellegrino, G., Kobayashi, E., Lina, J. M. & Grova, C. Comparison of the spatial resolution of source imaging techniques in high-density EEG and MEG. *NeuroImage* **157**, 531–544 (2017).

9. Frauscher, B. *et al.* Atlas of the normal intracranial electroencephalogram: neurophysiological awake activity in different cortical areas. *Brain* **141**, 1130–1144 (2018).

10. Donoghue, T. *et al.* Parameterizing neural power spectra into periodic and aperiodic components. *Nat Neurosci* **23**, 1655–1665 (2020).

11. Rolls, E. T., Huang, C.-C., Lin, C.-P., Feng, J. & Joliot, M. Automated anatomical labelling atlas 3. *NeuroImage* **206**, 116189 (2020).

12. Pellegrini, F., Delorme, A., Nikulin, V. & Haufe, S. Identifying good practices for detecting inter-regional linear functional connectivity from EEG. *NeuroImage* **277**, 120218 (2023).

13. Engel, A. K., Moll, C. K. E., Fried, I. & Ojemann, G. A. Invasive recordings from the human brain: clinical insights and beyond. *Nat Rev Neurosci* **6**, 35–47 (2005).

14. Fox, M. D. Mapping Symptoms to Brain Networks with the Human Connectome. *New England Journal of Medicine* **379**, 2237–2245 (2018).

15. Lofredi, R. *et al.* Subthalamic stimulation impairs stopping of ongoing movements. *Brain* **144**, 44–52 (2021).

16. Neudorfer, C. *et al.* Lead-DBS v3.0: Mapping deep brain stimulation effects to local anatomy and global networks. *NeuroImage* **268**, 119862 (2023).

17. Lofredi, R. *et al.* Subthalamic beta bursts correlate with dopamine-dependent motor symptoms in 106 Parkinson's patients. *npj Parkinsons Dis.* **9**, 1–9 (2023).

18. Hansen, J. Y. *et al.* Mapping neurotransmitter systems to the structural and functional organization of the human neocortex. *Nat Neurosci* **25**, 1569–1581 (2022).

19. Huang, C.-C., Rolls, E. T., Feng, J. & Lin, C.-P. An extended Human Connectome Project multimodal parcellation atlas of the human cortex and subcortical areas. *Brain Struct Funct* **227**, 763–778 (2022).

20. He, X. *et al.* Disrupted basal ganglia–thalamocortical loops in focal to bilateral tonic-clonic seizures. *Brain* **143**, 175–190 (2020).

21. Diedrichsen, J., Balsters, J. H., Flavell, J., Cussans, E. & Ramnani, N. A probabilistic MR atlas of the human cerebellum. *Neuroimage* **46**, 39–46 (2009).

22. Neumann, W.-J., Horn, A. & Kühn, A. A. Insights and opportunities for deep brain stimulation as a brain circuit intervention. *Trends in Neurosciences* **46**, 472–487 (2023).

23. Deffains, M. & Bergman, H. Parkinsonism-related β oscillations in the primate basal ganglia networks – Recent advances and clinical implications. *Parkinsonism & Related Disorders* **59**, 2–8 (2019).

24. Sharott, A. *et al.* Dopamine depletion increases the power and coherence of β-oscillations in the cerebral cortex and subthalamic nucleus of the awake rat. *European Journal of Neuroscience* **21**, 1413–1422 (2005).

25. Romo, R. & Schultz, W. Dopamine neurons of the monkey midbrain: contingencies of responses to active touch during self-initiated arm movements. *Journal of Neurophysiology* **63**, 592–606 (1990).

26. Patriarchi, T. *et al.* Ultrafast neuronal imaging of dopamine dynamics with designed genetically encoded sensors. *Science* **360**, eaat4422 (2018).

27. Coddington, L. T. & Dudman, J. T. Learning from Action: Reconsidering Movement Signaling in Midbrain Dopamine Neuron Activity. *Neuron* **104**, 63–77 (2019).

28. Miller, E. K., Lundqvist, M. & Bastos, A. M. Working Memory 2.0. *Neuron* **100**, 463–475 (2018).

29. Leventhal, D. K. *et al.* Basal Ganglia Beta Oscillations Accompany Cue Utilization. *Neuron* **73**, 523–536 (2012).

30. Meyer, L. The neural oscillations of speech processing and language comprehension: state of the art and emerging mechanisms. *European Journal of Neuroscience* **48**, 2609–2621 (2018).

31. Pierrieau, E., Berret, B., Lepage, J.-F. & Bernier, P.-M. From Motivation to Action: Action Cost Better Predicts Changes in Premovement Beta-Band Activity than Speed. *J. Neurosci.* **43**, 5264–5275 (2023).

32. Yin, Z. *et al.* Pathological pallidal beta activity in Parkinson's disease is sustained during sleep and associated with sleep disturbance. *Nat Commun* **14**, 5434 (2023).

33. Jabbi, M. *et al.* Convergent BOLD and Beta-Band Activity in Superior Temporal Sulcus and Frontolimbic Circuitry Underpins Human Emotion Cognition. *Cerebral Cortex* **25**, 1878–1888 (2015).

34. Clark, D. L., Brown, E. C., Ramasubbu, R. & Kiss, Z. H. T. Intrinsic Local Beta Oscillations in the Subgenual Cingulate Relate to Depressive Symptoms in Treatment-Resistant Depression. *Biological Psychiatry* **80**, e93–e94 (2016).

35. Schwerdt, H. N. *et al.* Dopamine and beta-band oscillations differentially link to striatal value and motor control. *Sci Adv* **6**, eabb9226 (2020).

36. Engel, A. K. & Fries, P. Beta-band oscillations—signalling the status quo? *Current Opinion in Neurobiology* **20**, 156–165 (2010).

37. Athalye, V. R., Carmena, J. M. & Costa, R. M. Neural reinforcement: re-entering and refining neural dynamics leading to desirable outcomes. *Curr Opin Neurobiol* **60**, 145–154 (2020).

38. Cavallo, A. & Wolf-Julian, N. *Dopaminergic reinforcement in the motor system: Implications for Parkinson's disease and deep brain stimulation.* https://www.authorea.com/users/667616/articles/667773-dopaminergic-reinforcement-in-the-motor-system-implications-for-parkinson-s-disease-and-deep-brain-stimulation?commit=b2d07512e5a5f31d444c5c0985d6ccd73f6cc3d6 (2023) doi:10.22541/au.169536617.79800198/v1.

39. Rossato, J. I., Bevilaqua, L. R. M., Izquierdo, I., Medina, J. H. & Cammarota, M. Dopamine Controls Persistence of Long-Term Memory Storage. *Science* **325**, 1017–1020 (2009).

40. Dunlop, B. W. & Nemeroff, C. B. The Role of Dopamine in the Pathophysiology of Depression. *Archives of General Psychiatry* **64**, 327–337 (2007).

41. Borton, D. A., Dawes, H. E., Worrell, G. A., Starr, P. A. & Denison, T. J. Developing Collaborative Platforms to Advance Neurotechnology and Its Translation. *Neuron* **108**, 286–301 (2020).

42. Neumann, W.-J., Gilron, R., Little, S. & Tinkhauser, G. Adaptive Deep Brain Stimulation: From Experimental Evidence Toward Practical Implementation. *Movement Disorders* **38**, 937–948 (2023).

43. Rosin, B. *et al.* Closed-Loop Deep Brain Stimulation Is Superior in Ameliorating Parkinsonism. *Neuron* **72**, 370–384 (2011).

44. Little, S. *et al.* Adaptive deep brain stimulation in advanced Parkinson disease. *Annals of Neurology* **74**, 449–457 (2013).

45. McNamara, C. G., Rothwell, M. & Sharott, A. Stable, interactive modulation of neuronal oscillations produced through brain-machine equilibrium. *Cell Reports* **41**, (2022).

46. Neumann, W.-J. *et al.* Cortico-pallidal oscillatory connectivity in patients with dystonia. *Brain* **138**, 1894–1906 (2015).

47. Tinkhauser, G. *et al.* The modulatory effect of adaptive deep brain stimulation on beta bursts in Parkinson's disease. *Brain* **140**, 1053–1067 (2017).

48. Setsompop, K. *et al.* Pushing the limits of in vivo diffusion MRI for the Human Connectome Project. *NeuroImage* **80**, 220–233 (2013).

49. Horn, A. & Kühn, A. A. Lead-DBS: A toolbox for deep brain stimulation electrode localizations and visualizations. *NeuroImage* **107**, 127–135 (2015).

50. Kaller, S. *et al.* Test–retest measurements of dopamine D1-type receptors using simultaneous PET/MRI imaging. *Eur J Nucl Med Mol Imaging* **44**, 1025–1032 (2017).

51. Alakurtti, K. *et al.* Long-Term Test–Retest Reliability of Striatal and Extrastriatal Dopamine D2/3 Receptor Binding: Study with [11C]Raclopride and High-Resolution PET. *J Cereb Blood Flow Metab* **35**, 1199–1205 (2015).

52. Smith, C. T. *et al.* Partial-volume correction increases estimated dopamine D2-like receptor binding potential and reduces adult age differences. *J Cereb Blood Flow Metab* **39**, 822–833 (2019).

53. Dukart, J. *et al.* Cerebral blood flow predicts differential neurotransmitter activity. *Sci Rep* **8**, 4074 (2018).

54. Sasaki, T. *et al.* Quantification of Dopamine Transporter in Human Brain Using PET with 18F-FE-PE2I. *Journal of Nuclear Medicine* **53**, 1065–1073 (2012).

55. Gómez, F. J. G., Huertas, I., Ramírez, J. A. L. & Solís, D. G. Elaboración de una plantilla de SPM para la normalización de imágenes de PET con 18F-DOPA. *Imagen Diagnóstica* **9**, 23–25 (2018).

Acknowledgements

Alessia Cavallo, Dr. Till Dembek, Dr. Konstantin Butenko and Dr. Garance Meyer for their support in the preparation of the manuscript.

Author contributions

WJN and MC formulated and conceptualised the experiments. Methods and data analysis were designed by WJN and MC and executed by MC, LW, JV and TM. Visualisations were done by MC, NR, RK and WJN. The project was supervised by WJN. The manuscript was written by MC and WJN.

Funding statement

The study was funded through the Hertie Network of Excellence in Clinical Neuroscience fellowship to WJN. WJN further received funding from the European Union (ERC, ReinforceBG, project 101077060), Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) – Project-ID 424778381 – TRR 295 and the Bundesministerium für Bildung und Forschung (BMBF, project FKZ01GQ1802).

Declaration of interests

WJN received honoraria for consulting from InBrain – Neuroelectronics that is a neurotechnology company and honoraria for talks from Medtronic that is a manufacturer of deep brain stimulation devices unrelated to this manuscript.