# Molecular response to multiple trace element contamination of the European sardine

Anaïs Beauvieux[1], Jean-Marc Fromentin[1], Claire Saraux[2], Diego Romero[3], Nathan Couffin[2,4], Adrien Brown[2,4], Luisa Metral[1], Fabrice Bertile[2,4], Quentin Schull[1]

1 MARBEC, Univ Montpellier, CNRS, Ifremer, IRD, Sète, France

2 Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France

3 Toxicology Department, Faculty of Veterinary Medicine, University of Murcia, 30100 Murcia, Spain

4 Infrastructure Nationale de Protéomique ProFI, FR2048 CNRS CEA, Strasbourg 67087, France

*Corresponding author
Correspondence: anais.beauvieux@gmail.com

**ABSTRACT**

In marine ecosystems, the presence of trace elements resulting from anthropogenic activities has raised concerns regarding their potential effects on marine organisms. This study delves into the intricate relationship between trace element contamination and the physiological responses of a key marine species in the Mediterranean Sea: the European sardine. Since 2008, this species has been experiencing a significant crisis in the region, prompting numerous studies to investigate the potential factors behind the dramatic decline in sardines' size, age, and body condition. However, thorough information on chemical contamination by trace elements and its physiological impact on this species was lacking. We found evidence for the accumulation of multiple elements in sardines, with a light East-West contamination gradient within the Gulf of Lions. While macro-physiological parameters (i.e. body condition) were not affected by contamination, pathways involved in cellular organization and response to stress were clearly upregulated, particularly in the liver, but also in muscle. In addition, a global upregulation in processes linked to the immune system, lipid homeostasis and oxidative stress was recorded in the liver. The associated energetic cost may add a substantial burden to sardines that already face multi-factorial constraints. This study also allows to pinpoint biomarkers of exposure and effects that may be important for monitoring Mediterranean sardine's health. The results of this study and particularly the complex changes in protein expression demonstrate the need for future studies to test the concomitant effects of multiple stressors acting simultaneously, including large scale contamination.

*Keywords:* Cocktail effect, Ecotoxicology, Gulf of Lions, *Sardina pilchardus*, Shotgun proteomic

**Introduction**

The Mediterranean Sea suffers from high levels of pollution, making it one of the most polluted seas in the world (Gabrielides 1995; Danovaro 2003; Sharma et al., 2021; Robledo Ardila et al., 2024). The sea's semi-enclosed nature contributes to the accumulation of pollutants, exacerbated by significant atmospheric inputs such as Saharan dust events, intense human activities (Bucchia et al., 2015; Pedrotti et al., 2016) and riverine outflows on the continental shelves (Durrieu de Madron et al., 2011). This is also intimately linked to the enhanced ability of pelagic food chains to bioaccumulate chemical elements in oligotrophic seas such as the Mediterranean Sea (Harmelin-Vivien et al., 2009; Chouvelon et al., 2018). Among natural and anthropogenic pollutants, trace elements are present in all environmental compartments. Some metals (e.g., copper, iron, zinc) are essential for maintaining the healthy function of the cells of all living organisms, particularly due to their role as components or co-factors of different enzymes (Wood et al., 2011). Nonetheless, such elements can become toxic when they exceed certain concentration thresholds, leading to important detrimental effects on the health of organisms (Plum et al., 2010; Dydak et al., 2011; Asaduzzaman et al., 2017). In addition, non-essential trace elements, such as lead, mercury, cadmium, arsenic, nickel, can cause several types of pathological damage at very low concentration (Friberg et al., 1979; Tchounwou et al., 2012; Jaishankar et al., 2014). The accumulation of these chemicals can influence enzymatic and metabolic activities, disrupting ion homeostasis, reducing growth and adversely affecting body condition and swimming performances (Hollis et al., 1999; Bervoets and Blust 2003; Bervoets et al., 2005). Metals are in fact inducers of oxidative stress in many animals (including fish), causing imbalances between the production of reactive oxygen species (ROS) (e.g. $H_2O_2$, $HO^{\cdot}$, $O_2^{-}$, $R^{\cdot}$, $ROO$) and cell antioxidant activity (Halliwell & Gutteridge, 2015). Increased ROS due to overproduction and/or the inability to destroy them may damage DNA structures and therefore alter the expression patterns of important proteins, hormones, enzymes, etc. (Morcillo et al., 2016; Javed et al., 2017). Metal contamination also modifies the expression of proteins (biomarkers) linked to cellular responses to stress and cell detoxification. The use of biomarkers, such as antioxidant enzymes (Pereira et al., 2013; Williams and Yoshida-Honmachi 2013), heat shock proteins (HSPs, Downs et al., 2006) and metallothioneins (Sakuragui et al., 2013; Williams and Yoshida-Honmachi 2013) to monitor metal pollution has been widely accepted in environmental and ecotoxicology programs.

While experimental data under controlled conditions taught us a lot on the molecular and physiological costs of single contaminants (Isani et al., 2009; Souid et al., 2015; Wang et al., 2013), the challenge that ecologists now face is to understand the effects of trace element mixtures. Indeed, wildlife species are continuously and increasingly exposed to a large number of different chemicals at the same time (Heys et al., 2016). Therefore, the ecotoxicological impacts of trace element mixtures are relatively less understood. Even at low concentrations, trace elements mixed in the environment can have a combined toxicological effect (Heys et al., 2016) forming "reactive chemical cocktails" that explain the synergistic effects of the combination of distinct elements (Kaushal et al., 2018). Considering the effects of only one pollutant at a time can therefore lead to misinterpretation of biomarker data (Celander, 2011).

The emergence of omics tools has provided a powerful means of analyzing complex and integrated responses to contaminants. These tools offer high sensitivity and excellent specificity in studying the molecular changes occurring in organisms (Denslow et al., 2005; Benninghoff 2007). Additionally, high-throughout shotgun proteomics (i.e. the direct and rapid analysis of the entire protein compartment within a complex mixture) provides a unique opportunity to comprehensively examine the expression of thousands of proteins in a specific tissue in a single experiment, using mass spectrometry and bioinformatics techniques instead of traditional biochemical methods (Sanchez et al., 2011; López-Pedrouso et al., 2020). This advance considerably improves our understanding at the protein level and further paves the way for identifying new biomarkers of exposure and effect, which can then be used to develop enhanced monitoring programs to better assess the impacts of pollutants on marine species (Apraiz et al., 2006; Benninghoff 2007).

We applied a quantitative proteomics approach to the European sardine (*Sardina pilchardus*) in the northwestern Mediterranean Sea. This small pelagic fish population has indeed shown a drastic decrease in size and body condition since 2008 (Van Beveren et al., 2014), resulting in a lower value on the stalls, which in turn has led to the collapse of the French small pelagic fish fisheries, which landed less than 200

tons in 2022 against 15,000 tons in the early 2000s (Van Beveren et al., 2016). The hypotheses of top-
down control through overfishing or natural predation or epizootic diseases have been refuted (Van
Beveren et al., 2016b, 2017, Queiros et al., 2018). The main hypothesis to explain the observed changes is
a modification of the quality and/or quantity of sardine preys (such as copepods, Brosset et al., 2016) due
to multifactorial environmental changes (SST, Upwelling, Stratification, Convection, WeMO, Chla
concentration etc. Feuilloley et al., 2020), thus limiting fish energy resources (bottom-up control, Brosset
et al., 2016; Saraux et al., 2019). However, little is known about a potential impact of contaminants on
Mediterranean sardines, since few ecotoxicological studies have been performed so far on small pelagic
fishes in the Gulf of Lions (but see COSTAS project; Tronczynski et al., 2013 and SUCHIMED campaign;
Bouchoucha 2021). In the Mediterranean Sea, surface-enriched concentrations of multiple metals (Co,
Cu, Ni and Zn), along with their significant negative correlation with salinity, suggest that concentrations
are influenced by the Rhône River plume (e.g. Cossa et al., 2017; Radakovitch et al., 2008). Without a
significant tide, the surface extension and main direction of this plume depend on wind and outflow
forcing conditions (Broche et al., 1998). In addition, given the sardines' considerable swimming ability, it
is uncertain how the spatial distribution of pollutants can overlap that of sardines. Hence, it is crucial to
obtain a thorough knowledge of the contamination pressures exerted on small pelagic species such as
sardines, as already highlighted by the United Nations Environment Programme, UNEP (Baker et al.,
2013).
The present study therefore focused on assessing in-situ trace element contamination in juvenile
European sardines in the Gulf of Lions (NW Mediterranean Sea) and on the impact of contaminant
cocktails on individual health using a proteomic approach. The specific objectives of this work were to: (i)
assess the trace element load in wild juvenile sardines and evaluate the spatial variability of this
contamination; (ii) investigate the impact of trace element mixtures on the proteome of liver (the main
organ implicated in xenobiotic metabolism as well as blood glucose regulation, protein synthesis, bile
production, vitamin and mineral storage, steroid metabolism, and immune function) and red muscle (an
organ with high metabolic capacity contributing to a major part of total muscle respiration rate, see
Teulier et al., 2019), and identify potential new biomarkers for both tissues that could serve as valuable
tools in the long-term monitoring of European sardines' health.
124

## Methods

### Study sites and sampling

We collected 105 juvenile sardines in the Gulf of Lions (northwestern Mediterranean Sea, Figure 1).
Specimens were collected in June–July 2021 during the yearly PELMED pelagic survey conducted by the
French Institute for the Exploitation of the Sea (Bourdeix & Hattab, 1985). Trawl locations were
dependent on the acoustic echoes detected, but covered a wide west-east gradient. Between 3 to 12
sardines per trawl were selected. Sagittal otolith pairs were removed from their otic cavity and read to
estimate age. Only age 0 individuals were selected for further analyses. All individuals were measured
(total length TL, in mm) usig a metal ruler (±0.1 mm) and weighed (total mass M, in g) using a digital
balance (±1 g). Their body condition was calculated using the Le Cren index Kn as estimated by Brosset et
al. (2015), which is commonly used as an indicator of general well-being:

$$Kn = \frac{WW}{0.00607 \times TL^{3.057}}$$

136 where WW is the total wet mass in g and TL the total length in cm.

**Figure 1.** Study area in the Gulf of Lions (northwestern Mediterranean Sea). Pelagic trawling locations for the 105 *S. pilchardus* juveniles in the Gulf are indicated by (★).

Liver and a piece of red muscle were carefully removed to avoid any protein contamination
(gloves, a hairnet and a mask were worn by the experimenter and ceramic tools washed with ethanol 96°
and MiliQ water), immediately frozen in a dry shipper containing liquid nitrogen (~-190 °C, ©CX100
Worthington Industries) on-board and later stored at -80 °C at the lab until proteomics analyses. The
white muscle on both sides of the fish was removed using cleansed ceramic tools to avoid any metal
contamination and stored at -40 °C on-board for the week at sea and then stored at -80° C at the lab until
trace element analyses (one side) and lipid content (other side). The red muscle was cleanly separated
from the white muscle by scraping the lateral skin stripped from the animal. White muscle samples were
taken from the epaxial mass.

**Body condition and reserve lipids**

Fish body condition was further investigated by determining white muscle reserve lipid content. Lipids
were extracted using roughly 0.1 g of muscle in a solvent mixture of chloroform-methanol 2 : 1, v/v
according to the method of Folch et al., (1957) and the content of each class of lipids was measured by
chromatography, using a Iatroscan as detailed in Sardenne et al., (2019). Triacylglycerols (TAG),
diacylglycerols (DAG as TAG precursors) and free fatty acids (FFA) were grouped together as reserve lipids
(Zhol et al., 1995; Tocher 2003; Lloret et al., 2013). The proportions of FFA were also checked to ensure
that the lipids had not been degraded during sample storage. Here, the proportions were 0.02% on
average (only one sample exhibited a concentration in FFA, whereas all the others showed no presence
of FFA), significantly below the 25% limit recommended by Parrish (1988).

**Trace element analysis**

All white muscle samples were lyophilized using a freeze dryer (®Leybold, Cologne, Germany). For
aluminium (Al), arsenic (As), beryllium (Be), bismuth (Bi), cadmium (Cd), chromium (Cr), copper (Cu),
lithium (Li), nickel (Ni), lead (Pb), rubidium (Rb), antimony (Sb), strontium (Sr), titanium (Ti), thallium (Tl)
and zinc (Zn) determination, the samples underwent a pre-treatment process where 0.5 grams of each
muscle sample were subjected to acid digestion using 4 mL of $HNO_3$ (69%) and 1 mL of $H_2O_2$ (33%) in
special Teflon reaction tubes within a microwave digestion system (UltraClave-Microwave Milestone®)
for 20 minutes at 220 °C, and then diluted to 10 mL with double deionized water (Milli-Q). Samples were

analyzed using inductively coupled plasma optical emission spectrometry (ICP-OES, ICAP 6500 Duo,
Thermo) to determine the levels of these elements. The detection limit was 0.001 mg/kg. Each sample
was read in duplicate and averaged. Based on UNE-EN ISO reference 11885, multi-element calibration
standards (SCP Science, in 4% nitric acid) were assembled with different concentrations of inorganic
elements. For calibration, see Romero et al. (2020).
Total mercury (Hg) content was measured using an atomic absorption spectrometer AMA254
Advanced Mercury Analyzer (Leco) without pre-treating or pre-concentrating the samples (wavelength =
253.65 nm, detection limit (DL) = 0.003 µg/g). The recovery rate for reference materials (Mercury ICP
Standard 1000 mg/L Hg, Merck) was above 95%.
176
**Quantitative proteomic analysis**
In the following proteomic approach, we opted to analyze a subset of 29 individuals who exhibited
the most distinct contamination profiles. This selection process involved assigning ranks to sardines
based on the level of each contaminant detected. The ranking ranged from 1 (indicating the lowest
contamination level) to 105 (representing the highest contamination level). The individuals were then
selected based on the total sum of these ranks. Consequently, we identified two groups, one comprising
14 sardines with the lowest contamination levels and the other consisting of 15 sardines with the highest
contamination levels (see Statistical analysis section for further details on individuals'selection).
Proteomic analyses of liver and red muscle were conducted as two separate experiments. The process of
sample preparation, nanoLC-MS/MS analysis, and mass spectrometry data analysis is described in detail
in ESM1. The procedure involved protein extraction and electrophoresis using SDS-PAGE, followed by in-
gel digestion with trypsin (Promega, Madison, WI, USA). The resulting peptides were extracted and
analyzed using a nanoUPLC system (nano-Acquity, Waters, Milford, MA, USA) coupled to a quadrupole-
Orbitrap hybrid mass spectrometer (Q-Exactive HF-X, Thermo Scientific, San Jose, CA, USA), controlled by
XCalibur software (v4.0.27.9; Thermo Fisher Scientific). The Q-Exactive HF-X was operated using a data-
dependent acquisition (DDA) strategy by selecting the Top-20 most intense ions in MS1 for fragmentation
in MS2. MS raw data processing was performed in MaxQuant (v2.0.3.1) (Cox et al., 2014), using
Andromeda algorithm to search peaklists against a protein database containing protein sequences from a
home-made annotation of the genome of  the European sardine (TaxID: 27697; Genbank assembly
accession: GCA_003604335.1; Assembly Name: UP_Spi). Only the proteins identified with at least two
peptides were retained. Protein quantification was performed using unique peptides only via the MaxLFQ
option implemented in MaxQuant (Cox et al., 2014) which is a label-free quantification (LFQ) method that
determines protein abundance based on the intensity of peptide signals. The mass spectrometry
proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol
et al., 2019) partner repository with the dataset identifiers PXD037276 (liver) and PXD037313 (red
muscle). QC-related measurements indicated stable performances of the analysis system all along the
two experiments, with median coefficients of variation (CV) of 0.47% (liver) and 0.56% (red muscle) for
retention times of iRT peptides over all injections. Median CVs of only 10.9% (liver) and 13.8% (red
muscle) were obtained for LFQ values of the proteins quantified from the repeated injections of
reference samples.
207
**Protein functional annotation**
To analyse our proteomic data from a functional point of view, we examined the functional
annotations of proteins listed by the AMIGO consortium (Gene Ontology, GO; http://geneontology.org/).
However, only one sardine protein is annotated in GO databases. We therefore took advantage of the
crucial evolutionary position of the spotted gar (*Lepisosteus oculatus*) between teleost fishes and
humans, which creates a very valuable bridge between their genomes (Braasch et al., 2016), and of the
fact that gar proteins are well annotated in GO databases.  We began by searching *L. oculatus* sequences
(UniprotKB, 22,463 protein entries; September 2022) for proteins homologous to *S. pilchardus* proteins
using       BLAST       searches       (FASTA       v36.1.4       program;       downloaded       from
http://fasta.bioch.virginia.edu/fasta_www2/fasta_down.shtml), and only the top BLAST hit for each
protein was retained. From *L. oculatus* protein sequences, the same strategy was employed to search for
human homologous proteins (TaxID: 9606; Reference proteome accesed in September 2022). The
relevance of the match between homologous proteins was checked manually and only 57 of 2505 and 22

of 1112 matches (liver and red muscle, respectively) could not be validated. An automatic extraction of
GO annotations was then performed using the MSDA software (Carapito et al., 2014).
223
**Statistical analyses**
All statistical analyses were performed with the statistical open source R software v.4.1.1 (R Core
Team, 2021). Element concentrations below Limit Of Detection (LOD, 0.001 mg/kg) were set to "LOD/$\sqrt{2}$"
as suggested by Verbovsek (2011) as the best substitution method.
Analyses comparing inorganic contamination patterns included only metals that were detected in at
least 50% of individuals out of the 105 collected. In order to consider possible relationships between
contamination and fish body size, we used a linear model (LM) between each contaminant concentration
and the individual total length. Normality and homoscedasticity of the residuals were verified using the
Shapiro's and Levene's tests, respectively. When distribution of the residuals did not follow a normal
distribution, a log transformation was applied prior to model fitting. If necessary (P < 0.05), size-corrected
contaminant concentration was defined as back-transformed residuals of the regression model.
To investigate relationships between contaminants and to evaluate a possible influence of the Rhône
River on trace element loads in sardines within the Gulf of Lions, we performed a Principal Component
Analysis (PCA) on scaled size-corrected contaminant concentrations of the 105 individuals and projected
distance from the Rhône River mouth on the PCA. We also assessed Spearman correlation between the
three first principal components (PCs) and the distance of the individuals from the Rhône River delta. To
evaluate a potential effect of inorganic contamination on individual's condition, we also performed LMs
between the first three PCs and body condition and reserve lipids.
A second PCA approach was performed using only the 29 individuals selected for proteomic analysis
as objects, while the descriptors were scaled size-corrected contaminant concentrations. The aim of this
analysis was to summarize all the information to investigate the relationships between contaminants and
individual proteomes. PCA axes (PC1, PC2, PC3) were retained as principal components reflecting
contaminant mixture 1, mixture 2, mixture 3 and used in further analyses.
Prior to statistical analysis, the proteomic datasets (liver and red muscle) were filtered to keep only
proteins that were expressed in at least 70% of individuals in each contamination group (common
threshold used in existing literature (Quque et al., 2023; Wang et al., 2020) leaving 2115 (out of 2505)
and 913 (out of 1112) expressed proteins in the liver and red muscle, respectively. Three and one
individuals were removed due to a high number of missing values for liver and muscle proteome analysis,
respectively (supported by the missing value heatmap and proteomic clustering isolation of these
individuals). Remaining missing values were imputed for each contamination group based on random
forest method (Jin et al., 2021; Kokla et al., 2019). Previous studies have shown that this method exhibits
strong performance and is particularly well-suited for label-free proteomic studies, where the underlying
reasons for missing data are not fully understood (Jin et al., 2021; Kokla et al., 2019). A log 2
transformation was used to normalize the protein intensities.
258
**Co-expression network and enrichment analysis**
*Gene co-expression network and hubproteins identification*
Weighted gene co-expression network analysis implemented in the WGCNA R package (Langfelder &
Horvath, 2008) was used for both tissues (liver and red muscle) to identify groups, or modules, of
proteins whose expression significantly correlated with inorganic contamination in white muscle. First, an
adjacency matrix for all pairs of proteins was constructed using the Spearman correlation raised to the
power (beta) of seven and six to approximate a scale-free network for liver and red muscle, respectively
(Langfelder & Horvath, 2008). The adjacency matrix was then transformed into a topological overlap
dissimilarity matrix and a combination of hierarchical clustering and a dynamic tree-cutting algorithm
were used to first define and then merge co-expressed modules of proteins. Proteins outside any module
(indicating low co-expression) were gathered in a grey module. The module's eigengene (i.e. first module
eigenvector: first axis of a principal component analysis conducted on the expression of all module
proteins) summarizes the expression of all proteins in that module. Then, we investigated whether the
eigengene of the modules was correlated with each contaminant mixture (identified through the PCA
analysis, see above), using Spearman correlation. Modules with a significant (P < 0.05) correlation and at
least R ≥ 0.4 were retained for further analysis (only the grey module, indicating low co-expression, for

muscle presented a significant correlation below 0.4). Positive and negative relationships reflect the up-
and down-regulation of the module with an increasing concentration of a contaminant mixture,
respectively. To explore the functional and physiological mechanisms associated with each module, we
then performed functional enrichment analyses.
279
*Functional enrichment analysis and pathway network*
For both tissues, enrichment analysis (biological process and cellular component) of the proteins
constituting each module was performed using Fisher's exact test using the '*GO_MWU*' package
particularly suitable for non-model organisms (https://github.com/z0on/GO_MWU) (Dixon et al., 2015).
The background used for each tissue consisted of the set of proteins identified in that specific tissue,
ensuring two distinct backgrounds for the analysis. This approach prevents biased enrichment from a
generic background, ensuring accurate determination of the functional significance of the identified
modules (Wijesooriya et al., 2022). Enriched GO terms are displayed in a dendrogram plot with distances
between terms reflecting the number of shared proteins. Additionally, we assessed the Spearman
correlation between each protein and the contaminant mixtures (PCA axes) to infer "gene significance"
(GS). Focusing on the top 10 enriched biological process terms (GO terms), these GS relationships were
compiled within a circle plot for each enriched pathway of the module. These plots highlight the up- or
down-regulation of the proteins and pathways in response to the contaminant mixtures.
293
294    *Network visualization*
295    ClueGO was used to illustrate overrepresented Gene Ontology terms. ClueGO is a Cytoscape plug-in
296    that visualizes the non-redundant biological terms for large numbers of proteins and integrates the GO
297    terms to create a GO/pathway network (Bindea et al., 2009). A network for each tissue was made in
298    order to visualize the pathways impacted by the three contaminant mixtures.
299
*Hubproteins identification*
For each module, GS was correlated with module membership (MM; Spearman correlation of protein
intensity versus the module eigengene) to identify proteins showing the highest degree of connectivity
within a module and with the considered contaminant mixture (hubproteins). The top 5 proteins with
simultaneously the highest MM and GS (rank sum) were selected as hubproteins in each module. Due to
their central position in the network, hubproteins are expected to play important biological roles within
their module and are considered as potential biomarkers.
307
*Functional comparison between tissues*
For each contamination mixture, the similarity of the functional response between tissues was
compared by plotting their respective GO term delta ranks against each other. GO term delta ranks
correspond to the difference between the mean ranks based on the GS of the proteins the GO term
contains *vs* the mean ranks of all the protein not included. Positive and negative delta ranks indicate that
the GO term tends to be regulated upwards or downwards, respectively. For a given pathway, the
strength of the relationship reflects the similarity of the enrichment between liver and red muscle. It is
important to note that these plots do not represent a formal statistical test, as the data points (gene
ontology categories) are not independent. Indeed, they often encompass overlapping sets of proteins,
nonetheless it allows identifying functional similarity or dissimilarity between functional enrichments.
318

319                                              **Results**

**Trace element burden across Gulf of Lions**
Thirteen trace elements (Al, As, Cu, Cr, Hg, Li, Ni, Pb, Rb, Sr, Ti, Tl and Zn, ESM2) were retained after
selection of contaminants detected (> LOD) in at least 50% of individuals (i.e. 53 individuals). Only three
elements (Pb, Rb and Tl) exhibited a significant relationship with fish length and were size-corrected.

While Tl displayed a negative relationship with length ($R^2$ = 0.16, P < 0.001), Pb and Rb had a positive
association ($R^2$ = 0.10, P = 0.009 and $R^2$ = 0.09, P = 0.001 respectively).
Considering the 105 sardines sampled across the Gulf of Lions within the same PCA, the first three
principal components explained 24.2%, 15.3% and 12.8% of the variance, respectively (around 52% in
total, ESM3a and Figure 2). The main contributing variables for the first axis (PC1) were Zn, Sr, Rb, Cu, Pb
and As, which all displayed positive contributions (ESM3b and Figure 2). Ni, Cr were positively related to
PC2 while, to a lesser extent, Tl was negatively loaded (ESM3c and Figure 2). Ti and Al were negatively
loaded along the third axis in opposition to Cr, Ni, Tl and Li (ESM3d and Figure 2). Highlighting the
distance of the various sampling sites from the mouth of the Rhône River within the same PCA showed a
slight east-west contamination gradient (Figure 2). PC1 and PC2 were negatively correlated with the
distance from the Rhône river, albeit slightly above significance levels (ESM4). Nonetheless, there was
considerable inter-individual variability in trace element contamination, which probably blurred this east-
west gradient (ESM4). The three first PCs displayed no significant correlation with individuals' body
condition (P = 0.47, P = 0.18 and P = 0.57 for PC1, PC2 and PC3, respectively), nor with the reserve lipid
content (P = 0.55, P = 0.66 and P = 0.79 for PC1, PC2 and PC3, respectively).

**Figure 2.** Biplot of PC1 vs PC2 (a) and PC1 vs PC3 (b) of the PCA built with size-corrected level of trace elements present in the 105 individuals sampled across the Gulf of Lionss. Each point represents an individual. Colour gradient indicates the distance from the Rhône river mouth.

Regarding the second PCA performed on the 29 individuals selected for proteomic analysis, the first
three principal components explained 33.3%, 16.9% and 13.6% of the total variance, respectively (around
64% in total, Figure 3 and ESM5). All 13 contaminants loaded positively on PC1 with 8 of them
contributing strongly. Further, PC1 clearly separated the 2 groups of individuals previously selected (high
contamination *vs* low contamination). Therefore, PC1 was used as a proxy for general inorganic
contamination level, which allows to work along a contamination gradient instead of categorical
contaminations. PC2 was mostly determined by a strong positive contribution of Ni and Cr, while PC3 was
determined by a strong positive contribution of Ti and Al and marginally by a negative contribution of Li.

The three PCs independently identified contaminant mixtures, which are listed in Table 1. We next used
proteomics to investigate the potential impact of these three contaminant mixtures on the health of the
Mediterranean sardine population.

**Figure 3**. Biplot of PC1 vs PC2 (a) and PC1 vs PC3 (b) of the PCA built with level of inorganic contaminants present in the 29 individuals selected for proteomic analysis. Each point represents an individual. Colours indicate contamination level of individuals (HI: High contamination and LI: Low contamination). The larger circles represent the barycenter of the individuals for a given group.

353 **Table 1.** Main variables that contribute to the first three principal components of the PCA on the 29 individuals selected for
354 proteomic analysis. Each PC represents a contaminant mixture.

| | Mixture 1 - PC1 | | Mixture 2 - PC2 | | Mixture 3 - PC3 | |
|---|---|---|---|---|---|---|
| | **-** | **+** | **-** | **+** | **-** | **+** |
| | none | Cu, Tl, Rb, Sr, Pb, Li, Zn, As | none | Ni, Cr | Li | Ti, Al |

**Physiological response to inorganic contamination**

Using a co-expression network analysis, we focused on the impact of metal contamination at the
proteome level on two key organs in fish (the liver and red muscle).

*Liver proteome*

We detected 9 distinct modules (WGCNA analysis) containing a total of 2115 proteins. We then
investigated the relationship between expression modules and the three metal mixtures to assess the
potential health effects of inorganic contaminant exposure. Both mixtures 1 and 3 displayed significant
positive correlations (P < 0.05, encompassing approximately 31% of the detected proteins) with the same

two co-expressed modules (turquoise and yellow, Figure 4), reflecting a general overexpression of
proteins within these 2 modules when mixtures 1 and 3 increased in concentration. Mixture 2 did not
display any relationship with any of the co-expressed modules.

**Figure 4.** Correlations between module eigengenes (columns) and trace element mixtures (rows) for liver proteome. Values in parentheses with module names indicate the number of proteins belonging to each module. Values in the cells are Spearman's correlation coefficients. Blank cells indicate non-significant correlation (P>0.05). *: P < 0.05, **: P < 0.01

The positive correlation of the turquoise module with mixture 1 (PC1, R = 0.52, P < 0.01) and mixture
3 (PC3, R = 0.46, P < 0.05) involved an overall upregulation of 170 GO terms (FDR < 0.01), mainly involved
in anatomical structure and development (e.g., epithelial cell development (GO:0002064), regulation of
anatomical structure size (GO:0090066), response to stimulus/stress (e.g. defense response
(GO:0006952), cellular response to DNA damage stimulus (GO:0006974), protein metabolism (e.g.
positive regulation of gene expression (GO:0010628), regulation of protein polymerization, transport and
cellular organization (e.g. cell development, regulation of cytoskeleton organization and cell cycle process
(spindle organization, cell cycle process, Figure 5 and ESM6a). Proteins from the turquoise module were
part of multiple subcellular compartments, such as the nuclear periphery, endocytic vesicles,
immunological synapses and the cytoskeleton (ESM6b). The positive correlation of the yellow module
with mixture 1 (P < 0.001) displayed a significant enrichment in 71 pathways mainly related to the
immune system, the metabolism of lipids and lipoproteins and transport (Figure 5 and ESM7a). Most of
these pathways were upregulated in response to an increase in mixture 1 (i.e. Cu, Rb, Sr, Tl, Zn, Li, Pb and
As). As for the cellular component analysis, proteins were mainly located in lumens (vesicle, endoplasmic,
endocytic) and were subunits of protein complexes (protein-lipid complex, hemoglobin complex, ESM7b).

**Figure 5.** Significant GO terms (FDR < 0.001) and ontological relationships in the liver. Colored circle size reflects the level of FDR-adjusted statistical significance. Terms enriched in the same module were grouped and presented in the same color. Each leading term, which has the highest significance, is indicated by bold font. Biological processes were grouped in larger pathways (large grey circles).

To identify potential biomarkers, hubproteins (top 5 proteins with the highest GS and MM) were
selected for each module (see Table 2). For the turquoise module, hubproteins were mainly implicated in
protein or lipid transport (Table 2). Interestingly, one of them, the serine/threonine-protein kinase mTOR,
was involved in multiple pathways such as cell development, cell structure and metabolism. Potential
biomarkers for the yellow module included 2 apolipoproteins with one presenting antioxidant activity
(i.e., the Apolipoprotein A-IV). The three other hubproteins were involved in diverse pathways such as
protein synthesis with the eukaryotic translation initiation factor 2 subunit 3, cytoskeleton and protein
transport with the spectrin beta chain and the DnaJ homolog subfamily C member 13 which is not only
implicated in protein transport but also exhibits chaperon function.

**Table 2.** Top 5 hubproteins selected for each co-expression module in the liver proteome. The name and UniprotKB ID are indicated for each of them as well as their main functions, their module, their expression pattern (up or down-regulation), contaminant mixture they correlate with and the module membership

| Name | UniprotKB ID | Module | Main function | Expression pattern | Principal Component | Module membership |
|---|---|---|---|---|---|---|
| Apolipoprotein B-100 isoform 3 | APOB | turquoise | Lipid transport and metabolism | ↑ up | PC1 and PC3 | 0.83 |
| Serine/threonine-protein kinase mTOR | MTOR | turquoise | Cell growth, Cell cycle progression, Metabolism, Cellular structure | ↑ up | PC1 and PC3 | 0.81 |
| Cytoplasmic dynein 1 heavy chain 1 isoform 5 | DYNC1H1 | turquoise | Intracellular motility: cell cycle and transport | ↑ up | PC1 and PC3 | 0.78 |
| U5 small nuclear ribonucleoprotein 200 kDa helicase | SNRNP200 | turquoise | mRNA processing/splicing | ↑ up | PC1 and PC3 | 0.89 |
| Cytoplasmic dynein 1 heavy chain 1 isoform 2 | DYNC1H1 | turquoise | Intracellular motility: cell cycle and transport | ↑ up | PC1 and PC3 | 0.80 |
| Apolipoprotein A-IV | APOA4 | yellow | Lipid transport and metabolism, Antioxidant capacity | ↑ up | PC1 and PC3 | 0.88 |
| Apolipoprotein B-100 isoform 1 | APOB | yellow | Lipid transport and metabolism | ↑ up | PC1 and PC3 | 0.79 |
| Eukaryotic translation initiation factor 2 subunit 3 | EIF2S3 | yellow | Translation | ↓ down | PC1 and PC3 | -0.77 |
| Spectrin beta chain, non-erythrocytic 1 | SPTBN1 | yellow | Cellular structure, Protein transport/localization | ↑ up | PC1 and PC3 | 0.84 |
| DnaJ homolog subfamily C member 13 | DNAJC13 | yellow | Chaperone, Protein transport | ↑ up | PC1 and PC3 | 0.88 |

392 *Muscle proteome*

393      The WGCNA analysis of the red muscle proteome, containing 913 proteins, resulted in four modules

394 namely blue (372 proteins), brown (119 proteins), turquoise (398 proteins), and grey (24 proteins).

**Figure 6.** Correlations between module eigengenes (columns) and trace element mixtures (rows) for the red muscle proteome. Values in parentheses with module names indicate the number of proteins belonging to each module. Values in the cells are Spearman's correlation coefficients. Blank cells indicate non-significant correlation (P > 0.05). *: P < 0.05, **: P < 0.01

395      Only the blue module, which accounts for 41% of all detected proteins in the red muscle proteome,

396 correlated (positively) with one contaminant mixture (i.e., mixture 1, Figure 6). This module was enriched

397 in cellular organization and ion transport pathways (Figure 7 and ESM8). According to the cellular

398 component enrichment analysis, proteins were mainly located in membranes (ESM8b).

**Figure 7.** Significant GO terms (FDR < 0.001) and ontological relationships in red muscle. Circle size indicates the level of FDR-adjusted statistical significance. Biological processes were grouped in larger pathways (large grey circles).

399   Two identified hubproteins for the blue module have a known function in the response to oxidative
400 stress (Gluthatione S-transferase and Hemoglobin subunit zeta), whereas two others are involved in
401 cellular structure and muscle contraction (myosin-binding protein H and Myomesin-2). The last identified
402 hubprotein was the ADP/ATP translocase 1 which is implicated in energy metabolism and apoptosic
403 signaling pathway (Table 3).

**Table 3.** Top 5 hubproteins selected for each co-expression module in the red muscle proteome. The name and UniprotKB ID are indicated for each of them as well as their main functions, their module, their expression pattern (up or down-regulation), contaminant mixture they correlate with and the module membership.

| Name | UniprotKB ID | Module | Main function | Expression pattern | Principal Component | Module membership |
|---|---|---|---|---|---|---|
| Glutathione S-transferase | LANCL1 | blue | Contaminant detoxification, Oxidative stress metabolism | ↓ down | PC1 | -0.92 |
| Hemoglobin subunit zeta | HBZ | blue | Gas transport, Oxidative stress metabolism | ↑ up | PC1 | 0.90 |
| Myosin-binding protein H | MYBPH | blue | Cell adhesion, Muscle contraction | ↓ down | PC1 | -0.92 |
| ADP/ATP translocase 1 | SLC25A4 | blue | Transmembrane transporter, Apoptosic signalling pathway | ↓ down | PC1 | -0.92 |
| Myomesin-2 | MYOM2 | blue | Cellular organization, Muscle contraction | ↓ down | PC1 | -0.91 |

405 *Comparison of liver vs red muscle proteome response*

406   Comparison of the delta-rank values obtained from GO enrichment analysis for both tissues against
407 mixture 1 revealed a significant positive correlation (R = 0.22, P < 0.001 ESM9a). Liver and muscle
408 enrichment for mixture 2 displayed no significant correlation (R = 0.02, P = 0.96, ESM9b), while
409 enrichment for mixture 3 revealed a significant but weak positive correlation between the two tissues (R

410     = 0.06, P < 0.01, ESM9c). This highlights that similar processes appear involved in liver and red muscle in
411     response to contaminant mixture 1 and, to a lesser extent, mixture 3.

## Discussion

Multiple studies have investigated the potential drivers of the drastic decrease in size, age and body
condition of sardines in the Gulf of Lions since 2008 (e.g. Brosset et al., 2015, 2016, 2017; Le Bourg et al.,
2015; Van Beveren et al., 2016, 2017; Feuilloley et al., 2020). However, thorough information on chemical
contamination by trace elements and its physiological impact on this species was lacking. We found
evidence for the accumulation of multiple elements in sardines, with a light east-west contamination
gradient within the Gulf of Lions. While macro-physiological parameters (i.e. body condition) were not
affected by contamination, pathways involved in cellular organization and response to stress were clearly
upregulated, particularly in the liver, but also in muscle. In addition, a global upregulation in processes
linked to the immune system and lipid homeostasis was recorded in the liver. We are proposing
biomarkers of the effects of trace element stress on health, which could serve as a starting point in
biomonitoring programs.
424
**Contamination load in the Mediterranean sardine of the Gulf of Lions**
The impact of contaminating loads from the Rhône River on the organisms living in the Gulf of Lions
has already been reported with regard to radionuclides, such as $^{210}$Po (Strady et al., 2015), heavy metals
(Cd, Pb and Hg for [60–200 μm] size fraction, Chouvelon et al., 2019) and organic contaminants, such as
Polychlorinated biphenyls (PCBs) in plankton (Alekseenko et al., 2018). In addition to the Rhône River
inputs, pollutant discharges are also likely to come from industrial and urban activities of the nearby
cities of Marseille and Fos-sur-Mer. Small pelagic planktivorous fish such as sardines in the eastern areas
of the Gulf of Lions might therefore be highly exposed to chemical contamination, notably via trophic
pathways (i.e. plankton). Interestingly, although large inter-individual variability in contamination was
recorded in our study, we noted a moderate east to west contamination gradient in the Gulf of Lions.
These results differ from previous work (Chouvelon et al., 2019) that did not detect a spatial pattern in
trace element contamination in the same species and location. These differences might be due to the
targeted tissue, as they considered contamination in the whole organism (liver, muscle, gonad and
remaining tissues) whereas we focused on muscle contamination. Indeed, factors such as tissue-specific
metabolic processes, affinity for contaminants and turnover rates can contribute to distinct
contamination patterns observed across differents tissues. Another reason for discrepancies between our
and previous studies might involve the fact that individuals were sampled at different life stages in those
former studies, whereas we focused on individuals in their first year of life (age 0). Indeed, confounding
effects with contaminant elimination through reproduction as well as bioaccumulation with age may
explain large variations and no clear spatial pattern (Bodiguel et al., 2009). Canli and Atli (2003)
highlighted the importance of fish size in determining the rate of physiological processes, as size may
influence the uptake, distribution, and elimination of trace elements. In our study, only three trace
elements (out of 13) displayed significant relationships with fish size, namely Pb, Rb and Tl. This lack of
relationship with size for most of trace elements could result from the complexity of bioaccumulation
processes involving interaction between various routes of absorption, excretion, passive release, and
metabolization (Joiris et al., 1999; Guo et al., 2016; Tang et al., 2017). Yet, the study of one-year-old
gilthead seabream juveniles (Beauvieux et al., 2024) showed comparable outcomes. This indicates that
when examining a single age group, size differences are likely minimal, simplifying the study of spatial
contamination. That said, exploring multiple age groups remains essential to uncover intricate
bioaccumulation patterns.
Trace metal concentrations in Mediterranean surface waters and top predators are generally higher
than in the Atlantic Ocean (Boyle et al., 1985; Cossa and Coquery 2005). This phenomenon can be largely
attributed to the Mediterranean's semi-enclosed geography, which promotes pollutant accumulation.
This accumulation is further exacerbated by significant atmospheric inputs, such as Saharan dust events,
intense human activities (Bucchia et al., 2015; Pedrotti et al., 2016), and riverine discharges on
continental shelves (Durrieu de Madron et al., 2011). Additionally, there is evidence suggesting that the
Mediterranean pelagic food webs may have a heightened capacity to bioaccumulate certain trace

elements, particularly Hg (Cossa and Coquery 2005; Harmelin-Vivien et al., 2009; Chouvelon et al., 2018).
This pattern also turned out to be true for As, Ni, Cr, Pb, Zn and Hg in sardines recorded in numerous
studies (considering only the European sardine in ESM10). Trace element concentrations found in the
present study were of a similar order of magnitude to those found in other studies focusing on the same
or close species of the Mediterranean, Atlantic or Indian Oceans, except for Li. Averaged Li levels were 32
times higher than those found by Guérin et al., (2011) on the Mediterranean sardine (but Li levels were
measured in only 4 individuals in this previous work) and 2 times higher than those found by Lozano-
Bilbao et al., (2019) on the European sardine in the Atlantic Ocean.
Similarly to findings in seabreams (Beauvieux et al., 2024), we identified a lack of monitoring for
several elements including Li, Rb, Sr, Ti and Tl in all locations (Indian and Atlantic Oceans and the
Mediterranean Sea, ESM10 and ESM11). This is partially explained by the absence of regulatory limits for
these elements in fish muscle (along with Cr and Ni), despite their increasing use in multiple areas
(industrial, agriculture, high-tech and medication industries) and despite their toxicity for organisms
(Exley et al., 1991; Authman 2011; Aslam and Yousafzai 2017; Blewett and Leonard 2017; Kumari et al.,
2017; Genchi et al., 2021).
The European regulation sets contamination thresholds for only three specific inorganic trace
elements, Pb, Cd and Hg. In contrast, other nations, such as Turkey and Greece have adopted thresholds
for a larger set of contaminants (for a comprehensive review, refer to Guerra-García et al., 2023).
Notably, trace element levels in sardine specimens from the present study remained below the threshold
values for fish consumption set by most international regulations except for As that exceeded these
limits. These findings emphasize the need for further efforts by the European commission and other
institutions to establish health limits not only for Arsenic but also for other nonessential and essential
elements.
Furthermore, it is important to consider the existence of organic and emerging pollutants (POPs,
HAPs, PFOS, PFAS, etc.) that are becoming increasingly pervasive worldwide (Magulova & Priceputu,
2016). Given their persistence, long-range transportability, biomagnification in food chains, and
bioaccumulation in humans and wildlife, their impact on individual health is becoming a growing concern
(Magulova & Priceputu, 2016). The combined effects of trace elements and POPs create a complex
chemical cocktail that can have synergistic or antagonistic impacts on marine life. Thus, there is a growing
need for additional research to fully understand the scope of contamination which requires considering
both trace elements and POPs.
493
**Commonly dysregulated pathways in the liver and red muscle in response to multiple-metal**
**contamination**
Liver and red muscle shared common proteome changes, such as cellular disorganization as well as
functions involved in transport and stress response (Figure 5 and 7). This is in line with the positive
correlation found between the functional enrichment of the two tissues in relation to mixture 1 and
mixture 3 (ESM9), highlighting similar responses (upregulation). However, we noted a much more diverse
response in the liver, with 17 times more pathways overexpressed compared to red muscle. Similar
results have been observed in seabreams (Beauvieux et al., 2024). This discrepancy in responses between
the two tissues could be related to the liver functions, which is the main organ responsible for
detoxification, transformation and storage of toxic compounds in fish (Bawuro et al., 2018; Beauvieux et
al., 2024).
Contaminant mixture 1 (Cu, Rb, Sr, Tl, Zn, Li, Pb and As) seemed to particularly impact cellular
organization in the two tissues, as reflected by an overexpression of multiple pathways involved in
cellular structure, intracellular transport and motility processes. In particular, two isoforms of cytoplasmic
dynein heavy chain seemed to be affected by mixture 1 in liver (along with mixture 3). Cytoplasmic
dyneins are central in intracellular trafficking, including macromolecular transport, chromosome
dynamics, and the cell cycle (Paschal et al., 1987; Kardon and Vale 2009; Bawuro et al., 2018). This is in
line with the significant downregulation in the pathway involved in mitotic spindle assembly that has
been also detected and could indicate that trace element exposure has genotoxic effects and may
repress hepatocyte division. Similarly, cytoplasmic dyneins (Table 2) were identified in bivalves as
responsive to contamination (Li & Wang, 2021; Sánchez-Marín et al., 2021), which made them great
candidate biomarkers of trace element contamination. Additionally, given that proper cellular

organization is crucial for organismal development, the abundance of upregulated pathways associated with anatomical structure and development may suggest a response to counteract contamination-induced disorganization. Histopathological alterations in hepatocytes have already been observed, revealing the disorganization of organelles within the cytoplasm (Moreira et al., 2003) and the presence of necrosis (Javed et al., 2017) in response to metal contamination. Since hepatocytes are involved in various aspects of intermediate metabolism, unfavourable consequences to physiology may be expected as well at the cellular and organismal level (Mela et al., 2007).

Red muscle proteome response to mixture 1 displayed fewer functional diversity response than that of the liver proteome, but it reflected an alteration of the biological structure of cells and, again, the cell cycle process, as highlighted by the upregulation of the pathway "mitotic cell cycle process". Accordingly, two of the five hubproteins identified were myosin and myomesin, which are key proteins in cell adhesion and muscle contraction. This is consistent with previous works that highlighted alterations of multiple myocytoskeleton proteins in response to contamination (Rodríguez-Ortega et al., 2003; Karim et al., 2011; Xu et al., 2019). As the contractile ability of myocytes resides in their highly organized cytoskeleton network, dysregulation here could greatly affect the locomotion of juvenile sardines, hindering their escape from predators and their foraging success.

The effects of trace element contamination on liver and muscle cytoskeleton are likely to be caused by oxidative stress (Epel et al., 2004), either through excess reactive oxygen species (ROS) generation (by redox-active metals) or reduction of antioxidant abundances and/or activities (redox-inactive metals) (Koivula & Eeva, 2010). In this study, we highlighted the upregulation of pathways such as "cellular oxidant detoxification" and "hydrogen peroxide catabolic process" in the liver and "reactive oxygen species metabolic process" in muscle. Consequently, two potential biomarkers of red muscle proteome in response to mixture 1 were identified (down-regulated gluthathione S-transferase and up-regulated hemoglobin subunit zeta, Table 3). They have both known functions in contaminant detoxification and antioxidant defences, respectively (Mollan et al., 2012). Gluthathione S-transferase was underexpressed in red muscle with increasing contamination (mixture 1), suggesting a redox-inactive potential of mixture 1 on this enzyme specifically but also highlighting that some canonical detoxification enzymes may be triggered or repressed by metal exposure, depending on their intrinsic nature (redox-active *vs* redox-inactive sensitivity).

Furthermore, trace element mixtures induced additional cellular responses aimed at safeguarding organisms from ROS-induced damage. Among these responses, a heat shock protein homologue (Dnaj homologue, Table 2) plays a crucial role by acting as cochaperone to the molecular chaperone (Hsp70). Its functions involve protein translation, promoting protein folding, translocation of polypeptides through membranes and degradation of damaged proteins (Mitra et al., 2009). In multicellular eukaryotes, programmed cell death, often referred to as apoptosis, is a prevalent stress response mechanism that comes into play when the level of stress surpasses the cell's ability to uphold genomic and macromolecular integrity. In this study, apoptosis signalling pathway involving ADP/ATP translocase and the serine/threonine-protein kinase mTOR (Table 3 and Table 2) appeared to be affected by trace element contamination. Previous transcriptomics and proteomics studies have reported alterations in apoptotic pathways in various fish species in response to environmental contaminants (Bohne-Kjersem et al., 2009; Leaver et al., 2010; Asker et al., 2013; Yadetie et al., 2013). The serine/threonine-protein kinase mTOR was overexpressed in response to trace element contamination in the liver, which is implicated in cellular death cascades along with lipid regulation and nutrient signaling. However, besides its abundance, post-translational modifications including phosphorylation, ubiquitination, acetylation, and glycosylation appears to be key regulators of mTOR signaling and should also be considered (Yin et al., 2021). The activation of the signalling network mTOR has been associated with organic and inorganic contaminants-induced apoptosis (Chen et al., 2008; McCuaig et al., 2020). Moreover, the ADP/ATP translocase is suggested to play a significant role in the process of mitochondrial permeabilization as it can switch its function from ADP/ATP exchange through inner mitochondrial membrane to pore-formation during apoptosis (Kumarswamy & Chandna, 2009). Therefore, these two proteins could be excellent candidates biomarker for trace element effects in liver in red muscle.

**Metabolism and immune system upregulation in liver**

Trace element contamination (mixtures 1 and 3) appeared to specifically affect the metabolism and
transport of lipids in the liver, as reflected by the overexpression of two apolipoproteins (hubproteins).
Lipids are an energy source and an essential component of cell membranes (phospholipids and
cholesterol). Besides, they also play a significant role as messengers in signal transduction pathways and
molecular recognition processes (Van Meer et al., 2008). Any changes in lipid metabolism would signal
impairment of these pathways. It has been reported that constant energy demand leads to mobilization
of triglycerides as they serve as lipid storage (Van Meer et al., 2008; Santos and Schulze 2012). The
upregulation of triglyceride metabolic process and transport underlined in this study could be an
evidence of their mobilization and use in repair or development of membranes damaged, for instance,
under increasing oxidative stress (Van Meer et al., 2008; Santos and Schulze 2012). Their mobilization and
transport may also be linked to low food intake or absorption due to gut damage or improper synthesis in
the liver under trace element stress. In the absence of further research to shed light on these regulation
patterns, we propose here two promising biomarkers that may indicate a dysregulation in lipid transport
and metabolism in response to contamination: apolipoprotein B-100 and apolipoprotein A-IV (see Table
2) that were found upregulated in our study, but also in seahorses and atlantic cod in response to heavy
metal and organic contaminant stress, respectively (Bohne-Kjersem et al., 2009; Liu et al., 2022). These
proteins are important regulators of triglyceride and cholesterol metabolism (Sacks, 2006). By studying
these potential biomarkers, we can gain valuable insights into the effects of metal contamination on
lipid-related processes.

The sardine liver proteome also presented a clear upregulation in multiple pathways involved in
immune processes. Regoli & Giuliani (2014) and Wang & Gallagher (2013) highlighted that such an
upregulation could be related to trace element accumulation in the liver and lead to increasing oxidative
stress. The fish immune system is known to be sensitive to environmental contamination (Si et al., 2019)
and altering immune homeostasis in fish could significantly constrain their survival and development.
Sustaining high levels of immune functions requires energy, which would be necessarily redirected from
other mains functions, such as growth, reproduction or maintenance (energy trade-off,Moret & Schmid-
Hempel, 2000).

In our experiment we did not find any correlation between body condition and contamination
mixtures. This is consistent with some other studies that investigated the relationship between metal
levels in the environment or fish tissues and body condition (Farag et al., 1998; Dethloff et al., 2001;
Lohner et al., 2001). The measurement of body mass may not sufficiently illustrate the costs of metal
exposure, since individuals can balance their resource budget and adjust their body condition over a
longer period (Daan et al., 1996, Beauvieux et al., 2022). The absence of relationship could also suggest
that body condition, although a great marker of nutritional stress  (Brosset et al., 2015), only declines
drastically when a high threshold of physiological stress is reached. Therefore, monitoring physiological
processes that are more sensitive and reflect progressive changes might allow to detect early warnings of
environmental challenges. Nonetheless, sardines from the Mediterranean Sea have shown a drastic
decline in body condition, size and life expectancy in the past two decades (Saraux et al., 2019). The
major driver proposed is the shift of sardine diet towards smaller planktonic prey observed in the Gulf of
Lions due to multi-factorial environmental changes (Feuillolley et al 2020, Brosset et al., 2016), leading to
lower foraging efficiency and a reallocation trade-off toward reproduction instead of survival (Beauvieux
et al., 2022; Queiros et al., 2019, 2024). Consequently, individuals at the end of the reproduction period
may rely on low energetic reserves to survive the end of the winter period. Adding any additional
energetic or immune burden, as those highlighted in response to contaminants in this study, might
emphasize their reduced chance of survival (Queiros et al., 2021).

## **Acknowledgements**

616    We are grateful to the technical and scientific crews of L'Europe for their work during PELMED survey.
617    Furthermore, we thank Sophie Prud'homme, Roberta Bettinetti and an anonymous reviewer for their
618    valuable inputs and comments on this manuscript.

## Data, scripts, code, and supplementary information availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol et al., 2019) partner repository with the dataset identifiers PXD037276 (liver) and PXD037313 (red muscle).

Scripts, figures 5 and 7 and other datasets are available online: https://doi.org/10.5281/zenodo.10683281

Supplementary materials are available online: https://doi.org/10.1101/2024.02.16.580673

## Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

## Funding

This work is part of the project ICIPOP granted by the "Agence Francaise de Developpement" (AFD). A. B. has a PhD grant from University of Montpellier – GAIA doctoral school.

## References

Alekseenko, E., Thouvenin, B., Tronczyński, J., Carlotti, F., Garreau, P., Tixier, C., & Baklouti, M. (2018). Modeling of PCB trophic transfer in the Gulf of Lions; 3D coupled model application. *Marine Pollution Bulletin*, *128*, 140–155. https://doi.org/10.1016/j.marpolbul.2018.01.008

Apraiz, I., Mi, J., & Cristobal, S. (2006). Identification of proteomic signatures of exposure to marine pollutants in mussels (*Mytilus edulis*). *Molecular and Cellular Proteomics*, *5*(7), 1274–1285. https://doi.org/10.1074/mcp.M500333-MCP200

Asaduzzaman, K., Khandaker, M. U., Binti Baharudin, N. A., Amin, Y. B. M., Farook, M. S., Bradley, D. A., & Mahmoud, O. (2017). Heavy metals in human teeth dentine: A bio-indicator of metals exposure and environmental pollution. *Chemosphere*, *176*, 221–230. https://doi.org/10.1016/j.chemosphere.2017.02.114

Asker, N., Kristiansson, E., Albertsson, E., Larsson, D. G. J., & Förlin, L. (2013). Hepatic transcriptome profiling indicates differential mRNA expression of apoptosis and immune related genes in eelpout (*Zoarces viviparus*) caught at Göteborg harbor, Sweden. *Aquatic Toxicology*, *130–131*, 58–67. https://doi.org/10.1016/j.aquatox.2012.12.017

Aslam, S., & Yousafzai, A. M. (2017). Chromium toxicity in fish: A review article. *~ 1483 ~ Journal of Entomology and Zoology Studies*, *5*(3), 1483–1488.

Authman, M. M. N. (2011). Environmental and experimental studies of aluminium toxicity on the liver of *Oreochromis niloticus* (Linnaeus, 1757) fish. *Life Science Journal*, *8*(4), 764–776.

Baker, E., Sevaldsen, P., & Kurvits, T. (2013). State and pressures of the marine and coastal Mediterranean environment. In *European Environment Agency* (Issue 5).

Bawuro, A. A., Voegborlo, R. B., & Adimado, A. A. (2018). Bioaccumulation of Heavy Metals in Some Tissues of Fish in Lake Geriyo, Adamawa State, Nigeria. *Journal of Environmental and Public Health*, *2018*. https://doi.org/10.1155/2018/1854892

Beauvieux, A., Fromentin, J.-M., Romero, D., Couffin, N., Brown, A., Metral, L., Bourjea, J., Bertile, F., & Schull,
Q. (2024). Molecular fingerprint of gilthead seabream physiology in response to pollutant mixtures in the
wild. *Environmental Pollution*, *340*, 122789. https://doi.org/10.1016/j.envpol.2023.122789

Beauvieux, A., Queiros, Q., Metral, L., Dutto, G., Gasset, E., Criscuolo, F., Fromentin, J. M., Saraux, C., & Schull,
Q. (2022). Energy allocation trade-offs between life-history traits in the Mediterranean sardine: an
ecophysiological approach. *Marine Ecology Progress Series*, *701*(Stearns 1989), 99–118.
https://doi.org/10.3354/meps14183

Benninghoff, A. D. (2007). Toxicoproteomics - The next step in the evolution of environmental biomarkers?
*Toxicological Sciences*, *95*(1), 1–4. https://doi.org/10.1093/toxsci/kfl157

Bervoets, L., & Blust, R. (2003). Metal concentrations in water, sediment and gudgeon (*Gobio gobio*) from a
pollution gradient: Relationship with fish condition factor. *Environmental Pollution*, *126*(1), 9–19.
https://doi.org/10.1016/S0269-7491(03)00173-8

Bervoets, L., Knaepkens, G., Eens, M., & Blust, R. (2005). Fish community responses to metal pollution.
*Environmental Pollution*, *138*(2), 338–349. https://doi.org/10.1016/j.envpol.2005.03.005

Bindea, G., Mlecnik, B., Hackl, H., Charoentong, P., Tosolini, M., Kirilovsky, A., Fridman, W.-H., Pagès, F.,
Trajanoski, Z., & Galon, J. (2009). ClueGO: a Cytoscape plug-in to decipher functionally grouped gene
ontology and pathway annotation networks. *Bioinformatics*, *25*(8), 1091–1093.
https://doi.org/10.1093/bioinformatics/btp101

Blewett, T. A., & Leonard, E. M. (2017). Mechanisms of nickel toxicity to fish and invertebrates in marine and
estuarine waters. *Environmental Pollution*, *223*, 311–322. https://doi.org/10.1016/j.envpol.2017.01.028

Bodiguel, X., Loizeau, V., Le Guellec, A. M., Roupsard, F., Philippon, X., & Mellon-Duval, C. (2009). Influence of
sex, maturity and reproduction on PCB and p,p'DDE concentrations and repartitions in the European
hake (*Merluccius merluccius*, L.) from the Gulf of Lions (N.W. Mediterranean). *Science of the Total
Environment*, *408*(2), 304–311. https://doi.org/10.1016/j.scitotenv.2009.10.004

Bohne-Kjersem, A., Skadsheim, A., Goksøyr, A., & Grøsvik, B. E. (2009). Candidate biomarker discovery in
plasma of juvenile cod (*Gadus morhua*) exposed to crude North Sea oil, alkyl phenols and polycyclic
aromatic hydrocarbons (PAHs). *Marine Environmental Research*, *68*(5), 268–277.
https://doi.org/10.1016/j.marenvres.2009.06.016

Bouchoucha, M. (2021). *SUCHI Med 2021*. https://doi.org/https://doi.org/10.17600/18001619

Bourdeix, J., & Hattab, T. (1985). *PELMED - PELAGIQUES MEDITERRANÉE*. https://doi.org/10.18142/19

Boyle, E. A., Chapnick, S. D., Bai, X. X., & Spivack, A. (1985). Trace metal enrichments in the Mediterranean Sea.
*Earth and Planetary Science Letters*, *74*(4), 405–419. https://doi.org/10.1016/S0012-821X(85)80011-X

Braasch, I., Gehrke, A. R., Smith, J. J., Kawasaki, K., Manousaki, T., Pasquier, J., Amores, A., Desvignes, T.,
Batzel, P., Catchen, J., Berlin, A. M., Campbell, M. S., Barrell, D., Martin, K. J., Mulley, J. F., Ravi, V., Lee, A.
P., Nakamura, T., Chalopin, D., … Postlethwait, J. H. (2016). The spotted gar genome illuminates
vertebrate evolution and facilitates human-teleost comparisons. *Nature Genetics*, *48*(4), 427–437.
https://doi.org/10.1038/ng.3526

Broche, P., Devenon, J. L., Forget, P., De Maistre, J. C., Naudin, J. J., & Cauwet, G. (1998). Experimental study of
the Rhone plume. Part I: Physics and dynamics. *Oceanologica Acta*, *21*(6), 725–738.
https://doi.org/10.1016/S0399-1784(99)80002-4

Brosset, P., Fromentin, J. M., Ménard, F., Pernet, F., Bourdeix, J. H., Bigot, J. L., Van Beveren, E., Pérez Roda, M.
A., Choy, S., & Saraux, C. (2015). Measurement and analysis of small pelagic fish condition: A suitable

method for rapid evaluation in the field. *Journal of Experimental Marine Biology and Ecology*, *462*, 90–97. https://doi.org/10.1016/j.jembe.2014.10.016

Brosset, P., Fromentin, J. M., Van Beveren, E., Lloret, J., Marques, V., Basilone, G., Bonanno, A., Carpi, P., Donato, F., Čikeš Keč, V., De Felice, A., Ferreri, R., Gašparević, D., Giráldez, A., Gücü, A., Iglesias, M., Leonori, I., Palomera, I., Somarakis, S., … Saraux, C. (2017). Spatio-temporal patterns and environmental controls of small pelagic fish body condition from contrasted Mediterranean areas. *Progress in Oceanography*, *151*(February), 149–162. https://doi.org/10.1016/j.pocean.2016.12.002

Brosset, P., Le Bourg, B., Costalago, D., BǍnaru, D., Van Beveren, E., Bourdeix, J. H., Fromentin, J. M., Ménard, F., & Saraux, C. (2016). Linking small pelagic dietary shifts with ecosystem changes in the Gulf of Lions. *Marine Ecology Progress Series*, *554*, 157–171. https://doi.org/10.3354/meps11796

Brosset, P., Ménard, F., Fromentin, J. M., Bonhommeau, S., Ulses, C., Bourdeix, J. H., Bigot, J. L., Van Beveren, E., Roos, D., & Saraux, C. (2015). Influence of environmental variability and age on the body condition of small pelagic fish in the Gulf of Lions. *Marine Ecology Progress Series*, *529*, 219–231. https://doi.org/10.3354/meps11275

Bucchia, M., Camacho, M., Santos, M. R. D., Boada, L. D., Roncada, P., Mateo, R., Ortiz-Santaliestra, M. E., Rodríguez-Estival, J., Zumbado, M., Orós, J., Henríquez-Hernández, L. A., García-Álvarez, N., & Luzardo, O. P. (2015). Plasma levels of pollutants are much higher in loggerhead turtle populations from the Adriatic Sea than in those from open waters (Eastern Atlantic Ocean). *Science of the Total Environment*, *523*, 161–169. https://doi.org/10.1016/j.scitotenv.2015.03.047

Carapito, C., Burel, A., Guterl, P., Walter, A., Varrier, F., Bertile, F., & Van Dorsselaer, A. (2014). MSDA, a proteomics software suite for in-depth Mass Spectrometry Data Analysis using grid computing. *Proteomics*, *14*(9), 1014–1019. https://doi.org/10.1002/pmic.201300415

Celander, M. C. (2011). Cocktail effects on biomarker responses in fish. *Aquatic Toxicology*, *105*(3–4), 72–77. https://doi.org/10.1016/j.aquatox.2011.06.002

Chen, L., Liu, L., Luo, Y., & Huang, S. (2008). MAPK and mTOR pathways are involved in cadmium-induced neuronal apoptosis. *Journal of Neurochemistry*, *105*(1), 251–261. https://doi.org/10.1111/j.1471-4159.2007.05133.x

Chouvelon, T., Cresson, P., Bouchoucha, M., Brach-Papa, C., Bustamante, P., Crochet, S., Marco-Miralles, F., Thomas, B., & Knoery, J. (2018). Oligotrophy as a major driver of mercury bioaccumulation in medium-to high-trophic level consumers: A marine ecosystem-comparative study. *Environmental Pollution*, *233*, 844–854. https://doi.org/10.1016/j.envpol.2017.11.015

Chouvelon, T., Strady, E., Harmelin-Vivien, M., Radakovitch, O., Brach-Papa, C., Crochet, S., Knoery, J., Rozuel, E., Thomas, B., Tronczynski, J., & Chiffoleau, J.-F. (2019). Patterns of trace metal bioaccumulation and trophic transfer in a phytoplankton-zooplankton-small pelagic fish marine food web. *Marine Pollution Bulletin*, *146*, 1013–1030. https://doi.org/10.1016/j.marpolbul.2019.07.047

Cossa, D., & Coquery, M. (2005). *The Mediterranean Mercury Anomaly, a Geochemical or a Biologocal Issue* (Vol. 5, Issue June, pp. 177–208). https://doi.org/10.1007/b107147

Cossa, D., Durrieu de Madron, X., Schäfer, J., Lanceleur, L., Guédron, S., Buscail, R., Thomas, B., Castelle, S., & Naudin, J. J. (2017). The open sea as the main source of methylmercury in the water column of the Gulf of Lions (Northwestern Mediterranean margin). *Geochimica et Cosmochimica Acta*, *199*, 222–237. https://doi.org/10.1016/j.gca.2016.11.037

Danovaro, R. (2003). Pollution threats in the Mediterranean Sea : An overview. *Chemistry and Ecology*, *19*(1), 15–32. https://doi.org/10.1080/0275754031000081467

Denslow, N. D., Knoebl, I., & Larkin, P. (2005). Chapter 3 Approaches in proteomics and genomics for eco-
toxicology. *Biochemistry and Molecular Biology of Fishes*, *6*(C), 85–116. https://doi.org/10.1016/S1873-
0140(05)80006-2

Dethloff, G. M., Bailey, H. C., & Maier, K. J. (2001). Effects of dissolved copper on select hematological,
biochemical, and immunological parameters of wild rainbow trout (*Oncorhynchus mykiss*). *Archives of*
*Environmental Contamination and Toxicology*, *40*(3), 371–380. https://doi.org/10.1007/s002440010185

Dixon, G. B., Davies, S. W., Aglyamova, G. V., Meyer, E., Bay, L. K., & Matz, M. V. (2015). Genomic determinants
of coral heat tolerance across latitudes. *Science*, *348*(6242), 1460–1462.
https://doi.org/10.1126/science.1261224

Downs, C. A., Richmond, R. H., Mendiola, W. J., Rougée, L., & Ostrander, G. K. (2006). Cellular physiological
effects of the MV Kyowa Violet fuel-oil spill on the hard coral, *Porites lobata*. *Environmental Toxicology*
*and Chemistry*, *25*(12), 3171–3180. https://doi.org/10.1897/05-509R1.1

Durrieu de Madron, X., Guieu, C., Sempéré, R., Conan, P., Cossa, D., D'Ortenzio, F., Estournel, C., Gazeau, F.,
Rabouille, C., Stemmann, L., Bonnet, S., Diaz, F., Koubbi, P., Radakovitch, O., Babin, M., Baklouti, M.,
Bancon-Montigny, C., Belviso, S., Bensoussan, N., … Verney, R. (2011). Marine ecosystems' responses to
climatic and anthropogenic forcings in the Mediterranean. *Progress in Oceanography*, *91*(2), 97–166.
https://doi.org/10.1016/j.pocean.2011.02.003

Dydak, U., Jiang, Y. M., Long, L. L., Zhu, H., Chen, J., Li, W. M., Edden, R. A. E., Hu, S., Fu, X., Long, Z., Mo, X. A.,
Meier, D., Harezlak, J., Aschner, M., Murdoch, J. B., & Zheng, W. (2011). In vivo measurement of brain
GABA concentrations by magnetic resonance spectroscopy in smelters occupationally exposed to
manganese. *Environmental Health Perspectives*, *119*(2), 219–224. https://doi.org/10.1289/ehp.1002192

Epel, E. S., Blackburn, E. H., Lin, J., Dhabhar, F. S., Adler, N. E., Morrow, J. D., & Cawthon, R. M. (2004).
Accelerated telomere shortening in response to life stress. *Proceedings of the National Academy of*
*Sciences of the United States of America*, *101*(49), 17312–17315.
https://doi.org/10.1073/pnas.0407162101

Exley, C., Chappell, J. S., & Birchall, J. D. (1991). A mechanism for acute aluminium toxicity in fish. *Journal of*
*Theoretical Biology*, *151*(3), 417–428. https://doi.org/10.1016/S0022-5193(05)80389-3

Farag, A. M., Woodward, D. F., Goldstein, J. N., Brumbaugh, W., & Meyer, J. S. (1998). Concentrations of
metals associated with mining waste in sediments, biofilm, benthic macroinvertebrates, and fish from
the Coeur d'Alene River Basin, Idaho. *Archives of Environmental Contamination and Toxicology*, *34*(2),
119–127. https://doi.org/10.1007/s002449900295

Feuilloley, G., Fromentin, J. M., Stemmann, L., Demarcq, H., Estournel, C., & Saraux, C. (2020). Concomitant
changes in the environment and small pelagic fish community of the Gulf of Lions. *Progress in*
*Oceanography*, *186*(June), 102375. https://doi.org/10.1016/j.pocean.2020.102375

Folch, J., Lees, M., & Stanley, G. H. S. (1957). A simple method for the isolation and purification of total lipides
from animal tissues. *Journal of Biological Chemistry*, *226*(1), 497–509. https://doi.org/10.1016/S0021-
9258(18)64849-5

Friberg, L., Nordberg, G. F., & Vouk, V. B. (1979). *Handbook of the toxicology of metals*. Elsevier/North-Holland
Biomedical Press.

Gabrielides, G. P. (1995). Pollution of the Mediterranean Sea. *Water Science and Technology*, *32*(9–10), 1–10.
https://doi.org/10.1016/0273-1223(96)00070-4

Genchi, G., Carocci, A., Lauria, G., Sinicropi, M. S., & Catalano, A. (2021). Thallium Use, Toxicity, and
Detoxification Therapy: An Overview. *Applied Sciences*, *11*(18), 8322.

786    https://doi.org/10.3390/app11188322

787 Guérin, T., Chekri, R., Vastel, C., Sirot, V., Volatier, J. L., Leblanc, J. C., & Noël, L. (2011). Determination of 20
788    trace elements in fish and other seafood from the French market. *Food Chemistry*, *127*(3), 934–942.
789    https://doi.org/10.1016/j.foodchem.2011.01.061

790 Guerra-García, J. M., Calero-Cano, S., Donázar-Aramendía, Í., Giráldez, I., Morales, E., Arechavala-Lopez, P., &
791    Cervera-Currado, J. L. (2023). Assessment of elemental composition in commercial fish of the Bay of
792    Cádiz, Southern Iberian Peninsula. *Marine Pollution Bulletin*, *187*(November 2022).
793    https://doi.org/10.1016/j.marpolbul.2022.114504

794 Guo, Z., Zhang, W., Du, S., Green, I., Tan, Q., & Zhang, L. (2016). Developmental patterns of copper
795    bioaccumulation in a marine fish model *Oryzias melastigma*. *Aquatic Toxicology*, *170*, 216–222.
796    https://doi.org/10.1016/j.aquatox.2015.11.026

797 Halliwell, B., & Gutteridge, J. M. C. (2015). *Free Radicals in Biology and Medicine*. Oxford University Press.
798    https://doi.org/10.1093/acprof:oso/9780198717478.001.0001

799 Harmelin-Vivien, M., Cossa, D., Crochet, S., Bănaru, D., Letourneur, Y., & Mellon-Duval, C. (2009). Difference of
800    mercury bioaccumulation in red mullets from the north-western Mediterranean and Black seas. *Marine
801    Pollution Bulletin*, *58*(5), 679–685. https://doi.org/10.1016/j.marpolbul.2009.01.004

802 Heys, K. A., Shore, R. F., Pereira, M. G., Jones, K. C., & Martin, F. L. (2016). Risk assessment of environmental
803    mixture effects. *RSC Advances*, *6*(53), 47844–47857. https://doi.org/10.1039/c6ra05406d

804 Hollis, L., McGeer, J. C., McDonald, D. G., & Wood, C. M. (1999). Cadmium accumulation, gill Cd binding,
805    acclimation, and physiological effects during long term sublethal Cd exposure in rainbow trout. *Aquatic
806    Toxicology*, *46*(2), 101–119. https://doi.org/10.1016/S0166-445X(98)00118-0

807 Isani, G., Andreani, G., Cocchioni, F., Fedeli, D., Carpené, E., & Falcioni, G. (2009). Cadmium accumulation and
808    biochemical responses in *Sparus aurata* following sub-lethal Cd exposure. *Ecotoxicology and
809    Environmental Safety*, *72*(1), 224–230. https://doi.org/10.1016/j.ecoenv.2008.04.015

810 Jaishankar, M., Tseten, T., Anbalagan, N., Mathew, B. B., & Beeregowda, K. N. (2014). Toxicity , mechanism and
811    health effects of some heavy metals. *Interdisciplinary Toxicology*, *7*(2), 60–72.
812    https://doi.org/10.2478/intox-2014-0009

813 Javed, M., Ahmad, M. I., Usmani, N., & Ahmad, M. (2017). Multiple biomarker responses (serum biochemistry,
814    oxidative stress, genotoxicity and histopathology) in *Channa punctatus* exposed to heavy metal loaded
815    waste water. *Scientific Reports*, *7*(1), 1675. https://doi.org/10.1038/s41598-017-01749-6

816 Joiris, C. R., Holsbeek, L., & Moatemri, N. L. (1999). Total and methylmercury in sardines Sardinella aurita and
817    *Sardina pilchardus* from Tunisia. *Marine Pollution Bulletin*, *38*(3), 188–192.
818    https://doi.org/10.1016/S0025-326X(98)00171-4

819 Kardon, J. R., & Vale, R. D. (2009). Regulators of the cytoplasmic dynein motor. *Nature Reviews Molecular Cell
820    Biology*, *10*(12), 854–865. https://doi.org/10.1038/nrm2804

821 Karim, M., Puiseux-Dao, S., & Edery, M. (2011). Toxins and stress in fish: Proteomic analyses and response
822    network. *Toxicon*, *57*(7–8), 959–969. https://doi.org/10.1016/j.toxicon.2011.03.018

823 Kaushal, S. S., Gold, A. J., Bernal, S., Johnson, T. A. N., Addy, K., Burgin, A., Burns, D. A., Coble, A. A., Hood, E.,
824    Lu, Y. H., Mayer, P., Minor, E. C., Schroth, A. W., Vidon, P., Wilson, H., Xenopoulos, M. A., Doody, T.,
825    Galella, J. G., Goodling, P., … Belt, K. T. (2018). Watershed 'chemical cocktails': forming novel elemental
826    combinations in Anthropocene fresh waters. *Biogeochemistry*, *141*(3), 281–305.
827    https://doi.org/10.1007/s10533-018-0502-6

Koivula, M. J., & Eeva, T. (2010). Metal-related oxidative stress in birds. *Environmental Pollution*, *158*(7), 2359–
2370. https://doi.org/10.1016/j.envpol.2010.03.013

Kumari, B., Kumar, V., Sinha, A. K., Ahsan, J., Ghosh, A. K., Wang, H., & DeBoeck, G. (2017). Toxicology of
arsenic in fish and aquatic systems. *Environmental Chemistry Letters*, *15*(1), 43–64.
https://doi.org/10.1007/s10311-016-0588-9

Kumarswamy, R., & Chandna, S. (2009). Putative partners in Bax mediated cytochrome-c release: ANT, CypD,
VDAC or none of them? *Mitochondrion*, *9*(1), 1–8. https://doi.org/10.1016/j.mito.2008.10.003

Langfelder, P., & Horvath, S. (2008). WGCNA: An R package for weighted correlation network analysis. *BMC
Bioinformatics*, *9*. https://doi.org/10.1186/1471-2105-9-559

Le Bourg, B., Bănaru, D., Saraux, C., Nowaczyk, A., Le Luherne, E., Jadaud, A., Bigot, J. L., & Richard, P. (2015).
Trophic niche overlap of sprat and commercial small pelagic teleosts in the Gulf of Lions (NW
Mediterranean Sea). *Journal of Sea Research*, *103*, 138–146.
https://doi.org/10.1016/j.seares.2015.06.011

Leaver, M. J., Diab, A., Boukouvala, E., Williams, T. D., Chipman, J. K., Moffat, C. F., Robinson, C. D., & George,
S. G. (2010). Hepatic gene expression in flounder chronically exposed to multiply polluted estuarine
sediment: Absence of classical exposure "biomarker" signals and induction of inflammatory, innate
immune and apoptotic pathways. *Aquatic Toxicology*, *96*(3), 234–245.
https://doi.org/10.1016/j.aquatox.2009.10.025

Li, Y., & Wang, W. X. (2021). Integrated transcriptomics and proteomics revealed the distinct toxicological
effects of multi-metal contamination on oysters. *Environmental Pollution*, *284*(June), 117533.
https://doi.org/10.1016/j.envpol.2021.117533

Liu, Y., Shang, D., Yang, Y., Cui, P., & Sun, J. (2022). Transcriptomic Analysis Provides Insights into Microplastic
and Heavy Metal Challenges in the Line Seahorse (*Hippocampus erectus*). *Fishes*, *7*(6), 338.
https://doi.org/10.3390/fishes7060338

Lloret, J., Shulman, G., & Love, R. M. (2013). Condition and Health Indicators of Exploited Marine Fishes. In
*Condition and Health Indicators of Exploited Marine Fishes*. John Wiley & Sons.
https://doi.org/10.1002/9781118752777

Lohner, T. W., Reash, R. J., Willet, V. E., & Rose, L. A. (2001). Assessment of tolerant sunfish populations
(*Lepomis* sp.) inhabiting selenium-laden coal ash effluents: 1. Hematological and population level
assessment. *Ecotoxicology and Environmental Safety*, *50*(3), 203–216.
https://doi.org/10.1006/eesa.2001.2097

López-Pedrouso, M., Varela, Z., Franco, D., Fernández, J. A., & Aboal, J. R. (2020). Can proteomics contribute to
biomonitoring of aquatic pollution? A critical review. In *Environmental Pollution* (Vol. 267, p. 115473).
Elsevier Ltd. https://doi.org/10.1016/j.envpol.2020.115473

Lozano-Bilbao, E., Díaz, Y., Lozano, G., Jurado-Ruzafa, A., Hardisson, A., Rubio, C., Jiménez, S., González-Weller,
D., & Gutiérrez, Á. J. (2019). Metal Content in Small Pelagic Fish in the North-West Africa. *Thalassas*,
*35*(2), 643–653. https://doi.org/10.1007/s41208-019-00141-7

McCuaig, L. M., Martyniuk, C. J., & Marlatt, V. L. (2020). Morphometric and proteomic responses of early-life
stage rainbow trout (*Oncorhynchus mykiss*) to the aquatic herbicide diquat dibromide. *Aquatic
Toxicology*, *222*, 105446. https://doi.org/10.1016/j.aquatox.2020.105446

Mela, M., Randi, M. A. F., Ventura, D. F., Carvalho, C. E. V., Pelletier, E., & Oliveira Ribeiro, C. A. (2007). Effects
of dietary methylmercury on liver and kidney histology in the neotropical fish *Hoplias malabaricus*.
*Ecotoxicology and Environmental Safety*, *68*(3), 426–435. https://doi.org/10.1016/j.ecoenv.2006.11.013

Mitra, A., Shevde, L. A., & Samant, R. S. (2009). Multi-faceted role of HSP40 in cancer. *Clinical and Experimental Metastasis*, *26*(6), 559–567. https://doi.org/10.1007/s10585-009-9255-x

Mollan, T. L., Khandros, E., Weiss, M. J., & Olson, J. S. (2012). Kinetics of α-globin binding to α-Hemoglobin Stabilizing Protein (AHSP) indicate preferential stabilization of hemichrome folding intermediate. *Journal of Biological Chemistry*, *287*(14), 11338–11350. https://doi.org/10.1074/jbc.M111.313247

Morcillo, P., Esteban, M., & Cuesta, A. (2016). Heavy metals produce toxicity, oxidative stress and apoptosis in the marine teleost fish SAF-1 cell line. *Chemosphere*, *144*, 225–233. https://doi.org/10.1016/j.chemosphere.2015.08.020

Moreira, C. M., Oliveira, E. M., Bonan, C. D., Sarkis, J. J. F., & Vassallo, D. V. (2003). Effects of mercury on myosin ATPase in the ventricular myocardium of the rat. *Comparative Biochemistry and Physiology - C Toxicology and Pharmacology*, *135*(3), 269–275. https://doi.org/10.1016/S1532-0456(03)00110-8

Moret, Y., & Schmid-Hempel, P. (2000). Survival for immunity: The price of immune system activation for bumblebee workers. *Science*, *290*(5494), 1166–1168. https://doi.org/10.1126/science.290.5494.1166

Parrish, C. C. (1988). Dissolved and particulate marine lipid classes: a review. *Marine Chemistry*, *23*(1–2), 17–40. https://doi.org/10.1016/0304-4203(88)90020-5

Paschal, B. M., Shpetner, H. S., & Vallee, R. B. (1987). MAP 1C is a microtubule-activated ATPase which translocates microtubules in vitro and has dynein-like properties. *Journal of Cell Biology*, *105*(3), 1273–1282. https://doi.org/10.1083/jcb.105.3.1273

Pedrotti, M. L., Petit, S., Elineau, A., Bruzaud, S., Crebassa, J. C., Dumontet, B., Martí, E., Gorsky, G., & Cózar, A. (2016). Changes in the floating plastic pollution of the mediterranean sea in relation to the distance to land. *PLoS ONE*, *11*(8), 1–14. https://doi.org/10.1371/journal.pone.0161581

Pereira, S., Pinto, A. L., Cortes, R., Fontaínhas-Fernandes, A., Coimbra, A. M., & Monteiro, S. M. (2013). Gill histopathological and oxidative stress evaluation in native fish captured in Portuguese northwestern rivers. *Ecotoxicology and Environmental Safety*, *90*, 157–166. https://doi.org/10.1016/j.ecoenv.2012.12.023

Perez-Riverol, Y., Csordas, A., Bai, J., Bernal-Llinares, M., Hewapathirana, S., Kundu, D. J., Inuganti, A., Griss, J., Mayer, G., Eisenacher, M., Pérez, E., Uszkoreit, J., Pfeuffer, J., Sachsenberg, T., Yilmaz, Ş., Tiwary, S., Cox, J., Audain, E., Walzer, M., … Vizcaíno, J. A. (2019). The PRIDE database and related tools and resources in 2019: Improving support for quantification data. *Nucleic Acids Research*, *47*(D1), D442–D450. https://doi.org/10.1093/nar/gky1106

Plum, L. M., Rink, L., & Hajo, H. (2010). The essential toxin: Impact of zinc on human health. *International Journal of Environmental Research and Public Health*, *7*(4), 1342–1365. https://doi.org/10.3390/ijerph7041342

Queiros, Q., Fromentin, J. M., Gasset, E., Dutto, G., Huiban, C., Metral, L., Leclerc, L., Schull, Q., McKenzie, D. J., & Saraux, C. (2019). Food in the sea: Size also matters for pelagic fish. *Frontiers in Marine Science*, *6*(JUL), 385. https://doi.org/10.3389/fmars.2019.00385

Queiros, Q., Mckenzie, D. J., Dutto, G., Killen, S., Saraux, C., & Schull, Q. (2024). Fish shrinking , energy balance and climate change. *Science of the Total Environment*, *906*, 167310. https://doi.org/10.1016/j.scitotenv.2023.167310

Radakovitch, O., Roussiez, V., Ollivier, P., Ludwig, W., Grenz, C., & Probst, J. L. (2008). Input of particulate heavy metals from rivers and associated sedimentary deposits on the Gulf of Lion continental shelf. *Estuarine, Coastal and Shelf Science*, *77*(2), 285–295. https://doi.org/10.1016/j.ecss.2007.09.028

Regoli, F., & Giuliani, M. E. (2014). Oxidative pathways of chemical toxicity and oxidative stress biomarkers in
marine organisms. *Marine Environmental Research*, *93*, 106–117.
https://doi.org/10.1016/j.marenvres.2013.07.006

Robledo Ardila, P. A., Álvarez-Alonso, R., Árcega-Cabrera, F., Durán Valsero, J. J., Morales García, R., Lamas-
Cosío, E., Oceguera-Vargas, I., & DelValls, A. (2024). Assessment and Review of Heavy Metals Pollution in
Sediments of the Mediterranean Sea. *Applied Sciences*, *14*(4). https://doi.org/10.3390/app14041435

Rodríguez-Ortega, M. J., Grøsvik, B. E., Rodríguez-Ariza, A., Goksøyr, A., & López-Barea, J. (2003). Changes in
protein expression profiles in bivalve molluscs (*Chamaelea gallina*) exposed to four model environmental
pollutants. *Proteomics*, *3*(8), 1535–1543. https://doi.org/10.1002/pmic.200300491

Sacks, F. M. (2006). The apolipoprotein story. *Atherosclerosis Supplements*, *7*(4), 23–27.
https://doi.org/10.1016/j.atherosclerosissup.2006.05.004

Sakuragui, M. M., Paulino, M. G., Pereira, C. D. S., Carvalho, C. S., Sadauskas-Henrique, H., & Fernandes, M. N.
(2013). Integrated use of antioxidant enzymes and oxidative damage in two fish species to assess
pollution in man-made hydroelectric reservoirs. *Environmental Pollution*, *178*, 41–51.
https://doi.org/10.1016/j.envpol.2013.02.032

Sanchez, B. C., Ralston-Hooper, K., & Sepúlveda, M. S. (2011). Review of recent proteomic applications in
aquatic toxicology. *Environmental Toxicology and Chemistry*, *30*(2), 274–282.
https://doi.org/10.1002/etc.402

Santos, C. R., & Schulze, A. (2012). Lipid metabolism in cancer. *FEBS Journal*, *279*(15), 2610–2623.
https://doi.org/10.1111/j.1742-4658.2012.08644.x

Saraux, C., Van Beveren, E., Brosset, P., Queiros, Q., Bourdeix, J.-H., Dutto, G., Gasset, E., Jac, C., Bonhommeau,
S., & Fromentin, J.-M. (2019). Small pelagic fish dynamics: A review of mechanisms in the Gulf of Lions.
*Deep Sea Research Part II: Topical Studies in Oceanography*, *159*, 52–61.
https://doi.org/10.1016/j.dsr2.2018.02.010

Sardenne, F., Bodin, N., Metral, L., Crottier, A., Le Grand, F., Bideau, A., Brisset, B., Bourjea, J., Saraux, C.,
Bonhommeau, S., Kerzérho, V., Bernard, S., & Rouyer, T. (2019). Effects of extraction method and
storage of dry tissue on marine lipids and fatty acids. *Analytica Chimica Acta*, *1051*(November), 82–93.
https://doi.org/10.1016/j.aca.2018.11.012

Sharma, S., Sharma, V., & Chatterjee, S. (2021). Microplastics in the Mediterranean Sea : Sources , Pollution
Intensity , Sea Health , and Regulatory Policies. *Frontiers in Marine Science*, *8*(May), 1–15.
https://doi.org/10.3389/fmars.2021.634934

Si, L. F., Wang, C. C., Guo, S. N., Zheng, J. L., & Xia, H. (2019). The lagged effects of environmentally relevant
zinc on non-specific immunity in zebrafish. *Chemosphere*, *214*, 85–93.
https://doi.org/10.1016/j.chemosphere.2018.09.050

Souid, G., Souayed, N., Yaktiti, F., & Maaroufi, K. (2015). Lead accumulation pattern and molecular biomarkers
of oxidative stress in seabream ( *Sparus aurata* ) under short-term metal treatment. *Drug and Chemical
Toxicology*, *38*(1), 98–105. https://doi.org/10.3109/01480545.2014.917091

Strady, E., Harmelin-Vivien, M., Chiffoleau, J. F., Veron, A., Tronczynski, J., & Radakovitch, O. (2015). 210Po and
210Pb trophic transfer within the phytoplankton-zooplankton-anchovy/sardine food web: A case study
from the Gulf of Lion (NW Mediterranean Sea). *Journal of Environmental Radioactivity*, *143*, 141–151.
https://doi.org/10.1016/j.jenvrad.2015.02.019

Tang, W. L., Evans, D., Kraemer, L., & Zhong, H. (2017). Body size-dependent Cd accumulation in the zebra
mussel *Dreissena polymorpha* from different routes. *Chemosphere*, *168*, 825–831.

https://doi.org/10.1016/j.chemosphere.2016.10.128

Tchounwou, P. B., Yedjou, C. G., Patlolla, A. K., & Sutton, D. J. (2012). Heavy Metal Toxicity and the
Environment. In A. Luch (Ed.), *Molecular, Clinical and Environmental Toxicology: Volume 3:*
*Environmental Toxicology* (pp. 133–164). Springer Basel. https://doi.org/10.1007/978-3-7643-8340-4_6

Teulier, L., Thoral, E., Queiros, Q., McKenzie, D. J., Roussel, D., Dutto, G., Gasset, E., Bourjea, J., & Saraux, C.
(2019). Muscle bioenergetics of two emblematic Mediterranean fish species: *Sardina pilchardus* and
*Sparus aurata*. *Comparative Biochemistry and Physiology -Part A : Molecular and Integrative Physiology*,
*235*, 174–179. https://doi.org/10.1016/j.cbpa.2019.06.008

Tocher, D. R. (2003). Metabolism and functions of lipids and fatty acids in teleost fish. In *Reviews in Fisheries*
*Science* (Vol. 11, Issue 2, pp. 107–184).  Taylor & Francis Group . https://doi.org/10.1080/713610925

Tournois, J., Darnaude, A. M., Ferraton, F., Aliaume, C., Mercier, L., & McKenzie, D. J. (2017). Lagoon nurseries
make a major contribution to adult populations of a highly prized coastal fish. *Limnology and*
*Oceanography*, *62*(3), 1219–1233. https://doi.org/10.1002/lno.10496

Tronczynski, J., Carlotti, F., Bodin, N., & Radakovitch, O. (2013). Contaminants bioaccumulation and
biomagnification in a short trophic system: phytoplankton, zooplankton, anchovy, sardine (COSTAS).
*Ciesm*.

Van Beveren, E., Fromentin, J.-M., Rouyer, T., Bonhommeau, S., Brosset, P., & Saraux, C. (2016). The fisheries
history of small pelagics in the Northern Mediterranean. *ICES Journal of Marine Science*, *73*(6), 1475–
1484. https://doi.org/10.1093/ICESJMS/FSW023

Van Beveren, E., Fromentin, J. M., Bonhommeau, S., Nieblas, A. E., Metral, L., Brisset, B., Jusup, M., Bauer, R.
K., Brosset, P., & Saraux, C. (2017). Predator–prey interactions in the face of management regulations:
Changes in Mediterranean small pelagic species are not due to increased tuna predation. *Canadian*
*Journal of Fisheries and Aquatic Sciences*, *74*(9), 1422–1430. https://doi.org/10.1139/cjfas-2016-0152

Van Beveren, E., Keck, N., Fromentin, J. M., Laurence, S., Boulet, H., Labrut, S., Baud, M., Bigarré, L., Brosset, P.,
& Saraux, C. (2016). Can pathogens alter the population dynamics of sardine in the NW Mediterranean?
*Marine Biology*, *163*(12), 240. https://doi.org/10.1007/s00227-016-3015-7

Van Meer, G., Voelker, D. R., & Feigenson, G. W. (2008). Membrane lipids: Where they are and how they
behave. *Nature Reviews Molecular Cell Biology*, *9*(2), 112–124. https://doi.org/10.1038/nrm2330

Wang, L., & Gallagher, E. P. (2013). Role of Nrf2 antioxidant defense in mitigating cadmium-induced oxidative
stress in the olfactory system of zebrafish. *Toxicology and Applied Pharmacology*, *266*(2), 177–186.
https://doi.org/10.1016/j.taap.2012.11.010

Wang, M., Wang, Y., Zhang, L., Wang, J., Hong, H., & Wang, D. (2013). Quantitative proteomic analysis reveals
the mode-of-action for chronic mercury hepatotoxicity to marine medaka (*Oryzias melastigma*). *Aquatic*
*Toxicology*, *130–131*, 123–131. https://doi.org/10.1016/j.aquatox.2013.01.012

Wijesooriya, K., Jadaan, S. A., Perera, K. L., Kaur, T., & Ziemann, M. (2022). Urgent need for consistent
standards in functional enrichment analysis. *PLOS Computational Biology*, *18*(3), 1–14.
https://doi.org/10.1371/journal.pcbi.1009935

Williams, C. R., & Yoshida-Honmachi, E. P. (2013). Effects of cadmium on olfactory mediated behaviors and
molecular biomarkers in coho salmon (*Oncorhynchus kisutch*). *Aquatic Toxicology*, *140–141*, 295–302.
https://doi.org/10.1016/j.aquatox.2013.06.010

Wood, C. ., Farell, A. ., & Brauner, C. . (2011). Copper of the Fish Physiology. In *Fish physiology* (pp. 55–133).

Xu, L., Lu, Z., Ji, C., Cong, M., Li, F., Shan, X., & Wu, H. (2019). Toxicological effects of As (V) in juvenile rockfish
*Sebastes schlegelii* by a combined metabolomic and proteomic approach. *Environmental Pollution*, *255*,
113333. https://doi.org/10.1016/j.envpol.2019.113333

Yadetie, F., Karlsen, O. A., Lanzén, A., Berg, K., Olsvik, P., Hogstrand, C., & Goksøyr, A. (2013). Global
transcriptome analysis of Atlantic cod (*Gadus morhua*) liver after in vivo methylmercury exposure
suggests effects on energy metabolism pathways. *Aquatic Toxicology*, *126*, 314–325.
https://doi.org/10.1016/j.aquatox.2012.09.013

Yin, S., Liu, L., & Gan, W. (2021). The Roles of Post-Translational Modifications on mTOR Signaling.
*International Journal of Molecular Sciences*, *22*(4). https://doi.org/10.3390/ijms22041784

Zhol, S., Ackman, R. G., & Morrison, C. (1995). Storage of lipids in the myosepta of Atlantic salmon (*Salmo
salar*). *Fish Physiology and Biochemistry*, *14*(2), 171–178. https://doi.org/10.1007/BF00002460

1008