Automated 3D analysis of social head-gaze behaviors in freely moving marmosets

Feng Xing[1,2], Alec G. Sheffield[1,2,3], Monika P. Jadi[2,3,5,†], Steve W.C Chang[2,4,5,6,†], and Anirvan S. Nandy[2,4,5,6,†]

[1]Inderdepartmental Neuroscience Program, Yale University, New Haven, CT

[2]Department of Neuroscience, Yale University, New Haven, CT

[3]Department of Psychiatry, Yale University, New Haven, CT

[4]Department of Psychology, Yale University, New Haven, CT

[5]Wu Tsai Institute, Yale University, New Haven, CT

[6]Kavli Institute for Neuroscience, Yale University, New Haven, CT

[†]senior authors, equal contribution

Correspondence: anirvan.nandy@yale.edu

Summary

Social communication relies on the ability to perceive and interpret the direction of others' attention, which is commonly conveyed through head orientation and gaze direction in both humans and non-human primates. However, traditional social gaze experiments in non-human primates require restraining head movements, which significantly limit their natural behavioral repertoire. Here, we developed a novel framework for accurately tracking facial features and three-dimensional head gaze orientations of multiple freely moving common marmosets (*Callithrix jacchus*). To accurately track the facial features of marmoset dyads in an arena, we adapted computer vision tools using deep learning networks combined with triangulation algorithms applied to the detected facial features to generate dynamic geometric facial frames in 3D space, overcoming common occlusion challenges. Furthermore, we constructed a virtual cone, oriented perpendicular to the facial frame, to model the head gaze directions. Using this framework, we were able to detect different types of interactive social gaze events, including partner-directed gaze and jointly-directed gaze to a shared spatial location. We observed clear effects of sex and familiarity on both interpersonal distance and gaze dynamics in marmoset dyads. Unfamiliar pairs exhibited more stereotyped patterns of arena occupancy, more sustained levels of social gaze across inter-animal distance, and increased gaze monitoring. On the other hand, familiar pairs exhibited higher levels of joint gazes. Moreover, males displayed significantly elevated levels of gazes toward females' faces and the surrounding regions irrespective of familiarity. Our study lays the groundwork for a rigorous quantification of primate behaviors in naturalistic settings.

Introduction

Primates, including humans, exhibit complex social structures and engage in rich interactions with members of their species, which are crucial for their survival and development. Among social stimuli, the face holds paramount importance with specialized neural systems (Deen et al., 2023; Hesse & Tsao, 2020) and is attentively prioritized by primates during much of their social interaction. Notably, the eyes garner the most attention among all facial features, playing a pivotal role in indicating the direction of others' attention and also possibly their intention (Dal Monte et al., 2015; Emery, 2000; Itier et al., 2007). Indeed, understanding and interpreting the gaze of fellow individuals is a fundamental attribute of the theory of mind (ToM) (Martin & Santos, 2016; Saxe & Kanwisher, 2003). While current studies of social gaze using pairs of rhesus macaques (*Macaca mulatta*) in a controlled laboratory setting (Dal Monte et al., 2016; Mosher et al., 2014; Ramezanpour & Thier 2020; Shepherd et al., 2006; Shepherd & Freiwald, 2018) provide valuable insights into social gaze behaviors, they are nevertheless limited in their ecological relevance.

To address these limitations, we turned to common marmosets (*Callithrix jacchus*), a highly prosocial primate species known for their social behavioral and cognitive similarities to humans (Miller et al., 2016). Marmosets are also a model system with increasing applications in computational ethology (Mitchell et al., 2014; Ngo et al., 2022). Like humans, they engage in cooperative breeding, a social system in which individuals care for offspring other than their own, usually at the expense of their reproduction (French, 1997; Solomon & French, 1997). Gaze directions, inferred from head orientation, hold crucial information about marmoset social interactions (Helney & Blazquez, 2011; Spadacenta et al., 2022). The emergence of computational ethology (Anderson & Perona, 2014; Datta, Anderson, Branson, Perona, & Leifer, 2019) has propelled the development of a host of computer vision tools using deep neural networks (e.g., OpenPose by Cao et al., 2017, DeepLabCut by Mathis et al., 2018, DANNCE by Dunn et al., 2021; SLEAP by Pereira et al., 2022). However, tracking head gaze direction in multiple freely moving marmosets poses a challenging problem, not yet solved by existing computer vision tools.

80   This problem is further complicated by the fact that accurately tracking gaze orientations
81   in primates requires three-dimensional information.

82
Here, we propose a novel framework based on a modified DeepLabCut pipeline,
capable of accurately detecting body parts of multiple marmosets in 2D space and
triangulating them in 3D space. By reconstructing the face frame with six facial points in
3D space, we can infer the head gaze of each animal across time. Importantly, marmosets
that are not head-restrained use rapid head movements for reorienting and visual
exploration (Pandey et al., 2020), and therefore the head direction serves as an excellent
proxy for gaze orientation in unrestrained marmosets. With this framework in place, we
investigated the gaze behaviors of male-female pairs of freely moving and interacting
marmosets to quantify their social gaze dynamics. We investigated these gaze dynamics
along the dimensions of sex and familiarity and found several key differences along both
of these important social dimensions, including increased partner gaze in males that is
modulated by familiarity, increased joint gaze among familiar pairs, and increased gaze
monitoring by males. This fully automated tracking system can thus serve as a powerful
tool for investigating primate group dynamics in naturalistic environments.

97
Results
98
99
Experimental setup and reconstruction of video images in 3D space
101
The experimental setup consisted of two arenas made with acrylic plates that allowed two
marmosets to visually interact with each other while being physically separate (Fig. 1A).
Each arena was 60.96 cm long, 30.48 cm wide, and 30.48 cm high. Five sides of the
arena, except the bottom, were transparent allowing a clear view of the animal subjects
under observation. The bottom side of the arena was perforated with 1-inch diameter
holes arranged in a hexagonal pattern to aid the animal's traction. The arenas were
mounted on a rigid frame made of aluminum building blocks, with the smaller sides facing
each other and were separated by a distance of 30.48 cm. A movable opaque divider was
placed between the arenas during intermittent breaks to prevent the animals from having
visual access to each other (Methods). Two monitors were attached to the aluminum
frame, one on each end, for displaying video or image stimuli to the animals. To capture

the whole experimental setup, two sets of four GoPro 8 cameras were attached to the
frame, where each set of cameras captured the view of one of the arenas.

After obtaining the intrinsic parameters of the cameras by calibration and the extrinsic
parameters of the cameras by L-frame analysis (see Methods), we established a world
coordinate system of each arena surrounded by the corresponding set of four cameras.
Crucially, the two independent world coordinate systems of the two arenas were
combined by measuring the distance between the two L-shaped frames and adding this
offset to one of the world coordinate systems.

With the established world coordinate system, any point captured by two or more
cameras could be triangulated into a common three-dimensional space. Thus, the
experimental setup was reconstructed into three-dimensional space by manually labeling
the vertices of the arenas and monitors in the image space captured by the cameras (Fig.
1B). The cameras on the monitor ends (marked as 'ML1', 'ML2', 'MR1', 'MR2' in Fig. 1B)
recorded both animal subjects, whereas the cameras in the middle (marked as 'OL1',
'OL2', 'OR1', 'OR2' in Fig. 1B) recorded only one animal subject.

Automatic detection of facial features of two marmosets

Six facial features – the two tufts, the central blaze, two eyes, and the mouth (Fig. 2A) –
were selected for automated tracking using a modified version of a deep neural network
(DeepLabCut, Mathis et al., 2018). The raw video from each camera was fed into the
network to compute probability heatmaps of each facial feature. We modified the original
method to detect features of two animals (Fig. 2B). After processing the raw video, two
locations with the highest probability over a threshold (95%) were picked from each
probability heatmap (Fig. 2B, *feature detection*). Since all the features from the same
animal should be clustered in image space, a K-means clustering algorithm (Fig. 2B, *initial*
clustering) was used on the candidate features with the constraint that one animal can
only have one unique feature (Fig. 2B, *refine clustering*); for example, one animal cannot
have two left eyes. After clustering, two clusters of features corresponding to the two
animals were obtained. To detect outliers that were not valid features, we first calculated
a distribution of within-cluster distances (Fig. 2B, *remove outliers*). Outliers were

determined as those points that had nearest-neighbor distances which were two standard
deviations above the average within-cluster distance, and were excluded from
subsequent analyses. Note that the above analyses were performed independently for
each video frame.
150
To establish temporal continuity across video frames and track animal identities, we
first calculated the centroid of each cluster (Fig. 2B, *calculate centroids*). Under the
heuristic that centroid trajectories corresponding to each individual animal are smoothly
continuous in space and time (i.e., there are no sudden jumps or reversals in centroid
location across frames at our sampling rate of 30Hz), we assigned identities to the
centroids thus enabling us to track identities over time (Fig. 2B, *establish identities*). The
facial features corresponding to a centroid inherited this identity (Fig. 2B, *cluster with*
identities).
159
160   Our method thus allowed us to accurately detect and track the facial features of two
161 marmosets in the arena (Fig. 2C), and in more general contexts such as in a home cage
162 with occlusions (Supplementary Video 1).
163
164 **Inferring head-gaze direction in 3D space**
165
In our experimental setup, the cameras in the middle unambiguously recorded only one
animal. The centroids of the facial features in the image space recorded by the middle
cameras were triangulated into three-dimensional space and were constrained to be
confined within the bounds of the arena. Any missing centroids were filled by interpolation
from neighboring frames from both the past and future time points. The triangulated
centroids acquired from cameras in the middle were then projected into the image space
of cameras on the monitor ends. Since both animals were recorded by the cameras on
the monitor ends, facial features from these camera views were detected with identities
assigned (as described in the 2D pipeline). In the image space of cameras on the monitor
ends, the cluster of facial feature points closer to the projected centroid and within the
bounds of the arena were kept for later triangulation.
177

178   The detected facial features captured by all four cameras for one animal were
179   subjected to triangulation. For each feature, results from all possible pairs of four cameras
180   were triangulated. All the triangulation results were averaged to yield the final coordinates
181   of the body part in three-dimensional space. Any missing features were filled by
182   interpolation from neighboring frames including the previous and future time points.

183
The six facial points constituted a semi-rigid geometric frame as the animal moves in
3D space ('face frame'; Fig. 2), allowing us to infer the animal's head-gaze direction as
follows. The gaze orientation was calculated as the normal to the facial plane defined by
the two eyes and the central blaze. The position of the ear tufts, which were behind this
facial plane, was used to determine gaze direction. Since marmoset saccade amplitudes
are largely restricted to 10 degrees (median less than 4 degrees) (Mitchell et al., 2014),
we modeled the head gaze as a virtual cone with a solid angle of 10 degrees ('gaze cone')
emanating from the facial plane (Fig. 3A). Notably, with multiple camera views, the face
frame can be reconstructed even when the face was invisible to one of the cameras, such
that the reconstructed face frame in 3D can be projected back into the image space to
validate the accuracy of the detection and reconstruction (Fig. 3B). We were thus able to
obtain the animal's continuous movement trajectory and the corresponding gaze direction
over time (Fig. 3C, Supplementary Video 2).

197
We first examined our method's ability to characterize gaze behaviors of freely moving
marmosets by presenting either video or image stimuli to individual animals on a monitor
screen (see Methods). Marmosets exhibited longer gaze duration to video stimuli
compared to image stimuli (Fig. S1A; Mann Whitney U Test, $p < 0.001$). However, this
difference was not caused by differences in gaze dispersion (Fig. S1B; Mann Whitney U
Test, ns). By examining the frequency of gaze events, we found that marmosets gazed at
the monitor more during the early period of video stimuli presentations compared to the
late period (Fig. S1C; Mann Whitney U Test, $p < 0.001$), while there was no such
difference for the image stimuli (Fig. S1C; Mann Whitney U Test, ns). There was also a
significant difference in gaze frequency between the early period of video presentations
compared to the same period for image presentations (Fig. S1C; Mann Whitney U Test,
$p < 10^{-10}$). Taken together, our results support that dynamic visual stimuli elicit greater

overt attention compared to static stimuli in marmosets, similar to macaques (Dal Monte
et al., 2016; Furl et al., 2012) and humans (Chevallier et al., 2015).

212
213   **Positional dynamics of marmoset dyads**
214
With this automated system in place, we recorded the behavior of four pairs of familiar
marmosets and four pairs of unfamiliar marmosets. Data from each pair was recorded in
one session consisting of ten five-minute free-viewing blocks interleaved with five-minute
breaks. Each pair consisted of a male and a female animal. The familiar pairs were cage
mates, while each member of an unfamiliar pair was from a different home cage with no
visual access to each other while in the colony. We first examined the movement
trajectories of marmoset dyads and used the centroids of the face frames across time to
represent the trajectories. For an example 5-minute segment (Fig. 4A), we observed that
the marmosets preferred to stay at the two ends of their respective arenas. This was
confirmed by a heatmap of projections of the trajectories on the plane parallel to the
vertical long side of the arenas ('XZ' plane). Furthermore, there were two hotspots along
the vertical axis in the heatmap of projections to the vertical short side plane ('YZ' plane),
suggesting that the animals' preferred body postures were either upright or crouched.

228
To quantify their positional dynamics, we examined the marginal distributions of the
movement trajectories along the horizontal ('X') and vertical ('Z') axes across all sessions
and grouped them along the dimensions of sex and familiarity (Fig. 4B). Along the X axis
(Fig. 4B, left), the distributions were slightly bimodal, with the main peak in the region
near to the inner edge of the arenas. Regardless of familiarity, male marmosets tended
to stay closer to the inner edge compared to females, as shown by the significant
differences in the distributions when X ranged from 0 to 150 mm. However, there were no
significant differences between the same sex members of familiar and unfamiliar pairs.
For the Z axis (Fig. 4B, right), the distributions were bimodal (Warren Sarle's bimodality
test), consistent with what we observed in the heatmaps indicating either upright or
crouched postures. The positional distributions along the Z axis were not different based
on sex or familiarity.
241

To further characterize the positional dynamics of the freely moving dyads, we

calculated the distance between the centroids of the pairs. We then examined the

distributions of the inter-animal distance along the X-axis separately for familiar and

unfamiliar pairs (Fig. 4C). The distributions were trimodal, and can be explained by the

bimodal distribution of movement trajectories of individual marmosets along the X-axis.

As mentioned above, marmosets tended to stay at the two ends of their arenas, and thus,

combinations of preferred positions at the two ends for the dyads (see insets in Fig. 4C)

resulted in the trimodal distribution. We termed these three peaks as 'Near', 'Intermediate',

and 'Far'. To quantify these distributions, we fitted the empirical data with mixture models

using maximum likelihood estimation (see Methods). The inter-animal distance for

unfamiliar pairs was best fitted by a mixture of three Gaussians while the distribution for

familiar pairs was best fitted by a mixture of Gamma and Gaussian distributions. The first

peak ('Near') of the familiar-pair distribution was best fitted by a Gamma distribution,

implying a higher degree of dispersion when familiar marmosets were close to each other.

Upon examining the temporal evolution of the inter-animal distance (within each 5-minute

viewing block), we detected that the inter-animal distance of unfamiliar pairs increased

over time, whereas this distance fluctuated over time for familiar pairs (Fig. 4D), further

indicating that the positional dynamics of marmoset dyads depended on familiarity.

260
261   **Social gaze dynamics of marmoset dyads**
262
We next investigated the interactive aspects of gaze behaviors in freely moving marmoset

dyads. The gaze interaction between two animals could be simplified as the relative

positions of two gaze cones in three-dimensional space (see Methods; Fig. 5A;

Supplementary Video 3). If the gaze cone of one animal intersected with the facial plane

of the second animal (but not vice versa), we termed it 'partner gaze'. If the gaze cones

of both animals intersected with that of the other's facial place, we termed it 'reciprocal

gaze'. In our dataset, the instances of reciprocal gaze were very low and were thus

excluded from further analysis. If the two cones intersected anywhere outside the facial

planes, we termed it 'joint gaze'. All other cases were regarded as 'no interaction' between

the two animals.

We analyzed the videos of marmoset dyads and identified stable gaze epochs by
thresholding the head velocity obtained from the centroids of the face frames (see
Methods). Stable epochs were categorized into gaze states based on the gaze event
types described above. We first analyzed the fraction of gaze states in the three position
ranges identified from the inter-animal distance analysis (Fig. 5B,C). We found that male
marmosets gazed more toward their partner females' faces regardless of familiarity ($p <$
0.01, χ^2 test). Fraction of male→female partner gaze incidents decreased with increasing
inter-animal distance for familiar pairs, while they remained constant in the case of
unfamiliar pairs (Fig. 5C). Moreover, females in unfamiliar pairs exhibited significantly (p
< 0.01, χ^2 test) higher partner gazes (female→male) compared to those in familiar pairs
(Fig. 5C). The total counts of social gaze states (joint gaze and partner gaze) were higher
for familiar pairs when they were near, but these decreased more dramatically with
increasing distance (Fig. 5B).

To investigate the dynamics of these gaze states, we computed state transition
probabilities among distinct gaze event types for familiar and unfamiliar dyads. We
applied a Markov chain model (see Methods) (Fig. 5D), in which the nodes were the gaze
states and the edges connecting the nodes represented the transitions between gaze
states. We first focused on the recurrent (self-transition) edges for the partner gaze states.
Recurrent edges indicate a transition back to the same stable gaze state after a break
likely due to physical movement, and reflect the robustness of the state despite movement.
In line with our previous results (Fig. 5B,C), males exhibit significantly higher (χ^2 test,
unfamiliar male vs unfamiliar female, $p < 10^{-10}$; familiar male vs familiar female, $p < 0.01$)
recurrent partner gazes compared to females, irrespective of familiarity (Fig. 5E).

A comparison of state transition probabilities across the dimension of familiarity
yielded several noteworthy findings (Fig. 5F). First, recurrent male partner gaze
(male→female) was significantly enhanced in unfamiliar pairs ($p < 0.05$, χ^2 test),
suggesting a heightened interest in unfamiliar females. Second, there was a higher
probability of transition from a female partner gaze to a male partner gaze in familiar pairs
compared to unfamiliar pairs, suggesting that familiar males have a greater awareness of
and tendency to reciprocate their partners' gaze ($p < 0.05$, χ^2 test). Third, there was a

higher probability of recurrent joint gazes in familiar pairs compared to unfamiliar pairs,
suggesting that familiar pairs explore common objects more than unfamiliar pairs ($p < 10^{-4}$, χ^2 test).
308

Monitoring others to anticipate their future actions is critical for successful social
interactions (Hari et al., 2015). In particular, successful interactive gaze exchanges
require constant monitoring of other's gaze. We analyzed the gaze distribution in the
surrounding region of a partner's face to estimate gaze monitoring tied to increased social
attention (Dal Monte et al., 2022). We quantified this by the distance between the centroid
of the partner's face-frame and the point of intersection of the gaze cone with the partner's
facial plane (Fig. 5G, left). Unfamiliar marmosets (both males and females) showed
significantly higher (Mann-Whitney U test, unfamiliar male vs familiar male, $p < 0.0001$;
unfamiliar female vs familiar female, $p < 0.001$) incidences of gaze toward the surrounding
region of the partner's face (Fig 5G, right; compare darker lines with the lighter lines).
Further, males exhibited markedly higher (Mann-Whitney U test, $p < 0.0001$) incidences
of gaze toward the partner females' face (Fig 5G, right; compare cyan lines with the
orange lines).
323

Overall, using our novel gaze tracking of freely moving marmoset dyads, we found
that both the social dimensions we examined – familiarity and sex – are significant
determinants of natural gaze dynamics among marmosets.

Discussion

In this study, we first presented a novel framework for the automated, markerless, and
identity-preserving tracking of 3D facial features of multiple marmosets. By building on
top of the deep-learning framework provided by DeepLabCut, we used a constellation of
cameras to overcome "blindspots" due to occlusion and imposed spatiotemporal
smoothness constraints on the detected features to establish and preserve identities
across time. The tracked facial features from each animal form a semi-rigid face frame as
the animal moves freely in 3D space, thereby allowing us to infer the animal's gaze

direction at each moment in time. It is important to reiterate that unrestrained marmosets
use rapid saccadic head-movements for reorienting (Pandey et al., 2020) and have a
limited amplitude range of saccadic eye-movements (Mitchell et al., 2014). Thus their
head direction serves as excellent proxy for gaze orientation in unrestrained conditions.
343
Primates are a highly visual species whose physical explorations of their environment
are not confined to two-dimensional surfaces. Gaze is a critical component of primate
social behavior and conveys important social signals such as interest, attention, and
emotion (Emery, 2000). Assessment of gaze is therefore important to understand non-
verbal communication and interpersonal dynamics. Our 3D gaze tracking approach was
able to capture both the positional and gaze dynamics of freely moving marmoset dyads
in a naturalistic context. We observed clear effects of sex and familiarity on both
interpersonal and gaze dynamics. Unfamiliar pairs exhibited more stereotyped patterns
of arena occupancy, more sustained levels of social gaze across distance, and increased
gaze monitoring, suggesting elevated levels of social attention compared to familiar pairs.
On the other hand, familiar pairs exhibited more recurrent joint gazes in the shared
environment compared to unfamiliar pairs. Familiar males also showed a higher tendency
to reciprocate their partner's gaze, suggesting a greater awareness of their partner's
social gaze state.
358
Supported by the natural ecology of marmosets (Yamamoto et al., 2014; Solomon &
French, 1997), we found dramatic sex differences in gaze behaviors, with males
exhibiting significantly elevated levels of gaze toward females' faces and the surrounding
regions irrespective of familiarity. It is important to note that dominance in marmosets is
not strictly determined by gender, as it can vary based on individual personalities and
intra-group social dynamics, although breeding females typically dominate social activity
within a group (Digby, 1995; Mustoe et al., 2023). While we have not explicitly controlled
for dominance in this study, whether part of the observed differences can be attributed to
dominance effects needs further exploration.
368
Gaze following plays a crucial role in social communication for humans and non-
human primates, allowing for joint attention (Emery et al., 1997; Brooks & Meltzoff, 2005;

Burkart & Heschl, 2006; Shepherd , 2010). Previous research demonstrated that head-
restrained marmosets exhibited preferential gazing toward marmoset face stimuli
observed by a conspecific in a quasi-reflexive manner during a free-choice task
(Spadacenta et al., 2019). Interestingly we did not find any differences in gaze-following
behaviors (transition from partner gaze to joint gaze) along the social dimensions we
tested here. Future investigation of such behaviors by manipulating social variables such
as dominance or kinship could provide a comprehensive understanding of gaze following
and joint attention in naturalistic behavioral contexts. The scarcity of reciprocal gazes in
our study may be attributed to the task-free experimental setup employed. Indeed, in
other joint action tasks requiring cooperation for rewards, marmosets actively engage in
reciprocal gaze behaviors (Miss & Burkart, 2018).

While we focused on the tracking of facial features in this study, our automated system
has the potential to extend to 3D whole-body tracking, encompassing limbs, tail, and the
main body features of marmosets. In our system, multiple cameras surrounding the arena
ensure that each body part of interest can be tracked through at least two cameras,
enabling triangulation in 3D space. Our current system uses a pre-trained ResNet model
(He et al., 2015) to track body parts of interest. However, considering the challenges
posed by whole-body tracking, such as interference from marmosets' fur that complicates
feature detection, the adoption of cutting-edge transformer networks like the vision
transformer model (Dosovitskiy et al., 2020) might significantly improve detection
performance. Such an advancement in tracking and reconstructing the entire marmoset
body frame would enable the analysis of such data using unsupervised learning
techniques (Berman et al., 2014; Calhoun et al., 2019) and thereby provide a deeper
understanding of primate social behavior.

In summary, our study lays the groundwork for a rigorous quantification of primate
behaviors in naturalistic settings. Not only does this allow us to gain deeper insights
beyond what is possible from field notes and observational studies, but it is also a key
first step to go beyond current reductionist paradigms and understanding the neural
dynamics underlying natural behaviors (Miller et al., 2022).

Acknowledgments

This research was supported by the National Institute of Mental Health (R21 120672, SWCC, ASN, MPJ), Simons Foundation Autism Research Initiative (SFARI 875855, SWCC, ASN, MPJ), Yale Orthwein Scholar Funds (ASN) and by the National Eye Institute core grant for vision research (P30 EY026878 to Yale University). We would like to thank the veterinary and husbandry staff at Yale for excellent animal care. We would like to thank Weikang Shi for helpful discussion on the manuscript.

Author contributions

ASN, SWCC & MPJ conceptualized the project. FX collected the data with assistance from AGS. FX analyzed the data. ASN supervised the project. FX, ASN, SWCC & MPJ wrote the manuscript.

Declaration of interests

The authors declare no competing interests.

Inclusion and Ethics

We support inclusive, diverse, and equitable conduct of research.

Figure captions

Figure 1. **Experimental setup and reconstruction in 3D space**.
(**A**) Two transparent acrylic arenas allowed marmosets to visually interact with each other. An opaque divider between the arenas was introduced intermittently to prevent visual access between animals. Two monitors on two ends were used to display video or image stimuli. Eight cameras surrounding the arenas ensured full coverage of both animals. LEDs at four positions were used to synchronize the video recordings across the set of cameras. (**B**) 3D reconstruction of the experimental setup. The two arenas are color-coded as orange and cyan. The cameras colored the same as the arenas indicate that they primarily record the marmoset in the corresponding arena. Two purple L-frames within the arenas were used to establish a world coordinate system in the reconstruction process. Two gray planes on both ends are the reconstructed monitors.

Figure 2. **Pipeline of detecting facial features of two marmosets**.
(**A**) Six facial features of the marmoset (face frame) are color-coded: right tuft (red), central blaze (yellow), left tuft (green), right eye (purple), left eye (blue), and mouth (magenta). (**B**) Feature tracking pipeline (right) with the corresponding illustration for each step across two adjacent video frames (left). At the end of the pipeline, the facial features are clustered with the identities assigned consistently across frames. Facial points are color-coded as in A. (**C**) Example frames of four steps in the pipeline shown in B. It can be seen clearly that the facial points are tracked and clustered accurately, and the identities are consistent across frames.

Figure 3. **3D Reconstruction of facial features and head gaze modeling**.
(**A**) The face frames of two marmosets are reconstructed in 3D using the tracked facial points in Fig. 2. A cone perpendicular to the face frame (gaze cone; 10-degree solid angle) is modeled as the head gaze. (**B**) Two example frames with the facial points projected from 3D space onto different camera views are shown. The left frame demonstrates that the facial points can be detected using information from other cameras, even if the face is invisible from that viewpoint. (**C**) Trajectory of the reconstructed face frame and the corresponding gaze cones across time.

Figure 4. **Positional dynamics of marmoset dyads**.

(**A**) Movement trajectories of the face frame centroids for a marmoset pair (orange for

female, cyan for male) in an example five-minute block. The heatmaps were calculated

using the projections of the trajectories to XY, YZ, and XZ planes. (**B**) Marginal

distributions of movement trajectories along the X and Z axes were calculated for all

marmosets and grouped by familiarity and sex (transparent colors for familiar pairs,

opaque colors for unfamiliar pairs). Black bars indicate significant differences between

pairs of distributions (Mann-Whitney U test, significance level at 5%). (**C**) Histograms of

inter-animal distance along the X axis show trimodal distributions for both familiar pairs

(gray) and unfamiliar pairs (black). The fitted red curve for the unfamiliar pairs is a tri-

Gaussian distribution, while the fitted red curve for the familiar pairs is a mixture of

Gamma and Gaussian distributions, with the first peak as the Gamma distribution. The

three regions were designated as 'Near', Intermediate', and 'Far'. The inset illustrates the

reason for this nomenclature. (**D**) Temporal evolution of inter-animal distance for

unfamiliar and familiar pairs (which each 5-minute viewing block). The central dark line is

the mean and the shaded area is the standard deviation. Black dots indicate significant

differences (Mann-Whitney U test, significance level at 5%).

Figure 5. **Live interactive gaze analysis of unfamiliar and familiar marmoset dyads**.

(**A**) Gaze type categorized based on the relative positions of the gaze cones. Joint gaze

is defined as two marmosets looking at the same location. A partner gaze is defined as

one animal looking at the other animal's face (but not vice versa). No interaction occurs

when the two gaze cones do not intersect. (**B**) Histograms of gaze count as a function of

inter-animal distance, shown separately for familiar and unfamiliar pairs. (**C**) Same data

as in B shown as pie charts of percentages in social gaze states. (**D**) Gaze state transition

diagrams for familiar and unfamiliar pairs. The nodes are the gaze states and the edges

connecting the nodes represent the transition between states. Edge colors indicate

transition probabilities. (**E**) Partner gaze self-transition probabilities for familiar and

unfamiliar pairs (χ^2 test). (**F**) Delta transition matrix between the unfamiliar pair and

familiar pair state transition diagrams. Transitions that are significantly different across

familiarity are marked by asterisks (χ^2 test, male to male, p < 0.05; female to male, p <

0.05; joint to joint, p < 0.0001). (**G**) Left, The schematic illustrates how gazing toward the

surrounding region of a partner's face area was measured. Right, Counts of gaze towards

the surrounding region of the partner's face by familiarity and sex. (Mann-Whitney U test,

*** means p < 0.001; **** means p < 0.0001)

Supplementary Figure 1. **Gaze behavior analysis of a single marmoset viewing**

stimuli on the monitor.

(**A**) Gaze duration for video stimuli is significantly higher compared to image stimuli (Mann

Whitney U Test, p < 0.001). (**B**) Left, Gaze dispersion is defined as the average distance

between the centers of the intersection of the gaze cone and the monitor within a gaze

epoch. There was no difference between the video and image stimuli for gaze dispersion

(Mann Whitney U Test, ns). (**C**) Left, Illustration of four gaze epochs (gray bars) to

repeated presentations of a stimulus and the gaze counts at different time points within

the duration of the presentation. Marmosets have more gazes in the early period than the

late period for the video stimuli (Mann Whitney U Test, p < 0.001), however, this is not the

case for the image stimuli (Mann Whitney U Test, ns). During the early period, marmosets

had significantly higher gaze counts for the video stimuli than the image stimuli (Mann

Whitney U Test, $p < 10^{-10}$).

Supplementary Video 1. Results from different processing stages of the facial features

detection pipeline are shown for two marmosets in their home cage.

Supplementary Video 2. 3D reconstruction of the face frame and the inferred gaze cone

across time for a single marmoset.

Supplementary Video 3. Categorization of gaze behavior epochs of two freely viewing

marmosets and transitions between the defined gaze states.

Methods

Camera calibration

All cameras (GoPro 8) were calibrated using an 8-by-9 black-white checkerboard. For each camera, the checkerboard was placed at various locations to sample the space of the camera's field of view. To achieve better calibration performance, the checkerboard was tilted and rotated to varying degrees thus producing a range of different views (Zhang, 2000). The corners of the checkerboard were automatically detected via a standard algorithm (detectCheckerboardPoints() function in the Image Processing and Computer Vision toolbox in MATLAB). The intrinsic parameters of each camera were estimated based on the data obtained from the checkerboard corner detection algorithm (estimateCameraParameters() function in Image Processing and Computer Vision toolbox in MATLAB).

L-frame analysis

L-shaped frames were used to obtain the extrinsic parameters of the cameras, the rotation matrix, and the transition vector (Timothy et al., 2021). The L-shaped frame was captured by four cameras that recorded one arena. Four points that were unevenly distributed on the L-shaped frame were manually labeled. The information of transformation from world coordinates to camera coordinates was then extracted based on the labeled result (cameraPoseToExtrinsics() function in Image Processing and Computer Vision toolbox in MATLAB).

Camera recording

GoPro 8 cameras were used and were simultaneously controlled via a Bluetooth remote control (The Remote by GoPro). Videos were recorded at 30 frames/sec with a linear lens. Frame resolution was set at 1920x1080 pixels. A circular polarizer filter was used to mitigate reflection artifacts.

Deep convolutional neural network (DCNN) model training

We used a modified version of DeepLabCut (Mathis et al., 2018) to perform automated markerless tracking of body parts of interest from two marmosets. The model was trained on 700 hand-labeled image frames extracted from videos of animals in their colony

settings. Each image frame was labeled with six facial points: the two tufts, the central

blaze, two eyes, and the mouth. The model was trained using GPUs on a large computing

cluster for 250,000 iterations until the loss reached a plateau.

Gaze cone calculation

At each time frame, the gaze orientation was calculated as the normal to the facial plane

('norm') defined by the two eyes and the central blaze. The position of the ear tufts, which

were behind this facial plane, was used to determine the direction of gaze. A gaze cone

was defined as a virtual cone of 10 degrees solid angle around this norm.

Head gaze velocity calculation and stable epoch identification

We used the change of the norm over consecutive time frames to calculate the head gaze

velocity:

$$v(t) = \frac{N(t+2) + N(t+1) - N(t-1) - N(t-2)}{6}$$

563
564 where v(t) is the velocity at time point t, N(t) is the norm at time point t.

566 We remove all time points where the head gaze velocity was larger than 0.1 in normalized

567 units. Segments no shorter than three consecutive time frames were identified as stable

568 epochs.

569
570 **Cone-monitor plane intersection**

571 We modified an existing method (Calinon & Billard, 2006; Sylvain, 2009) to determine the

572 elliptical intersection of a gaze cone and the finite plane defined by the monitor.

573
574 **Cone-facial plane intersection**

575 We used a numerical method to determine whether the gaze cone of one animal

576 intersected with the facial plane of the other. The facial plane was defined as the finite

577 triangular plane formed by three facial features: two eyes and mouth. Any point X in 3D

578 within the volume bounded by the cone satisfies the inequality:

$$\cos\theta - \frac{dot(coneDir,\ X - coneOrg)}{norm(X - coneOrg)} \leq 0$$

where θ is the solid angle of the gaze cone, *coneDir* is the direction vector of the gaze cone, *coneOrg* is the origin point of the gaze cone. The facial plane intersects with the cone if any point within the finite plane satisfies the inequality.

Cone-cone intersection

To calculate the cone-cone intersection, we used the same numerical method as above. If any point X in 3D simultaneously satisfied the following inequalities:

$$\cos\theta_1 - \frac{dot(coneDir_1,\ X - coneOrg_1)}{norm(X - coneOrg_1)} \leq 0$$

and

$$\cos\theta_2 - \frac{dot(coneDir_2,\ X - coneOrg_2)}{norm(X - coneOrg_2)} \leq 0$$

then the two cones were considered to be intersected. Subscripts in the above inequalities indicate the parameters of the two gaze cones under consideration.

Maximum likelihood estimation

We used the a maximum likelihood estimation method (mle() function in the Statistics and Machine Learning Toolbox in MATLAB) to fit a mixture of Gamma and Gaussian distributions.

Markov chain analysis

State transition matrices were obtained based on the behavioral data. These matrices were then used to generate the discrete Markov chains (dtmc() function in Econometrics Toolbox in MATLAB) and plotted (graphplot() function in MATLAB).

Warren Sarle's bimodality coefficient

612　Sarle's bimodality coefficient was used to test for bimodality. The coefficient was
613　calculated using publicly available MATLAB code based on the theory in Pfister et al.,
614　2013 .

616　# Experimental model and subject details

617
Animals

Nine adult marmosets were used in this study (four males, five females). Four familiar
male/female pairs were each from the same cage. Four unfamiliar male/female pairs were
selected from the nine animals such that each member of a pair were from different home
cages and did not have visual access to each other while in the colony. Animals were
kept in a colony maintained at around 75°F, 60% humidity and a 12h:12h light-dark cycle.

624
Single marmoset gazing at the monitor

A single freely moving marmoset was recorded by four cameras surrounding the arena.
Video or image stimuli were displayed at one of five locations (Center, Up, Down, Left and
Right) on the monitor (location chosen randomly). Each session contained only one
stimulus category (either video or image) and consisted of five blocks. Each block
consisted of ten five-second stimuli interleaved with ten five-second breaks. Each block
started with a white dot in the center of the screen on a black background lasting for one
second. At the end of the block, a juice reward (diluted condensed milk, condensed milk :
water = 1:7) was delivered with a syringe pump system (NE-500 programmable OEM
syringe pump from Pump Systems Inc.) along with an auditory cue.

635
Freely interacting marmoset dyads

Two freely moving marmosets, in separate arenas, were recorded by two sets of four
cameras surrounding the arenas. Each session consisted of ten five-minute free-viewing
blocks interleaved with nine five-minute breaks. A juice reward (diluted condensed milk,
condensed milk : water = 1:7) was delivered every minute through two syringe pump
systems during the free-viewing blocks. During the breaks, a divider was placed between
the two arenas that prevented the marmosets from seeing each other.

References

Anderson, David J., & Perona, P. (2014). Toward a Science of Computational Ethology. *Neuron, 84*(1), 18-31. doi:https://doi.org/10.1016/j.neuron.2014.09.005

Berman, G. J., Choi, D. M., Bialek, W., & Shaevitz, J. W. (2014). Mapping the stereotyped behaviour of freely moving fruit flies. *Journal of The Royal Society Interface, 11*(99), 20140672.

Brooks, R., & Meltzoff, A. N. (2005). The development of gaze following and its relation to language. *Dev Sci, 8*(6), 535-543. doi:10.1111/j.1467-7687.2005.00445.x

Burkart, J., & Heschl, A. (2006). Geometrical gaze following in common marmosets (Callithrix jacchus). *J Comp Psychol, 120*(2), 120-130. doi:10.1037/0735-7036.120.2.120

Cao, Z., Simon, T., Wei, S.-E., & Sheikh, Y. (2017). *Realtime multi-person 2d pose estimation using part affinity fields.* Paper presented at the Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition.

Chevallier, C., Parish-Morris, J., McVey, A., Rump, K. M., Sasson, N. J., Herrington, J. D., & Schultz, R. T. (2015). Measuring social attention and motivation in autism spectrum disorder using eye-tracking: Stimulus type matters. *Autism Res, 8*(5), 620-628. doi:10.1002/aur.1479

Dal Monte, O., Costa, V. D., Noble, P. L., Murray, E. A., & Averbeck, B. B. (2015). Amygdala lesions in rhesus macaques decrease attention to threat. *Nature Communications, 6*(1), 10161. doi:10.1038/ncomms10161

Dal Monte, O., Fan, S., Fagan, N. A., Chu, C. J., Zhou, M. B., Putnam, P. T., . . . Chang, S. W. C. (2022). Widespread implementations of interactive social gaze neurons in the primate prefrontal-amygdala networks. *Neuron, 110*(13), 2183-2197.e2187. doi:10.1016/j.neuron.2022.04.013

Dal Monte, O., Piva, M., Morris, J. A., & Chang, S. W. (2016). Live interaction distinctively shapes social gaze dynamics in rhesus macaques. *J Neurophysiol, 116*(4), 1626-1643. doi:10.1152/jn.00442.2016

Datta, S. R., Anderson, D. J., Branson, K., Perona, P., & Leifer, A. (2019). Computational Neuroethology: A Call to Action. *Neuron, 104*(1), 11-24. doi:https://doi.org/10.1016/j.neuron.2019.09.038

Deen, B., Schwiedrzik, C. M., Sliwa, J., & Freiwald, W. A. (2023). Specialized Networks for Social Cognition in the Primate Brain. *Annual Review of Neuroscience, 46*(1), 381-401. doi:10.1146/annurev-neuro-102522-121410

Digby, L. J. (1995). Social organization in a wild population ofCallithrix jacchus: II. Intragroup social behavior. *Primates, 36*(3), 361-375. doi:10.1007/BF02382859

Emery, N. J. (2000). The eyes have it: the neuroethology, function and evolution of social gaze. *Neurosci Biobehav Rev, 24*(6), 581-604. doi:10.1016/s0149-7634(00)00025-7

Emery, N. J., Lorincz, E. N., Perrett, D. I., Oram, M. W., & Baker, C. I. (1997). Gaze following and joint attention in rhesus monkeys (Macaca mulatta). *J Comp Psychol, 111*(3), 286-293. doi:10.1037/0735-7036.111.3.286

French, J. A. (1997). Proximate regulation of singular breeding in callitrichid primates. *Cooperative breeding in mammals*, 34-75.

Furl, N., Hadj-Bouziane, F., Liu, N., Averbeck, B. B., & Ungerleider, L. G. (2012). Dynamic and static facial expressions decoded from motion-sensitive areas in the macaque monkey. *J Neurosci, 32*(45), 15952-15962. doi:10.1523/jneurosci.1992-12.2012

Hari, R., Henriksson, L., Malinen, S., & Parkkonen, L. (2015). Centrality of Social Interaction in Human Brain Function. *Neuron, 88*(1), 181-193. doi:https://doi.org/10.1016/j.neuron.2015.09.022

Heiney, S. A., & Blazquez, P. M. (2011). Behavioral responses of trained squirrel and rhesus monkeys during oculomotor tasks. *Exp Brain Res, 212*(3), 409-416. doi:10.1007/s00221-011-2746-4

Hesse, J. K., & Tsao, D. Y. (2020). A new no-report paradigm reveals that face cells encode both consciously perceived and suppressed stimuli. *eLife, 9*, e58360. doi:10.7554/eLife.58360

Itier, R. J., Alain, C., Sedore, K., & McIntosh, A. R. (2007). Early face processing specificity: it's in the eyes! *J Cogn Neurosci, 19*(11), 1815-1826. doi:10.1162/jocn.2007.19.11.1815

Marshall, J. D., Klibaite, U., Gellis, A., Aldarondo, D. E., Ölveczky, B. P., & Dunn, T. W. (2021). The PAIR-R24M Dataset for Multi-animal 3D Pose Estimation. *bioRxiv*, 2021.2011.2023.469743. doi:10.1101/2021.11.23.469743

Martin, A., & Santos, L. R. (2016). What Cognitive Representations Support Primate Theory of Mind? *Trends Cogn Sci, 20*(5), 375-382. doi:10.1016/j.tics.2016.03.005

Mathis, A., Mamidanna, P., Cury, K. M., Abe, T., Murthy, V. N., Mathis, M. W., & Bethge, M. (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. *Nature neuroscience, 21*(9), 1281-1289.

Miller, C. T., Freiwald, W. A., Leopold, D. A., Mitchell, J. F., Silva, A. C., & Wang, X. (2016). Marmosets: A Neuroscientific Model of Human Social Behavior. *Neuron, 90*(2), 219-233. doi:10.1016/j.neuron.2016.03.018

Miller, C. T., Gire, D., Hoke, K., Huk, A. C., Kelley, D., Leopold, D. A., . . . Niell, C. M. (2022). Natural behavior is the language of the brain. *Curr Biol, 32*(10), R482-r493. doi:10.1016/j.cub.2022.03.031

Miss, F. M., & Burkart, J. M. (2018). Corepresentation during joint action in marmoset monkeys (Callithrix jacchus). *Psychological Science, 29*(6), 984-995.

Mitchell, J. F., Reynolds, J. H., & Miller, C. T. (2014). Active vision in marmosets: a model system for visual neuroscience. *J Neurosci, 34*(4), 1183-1194. doi:10.1523/jneurosci.3899-13.2014

Mosher, C. P., Zimmerman, P. E., & Gothard, K. M. (2014). Neurons in the monkey amygdala detect eye contact during naturalistic social interactions. *Curr Biol, 24*(20), 2459-2464. doi:10.1016/j.cub.2014.08.063

Mustoe, A. (2023). A tale of two hierarchies: Hormonal and behavioral factors underlying sex differences in social dominance in cooperative breeding callitrichids. *Horm Behav, 147*, 105293. doi:10.1016/j.yhbeh.2022.105293

Ngo, V., Gorman, J. C., De la Fuente, M. F., Souto, A., Schiel, N., & Miller, C. T. (2022). Active vision during prey capture in wild marmoset monkeys. *Curr Biol, 32*(15), 3423-3428.e3423. doi:10.1016/j.cub.2022.06.028

Pandey, S., Simhadri, S., & Zhou, Y. (2020). Rapid Head Movements in Common Marmoset Monkeys. *iScience, 23*(2), 100837. doi:10.1016/j.isci.2020.100837

Pereira, T. D., Tabris, N., Matsliah, A., Turner, D. M., Li, J., Ravindranath, S., . . . Murthy,
M. (2022). SLEAP: A deep learning system for multi-animal pose tracking. *Nature*
Methods, 19(4), 486-495. doi:10.1038/s41592-022-01426-1
Pfister, R., Schwarz, K. A., Janczyk, M., Dale, R., & Freeman, J. (2013). Good things
peak in pairs: a note on the bimodality coefficient. *Frontiers in Psychology, 4*.
doi:10.3389/fpsyg.2013.00700
Ramezanpour, H., & Thier, P. (2020). Decoding of the other's focus of attention by a
temporal cortex module. *Proc Natl Acad Sci U S A, 117*(5), 2663-2670.
doi:10.1073/pnas.1911269117
Saxe, R., & Kanwisher, N. (2003). People thinking about thinking people. The role of the
temporo-parietal junction in "theory of mind". *NeuroImage, 19*(4), 1835-1842.
doi:10.1016/s1053-8119(03)00230-1
Shepherd, S. (2010). Following Gaze: Gaze-Following Behavior as a Window into
Social Cognition. *Frontiers in Integrative Neuroscience, 4*.
doi:10.3389/fnint.2010.00005
Shepherd, S. V., Deaner, R. O., & Platt, M. L. (2006). Social status gates social attention
in monkeys. *Curr Biol, 16*(4), R119-120. doi:10.1016/j.cub.2006.02.013
Shepherd, S. V., & Freiwald, W. A. (2018). Functional Networks for Social
Communication in the Macaque Monkey. *Neuron, 99*(2), 413-420.e413.
doi:10.1016/j.neuron.2018.06.027
Solomon, N. G., & French, J. A. (1997). *Cooperative breeding in mammals*: Cambridge
University Press.
Spadacenta, S., Dicke, P. W., & Thier, P. (2022). A prosocial function of head-gaze
aversion and head-cocking in common marmosets. *Primates, 63*(5), 535-546.
doi:10.1007/s10329-022-00997-z
Yamamoto, M. E., Araujo, A., Arruda, M. d. F., Lima, A. K. M., Siqueira, J. d. O., &
Hattori, W. T. (2014). Male and female breeding strategies in a cooperative
primate. *Behavioural Processes, 109*, 27-33.
doi:https://doi.org/10.1016/j.beproc.2014.06.009
764

Figure 1. **Experimental setup and reconstruction in 3D space**.
(A) Two transparent acrylic arenas allowed marmosets to visually interact with each other. An opaque divider between the arenas was introduced intermittently to prevent visual access between animals. Two monitors on two ends were used to display video or image stimuli. Eight cameras surrounding the arenas ensured full coverage of both animals. LEDs at four positions were used to synchronize the video recordings across the set of cameras. (B) 3D reconstruction of the experimental setup. The two arenas are color-coded as orange and cyan. The cameras colored the same as the arenas indicate that they primarily record the marmoset in the corresponding arena. Two purple L-frames within the arenas were used to establish a world coordinate system in the reconstruction process. Two gray planes on both ends are the reconstructed monitors.

Figure 2. Pipeline of detecting facial features of two marmosets.
(A) Six facial features of the marmoset (face frame) are color-coded: right tuft (red), central blaze (yellow), left tuft (green), right eye (purple), left eye (blue), and mouth (magenta). (B) Feature tracking pipeline (right) with the corresponding illustration for each step across two adjacent video frames (left). At the end of the pipeline, the facial features are clustered with the identities assigned consistently across frames. Facial points are color-coded as in A. (C) Example frames of four steps in the pipeline shown in B. It can be seen clearly that the facial points are tracked and clustered accurately, and the identities are consistent across frames.

Figure 3. **3D Reconstruction of facial features and head gaze modeling**.
(A) The face frames of two marmosets are reconstructed in 3D using the tracked facial points in Fig. 2. A cone perpendicular to the face frame (gaze cone; 10-degree solid angle) is modeled as the head gaze. (B) Two example frames with the facial points projected from 3D space onto different camera views are shown. The left frame demonstrates that the facial points can be detected using information from other cameras, even if the face is invisible from that viewpoint. (C) Trajectory of the reconstructed face frame and the corresponding gaze cones across time.

Figure 4. **Positional dynamics of marmoset dyads**.

(A) Movement trajectories of the face frame centroids for a marmoset pair (orange for female, cyan for male) in an example five-minute block. The heatmaps were calculated using the projections of the trajectories to XY, YZ, and XZ planes. (B) Marginal distributions of movement trajectories along the X and Z axes were calculated for all marmosets and grouped by familiarity and sex (transparent colors for familiar pairs, opaque colors for unfamiliar pairs). Black bars indicate significant differences between pairs of distributions (Mann-Whitney U test, significance level at 5%). (C) Histograms of inter-animal distance along the X axis show trimodal distributions for both familiar pairs (gray) and unfamiliar pairs (black). The fitted red curve for the unfamiliar pairs is a tri-Gaussian distribution, while the fitted red curve for the familiar pairs is a mixture of Gamma and Gaussian distributions, with the first peak as the Gamma distribution. The three regions were designated as 'Near', Intermediate', and 'Far'. The inset illustrates the reason for this nomenclature. (D) Temporal evolution of inter-animal distance for unfamiliar and familiar pairs (which each 5-minute viewing block). The central dark line is the mean and the shaded area is the standard deviation. Black dots indicate significant differences (Mann-Whitney U test, significance level at 5%).

Figure 5. Live interactive gaze analysis of unfamiliar and familiar marmoset dyads.
(A) Gaze type categorized based on the relative positions of the gaze cones. Joint gaze is defined as two marmosets looking at the same location. A partner gaze is defined as one animal looking at the other animal's face (but not vice versa). No interaction occurs when the two gaze cones do not intersect. (B) Histograms of gaze count as a function of inter-animal distance, shown separately for familiar and unfamiliar pairs. (C) Same data as in B shown as pie charts of percentages in social gaze states. (D) Gaze state transition diagrams for familiar and unfamiliar pairs. The nodes are the gaze states and the edges connecting the nodes represent the transition between states. Edge colors indicate transition probabilities. (E) Partner gaze self-transition probabilities for familiar and unfamiliar pairs ($\chi 2$ test). (F) Delta transition matrix between the unfamiliar pair and familiar pair state transition diagrams. Transitions that are significantly different across familiarity are marked by asterisks ($\chi 2$ test, male to male, p < 0.05; female to male, p < 0.05; joint to joint, p < 0.0001). (G) Left, The schematic illustrates how gazing toward the surrounding region of a partner's face area was measured. Right, Counts of gaze towards the surrounding region of the partner's face by familiarity and sex. (Mann-Whitney U test, *** means p < 0.001; **** means p < 0.0001)

Supplementary Figure 1. **Gaze behavior analysis of a single marmoset viewing stimuli on the monitor**. (A) Gaze duration for video stimuli is significantly higher compared to image stimuli (Mann Whitney U Test, p < 0.001). (B) Left, Gaze dispersion is defined as the average distance between the centers of the intersection of the gaze cone and the monitor within a gaze epoch. There was no difference between the video and image stimuli for gaze dispersion (Mann Whitney U Test, ns). (C) Left, Illustration of four gaze epochs (gray bars) to repeated presentations of a stimulus and the gaze counts at different time points within the duration of the presentation. Marmosets have more gazes in the early period than the late period for the video stimuli (Mann Whitney U Test, p < 0.001), however, this is not the case for the image stimuli (Mann Whitney U Test, ns). During the early period, marmosets had significantly higher gaze counts for the video stimuli than the image stimuli (Mann Whitney U Test, p < 10^-10).