A genome-wide arrayed CRISPR screen identifies PLSCR1 as an intrinsic barrier

to SARS-CoV-2 entry that recent virus variants have evolved to resist

Authors

Jérémie Le Pen[1,*], Gabrielle Paniccia[1,*], Volker Kinast[2,3,#], Marcela Moncada-Velez[4,#], Alison
W. Ashbrook[1,#], Michael Bauer[1,#], H.-Heinrich Hoffmann[1,#], Ana Pinharanda[5,#], Inna Ricardo-Lax[1,#],
Ansgar F. Stenzel[1,6,#], Edwin A. Rosado-Olivieri[7,8,#], Kenneth H. Dinnon III[1], William C. Doyle[1,9],
Catherine A. Freije[1], Seon-Hui Hong[1], Danyel Lee[4,10,11], Tyler Lewy[1], Joseph M. Luna[1,12], Avery
Peace[1], Carltin Schmidt[4,13], William M. Schneider[1], Roni Winkler[1], Elaine Z. Yip[1], Chloe Larson[14],
Timothy McGinn[14], Miriam-Rose Menezes[14], Lavoisier Ramos-Espiritu[14], Priyam Banerjee[15], John
T. Poirier[16], Francisco J. Sànchez-Rivera[17,18], Aurélie Cobat[4,10,11], Qian Zhang[4,10,11], Jean-Laurent
Casanova[4,10,11,19,20], Thomas S. Carroll[21], J. Fraser Glickman[14], Eleftherios Michailidis[1,22,§], Brandon
Razooky[1,23,§], Margaret R. MacDonald[1,§], Charles M. Rice[1,§]

[*] Contributed equally to this work
[#] Contributed equally to this work
[§] Contributed equally to this work

Correspondence: Jérémie Le Pen: jlepen@rockefeller.edu; Charles M. Rice:
ricec@rockefeller.edu

Affiliations:
1. Laboratory of Virology and Infectious Disease, The Rockefeller University, New York, NY,
USA.
2. Department of Medical Microbiology and Virology, Carl von Ossietzky University
Oldenburg, Oldenburg, Germany.
3. Department for Molecular and Medical Virology, Faculty of Medicine, Ruhr University
Bochum, Bochum, Germany.

4. St Giles Laboratory of Human Genetics of Infectious Diseases, Rockefeller Branch, The

Rockefeller University, New York, NY, USA.

5. Department of Biological Sciences, Columbia University, New York, NY 10027, USA.

6. Present address: Department of Infectious Diseases, Molecular Virology, Heidelberg

University, Heidelberg, Germany.

7. Laboratory of Synthetic Embryology, The Rockefeller University, New York, NY, USA.

8. Present address: Department of Molecular Pathobiology, New York University.

9. Present address: Laboratory of Host-Pathogen Biology, The Rockefeller University, New

York, NY, USA.

10. Laboratory of Human Genetics of Infectious Diseases, Necker Branch, INSERM U1163,

Paris, France, EU.

11. Paris City University, Imagine Institute, Paris, France, EU.

12. Present address: Department of Biochemistry and Center for RNA Science and

Therapeutics, Case Western Reserve University, Cleveland, OH, USA.

13. Present address: Medical Faculty, Heinrich Heine University Dusseldorf, Germany.

14. Fisher Drug Discovery Resource Center, The Rockefeller University, New York, NY, USA.

15. Bio-Imaging Resource Center, The Rockefeller University, New York, NY, USA.

16. Laura and Isaac Perlmutter Cancer Center, New York University Grossman School of

Medicine, NYU Langone Health, New York, NY.

17. Department of Biology, Massachusetts Institute of Technology, Cambridge, 02142,

Massachusetts, USA.

18. David H. Koch Institute for Integrative Cancer Research, Massachusetts Institute of

Technology, Cambridge, 02142, Massachusetts, USA.

19. Department of Pediatrics, Necker Hospital for Sick Children, Paris, France, EU.

20. Howard Hughes Medical Institute, New York, NY, USA.

21. Bioinformatics Resource Center, The Rockefeller University, New York, NY, USA.

22. Present address: Laboratory of Biochemical Pharmacology, Department of Pediatrics,

Emory University School of Medicine, Atlanta, GA, USA.

23. Present address: National Resilience, Inc., La Jolla, CA, USA.

Abstract

Interferons (IFNs) play a crucial role in the regulation and evolution of host-virus interactions. Here, we conducted a genome-wide arrayed CRISPR knockout screen in the presence and absence of IFN to identify human genes that influence SARS-CoV-2 infection. We then performed an integrated analysis of genes interacting with SARS-CoV-2, drawing from a selection of 67 large-scale studies, including our own. We identified 28 genes of high relevance in both human genetic studies of COVID-19 patients and functional genetic screens in cell culture, with many related to the IFN pathway. Among these was the IFN-stimulated gene *PLSCR1*. PLSCR1 did not require IFN induction to restrict SARS-CoV-2 and did not contribute to IFN signaling. Instead, PLSCR1 specifically restricted spike-mediated SARS-CoV-2 entry. The PLSCR1-mediated restriction was alleviated by TMPRSS2 over-expression, suggesting that PLSCR1 primarily restricts the endocytic entry route. In addition, recent SARS-CoV-2 variants have adapted to circumvent the PLSCR1 barrier via currently undetermined mechanisms. Finally, we investigate the functional effects of PLSCR1 variants present in humans and discuss an association between PLSCR1 and severe COVID-19 reported recently.

Introduction

Viruses maintain a complex relationship with their host cells, co-opting host factors for their replication while being targeted by cellular defense mechanisms. Such cellular defenses include the interferon (IFN) pathway, where the infected cell senses foreign molecules and secretes IFN to trigger an antiviral state in neighboring cells [1].

Approximately 1-5% of critical COVID-19 patients have mutations that compromise the production of or response to type I IFNs, while an additional 15% possess autoantibodies that neutralize type I IFNs [2-8]. This highlights the essential role of type I IFN in the defense against the SARS-CoV-2 virus that caused the COVID-19 pandemic [9, 10]. Consequently, investigating IFN-stimulated genes (ISGs) is crucial to our understanding of the remarkable antiviral systems that evolved in nature and could enhance our preparedness for future pandemics.

Several recent studies have identified ISGs restricting SARS-CoV-2. Most of these studies involved gain-of-function genetic screens, over-expressing individual ISGs. The factors bone

marrow stromal cell antigen 2 (BST2), cholesterol 25-hydroxylase (CH25H), lymphocyte antigen 6

family member E (LY6E), 2'-5'-oligoadenylate synthetase 1 (OAS1), and receptor transporter

protein 4 (RTP4) were notably identified as SARS-CoV-2 antivirals in these studies [10-15]. One

advantage of the gain-of-function approach is that it circumvents potential genetic redundancies

between ISGs [16, 17]. However, this approach is biased towards ISGs that act autonomously

when over-expressed and does not mimic the cellular context of the IFN response, where

hundreds of genes and gene products are differentially regulated to establish an antiviral state.

To counter this limitation, two recent publications examined the effects of ISG loss of function in

IFN-treated cells. They conducted pooled CRISPR knockout (KO) screens in cells pre-treated with

IFN before SARS-CoV-2 infection [18, 19]. By sorting for cells with high SARS-CoV-2 viral load, they

identified SARS-CoV-2 restriction factors such as death domain associated protein (DAXX).

Here, we conducted a human whole-genome arrayed CRISPR KO screen to identify genes

that influence SARS-CoV-2 infection in cells with or without pretreatment with a low dose of IFN.

The arrayed approach, though logistically challenging, has advantages over the pooled format in

capturing both proviral and antiviral genes, genes affecting virus egress, and those coding for

secreted products that exert their impact on neighboring cells. It reliably captures genotype-

phenotype correlations while also unveiling the effects of single gene perturbation on cell growth

and death [20]. Additionally, the shorter culture time and lack of competition among cells with

different gene KO in the arrayed screen allow the inclusion of genes that would be depleted and

deemed essential in a pooled format [21]. The arrayed format thus enables the identification of

crucial cellular functions that may be co-opted by the virus or are vital for the cell's defense

against infection.

We then compiled a comprehensive list of genes interacting with SARS-CoV-2,

incorporating findings from our own screen as well as existing literature. This meta-analysis

revealed several host genes of interest, both previously described and novel. Notably, the ISG

product phospholipid scramblase 1 (PLSCR1) emerged as a prominent antiviral factor. PLSCR1 is

involved in several biological processes [22], including regulating the movement of phospholipids

between the two leaflets of a cell membrane (lipid scrambling) [23] and IFN signaling in the

context of virus infection [24].

Follow-up experiments revealed that PLSCR1 is a cell intrinsic factor that restricts spike-

mediated SARS-CoV-2 entry, independently of the IFN pathway, via currently undetermined

mechanisms. Our genetic screen data and meta-analysis provide a valuable resource to broaden

our understanding of coronavirus infection and innate immunity. Furthermore, we extend the

recent characterization of PLSCR1 as an antiviral against SARS-CoV-2 impacting COVID-19

outcomes (**Supp Fig. 1**)[19, 25, 26].

Results

A genome-wide arrayed CRISPR KO screen identifies known and novel factors

influencing SARS-CoV-2 infection.

While the liver is not the primary target organ of SARS-CoV-2 infection, human

hepatocellular carcinoma Huh-7.5 cells naturally express SARS-CoV-2 dependency factors,

including the receptor angiotensin converting enzyme 2 (ACE2), and proved unexpectedly useful

in SARS-CoV-2 research [19, 27-34]. Huh-7.5 cells are defective in virus sensing and do not

commonly produce IFN during infection [35]. We confirmed that Huh-7.5 cells do not induce ISG

expression during SARS-CoV-2 infection (**Fig 1A, Supp Tables 2 and 3**). This is likely due to a defect

upstream of IFN production, as these cells did induce ISG expression when treated with

recombinant IFN (**Fig 1B, Supp Tables 4 and 5**). Thus, Huh-7.5 cells are a convenient model for

studying controlled IFN responses during viral infection. Furthermore, IFN treatment restricted

SARS-CoV-2 (**Fig 1C**), indicating that some ISGs are effective in limiting SARS-CoV-2 infection in

Huh-7.5 cells. This allows us to study the functional landscape of SARS-CoV-2 restriction,

examining both intrinsic factors and those induced in response to IFN signaling. Similar to A549-

ACE2 cells [11, 36], Huh-7.5 cells do not express transmembrane serine protease 2 (TMPRSS2)

(**Supp Fig 1B**). As a result, SARS-CoV-2 entry is restricted to the endocytic route [37, 38].

Using these cells, we conducted a whole-genome arrayed CRISPR KO screen designed to

identify both SARS-CoV-2 proviral and antiviral genes, whose KO reduces or enhances SARS-CoV-

2 infection, respectively. In particular, we aimed to identify factors involved in the IFN response,

from IFN sensing to ISG induction, including effector ISGs that directly influence the virus life

cycle. With this arrayed approach performed in 384-well plates, the cells in each well received a

pool of four gRNAs targeting a single host gene. Each well was then either treated with 1 pM IFN-

α2a or left untreated before being infected with SARS-CoV-2, followed by SARS-CoV-2

nucleoprotein (N) immunofluorescence staining and high content microscopy (**Fig 1D**). This low

dose of IFN was chosen to mimic a cellular environment where IFN triggers an antiviral state

before infection. We anticipated that a saturating amount of IFN would lead to high ISG

transcription and functional redundancy between effectors, biasing hits towards factors in IFN

signaling. In contrast, a low dose of IFN, around the IC50, might enable identification of the

specific roles of individual effector ISGs.

We conducted the functional assays four days after KO to allow sufficient time for full

protein depletion, considering both cell division and protein half-life (**Supp Fig 2E**).

We included control gRNAs with known proviral and antiviral effects in our screening

plates. As expected, the SARS-CoV-2 receptor angiotensin-converting enzyme 2 (ACE2) [9]

behaved as a proviral gene with and without IFN pretreatment. The interferon-alpha/beta

receptor alpha chain 1 (IFNAR1) [39-41] was essential for the IFN-mediated restriction of SARS-

CoV-2. The ISG15 ubiquitin-like modifier (ISG15) has been previously described as a negative

regulator of the IFN response. ISG15 stabilizes ubiquitin-specific peptidase 18 (USP18) [42-44],

which in turn binds to the interferon-alpha/beta receptor alpha chain 2 (IFNAR2) and blocks signal

transduction [45, 46]. Accordingly, ISG15 was proviral in IFN-treated cells (**Fig 1E**).

Of the 16,790 screened genes, we selected 16,178 genes where KO did not lead to changes

in cellular fitness, as assessed by nuclei count (-2 ≤ z-score ≤ 2) (**Supp Fig 2B**). Of these, we

selected 12,119 genes expressed in three cell lines relevant for SARS-CoV-2 research (A549, Calu-

3, Huh-7.5 cells) and human lung cells, the primary target cell type *in vivo*, for downstream

analysis [47-49](Supp Table 6). We then binned the genes into two groups for data visualization,

depending on whether they were induced by IFN-α2a treatment in Huh-7.5 cells as determined

by mRNA-seq (log 2 FC ≥ 2 and padj ≤ 0.05) (**Supp Fig 2C, Supp Tables 7-9**).

We classified 448 genes with a SARS-CoV-2 infection z-score ≥ 2 as antiviral hits and 507

genes with a SARS-CoV-2 infection z-score ≤ -2 as proviral hits (**Supp Table 9**).

Our screen found known and previously unidentified host factors influencing SARS-CoV-2

infection (**Fig 1F**). As expected, positive regulators of IFN signaling, such as IFNAR1,2 [39-41],

interferon regulatory factor 9 (IRF9) [50, 51], Janus kinase 1 (JAK1) [52], and signal transducer and activator of transcription 2 (STAT2) [53, 54] were antiviral only in IFN pretreated cells. Known negative regulators of IFN signaling, such as ISG15 [42-44], suppressor of cytokine signaling 1 (SOCS1) [55], and USP18 [45, 46] were proviral only in IFN pretreated cells.

The receptor ACE2 was required for infection [9]. In our mRNA-seq analysis, IFN treatment was found to significantly upregulate ACE2 mRNA levels (**Supp Fig. 2C**). Prior studies indicate that IFN induces transcription of a truncated ACE2 isoform, rather than the full-length receptor for SARS-CoV-2 [56, 57]. The lysosomal cysteine protease cathepsin L (CTSL), required for SARS-CoV-2 spike protein activation during endocytosis [58-60], was also a proviral hit in our screens (**Fig 1F**).

The screen data likely contains false negatives. For example, STAT1 and tyrosine kinase 2 (TYK2) [61, 62] did not influence infection alongside other positive regulators of IFN signaling, which we attribute to the fact that some gRNAs in the library may have not efficiently directed Cas9 to cut at their respective target gene loci.

Collectively, the identification of known proviral and antiviral factors confirms the validity of our screening method.

We performed a Gene Set Enrichment Analysis (GSEA) to identify cellular pathways exhibiting proviral or antiviral properties in our screen. The full GSEA results, including the genes driving each pathway enrichment (so-called *leading edge*), can be found in **Supp Table 10**. Some top pathways ranked by adjusted p-value are summarized in **Supp Fig 2D**. Notably pathways associated with RNA pol II transcription and mRNA maturation, as well as pathways related to cellular respiration, exhibited antiviral activity independent of IFN (**Fig 2**). Surprisingly, RNA pol III transcription, in part driven by the genes RNA polymerase III subunit A (*POLR3A*) and RNA polymerase III subunit B (*POLR3B*), were critical to the antiviral response mediated by IFN. Conversely, factors involved in translation, such as eukaryotic translation initiation factor 3 subunits F and G (EIF3G and EIF3F), likely co-opted for producing viral proteins, were identified as proviral. Similarly, factors regulating cholesterol homeostasis, likely crucial for SARS-CoV-2 entry [14, 63], were also identified as proviral. For instance, the gene sterol regulatory element binding transcription factor 2 (*SREBF2*) was one of the top proviral genes (**Fig 2**).

Our arrayed CRISPR KO screen results constitute a valuable resource for research on

coronavirus infection specifically, as well as on innate immunity in general. These can be used to

help characterize human genes influencing SARS-CoV-2 infection and the IFN response.

The ISG PLSCR1 is associated with COVID-19 outcomes and exhibits antiviral effects

in functional SARS-CoV-2 genetic screens

To provide a thorough perspective on human genes that impact SARS-CoV-2 infection and

to place our arrayed CRISPR KO screen results within the context of existing research, we have

compiled a table that includes findings from a selection of 67 large-scale 'omic' studies related to

SARS-CoV-2. This compilation encompasses this study and 25 other functional genetic screens for

genes that influence SARS-CoV-2 infection [11-13, 15, 18, 19, 28, 31, 64-80], 24 human genetic

studies that correlate certain alleles with severe COVID-19 outcomes [5, 6, 8, 25, 26, 81-99], ten

publications detailing SARS-CoV-2 protein interactomes [100-109], six focusing on SARS-CoV-2

RNA interactomes [110-115], and one that examines proteins with altered phosphorylation states

in SARS-CoV-2-infected cells [116] **(Supp Tables 11-12 for the full, and summary tables,**

respectively). This table highlights the depth of research in publications addressing SARS-CoV-2

infection: genes reported in several independent large-scale studies are more credible candidates

for biological relevance (**Supp Fig 3**). As expected, genes associated with the IFN pathway, such

as *IFNAR2*, *OAS1*, and *ZC3HAV1/ZAP*, frequently emerged as significant in SARS-CoV-2 studies.

The overlap of gene hits influencing SARS-CoV-2 between our arrayed screen and fifteen

published whole-genome pooled screens [19, 31, 64-67, 69-71, 74, 75, 77-80] was higher than

expected by chance (**Supp. Fig 4**). This finding indicates that genetic screens conducted with

different methods (e.g., CRISPR activation or CRISPR KO, pooled or arrayed format) and in various

cellular contexts (e.g., Huh-7.5, A549-ACE2, Calu-3, with or without IFN pretreatment) exhibit

both specificity and significant overlap. A pathway analysis of the hits from our arrayed screen,

alongside hits from pooled screens, is available in **Supp Table 13**.

We focused on 28 genes identified in both human genetic studies of COVID-19 patients

and in functional genetic screens in cell culture, including our own (**Fig 3**). These genes are likely

to have significant physiological relevance and to be well-suited for mechanistic studies in cell

culture. Among these, the ISG *PLSCR1* stood out, being identified as one of the most potent antiviral genes in our screen (**Fig 1F**). *PLSCR1* variants have been linked to severe COVID-19 in a recent GWAS (listed in **Table 1**)[25, 26]. This was attributed to a role of PLSCR1 in regulating the IFN response in COVID-19 patients [26]. Indeed, a pioneering study showed that PLSCR1 potentiates the transcriptional response to IFN-β treatment in human ovarian carcinoma Hey1B cells [24]. However, PLSCR1 surprisingly appeared as a potent SARS-CoV-2 antiviral even in the absence of IFN in our screens, suggesting a cell intrinsic, IFN-independent function. In other words, baseline levels of PLSCR1 may be sufficient to restrict SARS-CoV-2, and IFN pretreatment could simply enhance this effect by elevating cellular PLSCR1 levels.

Intrinsic PLSCR1 restricts SARS-CoV-2 independently of the IFN pathway.

To better characterize the function of PLSCR1 during SARS-CoV-2 infection, we generated and validated by western blot (WB) PLSCR1 KO bulk Huh-7.5 and A549-ACE2 lines (**Supp Fig 5A**). As observed in the arrayed screen (**Supp Fig 2B**), PLSCR1 KO cells were viable (**Supp Fig 5B**). PLSCR1 depletion increased susceptibility to SARS-CoV-2 independently of IFN pretreatment (**Fig 4A**). Cell treatment with a JAK-STAT inhibitor, which effectively abrogated IFN signaling, confirmed that intrinsic PLSCR1 limits SARS-CoV-2 infection independently of the IFN signaling pathway (**Fig 4A**). SARS-CoV-2 susceptibility of PLSCR1 KO cells was reversed by the ectopic expression of PLSCR1 (**Fig 4B, Supp Fig 5C**). Interestingly, while PLSCR1 tagged with an N-terminal FLAG tag could rescue, PLSCR1 tagged with a C-terminal FLAG tag could not. The C-terminus of the protein is extracellular, and previous research suggests that this region is important for the protein's scramblase activity and Ca^{2+} binding [117]. It is possible that the addition of this FLAG-tag impaired Ca^{2+} binding, affected PLSCR1's localization at the plasma membrane, or otherwise disrupted the structure of this region, thereby abolishing PLSCR1's antiviral ability. We co-cultured PLSCR1 reconstituted cells and PLSCR1 KO cells in the same well and infected them with SARS-CoV-2. A higher proportion of PLSCR1 KO than PLSCR1 reconstituted cells were positive for SARS-CoV-2 indicating that PLSCR1 acts in a cell autonomous manner (**Fig 4C**). Overall, our data indicate that intrinsic PLSCR1 restricts SARS-CoV-2 in cell culture, even in the absence of IFN. Given that

PLSCR1 mRNA is constitutively expressed in SARS-CoV-2 target cells (**Supp Fig 5D**) [118], its intrinsic antiviral function may also be effective *in vivo*.

IFN signaling is unaffected by the loss of PLSCR1 in A549-ACE2 and Huh-7.5 cells.

PLSCR1 has been shown to potentiate ISG transcription in IFN-treated Hey1B cells [24]. We thus hypothesized PLSCR1 might enhance the type I IFN response in A549-ACE2 and Huh-7.5 cells. We investigated PLSCR1's role in the IFN response by infecting Huh-7.5 cells with chikungunya virus (CHIKV), which is unaffected by PLSCR1 KO without IFN (**Supp Fig 6A**). PLSCR1 depletion did not functionally affect the antiviral effects of IFN treatment (**Supp Fig 6B**). Furthermore, IFN treatment induced *OAS1* and *IFI6*, two ISGs known to restrict SARS-CoV-2 [11, 13, 18, 95, 96, 98, 99], to a similar extent in both WT and PLSCR1 KO cells, indicating that the IFN signaling pathway was unaffected by PLSCR1 depletion (**Supp Fig 6C-J**). Finally, PLSCR1 depletion did not alter basal ISG transcription in the absence of IFN (**Supp Fig 7, Supp Tables 14 and 15**).

These findings indicate that PLSCR1 limits SARS-CoV-2 infection independently of the IFN signaling pathway in A549-ACE2 and Huh-7.5 cells.

PLSCR1 restricts SARS-CoV-2 entry.

We hypothesized that PLSCR1 directly targets and inhibits a specific step of the SARS-CoV-2 life cycle. PLSCR1 primarily localized at the plasma membrane in Huh-7.5 cells (**Fig 5A**). Furthermore, PLSCR1 depletion led to increased SARS-CoV-2 foci formation (**Fig 5B,C**), and PLSCR1 KO cells did not show increased susceptibility to a SARS-CoV-2 replicon system that bypasses entry (**Fig 5D**) [119]. In contrast, a single-cycle, replication-defective human immunodeficiency virus type-1 (HIV-1) particles pseudo-typed with SARS-CoV-2 spike showed enhanced entry in PLSCR1 depleted cells (**Fig 5E, F**) [120]. This data indicates that PLSCR1 restricts SARS-CoV-2 spike-mediated virion entry. Over-expression of TMPRSS2 lifted the PLSCR1-mediated restriction of authentic SARS-CoV-2 (**Fig 5G-I**) and of SARS-CoV-2 spike pseudo-typed particles (**Fig 5J-M**), indicating that PLSCR1 primarily restricts the endosomal entry route.

Other ISG products have been described to restrict SARS-CoV-2 entry (**Supp Fig 1**) [1], notably: CH25H promotes cholesterol sequestration in lipid droplets, decreasing the pool of

accessible cholesterol required for virus-cell membrane fusion [28,29]; LY6E blocks virus-cell

membrane fusion via currently undetermined mechanisms [121]; nuclear receptor coactivator 7

(NCOA7) over-acidifies the lysosome, leading to viral antigen degradation by lysosomal proteases

[31,32]; and interferon induced transmembrane protein 2 (IFITM2) blocks pH- and cathepsin-

dependent SARS-CoV-2 virus-cell membrane fusion in the endosome [122]. The aforementioned

ISGs were not hits in our CRISPR KO screen for antiviral genes in IFN-treated cells. This may be

due to functional redundancies among them. We found that over-expression of any of the

aforementioned ISGs restricted SARS-CoV-2 in PLSCR1 KO cells, indicating that they do not need

PLSCR1 for their antiviral activity (**Supp Fig 8A**). PLSCR1 is unlikely to require CH25H, LY6E, or

IFITM2 for its function as these are expressed at minimal levels in Huh-7.5 cells without IFN pre-

treatment (**Supp Fig 8B,C, Supp Tables 4, 5, 16, and 17**). We cannot exclude an association

between PLSCR1 and NCOA7, although the latter was not hit in our screen.

In addition to SARS-CoV-2, we also evaluated the antiviral activity of PLSCR1 against ten

viruses that utilize endosomal entry: CHIKV, human parainfluenza virus (hPIV), herpes simplex

virus 1 (HSV-1), influenza A virus (IAV), human coronavirus OC43 (hCoV-OC43), human

coronavirus NL63 (hCoV-NL63), human coronavirus 229E (hCoV-229E), Sindbis virus (SINV),

Venezuelan equine encephalitis virus (VEEV), and vesicular stomatitis virus (VSV). Only SARS-CoV-

2 showed a notable susceptibility to PLSCR1's inhibitory effects (**Supp Fig 9**).

One hypothesis for PLSCR1's specificity for SARS-CoV-2 is that it alters the surface levels

of its receptor, ACE2. However, flow cytometry on live cells did not show a significant effect of

PLSCR1 on ACE2 surface levels in A549-ACE2 cells (**Supp Fig 10**). The precise mechanism of action

of PLSCR1 remains undetermined.

Recent variants of SARS-CoV-2 are less restricted by PLSCR1

During the COVID-19 pandemic, SARS-CoV-2 variants evolved from the initial strain,

showing increased immune evasion and transmissibility [123-125]. To examine if these variants

could circumvent the antiviral action of PLSCR1, we infected WT and PLSCR1 KO Huh-7.5 cells with

an early strain isolated in July 2020 (NY-RU-NY1, subsequently referred to as "parental") and the

Beta (B.1.352), Delta (B.1.617.2), Omicron (BA.5), and Omicron (XBB.1.5) variants. PLSCR1

continued to restrict these later variants when examining the percentage of infected cells 24
hours post-infection (**Figs 6A-E**).

To determine if the magnitude of PLSCR1 restriction was the same for the parental SARS-
CoV-2 strain and recent variants, we plotted the percentage of infection data from **Figs 6A-E** as a
ratio of PLSCR1 WT to KO (**Fig 6F**). Recent SARS-CoV-2 variants showed reduced differences in
infection rates between PLSCR1 WT and KO cells than the parental SARS-CoV-2 strain. The
diminished difference in sensitivity between PLSCR1 WT and KO cells was most pronounced with
Omicron BA.5 and its descendant, XBB.1.5 (**Fig 6F**).

To examine this further, we infected PLSCR1 WT and KO Huh-7.5 cells with approximately
50 focus-forming units (FFU) per well for different SARS-CoV-2 strains (**Figs 6G and 6H**). In line
with **Fig 6F,** the data indicate that the difference in virus susceptibility between PLSCR1 WT and
KO cells is lower for more recent variants such as Beta (B.1.352) and especially Omicron (XBB.1.5)
compared with the parental strain (**Fig 6H**).

Finally, we quantified virus production over an infectious time course for the parental
SARS-CoV-2 strain versus Omicron (XBB.1.5) in Huh-7.5 PLSCR1 WT and KO cells (**Figs 6I and J**). As
expected, parental SARS-CoV-2 replicated with faster kinetics upon PLSCR1 depletion. In contrast,
Omicron (XBB.1.5) replication was not affected by PLSCR1 depletion. Our data suggest that
PLSCR1 restricts newer SARS-CoV-2 variants less efficiently in Huh-7.5 cells. This could be due to
adaptation of the recent variants to directly antagonize PLSCR1 and/or utilize an alternative entry
route in Huh-7.5 cells that is TMPRSS2-independent and invulnerable to PLSCR1.

Association between PLSCR1 variants and severe COVID-19

PLSCR1 encodes a 318 amino acid protein containing a palmitoylation motif and a
transmembrane domain which regulate its plasma membrane localization, and a nuclear
localization signal (NLS) and transcriptional activation domain thought to be important for its
nuclear functions (**Fig 7A**)[22, 126-131].

A recent GWAS has identified an association between PLSCR1 variants and severe COVID-
19 outcomes, reporting an odds ratio of approximately 1.2 and a p-value of approximately 10^{-8}

(**Table 1**)[25, 26]. In other words, the GWAS suggests that PLSCR1 has a small, but significant effect

on severe COVID-19 risks.

GWAS typically identify variants associated with increased odds of a disease, but these

variants are not necessarily causative. Among the PLSCR1 variants identified in the COVID-19

GWAS cited above, only p.His262Tyr (rs343320) results in a protein-coding change, located in the

NLS (**Fig 7A**, **Table 1**). This raises the hypothesis that p.His262Tyr might alter PLSCR1's antiviral

function. However, we cannot dismiss the possibility that (i) some non-coding variants identified

in the GWAS could influence the regulation of PLSCR1 mRNA, potentially leading to functional

outcomes, and (ii) the GWAS might have missed other nonsynonymous variants besides

p.His262Tyr that could potentially impact PLSCR1 function.

To assess the relative importance of the p.His262Tyr variant in humans, we examined the

full collection of PLSCR1 variants reported in gnomAD, a human population reference database

(**Fig. 7B**) [132, 133]. The p.His262Tyr variant is the most common nonsynonymous variant in this

gene, with a minor allele frequency (MAF) of 0.07, MAFmax of 0.1, and 4,348 homozygous

individuals in gnomAD. Another variant, resulting in a frameshift mutation, p.Ile110fs

(rs749938276), is the most common predicted loss-of-function (pLOF) of PLSCR1 in gnomAD (MAF

of 0.0003, MAFmax of 0.01, 1 homozygous individual), indicating that PLSCR1 is not essential.

We aimed to investigate the functional effects of the p.His262Tyr and p.Ile110fs variants

in cell culture, along with two other relatively common nonsynonymous variants, p.Gly20Val

(rs79551579, MAF of 0.002, MAFmax of 0.04) and p.Ala29Thr (rs41267859, MAF of 0.01, MAFmax

of 0.04). For this, we stably and ectopically expressed the PLSCR1 variants of interest in PLSCR1

KO A549-ACE2 cells using a lentiviral vector [134]. The variant p.Ile110fs did not rescue PLSCR1

KO, confirming it as a loss-of-function mutation (**Fig 7C**). The p.His262Tyr variant, expressed at

similar levels to WT PLSCR1 in this system (**Supp Fig. 11A**), behaved as a hypomorphic allele,

providing only partial rescue of PLSCR1 antiviral function (**Figs. 7C and D**). Additionally,

introducing PLSCR1 p.His262Tyr into PLSCR1 WT A549-ACE2 cells increased their susceptibility to

SARS-CoV-2 infection (**Fig 7D**), suggesting a dominant effect. However, this effect might be

attributed to the overexpression of PLSCR1 p.His262Tyr from the transgene, compared to the

natural expression levels of PLSCR1 WT from the endogenous locus. To counter this, we examined

patient-derived SV40-immortalized fibroblasts expressing ACE2 that were heterozygous for p.His262Tyr. These cells were already present in our collection, originating from a female tuberculosis patient from Turkey (JL0192) and a female herpes simplex encephalitis patient from France (JL0480). The origin of these cells is serendipitous; our goal was not to link these diseases to PLSCR1 but rather to infect cells carrying the His262Tyr variant with SARS-CoV-2. Cells heterozygous for p.His262Tyr were hyper-susceptible to SARS-CoV-2 infection compared to PLSCR1 WT control SV40-fibroblasts (**Fig 7E**), further suggesting that p.His262Tyr is dominant. We cannot formally rule out that the examined SV40-fibroblasts may carry other mutations influencing SARS-CoV-2 infection.

Since the p.His262Tyr mutation is located in the NLS region, we hypothesized that it could impair PLSCR1's nuclear localization. However, PLSCR1 WT was primarily localized in the cytoplasm in untreated A549-ACE2 cells (**Fig 5A**), in IFN-treated cells (**Supp Fig 11B**), and in SARS-CoV-2 infected cells (**Supp Fig 11C**). Similarly, PLSCR1 His262Tyr was also enriched in the cytoplasm (**Supp Fig 12**). This suggests that the palmitoylation motif, known to be dominant over the NLS [129], dictates the cytoplasmic localization of PLSCR1. Therefore, we propose that the p.His262Tyr mutation affects PLSCR1 through a mechanism other than altering its nuclear localization.

Our data collectively highlight PLSCR1's function in restricting SARS-CoV-2 entry in cell culture, thereby clarifying the association between PLSCR1 variants and severe COVID-19 outcomes [25, 26]. Future human genetic studies are crucial for determining if certain PLSCR1 variants in the population cause increased risks of severe COVID-19. Our results highlight the variant p.His262Tyr as a potential causative candidate, as it caused increased SARS-CoV-2 infection in cell culture.

Discussion

Here, we conducted an unbiased arrayed CRISPR KO screen on Huh-7.5 cells infected with SARS-CoV-2. The screen revealed novel aspects of SARS-CoV-2 and IFN biology while also confirming previously known facets. Pathways related to mRNA transcription and maturation were identified as antiviral. This observation may stem from the conflict between the host cell

and SARS-CoV-2, where the host attempts to export mRNAs from the nucleus to facilitate antiviral

responses while the virus replicates in the cytoplasm, impeding nuclear export [135-138]. RNA

Pol III transcription was specifically essential for the IFN-mediated antiviral response, through

mechanisms that are yet to be determined. Interestingly, inborn errors in POLR3A and POLR3C

have been previously described in patients with severe varicella zoster virus infections [139].

Cellular respiration was identified as a key IFN-independent antiviral pathway. Furthermore,

mitophagy was identified as proviral. This may indicate the infected cell's increased demand for

energy and ATP to combat the virus. Alternatively, cellular respiration may have other, yet-to-be-

identified, IFN-independent antiviral roles. Conversely, translation and cholesterol homeostasis

emerged as the foremost proviral pathways. These findings underscore the complex, dualistic

nature of the interactions between SARS-CoV-2 and host cells.

Our screen notably identified the ISG zinc-finger antiviral protein (*ZC3HAV1/ZAP*) as a

proviral factor in IFN-treated cells. Initially, ZC3HAV1/ZAP gained attention as an antiviral factor

that targets the SARS-CoV-2 RNA genome [115] and prevents programmed ribosomal

frameshifting [140]. Yet, a recent study demonstrated that ZC3HAV1/ZAP also promotes the

formation of SARS-CoV-2 non-structural proteins 3 and 4-induced double-membrane vesicles,

essential for virus replication [72]. SARS-CoV-2 may have adapted to exploit certain ISG products,

such as ZC3HAV1/ZAP, within the cellular environment it encounters. It is still unclear if the

seemingly contradictory roles of ZC3HAV1/ZAP – both proviral and antiviral – are caused by

distinct isoforms.

Many other ISG products influenced SARS-CoV-2 infection, PLSCR1 being the most potent

restriction factor. PLSCR1 did not influence ISG induction as previously reported [24], but rather

inhibited spike-mediated SARS-CoV-2 entry through the endocytic route. Our results corroborate

a recent study from Xu et al [19] and provide an explanation for the enrichment for PLSCR1 SNPs

observed in a GWAS on severe COVID-19 [25, 26].

The molecular mechanisms of PLSCR1-mediated restriction of SARS-CoV-2 entry remain

to be elucidated. PLSCR1 could be altering the lipid composition at the contact site between the

virus and endosomal membranes, akin to the ISG interferon induced transmembrane protein 3

(IFITM3) for influenza A virus [141-144]. PLSCR1 was first identified as a Ca^{2+}-dependent

phospholipid scramblase [23], but it is unclear whether PLSCR1 depletion affects the bidirectional

movement of phospholipids *in vivo* [145-147]. A C-terminal FLAG-tag abolished the antiviral

ability of reconstituted PLSCR1, possibly by interfering with the function of the Ca^{2+} binding

domain. In contrast, inhibiting PLSCR1's phospholipid scramblase activity did not alleviate SARS-

CoV-2 restriction [19].

PLSCR1 is one of a few ISGs known to restrict SARS-CoV-2 entry (**Supp Fig 1A**)[1, 148], with

possibly more yet to be discovered. Future research should investigate the associations and

combinatorial effects between antiviral ISGs, which could lead to key mechanistic insights into

their mode of action.

Intriguingly, PLSCR1 specifically restricted SARS-CoV-2 in Huh-7.5 and A549-ACE2 cells but

it did not show similar inhibitory effects on other viruses that enter cells via endocytosis. Future

studies will investigate the mechanisms behind this specificity. PLSCR1 has been described to

inhibit a range of viruses in various cell lines, such as encephalomyocarditis virus, vesicular

stomatitis virus, Epstein-Barr virus, hepatitis B virus, hepatitis C virus, human cytomegalovirus,

human immunodeficiency virus 1, human T-cell lymphotropic virus type 1, and influenza A virus

[24, 149-155]. It has been proposed that PLSCR1 directly binds viral proteins and impairs their

functions to restrict the non-coronaviruses cited above, reviewed in [156]. However, it seems

unlikely that PLSCR1 has evolved to interact directly with such a diverse set of viral proteins. An

alternative explanation is that diverse viral proteins convergently evolved to bind PLSCR1 as a

mechanism of immune evasion. For example, by altering PLSCR1's subcellular localization.

Meanwhile, overexpressing PLSCR1 in cell culture could act *like a sponge*, absorbing these viral

proteins and thereby hindering viral function. We searched for PLSCR1 interactions in ten SARS-

CoV-2 proteins interactome studies, relying on ectopic expression of individual viral proteins [100-

102, 104-109, 157], and no interaction was reported in two or more independent studies. Two

interactions were reported in a single study: (i) PLSCR1-ORF7b [102], and (ii) PLSCR1-ORF8 [105],

both by proximity biotinylation, which was less stringent compared to affinity purification-mass

spectrometry (AP-MS) and yeast two-hybrid (Y2H) techniques (**Supp Fig 13**). To date, there is no

strong evidence of a direct interaction between a SARS-CoV-2 protein and PLSCR1, although we

cannot rule out that such interactions may occur or even appear in the future as SARS-CoV-2

evolves.

Previous research suggests that Omicron has developed increased resistance to IFN [158,

159], a trait associated with its highly-mutated spike protein [160]. Here, we show that recent

SARS-CoV-2 variants, including Omicron BA.5 and XBB.1.5, exhibit reduced sensitivity to PLSCR1-

mediated restriction compared to the New York 2020 strain, which served as a reference in our

study (**Fig 6**). Evasion from PLSCR1 may thus have contributed to Omicron's increased resistance

to IFN.

The original SARS-CoV-2 strain can enter cells through both TMPRSS2-dependent fusion

near the cell surface and clathrin-mediated endocytosis, with a preference for the former *in vivo*

[38]. ISGs that primarily restrict SARS-CoV-2 endocytosis, such as NCOA7, which perturbs

lysosome acidification [161, 162], and PLSCR1, may play a role in constraining the original strain

to TMPRSS2-dependent entry. Interestingly, the more recent Omicron variant shows both

increased evasion from PLSCR1 restriction **(Fig 6)** and an acquired preference for TMPRSS2-

independent entry. This occurs either via endocytosis or TMPRSS2-independent fusion near the

cell surface, utilizing metalloproteases to cleave the spike protein [160, 163, 164]. We postulate

that Omicron evolved its entry pathway to evade PLSCR1-mediated restriction while shifting its

cell tropism to the upper airways and away from TMPRSS2-expressing cells [163, 165-169].

Future research should investigate: (i) mechanism(s) by which Omicron evades PLSCR1

restriction and whether this evasion is primarily due to mutations in the spike protein, (ii) the

association between PLSCR1-mediated restriction and the distinct entry mechanisms utilized by

SARS-CoV-2 variants, (iii) how various intrinsic factors that restrict virus entry, such as PLSCR1,

have influenced SARS-CoV-2 entry routes, evolution, and cell tropism. Understanding these

aspects of PLSCR1 restriction could provide mechanistic insight into broad strategies of PLSCR1

evasion employed by both newer SARS-CoV-2 variants as well as viruses from other families that

were unaffected by PLSCR1 KO. Subversion of PLSCR1 restriction may serve as a useful immune

evasion strategy that could be shared by diverse viruses with longer evolutionary selection than

SARS-CoV-2.

Several PLSCR1 variants were enriched in a GWAS on severe COVID-19, with a relatively

low odds ratio of approximately 1.2 [25, 26]. This modest odds ratio likely reflects the complex

redundancies within antiviral defenses, from innate immunity featuring multiple effector ISGs

that restrict SARS-CoV-2 [10-15, 18, 19], to adaptive immunity [170]. For example, we show that

the ISGs CH25H [28,29], IFITM2 [122], LY6E [121], and NCOA7 [31,32] still function to restrict

SARS-CoV-2 in PLSCR1-depleted cells (**Supp Fig 8**). However, for these very reasons, the

identification of PLSCR1 variants in the GWAS remains noteworthy. Of these enriched variants,

only PLSCR1 p.His262Tyr (rs343320) resulted in a protein-coding change. Our findings indicate

that p.His262Tyr exhibits a hypomorphic and dominant effect in cell culture, leading to increased

SARS-CoV-2 infection. Future research should aim to ascertain whether p.His262Tyr, or potentially

other PLSCR1 variants, are directly responsible for elevated risks of severe COVID-19 in patients.

Our findings show that baseline levels of PLSCR1 are effective in limiting SARS-CoV-2

infection. This is in line with other studies where ISGs like *DAXX* and *LY6E* were shown to inhibit

SARS-CoV-2 independently of IFN [18, 121, 171]. mRNA-seq analyses of Huh-7.5 cells and primary

human hepatocytes, as well as data from the GTEx consortium on various human tissues [49,

172], revealed that many ISGs are constitutively expressed, even without IFN stimulation (**Supp**

Fig 14). This supports the idea that the IFN-induced antiviral state results more from enhanced

expression of antiviral genes rather than a binary ON/OFF switch. In future studies, it will be

interesting to explore whether intrinsically expressed ISGs also carry out cellular functions beyond

pathogen defense.

Materials and Methods

Plasmids, oligos, and primers

The plasmids, gene fragments, and primers used in this study are listed in **Supp Tables 18,**

19, and 20, respectively.

Cell Lines

Huh-7.5 (human hepatocellular carcinoma; *H. sapiens*; sex: male) [173], Huh-7.5-Cas9 [31], A549-ACE2 (human lung carcinoma; *H. sapiens*; sex: male; generously provided by the laboratory of Brad R. Rosenberg) [11], A549-ACE2-TMPRSS2 [31], Lenti-X 293T (*H. sapiens*; sex: female: Takara, cat. #632180), Caco2 (*H. sapiens*; sex: male; ATCC, cat. HTB-37), Vero E6 (*Chlorocebus sabaeus*; sex: female; kidney epithelial cells, ATCC cat. #CRL-1586), BHK-21 (*Mesocricetus auratus*; sex: unspecified; kidney fibroblasts) cells, and SV40-Fibroblasts were cultured in Dulbecco's Modified Eagle Medium (DMEM, Fisher Scientific, cat. #11995065) supplemented with 0.1 mM nonessential amino acids (NEAA, Fisher Scientific, cat. #11140076) and 10% fetal bovine serum (FBS, HyClone Laboratories, Lot. #KTH31760) at 37°C and 5% CO2. All cell lines tested negative for mycoplasma.

Virus stocks

CHIKV-181/25-mKate2: the infectious clone was a kind gift from Mark Heise (University of North Carolina, USA) [not published yet]. 20 μg of infectious clone DNA was linearized with NotI-HF at 37°C overnight. Complete digestion was confirmed by running a sample of the digested DNA on a 1% agarose gel. After confirmation, linearized DNA was cleaned via phenol-chloroform extraction, and then ethanol precipitated. The precipitated DNA was resuspended in 20 μL of RNase-free H_2O and in vitro transcribed with an SP6 mMessage mMachine In Vitro Transcription Kit (ThermoFisher, cat. AM1340). The generated RNA was electroporated into 1.2 x 10^7 BHK-21 cells, and the produced virus was harvested once approximately 80% of the electroporated cells lifted or showed signs of cytopathic effects and 100% of the cells were positive for mKate2 signal. The titer of the virus was 8.5 x 10^6 focus-forming units (FFU) on Huh-7.5 cells.

hCoV-NL63: was generously provided by Volker Thiel (University of Bern) and amplified at 33°C in Huh-7.5 cells as in [31].

hCoV-OC43: was obtained from ZeptoMetrix (cat. #0810024CF) and amplified at 33°C in Huh-7.5 cells as in [31].

hPIV3-GFP [174]: stock (based on strain JS) grown in VeroE6 cells as in [175].

HSV-1-GFP: stock made by passage on VeroE6 cells. 2 x 10^7 cells seeded in a T175 flask

were infected at an MOI of 0.01 PFU/ml of HSV-1-GFP virus engineered and provided by Ian Mohr

[176]. After a one-hour incubation at 37°C, the inoculum was removed, and 20 ml of DMEM

supplemented to contain 10% FBS and NEAA was added. Cells were incubated at 37°C for 24 h or

until CPE was evident. Cell supernatant containing progeny virus was harvested and titrated on

Vero E6 cells (2.4% avicel, fix 2 dpi) at 2.4 x 10^8 PFU/ml.

IAV WSN (H1N1): was generated in MDCK cells. Cells were inoculated at MOI 0.01 in

DMEM supplemented with NEAA, 0.2% BSA, 0.1% FCS, 50 mM Hepes, and 1 µg/ml TPCK-trypsin.

Virus-containing culture supernatant was harvested at 52 h post-infection and cleared by

centrifugation.

SARS-CoV-2: unless otherwise stated, the isolate SARS-CoV-2/human/USA/NY-RU-

NY1/2020 was used in this study [177]. The virus was sourced from the saliva of a deidentified

patient in New York City, collected on July 28, 2020. Its sequence is publicly accessible (GenBank

OM345241). The virus isolate was amplified in Caco-2 cells. The passage 3 stock employed had a

titer of 3.4 × 10^6 PFU/ml, as measured on Vero E6 cells using a 1% methylcellulose overlay,

according to previously described methods [178]. The Beta (B.1.351), Delta (B.1.617.2), Omicron

BA.5, and Omicron XBB.1.5 variants were obtained from BEI resources (cat. # NR-54008, NR-

55611, NR-58616, and NR-59104, respectively), amplified in Vero E6 cells engineered to stably

express TMPRSS2, and titer was determined as described above.

SINV Toto1101 [179]: expressing an nsP3-mScarletI fusion reporter was generated by

cloning the sequence encoding mScarletI in frame into a unique SpeI restriction site in the

pToto1101 infectious clone plasmid as previously described [180]. *In vitro* transcribed, capped

RNA was generated from the pToto1101-nsP3-mScarletI plasmid (Invitrogen mMessage

mMachine SP6 kit, AM1340) and electroporated into BHK-J cells, a derivative of BHK-21 cells

(ATCC, CCL-10) as previously described [180]. 24 hours post electroporation, centrifuge clarified

supernatants were aliquoted and stored at -80C. BHK-J cells were cultured and virus stocks

generated in MEM supplemented with 7.5% FBS.

VEEV-dsEGFP [16, 181]: the infectious clone plasmid was linearized (MluI) and transcribed

in vitro using an mMessage mMachine SP6 transcription kit (Ambion). BHK-21 cells were

electroporated with viral RNA, and supernatant containing progeny virus was harvested after
incubation at 37°C for 30 h or until CPE was evident. Virus was titrated by plaque assay on BHK-
21 cells (2.4% avicel, fix 2 dpi). BHK-21: 1.45 x 10^9 FPU/ml.

VSV-GFP [182]: grown in BHK-21 cells as in [175].

YFV 17D: was generated via transfection of Huh-7.5 with in vitro transcribed RNA from
pACNR-FLYF-17D plasmid as described in [175].

mRNA-seq

mRNA-seq on SARS-CoV-2-infected cells

Cell culture and infection: 75,500 Huh-7.5 cells or 150,000 Calu-3 cells were seeded in
each well of a 12-well plate with 1 mL media. Media: DMEM with 5% FBS and 1% NEAA for Huh-
7.5 cells or EMEM (ATCC, 30-2003) with 10% FBS for Calu-3 cells. The next day, cells were infected
by removing 500 µL of media and adding 500 µL of media with SARS-CoV-2 strain USA-WA1/2020
(BEI Resources, NR-52281) at 5,000 PFU/well (virus titer determined in Huh-7.5 cells). After one
day, the wells were washed with PBS and cells were harvested in 1 mL TRIzol (Invitrogen, cat.
15596-018). N = 3 replicates (separate wells) per sample.

RNA extraction: 2 ml MaXtract High Density tubes (Qiagen, 129056) were centrifugated
at 12,000–16,000 x g for 20-30 second centrifugation. A volume of 750 µL TRIzol-prepared sample
was combined with 150 µL chloroform in these tubes and hand-shaken vigorously. Phase
separation was accomplished by centrifugation at 1500 x g for 5 min at 4°C. The aqueous phase
was then mixed with 400 µL ethanol 95-100% in a separate tube. These preparations were then
transferred to Zymo Research RNA clean and concentrator-25 kit columns (Zymo Research, cat.
R1018) and subjected to multiple wash and centrifugation steps as recommended by the
manufacturer. An in-column DNase I treatment was performed using Qiagen DNase (Qiagen,
79254). Finally, RNA was eluted with 50 µL DNase/RNase-Free water and stored at -80°C.

Sequencing: Poly-A enriched libraries were made using the TruSeq stranded mRNA LT kit
(Illumina, Cat# 20020594) and sequenced on a NovaSeq SP with PE150 read length.

 mRNA-seq on IFN-treated cells

Cell culture and treatment: A549 cells: 300,000 cells were seeded in each well of a 6-well

plate with 3 mL of DMEM supplemented with 10% FBS and 1% NEAA. The following day, the media

was aspirated and replaced with 2 mL of DMEM with or without 0.5 nM IFN-α2a (PBL, cat. 11101-

2), and the cells were incubated at 37°C. After 24 hours, the cells were harvested in 500 µL TRIzol.

N = 3 replicates (separate wells treated on the same day) per sample. **Calu-3 cells**: 200,000 cells

were seeded in each well of a 12-well plate with 1 mL media. Two days later, the media was

replaced with EMEM (ATCC, cat. 30-2003) with 10% FBS with 0.5 IFN-α2a (PBL, cat. 11101-2) and

incubated at 37°C. 24 h later, cells were harvested in 500 µL TRIzol. N = 3 replicates (separate

wells treated on the same day) per sample. **Huh-7.5 cells**: 75,500 cells were seeded in each well

of a 12-well plate with 1 mL media. Two days later, the media was replaced with 1 mL of DMEM

with 5%FBS, 1% NEAA with 0.5 nM IFN-α2a (PBL, cat. 11101-2) and incubated at 37°C. 24 h later,

cells were harvested in 500 µL TRIzol. N = 3 replicates (separate wells treated on the same day)

per sample.

RNA extraction as described above.

Sequencing: Poly A-enriched libraries were made using the NEBNext Ultra II RNA Library

Prep Kit for Illumina (NEB, cat. E7770) and sequenced on a NovaSeq SP with PE150 read length.

mRNA-seq on PLSCR1 KO cells

Cell culture and CRISPR KO: 30,000 Huh-7.5 cells were seeded in five wells of a 24-well

plate with 480 uL media. The cells were reverse transfected with 120 uL of a transfection mixture

composed of 250 nM of pooled anti-PLSCR1 or non-targeting Edit-R crRNAs from Horizon

Discovery (cat. CM-003729-01-0002, CM-003729-02-0002, CM-003729-03-0002, and CM-

003729-04-0002 or U-007501-01-05, U-007502-01-05, U-007503-01-05, and U-007504-01-05,

respectively) which had been resuspended with an equimolar amount of Edit-R tracrRNA

(Horizon, cat. U-002005-20) and a 1:200 dilution of Dharmafect 4 (Horizon, cat. T-2004-01). The

following day, the media was changed, and the cells were progressively scaled up to a 6-well plate

over the next 4 days. When the cells were confluent in the 6-well plate, the media was removed

from four of the wells. They were then washed with 1x PBS (cat. 14190-144) and lysed with 1 mL

TRIzol (Life Technologies, cat. 15596-018) for 5 minutes at room temperature before transferring

to an Eppendorf tube and freezing at -80°C to await RNA extraction. The remaining well was lysed

with 300 µL of RIPA buffer (Thermo cat. 89900) supplemented with 1x protease inhibitor (Thermo

cat. 87786) and 1x EDTA and prepared for western blot as described below, in the "Western Blots"

section.

RNA extraction as described above.

Sequencing: Poly A-enriched libraries were made using the NEBNext Ultra II RNA Library

Prep Kit for Illumina (NEB, cat. E7770) and sequenced on a NovaSeq SP with PE150 read length.

mRNA-seq analysis

mRNA-seq reads were first quality-filtered and adapter-trimmed using Trim Galore with

parameters -q 20 -e 0.1 --length 20 --paired and Cutadapt. Reads were then mapped to the human

genome GRCh38 or to a combined SARS-CoV-2 MN985325.1/human genome GRCh38 using STAR

[183] with settings including --runThreadN 8 --outFilterMultimapNmax 1 --twopassMode Basic.

Feature counting was performed using the featureCounts function from the Rsubread package

[184], with strandness specified depending on the sequencing and other parameters as default.

The resulting counts were imported into a DESeqDataSet object using the DESeq2 package [185]

with a design formula of ~Group. Size factors were estimated and normalized counts were

extracted and saved. Differential expression analysis was performed using DESeq with the created

DESeqDataSet object, contrasted by sample groups, cooksCutoff and independentFiltering

disabled, and otherwise default parameters.

TMPRSS2 western blot

Cell lysates were prepared in RIPA buffer (150 mM NaCl, 1% NP-40, 0.5% DOC, 0.1% SDS,

50 mM Tris-HCl [pH 7.4] with addition of Halt™ Protease and Phosphatase Inhibitor Cocktail

[ThermoFisher: 78440]) and incubated on ice for 30 min. Lysates were clarified by centrifugation

at 16,000 x g for 10 min at 4°C. Protein concentration was determined by bicinchoninic (BCA)

protein assay (Pierce BCA Protein Assay Kit, ThermoFisher Scientific: 23227) and samples were

resolved on NuPAGE 4%–12% Bis-Tris gels (Invitrogen) followed by transfer onto 0.4 µm

nitrocellulose membranes. Membranes were incubated in blocking buffer (TBS + 5% milk) and

incubated with primary antibody prepared in blocking buffer with 0.1% Tween-20. Membranes

were washed 3X with TBS + 0.1% Tween-20 and incubated with LI-COR (Lincoln, NE, USA) IRDye®

680RD or IRDye® 800CW secondary antibodies. Membranes were imaged using an Azure (Dublin,

CA, USA) 600 imaging system.

Unbiased arrayed CRISPR KO screening

Screen overview

The content of each gRNA 384-well plate constituting the whole-genome library (61

library plates total) was transfected to 16 assay 384-well plates (976 assay plates total). Positive

and negative control gene gRNAs were incorporated into vacant wells of each assay plate as

described below. Huh-7.5-Cas9 cells were subsequently seeded into these assay plates. The 16

assay plates served as replicates for three distinct experimental conditions: 4 replicates for mock

treatment followed by mock infection, 5 replicates for IFN-α2a treatment followed by SARS-CoV-

2 infection, and 7 replicates for mock treatment followed by SARS-CoV-2 infection. Each day, three

library 384-well plates were processed, along with their corresponding 48 assay plates. The full

gRNA library, distributed across 61 384-well plates, was completed over a span of 21 days. For

each set of plates, cell seeding was conducted on day 0, IFN-α2a treatment on day 4, SARS-CoV-

2 infection on day 5, and cell fixation on day 6.

gRNA library preparation

A 0.1 nmol Edit-R Human Whole Genome crRNA Library (Horizon, cat. GP-005005-01)

containing four crRNAs per gene and one gene per well (total 0.1 nmol crRNA/well) was

resuspended in 80 μL of a 1.25 μM tracrRNA (Horizon, cat. U-002005-1000) 10 mM Tris-HCL pH

7.4 solution to create a 1.25 μM gRNA solution. The library was then aliquoted in 10 mM Tris-HCL

pH 7.4 in several 96-well plate and 384-well plate copies using a Tecan Freedom EVO liquid

handler. A single-use library copy containing a 40 μL/well of a 312.5 nM gRNA solution in the 384-

well plate format was used in this study.

gRNA reverse transfection (day 0)

In each well of the 384-well assay plates, 40 μL of a transfection solution was prepared by combining 2% DharmaFect-4 transfection reagent (Horizon, cat. GP-T-2004-07A) in Opti-MEM (Gibco, cat. 31985070). This was added to 40 μL of a 312.5 nM gRNA library using a Thermofisher Multidrop Reagent Dispenser, yielding an 80 μL/well transfection mixture. The mixture was left to incubate at room temperature for 20 minutes. Simultaneously, assay plates were preloaded with 11 μL/well of serum-free media, which was formulated from DMEM, 1X Antibiotic-Antimycotic solution (Gibco, cat. 15240-062), and 1X NEAA, dispensed via a Thermofisher Multidrop Reagent Dispenser. Subsequently, 4 μL/well of the transfection mixture was dispensed into each of the assay plates (16 assay plates per library plate) using a Tecan Freedom EVO liquid handler. During this time, Huh-7.5 cells were prepared in media containing 25% FBS, 1X Antibiotic-Antimycotic solution, and 1X NEAA. A volume of 10 μL cells/well was added to the assay plates, again using a Thermofisher Multidrop Reagent Dispenser. Ultimately, each well contained 1,250 cells in a 25 μL final volume, with a composition of 25 nM gRNA, 10% FBS, 0.8X Antibiotic-Antimycotic, and 0.8X NEAA. Plates were then span at 200 g for 5 minutes. To minimize evaporation, plates were sealed with Breathe-Easy sealing membranes (Sigma-Aldrich, cat. Z380059) and placed in humid chambers constructed from a 245 mm x 245 mm dish containing a paper towel moistened with 15 mL of 1X Antibiotic-Antimycotic solution. Four assay plates were placed in each humid chamber and incubated at 37°C.

IFN-α2a treatment (day 1)

Each well received 5 μL of IFN-α2a (PBL, cat. 11101-2) in media (DMEM, 20% FBS, 1X Antibiotic-Antimycotic solution, 1X NEAA), using a Thermofisher Multidrop Reagent Dispenser, for a final concentration of 1 pM IFN-α2a in a final volume of 30 μL. Plates were then span at 200 g for 5 minutes and incubated at 37°C.

SARS-CoV-2 infection (day 5)

Each well received 212.5 PFU SARS-CoV-2 virus (titer determined on Vero E6 cells, see Virus Stocks section above) diluted in 5 μL of media (DMEM, 20% FBS, 1X Antibiotic-Antimycotic solution, 1X NEAA) for a final volume of 35 μL in the BSL3. Plates were then span at 200 g for 5 minutes and incubated at 37°C.

 Fixing (day 6)

 Each well received 50 μL of 20% neutral buffered formalin (Azer Scientific, cat. 20NBF-4-

 G) and plates were incubated overnight. The formalin mixture was then removed and each well

 received 50 μL of PBS.

 IF staining

 For IF staining of SARS-CoV-2 infected cells in the arrayed CRISPR KO screen (**Figs 1D-F)**, as

 well as some focused experiments (**Fig 4**, **Fig 5G-I**, **Figs 7 C and E**, **Supp Fig 8**, **Supp Fig 9F**): the

 following solutions were prepared for both 96-well plate (96-wp) and 384-well plate (384-wp):

 PBS (Phosphate Buffered Saline), Perm Solution: Comprised of PBS with an added concentration

 of 0.1% Triton X100, Blocking Solution: PBS was mixed with 1% BSA. This solution was prepared a

 day in advance and filtered before use, PBST: PBS with 0.1% of Tween 20, Primary Antibody

 Solution: Genetex anti SARS-CoV-2 N poly rabbit antibody (GTX135357) at a dilution of 1:3000,

 Secondary Antibody Solution: AF647 anti-rabbit antibodies at a dilution of 1:3000 and Hoechst

 33342 (10 mg/ml) at 1:10,000. Plates were stained on a Biotek EL406 Microplate Washer

 Dispenser using the following steps: 1. Priming: The washer was primed with 200 ml of each

 buffer: PBS, Perm Solution, Blocking Solution, and PBST. 2. First Washing Phase: Contents of the

 plates were aspirated. Plates were then washed with 50 μL/well (384-wp) or 200 μL/well (96-wp)

 of Perm Solution, followed by a slow shake for 3 seconds. 3. Permeabilization: A delay of

 approximately 1 minute was implemented for permeabilization, in addition to the time required

 to process all the plates (around 1 minute per plate). 4. Second Washing Phase: Plates were

 washed with 50 μL/well (384-wp) or 200 μL/well (96-wp) of PBS. Subsequently, 50 μL/well (384-

 wp) or 200 μL/well (96-wp) of Blocking Solution was added to the plates, followed by a slow shake

 for 3 seconds. 5. Blocking, autoclean, and Primary Antibody Priming: The washer was set to

 undergo an autoclean cycle with PBS for 30 minutes. Simultaneously, the syringe containing the

 Primary Antibody Solution was primed with 16 ml. 6. Third Washing Phase and First Antibody

 Dispensing: After aspirating the contents of the plates, 15 μL/well (384-wp) or 60 μL/well (96-wp)

 from the Primary Antibody Solution was added, followed by a slow shake for 3 seconds. 7. Primary

 Antibody Incubation, autoclean, and Secondary Antibody Priming: The washer was subjected to

 another autoclean cycle using PBS for 2 hours and 5 minutes. The syringe containing the

Secondary Antibody Solution was primed with 16 ml during this period. 8. Fourth Washing Phase

and Second Antibody Dispensing: Plates were washed with 50 μL/well (384-wp) or 200 μL/well

(96-wp) of PBST, followed by a 2-second slow shake and aspiration. Then, 15 μL/well (384-wp) or

60 μL/well (96-wp) from the Secondary Antibody Solution was added, accompanied by a 3-second

slow shake. 9. Secondary Antibody Incubation and autoclean: An autoclean cycle with PBS was

initiated and lasted for 1 hour. 10. Final Washing Phase: Plates were washed with 50 μL/well (384-

wp) or 200 μL/well (96-wp) of PBST. This was followed by two consecutive washes with 50 μL/well

(384-wp) or 200 μL/well (96-wp) of PBS, incorporating a 2-second slow shake in each cycle. Finally,

plates were left with 50 μL/well (384-wp) or 200 μL/well (96-wp) of PBS.

Imaging

Plates were imaged with a ImageXpress micro-XL and analyzed with MetaXpress

(Molecular Devices).

Analysis

Analysis was conducted in R.

Data Omission: We excluded five library plates, constituting 8% of the total library, due to

insufficient infection levels for accurate quantification.

Normalization: Two variables were subject to normalization—percentage of SARS-CoV-2

positive cells and the count of nuclei. The normalization steps were applied separately for the

three screening conditions: mock treatment followed by mock infection, IFN-α2a treatment

followed by SARS-CoV-2 infection, and mock treatment followed by SARS-CoV-2 infection. Data

was first Z-scale normalized within assay plates:

$$Scale(x) = \frac{x - mean(x)}{sd(x)}$$

And then Z-scale normalized per row and per column to remove any spatial effects.

Statistics: a robust statistic accounting for technical and biological variability was applied

using the below formula within the replicates of each gene**:**

$$Stat\ score(x) = \frac{mean(x)}{sd(x)}$$

This statistic was further standardized by Z-scaling across all genes to produce our final z-

score.

Exclusion of genes influencing cell proliferation: 224 genes with nuclei count z-score ≥ 2

and 388 genes with nuclei count z-score ≤ 2 in the mock treatment followed by mock infection

condition were deemed to influence cell proliferation and excluded from subsequent analyses.

Exclusion of genes not expressed in cell lines of interest (A549, Calu-3, Huh-7.5 cells) and

in human lung cells. Expression data from cell lines from [47, 48]. Expression data from tissues

from [49]. Genes were considered expressed if they had at least one read count within exons.

Gene set enrichment analysis

For the pathway analysis in **Supp Fig 2E**, we leveraged the FGSEA package [186] to perform

Gene Set Enrichment Analysis (GSEA) using gene sets found in the Molecular Signatures Database

(MSigDB) [187]: Reactome [188], KEGG [189], Wikipathways [190], Pathway Interaction Database

[191], and Biocarta [192]. The analysis was conducted separately for two conditions: IFN-α

pretreated SARS-CoV-2 infection and non-pretreated SARS-CoV-2 infection. We attributed a score

to each pathway for both conditions:

$Score = -log10(padj) \times sign(Normalized\ Enrichment\ Score)$

Each pathway was then attributed to one of nine quadrants (as in **Fig 1F**) based on its

score in the IFN-α pretreated condition (axis x) versus non-pretreated condition (axis y), using

padj ≤ 0.05 as a cutoff.

Compilation of published large-scale omic studies on SARS-CoV-2

As a rule, we listed the genes classified as 'hits' by the authors of the respective studies.

Below are some exceptions or clarifications:

Functional genetic screens

Baggen, et al. [80]: we used the "low stringency adjusted" analysis in Suppl Table 11.

Proviral: p_value_neg ≤ 0.05 and log2 FC ≥ 1. Antiviral: p_value_pos ≤ 0.05 and log2 FC ≤ -1. We

also used "High stringency" analysis in Suppl Table 7. Proviral: Gene is TMEM106B (log2 FC = 3.8

and p_value_neg = 0.08) or p_value_neg ≤ 0.05 and log2 FC ≥ 1 (no gene matched this criteria).

Antiviral: p_value_pos ≤ 0.05 and log2 FC ≤ -1 (no gene matched this criteria). **Biering, et al. [79]:** in Supplementary Table 1, in Tab 1: LOF-enriched screen analysis, for proviral genes, we used FDR ≤ 0.05. In Tab 2: GOF-depleted screen analysis, proviral: FDR ≤ 0.05. In Tab 3: GOF-enriched screen analysis, for antiviral genes, we used FDR ≤ 0.05. **Chan, et al. [78]:** in Multimedia component 6, for Vero E6 (T16); UM-UC-4 (T23); HEK293+A+T (T12); HuH-7 (T15) and Calu-3 (T43), we considered gene as hits at FDR < 0.1 (as in Figure 4A). We listed the genes as proviral if differential ≥ 0 or antiviral if differential ≤ 0. **Daniloski, et al. [77]:** we used Table S1. FDR MOI1 ≤ 0.05 or FDR MOI3 ≤ 0.05. **Danziger, et al. [11]:** in S1 Table, we used the genes annotated as proviral or antiviral by the authors. **Gordon, et al. [76]:** for A549 +ACE2 in Table S6 or Caco2 in Table S7, for proviral genes, we used Averaged z-scores ≤ 2 and for antiviral genes, we used Averaged z-scores ≥ 2. **Grodzki, et al. [75]:** for VeroE6 in additional file 4, tab 4, we used FDR <0.25. For HEK293T +Cas9 Study 1 in additional file 6, tab17, we used FDR <0.25. For HEK293T +Cas9 Study 1 in additional file 7, tab7, we used FDR <0.25. **Hoffmann, et al. [28]:** for 37°C (Table S1E) and for 33°C (Table S1C), we selected proviral genes if FDR ≤ 0.05 and z-score ≥ 0 and antiviral genes if FDR ≤ 0.05 and z-score ≤ 0. **Hossain, et al. [69]:** in Figure 3E, we selected the top 15 genes in the spike-mNG axis by negative log robust rank aggregation. **Israeli, et al. [74]:** we used Supplementary Data 1. **Kaur, et al. [15]:** we used the genes labelled in Figure 1. **Le Pen et al. (this study):** we used z-score ≥ 2 for antiviral genes and ≤ 2 for proviral genes, see Unbiased arrayed CRISPR KO Screen analysis section above for more details. **Loo, et al. [73]:** we used the genes labelled in Figure 2. **Mac Kain, et al. [18]:** in Electronic Supplementary Material 5, for antiviral genes, we used the filter: pos|rank ≤13, for proviral genes, we used neg|rank ≤13. **Martin-Sancho, et al. [12]:** we used Table S3, "Lentivirus validated hits". **Pahmeier, et al. [72]:** we used the genes labelled in Figure 6. **Rebendenne, et al. [71]:** for Calu3_Gattinara, we used for proviral genes: residual_z-score_avg ≥ 2.5 and for antiviral genes: residual_z-score_avg ≤ 2.5 (no gene). For VeroE6, proviral: residual_z-score_avg ≥ 2.5, and antiviral: residual_z-score_avg ≤ 2.5. For Caco2, proviral: residual_z-score_avg ≥ 2.5 and antiviral: residual_z-score_avg ≤ 2.5. For Calu3_Calabrese, proviral: residual_z-score_avg ≤ 2 and antiviral: residual_z-score_avg ≥ 2. **Rehfeld, et al. [70]:** In Table S1, we considered genes as hits for PRF-1 top eGFP-mCh or PRF-1 bottom eGFP-mCh if FDR ≤ 0.05. **Schneider, et al. [31]:** for both 37°C (Table_S1A) and 33°C (Table_S1B), we listed the gene as

proviral if FDR ≤ 0.05 and z-score ≥ 0 and antiviral if FDR ≤ 0.05 and z-score ≤ 0. **Wang, et al. [67]:**

we used Table S1. Proviral: Enrichment score ≤ 10^(-4). **Wei, et al. 2021 [66]:** we used Table S1.

For proviral genes, we used Cas9-v1 Avg. ≥ 2.5 & Cas9-v2 Avg. ≥ 2.5. Average between Cas9-v1

Avg. and Cas9-v2 Avg. is given in the table. For antiviral genes, we used Cas9-v1 Avg. ≤ -2.5 &

Cas9-v2 Avg. ≤ -2.5. Average between Cas9-v1 Avg. and Cas9-v2 Avg. is given in the table. **Wei, et**

al. 2023 [65]: in Table S1, for Day 7 or Day 14, we used fdr ≤ 0.05 for positive regulators of

ribosomal frameshifting or negative regulators of ribosomal frameshifting. **Wickenhagen, et al.**

[13]: we used the genes labelled in Figure 1B. **Xu, et al. [19]:** in Huh-7.5 or A549-ACE2 cells, in

untreated or in IFN-gamma treatment, we used Log10 p-value (mNG-High vs. mNG-Low

Enrichment) ≥ 3. **Zhu et al. [64]:** In Supplementary Data 1 , SARS-CoV-2 WT and 2 VOCs tested,

for proviral genes, we used pos.score_wt ≤ 0.0005 or pos.score_alpha ≤ 0.0005 or pos.score_beta

≤ 0.0005.

Human genetic studies

Degenhardt, et al. [90]: we used Table 2 and added KANSL1 and TAC4, based on new

analysis by Pairo-Castineira et al. 2023 [25]. **Kousathanas, et al. [26]:** we used Table 1 from Pairo-

Castineira, et al. 2023 [25]. **Pairo-Castineira, et al. 2021 [98]**: we used Table 1 from Pairo-

Castineira, et al. 2023 [25]. **Pairo-Castineira, et al. 2023 [25]**: we only considered variants near

annotated genes (i.e., we excluded rs1073165). **Roberts, et al. [83]:** we used Figure 2. **Zhou, et**

al. [95]: we used Table 1 and the p-values from COVID-19 hospitalization (European ancestry

only).

SARS-CoV-2 protein interactomes

Davies, et al. [109], Laurent, et al. [102], Li, et al. [108], Samavarch-Tehrani, et al. [100],

St-Germain, et al. [103], Stukalov, et al. [107]: we used the Supplementay Table 3 from [27].

Gordon, et al. [101]: we used the genes listed in Table S2. Liu, et al. [104] and May, et al. [105]:

we used the genes listed in May, et al. [105] Table S5-new. Zhou, et al. [106]: we used the genes

listed in Table S1. "SARS-CoV-2-human protein-protein interactions identified in this study."

SARS-CoV-2 RNA interactomes

Flynn, et al. [114], Kamel, et al. [113], Labeau, et al. [112], Lee, et al. [115], Schmidt, et al.

2021 [111]: we used the Supplementay Table 3 from [27]. Schmidt, et al. 2023 [110]: we used

Table S2, "Huh-7 interactome comparison tab", genes listed in the following categories: "Huh-7
gRNA FDR5 HS" and "Huh-7 sgmRNA FDR5 HS".

Altered phosphorylation states in SARS-CoV-2-infected cells.

Bouhaddou, et al. [116]: for Vero E6 cells, we used Table S1, tab 1 "PhosphoDataFull" and
filtered for adj.pvalue ≤ 0.05 & log2FC ≥ 1 or adj.pvalue ≤ 0.05 & log2FC ≤ -1 in at least three
different time points.

Generation of PLSCR1 KO cells

CRISPR KO

KO Huh-7.5 and A549-ACE2 cells were generated using two anti-PLSCR1 Edit-R crRNAs
from Horizon Discovery (cat. CM-003729-02-0002 and CM-003729-04-0002) or non-targeting
controls (cat. U-007501-01-05 and U-007502-01-05) resuspended with an equimolar amount of
Edit-R tracrRNA (Horizon, cat. U-002005-20) to form sgRNAs. The sgRNAs were then co-
transfected with Cas9-mKate2 mRNA (Horizon, cat. CAS12218) according to the manufacturer's
protocol. 24 to 48 hours after transfection, cells were examined for mKate2 signal and FACS sorted
into bulk and single cell populations, gating on mKate2 signal. Bulk and single cell populations
were then assessed for PLSCR1 expression by western blot to confirm KO.

Amplicon sequencing

Genomic DNA was isolated from a frozen cell pellet using the Qiagen DNeasy kit (Qiagen,
cat. 69504) and treated with RNAse A in the optional RNA digestion step. The region of interest
was then amplified using Q5 2x mastermix (New England Biolabs, cat. M0492S), 500 ng of
template DNA, 0.5 μM of forward and reverse primers, and the following PCR conditions: 98°C
for 30 seconds, followed by 98°C for 5 seconds, 64°C for 15 seconds, and 72°C for 20 seconds,
repeating those steps 30 times before holding at 72°C for 2 minutes. The primers used when
amplifying PLSCR1 genomic DNA from WT and KO Huh-7.5 and A549+ACE2 cells were RU-O-
32687 (5' AACATAGAGGTGATTATGATTTCGTCT) and RU-O-32526 (5'
GGAGGAGCTTGGATTTCTATCTAC). PCR reactions were run on a 1% agarose gel to confirm
amplification. Amplicons were purified with a Zymo DNA clean and concentrator kit (Zymo, cat.
D4013) before sending to Genewiz for amplicon sequencing.

Western Blots

Cell pellets were collected and lysed in RIPA buffer (Thermo, catalog number 89900) with 1x Halt protease inhibitor cocktail and 1x EDTA (Thermo, catalog number 87786). Cell lysates were spun down in a refrigerated centrifuge at 15,000 g at 4°C for 15 minutes to pellet any cell debris, and the supernatant was collected and transferred to another tube. The collected samples were quantified by BCA assay (Thermo Scientific, cat. #23225). Before loading into the gel, we added sample buffer (Thermo, catalog number NP0007) with β-mercaptoethanol and heated the sample at 95°C for 10 minutes. Samples were allowed to cool back to room temperature before loading into 12% Bis-Tris 1.0 mm gels (Invitrogen, cat. #NP0321BOX). Proteins were electrophoretically transferred onto nitrocellulose membranes. Membranes were blocked with 5% fat-free milk in 1X TBS (Thermo, catalog number NP00061) and then incubated with primary antibody at 4°C overnight in 5% fat-free milk in 1x TBS with 0.5% Tween-20 (TBST). Primary antibody: rabbit anti-PLSCR1 polyclonal antibody (Proteintech, cat. #11582-1-AP) and mouse anti-β-actin antibody (Millipore Sigma, cat. A5316-100UL) as a loading control. After incubation, membranes were washed three times with 1x TBST and then incubated with fluorescently conjugated secondary antibodies for 2 hours at room temperature. Secondary antibodies: LI-COR IRDye goat anti-rabbit 800 and goat anti-mouse 680 (LI-COR cat. 926-32211 and 926-68070, respectively). Membranes were washed three times with 1X TBST, once with 1X TBS, then imaged on an Azure 600. For the western blot in Supplementary Figure 4, this protocol was modified slightly: proteins were electrophoretically transferred onto 0.22 µm polyvinylidene difluoride (PVDF) membranes, incubated with a primary antibody solution of rabbit anti-PLSCR1 polyclonal antibody (Proteintech, cat. #11582-1-AP) and polyclonal rabbit anti-RPS11 antibody (Abcam, cat. ab157101), a secondary antibody solution of goat anti-rabbit HRP (Invitrogen, cat. 31462) and visualized using a SuperSignal West Femto Maximum Sensitivity Substate kit (Thermo, cat. #34096).

Cell viability assay

4,000 A549-ACE2 cells/well or 8,000 Huh-7.5 cells/well were seeded on day 0 in 100 µL media (DMEM, 10% FBS, 1X NEAA, 1X Penicillin-Streptomycin) in a 96-well plate. The next day,

blasticidin selection was added as indicated in the figure to serve as a control for reduced cell
viability. On day 4, cell viability was assessed by resazurin assay (Abcam, cat. ab129732) according
to the manufacturer's protocol.

JAK-STAT inhibitor treatment

InSolution (Millipore, cat. 420097-500UG) was used according to the manufacturer's
instructions.

Titration of IFN-a2a in CHIKV-infected cells

6,000 Huh-7.5 cells/well were seeded in 100 µL media (DMEM, 10% FBS, 1X NEAA). The
following day, we treated cells with one of twelve concentrations of IFN-a2a (PBL, cat. 11101-2):
64 pM, 32 pM, 16 pM, 8 pM, 4 pM, 2 pM, 1 pM, 0.5 pM, 0.25 pM, 0.125 pM, 0.0625 pM, and 0
pM. The following day, the cells were infected with 2 µL of CHIKV-181/25-mKate2 (approximately
17,000 FFU per well, titer determined on Huh-7.5 cells) and fixed after 12 hours. Plates were
stained with a 1:1000 dilution of Hoechst for at least 10 minutes before washing with PBS and
imaging for mKate2 signal.

RT-qPCRs on ISGs

Huh-7.5 and A549-ACE2 cells were seeded at densities of 36,000 or 18,000 cells/well,
respectively, in 500 µL of media (DMEM, 10% FBS, 1X NEAA) in 24-well plates. The following day,
a dilution series of IFN-α2a (PBL, cat. 11101-2) or IFN-β (PBL, cat. #11415) was prepared (64 pM,
32 pM, 16 pM, 8 pM, 4 pM, 2 pM, 1 pM, 0.5 pM, 0.25 pM, 0.125 pM, 0.0625 pM, and 0 pM) and
50 µL of each dilution added to the cells in duplicate. After 24 hours, the media was removed and
the cells were washed with 1 mL of ice-cold PBS. 200 µL of RNA Lysis Buffer (Zymo Research, cat.
#R1060-1-100) was added to the cells, and the plates were frozen at -20°C before RNA isolation.
RNA was extracted using the Zymo Quick RNA 96-kit (Zymo Research, cat. R1052) including
DNaseI treatment, followed by cDNA synthesis using the SuperScript™ IV VILO™ Master Mix
(Invitrogen, cat. 11756050) according to manufacturers' instructions. qPCRs were conducted on

a QuantStudio 3 cycler using the Taqman Fast Advance master mix (Life Technologies Corporation,

cat. 4444965) and the following assays: *RPS11* (ThermoFisher 4331182; Hs01574200_gH), *IFI6*

(ThermoFisher 4331182; Hs00242571_m1), *OAS1* (ThermoFisher 4331182; Hs00973635_m1).

IFI6 and *OAS1* were normalized to *RPS11* mRNA levels using the deltaCt method [193].

PLSCR1 subcellular localization

IF staining

A549-ACE2 cells were plated onto #1.5, 12mm glass coverslips (Fisher Scientific, cat.

#1254581) placed at the bottom of the wells of a 24-well plate. When confluent, the cells were

fixed, permeabilized with 1% Triton X-100 for 5 minutes and blocked for 1 hour at room

temperature with 1 mL of PBS-BGT (1x PBS with 0.5% bovine serum albumin, 0.1% glycine, 0.05%

Tween 20). Afterward, the cells were incubated in a 1:500 dilution of 4D2 mouse anti-PLSCR1

antibody (Millipore Sigma, cat. #MABS483) in PBS-BG (1x PBS with 0.5% bovine serum albumin

and 0.1% glycine) overnight at 4C with rocking. The cells were then washed twice with PBS-BGT

before incubation with a secondary antibody solution of 1:1000 anti-mouse 588 (ThermoFisher,

cat. #A-11001) and 1:1000 Hoechst dye (ThermoFisher, cat. #62249) in PBS-BG for two hours at

room temperature, followed by three washes with PBS-BGT. For those images where cells were

treated with IFN shown in **Supp Fig 11B**, A549-ACE2 cells were treated with 20 pM of IFN-α2a

(PBL, cat. 11101-2) for 48 hours before harvest, then fixed and stained as described above. For

those images where cells were infected with SARS-CoV-2 shown in **Supp Fig 11C**, A549-ACE2 cells

were infected with approximately 3,400,000 PFU of SARS-CoV-2 (titer determined on Vero E6

cells) diluted in 200 μL Opti-MEM to achieve almost 100% infection. After 24 hours, cells were

fixed and stained as described above, with slight modification to the antibody solutions. Primary

antibody solution: 1:500 4D2 mouse anti-PLSCR1 and 1:3000 rabbit anti-SARS-CoV-2 N (Genetex,

cat. GTX135357) in PBS-BG. Secondary antibody solution: 1:1000 goat anti-mouse 594, 1:1000

goat anti-rabbit 647, and 1:1000 Hoechst dye.

Imaging

The coverslips were mounted onto slides (Fisher Scientific, cat. #1255015) with Invitrogen

ProLong Gold Antifade Mountant (Fisher Scientific, cat. # P36930). The slides were allowed to

cure for 24 hours before the edges of the coverslips were sealed. For **Fig 5A**, the cells were imaged

by confocal microscopy. Confocal images were acquired using Zeiss Zen Blue (v3.5) software on a

LSM 980 point scanning confocal microscope (Zeiss) hooked to a Axio Observer.Z1 / 7 stand

equipped with C Plan-Apochromat 63X/1.40 oil (RI:1.518 at 23°C) objective lens (Zeiss). CW

excitation laser lines 405 nm and 488 nm were used to excite the fluorescence of DAPI and AF488

labeled samples. Emitted fluorescence were spectrally grated (410-483 nm for DAPI, 499-552 nm

for AF488) to avoid fluorescence bleed through and were detected in MA-PMT (DAPI), and GaAsP-

PMT (AF488). The confocal pinhole was set to 1AU for AF488, and the detector master gains were

set within the linear range of detection (550-750V). For the IFN-treated and SARS-CoV-2 infected

cells shown in **Supp Fig 11B,C**, these were mounted as described previously but imaged via

widefield imaging. Widefield imaging was performed in the Rockefeller University's Bio-Imaging

Resource Center, RRID:SCR_017791, on a DeltaVision Image Restoration Microscope (Applied

Precision, now Leica Microsystems) using inverted IX-70 stand (Olympus, Evident Scientific) and a

60X NA 1.42 oil immersion objective (Olympus), and filter sets for DAPI (Ex: 390 ± 18, Em: 435 ±

48), Alexa Fluor 594 (Ex: 575 ± 25, Em: 632 ± 60), Alexa Fluor 647 (Ex: 575 ± 25, Em: 676 ± 34).

Images were acquired on a pco.edge sCMOS camera, and were subsequently deconvolved using

the Deltavision image acquisition software (SoftWorx). Scanned images for both confocal and

widefield imaging were saved as .czi files.

Focus-forming assay on SARS-CoV-2-infected cells

In **Fig 5B**, Huh-7.5 and A549-ACE2 cells were cultured in media (DMEM with 5% FBS) and

seeded at densities of 2×10^5 and 1×10^5 cells per well, respectively, in collagen-coated 12-well

plates to reach 80-90% confluency by the day of infection. A 1:10 serial dilution of virus stock was

made in Opti-MEM in five separate tubes. Media was aspirated from the cells, and the wells were

washed with 1 ml of PBS before adding 200 µL of each virus dilution to the cells in triplicate. Plates

were incubated at 37°C with 5% CO2 for 1 hour, rocking every 15 minutes for even virus

distribution. A 1% methylcellulose overlay medium was prepared and mixed with complete

growth media at 37°C; 2 ml of this overlay was added to each well after removing the virus

inoculum. Plates were then incubated at 37°C with 5% CO2 for 48 for Huh-7.5 cells or 72 hours

for A549-ACE2 cells. Cells were then fixed in final 10% neutral buffered formalin and IF stained as

described in the Unbiased Arrayed CRISPR screen section. PLSCR1 KO and WT cells were

compared at similar virus dilutions.

In **Fig 6G**, the above protocol was followed to titer SARS-CoV-2 strains on Huh-7.5 PLSCR1

KO cells. Then, Huh-7.5 WT and KO cells were seeded at 2×10^5 cells per well in 1 mL of media

(DMEM, 10% FBS, 1X NEAA) in 12-well plates to reach 80-90% confluency the next day. Media

was aspirated from the cells, the wells were washed with 1mL of PBS, and then the cells were

infected with approximately 50 FFU of SARS-CoV-2 (for each strain) as determined on PLSCR1 KO

Huh-7.5 cells diluted in 200 µL of Opti-MEM. Plates were then incubated, overlayed with

methylcellulose, fixed, and stained as described above.

1006

Transfection with SARS-CoV-2 replicon system

The SARS-CoV-2 replicon and the method for electroporation has been described

previously [119]. Briefly, 6×10^6 Huh-7.5 WT and PLSCR1 KO cells were electroporated at 710 V

with 2 µg of SARS-CoV-2 N mRNA and 5 µg of replicon RNA. The cells rested for 10 minutes at

room temperature before resuspending to a concentration of 300,000 cells/mL and plating 100

µL of cells into each well of a 96-well plate. After 24 hours, supernatant was collected from the

replicon-transfected cells and assayed for *Renilla* luciferase activity according to kit instructions

(Promega, cat. E2810).

1015

Transduction with SARS-CoV-2 spike/VSV-G-pseudotyped, single-cycle, replication-

defective HIV-1 viruses.

Virus preparation

SARS-CoV-2 spike/VSV-G-pseudotyped, single-cycle, replication-defective HIV-1 viruses

(pCCNanoLuc/GFP) were prepared as in [120]. Plasmids were a kind gift of Theodora Hatziioannou

and Paul D. Bieniasz (The Rockefeller University, NY, USA) [120, 194, 195]. One day before the

transfection, 4×10^6 293T cells were seeded in a 10 cm dish. One hour prior to transfection, the

growth media in the dish was replaced with 9 mL of fresh media containing 2% serum. A 1,000

µL transfection mixture was prepared, comprising the diluent (a 150 mM NaCl solution prepared

with sterile cell culture water), 5 µg of HIV GP plasmid, 5 µg of pCLG plasmid, and either 2.5 µg of

SARS-CoV-2 spike Δ19 or 1 µg of pHCMV.G plasmid, ensuring the total plasmid content did not

exceed 12.5 µg. After brief vortexing, 50 µL of PEI (1 mg/mL, Polysciences cat. 23966) was added

to achieve a 1:4 DNA/PEI ratio. The mixture was vortexed for 5 seconds and then allowed to sit

for 20 minutes in a hooded environment. Following gentle mixing by pipetting, 1 mL of the

transfection mixture was added to the 10 cm dish. Media was changed 12 hours post-transfection,

and the supernatant was harvested and filtered through a 0.2-micron filter 48 hours post-

transfection, then stored at -80°C.

Transduction of PLSCR1 KO or WT A549-ACE2 cells

Seeded in 96-well plates, 6,000 A549-ACE2 cells per well were cultured in 100 µL of media.

After two days, either 10 µL of SARS-CoV-2 spike pseudotyped virus or 0.01 µL of VSV-G

pseudotyped virus were diluted in a final volume of 100 µL of media and added to the wells to

yield comparable NanoLuc signals. Plates were then spun at 200 g for 5 minutes and incubated at

37°C. Two days post-infection, the media was aspirated and replaced with 50 µL of NanoGlo

solution, sourced from the Promega Nano-Glo Luciferase Assay kit (Promega, N1110), with a

substrate to buffer ratio of 1:100. NanoLuc signal was subsequently quantified using a Fluostar

Omega plate reader.

Transduction of siRNA-treated HEK293T cells

Seeded in 96-well plates, 1,600 HEK293T, HEK293T-ACE2, or HEK293T-ACE2-TMPRSS2 cells

were cultured in 80 µL of media. The next day, a 20 µL transfection mixture made of Opti-MEM,

1% DharmaFECT1 (Horizon, T-2001-03), and 250 nM siRNA, PLSCR1 ON-TARGETplus SMARTpool

siRNA (Horizon, cat. L-003729-00-0005) or non-targeting control (Horizon, cat. D-001810-10-05)

was added to the cells. The final concentration of siRNAs was 25 nM. After two days, either 2 µL

of SARS-CoV-2 spike pseudotyped virus or 0.2 µL of VSV-G pseudotyped virus were diluted in a

final volume of 100 µL of media and added to the wells to yield comparable NanoLuc signals.

Plates were then span at 200 g for 5 minutes and incubated at 37°C. Two days after infection,

NanoLuc signal was quantified as described in the "Infection of PLSCR1 KO or WT cells" section.

1052

Infection of siRNA-treated A549 cells with SARS-CoV-2

Seeded in 96-well plates, 1,000 A549, A549-ACE2, or A549-ACE2-TMPRSS2 cells were

cultured in 80 µL of media. On the same day, a 20 µL transfection mixture made of Opti-MEM, 1%

DharmaFECT1 (Horizon, T-2001-03), and 250 nM siRNA, PLSCR1 ON-TARGETplus SMARTpool

siRNA (Horizon, cat. L-003729-00-0005) or non-targeting control (Horizon, cat. D-001810-10-05)

was added to the cells. The final concentration of siRNAs was 25 nM. Three days after

transfection, the cells were infected by adding 34,000 PFU of SARS-CoV-2 (titer determined on

Vero E6 cells) diluted in 10 µL media to each well. Plates were then spun at 200 g for 5 minutes

and incubated at 37°C. Staining and readout as described above in the "Unbiased arrayed CRISPR

KO screening" section.

1063

Pan-virus infection of PLSCR1 KO cells

A549-ACE2 cells were seeded at a density of 6,000 cells/well in 96-well plates in 90 µL

media. The following day, 10 µL diluted virus was added to each well. Virus concentrations as

follow: CHIKV-mKate, 0.05 µL virus stock per well (titer 8.5 x 10^6 PFU/mL determined in Huh-7.5

cells); hCoV-NL63, 10 µL virus stock per well (titer 1.4 x 10^5 PFU/mL); hCoV-OC43, 10 µL virus stock

per well (titer 1.06 x 10^7 PFU/mL); hPIV-GFP, 0.05 µL virus stock per well; HSV1-GFP, 0.5 µL virus

stock per well (titer 2.4 x 10^8 PFU/mL determined on Vero E6 cells); IAV WSN, 0.5 µL virus stock

per well; SARS-CoV-2, 0.5 µL virus stock per well (titer 3.4 x 10^6 PFU/mL determined on Vero E6

cells); SINV-Toto1101-mScarletI, 10 µL virus stock per well; VEEV-EGFP, 0.005 µL virus stock per

well (titer 1.45 x 10^9 PFU/mL determined on BHK-21); VSV-GFP, 0.05 µL virus stock per well;

YFV_17D, 5 µL virus stock per well. The cells were fixed, stained and imaged as described in the

Unbiased arrayed CRISPR KO screen section. Fluorescent viruses were not stained: the fluorescent

signal was used as a reporter. We used the following primary antibodies when applicable: anti-

dsRNA (J2) mouse (Nordic MUbio, cat. 10010200) diluted 1:500 was used for hCoV-NL63 and

hCoV-OC43, anti-IAV mouse (Millipore, cat. MAB8257) diluted 1:3000, anti-YFV mouse (Santa

Cruz Biotechnology, cat# sc-58083) diluted 1:500, anti-SARS2-S rabbit (Genetex, cat. GTX135357)

diluted 1:3000.

Huh-7.5 cells were transfected with a 1:200 dilution of Dharmafect 4 (Horizon, cat. T-2004-

01) and 25 nM ON-TARGETplus SMARTpool siRNAs (Horizon Discovery) in 96-well plates. The cells

were infected three days after siRNA transfection with hCoV-NL63 or hCoV-OC43 or four days

after siRNA transfection with SARS-CoV-2 or hCoV-229E. IF and imaging as described above.

1085

Measuring extracellular ACE2 expression using flow cytometry

A549-ACE2 cells were detached from the wells using Accutase (Innovative Cell

Technologies, Inc., Cat: AT-104) for 5 minutes at 37°C. Following detachment, cells were rinsed

once with PBS and stained with LIVE/DEAD Fixable Aqua (Invitrogen, Cat: L34966, diluted 1:1000

in PBS) for 15 minutes at room temperature in the dark. After staining, cells were rinsed once

more with PBS. Then, cells were incubated with FcR blocking reagent (Miltenyi Biotec, 1:50) and

ACE2-AF488 (Sino Biological, Cat: 10108-MM37-F-100, 1:30) in FACS buffer (2% FBS in PBS) for 30

min at 4°C in the dark. Following incubation, cells were washed twice with FACS buffer and

subjected to flow cytometry analysis. Samples were acquired on an LSRII flow cytometer (BD

Biosciences), and the results were analyzed with FlowJo software (Tree Star). Before analyzing cell

surface ACE2 expression, gating procedures were implemented to exclude dead cells and

doublets from the analysis.

1098

SARS-CoV-2 variant infections

For **Figure 6A-E**, Huh-7.5 cells were plated at 7,500 cells/well in 100 µL of media (DMEM,

10% FBS, 1X NEAA) in 96-well plates. The next day, cells were infected with quantities of virus that

yielded comparable percent infections in the WT cells, then spun at 200 g for 5 minutes and

incubated at 33°C. The quantities of virus used were as follows: 0.1 µL/well for parental, 0.05

µL/well for beta, 1 µL/well for delta, 0.5 µL/well for Omicron BA.5, and 0.05 µL/well for Omicron

XBB.1.5. The infected cells were fixed after 24 hours and stained as described previously.

1106

Growth curves using the parental SARS-CoV-2 strain and Omicron (XBB.1.5)

Huh-7.5 WT and PLSCR1 KO cells were plated at 100,000 cells/well in 1mL media (DMEM,

10% FBS, 1X NEAA) in a 24-well plate. The following day, cells were infected at MOI 0.01 (as titered

on PLSCR1 KO Huh-7.5 cells) with the New York 2020 isolate of SARS-CoV-2 ("parental") and SARS-

CoV-2 Omicron XBB.1.5. Plates were incubated for 1hr at 37°C. Inoculum was removed and saved

as the 0hr timepoint. Cells were washed twice with PBS. Fresh media was added, then collected

as the 1hr timepoint. Fresh media was added again, and the cells were incubated at 33°C. All

supernatant was collected from the cells again at 24 hours, 48 hours, and 72 hours, and was

replaced with fresh media. Collected supernatant was frozen at -80°C until titering. Supernatants

were titered by plaque assay as follows: Vero E6 cells were plated at 200,000 cells/well in 12-well

plates. The following day, collected supernatant was tenfold serially diluted, the media removed

from the plates, and 300 µL diluted virus added. Each sample of supernatant was titered in

duplicate. Cells were incubated with virus for 1 hour at 37°C, then inoculum was removed and

1mL of 1% methylcellulose (supplemented with 5% FBS). Plates were incubated for 3 days at 37°C,

then fixed with 2mL of 7% formaldehyde and stained with crystal violet before being rinsed with

water, left to dry overnight, and the plaques counted.

1123

Generating and infecting cells overexpressing ISGs (Including PLSCR1)

Plasmid sources

WT PLSCR1, IFITM2, and LY6E over-expression lentiviral plasmids originated from the

SCRPSY library [196]. CH25H originated from the Thermo Scientific Precision LentiORF Collection.

RRL.sin.cPPT.CMV/Flag-NCOA7 variant 6.IRES-puro.WPRE (CG494) was a gift from Caroline Goujon

(Addgene plasmid # 139447) [161].

Plasmid cloning

N-terminal 3x FLAG-tagged PLSCR1, C-terminal 3x FLAG-tagged PLSCR1, and PLSCR1

H262Y in the SCRPSY vector, used in **Figs 4B and 7D**, were generated by designing and ordering

large dsDNA gene blocks of PLSCR1 that contained the desired mutations from IDT. These gene

blocks were cloned into the PLSCR1-SCRPSY vector [196] and confirmed by sequencing using

Plasmidsaurus (see **Supp Table 18** for sequences).

PLSCR1 variants of interest (**Fig 7C and Supp Fig 12**) were cloned by site directed

mutagenesis using the NEB Quick Protocol for Q5® Site-Directed Mutagenesis Kit (cat. E0552)

according to the manufacturer's instructions. For G20V (rs79551579), we used the primers

JLP621-O-01/02; for A29T (rs41267859), we used the primers JLP621-O-03/04; for I110fs

rs749938276, we used the primers JLP621-O-07/08; and for H262Y (rs343320), we used the

primers JLP621-O-09/10. Primer sequences are listed in **Supp Table 20**.

Lentivirus production

Lentivirus were generated in Lenti-X 293T cells by transfecting 200 ng VSV-G plasmid, 700

ng Gag-Pol plasmid, and 1100 ng plasmid of interest with lipofectamine 2000 in DMEM

supplemented with 5% FBS. Media was changed 4-6 hours later, and lentivirus harvested at 24

and 48 hours. Lentivirus-containing supernatants from both timepoints were pooled, then filtered

through a 0.45 µM filter before aliquoting into 2 mL tubes and freezing at -80°C until use.

Cell transduction

0.3 million cells were transduced in suspension in a 12-well plate using a serial dilution of

each lentivirus in DMEM (Fisher Scientific, cat. #11995065) supplemented with 5% FBS (HyClone

Laboratories, Lot. #KTH31760) and 4 µg/mL polybrene (Millipore, cat. #TR-1003-G). The cells

were then spinoculated at 37°C for 1 hour at 1000 x g. The following day, the cells were split into

two 6-well plates. After 24 hours, one of the duplicates was treated with 2 µg/mL puromycin to

select for transduced cells. Further experiments were carried out using the cells that had

approximately 30% transduction before selection.

SARS-CoV-2 infection

In **Fig 7D**, Huh-7.5 and A549+ACE2 cells were plated at 6,000 cells/well and 3,000

cells/well, respectively, in 100 µL of media (DMEM, 10% FBS, 1X NEAA) in 96-well plates. The

following day, cells were treated with IFN (10pM for Huh-7.5 cells and 20pM for A549+ACE2 cells).

On the third day, the Huh-7.5 cells were infected by adding 10 µL of media containing 0.1 µL of

virus (titer of 3.4×10^6 PFU/ml, as measured on Vero E6 cells) to each well and the A549+ACE2

cells were infected by adding 10 µL of media containing 1 µL of virus to each well. After adding

the virus, the plates were spun at 200 g for 5 minutes and incubated at 37°C. Plates were

harvested the next day by fixing and staining as described above

In **Fig 7C** and **Supp Figs 8A and 12**, A549+ACE2 cells were plated at 7,000 cells/well in 100

µL of media (DMEM, 10% FBS, 1X NEAA) in 96-well plates. The following day, cells were infected

by adding 10 μL of media containing 1 μL of virus (titer of 3.4 × 106 PFU/ml, as measured on Vero

E6 cells) to each well as described above.

IF staining and imaging

In **Figs 7C-E and Supp Figs 8A and 12**, cells were stained for IF as described in the *Unbiased*

arrayed CRISPR KO screening section, with different primary and secondary antibody solutions.

The primary antibody solution was a 1:3000 dilution of rabbit anti-SARS-CoV-2 nucleocapsid

polyclonal antibody (Genetex, cat. #GTX135357) and a 1:500 dilution of 4D2 mouse anti-PLSCR1

antibody (Millipore Sigma, cat. #MABS483) in PBS, and the secondary antibody solution was

1:1000 goat anti-rabbit 594 (ThermoFisher, cat. #A-11012) or 1:1000 goat anti-rabbit Alexa

Fluor™v 647 (ThermoFisher, cat. #A-21245), 1:1000 goat anti-mouse Alexa Fluor™ 488

(Thermofisher, cat. #A-11001), and 1:1000 Hoechst dye (ThermoFisher, cat. #62249) in PBST.

1178

SARS-CoV-2 infection of human SV40-fibroblasts-ACE2

Patients

Written informed consent was obtained in the country of residence of the patients, in

accordance with local regulations, and with institutional review board (IRB) approval.

Experiments were conducted in the United States in accordance with local regulations and with

the approval of the IRB. Approval was obtained from the Rockefeller University Institutional

Review Board in New York, USA (protocol no. JCA-0700).

PLSCR1 H262Y was genotyped using the primers JLP616-O-1-4 listed in **Supp Table 20**.

Generation of human SV40-fibroblasts ACE2 stable cell lines

ACE2 cDNA was inserted with In-Fusion cloning kit (Takara Bio) and using the XhoI and

BamHI restriction sites into linearized pTRIP-SFFV-CD271-P2A in accordance with the

manufacturers' instructions. We checked the entire sequence of the *ACE2* cDNA in the plasmid

by Sanger sequencing. Then, HEK293T cells were dispensed into a six-well plate at a density of

8 x10^5 cells per well. On the next day, cells were transfected with pCMV-VSV-G (0.2 μg), pHXB2-

env (0.2 μg; NIH-AIDS Reagent Program; 1069), psPAX2 (1 μg; Addgene plasmid no. 12260) and

pTRIP-SFFV-CD271-P2A-ACE2 (1.6 μg) in Opti-MEM (Gibco; 300 μL) containing X-tremeGene-9

(Sigma Aldrich; 10 µL) according to the manufacturers' protocol. After transfection for 6 h, the

medium was replaced with 3 mL fresh culture medium, and the cells were incubated for a further

24 h for the production of lentiviral particles. The viral supernatant was collected and passed

through a syringe filter with 0.2 µm pores (Pall) to remove debris. Protamine sulfate (Sigma;

10 µg/mL) was added to the supernatant, which was then used immediately or stored at −80 °C

until use.

For the transduction of SV40-fibroblasts with ACE2, 5 x10^5 cells per well were seeded in

six-well plates. Viral supernatant was added (500 µL per well). The cells were then further

incubated for 48 h at 37 °C. Cells were keep in culture and after 8 days, transduction efficiency

was evaluated by CD271 surface staining with CD271 AlexaFluor 647, 1:200 dilution (BD

Pharmigen 560326). MACS-sorting was performed with CD271 positive selection beads (Miltenyi

Biotec) if the proportion of CD271-positive cells was below 80% [8, 177].

Infection

5,000 cells per well were seeded in a 96-well plate and infected the next day with SARS-

CoV-2 at MOI = 0.05, using a titer determined on Vero E6 cells. Cells were fixed at 2 dpi, stained

and imaged as described in the *Unbiased arrayed CRISPR KO screen* section.

1211

Acknowledgments

We thank Georgia McClain for reading and editing this manuscript. We thank the staff at

the Laboratory of Virology and Infectious Disease: Ellen Castillo, Michela De Santis, Arnella Norris,

Aileen O'Connell, Santa Maria Pecoraro Di Vittorio, Glen Santiago, and Sonia Shirley. Ching-Wen

Chang and Lihong Liu (Columbia University, NY, USA), and Theodora Hatziioannou, Viren Baharani

and Paul D. Bieniasz (The Rockefeller University, NY, USA) generously provided plasmids,

instructions to generate SARS-CoV-2 spike-pseudotyped, single-cycle, replication-defective HIV-1

viruses [120, 194, 195], and assisted us with data interpretation. Oded Danziger and Brad R.

Rosenberg (Department of Microbiology at the Icahn School of Medicine at Mount Sinai, NY, USA)

kindly provided the A549-ACE2 cells [11] used in this study. FACS was conducted at the Flow

Cytometry Resource Center at Rockefeller University. mRNA-seq was performed by the Genomics

Resource Center at The Rockefeller University and by Novogene. Confocal microscopy was

performed in the Rockefeller University's Bio-Imaging Resource Center, RRID:SCR_017791. We

thank Ankit Patel, Sales Manager at Horizon Discovery.

Trim Galore was developed at The Babraham Institute by @FelixKrueger, now part of Altos

Labs. The Genotype-Tissue Expression (GTEx) Project [49] was supported by the Common Fund of

the Office of the Director of the National Institutes of Health, and by NCI, NHGRI, NHLBI, NIDA,

NIMH, and NINDS. The data used for the analyses described in this manuscript were obtained

from   the   EMBL-European   Bioinformatics   Institute   portal,

"https://www.ebi.ac.uk/gxa/experiments/E-MTAB-5214/Results", on October 1st, 2020.

Work in the Laboratory of Virology and Infectious Disease was supported by NIH grants

P01AI138398-S1, 2U19AI111825, R01AI091707-10S1, and R01AI161444; a George Mason

University Fast Grant; the G. Harold and Leila Y. Mathers Charitable Foundation; the Meyer

Foundation; the Pilot Project Robertson Therapeutic Development Fund at The Rockefeller

University; and the Bawd Foundation. The Laboratory of Human Genetics of Infectious Diseases

is supported by the Howard Hughes Medical Institute, the St. Giles Foundation, NIH

(R01AI163029, UL1TR001866), the French National Research Agency (ANR) under the

"Investments for the Future" program (ANR-10-IAHU-01), the Integrative Biology of Emerging

Infectious Diseases Laboratory of Excellence (ANR-10-LABX-62-IBEID), the French Foundation for

Medical Research (FRM) (EQU201903007798), ANR GENVIR (ANR-20-CE93-003) project, the ANR-

RHU COVIFERON Program (ANR-21-RHUS-08), the French Foundation for Medical Research (FRM)

(EQU201903007798), the European Union's Horizon 2020 research and innovation program

under grant agreement No. 824110 (EASI-genomics), the HORIZON-HLTH-2021-DISEASE-04

program under grant agreement 01057100 (UNDINE), the Square Foundation, William E. Ford,

General Atlantic's Chairman and Chief Executive Officer, Gabriel Caillaux, General Atlantic's Co-

President, Managing Director and Head of Business at EMEA, and the General Atlantic

Foundation. J.L.P. was supported by the Francois Wallace Monahan Postdoctoral Fellowship at

The Rockefeller University and the European Molecular Biology Organization Long-Term

Fellowship (ALTF 380-2018). G.P. was supported by the James H. Gilliam Fellowship for Advanced

Study from the Howard Hughes Medical Institute and the Graduate Research Fellowship Program

from the National Science Foundation (FAIN 1946429). M.B. was supported by a Swiss National

Science Foundation fellowship (P500PB_203007).

1254

Financial Disclosure

1255

1256 The funders had no role in study design, data collection and analysis, decision to publish,

1257 or preparation of the manuscript.

1258

Data and code availability

1259

mRNA-seq data is available on the Sequence Read Archive (SRA), hosted by the National

Center for Biotechnology Information (NCBI), under BioProject PRJNA1138251 and BioSamples

SAMN42689767-SAMN42689770 corresponding to **Supp Tables 2 and 3**, BioSamples

SAMN42689771- SAMN42689774 corresponding to **Supp Tables 4 and 5**, BioSamples

SAMN42689775- SAMN42689776 corresponding to **Supp Tables 14 and 15**, BioSamples

SAMN42689777- SAMN42689778 corresponding to **Supp Tables 16 and 17**.

Code and Supporting Information are available on Dryad (DOI:

10.5061/dryad.6q573n65k).

Tables

1268

1269 Table 1. PLSCR1 variants associated with severe COVID-19 in a GWAS [25, 26]

1270

GWAS p-value	rsID	Genomic Coordinate (GRCh38)	Nucleotide Change	Gene	Functional Consequence
7.52E-07	rs116553931	chr3:146430956	C:T	PLSCR2	intron variant
1.08E-07	rs454645	chr3:146514682	C:T	PLSCR1	Downstream

P-value	rsID	Position	Alleles	Gene	transcript variant
5.43E-08	rs343320	chr3:146517122	G:A	PLSCR1	His262Tyr
8.21E-08	rs343318	chr3:146518204	T:C	PLSCR1	intron variant
1.52E-07	rs343317	chr3:146518374	A:G	PLSCR1	intron variant
1.00E-07	rs186910	chr3:146520241	A:G	PLSCR1	intron variant
1.13E-07	rs173150	chr3:146520256	A:T	PLSCR1	intron variant
1.35E-07	rs71302408	chr3:146520389	T:C	PLSCR1	intron variant
7.06E-08	rs343316	chr3:146521151	A:G	PLSCR1	intron variant
4.64E-08	rs343314	chr3:146522652	C:T	PLSCR1	intron variant
7.46E-08	rs343312	chr3:146522970	G:A	PLSCR1	intron variant

1271

Figure Legends

Figure 1. An unbiased arrayed CRISPR KO screen identifies IFN-dependent and IFN-independent genes influencing SARS-CoV-2 infection.

A. mRNA-seq comparison between Huh-7.5 and Calu-3 cells, focusing on a subset of 224 ISGs, in response to 24 h SARS-CoV-2 infection MOI 0.03. Red diamond, PLSCR1 RNA level. Viral RNA levels were comparable in both cell lines (not shown). ****, $p \leq 0.0001$; two-tailed t-test.

B.　Cells were treated with 0.5 nM IFN-α2a for 24 h. mRNA-seq analysis as in **(A)**.

C.　Huh-7.5-Cas9 cells were pretreated with different amounts of IFN-α2a, then infected with

SARS-CoV-2 for 24 h followed by IF staining for SARS-CoV-2 N protein; n = 6 separate wells

infected on the same day; error bars represent SEM; ****, $p \leq 0.0001$; two-tailed t-test.

D.　Diagram of the arrayed CRISPR KO screen method.

E.　Median SARS-CoV-2 positive cell percentages determined by IF staining for the indicated

control genes. The control genes were distributed across 28 separate wells (without IFN

pretreatment) or across 20 separate wells (with IFN pretreatment) for each screened

library plate. NTC: non-targeting control.

F.　The virus level (percentage of infected cells normalized and z-score calculated) is plotted

for 24 h 0 pM (y axis) or 1 pM (x axis) IFN-α2a pretreatment followed by 24 h infection (n

≥ 5). The genes were categorized as ISG or other based on mRNA-seq of IFN-α2a-treated

cells as in **(B)**. ISGs were defined by a fold change ≥ 2 and padj ≤ 0.05 in the IFN-treatment

versus untreated pairwise comparison.

The data underlying this Figure can be found in **Supp Table 1**.

1294

1295

Figure 2. Cellular pathways influencing SARS-CoV-2 infection.

GSEA was performed on the screen data (**Supp Table 10**). The individual genes for some

of the top pathways are shown here. The full names of the plotted pathways are as

follows: translation initiation, "REACTOME EUKARYOTIC TRANSLATION INITIATION";

negative rRNA regulation, "REACTOME NEGATIVE EPIGENETIC REGULATION OF RRNA

EXPRESSION"; cholesterol biosynthesis, "REACTOME REGULATION OF CHOLESTEROL

BIOSYNTHESIS BY SREBP SREBF"; pre-mRNA processing, "REACTOME PROCESSING OF

CAPPED INTRON CONTAINING PRE MRNA"; cellular respiration, "WP ELECTRON

TRANSPORT CHAIN OXPHOS SYSTEM IN MITOCHONDRIA"; RNA Pol III transcription,

"REACTOME RNA POLYMERASE III TRANSCRIPTION INITIATION FROM TYPE 3 PROMOTER."

Pathways are from the Reactome [188] and Wikipathways [190] databases.

The data underlying this Figure can be found in **Supp Table 1**.

1308

Figure 3. Human genes significant in human genetics studies on COVID-19 patients

and in functional genetic screens in cell culture

Overlap between genes considered significant by human genetics studies on COVID-19

patients [5, 6, 8, 25, 26, 81-99] and functional genetic screens for genes influencing SARS-

CoV-2 infection in cell culture [11-13, 15, 18, 19, 28, 31, 64-80]. The full list of genes is

available in **Supp Table 11**, and a summary is in **Supp Table 12**. Please note that not all

genes represented here will influence COVID-19 outcomes. We apologize to the many

colleagues whose work was not cited and discussed.

1317

Figure 4. PLSCR1 is a highly effective anti-SARS-CoV-2 effector ISG contributing to

intrinsic immunity in the absence of IFN.

A. Cells were pretreated with a JAK-STAT inhibitor (InSolution 1 µM) for 2 h, followed by IFN-

α2a (10 pM Huh-7.5 or 20 µM A549-ACE2) for 24 h and were infected with SARS-CoV-2 for

24 h followed by IF staining for viral N protein. Huh-7.5 infection using an MOI of 0.5 (titer

determined by focus forming assay on Huh-7.5 WT cells). A549-ACE2 infection using an

MOI of 0.01 (titer determined by focus forming assay on A549-ACE2 WT cells). The

percentage of SARS-CoV-2 positive cells is plotted. n = 4 separate wells infected on the

same day.

B. Cells were reconstituted with the indicated proteins by stable transduction with

lentiviruses, then infected as in **(A)**. n = 4 separate wells infected on the same day.

C. Cells were co-cultured as indicated (50:50 mix), then infected as in **(A)** and the % infection

of each cell type was determined. n = 4 separate wells infected on the same day; ****, p

≤ 0.0001; two-tailed t-test.

The data underlying this Figure can be found in **Supp Table 1**.

1333

 Figure 5. PLSCR1 restricts spike-mediated SARS-CoV-2 entry.

A. A549-ACE2 cells were IF stained using an anti-PLSCR1 antibody (white) and Hoechst
33342 nuclear staining (blue) and imaged at 63 X magnification on a confocal
microscope.

B. Focus forming assays: SARS-CoV-2 N IF (red) and Hoechst 33342 nuclear staining (blue)
on similarly infected WT or PLSCR1 KO Huh-7.5 and A549-ACE2 cells after 2 and 3 d,
respectively.

C. Quantification of **(B)**.

D. Huh-7.5 WT and PLSCR1 KO cells electroporated with SARS-CoV-2 replicon which
produces a secreted luciferase. Luciferase activity assayed 24 hours after
electroporation. n = 36 separate wells from a single electroporation event. Error bars
represent sd; ns = non-significant; two-tailed t-test.

E-F. Transduction of A549-ACE2 cells with an HIV-based replicon expressing the
nanoluciferase pseudotyped with VSV-G or SARS-CoV-2 spike, respectively. n = 5
separate wells transduced on the same day. Nanoluciferase signal measured 2 dpi. Error
bars represent sd; ns, non-significant; **, $p \leq 0.01$; two-tailed t-test.

G-I. A549 cells WT, expressing ACE2, or expressing ACE2-TMPRSS2 as indicated were
transfected with PLSCR1 or non-template control (NTC) siRNAs as indicated for 3 d and
infected with SARS-CoV-2 for 1 d. SARS-CoV-2 N was stained by IF and the percentage of
positive cells was determined by imaging. n = 6 separate wells infected on the same day.
Error bars represent sd; ns, non-significant; **, $p \leq 0.01$; ***, $p \leq 0.001$; two-tailed t-
test.

J-M. HEK293T cells expressing ACE2 or ACE2-TMPRSS2, as indicated, were transfected with
siRNA knockdown of PLSCR1 or non-template control (NTC), as indicated, for 3 d and
transduced with an HIV-based replicon expressing the nanoluciferase pseudotyped with
VSV-G or SARS-CoV-2 spike, as indicated for 2 d. n = 3 separate wells transduced on the
same day. Error bars represent sd; ns, non-significant; **, $p \leq 0.01$; two-tailed t-test.

The data underlying this Figure can be found in **Supp Table 1**.

1362

Figure 6. Newer variants of SARS-CoV-2 are less restricted by PLSCR1

A-E. Infection of Huh-7.5 cells with SARS-CoV-2 (parental) or its descendant variants, Beta,

Delta, Omicron BA.5, and Omicron XBB.1.5 for 24 hours. SARS-CoV-2 N was stained by

IF and the percentage of positive cells determined by imaging. n = 10 separate wells

infected on the same day. Error bars represent sd; ****, $p < 0.0001$; two-tailed t-test.

F.   Ratio of WT/KO percent infection from A-E. Error bars represent sd. ns, non-significant;

***, $p \leq 0.001$; ****, $p \leq 0.0001$; one-way ANOVA.

G.   Focus forming assay on Huh-7.5 WT and PLSCR1 KO cells infected with approximately 50

FFU of SARS-CoV-2 variants as indicated. FFUs were determined on PLSCR1 KO Huh-7.5

cells. Representative images.

H.   Foci from (**G**) were counted, and then a ratio of WT-to-KO plotted for each SARS-CoV-2

variant. n = 6 separate wells infected on the same day. Error bars represent sd. **, $p \leq$

0.01; one-way ANOVA.

I.   Virus production over an infectious time course (growth curve) for the parental SARS-

CoV-2 strain. n = 2 to 3 separate wells infected on the same day. Error bars represent

sd. ns, non-significant, *, $p \leq 0.05$, ***, $p \leq 0.001$, two-tailed t-test on the log10-

transformed values.

J.   As in (**I**) with the Omicron (XBB.1.5) strain.

The data underlying this Figure can be found in **Supp Table 1**.

1382

Figure 7. PLSCR1 p.His262Tyr, which associates with severe COVID-19, leads to

higher SARS-CoV-2 infection in cell culture.

A.   Protein diagram of PLSCR1. Domain coordinates from UniProt [197].

B.   CADD MAF plot showing the common variants of the PLSCR1 gene reported in gnomAD

[132, 133]. The plot displays the Combined Annotation Dependent Depletion (CADD)

scores, indicating the predicted deleteriousness of each variant, against the Minor Allele

Frequency (MAF). Fs, frameshift.

C.   A549-ACE2 PLSCR1 KO cells were transduced to stably and ectopically express the

indicated PLSCR1 variants. The cells were then infected for 24 h with SARS-CoV-2. SARS-

CoV-2 N was stained by IF and the percentage of positive cells determined by imaging. n

= 4 separate wells infected on the same day. Error bars represent sd; **, p ≤ 0.01; ****, p

< 0.0001; one-way ANOVA. Fs, frameshift.

D. A549-ACE2 cells, WT and PLSCR1 KO, stably expressing N-terminal FLAG-tagged Firefly

Luciferase (Fluc), N-terminal FLAG-tagged PLSCR1, or N-terminal FLAG-tagged PLSCR1

H262Y mutant and infected with SARS-CoV-2 for 24 hours. SARS-CoV-2 N was stained by

IF and the percentage of positive cells determined by imaging. n = 15 separate wells

infected on the same day. Error bars represent sd; ****, p ≤ 0.0001; two-tailed t-test.

E. SV40-Fibroblast-ACE2 cells, genotype as indicated, infected for 24 hours with SARS-CoV-2.

n = 8 separate wells infected on the same day. ns, non-significant; *, p ≤ 0.05; **, p ≤ 0.01;

****, p ≤ 0.0001; one-way ANOVA.

The data underlying this Figure can be found in **Supp Table 1**.

1404

Supplementary Figure Legends

1405

Supplementary Figure 1.

1406

A. A schematic of SARS-CoV-2 entry and the sites where known ISG entry restriction factors

function is shown, adapted from [1]. CD74 suppresses endolysosomal cathepsins, enzymes

that process certain viral glycoproteins to make them fusion-competent [148, 198]. CH25H

facilitates the sequestration of accessible cholesterol, which results in decreased virus-cell

membrane fusion and viral entry [14, 63]. NCOA7 accelerates the acidification of the

lysosome, leading to the degradation of viral antigens [161, 162]. The role of IFITM1-3

proteins in SARS-CoV-2 entry has been the subject of numerous studies, which have

sometimes yielded varying conclusions [1, 122, 199-205]. However, it appears that (i)

IFITM2 restricts WT SARS-CoV-2 endosomal entry [122], (ii) recent SARS-CoV-2 variants

may evade IFITM2 [199], and (iii) IFITM3 KO mice are hyper-susceptible to WT SARS-CoV-2

[206]. LY6E and PLSCR1 restrict virus-cell membrane fusion at the endosome through

unknown mechanisms, see [19, 121] and this study.

B. TMPRSS2 western blot in Huh-7.5-Cas9 cells, both WT and TMPRSS2 KO, conducted as in
the whole-genome arrayed screen (**Fig 1**), and in A549-ACE2 (A549 + A) and A549-ACE2-
TMPRSS2 (A549 + A/T) cells.

1422

Supplementary Figure 2.
A. Model of protein levels of the gene product over time after complete KO, considering the
Huh-7.5 cell division rate (1.7 times per day) and protein half-lives in hepatocytes from
[207].
B. Nuclei count (z-score) in arrayed genetic screen without SARS-CoV-2 infection. Examples
of VCP (essential gene) and PLSCR1 (SARS-CoV-2 antiviral hit) are plotted.
C. Volcano plot of Huh-7.5 cells mRNA-seq treated with 0.5 nM IFN-α2a for 24 h as in **Fig 1B**.
D. Gene set enrichment analysis conducted on the arrayed CRISPR KO screens data
represented in **Figs 1 and 2**. Description of the top pathways ranked by p-value for each
quadrant. Databases: [1]Reactome; [2]WikiPathways; [3]Pathway Interaction Database; [4]KEGG;
[5]Biocarta.
The data underlying this Figure can be found in **Supp Table 1**.

1435

Supplementary Figure 3
A. Occurrence of human genes interacting with SARS-CoV-2 drawn from a selection of 67
large-scale studies. The occurrence reflects the number of independent studies finding
each gene as significant.
B. Upset plot on data as in **(A)**, showing the overlap in significant genes in large-scale SARS-
CoV-2 studies by category.

1442

1443   Supplementary Figure 4.
1444   Cross comparison between proviral and antiviral hits reported in this study and fifteen
1445   published whole-genome pooled screens [19, 31, 64-67, 69-71, 74, 75, 77-80]. P-value:
1446   Fisher's exact test.

1447

Supplementary Figure 5.

A. Western blot on PLSCR1 KO cells against PLSCR1 (green) and ß-actin (red).

B. Cells as indicated were seeded at similar density, treated or not with Blasticidin (used here

as a control to decrease cell viability), and cultured for 4 days before resazurin cell viability

assay. n = 4 independent wells.

C. Western blot on PLSCR1 WT and KO A549-ACE2 cells against PLSCR1 (green) and ß-actin

(red).

D. Single-cell RNA-seq data from [118], showing the *PLSCR1* mRNA expression in some SARS-

CoV-2 target cells.

The data underlying this Figure can be found in **Supp Table 1**.

1458

Supplementary Figure 6.

A. Huh-7.5 WT and PLSCR1 KO cells were infected with 17,000 FFU of CHIKV 181/25 mKate2

for 12 or 24 hours, then fixed, IF stained for nuclei, and the percentage of positive cells

determined by imaging for mKate2 reporter signal. n = 12 separate wells infected on the

same day. Error bars represent sd. ns, non-significant; two-way ANOVA.

B. Huh-7.5 cells, PLSCR1 KO as indicated, were pretreated with different amounts of IFN-α2a,

then infected with 17,000 FFU of CHIKV-mKate for 12 h; n = 7 separate wells infected on

the same day; error bars represent sd.

C-J. Cells were treated for 24 h by IFN, as indicated, followed by RT-qPCRs on ISGs.

The data underlying this Figure can be found in **Supp Table 1**.

1469

Supplementary Figure 7.

A. Western blot against PLSCR1 and RPS11. Cas-9-expressing Huh-7.5 cells were transfected

with 4-gRNA pools targeting PLSCR1 or non-template control (as indicated) and cells were

in culture for 7 d.

B. Quantification of bands intensity in **(A)**.

C. mRNA-seq on PLSCR1 KO Huh-7.5 cells as in **(A,B)**.

The data underlying this Figure can be found in **Supp Table 1**.

1477

Supplementary Figure 8.

A. A549-ACE2 PLSCR1 KO cells were transduced to stably and ectopically express the

indicated genes. The cells were then infected for 24 h with parental SARS-CoV-2. SARS-

CoV-2 N was stained by IF and the percentage of positive cells determined by imaging. n

= 4 separate wells infected on the same day. Error bars represent sd; **, $p \leq 0.01$; ****, p

< 0.0001; one-way ANOVA.

B. Normalised reads count from mRNA-seq on A549 cells treated or not with 0.5 nM IFN-α2a

for 24 h.

C. Normalised reads count from mRNA-seq on Huh-7.5 cells treated or not with 0.5 nM IFN-

α2a for 24 h.

The data underlying this Figure can be found in **Supp Table 1**.

1489

Supplementary Figure 9.

A-I. Infection of A549-ACE2 cells with viruses as indicated. IF staining was used for IAV WSN,

hCoV-OC43 and SARS-CoV-2, otherwise a fluorescent reporter was used. Percentage of

virus positive cells was determined by imaging. n = six separate wells infected on the

same day, error bars represent SD; ns, non-significant; **, $p \leq 0.01$; ***, $p \leq 0.001$; ****,

$p \leq 0.0001$; ****, $p \leq 0.0001$; two-tailed t-test.

J-M. Infection of Huh-7.5 cells with viruses as indicated. siRNA knockdown of PLSCR1 vs non-

template control (NTC). n = four to sixteen separate wells infected on the same day; ns,

non-significant; ****, $p \leq 0.0001$; two-tailed t-test.

The data underlying this Figure can be found in **Supp Table 1**.

1500

1501

Supplementary Tables

Available on Dryad (DOI: 10.5061/dryad.6q573n65k).

Supplementary Table 1.

Data underlying Figures 1ABCEF, 2, 4ABC, 5CDEFGHIJKLM, 6ABCDEFHIJ, 7BCDE, S2ABC, S5BD, S6ABCDEFGHIJ, S7BC, S8ABC, S9ABCDEFGHIJKLM, S14

Supplementary Table 2.

mRNA-seq on SARS-CoV-2 infected Calu-3 and Huh-7.5 cells, Normalized reads, related to **Fig 1A**.

Supplementary Table 3.

mRNA-seq on SARS-CoV-2 infected Calu-3 and Huh-7.5 cells, Differential Gene Expression.

Supplementary Table 4.

mRNA-seq on IFN-treated Calu-3 and Huh-7.5 cells, Normalized reads, related to **Fig 1B**, and **Supp Figs 8C and 13.**

Supplementary Table 5.

mRNA-seq on IFN-treated Calu-3 and Huh-7.5 cells, Differential Gene Expression, related to **Fig 1F** and **Supp Fig 2C**.

Supplementary Table 6.

Genes expressed in three cell lines relevant for SARS-CoV-2 research (A549, Calu-3, Huh-7.5 cells) and human lung cells [47-49].

1556   Supplementary Table 7.

1557   Arrayed CRISPR KO screen, raw data.

1558

Supplementary Table 8.

Arrayed CRISPR KO screen, analyzed data. Related to **Fig. 1F** and **Supp Fig 2B**. Columns are

as follow: *Stat.W1_SARS2_IFN_0pM*: cell growth z-score determined by nuclear staining in

samples with no interferon pretreatment and with SARS-CoV-2 infection.

Stat.W1_SARS2_IFN_1pM: cell growth z-score determined by nuclear staining in samples with

interferon pretreatment and with SARS-CoV-2 infection. *Stat.W1_Uninfected_IFN_0pM*: cell

growth z-score determined by nuclear staining in samples with no interferon pretreatment and

no SARS-CoV-2 infection. *Stat.W2_SARS2_IFN_0pM*: SARS-CoV-2 infection z-score in samples with

no interferon pretreatment and with SARS-CoV-2 infection. *Stat.W2_SARS2_IFN_1pM*: SARS-CoV-

2 infection z-score in samples with interferon pretreatment and with SARS-CoV-2 infection.

Stat.W2_Uninfected_IFN_0pM: SARS-CoV-2 infection z-score in samples with no interferon

pretreatment and no SARS-CoV-2 infection. Index: random index used for plotting. RNAseq:

classification of the gene as an interferon-stimulated gene ("ISG") or not ("NS") based on the

accompanying mRNA-seq data.

1573

1574   Supplementary Table 9.

1575   Arrayed CRISPR KO screen, summary table.

1576

1577   Supplementary Table 10.

1578   GSEA on arrayed CRISPR KO screen. Related to **Supp Fig 2D**.

1579

1580   Supplementary Table 11.

1581   Consolidated list of human genes classified as hits in selected SARS-CoV-2 studies, full

1582   table. Related to **Fig 3** and **Supp Fig 3.**

1583

1584   Supplementary Table 12.

1585   Consolidated list of human genes classified as hits in selected SARS-CoV-2 studies,

1586 summary table. Related to **Fig 3** and **Supp Fig 3.**

1587

1588   Supplementary Table 13.

1589   Pathway analysis of hits from our arrayed screen alongside fifteen unbiased pooled

1590 screens [19, 31, 64-67, 69-71, 74, 75, 77-80] using the STRING database [208]. Related to **Supp**

1591 **Fig 4**.

1592

1593   Supplementary Table 14.

1594   mRNA-seq on PLSCR1 KO Huh-7.5 cells, Normalized reads, related to **Supp Fig 7C**.

1595

1596   Supplementary Table 15.

1597   mRNA-seq on PLSCR1 KO Huh-7.5 cells, Differential Gene Expression, related to **Supp Fig**

1598 **7C**.

1599

1600   Supplementary Table 16.

1601   mRNA-seq on IFN-treated A549 cells, Normalized reads, related to **Supp Fig 8B**.

1602

1603   Supplementary Table 17.

1604   mRNA-seq on IFN-treated A549 cells, Differential Gene Expression, related to **Supp Fig 8B**.

1605

1606   Supplementary Table 18.

1607   Plasmids used in this study.

1608

1609   Supplementary Table 19.

1610   Gene fragments used in this study.

1611

1612   Supplementary Table 20.

1613   Primers used in this study.

1614

1615   ## Supporting Information

1616   Available on Dryad (DOI: 10.5061/dryad.6q573n65k).

1617

1618   ### Supporting Information 1

1619   Raw images, related to **Supp Figs 1B, 5AC, 7A, 11**.

1620

1621   ### Supporting Information 2

1622   Flow cytometry gating strategy, related to **Supp Fig 10**.

1623

1624   ### Supporting Information 3

1625   Flow cytometry raw data, related to **Supp Fig 10**.

1626

1627   ### Supporting Information 4

1628   mRNA-seq raw reads, related to **Supp Tables 2, 3, 4, 5, 14, 15, 16, 17**.

1629

References

1.   Le Pen J, Rice CM. The antiviral state of the cell: lessons from SARS-CoV-2. Curr Opin
Immunol. 2024;87:102426. Epub 20240523. doi: 10.1016/j.coi.2024.102426. PubMed PMID:
38795501.
2.   Bastard P, Zhang Q, Zhang SY, Jouanguy E, Casanova JL. Type I interferons and SARS-CoV-
2: from cells to organisms. Curr Opin Immunol. 2022;74:172-82. Epub 20220125. doi:
10.1016/j.coi.2022.01.003.   PubMed   PMID: 35149239;   PubMed   Central   PMCID:
PMCPMC8786610.
3.   Bastard P, Gervais A, Le Voyer T, Rosain J, Philippot Q, Manry J, et al. Autoantibodies
neutralizing type I IFNs are present in ~4% of uninfected individuals over 70 years old and account
for   ~20%   of   COVID-19   deaths.   Sci   Immunol.   2021;6(62).   Epub   2021/08/21.   doi:
10.1126/sciimmunol.abl4340. PubMed PMID: 34413139.

4.   Lopez J, Mommert M, Mouton W, Pizzorno A, Brengel-Pesce K, Mezidi M, et al. Early nasal
type I IFN immunity against SARS-CoV-2 is compromised in patients with autoantibodies against
type I IFNs. J Exp Med. 2021;218(10). Epub 20210806. doi: 10.1084/jem.20211211. PubMed
PMID: 34357402; PubMed Central PMCID: PMCPMC8352718.

5.   Asano T, Boisson B, Onodi F, Matuozzo D, Moncada-Velez M, Maglorius Renkilaraj MRL, et
al. X-linked recessive TLR7 deficiency in ~1% of men under 60 years old with life-threatening
COVID-19. Sci Immunol. 2021;6(62). doi: 10.1126/sciimmunol.abl4348. PubMed PMID:
34413140; PubMed Central PMCID: PMCPMC8532080.

6.   Zhang Q, Bastard P, Liu Z, Le Pen J, Moncada-Velez M, Chen J, et al. Inborn errors of type I
IFN immunity in patients with life-threatening COVID-19. Science. 2020;370(6515). Epub
20200924. doi: 10.1126/science.abd4570. PubMed PMID: 32972995; PubMed Central PMCID:
PMCPMC7857407.

7.   Bastard P, Rosen LB, Zhang Q, Michailidis E, Hoffmann HH, Zhang Y, et al. Autoantibodies
against type I IFNs in patients with life-threatening COVID-19. Science. 2020;370(6515). Epub
20200924. doi: 10.1126/science.abd4585. PubMed PMID: 32972996; PubMed Central PMCID:
PMCPMC7857397.

8.   Zhang Q, Matuozzo D, Le Pen J, Lee D, Moens L, Asano T, et al. Recessive inborn errors of
type I IFN immunity in children with COVID-19 pneumonia. J Exp Med. 2022;219(8). Epub
20220616. doi: 10.1084/jem.20220131. PubMed PMID: 35708626; PubMed Central PMCID:
PMCPMC9206114.

9.   Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak associated
with a new coronavirus of probable bat origin. Nature. 2020;579(7798):270-3. Epub 20200203.
doi: 10.1038/s41586-020-2012-7. PubMed PMID: 32015507; PubMed Central PMCID:
PMCPMC7095418.

10.   Savan R, Gale M, Jr. Innate immunity and interferon in SARS-CoV-2 infection outcome.
Immunity. 2023;56(7):1443-50. doi: 10.1016/j.immuni.2023.06.018. PubMed PMID: 37437537;
PubMed Central PMCID: PMCPMC10361255.

11.   Danziger O, Patel RS, DeGrace EJ, Rosen MR, Rosenberg BR. Inducible CRISPR activation
screen for interferon-stimulated genes identifies OAS1 as a SARS-CoV-2 restriction factor. PLoS
Pathog. 2022;18(4):e1010464. Epub 20220414. doi: 10.1371/journal.ppat.1010464. PubMed
PMID: 35421191; PubMed Central PMCID: PMCPMC9041830.

12.   Martin-Sancho L, Lewinski MK, Pache L, Stoneham CA, Yin X, Becker ME, et al. Functional
landscape of SARS-CoV-2 cellular restriction. Mol Cell. 2021;81(12):2656-68 e8. Epub 20210413.
doi: 10.1016/j.molcel.2021.04.008. PubMed PMID: 33930332; PubMed Central PMCID:
PMCPMC8043580.

13.   Wickenhagen A, Sugrue E, Lytras S, Kuchi S, Noerenberg M, Turnbull ML, et al. A prenylated
dsRNA sensor protects against severe COVID-19. Science. 2021;374(6567):eabj3624. Epub
20211029. doi: 10.1126/science.abj3624. PubMed PMID: 34581622.

14.   Zang R, Case JB, Yutuc E, Ma X, Shen S, Gomez Castro MF, et al. Cholesterol 25-hydroxylase
suppresses SARS-CoV-2 replication by blocking membrane fusion. Proc Natl Acad Sci U S A.
2020;117(50):32105-13. Epub 20201125. doi: 10.1073/pnas.2012197117. PubMed PMID:
33239446; PubMed Central PMCID: PMCPMC7749331.

15. Kaur R, Tada T, Landau NR. Restriction of SARS-CoV-2 replication by receptor transporter
protein 4 (RTP4). mBio. 2023;14(4):e0109023. Epub 20230629. doi: 10.1128/mbio.01090-23.
PubMed PMID: 37382452; PubMed Central PMCID: PMCPMC10470548.

16. Schoggins JW, Wilson SJ, Panis M, Murphy MY, Jones CT, Bieniasz P, et al. A diverse range
of gene products are effectors of the type I interferon antiviral response. Nature.
2011;472(7344):481-5. Epub 20110410. doi: 10.1038/nature09907. PubMed PMID: 21478870;
PubMed Central PMCID: PMCPMC3409588.

17. Schoggins JW. Interferon-Stimulated Genes: What Do They All Do? Annu Rev Virol.
2019;6(1):567-84. Epub 2019/07/10. doi: 10.1146/annurev-virology-092818-015756. PubMed
PMID: 31283436.

18. Mac Kain A, Maarifi G, Aicher SM, Arhel N, Baidaliuk A, Munier S, et al. Identification of
DAXX as a restriction factor of SARS-CoV-2 through a CRISPR/Cas9 screen. Nat Commun.
2022;13(1):2442. Epub 20220504. doi: 10.1038/s41467-022-30134-9. PubMed PMID: 35508460;
PubMed Central PMCID: PMCPMC9068693.

19. Xu D, Jiang W, Wu L, Gaudet RG, Park ES, Su M, et al. PLSCR1 is a cell-autonomous defence
factor against SARS-CoV-2 infection. Nature. 2023;619(7971):819-27. Epub 20230712. doi:
10.1038/s41586-023-06322-y. PubMed PMID: 37438530; PubMed Central PMCID:
PMCPMC10371867.

20. Kim HS, Lee K, Kim SJ, Cho S, Shin HJ, Kim C, et al. Arrayed CRISPR screen with image-based
assay reliably uncovers host genes required for coxsackievirus infection. Genome Res.
2018;28(6):859-68. Epub 20180430. doi: 10.1101/gr.230250.117. PubMed PMID: 29712754;
PubMed Central PMCID: PMCPMC5991512.

21. Cross BC, Lawo S, Archer CR, Hunt JR, Yarker JL, Riccombeni A, et al. Increasing the
performance of pooled CRISPR-Cas9 drop-out screening. Sci Rep. 2016;6:31782. Epub 20160822.
doi: 10.1038/srep31782. PubMed PMID: 27545104; PubMed Central PMCID: PMCPMC4992892.

22. Kodigepalli KM, Bowers K, Sharp A, Nanjundan M. Roles and regulation of phospholipid
scramblases. FEBS Lett. 2015;589(1):3-14. Epub 20141203. doi: 10.1016/j.febslet.2014.11.036.
PubMed PMID: 25479087.

23. Zhou Q, Zhao J, Stout JG, Luhm RA, Wiedmer T, Sims PJ. Molecular cloning of human
plasma membrane phospholipid scramblase. A protein mediating transbilayer movement of
plasma membrane phospholipids. J Biol Chem. 1997;272(29):18240-4. doi:
10.1074/jbc.272.29.18240. PubMed PMID: 9218461.

24. Dong B, Zhou Q, Zhao J, Zhou A, Harty RN, Bose S, et al. Phospholipid scramblase 1
potentiates the antiviral activity of interferon. J Virol. 2004;78(17):8983-93. doi:
10.1128/JVI.78.17.8983-8993.2004. PubMed PMID: 15308695; PubMed Central PMCID:
PMCPMC506946.

25. Pairo-Castineira E, Rawlik K, Bretherick AD, Qi T, Wu Y, Nassiri I, et al. GWAS and meta-
analysis identifies 49 genetic variants underlying critical COVID-19. Nature. 2023;617(7962):764-
8. Epub 20230517. doi: 10.1038/s41586-023-06034-3. PubMed PMID: 37198478; PubMed
Central PMCID: PMCPMC10208981.

26. Kousathanas A, Pairo-Castineira E, Rawlik K, Stuckey A, Odhams CA, Walker S, et al. Whole-
genome sequencing reveals host factors underlying critical COVID-19. Nature.

2022;607(7917):97-103. Epub 20220307. doi: 10.1038/s41586-022-04576-6. PubMed PMID:
35255492; PubMed Central PMCID: PMCPMC9259496.

27.	Baggen J, Vanstreels E, Jansen S, Daelemans D. Cellular host factors for SARS-CoV-2
infection. Nat Microbiol. 2021;6(10):1219-32. Epub 20210901. doi: 10.1038/s41564-021-00958-
0. PubMed PMID: 34471255.

28.	Hoffmann HH, Sanchez-Rivera FJ, Schneider WM, Luna JM, Soto-Feliciano YM, Ashbrook
AW, et al. Functional interrogation of a SARS-CoV-2 host protein interactome identifies unique
and shared coronavirus host factors. Cell Host Microbe. 2021;29(2):267-80 e5. Epub 20201216.
doi: 10.1016/j.chom.2020.12.009. PubMed PMID: 33357464; PubMed Central PMCID:
PMCPMC7833927.

29.	Li Q, Wu J, Nie J, Zhang L, Hao H, Liu S, et al. The Impact of Mutations in SARS-CoV-2 Spike
on Viral Infectivity and Antigenicity. Cell. 2020;182(5):1284-94 e9. Epub 20200717. doi:
10.1016/j.cell.2020.07.012. PubMed PMID: 32730807; PubMed Central PMCID:
PMCPMC7366990.

30.	Riva L, Yuan S, Yin X, Martin-Sancho L, Matsunaga N, Pache L, et al. Discovery of SARS-CoV-
2 antiviral drugs through large-scale compound repurposing. Nature. 2020;586(7827):113-9.
Epub 20200724. doi: 10.1038/s41586-020-2577-1. PubMed PMID: 32707573; PubMed Central
PMCID: PMCPMC7603405.

31.	Schneider WM, Luna JM, Hoffmann HH, Sanchez-Rivera FJ, Leal AA, Ashbrook AW, et al.
Genome-Scale Identification of SARS-CoV-2 and Pan-coronavirus Host Factor Networks. Cell.
2021;184(1):120-32 e14. Epub 20201209. doi: 10.1016/j.cell.2020.12.006. PubMed PMID:
33382968; PubMed Central PMCID: PMCPMC7796900.

32.	Xia S, Lan Q, Su S, Wang X, Xu W, Liu Z, et al. The role of furin cleavage site in SARS-CoV-2
spike protein-mediated membrane fusion in the presence or absence of trypsin. Signal Transduct
Target Ther. 2020;5(1):92. Epub 20200612. doi: 10.1038/s41392-020-0184-0. PubMed PMID:
32532959; PubMed Central PMCID: PMCPMC7289711.

33.	Xia S, Liu M, Wang C, Xu W, Lan Q, Feng S, et al. Inhibition of SARS-CoV-2 (previously 2019-
nCoV) infection by a highly potent pan-coronavirus fusion inhibitor targeting its spike protein that
harbors a high capacity to mediate membrane fusion. Cell Res. 2020;30(4):343-55. Epub
20200330. doi: 10.1038/s41422-020-0305-x. PubMed PMID: 32231345; PubMed Central PMCID:
PMCPMC7104723.

34.	Yan Y, Chang L, Wang L. Laboratory testing of SARS-CoV, MERS-CoV, and SARS-CoV-2 (2019-
nCoV): Current status, challenges, and countermeasures. Rev Med Virol. 2020;30(3):e2106. Epub
20200417. doi: 10.1002/rmv.2106. PubMed PMID: 32302058; PubMed Central PMCID:
PMCPMC7235496.

35.	Sumpter R, Jr., Loo YM, Foy E, Li K, Yoneyama M, Fujita T, et al. Regulating intracellular
antiviral defense and permissiveness to hepatitis C virus RNA replication through a cellular RNA
helicase, RIG-I. J Virol. 2005;79(5):2689-99. doi: 10.1128/JVI.79.5.2689-2699.2005. PubMed
PMID: 15708988; PubMed Central PMCID: PMCPMC548482.

36.	Matsuyama S, Nao N, Shirato K, Kawase M, Saito S, Takayama I, et al. Enhanced isolation
of SARS-CoV-2 by TMPRSS2-expressing cells. Proc Natl Acad Sci U S A. 2020;117(13):7001-3. Epub
20200312. doi: 10.1073/pnas.2002589117. PubMed PMID: 32165541; PubMed Central PMCID:
PMCPMC7132130.

37.   Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, et al. SARS-
CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease
Inhibitor. Cell. 2020;181(2):271-80 e8. Epub 20200305. doi: 10.1016/j.cell.2020.02.052. PubMed
PMID: 32142651; PubMed Central PMCID: PMCPMC7102627.

38.   Jackson CB, Farzan M, Chen B, Choe H. Mechanisms of SARS-CoV-2 entry into cells. Nat
Rev Mol Cell Biol. 2022;23(1):3-20. Epub 20211005. doi: 10.1038/s41580-021-00418-x. PubMed
PMID: 34611326; PubMed Central PMCID: PMCPMC8491763.

39.   de Weerd NA, Samarajiwa SA, Hertzog PJ. Type I interferon receptors: biochemistry and
biological functions. J Biol Chem. 2007;282(28):20053-7. Epub 20070514. doi:
10.1074/jbc.R700006200. PubMed PMID: 17502368.

40.   de Weerd NA, Nguyen T. The interferons and their receptors--distribution and regulation.
Immunol Cell Biol. 2012;90(5):483-91. Epub 20120313. doi: 10.1038/icb.2012.9. PubMed PMID:
22410872; PubMed Central PMCID: PMCPMC7165917.

41.   Uze G, Schreiber G, Piehler J, Pellegrini S. The receptor of the type I interferon family. Curr
Top Microbiol Immunol. 2007;316:71-95. doi: 10.1007/978-3-540-71329-6_5. PubMed PMID:
17969444.

42.   Perng YC, Lenschow DJ. ISG15 in antiviral immunity and beyond. Nat Rev Microbiol.
2018;16(7):423-39. doi: 10.1038/s41579-018-0020-5. PubMed PMID: 29769653; PubMed Central
PMCID: PMCPMC7097117.

43.   Speer SD, Li Z, Buta S, Payelle-Brogard B, Qian L, Vigant F, et al. ISG15 deficiency and
increased viral resistance in humans but not mice. Nat Commun. 2016;7:11496. Epub 20160519.
doi: 10.1038/ncomms11496. PubMed PMID: 27193971; PubMed Central PMCID:
PMCPMC4873964.

44.   Zhang X, Bogunovic D, Payelle-Brogard B, Francois-Newton V, Speer SD, Yuan C, et al.
Human intracellular ISG15 prevents interferon-alpha/beta over-amplification and auto-
inflammation. Nature. 2015;517(7532):89-93. Epub 20141012. doi: 10.1038/nature13801.
PubMed PMID: 25307056; PubMed Central PMCID: PMCPMC4303590.

45.   Malakhova OA, Kim KI, Luo JK, Zou W, Kumar KG, Fuchs SY, et al. UBP43 is a novel regulator
of interferon signaling independent of its ISG15 isopeptidase activity. EMBO J. 2006;25(11):2358-
67. Epub 20060518. doi: 10.1038/sj.emboj.7601149. PubMed PMID: 16710296; PubMed Central
PMCID: PMCPMC1478183.

46.   Francois-Newton V, Magno de Freitas Almeida G, Payelle-Brogard B, Monneron D, Pichard-
Garcia L, Piehler J, et al. USP18-based negative feedback control is induced by type I and type III
interferons and specifically inactivates interferon alpha response. PLoS One. 2011;6(7):e22200.
Epub 20110714. doi: 10.1371/journal.pone.0022200. PubMed PMID: 21779393; PubMed Central
PMCID: PMCPMC3136508.

47.   Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The Cancer
Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. Nature.
2012;483(7391):603-7. Epub 20120328. doi: 10.1038/nature11003. PubMed PMID: 22460905;
PubMed Central PMCID: PMCPMC3320027.

48.   Klijn C, Durinck S, Stawiski EW, Haverty PM, Jiang Z, Liu H, et al. A comprehensive
transcriptional portrait of human cancer cell lines. Nat Biotechnol. 2015;33(3):306-12. Epub
20141208. doi: 10.1038/nbt.3080. PubMed PMID: 25485619.

49.   Consortium GT. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis:
multitissue gene regulation in humans. Science. 2015;348(6235):648-60. Epub 20150507. doi:
10.1126/science.1262110. PubMed PMID: 25954001; PubMed Central PMCID:
PMCPMC4547484.

50.   Horvath CM, Stark GR, Kerr IM, Darnell JE, Jr. Interactions between STAT and non-STAT
proteins in the interferon-stimulated gene factor 3 transcription complex. Mol Cell Biol.
1996;16(12):6957-64. doi: 10.1128/MCB.16.12.6957. PubMed PMID: 8943351; PubMed Central
PMCID: PMCPMC231699.

51.   Veals SA, Schindler C, Leonard D, Fu XY, Aebersold R, Darnell JE, Jr., et al. Subunit of an
alpha-interferon-responsive transcription factor is related to interferon regulatory factor and Myb
families of DNA-binding proteins. Mol Cell Biol. 1992;12(8):3315-24. doi:
10.1128/mcb.12.8.3315-3324.1992. PubMed PMID: 1630447; PubMed Central PMCID:
PMCPMC364572.

52.   Darnell JE, Jr., Kerr IM, Stark GR. Jak-STAT pathways and transcriptional activation in
response to IFNs and other extracellular signaling proteins. Science. 1994;264(5164):1415-21.
doi: 10.1126/science.8197455. PubMed PMID: 8197455.

53.   Schindler C, Strehlow I. Cytokines and STAT signaling. Adv Pharmacol. 2000;47:113-74.
doi: 10.1016/s1054-3589(08)60111-8. PubMed PMID: 10582086.

54.   Levy DE, Darnell JE, Jr. Stats: transcriptional control and biological impact. Nat Rev Mol
Cell Biol. 2002;3(9):651-62. doi: 10.1038/nrm909. PubMed PMID: 12209125.

55.   Naka T, Narazaki M, Hirata M, Matsumoto T, Minamoto S, Aono A, et al. Structure and
function of a new STAT-induced STAT inhibitor. Nature. 1997;387(6636):924-9. doi:
10.1038/43219. PubMed PMID: 9202127.

56.   Ziegler CGK, Allon SJ, Nyquist SK, Mbano IM, Miao VN, Tzouanas CN, et al. SARS-CoV-2
Receptor ACE2 Is an Interferon-Stimulated Gene in Human Airway Epithelial Cells and Is Detected
in Specific Cell Subsets across Tissues. Cell. 2020;181(5):1016-35 e19. Epub 20200427. doi:
10.1016/j.cell.2020.04.035. PubMed PMID: 32413319; PubMed Central PMCID:
PMCPMC7252096.

57.   Onabajo OO, Banday AR, Stanifer ML, Yan W, Obajemu A, Santer DM, et al. Interferons and
viruses induce a novel truncated ACE2 isoform and not the full-length SARS-CoV-2 receptor. Nat
Genet. 2020;52(12):1283-93. Epub 20201019. doi: 10.1038/s41588-020-00731-9. PubMed PMID:
33077916; PubMed Central PMCID: PMCPMC9377523.

58.   Zhao MM, Yang WL, Yang FY, Zhang L, Huang WJ, Hou W, et al. Cathepsin L plays a key role
in SARS-CoV-2 infection in humans and humanized mice and is a promising target for new drug
development. Signal Transduct Target Ther. 2021;6(1):134. Epub 20210327. doi: 10.1038/s41392-
021-00558-8. PubMed PMID: 33774649; PubMed Central PMCID: PMCPMC7997800.

59.   Shang J, Wan Y, Luo C, Ye G, Geng Q, Auerbach A, et al. Cell entry mechanisms of SARS-
CoV-2. Proc Natl Acad Sci U S A. 2020;117(21):11727-34. Epub 20200506. doi:
10.1073/pnas.2003138117. PubMed PMID: 32376634; PubMed Central PMCID:
PMCPMC7260975.

60.   Liu T, Luo S, Libby P, Shi GP. Cathepsin L-selective inhibitors: A potentially promising
treatment for COVID-19 patients. Pharmacol Ther. 2020;213:107587. Epub 20200526. doi:

10.1016/j.pharmthera.2020.107587. PubMed PMID: 32470470; PubMed Central PMCID:
PMCPMC7255230.
61.	Pellegrini S, John J, Shearer M, Kerr IM, Stark GR. Use of a selectable marker regulated by
alpha interferon to obtain mutations in the signaling pathway. Mol Cell Biol. 1989;9(11):4605-12.
doi: 10.1128/mcb.9.11.4605-4612.1989. PubMed PMID: 2513475; PubMed Central PMCID:
PMCPMC363606.
62.	Velazquez L, Fellous M, Stark GR, Pellegrini S. A protein tyrosine kinase in the interferon
alpha/beta signaling pathway. Cell. 1992;70(2):313-22. doi: 10.1016/0092-8674(92)90105-l.
PubMed PMID: 1386289.
63.	Wang S, Li W, Hui H, Tiwari SK, Zhang Q, Croker BA, et al. Cholesterol 25-Hydroxylase
inhibits SARS-CoV-2 and other coronaviruses by depleting membrane cholesterol. EMBO J.
2020;39(21):e106057. Epub 20201005. doi: 10.15252/embj.2020106057. PubMed PMID:
32944968; PubMed Central PMCID: PMCPMC7537045.
64.	Zhu Y, Feng F, Hu G, Wang Y, Yu Y, Zhu Y, et al. A genome-wide CRISPR screen identifies host
factors that regulate SARS-CoV-2 entry. Nat Commun. 2021;12(1):961. Epub 20210211. doi:
10.1038/s41467-021-21213-4. PubMed PMID: 33574281; PubMed Central PMCID:
PMCPMC7878750.
65.	Wei LH, Sun Y, Guo JU. Genome-wide CRISPR screens identify noncanonical translation
factor eIF2A as an enhancer of SARS-CoV-2 programmed -1 ribosomal frameshifting. Cell Rep.
2023;42(8):112987. Epub 20230814. doi: 10.1016/j.celrep.2023.112987. PubMed PMID:
37581984.
66.	Wei J, Alfajaro MM, DeWeirdt PC, Hanna RE, Lu-Culligan WJ, Cai WL, et al. Genome-wide
CRISPR Screens Reveal Host Factors Critical for SARS-CoV-2 Infection. Cell. 2021;184(1):76-91 e13.
Epub 20201020. doi: 10.1016/j.cell.2020.10.028. PubMed PMID: 33147444; PubMed Central
PMCID: PMCPMC7574718.
67.	Wang R, Simoneau CR, Kulsuptrakul J, Bouhaddou M, Travisano KA, Hayashi JM, et al.
Genetic Screens Identify Host Factors for SARS-CoV-2 and Common Cold Coronaviruses. Cell.
2021;184(1):106-19 e14. Epub 20201209. doi: 10.1016/j.cell.2020.12.004. PubMed PMID:
33333024; PubMed Central PMCID: PMCPMC7723770.
68.	Song J, Chow RD, Pena-Hernandez MA, Zhang L, Loeb SA, So EY, et al. LRRC15 inhibits SARS-
CoV-2 cellular entry in trans. PLoS Biol. 2022;20(10):e3001805. Epub 20221013. doi:
10.1371/journal.pbio.3001805. PubMed PMID: 36228039; PubMed Central PMCID:
PMCPMC9595563.
69.	Hossain MS, Kerkvliet JG, Hoppe AD. Whole genome CRISPR screening strategy to identify
genes contributing to SARS-CoV-2 spike and VSV-G mediated entry. J Med Virol.
2023;95(9):e29087. doi: 10.1002/jmv.29087. PubMed PMID: 37707319.
70.	Rehfeld F, Eitson JL, Ohlson MB, Chang TC, Schoggins JW, Mendell JT. CRISPR screening
reveals a dependency on ribosome recycling for efficient SARS-CoV-2 programmed ribosomal
frameshifting and viral replication. Cell Rep. 2023;42(2):112076. Epub 20230130. doi:
10.1016/j.celrep.2023.112076. PubMed PMID: 36753415; PubMed Central PMCID:
PMCPMC9884621.
71.	Rebendenne A, Roy P, Bonaventure B, Chaves Valadao AL, Desmarets L, Arnaud-Arnould
M, et al. Bidirectional genome-wide CRISPR screens reveal host factors regulating SARS-CoV-2,

MERS-CoV and seasonal HCoVs. Nat Genet. 2022;54(8):1090-102. Epub 20220725. doi:
10.1038/s41588-022-01110-2. PubMed PMID: 35879413.

72. Pahmeier F, Lavacca TM, Goellner S, Neufeldt CJ, Prasad V, Cerikan B, et al. Identification
of host dependency factors involved in SARS-CoV-2 replication organelle formation through
proteomics and ultrastructural analysis. J Virol. 2023:e0087823. Epub 20231031. doi:
10.1128/jvi.00878-23. PubMed PMID: 37905840.

73. Loo L, Waller MA, Moreno CL, Cole AJ, Stella AO, Pop OT, et al. Fibroblast-expressed
LRRC15 is a receptor for SARS-CoV-2 spike and controls antiviral and antifibrotic transcriptional
programs. PLoS Biol. 2023;21(2):e3001967. Epub 20230209. doi: 10.1371/journal.pbio.3001967.
PubMed PMID: 36757924; PubMed Central PMCID: PMCPMC9910744.

74. Israeli M, Finkel Y, Yahalom-Ronen Y, Paran N, Chitlaru T, Israeli O, et al. Genome-wide
CRISPR screens identify GATA6 as a proviral host factor for SARS-CoV-2 via modulation of ACE2.
Nat Commun. 2022;13(1):2237. Epub 20220425. doi: 10.1038/s41467-022-29896-z. PubMed
PMID: 35469023; PubMed Central PMCID: PMCPMC9039069.

75. Grodzki M, Bluhm AP, Schaefer M, Tagmount A, Russo M, Sobh A, et al. Genome-scale
CRISPR screens identify host factors that promote human coronavirus infection. Genome Med.
2022;14(1):10. Epub 20220127. doi: 10.1186/s13073-022-01013-1. PubMed PMID: 35086559;
PubMed Central PMCID: PMCPMC8792531.

76. Gordon DE, Hiatt J, Bouhaddou M, Rezelj VV, Ulferts S, Braberg H, et al. Comparative host-
coronavirus protein interaction networks reveal pan-viral disease mechanisms. Science.
2020;370(6521). Epub 20201015. doi: 10.1126/science.abe9403. PubMed PMID: 33060197;
PubMed Central PMCID: PMCPMC7808408.

77. Daniloski Z, Jordan TX, Wessels HH, Hoagland DA, Kasela S, Legut M, et al. Identification
of Required Host Factors for SARS-CoV-2 Infection in Human Cells. Cell. 2021;184(1):92-105 e16.
Epub 20201024. doi: 10.1016/j.cell.2020.10.030. PubMed PMID: 33147445; PubMed Central
PMCID: PMCPMC7584921.

78. Chan K, Farias AG, Lee H, Guvenc F, Mero P, Brown KR, et al. Survival-based CRISPR genetic
screens across a panel of permissive cell lines identify common and cell-specific SARS-CoV-2 host
factors. Heliyon. 2023;9(1):e12744. Epub 20221230. doi: 10.1016/j.heliyon.2022.e12744.
PubMed PMID: 36597481; PubMed Central PMCID: PMCPMC9800021.

79. Biering SB, Sarnik SA, Wang E, Zengel JR, Leist SR, Schafer A, et al. Genome-wide
bidirectional CRISPR screens identify mucins as host factors modulating SARS-CoV-2 infection. Nat
Genet. 2022;54(8):1078-89. Epub 20220725. doi: 10.1038/s41588-022-01131-x. PubMed PMID:
35879412; PubMed Central PMCID: PMCPMC9355872.

80. Baggen J, Persoons L, Vanstreels E, Jansen S, Van Looveren D, Boeckx B, et al. Genome-
wide CRISPR screening identifies TMEM106B as a proviral host factor for SARS-CoV-2. Nat Genet.
2021;53(4):435-44. Epub 20210308. doi: 10.1038/s41588-021-00805-2. PubMed PMID:
33686287.

81. van der Made CI, Simons A, Schuurs-Hoeijmakers J, van den Heuvel G, Mantere T, Kersten
S, et al. Presence of Genetic Variants Among Young Men With Severe COVID-19. JAMA.
2020;324(7):663-73. doi: 10.1001/jama.2020.13719. PubMed PMID: 32706371; PubMed Central
PMCID: PMCPMC7382021.

82. Shelton JF, Shastri AJ, Ye C, Weldon CH, Filshtein-Sonmez T, Coker D, et al. Trans-ancestry
analysis reveals genetic and nongenetic associations with COVID-19 susceptibility and severity.
Nat Genet. 2021;53(6):801-8. Epub 20210422. doi: 10.1038/s41588-021-00854-7. PubMed
PMID: 33888907.
83. Roberts GHL, Partha R, Rhead B, Knight SC, Park DS, Coignet MV, et al. Expanded COVID-
19 phenotype definitions reveal distinct patterns of genetic association and protective effects.
Nat Genet. 2022;54(4):374-81. Epub 20220411. doi: 10.1038/s41588-022-01042-x. PubMed
PMID: 35410379.
84. Pietzner M, Chua RL, Wheeler E, Jechow K, Willett JDS, Radbruch H, et al. ELF5 is a
potential respiratory epithelial cell-specific risk gene for severe COVID-19. Nat Commun.
2022;13(1):4484. Epub 20220815. doi: 10.1038/s41467-022-31999-6. PubMed PMID: 35970849;
PubMed Central PMCID: PMCPMC9378714.
85. Latini A, Agolini E, Novelli A, Borgiani P, Giannini R, Gravina P, et al. COVID-19 and Genetic
Variants of Protein Involved in the SARS-CoV-2 Entry into the Host Cells. Genes (Basel).
2020;11(9). Epub 20200827. doi: 10.3390/genes11091010. PubMed PMID: 32867305; PubMed
Central PMCID: PMCPMC7565048.
86. Horowitz JE, Kosmicki JA, Damask A, Sharma D, Roberts GHL, Justice AE, et al. Genome-
wide analysis provides genetic evidence that ACE2 influences COVID-19 risk and yields risk scores
associated with severe disease. Nat Genet. 2022;54(4):382-92. Epub 20220303. doi:
10.1038/s41588-021-01006-7. PubMed PMID: 35241825; PubMed Central PMCID:
PMCPMC9005345.
87. Garcia-Garcia A, Perez de Diego R, Flores C, Rinchai D, Sole-Violan J, Deya-Martinez A, et
al. Humans with inherited MyD88 and IRAK-4 deficiencies are predisposed to hypoxemic COVID-
19 pneumonia. J Exp Med. 2023;220(5). Epub 20230303. doi: 10.1084/jem.20220170. PubMed
PMID: 36880831; PubMed Central PMCID: PMCPMC9998661.
88. Fallerini C, Daga S, Mantovani S, Benetti E, Picchiotti N, Francisci D, et al. Association of
Toll-like receptor 7 variants with life-threatening COVID-19 disease in males: findings from a
nested case-control study. Elife. 2021;10. Epub 20210302. doi: 10.7554/eLife.67569. PubMed
PMID: 33650967; PubMed Central PMCID: PMCPMC7987337.
89. Severe Covid GG, Ellinghaus D, Degenhardt F, Bujanda L, Buti M, Albillos A, et al.
Genomewide Association Study of Severe Covid-19 with Respiratory Failure. N Engl J Med.
2020;383(16):1522-34. Epub 20200617. doi: 10.1056/NEJMoa2020283. PubMed PMID:
32558485; PubMed Central PMCID: PMCPMC7315890.
90. Degenhardt F, Ellinghaus D, Juzenas S, Lerga-Jaso J, Wendorff M, Maya-Miles D, et al.
Detailed stratified GWAS analysis for severe COVID-19 in four European populations. Hum Mol
Genet. 2022;31(23):3945-66. doi: 10.1093/hmg/ddac158. PubMed PMID: 35848942; PubMed
Central PMCID: PMCPMC9703941.
91. D'Antonio M, Nguyen JP, Arthur TD, Matsui H, Initiative C-HG, D'Antonio-Chronowska A,
et al. SARS-CoV-2 susceptibility and COVID-19 disease severity are associated with genetic
variants affecting gene expression in a variety of tissues. Cell Rep. 2021;37(7):110020. Epub
20211103. doi: 10.1016/j.celrep.2021.110020. PubMed PMID: 34762851; PubMed Central
PMCID: PMCPMC8563343.

92. Cruz R, Diz-de Almeida S, Lopez de Heredia M, Quintela I, Ceballos FC, Pita G, et al. Novel
genes and sex differences in COVID-19 severity. Hum Mol Genet. 2022;31(22):3789-806. doi:
10.1093/hmg/ddac132. PubMed PMID: 35708486; PubMed Central PMCID: PMCPMC9652109.

93. Initiative C-HG. A first update on mapping the human genetic architecture of COVID-19.
Nature. 2022;608(7921):E1-E10. Epub 20220803. doi: 10.1038/s41586-022-04826-7. PubMed
PMID: 35922517; PubMed Central PMCID: PMCPMC9352569.

94. Andolfo I, Russo R, Lasorsa VA, Cantalupo S, Rosato BE, Bonfiglio F, et al. Common variants
at 21q22.3 locus influence MX1 and TMPRSS2 gene expression and susceptibility to severe COVID-
19. iScience. 2021;24(4):102322. Epub 20210317. doi: 10.1016/j.isci.2021.102322. PubMed
PMID: 33748697; PubMed Central PMCID: PMCPMC7968217.

95. Zhou S, Butler-Laporte G, Nakanishi T, Morrison DR, Afilalo J, Afilalo M, et al. A
Neanderthal OAS1 isoform protects individuals of European ancestry against COVID-19
susceptibility and severity. Nat Med. 2021;27(4):659-67. Epub 20210225. doi: 10.1038/s41591-
021-01281-1. PubMed PMID: 33633408.

96. Huffman JE, Butler-Laporte G, Khan A, Pairo-Castineira E, Drivas TG, Peloso GM, et al.
Multi-ancestry fine mapping implicates OAS1 splicing in risk of severe COVID-19. Nat Genet.
2022;54(2):125-7. Epub 20220113. doi: 10.1038/s41588-021-00996-8. PubMed PMID: 35027740;
PubMed Central PMCID: PMCPMC8837537.

97. Initiative C-HG. Mapping the human genetic architecture of COVID-19. Nature.
2021;600(7889):472-7. Epub 20210708. doi: 10.1038/s41586-021-03767-x. PubMed PMID:
34237774; PubMed Central PMCID: PMCPMC8674144.

98. Pairo-Castineira E, Clohisey S, Klaric L, Bretherick AD, Rawlik K, Pasko D, et al. Genetic
mechanisms of critical illness in COVID-19. Nature. 2021;591(7848):92-8. Epub 20201211. doi:
10.1038/s41586-020-03065-y. PubMed PMID: 33307546.

99. Banday AR, Stanifer ML, Florez-Vargas O, Onabajo OO, Papenberg BW, Zahoor MA, et al.
Genetic regulation of OAS1 nonsense-mediated decay underlies association with COVID-19
hospitalization in patients of European and African ancestries. Nat Genet. 2022;54(8):1103-16.
Epub 20220714. doi: 10.1038/s41588-022-01113-z. PubMed PMID: 35835913; PubMed Central
PMCID: PMCPMC9355882.

100. Payman S-T, Hala A, James DRK, Audrey A, Reuben S, Zhen-Yuan L, et al. A SARS-CoV-2 –
host proximity interactome. bioRxiv. 2020:2020.09.03.282103. doi: 10.1101/2020.09.03.282103.

101. Gordon DE, Jang GM, Bouhaddou M, Xu J, Obernier K, White KM, et al. A SARS-CoV-2
protein interaction map reveals targets for drug repurposing. Nature. 2020;583(7816):459-68.
Epub 20200430. doi: 10.1038/s41586-020-2286-9. PubMed PMID: 32353859; PubMed Central
PMCID: PMCPMC7431030.

102. Estelle MNL, Yorgos S, Anastassia K, Jean-Pascal G, Payman Samavarchi T, Dae-Kyum K, et
al. Global BioID-based SARS-CoV-2 proteins proximal interactome unveils novel ties between viral
polypeptides and host factors involved in multiple COVID19-associated mechanisms. bioRxiv.
2020:2020.08.28.272955. doi: 10.1101/2020.08.28.272955.

103. St-Germain JR, Astori A, Samavarchi-Tehrani P, Abdouni H, Macwan V, Kim D-K, et al. A
SARS-CoV-2 BioID-based virus-host membrane protein interactome and virus peptide
compendium: new proteomics resources for COVID-19 research. bioRxiv.
2020:2020.08.28.269175. doi: 10.1101/2020.08.28.269175.

104. Liu X, Huuskonen S, Laitinen T, Redchuk T, Bogacheva M, Salokas K, et al. SARS-CoV-2-host proteome interactions for antiviral drug discovery. Mol Syst Biol. 2021;17(11):e10396. doi: 10.15252/msb.202110396. PubMed PMID: 34709727; PubMed Central PMCID: PMCPMC8552907.

105. May DG, Martin-Sancho L, Anschau V, Liu S, Chrisopulos RJ, Scott KL, et al. A BioID-Derived Proximity Interactome for SARS-CoV-2 Proteins. Viruses. 2022;14(3). Epub 20220315. doi: 10.3390/v14030611. PubMed PMID: 35337019; PubMed Central PMCID: PMCPMC8951556.

106. Zhou Y, Liu Y, Gupta S, Paramo MI, Hou Y, Mao C, et al. A comprehensive SARS-CoV-2-human protein-protein interactome reveals COVID-19 pathobiology and potential host therapeutic targets. Nat Biotechnol. 2023;41(1):128-39. Epub 20221010. doi: 10.1038/s41587-022-01474-0. PubMed PMID: 36217030; PubMed Central PMCID: PMCPMC9851973.

107. Stukalov A, Girault V, Grass V, Karayel O, Bergant V, Urban C, et al. Multilevel proteomics reveals host perturbations by SARS-CoV-2 and SARS-CoV. Nature. 2021;594(7862):246-52. Epub 20210412. doi: 10.1038/s41586-021-03493-4. PubMed PMID: 33845483.

108. Li J, Guo M, Tian X, Wang X, Yang X, Wu P, et al. Virus-Host Interactome and Proteomic Survey Reveal Potential Virulence Factors Influencing SARS-CoV-2 Pathogenesis. Med. 2021;2(1):99-112 e7. Epub 20200721. doi: 10.1016/j.medj.2020.07.002. PubMed PMID: 32838362; PubMed Central PMCID: PMCPMC7373048.

109. Davies JP, Almasy KM, McDonald EF, Plate L. Comparative Multiplexed Interactomics of SARS-CoV-2 and Homologous Coronavirus Nonstructural Proteins Identifies Unique and Shared Host-Cell Dependencies. ACS Infect Dis. 2020;6(12):3174-89. Epub 20201202. doi: 10.1021/acsinfecdis.0c00500. PubMed PMID: 33263384; PubMed Central PMCID: PMCPMC7724760.

110. Schmidt N, Ganskih S, Wei Y, Gabel A, Zielinski S, Keshishian H, et al. SND1 binds SARS-CoV-2 negative-sense RNA and promotes viral RNA synthesis through NSP9. Cell. 2023;186(22):4834-50 e23. Epub 20231003. doi: 10.1016/j.cell.2023.09.002. PubMed PMID: 37794589; PubMed Central PMCID: PMCPMC10617981.

111. Schmidt N, Lareau CA, Keshishian H, Ganskih S, Schneider C, Hennig T, et al. The SARS-CoV-2 RNA-protein interactome in infected human cells. Nat Microbiol. 2021;6(3):339-53. Epub 20201221. doi: 10.1038/s41564-020-00846-z. PubMed PMID: 33349665; PubMed Central PMCID: PMCPMC7906908.

112. Labeau A, Fery-Simonian L, Lefevre-Utile A, Pourcelot M, Bonnet-Madin L, Soumelis V, et al. Characterization and functional interrogation of the SARS-CoV-2 RNA interactome. Cell Rep. 2022;39(4):110744. doi: 10.1016/j.celrep.2022.110744. PubMed PMID: 35477000; PubMed Central PMCID: PMCPMC9040432.

113. Kamel W, Noerenberg M, Cerikan B, Chen H, Jarvelin AI, Kammoun M, et al. Global analysis of protein-RNA interactions in SARS-CoV-2-infected cells reveals key regulators of infection. Mol Cell. 2021;81(13):2851-67 e7. Epub 20210524. doi: 10.1016/j.molcel.2021.05.023. PubMed PMID: 34118193; PubMed Central PMCID: PMCPMC8142890.

114. Flynn RA, Belk JA, Qi Y, Yasumoto Y, Wei J, Alfajaro MM, et al. Discovery and functional interrogation of SARS-CoV-2 RNA-host protein interactions. Cell. 2021;184(9):2394-411 e16. Epub 20210311. doi: 10.1016/j.cell.2021.03.012. PubMed PMID: 33743211; PubMed Central PMCID: PMCPMC7951565.

115. Lee S, Lee YS, Choi Y, Son A, Park Y, Lee KM, et al. The SARS-CoV-2 RNA interactome. Mol
Cell. 2021;81(13):2838-50 e6. Epub 20210427. doi: 10.1016/j.molcel.2021.04.022. PubMed
PMID: 33989516; PubMed Central PMCID: PMCPMC8075806.
116. Bouhaddou M, Memon D, Meyer B, White KM, Rezelj VV, Correa Marrero M, et al. The
Global Phosphorylation Landscape of SARS-CoV-2 Infection. Cell. 2020;182(3):685-712 e19. Epub
20200628. doi: 10.1016/j.cell.2020.06.034. PubMed PMID: 32645325; PubMed Central PMCID:
PMCPMC7321036.
117. Sanchez-Magraner L, Posada IM, Andraka N, Contreras FX, Viguera AR, Guerin DM, et al.
The C-terminal transmembrane domain of human phospholipid scramblase 1 is essential for the
protein flip-flop activity and Ca(2)(+)-binding. J Membr Biol. 2014;247(2):155-65. Epub 20131217.
doi: 10.1007/s00232-013-9619-7. PubMed PMID: 24343571.
118. Abdulla S, Aevermann B, Assis P, Badajoz S, Bell SM, Bezzi E, et al. CZ CELL×GENE Discover:
A single-cell data platform for scalable exploration, analysis and modeling of aggregated data.
bioRxiv. 2023:2023.10.30.563174. doi: 10.1101/2023.10.30.563174.
119. Ricardo-Lax I, Luna JM, Thao TTN, Le Pen J, Yu Y, Hoffmann HH, et al. Replication and single-
cycle delivery of SARS-CoV-2 replicons. Science. 2021;374(6571):1099-106. Epub 20211014. doi:
10.1126/science.abj8430. PubMed PMID: 34648371; PubMed Central PMCID: PMCPMC9007107.
120. Schmidt F, Weisblum Y, Muecksch F, Hoffmann HH, Michailidis E, Lorenzi JCC, et al.
Measuring SARS-CoV-2 neutralizing antibody activity using pseudotyped and chimeric viruses. J
Exp Med. 2020;217(11). doi: 10.1084/jem.20201181. PubMed PMID: 32692348; PubMed Central
PMCID: PMCPMC7372514.
121. Pfaender S, Mar KB, Michailidis E, Kratzel A, Boys IN, V'Kovski P, et al. LY6E impairs
coronavirus fusion and confers immune control of viral disease. Nat Microbiol. 2020;5(11):1330-
9. Epub 20200723. doi: 10.1038/s41564-020-0769-y. PubMed PMID: 32704094; PubMed Central
PMCID: PMCPMC7916999.
122. Winstone H, Lista MJ, Reid AC, Bouton C, Pickering S, Galao RP, et al. The Polybasic
Cleavage Site in SARS-CoV-2 Spike Modulates Viral Sensitivity to Type I Interferon and IFITM2. J
Virol. 2021;95(9). Epub 20210412. doi: 10.1128/JVI.02422-20. PubMed PMID: 33563656;
PubMed Central PMCID: PMCPMC8104117.
123. Thorne LG, Bouhaddou M, Reuschl AK, Zuliani-Alvarez L, Polacco B, Pelin A, et al. Evolution
of enhanced innate immune evasion by SARS-CoV-2. Nature. 2022;602(7897):487-95. Epub
20211223. doi: 10.1038/s41586-021-04352-y. PubMed PMID: 34942634; PubMed Central
PMCID: PMCPMC8850198.
124. Markov PV, Ghafari M, Beer M, Lythgoe K, Simmonds P, Stilianakis NI, et al. The evolution
of SARS-CoV-2. Nature Reviews Microbiology. 2023;21(6):361-79. doi: 10.1038/s41579-023-
00878-2.
125. Carabelli AM, Peacock TP, Thorne LG, Harvey WT, Hughes J, de Silva TI, et al. SARS-CoV-2
variant biology: immune escape, transmission and fitness. Nature Reviews Microbiology.
2023;21(3):162-77. doi: 10.1038/s41579-022-00841-7.
126. Huang P, Liao R, Chen X, Wu X, Li X, Wang Y, et al. Nuclear translocation of PLSCR1 activates
STAT1 signaling in basal-like breast cancer. Theranostics. 2020;10(10):4644-58. Epub 20200325.
doi: 10.7150/thno.43150. PubMed PMID: 32292520; PubMed Central PMCID: PMCPMC7150476.

127.   Nanjundan M, Sun J, Zhao J, Zhou Q, Sims PJ, Wiedmer T. Plasma membrane phospholipid
scramblase 1 promotes EGF-dependent activation of c-Src through the epidermal growth factor
receptor. J Biol Chem. 2003;278(39):37413-8. Epub 20030718. doi: 10.1074/jbc.M306182200.
PubMed PMID: 12871937.

128.   Ben-Efraim I, Zhou Q, Wiedmer T, Gerace L, Sims PJ. Phospholipid scramblase 1 is imported
into the nucleus by a receptor-mediated pathway and interacts with DNA. Biochemistry.
2004;43(12):3518-26. doi: 10.1021/bi0356911. PubMed PMID: 15035622.

129.   Chen MH, Ben-Efraim I, Mitrousis G, Walker-Kopp N, Sims PJ, Cingolani G. Phospholipid
scramblase 1 contains a nonclassical nuclear localization signal with unique binding site in
importin   alpha.   J   Biol   Chem.   2005;280(11):10599-606.   Epub   20041217.   doi:
10.1074/jbc.M413194200. PubMed PMID: 15611084.

130.   Wiedmer T, Zhao J, Nanjundan M, Sims PJ. Palmitoylation of phospholipid scramblase 1
controls its distribution between nucleus and plasma membrane. Biochemistry. 2003;42(5):1227-
33. doi: 10.1021/bi026679w. PubMed PMID: 12564925.

131.   Zhao J, Zhou Q, Wiedmer T, Sims PJ. Palmitoylation of phospholipid scramblase is required
for   normal   function   in   promoting   Ca2+-activated   transbilayer   movement   of   membrane
phospholipids.   Biochemistry.   1998;37(18):6361-6.   doi: 10.1021/bi980218m.   PubMed   PMID:
9572851.

132.   Chen S, Francioli LC, Goodrich JK, Collins RL, Kanai M, Wang Q, et al. A genomic mutational
constraint map using variation in 76,156 human genomes. Nature. 2024;625(7993):92-100. doi:
10.1038/s41586-023-06045-0.

133.   Zhang P, Bigio B, Rapaport F, Zhang SY, Casanova JL, Abel L, et al. PopViz: a webserver for
visualizing minor allele frequencies and damage prediction scores of human genetic variations.
Bioinformatics.   2018;34(24):4307-9.   doi: 10.1093/bioinformatics/bty536.   PubMed   PMID:
30535305; PubMed Central PMCID: PMCPMC6289133.

134.   Dittmann M, Hoffmann HH, Scull MA, Gilmore RH, Bell KL, Ciancanelli M, et al. A serpin
shapes   the   extracellular   environment   to   prevent   influenza   A   virus   maturation.   Cell.
2015;160(4):631-43. doi: 10.1016/j.cell.2015.01.040. PubMed PMID: 25679759; PubMed Central
PMCID: PMCPMC4328142.

135.   Burke JM, St Clair LA, Perera R, Parker R. SARS-CoV-2 infection triggers widespread host
mRNA decay leading to an mRNA export block. RNA. 2021;27(11):1318-29. Epub 20210727. doi:
10.1261/rna.078923.121. PubMed PMID: 34315815; PubMed Central PMCID: PMCPMC8522697.

136.   Zhang K, Miorin L, Makio T, Dehghan I, Gao S, Xie Y, et al. Nsp1 protein of SARS-CoV-2
disrupts the mRNA export machinery to inhibit host gene expression. Sci Adv. 2021;7(6). Epub
20210205.   doi: 10.1126/sciadv.abe7386.   PubMed   PMID: 33547084;   PubMed   Central   PMCID:
PMCPMC7864571.

137.   Kimura I, Konno Y, Uriu K, Hopfensperger K, Sauter D, Nakagawa S, et al. Sarbecovirus ORF6
proteins hamper induction of interferon signaling. Cell Rep. 2021;34(13):108916. Epub 20210312.
doi: 10.1016/j.celrep.2021.108916.   PubMed   PMID: 33765414;   PubMed   Central   PMCID:
PMCPMC7953434.

138.   Setaro AC, Gaglia MM. All hands on deck: SARS-CoV-2 proteins that block early anti-viral
interferon   responses.   Curr   Res   Virol   Sci.   2021;2:100015.   Epub   20211112.   doi:

10.1016/j.crviro.2021.100015. PubMed PMID: 34786565; PubMed Central PMCID: PMCPMC8588586.

139.   Ogunjimi B, Zhang SY, Sorensen KB, Skipper KA, Carter-Timofte M, Kerner G, et al. Inborn errors in RNA polymerase III underlie severe varicella zoster virus infections. J Clin Invest. 2017;127(9):3543-56. Epub 20170807. doi: 10.1172/JCI92280. PubMed PMID: 28783042; PubMed Central PMCID: PMCPMC5669568.

140.   Zimmer MM, Kibe A, Rand U, Pekarek L, Ye L, Buck S, et al. The short isoform of the host antiviral protein ZAP acts as an inhibitor of SARS-CoV-2 programmed ribosomal frameshifting. Nat Commun. 2021;12(1):7193. Epub 20211210. doi: 10.1038/s41467-021-27431-0. PubMed PMID: 34893599; PubMed Central PMCID: PMCPMC8664833.

141.   Klein S, Golani G, Lolicato F, Lahr C, Beyer D, Herrmann A, et al. IFITM3 blocks influenza virus entry by sorting lipids and stabilizing hemifusion. Cell Host Microbe. 2023. Epub 20230329. doi: 10.1016/j.chom.2023.03.005. PubMed PMID: 37003257.

142.   Guo X, Steinkuhler J, Marin M, Li X, Lu W, Dimova R, et al. Interferon-Induced Transmembrane Protein 3 Blocks Fusion of Diverse Enveloped Viruses by Altering Mechanical Properties of Cell Membranes. ACS Nano. 2021;15(5):8155-70. Epub 20210303. doi: 10.1021/acsnano.0c10567. PubMed PMID: 33656312; PubMed Central PMCID: PMCPMC8159881.

143.   Suddala KC, Lee CC, Meraner P, Marin M, Markosyan RM, Desai TM, et al. Interferon-induced transmembrane protein 3 blocks fusion of sensitive but not resistant viruses by partitioning into virus-carrying endosomes. PLoS Pathog. 2019;15(1):e1007532. Epub 20190114. doi: 10.1371/journal.ppat.1007532. PubMed PMID: 30640957; PubMed Central PMCID: PMCPMC6347298.

144.   Desai TM, Marin M, Chin CR, Savidis G, Brass AL, Melikyan GB. IFITM3 restricts influenza A virus entry by blocking the formation of fusion pores following virus-endosome hemifusion. PLoS Pathog. 2014;10(4):e1004048. Epub 20140403. doi: 10.1371/journal.ppat.1004048. PubMed PMID: 24699674; PubMed Central PMCID: PMCPMC3974867.

145.   Fadeel B, Gleiss B, Hogstrand K, Chandra J, Wiedmer T, Sims PJ, et al. Phosphatidylserine exposure during apoptosis is a cell-type-specific event and does not correlate with plasma membrane phospholipid scramblase expression. Biochem Biophys Res Commun. 1999;266(2):504-11. doi: 10.1006/bbrc.1999.1820. PubMed PMID: 10600532.

146.   Zhou Q, Zhao J, Al-Zoghaibi F, Zhou A, Wiedmer T, Silverman RH, et al. Transcriptional control of the human plasma membrane phospholipid scramblase 1 gene is mediated by interferon-alpha. Blood. 2000;95(8):2593-9. PubMed PMID: 10753839.

147.   Zhou Q, Zhao J, Wiedmer T, Sims PJ. Normal hemostasis but defective hematopoietic response to growth factors in mice deficient in phospholipid scramblase 1. Blood. 2002;99(11):4030-8. doi: 10.1182/blood-2001-12-0271. PubMed PMID: 12010804.

148.   Majdoul S, Compton AA. Lessons in self-defence: inhibition of virus entry by intrinsic immunity. Nat Rev Immunol. 2022;22(6):339-52. Epub 20211013. doi: 10.1038/s41577-021-00626-8. PubMed PMID: 34646033; PubMed Central PMCID: PMCPMC8511856.

149.   Kusano S, Eizuru Y. Human phospholipid scramblase 1 interacts with and regulates transactivation of HTLV-1 Tax. Virology. 2012;432(2):343-52. Epub 20120711. doi: 10.1016/j.virol.2012.06.019. PubMed PMID: 22789739.

150.   Kusano S, Eizuru Y. Interaction of the phospholipid scramblase 1 with HIV-1 Tat results in
the repression of Tat-dependent transcription. Biochem Biophys Res Commun. 2013;433(4):438-
44. Epub 20130315. doi: 10.1016/j.bbrc.2013.02.098. PubMed PMID: 23501106.

151.   Sadanari H, Takemoto M, Ishida T, Otagiri H, Daikoku T, Murayama T, et al. The Interferon-
Inducible Human PLSCR1 Protein Is a Restriction Factor of Human Cytomegalovirus. Microbiol
Spectr. 2022;10(1):e0134221. Epub 2022/02/10. doi: 10.1128/spectrum.01342-21. PubMed
PMID: 35138119; PubMed Central PMCID: PMCPMC8826943.

152.   Metz P, Dazert E, Ruggieri A, Mazur J, Kaderali L, Kaul A, et al. Identification of type I and
type II interferon-induced effectors controlling hepatitis C virus replication. Hepatology.
2012;56(6):2082-93. Epub 20121014. doi: 10.1002/hep.25908. PubMed PMID: 22711689.

153.   Yang J, Zhu X, Liu J, Ding X, Han M, Hu W, et al. Inhibition of Hepatitis B virus replication
by phospholipid scramblase 1 in vitro and in vivo. Antiviral Res. 2012;94(1):9-17. Epub 20120209.
doi: 10.1016/j.antiviral.2012.01.010. PubMed PMID: 22342889.

154.   Kusano S, Ikeda M. Interaction of phospholipid scramblase 1 with the Epstein-Barr virus
protein BZLF1 represses BZLF1-mediated lytic gene transcription. J Biol Chem.
2019;294(41):15104-16. Epub 2019/08/23. doi: 10.1074/jbc.RA119.008193. PubMed PMID:
31434743; PubMed Central PMCID: PMCPMC6791327.

155.   Luo W, Zhang J, Liang L, Wang G, Li Q, Zhu P, et al. Phospholipid scramblase 1 interacts
with influenza A virus NP, impairing its nuclear import and thereby suppressing virus replication.
PLoS Pathog. 2018;14(1):e1006851. Epub 20180119. doi: 10.1371/journal.ppat.1006851.
PubMed PMID: 29352288; PubMed Central PMCID: PMCPMC5792031.

156.   Dal Col J, Lamberti MJ, Nigro A, Casolaro V, Fratta E, Steffan A, et al. Phospholipid
scramblase 1: a protein with multiple functions via multiple molecular interactors. Cell Commun
Signal. 2022;20(1):78. Epub 2022/06/02. doi: 10.1186/s12964-022-00895-3. PubMed PMID:
35650588.

157.   Jonathan RS-G, Audrey A, Payman S-T, Hala A, Vinitha M, Dae-Kyum K, et al. A SARS-CoV-
2 BioID-based virus-host membrane protein interactome and virus peptide compendium: new
proteomics resources for COVID-19 research. bioRxiv. 2020:2020.08.28.269175. doi:
10.1101/2020.08.28.269175.

158.   Guo K, Barrett BS, Morrison JH, Mickens KL, Vladar EK, Hasenkrug KJ, et al. Interferon
resistance of emerging SARS-CoV-2 variants. Proc Natl Acad Sci U S A.
2022;119(32):e2203760119. Epub 20220722. doi: 10.1073/pnas.2203760119. PubMed PMID:
35867811; PubMed Central PMCID: PMCPMC9371743.

159.   Shalamova L, Felgenhauer U, Wilhelm J, Schaubmar AR, Buttner K, Schoen A, et al.
Omicron variant of SARS-CoV-2 exhibits an increased resilience to the antiviral type I interferon
response. PNAS Nexus. 2022;1(2):pgac067. Epub 20220523. doi: 10.1093/pnasnexus/pgac067.
PubMed PMID: 36713328; PubMed Central PMCID: PMCPMC9802332.

160.   Shi G, Li T, Lai KK, Johnson RF, Yewdell JW, Compton AA. Omicron Spike confers enhanced
infectivity and interferon resistance to SARS-CoV-2 in human nasal tissue. Nat Commun.
2024;15(1):889. Epub 20240130. doi: 10.1038/s41467-024-45075-8. PubMed PMID: 38291024;
PubMed Central PMCID: PMCPMC10828397.

161.   Doyle T, Moncorge O, Bonaventure B, Pollpeter D, Lussignol M, Tauziet M, et al. The
interferon-inducible isoform of NCOA7 inhibits endosome-mediated viral entry. Nat Microbiol.

2018;3(12):1369-76. Epub 20181126. doi: 10.1038/s41564-018-0273-9. PubMed PMID:
30478388; PubMed Central PMCID: PMCPMC6329445.

162.   Khan H, Winstone H, Jimenez-Guardeno JM, Graham C, Doores KJ, Goujon C, et al.
TMPRSS2 promotes SARS-CoV-2 evasion from NCOA7-mediated restriction. PLoS Pathog.
2021;17(11):e1009820. Epub 20211122. doi: 10.1371/journal.ppat.1009820. PubMed PMID:
34807954; PubMed Central PMCID: PMCPMC8648102.

163.   Willett BJ, Grove J, MacLean OA, Wilkie C, De Lorenzo G, Furnon W, et al. SARS-CoV-2
Omicron is an immune escape variant with an altered cell entry pathway. Nat Microbiol.
2022;7(8):1161-79. Epub 20220707. doi: 10.1038/s41564-022-01143-7. PubMed PMID:
35798890; PubMed Central PMCID: PMCPMC9352574.

164.   Peacock TP, Brown JC, Zhou J, Thakur N, Sukhova K, Newman J, et al. The altered entry
pathway and antigenic distance of the SARS-CoV-2 Omicron variant map to separate domains of
spike protein. bioRxiv. 2022:2021.12.31.474653. doi: 10.1101/2021.12.31.474653.

165.   Shuai H, Chan JF, Hu B, Chai Y, Yuen TT, Yin F, et al. Attenuated replication and pathogenicity
of SARS-CoV-2 B.1.1.529 Omicron. Nature. 2022;603(7902):693-9. Epub 20220121. doi:
10.1038/s41586-022-04442-5. PubMed PMID: 35062016.

166.   Hui KPY, Ho JCW, Cheung MC, Ng KC, Ching RHH, Lai KL, et al. SARS-CoV-2 Omicron variant
replication in human bronchus and lung ex vivo. Nature. 2022;603(7902):715-20. Epub 20220201.
doi: 10.1038/s41586-022-04479-6. PubMed PMID: 35104836.

167.   Zhao H, Lu L, Peng Z, Chen LL, Meng X, Zhang C, et al. SARS-CoV-2 Omicron variant shows
less efficient replication and fusion activity when compared with Delta variant in TMPRSS2-
expressed   cells.   Emerg   Microbes   Infect.   2022;11(1):277-83.   doi:
10.1080/22221751.2021.2023329. PubMed PMID: 34951565; PubMed Central PMCID:
PMCPMC8774049.

168.   Meng B, Abdullahi A, Ferreira I, Goonawardane N, Saito A, Kimura I, et al. Altered TMPRSS2
usage by SARS-CoV-2 Omicron impacts infectivity and fusogenicity. Nature. 2022;603(7902):706-
14. Epub 20220201. doi: 10.1038/s41586-022-04474-x. PubMed PMID: 35104837; PubMed
Central PMCID: PMCPMC8942856.

169.   Fan Y, Li X, Zhang L, Wan S, Zhang L, Zhou F. SARS-CoV-2 Omicron variant: recent progress
and future perspectives. Signal Transduct Target Ther. 2022;7(1):141. Epub 20220428. doi:
10.1038/s41392-022-00997-x. PubMed PMID: 35484110; PubMed Central PMCID:
PMCPMC9047469.

170.   Sette A, Crotty S. Adaptive immunity to SARS-CoV-2 and COVID-19. Cell. 2021;184(4):861-
80. Epub 20210112. doi: 10.1016/j.cell.2021.01.007. PubMed PMID: 33497610; PubMed Central
PMCID: PMCPMC7803150.

171.   Mar KB, Van Dyke MC, Lopez AH, Eitson JL, Fan W, Hanners NW, et al. LY6E protects mice
from pathogenic effects of murine coronavirus and SARS-CoV-2. bioRxiv. 2023. Epub 20230126.
doi: 10.1101/2023.01.25.525551. PubMed PMID: 36747632; PubMed Central PMCID:
PMCPMC9900800.

172.   Consortium GT. The Genotype-Tissue Expression (GTEx) project. Nat Genet.
2013;45(6):580-5. doi: 10.1038/ng.2653. PubMed PMID: 23715323; PubMed Central PMCID:
PMCPMC4010069.

173.   Blight KJ, McKeating JA, Rice CM. Highly permissive cell lines for subgenomic and genomic
hepatitis C virus RNA replication. J Virol. 2002;76(24):13001-14. doi: 10.1128/jvi.76.24.13001-
13014.2002. PubMed PMID: 12438626; PubMed Central PMCID: PMCPMC136668.
174.   Zhang L, Bukreyev A, Thompson CI, Watson B, Peeples ME, Collins PL, et al. Infection of
ciliated cells by human parainfluenza virus type 3 in an in vitro model of human airway epithelium.
J Virol. 2005;79(2):1113-24. doi: 10.1128/JVI.79.2.1113-1124.2005. PubMed PMID: 15613339;
PubMed Central PMCID: PMCPMC538579.
175.   Hoffmann HH, Schneider WM, Rozen-Gagnon K, Miles LA, Schuster F, Razooky B, et al.
TMEM41B Is a Pan-flavivirus Host Factor. Cell. 2021;184(1):133-48 e20. Epub 20201209. doi:
10.1016/j.cell.2020.12.005.   PubMed   PMID: 33338421;   PubMed   Central   PMCID:
PMCPMC7954666.
176.   Benboudjema L, Mulvey M, Gao Y, Pimplikar SW, Mohr I. Association of the herpes simplex
virus type 1 Us11 gene product with the cellular kinesin light-chain-related protein PAT1 results
in   the   redistribution   of   both   polypeptides.   J   Virol.   2003;77(17):9192-203.   doi:
10.1128/jvi.77.17.9192-9203.2003.   PubMed   PMID: 12915535;   PubMed   Central   PMCID:
PMCPMC187382.
177.   Lee D, Le Pen J, Yatim A, Dong B, Aquino Y, Ogishi M, et al. Inborn errors of OAS-RNase L
in   SARS-CoV-2-related   multisystem   inflammatory   syndrome   in   children.   Science.
2023;379(6632):eabo3627. Epub 20230210. doi: 10.1126/science.abo3627. PubMed PMID:
36538032.
178.   Mendoza EJ, Manguiat K, Wood H, Drebot M. Two Detailed Plaque Assay Protocols for the
Quantification of Infectious SARS-CoV-2. Curr Protoc Microbiol. 2020;57(1):ecpmc105. doi:
10.1002/cpmc.105. PubMed PMID: 32475066; PubMed Central PMCID: PMCPMC7300432.
179.   Polo JM, Davis NL, Rice CM, Huang HV, Johnston RE. Molecular analysis of Sindbis virus
pathogenesis in neonatal mice by using virus recombinants constructed in vitro. J Virol.
1988;62(6):2124-33. doi: 10.1128/JVI.62.6.2124-2133.1988. PubMed PMID: 2835514; PubMed
Central PMCID: PMCPMC253309.
180.   Bick MJ, Carroll JW, Gao G, Goff SP, Rice CM, MacDonald MR. Expression of the zinc-finger
antiviral   protein   inhibits   alphavirus   replication.   J   Virol.   2003;77(21):11555-62.   doi:
10.1128/jvi.77.21.11555-11562.2003.   PubMed   PMID: 14557641;   PubMed   Central   PMCID:
PMCPMC229374.
181.   Atasheva S, Kim DY, Akhrymuk M, Morgan DG, Frolova EI, Frolov I. Pseudoinfectious
Venezuelan   equine   encephalitis   virus: a   new   means   of   alphavirus   attenuation.   J   Virol.
2013;87(4):2023-35. Epub 20121205. doi: 10.1128/JVI.02881-12. PubMed PMID: 23221545;
PubMed Central PMCID: PMCPMC3571482.
182.   Dalton KP, Rose JK. Vesicular stomatitis virus glycoprotein containing the entire green
fluorescent protein on its cytoplasmic domain is incorporated efficiently into virus particles.
Virology. 2001;279(2):414-21. doi: 10.1006/viro.2000.0736. PubMed PMID: 11162797.
183.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast universal
RNA-seq   aligner.   Bioinformatics.   2013;29(1):15-21.   Epub   20121025.   doi:
10.1093/bioinformatics/bts635.   PubMed   PMID: 23104886;   PubMed   Central   PMCID:
PMCPMC3530905.

184.	Liao Y, Smyth GK, Shi W. The R package Rsubread is easier, faster, cheaper and better for
alignment and quantification of RNA sequencing reads. Nucleic Acids Res. 2019;47(8):e47. doi:
10.1093/nar/gkz114. PubMed PMID: 30783653; PubMed Central PMCID: PMCPMC6486549.

185.	Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
seq data with DESeq2. Genome Biol. 2014;15(12):550. doi: 10.1186/s13059-014-0550-8. PubMed
PMID: 25516281; PubMed Central PMCID: PMCPMC4302049.

186.	Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A. Fast gene set
enrichment analysis. bioRxiv. 2021:060012. doi: 10.1101/060012.

187.	Liberzon A, Subramanian A, Pinchback R, Thorvaldsdottir H, Tamayo P, Mesirov JP.
Molecular signatures database (MSigDB) 3.0. Bioinformatics. 2011;27(12):1739-40. Epub
20110505. doi: 10.1093/bioinformatics/btr260. PubMed PMID: 21546393; PubMed Central
PMCID: PMCPMC3106198.

188.	Fabregat A, Jupe S, Matthews L, Sidiropoulos K, Gillespie M, Garapati P, et al. The
Reactome Pathway Knowledgebase. Nucleic Acids Res. 2018;46(D1):D649-D55. doi:
10.1093/nar/gkx1132. PubMed PMID: 29145629; PubMed Central PMCID: PMCPMC5753187.

189.	Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids Res.
2000;28(1):27-30. doi: 10.1093/nar/28.1.27. PubMed PMID: 10592173; PubMed Central PMCID:
PMCPMC102409.

190.	Agrawal A, Balci H, Hanspers K, Coort SL, Martens M, Slenter DN, et al. WikiPathways 2024:
next generation pathway database. Nucleic Acids Res. 2024;52(D1):D679-D89. doi:
10.1093/nar/gkad960. PubMed PMID: 37941138; PubMed Central PMCID: PMCPMC10767877.

191.	Schaefer CF, Anthony K, Krupa S, Buchoff J, Day M, Hannay T, et al. PID: the Pathway
Interaction Database. Nucleic Acids Res. 2009;37(Database issue):D674-9. Epub 20081002. doi:
10.1093/nar/gkn653. PubMed PMID: 18832364; PubMed Central PMCID: PMCPMC2686461.

192.	Nishimura D. BioCarta. Biotech Software & Internet Report: The Computer Software
Journal for Scient. 2001;2(3):117-20.

193.	Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method.
Nat Protoc. 2008;3(6):1101-8. doi: 10.1038/nprot.2008.73. PubMed PMID: 18546601.

194.	Witte L, Baharani VA, Schmidt F, Wang Z, Cho A, Raspe R, et al. Epistasis lowers the genetic
barrier to SARS-CoV-2 neutralizing antibody escape. Nat Commun. 2023;14(1):302. Epub
20230119. doi: 10.1038/s41467-023-35927-0. PubMed PMID: 36653360; PubMed Central
PMCID: PMCPMC9849103.

195.	Wang Z, Muecksch F, Raspe R, Johannsen F, Turroja M, Canis M, et al. Memory B cell
development elicited by mRNA booster vaccinations in the elderly. J Exp Med. 2023;220(9). Epub
20230627. doi: 10.1084/jem.20230668. PubMed PMID: 37368240; PubMed Central PMCID:
PMCPMC10300432.

196.	Dittmann M, Hoffmann HH, Scull MA, Gilmore RH, Bell KL, Ciancanelli M, et al. A serpin
shapes the extracellular environment to prevent influenza A virus maturation. Cell.
2015;160(4):631-43. doi: 10.1016/j.cell.2015.01.040. PubMed PMID: 25679759; PubMed Central
PMCID: PMC4328142.

197.	UniProt C. UniProt: the Universal Protein Knowledgebase in 2023. Nucleic Acids Res.
2023;51(D1):D523-D31. doi: 10.1093/nar/gkac1052. PubMed PMID: 36408920; PubMed Central
PMCID: PMCPMC9825514.

198. Bruchez A, Sha K, Johnson J, Chen L, Stefani C, McConnell H, et al. MHC class II
transactivator CIITA induces cell resistance to Ebola virus and SARS-like coronaviruses. Science.
2020;370(6513):241-7. Epub 20200827. doi: 10.1126/science.abb3753. PubMed PMID:
32855215; PubMed Central PMCID: PMCPMC7665841.

199. Lista MJ, Winstone H, Wilson HD, Dyer A, Pickering S, Galao RP, et al. The P681H Mutation
in the Spike Glycoprotein of the Alpha Variant of SARS-CoV-2 Escapes IFITM Restriction and Is
Necessary for Type I Interferon Resistance. J Virol. 2022;96(23):e0125022. Epub 20221109. doi:
10.1128/jvi.01250-22. PubMed PMID: 36350154; PubMed Central PMCID: PMCPMC9749455.

200. Shi G, Kenney AD, Kudryashova E, Zani A, Zhang L, Lai KK, et al. Opposing activities of IFITM
proteins in SARS-CoV-2 infection. EMBO J. 2021;40(3):e106501. Epub 20201221. doi:
10.15252/embj.2020106501. PubMed PMID: 33270927; PubMed Central PMCID:
PMCPMC7744865.

201. Prelli Bozzo C, Nchioua R, Volcic M, Koepke L, Kruger J, Schutz D, et al. IFITM proteins
promote SARS-CoV-2 infection and are targets for virus inhibition in vitro. Nat Commun.
2021;12(1):4584. Epub 20210728. doi: 10.1038/s41467-021-24817-y. PubMed PMID: 34321474;
PubMed Central PMCID: PMCPMC8319209.

202. Peacock TP, Goldhill DH, Zhou J, Baillon L, Frise R, Swann OC, et al. The furin cleavage site
in the SARS-CoV-2 spike protein is required for transmission in ferrets. Nat Microbiol.
2021;6(7):899-909. Epub 20210427. doi: 10.1038/s41564-021-00908-w. PubMed PMID:
33907312.

203. Nchioua R, Schundner A, Kmiec D, Prelli Bozzo C, Zech F, Koepke L, et al. SARS-CoV-2
Variants of Concern Hijack IFITM2 for Efficient Replication in Human Lung Cells. J Virol.
2022;96(11):e0059422. Epub 20220511. doi: 10.1128/jvi.00594-22. PubMed PMID: 35543509;
PubMed Central PMCID: PMCPMC9175628.

204. Xu F, Wang G, Zhao F, Huang Y, Fan Z, Mei S, et al. IFITM3 Inhibits SARS-CoV-2 Infection
and Is Associated with COVID-19 Susceptibility. Viruses. 2022;14(11). Epub 20221118. doi:
10.3390/v14112553. PubMed PMID: 36423162; PubMed Central PMCID: PMCPMC9692367.

205. Shi G, Chiramel AI, Li T, Lai KK, Kenney AD, Zani A, et al. Rapalogs downmodulate intrinsic
immunity and promote cell entry of SARS-CoV-2. J Clin Invest. 2022;132(24). Epub 20221215. doi:
10.1172/JCI160766. PubMed PMID: 36264642; PubMed Central PMCID: PMCPMC9753997.

206. Kenney AD, Zani A, Kawahara J, Eddy AC, Wang XL, Mahesh KC, et al. Interferon-induced
transmembrane protein 3 (IFITM3) limits lethality of SARS-CoV-2 in mice. EMBO Rep.
2023;24(4):e56660. Epub 20230307. doi: 10.15252/embr.202256660. PubMed PMID: 36880581;
PubMed Central PMCID: PMCPMC10074051.

207. Mathieson T, Franken H, Kosinski J, Kurzawa N, Zinn N, Sweetman G, et al. Systematic
analysis of protein turnover in primary cells. Nat Commun. 2018;9(1):689. Epub 2018/02/17. doi:
10.1038/s41467-018-03106-1. PubMed PMID: 29449567; PubMed Central PMCID:
PMCPMC5814408.

208. Szklarczyk D, Kirsch R, Koutrouli M, Nastou K, Mehryary F, Hachilif R, et al. The STRING
database in 2023: protein-protein association networks and functional enrichment analyses for
any sequenced genome of interest. Nucleic Acids Res. 2023;51(D1):D638-D46. doi:
10.1093/nar/gkac1000. PubMed PMID: 36370105; PubMed Central PMCID: PMCPMC9825434.
2406

Figure 1

Figure 2

Figure 3

SARS-CoV-2 functional genetic screens (26 publications)

COVID-19 human genetics studies (24 publications)

2,603 genes

113 genes

28 genes

PROVIRAL

Virus entry
ACE2*   TMPRSS2

Cell adhesion and immune signaling
ICAM3   PCDH7

Intracellular trafficking and signaling
RAB2A

Guanyl-nucleotide exchange
ARHGEF38

IFN signaling
IRF1

Ion channels and membrane potential
KCNC3

Transcriptional regulation
FOXP4   ZBTB32

Cell metabolism and redox homeostasis
CAT*   FDX2*

ANTIVIRAL

Protective mucous barrier
MUC1*

Cell adhesion and immune signaling
ICAM1*

TLR7 signaling
TLR7
MYD88   UNC93B1

IFN signaling
IFNAR1*   IFNAR2*
JAK1*
STAT2*

IFN-stimulated genes
OAS1   OAS2   OAS3   PLSCR1*

Transcriptional regulation
KAT7*   ZBTB11*

RNA splicing
CLK2

* Hit in arrayed CRISPR KO screen.

Figure 4

Figure 5

Figure 6

Figure 7

Supplementary figure 1

A

B

Supplementary figure 2

Supplementary figure 3

Supplementary figure 4

A

B

C

Supplementary figure 5

A

B

C

D

	Cell Count	PLSCR1
Lung	**2.7m**	
respiratory goblet cell	7.2k	
nasal mucosa goblet cell	3.9k	
bronchial goblet cell	557	
bronchial epithelial cell	1.1k	
epithelial cell of alveolus of lung	261.3k	
lung ciliated cell	8.7k	
Nose	**271.1k**	
respiratory goblet cell	100.3k	
basal cell of epithelium of trachea	28.3k	
nasal mucosa goblet cell	34.4k	

Supplementary figure 6

Supplementary figure 7

A

Marker NTC gRNAs PLSCR1 gRNAs Marker

PLSCR1

RPS11

B

Band Intensity Normalized
to RPS11 Control

1.5

1.0

0.5

0.0

NTC PLSCR1

gRNA

C

PLSCR1 vs NTC gRNAs
mRNA-seq

- log10 p-value adjusted

PLSCR1

NPIPB13

0.1 0.5 2.0 10.0

Fold change

Supplementary figure 8

Supplementary figure 9

Supplementary figure 10

A549 no staining

A549

A549-ACE2 CTR no staining

A549-ACE2 CTR

A549-ACE2 PLSCR1 KO no staining

A549-ACE2 PLSCR1 KO

-10^3 0 10^3 10^4

ACE2-FITC

Supplementary figure 11

A

Actin

PLSCR1

WT background

KO background

B

DAPI PLSCR1

PLSCR1

10 μm

10 μm

C

DAPI PLSCR1 SARS-CoV-2 N

PLSCR1 SARS-CoV-2 N

10 μm

10 μm

PLSCR1

SARS-CoV-2 N

10 μm

10 μm

Supplementary figure 12

Supplementary figure 13

A

B

Supplementary figure 14