A Genome-Wide Arrayed CRISPR Screen Reveals PLSCR1 as an Intrinsic Barrier to SARS-CoV-2 Entry

Authors

Jérémie Le Pen[1,*], Gabrielle Paniccia[1,*], Michael Bauer[1,#], H.-Heinrich Hoffmann[1,#], Volker Kinast[2,3,#], Marcela Moncada-Velez[4,#], Ana Pinharanda[5,#], Inna Ricardo-Lax[1,#], Ansgar F. Stenzel[1,6,#], Edwin A. Rosado-Olivieri[7,8,#], Alison W. Ashbrook[1], Kenneth H. Dinnon[1], William C. Doyle[1,9], Catherine A. Freije[1], Seon-Hui Hong[1], Danyel Lee[4,10,11], Tyler Lewy[1], Joseph M. Luna[1,12], Avery Peace[1], Carltin Schmidt[4], William M. Schneider[1], Roni Winkler[1], Chloe Larson[13], Timothy McGinn[13], Miriam-Rose Menezes[13], Lavoisier Ramos-Espiritu[13], Priyam Banerjee[14], John T. Poirier[15], Francisco J. Sànchez-Rivera[16,17], Qian Zhang[4], Jean-Laurent Casanova[4,10,11,18,19], Thomas S. Carroll[20], J. Fraser Glickman[13], Eleftherios Michailidis[1,21,§], Brandon Razooky[1,22,§], Margaret R. MacDonald[1,§], Charles M. Rice[1,§]

[*] Contributed equally to this work
[#] Contributed equally to this work
[§] Contributed equally to this work

Correspondence: Jérémie Le Pen: jlepen@rockefeller.edu; Charles M. Rice: ricec@rockefeller.edu

Affiliations:
1. Laboratory of Virology and Infectious Disease, The Rockefeller University, New York, NY, USA.
2. Department of Medical Microbiology and Virology, Carl von Ossietzky University Oldenburg, Oldenburg, Germany.
3. Department for Molecular and Medical Virology, Faculty of Medicine, Ruhr University Bochum, Bochum, Germany.
4. St Giles Laboratory of Human Genetics of Infectious Diseases, Rockefeller Branch, The Rockefeller University, New York, NY, USA.
5. Department of Biological Sciences, Columbia University, New York, NY 10027, USA.
6. Present address: Department of Infectious Diseases, Molecular Virology, Heidelberg University, Heidelberg, Germany.
7. Laboratory of Synthetic Embryology, The Rockefeller University, New York, NY, USA.
8. Present address: Department of Molecular Pathobiology, New York University.
9. Present address: Laboratory of Host-Pathogen Biology, The Rockefeller University, New York, NY, USA.

10. Laboratory of Human Genetics of Infectious Diseases, Necker Branch, INSERM U1163, Paris,
France, EU.
11. University of Paris, Imagine Institute, Paris, France, EU.
12. Present address: Department of Biochemistry and Center for RNA Science and Therapeutics,
Case Western Reserve University, Cleveland, OH, USA.
13. Fisher Drug Discovery Resource Center, The Rockefeller University, New York, NY, USA.
14. Bio-Imaging Resource Center, The Rockefeller University, New York, NY, USA.
15. Laura and Isaac Perlmutter Cancer Center, New York University Grossman School of Medicine,
NYU Langone Health, New York, NY.
16. Department of Biology, Massachusetts Institute of Technology, Cambridge, 02142,
Massachusetts, USA.
17. David H. Koch Institute for Integrative Cancer Research, Massachusetts Institute of
Technology, Cambridge, 02142, Massachusetts, USA.
18. Department of Pediatrics, Necker Hospital for Sick Children, Paris, France, EU.
19. Howard Hughes Medical Institute, New York, NY, USA.
20. Bioinformatics Resource Center, The Rockefeller University, New York, NY, USA.
21. Present address: Laboratory of Biochemical Pharmacology, Department of Pediatrics, Emory
University School of Medicine, Atlanta, GA, USA.
22. Present address: National Resilience, Inc., La Jolla, CA, USA.

Abstract

Interferons (IFNs) play a crucial role in the regulation and evolution of host-virus
interactions. Here, we conducted a genome-wide arrayed CRISPR knockout screen in the presence
and absence of IFN to identify human genes that influence SARS-CoV-2 infection. We then
performed an integrated analysis of genes interacting with SARS-CoV-2, drawing from a selection
of 67 large-scale studies, including our own. We identified 28 genes of high relevance in both
human genetic studies of COVID-19 patients and functional genetic screens in cell culture, with
many related to the IFN pathway. Among these was the IFN-stimulated gene *PLSCR1*. PLSCR1 did
not require IFN induction to restrict SARS-CoV-2 and did not contribute to IFN signaling. Instead,
PLSCR1 specifically restricted spike-mediated SARS-CoV-2 entry. The PLSCR1-mediated restriction
was alleviated by TMPRSS2 over-expression, suggesting that PLSCR1 primarily restricts the
endocytic entry route. In addition, recent SARS-CoV-2 variants have adapted to circumvent the
PLSCR1 barrier via currently undetermined mechanisms. Our study contributes to understanding
the association between PLSCR1 variants and severe COVID-19 cases reported in a recent GWAS.

Introduction

Viruses maintain a complex relationship with their host cells, co-opting host factors for
their replication while being targeted by cellular defense mechanisms. Such cellular defenses
include the interferon (IFN) pathway, where the infected cell senses foreign molecules and
secretes IFN to trigger an antiviral state in neighboring cells.

Approximately 1-5% of critical COVID-19 patients have mutations that compromise the
production of or response to type I IFNs, while an additional 15% possess autoantibodies that
neutralize type I IFNs [1-7]. This highlights the essential role of type I IFN in the defense against

the SARS-CoV-2 virus that caused the COVID-19 pandemic [8, 9]. Consequently, investigating IFN-
stimulated genes (ISGs) is crucial to our understanding of the remarkable antiviral systems that
evolved in nature. This knowledge could enhance our preparedness for future pandemics.

Several recent studies have identified ISGs restricting SARS-CoV-2. Most of these studies
involved gain-of-function genetic screens, over-expressing individual ISGs. The factors bone
marrow stromal cell antigen 2 (BST2), cholesterol 25-hydroxylase (CH25H), lymphocyte antigen 6
family member E (LY6E), 2'-5'-oligoadenylate synthetase 1 (OAS1), and receptor transporter
protein 4 (RTP4) were notably identified as SARS-CoV-2 antivirals in these studies [9-14]. One
advantage of the gain-of-function approach is that it circumvents potential genetic redundancies
between ISGs [15, 16]. However, this approach is biased towards ISGs that act autonomously
when over-expressed and does not mimic the cellular context of the IFN response, where
hundreds of genes and gene products are differentially regulated to establish an antiviral state.
To counter this limitation, two recent publications examined the effects of ISG loss of function in
IFN-treated cells. They conducted pooled CRISPR knockout (KO) screens in cells pre-treated with
IFN before SARS-CoV-2 infection [17, 18]. By sorting for cells with high SARS-CoV-2 viral load, they
identified SARS-CoV-2 restriction factors such as death domain associated protein (DAXX).

Here, we conducted a human whole-genome arrayed CRISPR KO screen to identify genes
that influence SARS-CoV-2 infection in cells with or without pretreatment with a low dose of IFN.
The arrayed approach, though logistically challenging, has advantages over the pooled format in
capturing both proviral and antiviral genes, genes affecting virus egress, and those coding for
secreted products that exert their impact on neighboring cells. It reliably captures genotype-
phenotype correlations while also unveiling the effects of single gene perturbation on cell growth
and death [19]. We then compiled a comprehensive list of genes interacting with SARS-CoV-2,
incorporating findings from our own screen as well as existing literature. This meta-analysis
revealed several host genes of interest, both previously described and novel. Notably, the ISG
product phospholipid scramblase 1 (PLSCR1) emerged as a prominent antiviral factor. PLSCR1 is
involved in several biological processes [20], including regulating the movement of phospholipids
between the two leaflets of a cell membrane (lipid scrambling) [21] and IFN signaling in the
context of virus infection [22]. Follow-up experiments revealed that PLSCR1 is a cell intrinsic factor
that restricts spike-mediated SARS-CoV-2 entry, independently of the IFN pathway, via currently
undetermined mechanisms. Our genetic screen data and meta-analysis provide a valuable
resource to broaden our understanding of coronavirus infection and innate immunity.
Furthermore, we extend the recent characterization of PLSCR1 as an antiviral against SARS-CoV-
2 impacting COVID-19 outcomes (**Fig. 1**)[18, 23, 24].

Results

A genome-wide arrayed CRISPR KO screen identifies known and novel factors influencing SARS-CoV-2 infection.

While the liver is not the primary target organ of SARS-CoV-2 infection, human
hepatocellular carcinoma Huh-7.5 cells naturally express SARS-CoV-2 dependency factors,
including the receptor angiotensin converting enzyme 2 (ACE2), and proved unexpectedly useful
in SARS-CoV-2 research [18, 25-32]. Huh-7.5 cells do not produce IFN during SARS-CoV-2 infection,

but they do induce an antiviral state in response to IFN pretreatment (**Fig 2A-2C, Supp Tables 1-**
4), making them a convenient model to control the exposure of the cells to IFN.

Using these cells, we conducted a whole-genome arrayed CRISPR KO screen designed to
identify both SARS-CoV-2 proviral and antiviral genes, whose KO reduces or enhances SARS-CoV-
2 infection, respectively. In particular, we aimed to identify factors involved in the IFN response,
from IFN sensing to ISG induction, including effector ISGs that directly influence the virus life
cycle. With this arrayed approach performed in 384-well plates, the cells in each well received a
pool of four gRNAs targeting a single host gene. Each well was then either treated with 1 pM IFN-
α2a or left untreated before being infected with SARS-CoV-2, followed by SARS-CoV-2
nucleoprotein (N) immunofluorescence staining and high content microscopy (**Fig 2D**). This low
dose of IFN was chosen to mimic a cellular environment where IFN triggers an antiviral state
before infection. We anticipated that a saturating amount of IFN would lead to high ISG
transcription and functional redundancy between effectors, biasing hits towards factors in IFN
signaling. In contrast, a low dose of IFN, around the IC50, might enable identification of the
specific roles of individual effector ISGs.

Of the 16,790 screened genes, we selected 16,178 genes where KO did not lead to changes
in cellular fitness assessed by nuclei count (-2 ≤ z-score ≤ 2)(**Supp Fig 1A**). Of these, we selected
12,119 genes expressed in three cell lines relevant for SARS-CoV-2 research (A549, Calu-3, Huh-
7.5 cells) and human lung cells, the primary target cell type *in vivo*, for downstream analysis [33-
35]. We then binned the genes into two groups for data visualization, depending on whether they
were induced by IFN-α2a treatment in Huh-7.5 cells as determined by mRNA-seq (log 2 FC ≥ 2
and padj ≤ 0.05) (**Supp Fig 1B, Supp Tables 5-7**).

Our screen found known and previously unidentified host factors influencing SARS-CoV-2
infection (**Fig 2E**). As expected, positive regulators of IFN signaling, such as interferon-alpha/beta
receptor alpha chain 1 and 2 (IFNAR1,2) [36-38], interferon regulatory factor 9 (IRF9) [39, 40],
Janus kinase 1 (JAK1) [41], and signal transducer and activator of transcription 2 (STAT2) [42, 43]
were antiviral only in IFN pretreated cells. Known negative regulators of IFN signaling, such as
ISG15 ubiquitin-like modifier (ISG15) [44-46], suppressor of cytokine signaling 1 (SOCS1) [47], and
ubiquitin specific peptidase 18 (USP18) [48, 49] were proviral only in IFN pretreated cells.

The SARS-CoV-2 receptor ACE2 was confirmed as proviral with or without IFN
pretreatment. In our mRNA-seq analysis, IFN treatment was found to significantly upregulate
ACE2 mRNA levels (**Supp Fig. 1B**). Prior studies indicate that IFN induces transcription of a
truncated ACE2 isoform, rather than the full-length receptor for SARS-CoV-2 [50, 51].

The lysosomal cysteine protease cathepsin L (CTSL), required for SARS-CoV-2 spike protein
activation [52-54], was a proviral hit in our screens. In contrast, KO of the cell-surface
transmembrane serine protease 2 (TMPRSS2) did not influence infection, suggesting that SARS-
CoV-2 particles primarily enter Huh-7.5 cells through the endocytic pathway that does not depend
on TMPRSS2 (**Fig 1**) [55, 56].

The screen data likely contains false negatives. For example, STAT1 and tyrosine kinase 2
(TYK2) [57, 58] did not influence infection alongside other positive regulators of IFN signaling,
which we attribute to the fact that some gRNAs in the library may have not efficiently directed
Cas9 to cut at their respective target gene loci.

Collectively, the identification of known proviral and antiviral factors confirms the validity
of our screening method.

We performed a Gene Set Enrichment Analysis (GSEA) to identify cellular pathways
exhibiting proviral or antiviral properties in our screen. The full GSEA results, including the genes
driving each pathway enrichment (so-called *leading edge*), can be found in **Supp Table 8**. Some
top pathways ranked by adjusted p-value are summarized in **Fig 2F**. Notably pathways associated
with RNA pol II transcription and mRNA maturation, as well as pathways related to cellular
respiration, exhibited antiviral activity independent of IFN. Surprisingly, pathways associated with
RNA pol III transcription, in part driven by the genes RNA polymerase III subunit A (*POLR3A*) and
RNA polymerase III subunit B (*POLR3B*), were critical to the antiviral response mediated by IFN.
Conversely, factors involved in translation, such as eukaryotic translation initiation factor 3
subunits F and G (EIF3G and EIF3F), likely co-opted for producing viral proteins, were identified
as proviral. Similarly, factors regulating cholesterol homeostasis, likely crucial for SARS-CoV-2
entry [13, 59], were also identified as proviral. For instance, the gene sterol regulatory element
binding transcription factor 2 (*SREBF2*) was one of the top proviral genes (**Fig 2E**).
Our arrayed CRISPR KO screen results thus constitute a valuable resource for research on
coronavirus infection and innate immunity. These can be used to help characterize human genes
influencing SARS-CoV-2 infection and the IFN response.
178
The ISG PLSCR1 is associated with COVID-19 outcomes and exhibits antiviral effects
in functional SARS-CoV-2 genetic screens
To provide a thorough perspective on human genes that impact SARS-CoV-2 infection and
to place our arrayed CRISPR KO screen results within the context of existing research, we have
compiled a table that includes findings from a selection of 67 large-scale 'omic' studies related to
SARS-CoV-2. This compilation encompasses this study and 25 other functional genetic screens for
genes that influence SARS-CoV-2 infection [10-12, 14, 17, 18, 26, 29, 60-76], 24 human genetic
studies that correlate certain alleles with severe COVID-19 outcomes [4, 5, 7, 23, 24, 77-95], ten
publications detailing SARS-CoV-2 protein interactomes [96-105], six focusing on SARS-CoV-2 RNA
interactomes [106-111], and one that examines proteins with altered phosphorylation states in
SARS-CoV-2-infected cells [112] **(Supp Tables 9-10 for the full, and summary tables,**
respectively). This table highlights the depth of research in publications addressing SARS-CoV-2
infection: genes reported in several independent large-scale studies are more credible candidates
for biological relevance (**Supp Fig 2**). As expected, genes associated with the IFN pathway, such
as *IFNAR2*, *OAS1*, and *ZC3HAV1/ZAP*, frequently emerged as significant in SARS-CoV-2 studies.
We focused on 28 genes identified in both human genetic studies of COVID-19 patients
and in functional genetic screens in cell culture, including our own (**Fig 3**). These genes are likely
to have significant physiological relevance and to be well-suited for mechanistic studies in cell
culture. Among these, the ISG *PLSCR1* stood out, being identified as one of the most potent
antiviral genes in our screen (**Fig 2E**). *PLSCR1* variants have been linked to severe COVID-19 in a
recent GWAS (listed in **Table 1**)[23, 24]. This was attributed to a role of PLSCR1 in regulating the
IFN response in COVID-19 patients [24]. Indeed, a pioneering study showed that PLSCR1
potentiates the transcriptional response to IFN-β treatment in human ovarian carcinoma Hey1B
cells [22]. However, PLSCR1 surprisingly appeared as a potent SARS-CoV-2 antiviral even in the
absence of IFN in our screens, suggesting a cell intrinsic, IFN-independent function. In other
words, baseline levels of PLSCR1 may be sufficient to restrict SARS-CoV-2, and IFN pretreatment
could simply enhance this effect by elevating cellular PLSCR1 levels.

Intrinsic PLSCR1 restricts SARS-CoV-2 independently of the IFN pathway.

To better characterize the function of PLSCR1 during SARS-CoV-2 infection, we generated
and validated by western blot (WB) PLSCR1 KO bulk Huh-7.5 and A549-ACE2 lines (**Supp Fig 3A**).
As observed in the arrayed screen (**Supp Fig 1A**), PLSCR1 KO cells were viable (**Supp Fig 3B**).
PLSCR1 depletion increased susceptibility to SARS-CoV-2 independently of IFN pretreatment (**Fig**
4A). Cell treatment with a JAK-STAT inhibitor, which effectively abrogated IFN signaling, confirmed
that intrinsic PLSCR1 limits SARS-CoV-2 infection independently of the IFN signaling pathway (**Fig**
4A). SARS-CoV-2 susceptibility of PLSCR1 KO cells was reversed by the ectopic expression of
PLSCR1 (**Fig 4B, Supp Fig 3C**). Interestingly, while PLSCR1 tagged with an N-terminal FLAG tag
could rescue, PLSCR1 tagged with a C-terminal FLAG tag could not. The C-terminus of the protein
is extracellular, and previous research suggests that this region is important for the protein's
scramblase activity and Ca^{2+} binding [113]. It is possible that the addition of this FLAG-tag
impaired Ca^{2+} binding, affected PLSCR1's localization at the plasma membrane, or otherwise
disrupted the structure of this region, thereby abolishing PLSCR1's antiviral ability. We co-cultured
PLSCR1 reconstituted cells and PLSCR1 KO cells in the same well and infected them with SARS-
CoV-2. A higher proportion of PLSCR1 KO than PLSCR1 reconstituted cells were positive for SARS-
CoV-2 indicating that PLSCR1 acts in a cell autonomous manner (**Fig 4C**). Altogether, these data
suggest that intrinsic PLSCR1 contributes to the restriction of SARS-CoV-2, even without IFN.
224

IFN signaling is unaffected by the loss of PLSCR1 in A549-ACE2 and Huh-7.5 cells.

PLSCR1 has been shown to potentiate ISG transcription in IFN-treated Hey1B cells [22].
We thus hypothesized PLSCR1 might enhance the type I IFN response in A549-ACE2 and Huh-7.5
cells. We investigated PLSCR1's role in the IFN response by infecting Huh-7.5 cells with
chikungunya virus (CHIKV), which is unaffected by PLSCR1 KO without IFN (**Fig 5A**). PLSCR1
depletion did not functionally affect the antiviral effects of IFN treatment (**Fig 5B**). Furthermore,
IFN treatment induced *OAS1* and *IFI6*, two ISGs known to restrict SARS-CoV-2 [10, 12, 17, 91, 92,
94, 95], to a similar extent in both WT and PLSCR1 KO cells, indicating that the IFN signaling
pathway was unaffected by PLSCR1 depletion (**Fig 5C-J**). Finally, PLSCR1 depletion did not alter
basal ISG transcription in the absence of IFN (**Supp Fig 4**).
These findings indicate that PLSCR1 limits SARS-CoV-2 infection independently of the IFN
signaling pathway in A549-ACE2 and Huh-7.5 cells.
237

PLSCR1 restricts SARS-CoV-2 entry.

We hypothesized that PLSCR1 directly targets and inhibits a specific step of the SARS-CoV-
2 life cycle. PLSCR1 primarily localized at the plasma membrane in Huh-7.5 cells (**Fig 6A**).
Furthermore, PLSCR1 depletion led to increased SARS-CoV-2 foci formation (**Fig 6B,C**), and PLSCR1
KO cells did not show increased susceptibility to a SARS-CoV-2 replicon system that bypasses entry
(**Fig 6D**) [114]. In contrast, a single-cycle, replication-defective human immunodeficiency virus
type-1 (HIV-1) particles pseudo-typed with SARS-CoV-2 spike showed enhanced entry in PLSCR1
depleted cells (**Fig 6E, F**) [115]. This data indicates that PLSCR1 restricts SARS-CoV-2 spike-
mediated virion entry. Over-expression of TMPRSS2 lifted the PLSCR1-mediated restriction of
authentic SARS-CoV-2 (**Fig 6G-I**) and of SARS-CoV-2 spike pseudo-typed particles (**Fig 6J-M**),

indicating that PLSCR1 primarily restricts the endosomal entry route. It is possible that the TMPRSS2-dependent entry near the cell surface provides SARS-CoV-2 with some level of evasion from PLSCR1 restriction, as recently described for the ISG NCOA7 [116].

In addition to SARS-CoV-2, we also evaluated the antiviral activity of PLSCR1 against ten viruses that utilize endosomal entry: CHIKV, human parainfluenza virus (hPIV), herpes simplex virus 1 (HSV-1), influenza A virus (IAV), human coronavirus OC43 (hCoV-OC43), human coronavirus NL63 (hCoV-NL63), human coronavirus 229E (hCoV-229E), Sindbis virus (SINV), Venezuelan equine encephalitis virus (VEEV), and vesicular stomatitis virus (VSV). Only SARS-CoV-2 showed a notable susceptibility to PLSCR1's inhibitory effects (**Supp Fig 5**).

Recent variants of SARS-CoV-2 are less restricted by PLSCR1

During the COVID-19 pandemic, SARS-CoV-2 variants evolved from the initial strain, showing increased immune evasion and transmissibility [117-119]. To examine if these variants could circumvent the antiviral action of PLSCR1, we infected WT and PLSCR1 KO Huh-7.5 cells with an early strain isolated in July 2020 (NY-RU-NY1) and the Beta (B.1.352), Delta (B.1.617.2), Omicron (BA.5), and Kraken (XBB.1.5) variants **(Fig 7A-E)**. Although PLSCR1 continued to restrict these later variants, the discrepancy in infection rates between WT and PLSCR1 KO cells diminished compared to the original strain, notably with the Omicron and its descendant, Kraken **(Fig 7F)**. We then infected WT and PLSCR1 KO Huh-7.5 cells with the same focus-forming units (FFU) of each virus, as determined in PLSCR1 KO cells, and conducted a focus forming assay. The ratio of foci in WT cells relative to KO cells increased for later variants, especially for the Omicron subvariant Kraken (XBB.1.5) **(Fig 7G)**. Our data suggests a diminished efficacy of PLSCR1 in restricting the newer SARS-CoV-2 variants.

Association between PLSCR1 variants and severe COVID-19

PLSCR1 encodes a 318 amino acid protein containing a palmitoylation motif and a transmembrane domain which regulate its plasma membrane localization, and a nuclear localization signal (NLS) and transcriptional activation domain thought to be important for its nuclear functions (**Fig 8A**)[20, 120-125].

A recent GWAS has identified an association between PLSCR1 variants and severe COVID-19 outcomes, reporting an odds ratio of approximately 1.2 and a p-value of approximately 10^{-8} (**Table 1**)[23, 24]. In other words, the GWAS suggests that PLSCR1 has a small, but significant effect on severe COVID-19 risks.

GWAS typically identify variants associated with increased odds of a disease, but these variants are not necessarily causative. Among the PLSCR1 variants identified in the COVID-19 GWAS cited above, only rs343320 results in a protein-coding change, specifically His262Tyr, located in the NLS (**Fig 8A**, **Table 1**). Although we cannot dismiss the possibility that (i) some non-coding variants identified in the GWAS could influence the regulation of *PLSCR1* mRNA, potentially leading to functional outcomes, and (ii) the GWAS might have missed nonsynonymous variants impacting PLSCR1 function, we thought to investigate the functional effects of the His262Tyr variant in cell culture. We ectopically expressed PLSCR1 His262Tyr or PLSCR1 WT in A549-ACE2 cells from a lentiviral vector [126]. PLSCR1 His262Tyr and PLSCR1 WT were expressed at similar levels in this system (**Supp Fig 6**). PLSCR1 His262Tyr did not fully rescue the PLSCR1 KO

(**Fig 8B**), indicating that the His262Tyr variant is hypomorphic. Additionally, introducing PLSCR1
His262Tyr into cells already expressing PLSCR1 WT increased their susceptibility to SARS-CoV-2
infection (**Fig 8B**), suggesting a dominant effect. However, this effect might be attributed to the
overexpression of PLSCR1 His262Tyr from the transgene, compared to the natural expression
levels of PLSCR1 WT from the endogenous locus. To counter this, we examined patient-derived
SV40-immortalized fibroblasts expressing ACE2 that were heterozygous for His262Tyr. These cells
were hyper-susceptible to SARS-CoV-2 infection compared to PLSCR1 WT control SV40-fibroblasts
(**Fig 8C**), further suggesting that His262Tyr is dominant. We cannot formally rule out that the
examined SV40-fibroblasts may carry other mutations influencing SARS-CoV-2 infection.
Our data collectively highlight PLSCR1's function in restricting SARS-CoV-2 entry in cell
culture, thereby clarifying the association between PLSCR1 variants and severe COVID-19
outcomes [23, 24]. Future human genetic studies are crucial for determining if certain PLSCR1
variants in the population cause increased risks of severe COVID-19. Our results highlight the
variant rs343320 (His262Tyr) as a potential causative candidate, as it caused increased SARS-CoV-
2 infection in cell culture.

Discussion

Here, we conducted an unbiased arrayed CRISPR KO screen on Huh-7.5 cells infected with
SARS-CoV-2. The screen revealed novel aspects of SARS-CoV-2 and IFN biology while also
confirming previously known facets. Pathways related to mRNA transcription and maturation
were identified as antiviral. This observation may stem from the conflict between the host cell
and SARS-CoV-2, where the host attempts to export mRNAs from the nucleus to facilitate antiviral
responses while the virus replicates in the cytoplasm, impeding nuclear export [127-130]. RNA
Pol III transcription was specifically essential for the IFN-mediated antiviral response, through
mechanisms that are yet to be determined. Interestingly, inborn errors in POLR3A and POLR3C
have been previously described in patients with severe varicella zoster virus infections [131].
Cellular respiration was identified as a key IFN-independent antiviral pathway. Furthermore,
mitophagy was identified as proviral. This may indicate the infected cell's increased demand for
energy and ATP to combat the virus. Alternatively, cellular respiration may have other, yet-to-be-
identified, IFN-independent antiviral roles. Conversely, translation and cholesterol homeostasis
emerged as the foremost proviral pathways. These findings underscore the complex, dualistic
nature of the interactions between SARS-CoV-2 and host cells.
Our screen notably identified the ISG zinc-finger antiviral protein (*ZC3HAV1/ZAP*) as a
proviral factor in IFN-treated cells. Initially, ZC3HAV1/ZAP gained attention as an antiviral factor
that targets the SARS-CoV-2 RNA genome [111] and prevents programmed ribosomal
frameshifting [132]. Yet, a recent study demonstrated that ZC3HAV1/ZAP also promotes the
formation of SARS-CoV-2 non-structural proteins 3 and 4-induced double-membrane vesicles,
essential for virus replication [68]. SARS-CoV-2 may have adapted to exploit certain ISG products,
such as ZC3HAV1/ZAP, within the cellular environment it encounters. It is still unclear if the
seemingly contradictory roles of ZC3HAV1/ZAP – both proviral and antiviral – are caused by
distinct isoforms.
Many other ISG products influenced SARS-CoV-2 infection, PLSCR1 being the most potent
restriction factor. PLSCR1 did not influence ISG induction as previously reported [22], but rather
inhibited spike-mediated SARS-CoV-2 entry through the endocytic route. Our results corroborate

a recent study from Xu et al [18] and provide an explanation for the enrichment for PLSCR1 SNPs observed in a GWAS on severe COVID-19 [23, 24].

The molecular mechanisms of PLSCR1-mediated restriction of SARS-CoV-2 entry remain to be elucidated. PLSCR1 could be altering the lipid composition at the contact site between the virus and endosomal membranes, akin to the ISG IFITM3 for influenza A virus [133-136]. PLSCR1 was first identified as a Ca^{2+}-dependent phospholipid scramblase [21], but it is unclear whether PLSCR1 depletion affects the bidirectional movement of phospholipids *in vivo* [137-139]. A C-terminal FLAG-tag abolished the antiviral ability of reconstituted PLSCR1, possibly by interfering with the function of the Ca^{2+} binding domain. In contrast, inhibiting PLSCR1's phospholipid scramblase activity did not alleviate SARS-CoV-2 restriction [18].

Intriguingly, PLSCR1 specifically restricted SARS-CoV-2 in Huh-7.5 and A549-ACE2 cells but it did not show similar inhibitory effects on other viruses that enter cells via endocytosis. Future studies will investigate the mechanisms behind this specificity. PLSCR1 has been described to inhibit a range of viruses in various cell lines, such as encephalomyocarditis virus, vesicular stomatitis virus, Epstein-Barr virus, hepatitis B virus, hepatitis C virus, human cytomegalovirus, human immunodeficiency virus 1, human T-cell lymphotropic virus type 1, and influenza A virus [22, 140-146]. It has been proposed that PLSCR1 directly binds viral proteins and impairs their functions to restrict the non-coronaviruses cited above, reviewed in [147]. However, it seems unlikely that PLSCR1 has evolved to interact directly with such a diverse set of viral proteins. An alternative explanation is that diverse viral proteins convergently evolved to bind PLSCR1 as a mechanism of immune evasion. Meanwhile, overexpressing PLSCR1 in cell culture could act *like a sponge*, absorbing these viral proteins and thereby hindering viral function. We searched for PLSCR1 interactions in ten SARS-CoV-2 proteins interactome studies, relying on ectopic expression of individual viral proteins [96-98, 100-105, 148], and no interaction was reported in two or more independent studies. Two interactions were reported in a single study: (i) PLSCR1-ORF7b [98], and (ii) PLSCR1-ORF8 [101], both by proximity biotinylation, which was less stringent compared to affinity purification-mass spectrometry (AP-MS) and yeast two-hybrid (Y2H) techniques (**Supp Fig 7**). To date, there is no strong evidence of a direct interaction between a SARS-CoV-2 protein and PLSCR1, although we cannot rule out that such interactions may occur or even appear in the future as SARS-CoV-2 evolves.

Recent SARS-CoV-2 variants, including Omicron (BA.5) and Kraken (XBB.1.5), showed reduced sensitivity to PLSCR1-mediated restriction compared to the New York 2020 strain which served as a reference in our study. Previous research suggests that Omicron, in particular, has developed increased resistance to IFN [149, 150], a trait associated with its highly-mutated spike protein [151]. Omicron's relative resistance to IFN may be due to alternative entry routes that alleviate the restrictions from antiviral ISGs targeting endocytosis, such as PLSCR1 (**Fig 1**). While Omicron favors endocytic entry over TMPRSS-2-dependent entry near the cell-surface [152, 153], recent findings suggest that Omicron can also utilize cellular metalloproteinases for near cell-surface entry, enhancing its infectivity in nasal epithelia [151]. Future research should investigate how SARS-CoV-2 variants evade PLSCR1 and whether this evasion is primarily due to mutations in the spike protein facilitating alternative entry mechanisms or to other factors.

Several PLSCR1 variants were enriched in a GWAS on severe COVID-19, with a relatively low odds ratio of approximately 1.2 [23, 24]. Considering the complex redundancies within antiviral defenses, from innate immunity featuring multiple effector ISGs that restrict SARS-CoV-

2 [9-14, 17, 18], to adaptive immunity [154], the modest odds ratio associated with a single
effector ISG not involved in IFN signaling may not be unexpected. However, for these very reasons,
the identification of PLSCR1 in the GWAS remains noteworthy. Of these enriched PLSCR1 variants,
only rs343320 resulted in a protein-coding change, His262Tyr. Our findings indicate that His262Tyr
exhibits a hypomorphic and dominant effect in cell culture, leading to increased SARS-CoV-2
infection. Future research should aim to ascertain whether rs343320, or potentially other PLSCR1
variants, are directly responsible for elevated risks of severe COVID-19 in patients. It is also
noteworthy to mention the presence of a loss-of-expression/loss-of-function variant in PLSCR1,
specifically p.Ile110AsnfsTer6 (rs749938276). This variant exhibits a minor allele frequency (MAF)
of 0.0003 in the general population, yet it is notably more prevalent among the Ashkenazi Jewish
population, with a maximum MAF (MAFmax) of 0.01. The identification of at least one
homozygous individual in the gnomAD database underscores the viability of individuals deficient
in PLSCR1 [155]. This suggests a potential enrichment of this variant within specific populations,
an aspect that merits further epidemiological and functional exploration to understand its impact
on susceptibility to SARS-CoV-2.

Our findings show that baseline levels of PLSCR1 are effective in limiting SARS-CoV-2
infection. This is in line with other studies where ISGs like *DAXX* and *LY6E* were shown to inhibit
SARS-CoV-2 independently of IFN [17, 156, 157]. mRNA-seq analyses of Huh-7.5 cells and primary
human hepatocytes, as well as data from the GTEx consortium on various human tissues [35,
158], revealed that many ISGs are constitutively expressed, even without IFN stimulation (**Supp**
Fig 8). This supports the idea that the IFN-induced antiviral state results more from enhanced
expression of antiviral genes rather than a binary ON/OFF switch. In future studies, it will be
interesting to explore whether intrinsically expressed ISGs also carry out cellular functions beyond
pathogen defense.

Materials and Methods

Plasmids, oligos, and primers

405   The plasmids, gene fragments, and primers used in this study are listed in **Supp Tables**
406 **11, 12, and 13**, respectively.

Cell Lines

Huh-7.5 (human hepatocellular carcinoma) [159], A549-ACE2 (human lung carcinoma,
generously provided by the laboratory of Brad R. Rosenberg), Lenti-X 293T (Takara, cat. #632180),
Caco2, Vero E6 (*Chlorocebus sabaeus* kidney epithelial cells, ATCC cat. #CRL-1586), BHK-21
(hamster kidney) cells, and SV40-Fribroblasts were cultured in Dulbecco's Modified Eagle Medium
(DMEM, Fisher Scientific, cat. #11995065) supplemented with 0.1 mM nonessential amino acids
(NEAA, Fisher Scientific, cat. #11140076) and 10% fetal bovine serum (FBS, HyClone Laboratories,
Lot. #KTH31760) at 37°C and 5% CO2. All cell lines tested negative for mycoplasma.

Virus stocks

418   **CHIKV-181/25-mKate2:** the infectious clone was a kind gift from Mark Heise (University
419 of North Carolina, USA) [not published yet]. 20 µg of infectious clone DNA was linearized with
420 NotI-HF at 37°C overnight. Complete digestion was confirmed by running a sample of the digested

DNA on a 1% agarose gel. After confirmation, linearized DNA was cleaned via phenol-chloroform extraction, and then ethanol precipitated. The precipitated DNA was resuspended in 20 µL of RNase-free H_2O and in vitro transcribed with an SP6 mMessage mMachine In Vitro Transcription Kit (ThermoFisher, cat. AM1340). The generated RNA was electroporated into 1.2×10^7 BHK-21 cells, and the produced virus was harvested once approximately 80% of the electroporated cells lifted or showed signs of cytopathic effects and 100% of the cells were positive for mKate2 signal. The titer of the virus was 8.5×10^6 focus-forming units (FFU) on Huh-7.5 cells.

hCoV-NL63: was generously provided by Volker Thiel (University of Bern) and amplified at 33°C in Huh-7.5 cells as in [29].

hCoV-OC43: was obtained from ZeptoMetrix (cat. #0810024CF) and amplified at 33°C in Huh-7.5 cells as in [29].

hPIV3-GFP [160]: stock (based on strain JS) grown in VeroE6 cells as in [161].

HSV-1-GFP: stock made by passage on VeroE6 cells. 2×10^7 cells seeded in a T175 flask were infected at an MOI of 0.01 PFU/ml of HSV-1-GFP virus engineered and provided by Ian Mohr [162]. After a one-hour incubation at 37°C, the inoculum was removed, and 20 ml of DMEM supplemented to contain 10% FBS and NEAA was added. Cells were incubated at 37°C for 24 h or until CPE was evident. Cell supernatant containing progeny virus was harvested and titrated on Vero E6 cells (2.4% avicel, fix 2 dpi) at 2.4×10^8 PFU/ml.

IAV WSN (H1N1): was generated in MDCK cells. Cells were inoculated at MOI 0.01 in DMEM supplemented with NEAA, 0.2% BSA, 0.1% FCS, 50 mM Hepes, and 1 µg/ml TPCK-trypsin. Virus-containing culture supernatant was harvested at 52 h post-infection and cleared by centrifugation.

SARS-CoV-2: unless otherwise stated, the isolate SARS-CoV-2/human/USA/NY-RU-NY1/2020 was used in this study [163]. The virus was sourced from the saliva of a deidentified patient in New York City, collected on July 28, 2020. Its sequence is publicly accessible (GenBank OM345241). The virus isolate was amplified in Caco-2 cells. The passage 3 stock employed had a titer of 3.4×10^6 PFU/ml, as measured on Vero E6 cells using a 1% methylcellulose overlay, according to previously described methods [164]. The Beta (B.1.351), Delta (B.1.617.2), Omicron (BA.5), and Kraken (XBB.1.5) variants were obtained from BEI resources (cat. # NR-54008, NR-55611, NR-58616, and NR-59104, respectively), amplified in Vero E6 cells engineered to stably express TMPRSS2, and titer was determined as described above.

SINV Toto1101 [165]: expressing an nsP3-mScarletI fusion reporter was generated by cloning the sequence encoding mScarletI in frame into a unique SpeI restriction site in the pToto1101 infectious clone plasmid as previously described [166]. *In vitro* transcribed, capped RNA was generated from the pToto1101-nsP3-mScarletI plasmid (Invitrogen mMessage mMachine SP6 kit, AM1340) and electroporated into BHK-J cells, a derivative of BHK-21 cells (ATCC, CCL-10) as previously described [166]. 24 hours post electroporation, centrifuge clarified supernatants were aliquoted and stored at -80C. BHK-J cells were cultured and virus stocks generated in MEM supplemented with 7.5% FBS.

VEEV-dsEGFP [15, 167]: the infectious clone plasmid was linearized (MluI) and transcribed *in vitro* using an mMessage mMachine SP6 transcription kit (Ambion). BHK-21 cells were electroporated with viral RNA, and supernatant containing progeny virus was harvested after incubation at 37°C for 30 h or until CPE was evident. Virus was titrated by plaque assay on BHK-21 cells (2.4% avicel, fix 2 dpi). BHK-21: 1.45×10^9 FPU/ml.

VSV-GFP [168]: grown in BHK-21 cells as in [161].

YFV 17D: was generated via transfection of Huh-7.5 with in vitro transcribed RNA from
pACNR-FLYF-17D plasmid as described in [161].

mRNA-seq

mRNA-seq on SARS-CoV-2-infected cells

Cell culture and infection: 75,500 Huh-7.5 cells or 150,000 Calu-3 cells were seeded in
each well of a 12-well plate with 1 mL media. Media: DMEM with 5% FBS and 1% NEAA for Huh-
7.5 cells or EMEM (ATCC, 30-2003) with 10% FBS for Calu-3 cells. The next day, cells were infected
by removing 500 µL of media and adding 500 µL of media with SARS-CoV-2 strain USA-WA1/2020
(BEI Resources, NR-52281) at 5,000 PFU/well (virus titer determined in Huh-7.5 cells). After one
day, the wells were washed with PBS and cells were harvested in 1 mL TRIzol (Invitrogen, cat.
15596-018). N = 3 replicates (separate wells) per sample.

RNA extraction: 2 ml MaXtract High Density tubes (Qiagen, 129056) were centrifugated
at 12,000–16,000 x g for 20-30 second centrifugation. A volume of 750 µL TRIzol-prepared sample
was combined with 150 µL chloroform in these tubes and hand-shaken vigorously. Phase
separation was accomplished by centrifugation at 1500 x g for 5 min at 4°C. The aqueous phase
was then mixed with 400 µL ethanol 95-100% in a separate tube. These preparations were then
transferred to Zymo Research RNA clean and concentrator-25 kit columns (Zymo Research, cat.
R1018) and subjected to multiple wash and centrifugation steps as recommended by the
manufacturer. An in-column DNase I treatment was performed using Qiagen DNase (Qiagen,
79254). Finally, RNA was eluted with 50 µL DNase/RNase-Free water and stored at -80°C.

Sequencing: Poly-A enriched libraries were made using the TruSeq stranded mRNA LT kit
(Illumina, Cat# 20020594) and sequenced on a NovaSeq SP with PE150 read length.

mRNA-seq on IFN-treated cells

Cell culture and treatment: 75,500 Huh-7.5 or 200,000 Calu-3 cells were seeded in each
well of a 12-well plate with 1 mL media. Two days later, the media was replaced with 1 mL of
DMEM with 5%FBS, 1% NEAA in Huh-7.5 or EMEM (ATCC, cat. 30-2003) with 10% FBS for Calu-3
cells with IFN-α2a (PBL, cat. 11101-2) and incubated at 37°C. 24 h later, cells were harvested in
500 µL TRIzol. N = 3 replicates (separate wells) per sample.

RNA extraction as described above.

Sequencing: Poly A-enriched libraries were made using the NEBNext Ultra II RNA Library
Prep Kit for Illumina (NEB, cat. E7770) and sequenced on a NovaSeq SP with PE150 read length.

mRNA-seq on PLSCR1 KO cells

Cell culture and CRISPR KO: 30,000 Huh-7.5 cells were seeded in five wells of a 24-well
plate with 480 uL media. The cells were reverse transfected with 120 uL of a transfection mixture
composed of 250 nM of pooled anti-PLSCR1 or non-targeting Edit-R crRNAs from Horizon
Discovery (cat. CM-003729-01-0002, CM-003729-02-0002, CM-003729-03-0002, and CM-
003729-04-0002 or U-007501-01-05, U-007502-01-05, U-007503-01-05, and U-007504-01-05,
respectively) which had been resuspended with an equimolar amount of Edit-R tracrRNA
(Horizon, cat. U-002005-20) and a 1:200 dilution of Dharmafect 4 (Horizon, cat. T-2004-01). The

following day, the media was changed, and the cells were progressively scaled up to a 6-well plate
over the next 4 days. When the cells were confluent in the 6-well plate, the media was removed
from four of the wells. They were then washed with 1x PBS (cat. 14190-144) and lysed with 1 mL
TRIzol (Life Technologies, cat. 15596-018) for 5 minutes at room temperature before transferring
to an Eppendorf tube and freezing at -80°C to await RNA extraction. The remaining well was lysed
with 300 µL of RIPA buffer (Thermo cat. 89900) supplemented with 1x protease inhibitor (Thermo
cat. 87786) and 1x EDTA and prepared for western blot as described below, in the "Western Blots"
section.

RNA extraction as described above.

Sequencing: Poly A-enriched libraries were made using the NEBNext Ultra II RNA Library
Prep Kit for Illumina (NEB, cat. E7770) and sequenced on a NovaSeq SP with PE150 read length.

mRNA-seq analysis

mRNA-seq reads were first quality-filtered and adapter-trimmed using Trim Galore with
parameters -q 20 -e 0.1 --length 20 --paired and Cutadapt. Reads were then mapped to the human
genome GRCh38 or to a combined SARS-CoV-2 MN985325.1/human genome GRCh38 using STAR
[169] with settings including --runThreadN 8 --outFilterMultimapNmax 1 --twopassMode Basic.
Feature counting was performed using the featureCounts function from the Rsubread package
[170], with strandness specified depending on the sequencing and other parameters as default.
The resulting counts were imported into a DESeqDataSet object using the DESeq2 package [171]
with a design formula of ~Group. Size factors were estimated and normalized counts were
extracted and saved. Differential expression analysis was performed using DESeq with the created
DESeqDataSet object, contrasted by sample groups, cooksCutoff and independentFiltering
disabled, and otherwise default parameters.

Unbiased arrayed CRISPR KO screening

Screen overview

The content of each gRNA 384-well plate constituting the whole-genome library (61
library plates total) was transfected to 16 assay 384-well plates (976 assay plates total). Positive
and negative control gene gRNAs were incorporated into vacant wells of each assay plate as
described below. Huh-7.5 cells were subsequently seeded into these assay plates. The 16 assay
plates served as replicates for three distinct experimental conditions: 4 replicates for mock
treatment followed by mock infection, 5 replicates for IFN-α2a treatment followed by SARS-CoV-
2 infection, and 7 replicates for mock treatment followed by SARS-CoV-2 infection. Each day, three
library 384-well plates were processed, along with their corresponding 48 assay plates. The full
gRNA library, distributed across 61 384-well plates, was completed over a span of 21 days. For
each set of plates, cell seeding was conducted on day 0, IFN-α2a treatment on day 4, SARS-CoV-
2 infection on day 5, and cell fixation on day 6.

gRNA library preparation

A 0.1 nmol Edit-R Human Whole Genome crRNA Library (Horizon, cat. GP-005005-01)
containing four crRNAs per gene and one gene per well (total 0.1 nmol crRNA/well) was
resuspended in 80 µL of a 1.25 µM tracrRNA (Horizon, cat. U-002005-1000) 10 mM Tris-HCL pH
7.4 solution to create a 1.25 µM gRNA solution. The library was then aliquoted in 10 mM Tris-HCL

pH 7.4 in several 96-well plate and 384-well plate copies using a Tecan Freedom EVO liquid
handler. A single-use library copy containing a 40 μL/well of a 312.5 nM gRNA solution in the 384-
well plate format was used in this study.

gRNA reverse transfection (day 0)
In each well of the 384-well assay plates, 40 μL of a transfection solution was prepared by
combining 2% DharmaFect-4 transfection reagent (Horizon, cat. GP-T-2004-07A) in Opti-MEM
(Gibco, cat. 31985070). This was added to 40 μL of a 312.5 nM gRNA library using a Thermofisher
Multidrop Reagent Dispenser, yielding an 80 μL/well transfection mixture. The mixture was left
to incubate at room temperature for 20 minutes. Simultaneously, assay plates were preloaded
with 11 μL/well of serum-free media, which was formulated from DMEM, 1X Antibiotic-
Antimycotic solution (Gibco, cat. 15240-062), and 1X NEAA, dispensed via a Thermofisher
Multidrop Reagent Dispenser. Subsequently, 4 μL/well of the transfection mixture was dispensed
into each of the assay plates (16 assay plates per library plate) using a Tecan Freedom EVO liquid
handler. During this time, Huh-7.5 cells were prepared in media containing 25% FBS, 1X Antibiotic-
Antimycotic solution, and 1X NEAA. A volume of 10 μL cells/well was added to the assay plates,
again using a Thermofisher Multidrop Reagent Dispenser. Ultimately, each well contained 1,250
cells in a 25 μL final volume, with a composition of 25 nM gRNA, 10% FBS, 0.8X Antibiotic-
Antimycotic, and 0.8X NEAA. Plates were then span at 200 g for 5 minutes. To minimize
evaporation, plates were sealed with Breathe-Easy sealing membranes (Sigma-Aldrich, cat.
Z380059) and placed in humid chambers constructed from a 245 mm x 245 mm dish containing
a paper towel moistened with 15 mL of 1X Antibiotic-Antimycotic solution. Four assay plates were
placed in each humid chamber and incubated at 37°C.

IFN-α2a treatment (day 1)
Each well received 5 μL of IFN-α2a (PBL, cat. 11101-2) in media (DMEM, 20% FBS, 1X
Antibiotic-Antimycotic solution, 1X NEAA), using a Thermofisher Multidrop Reagent Dispenser,
for a final concentration of 1 pM IFN-α2a in a final volume of 30 μL. Plates were then span at 200
g for 5 minutes and incubated at 37°C.

SARS-CoV-2 infection (day 5)
Each well received 212.5 PFU SARS-CoV-2 virus (titer determined on Vero E6 cells, see
Virus Stocks section above) diluted in 5 μL of media (DMEM, 20% FBS, 1X Antibiotic-Antimycotic
solution, 1X NEAA) for a final volume of 35 μL in the BSL3. Plates were then span at 200 g for 5
minutes and incubated at 37°C.

Fixing (day 6)
Each well received 50 μL of 20% neutral buffered formalin (Azer Scientific, cat. 20NBF-4-
G) and plates were incubated overnight. The formalin mixture was then removed and each well
received 50 μL of PBS.

IF staining
For IF staining of SARS-CoV-2 infected cells in the arrayed CRISPR KO screen (**Fig 1E**), as
well as some focused experiments (**Fig 3B**, **Fig 3C**, **Fig 4B,C**, **Fig 8C**): the following solutions were
prepared for both 96-well plate (96-wp) and 384-well plate (384-wp): PBS (Phosphate Buffered
Saline), Perm Solution: Comprised of PBS with an added concentration of 0.1% Triton X100,
Blocking Solution: PBS was mixed with 1% BSA. This solution was prepared a day in advance and
filtered before use, PBST: PBS with 0.1% of Tween 20, Primary Antibody Solution: Genetex anti

SARS-CoV-2 N poly rabbit antibody (GTX135357) at a dilution of 1:3000, Secondary Antibody
Solution: AF647 anti-rabbit antibodies at a dilution of 1:3000 and Hoechst 33342 (10 mg/ml) at
1:10,000. Plates were stained on a Biotek EL406 Microplate Washer Dispenser using the following
steps: 1. Priming: The washer was primed with 200 ml of each buffer: PBS, Perm Solution, Blocking
Solution, and PBST. 2. First Washing Phase: Contents of the plates were aspirated. Plates were
then washed with 50 µL/well (384-wp) or 200 µL/well (96-wp) of Perm Solution, followed by a
slow shake for 3 seconds. 3. Permeabilization: A delay of approximately 1 minute was
implemented for permeabilization, in addition to the time required to process all the plates
(around 1 minute per plate). 4. Second Washing Phase: Plates were washed with 50 µL/well (384-
wp) or 200 µL/well (96-wp) of PBS. Subsequently, 50 µL/well (384-wp) or 200 µL/well (96-wp) of
Blocking Solution was added to the plates, followed by a slow shake for 3 seconds. 5. Blocking,
autoclean, and Primary Antibody Priming: The washer was set to undergo an autoclean cycle with
PBS for 30 minutes. Simultaneously, the syringe containing the Primary Antibody Solution was
primed with 16 ml. 6. Third Washing Phase and First Antibody Dispensing: After aspirating the
contents of the plates, 15 µL/well (384-wp) or 60 µL/well (96-wp) from the Primary Antibody
Solution was added, followed by a slow shake for 3 seconds. 7. Primary Antibody Incubation,
autoclean, and Secondary Antibody Priming: The washer was subjected to another autoclean
cycle using PBS for 2 hours and 5 minutes. The syringe containing the Secondary Antibody
Solution was primed with 16 ml during this period. 8. Fourth Washing Phase and Second Antibody
Dispensing: Plates were washed with 50 µL/well (384-wp) or 200 µL/well (96-wp) of PBST,
followed by a 2-second slow shake and aspiration. Then, 15 µL/well (384-wp) or 60 µL/well (96-
wp) from the Secondary Antibody Solution was added, accompanied by a 3-second slow shake.
9. Secondary Antibody Incubation and autoclean: An autoclean cycle with PBS was initiated and
lasted for 1 hour. 10. Final Washing Phase: Plates were washed with 50 µL/well (384-wp) or 200
µL/well (96-wp) of PBST. This was followed by two consecutive washes with 50 µL/well (384-wp)
or 200 µL/well (96-wp) of PBS, incorporating a 2-second slow shake in each cycle. Finally, plates
were left with 50 µL/well (384-wp) or 200 µL/well (96-wp) of PBS.

Imaging
Plates were imaged with a ImageXpress micro-XL and analyzed with MetaXpress
(Molecular Devices).

Analysis
Analysis was conducted in R.

Data Omission: We excluded five library plates, constituting 8% of the total library, due to
insufficient infection levels for accurate quantification.

Normalization: Two variables were subject to normalization—percentage of SARS-CoV-2
positive cells and the count of nuclei. The normalization steps were applied separately for the
three screening conditions: mock treatment followed by mock infection, IFN-α2a treatment
followed by SARS-CoV-2 infection, and mock treatment followed by SARS-CoV-2 infection. Data
was first Z-scale normalized within assay plates:

$$Scale(x) = \frac{x - mean(x)}{sd(x)}$$

And then Z-scale normalized per row and per column to remove any spatial effects.

Statistics: a robust statistic accounting for technical and biological variability was applied
using the below formula within the replicates of each gene**:**

637
$$Stat\ score(x) = \frac{mean(x)}{sd(x)}$$

This statistic was further standardized by Z-scaling across all genes to produce our final z-
score.

Exclusion of genes influencing cell proliferation: 224 genes with nuclei count z-score ≥ 2
and 388 genes with nuclei count z-score ≤ 2 in the mock treatment followed by mock
infection condition were deemed to influence cell proliferation and excluded from
subsequent analyses.

Exclusion of genes not expressed in cell lines of interest (A549, Calu-3, Huh-7.5 cells) and
in human lung cells. Expression data from cell lines from [33, 34]. Expression data from tissues
from [35]. Genes were considered expressed if they had at least one read count within exons.

Gene set enrichment analysis

For the pathway analysis, we leveraged the FGSEA package [172] to perform Gene Set
Enrichment Analysis (GSEA) using gene sets found in the Molecular Signatures Database (MSigDB)
[173]: Reactome [174], KEGG [175], Wikipathways [176], Pathway Interaction Database [177], and
Biocarta [178]. The analysis was conducted separately for two conditions: IFN-α pretreated SARS-
CoV-2 infection and non-pretreated SARS-CoV-2 infection. We attributed a score to each pathway
for both conditions:

655
$$Score = -log10(padj) \times sign(Normalized\ Enrichment\ Score)$$

656   Each pathway was then attributed to one of nine quadrants (as in **Fig 2F**) based on its
657   score in the IFN-α pretreated condition (axis x) versus non-pretreated condition (axis y), using
658   padj ≤ 0.05 as a cutoff.

Compilation of published large-scale omic studies on SARS-CoV-2

661   As a rule, we listed the genes classified as 'hits' by the authors of the respective studies.
662   Below are some exceptions or clarifications:

Functional genetic screens

Baggen, et al. [76]: we used the "low stringency adjusted" analysis in Suppl Table 11.
Proviral: p_value_neg ≤ 0.05 and log2 FC ≥ 1. Antiviral: p_value_pos ≤ 0.05 and log2 FC ≤ -1. We
also used "High stringency" analysis in Suppl Table 7. Proviral: Gene is TMEM106B (log2 FC = 3.8
and p_value_neg = 0.08) or p_value_neg ≤ 0.05 and log2 FC ≥ 1 (no gene matched this criteria).
Antiviral: p_value_pos ≤ 0.05 and log2 FC ≤ -1 (no gene matched this criteria). **Biering, et al. [75]:**
in Supplementary Table 1, in Tab 1: LOF-enriched screen analysis, for proviral genes, we used FDR
≤ 0.05. In Tab 2: GOF-depleted screen analysis, proviral: FDR ≤ 0.05. In Tab 3: GOF-enriched screen
analysis, for antiviral genes, we used FDR ≤ 0.05. **Chan, et al. [74]:** in Multimedia component 6,
for Vero E6 (T16); UM-UC-4 (T23); HEK293+A+T (T12); HuH-7 (T15) and Calu-3 (T43), we
considered gene as hits at FDR < 0.1 (as in Figure 4A). We listed the genes as proviral if differential
≥ 0 or antiviral if differential ≤ 0. **Daniloski, et al. [73]:** we used Table S1. FDR MOI1 ≤ 0.05 or FDR
MOI3 ≤ 0.05. **Danziger, et al. [10]:** in S1 Table, we used the genes annotated as proviral or antiviral
by the authors. **Gordon, et al. [72]:** for A549 +ACE2 in Table S6 or Caco2 in Table S7, for proviral

genes, we used Averaged z-scores ≤ 2 and for antiviral genes, we used Averaged z-scores ≥ 2. **Grodzki, et al. [71]:** for VeroE6 in additional file 4, tab 4, we used FDR <0.25. For HEK293T +Cas9 Study 1 in additional file 6, tab17, we used FDR <0.25. For HEK293T +Cas9 Study 1 in additional file 7, tab7, we used FDR <0.25. **Hoffmann, et al. [26]:** for 37°C (Table S1E) and for 33°C (Table S1C), we selected proviral genes if FDR ≤ 0.05 and z-score ≥ 0 and antiviral genes if FDR ≤ 0.05 and z-score ≤ 0. **Hossain, et al. [65]:** in Figure 3E, we selected the top 15 genes in the spike-mNG axis by negative log robust rank aggregation. **Israeli, et al. [70]:** we used Supplementary Data 1. **Kaur, et al. [14]:** we used the genes labelled in Figure 1. **Le Pen et al. (this study):** we used z-score ≥ 2 for antiviral genes and ≤ 2 for proviral genes, see Unbiased arrayed CRISPR KO Screen analysis section above for more details. **Loo, et al. [69]:** we used the genes labelled in Figure 2. **Mac Kain, et al. [17]:** in Electronic Supplementary Material 5, for antiviral genes, we used the filter: pos|rank ≤13, for proviral genes, we used neg|rank ≤13. **Martin-Sancho, et al. [11]:** we used Table S3, "Lentivirus validated hits". **Pahmeier, et al. [68]:** we used the genes labelled in Figure 6. **Rebendenne, et al. [67]:** for Calu3_Gattinara, we used for proviral genes: residual_z-score_avg ≥ 2.5 and for antiviral genes: residual_z-score_avg ≤ 2.5 (no gene). For VeroE6, proviral: residual_z-score_avg ≥ 2.5, and antiviral: residual_z-score_avg ≤ 2.5. For Caco2, proviral: residual_z-score_avg ≥ 2.5 and antiviral: residual_z-score_avg ≤ 2.5. For Calu3_Calabrese, proviral: residual_z-score_avg ≤ 2 and antiviral: residual_z-score_avg ≥ 2. **Rehfeld, et al. [66]:** In Table S1, we considered genes as hits for PRF-1 top eGFP-mCh or PRF-1 bottom eGFP-mCh if FDR ≤ 0.05. **Schneider, et al. [29]:** for both 37°C (Table_S1A) and 33°C (Table_S1B), we listed the gene as proviral if FDR ≤ 0.05 and z-score ≥ 0 and antiviral if FDR ≤ 0.05 and z-score ≤ 0. **Wang, et al. [63]:** we used Table S1. Proviral: Enrichment score ≤ 10^(-4). **Wei, et al. 2021 [62]:** we used Table S1. For proviral genes, we used Cas9-v1 Avg. ≥ 2.5 & Cas9-v2 Avg. ≥ 2.5. Average between Cas9-v1 Avg. and Cas9-v2 Avg. is given in the table. For antiviral genes, we used Cas9-v1 Avg. ≤ -2.5 & Cas9-v2 Avg. ≤ -2.5. Average between Cas9-v1 Avg. and Cas9-v2 Avg. is given in the table. **Wei, et al. 2023 [61]:** in Table S1, for Day 7 or Day 14, we used fdr ≤ 0.05 for positive regulators of ribosomal frameshifting or negative regulators of ribosomal frameshifting. **Wickenhagen, et al. [12]:** we used the genes labelled in Figure 1B. **Xu, et al. [18]:** in Huh-7.5 or A549-ACE2 cells, in untreated or in IFN-gamma treatment, we used Log10 p-value (mNG-High vs. mNG-Low Enrichment) ≥ 3. **Zhu et al. [60]:** In Supplementary Data 1 , SARS-CoV-2 WT and 2 VOCs tested, for proviral genes, we used pos.score_wt ≤ 0.0005 or pos.score_alpha ≤ 0.0005 or pos.score_beta ≤ 0.0005.

Human genetic studies

Degenhardt, et al. [86]: we used Table 2 and added KANSL1 and TAC4, based on new analysis by Pairo-Castineira et al. 2023 [23]. **Kousathanas, et al. [24]:** we used Table 1 from Pairo-Castineira, et al. 2023 [23]. **Pairo-Castineira, et al. 2021 [94]**: we used Table 1 from Pairo-Castineira, et al. 2023 [23]. **Pairo-Castineira, et al. 2023 [23]:** we only considered variants near annotated genes (i.e., we excluded rs1073165). **Roberts, et al. [79]:** we used Supplementary Table 4. **Zhou, et al. [91]:** we used Table 1 and the p-values from COVID-19 hospitalization (European ancestry only).

SARS-CoV-2 protein interactomes

Davies, et al. [105], Laurent, et al. [98], Li, et al. [104], Samavarch-Tehrani, et al. [96], St-Germain, et al. [99], Stukalov, et al. [103]: we used the Supplementay Table 3 from [25]. **Gordon,**

720 **et al. [97]:** we used the genes listed in Table S2. **Liu, et al. [100] and May, et al. [101]:** we used
721 the genes listed in May, et al. [101] Table S5-new. **Zhou, et al. [102]:** we used the genes listed in
722 Table S1. "SARS-CoV-2-human protein-protein interactions identified in this study."

SARS-CoV-2 RNA interactomes

724 **Flynn, et al. [110], Kamel, et al. [109], Labeau, et al. [108], Lee, et al. [111], Schmidt, et**
725 **al. 2021 [107]:** we used the Supplementay Table 3 from [25]**. Schmidt, et al. 2023 [106]:** we used
726 Table S2, "Huh-7 interactome comparison tab", genes listed in the following categories: "Huh-7
727 gRNA FDR5 HS" and "Huh-7 sgmRNA FDR5 HS".

Altered phosphorylation states in SARS-CoV-2-infected cells.

729 **Bouhaddou, et al. [112]:** for Vero E6 cells, we used Table S1, tab 1 "PhosphoDataFull" and
730 filtered for adj.pvalue ≤ 0.05 & log2FC ≥ 1 or adj.pvalue ≤ 0.05 & log2FC ≤ -1 in at least three
731 different time points.

Generation of PLSCR1 KO cells

CRISPR KO

KO Huh-7.5 and A549-ACE2 cells were generated using two anti-PLSCR1 Edit-R crRNAs
from Horizon Discovery (cat. CM-003729-02-0002 and CM-003729-04-0002) or non-targeting
controls (cat. U-007501-01-05 and U-007502-01-05) resuspended with an equimolar amount of
Edit-R tracrRNA (Horizon, cat. U-002005-20) to form sgRNAs. The sgRNAs were then co-
transfected with Cas9-mKate2 mRNA (Horizon, cat. CAS12218) according to the manufacturer's
protocol. 24 to 48 hours after transfection, cells were examined for mKate2 signal and FACS sorted
into bulk and single cell populations, gating on mKate2 signal. Bulk and single cell populations
were then assessed for PLSCR1 expression by western blot to confirm KO.

Amplicon sequencing

Genomic DNA was isolated from a frozen cell pellet using the Qiagen DNeasy kit (Qiagen,
cat. 69504) and treated with RNAse A in the optional RNA digestion step. The region of interest
was then amplified using Q5 2x mastermix (New England Biolabs, cat. M0492S), 500 ng of
template DNA, 0.5 µM of forward and reverse primers, and the following PCR conditions: 98°C
for 30 seconds, followed by 98°C for 5 seconds, 64°C for 15 seconds, and 72°C for 20 seconds,
repeating those steps 30 times before holding at 72°C for 2 minutes. The primers used when
amplifying PLSCR1 genomic DNA from WT and KO Huh-7.5 and A549+ACE2 cells were RU-O-
32687 (5' AACATAGAGGTGATTATGATTTCGTCT) and RU-O-32526 (5'
GGAGGAGCTTGGATTTCTATCTAC). PCR reactions were run on a 1% agarose gel to confirm
amplification. Amplicons were purified with a Zymo DNA clean and concentrator kit (Zymo, cat.
D4013) before sending to Genewiz for amplicon sequencing.

Western Blots

Cell pellets were collected and lysed in RIPA buffer (Thermo, catalog number 89900) with
1x Halt protease inhibitor cocktail and 1x EDTA (Thermo, catalog number 87786). Cell lysates were
spun down in a refrigerated centrifuge at 15,000 g at 4°C for 15 minutes to pellet any cell debris,
and the supernatant was collected and transferred to another tube. The collected samples were
quantified by BCA assay (Thermo Scientific, cat. #23225). Before loading into the gel, we added
sample buffer (Thermo, catalog number NP0007) with β-mercaptoethanol and heated the sample
at 95°C for 10 minutes. Samples were allowed to cool back to room temperature before loading

into 12% Bis-Tris 1.0 mm gels (Invitrogen, cat. #NP0321BOX). Proteins were electrophoretically
transferred onto nitrocellulose membranes. Membranes were blocked with 5% fat-free milk in 1X
TBS (Thermo, catalog number NP00061) and then incubated with primary antibody at 4°C
overnight in 5% fat-free milk in 1x TBS with 0.5% Tween-20 (TBST). Primary antibody: rabbit anti-
PLSCR1 polyclonal antibody (Proteintech, cat. #11582-1-AP) and mouse anti-β-actin antibody
(Millipore Sigma, cat. A5316-100UL) as a loading control. After incubation, membranes were
washed three times with 1x TBST and then incubated with fluorescently conjugated secondary
antibodies for 2 hours at room temperature. Secondary antibodies: LI-COR IRDye goat anti-rabbit
800 and goat anti-mouse 680 (LI-COR cat. 926-32211 and 926-68070, respectively). Membranes
were washed three times with 1X TBST, once with 1X TBS, then imaged on an Azure 600. For the
western blot in Supplementary Figure 4, this protocol was modified slightly: proteins were
electrophoretically transferred onto 0.22 µm polyvinylidene difluoride (PVDF) membranes,
incubated with a primary antibody solution of rabbit anti-PLSCR1 polyclonal antibody
(Proteintech, cat. #11582-1-AP) and polyclonal rabbit anti-RPS11 antibody (Abcam, cat.
ab157101), a secondary antibody solution of goat anti-rabbit HRP (Invitrogen, cat. 31462) and
visualized using a SuperSignal West Femto Maximum Sensitivity Substate kit (Thermo, cat.
#34096).
780

Cell viability assay
4,000 A549-ACE2 cells/well or 8,000 Huh-7.5 cells/well were seeded on day 0 in 100 µL
media (DMEM, 10% FBS, 1X NEAA, 1X Penicillin-Streptomycin) in a 96-well plate. The next day,
blasticidin selection was added as indicated in the figure to serve as a control for reduced cell
viability. On day 4, cell viability was assessed by resazurin assay (Abcam, cat. ab129732) according
to the manufacturer's protocol.
787

788 JAK-STAT inhibitor treatment
789   InSolution (Millipore, cat. 420097-500UG) was used according to the manufacturer's
790 instructions.
791

Titration of IFN-a2a in CHIKV-infected cells
6,000 Huh-7.5 cells/well were seeded in 100 µL media (DMEM, 10% FBS, 1X NEAA). The
following day, we treated cells with one of twelve concentrations of IFN-a2a (PBL, cat. 11101-2):
64 pM, 32 pM, 16 pM, 8 pM, 4 pM, 2 pM, 1 pM, 0.5 pM, 0.25 pM, 0.125 pM, 0.0625 pM, and 0
pM. The following day, the cells were infected with 2 µL of CHIKV-181/25-mKate2 (approximately
17,000 FFU per well, titer determined on Huh-7.5 cells) and fixed after 12 hours. Plates were
stained with a 1:1000 dilution of Hoechst for at least 10 minutes before washing with PBS and
imaging for mKate2 signal.
800

RT-qPCRs on ISGs
Huh-7.5 and A549-ACE2 cells were seeded at densities of 36,000 or 18,000 cells/well,
respectively, in 500 µL of media (DMEM, 10% FBS, 1X NEAA) in 24-well plates. The following day,
a dilution series of IFN-α2a (PBL, cat. 11101-2) or IFN-β (PBL, cat. #11415) was prepared (64 pM,
32 pM, 16 pM, 8 pM, 4 pM, 2 pM, 1 pM, 0.5 pM, 0.25 pM, 0.125 pM, 0.0625 pM, and 0 pM) and

50 μL of each dilution added to the cells in duplicate. After 24 hours, the media was removed and
the cells were washed with 1 mL of ice-cold PBS. 200 μL of RNA Lysis Buffer (Zymo Research, cat.
#R1060-1-100) was added to the cells, and the plates were frozen at -20°C before RNA isolation.
RNA was extracted using the Zymo Quick RNA 96-kit (Zymo Research, cat. R1052) including
DNaseI treatment, followed by cDNA synthesis using the SuperScript™ IV VILO™ Master Mix
(Invitrogen, cat. 11756050) according to manufacturers' instructions. qPCRs were conducted on
a QuantStudio 3 cycler using the Taqman Fast Advance master mix (Life Technologies Corporation,
cat. 4444965) and the following assays: *RPS11* (ThermoFisher 4331182; Hs01574200_gH), *IFI6*
(ThermoFisher 4331182; Hs00242571_m1), *OAS1* (ThermoFisher 4331182; Hs00973635_m1).
IFI6 and *OAS1* were normalized to *RPS11* mRNA levels using the deltaCt method [179].
816
PLSCR1 subcellular localization
IF staining
A549-ACE2 cells were plated onto #1.5, 12mm glass coverslips (Fisher Scientific, cat.
#1254581) placed at the bottom of the wells of a 24-well plate. When confluent, the cells were
fixed, permeabilized with 1% Triton X-100 for 5 minutes and blocked for 1 hour at room
temperature with 1 mL of PBS-BGT (1x PBS with 0.5% bovine serum albumin, 0.1% glycine, 0.05%
Tween 20). Afterward, the cells were incubated in a 1:500 dilution of 4D2 mouse anti-PLSCR1
antibody (Millipore Sigma, cat. #MABS483) in PBS-BG (1x PBS with 0.5% bovine serum albumin
and 0.1% glycine) overnight at 4C with rocking. The cells were then washed twice with PBS-BGT
before incubation with a secondary antibody solution of 1:1000 anti-mouse 588 (ThermoFisher,
cat. #A-11001) and 1:1000 Hoechst dye (ThermoFisher, cat. #62249) in PBS-BG for two hours at
room temperature, followed by three washes with PBS-BGT.
Imaging
The coverslips were mounted onto slides (Fisher Scientific, cat. #1255015) with Invitrogen
ProLong Gold Antifade Mountant (Fisher Scientific, cat. # P36930). The slides were allowed to
cure for 24 hours before the edges of the coverslips were sealed, and the cells were imaged by
confocal microscopy. Confocal images were acquired using Zeiss Zen Blue (v3.5) software on a
LSM 980 point scanning confocal microscope (Zeiss) hooked to a Axio Observer.Z1 / 7 stand
equipped with C Plan-Apochromat 63X/1.40 oil (RI:1.518 at 23°C) objective lens (Zeiss). CW
excitation laser lines 405 nm and 488 nm were used to excite the fluorescence of DAPI and AF488
labeled samples. Emitted fluorescence were spectrally grated (410-483 nm for DAPI, 499-552 nm
for AF488) to avoid fluorescence bleed through and were detected in MA-PMT (DAPI), and GaAsP-
PMT (AF488). The confocal pinhole was set to 1AU for AF488, and the detector master gains were
set within the linear range of detection (550-750V). Scanned images were saved as .czi files.
841
Focus-forming assay on SARS-CoV-2-infected cells
In **Fig 6B**, Huh-7.5 and A549-ACE2 cells were cultured in media (DMEM with 5% FBS) and
seeded at densities of 2×10^5 and 1×10^5 cells per well, respectively, in collagen-coated 12-well
plates to reach 80-90% confluency by the day of infection. A 1:10 serial dilution of virus stock was
made in Opti-MEM in five separate tubes. Media was aspirated from the cells, and the wells were
washed with 1 ml of PBS before adding 200 μL of each virus dilution to the cells in triplicate. Plates
were incubated at 37°C with 5% CO2 for 1 hour, rocking every 15 minutes for even virus

distribution. A 1% methylcellulose overlay medium was prepared and mixed with complete
growth media at 37°C; 2 ml of this overlay was added to each well after removing the virus
inoculum. Plates were then incubated at 37°C with 5% CO2 for 48 for Huh-7.5 cells or 72 hours
for A549-ACE2 cells. Cells were then fixed in final 10% neutral buffered formalin and IF stained as
described in the Unbiased Arrayed CRISPR screen section. PLSCR1 KO and WT cells were
compared at similar virus dilutions.

In **Fig 7G**, the above protocol was followed to titer SARS-CoV-2 strains on Huh-7.5 PLSCR1
KO cells. Then, Huh-7.5 WT and KO cells were seeded at 2×10^5 cells per well in 1 mL of media
(DMEM, 10% FBS, 1X NEAA) in 12-well plates to reach 80-90% confluency the next day. Media
was aspirated from the cells, the wells were washed with 1mL of PBS, and then the cells were
infected with 50 FFU of SARS-CoV-2 (for each strain) diluted in 200 µL of Opti-MEM. Plates were
then incubated, overlayed with methylcellulose, fixed, and stained as described above.

Transfection with SARS-CoV-2 replicon system

The SARS-CoV-2 replicon and the method for electroporation has been described
previously [114]. Briefly, 6×10^6 Huh-7.5 WT and PLSCR1 KO cells were electroporated at 710 V
with 2 µg of SARS-CoV-2 N mRNA and 5 µg of replicon RNA. The cells rested for 10 minutes at
room temperature before resuspending to a concentration of 300,000 cells/mL and plating 100
µL of cells into each well of a 96-well plate. After 24 hours, supernatant was collected from the
replicon-transfected cells and assayed for *Renilla* luciferase activity according to kit instructions
(Promega, cat. E2810).

Infection with SARS-CoV-2 spike/VSV-G-pseudotyped, single-cycle, replication-defective
HIV-1 viruses.

Virus preparation

SARS-CoV-2 spike/VSV-G-pseudotyped, single-cycle, replication-defective HIV-1 viruses
(pCCNanoLuc/GFP) were prepared as in [115]. Plasmids were a kind gift of Theodora Hatziioannou
and Paul D. Bieniasz (The Rockefeller University, NY, USA) [115]. One day before the transfection,
4×10^6 293T cells were seeded in a 10 cm dish. One hour prior to transfection, the growth media
in the dish was replaced with 9 mL of fresh media containing 2% serum. A 1,000 µL transfection
mixture was prepared, comprising the diluent (a 150 mM NaCl solution prepared with sterile cell
culture water), 5 µg of HIV GP plasmid, 5 µg of pCLG plasmid, and either 2.5 µg of SARS-CoV-2
spike Δ19 or 1 µg of pHCMV.G plasmid, ensuring the total plasmid content did not exceed 12.5
µg. After brief vortexing, 50 µL of PEI (1 mg/mL, Polysciences cat. 23966) was added to achieve a
1:4 DNA/PEI ratio. The mixture was vortexed for 5 seconds and then allowed to sit for 20 minutes
in a hooded environment. Following gentle mixing by pipetting, 1 mL of the transfection mixture
was added to the 10 cm dish. Media was changed 12 hours post-transfection, and the supernatant
was harvested and filtered through a 0.2-micron filter 48 hours post-transfection, then stored at
-80°C.

Infection of PLSCR1 KO or WT cells

Seeded in 96-well plates, 6,000 A549+ACE2 cells per well were cultured in 100 µL of
media. After two days, either 10 µL of SARS-CoV-2 spike pseudotyped virus or 0.01 µL of VSV-G
pseudotyped virus were diluted in a final volume of 100 µL of media and added to the wells to

yield comparable NanoLuc signals. Plates were then spun at 200 g for 5 minutes and incubated at 37°C. Two days post-infection, the media was aspirated and replaced with 50 µL of NanoGlo solution, sourced from the Promega Nano-Glo Luciferase Assay kit (Promega, N1110), with a substrate to buffer ratio of 1:100. NanoLuc signal was subsequently quantified using a Fluostar Omega plate reader.

Infection of siRNA-treated cells

Seeded in 96-well plates, 1,600 HEK293T, HEK293T-ACE2, or HEK293T-ACE2-TMPRSS2 cells were cultured in 80 µL of media. The next day, a 20 µL transfection mixture made of Opti-MEM, 1% DharmaFECT1 (Horizon, T-2001-03), and 250 nM siRNA, PLSCR1 ON-TARGETplus SMARTpool siRNA (Horizon, cat. L-003729-00-0005) or non-targeting control (Horizon, cat. D-001810-10-05) was added to the cells. The final concentration of siRNAs was 25 nM. After two days, either 2 µL of SARS-CoV-2 spike pseudotyped virus or 0.2 µL of VSV-G pseudotyped virus were diluted in a final volume of 100 µL of media and added to the wells to yield comparable NanoLuc signals. Plates were then span at 200 g for 5 minutes and incubated at 37°C. Two days after infection, NanoLuc signal was quantified as described in the "Infection of PLSCR1 KO or WT cells" section.

Infection of siRNA-treated cells with SARS-CoV-2

Seeded in 96-well plates, 1,000 A549, A549-ACE2, or A549-ACE2-TMPRSS2 cells were cultured in 80 µL of media. On the same day, a 20 µL transfection mixture made of Opti-MEM, 1% DharmaFECT1 (Horizon, T-2001-03), and 250 nM siRNA, PLSCR1 ON-TARGETplus SMARTpool siRNA (Horizon, cat. L-003729-00-0005) or non-targeting control (Horizon, cat. D-001810-10-05) was added to the cells. The final concentration of siRNAs was 25 nM. Three days after transfection, the cells were infected by adding 34,000 PFU of SARS-CoV-2 (titer determined on Vero E6 cells) diluted in 10 µL media to each well. Plates were then spun at 200 g for 5 minutes and incubated at 37°C. Staining and readout as described above in the "Unbiased arrayed CRISPR KO screening" section.

Pan-virus infection of PLSCR1 KO cells

A549-ACE2 cells were seeded at a density of 6,000 cells/well in 96-well plates in 90 µL media. The following day, 10 µL diluted virus was added to each well. Virus concentrations as follow: CHIKV-mKate, 0.05 µL virus stock per well (titer 8.5 x 10^6 PFU/mL determined in Huh-7.5 cells); hCoV-NL63, 10 µL virus stock per well (titer 1.4 x 10^5 PFU/mL); hCoV-OC43, 10 µL virus stock per well (titer 1.06 x 10^7 PFU/mL); hPIV-GFP, 0.05 µL virus stock per well; HSV1-GFP, 0.5 µL virus stock per well (titer 2.4 x 10^8 PFU/mL determined on Vero E6 cells); IAV WSN, 0.5 µL virus stock per well; SARS-CoV-2, 0.5 µL virus stock per well (titer 3.4 x 10^6 PFU/mL determined on Vero E6 cells); SINV-Toto1101-mScarletI, 10 µL virus stock per well; VEEV-EGFP, 0.005 µL virus stock per well (titer 1.45 x 10^9 PFU/mL determined on BHK-21); VSV-GFP, 0.05 µL virus stock per well; YFV_17D, 5 µL virus stock per well. The cells were fixed, stained and imaged as described in the *Unbiased arrayed CRISPR KO screen* section. Fluorescent viruses were not stained: the fluorescent signal was used as a reporter. We used the following primary antibodies when applicable: anti-dsRNA (J2) mouse (Nordic MUbio, cat. 10010200) diluted 1:500 was used for hCoV-NL63 and hCoV-OC43, anti-IAV mouse (Millipore, cat. MAB8257) diluted 1:3000, anti-YFV mouse (Santa

Cruz Biotechnology, cat# sc-58083) diluted 1:500, anti-SARS2-S rabbit (Genetex, cat. GTX135357)
diluted 1:3000.

Huh-7.5 cells were transfected with a 1:200 dilution of Dharmafect 4 (Horizon, cat. T-2004-
01) and 25 nM ON-TARGETplus SMARTpool siRNAs (Horizon Discovery) in 96-well plates. The cells
were infected three days after siRNA transfection with hCoV-NL63 or hCoV-OC43 or four days
after siRNA transfection with SARS-CoV-2 or hCoV-229E. IF and imaging as described above.

Reconstitution of WT and mutant PLSCR1

Plasmid cloning

N-terminal 3x FLAG-tagged PLSCR1, C-terminal 3x FLAG-tagged PLSCR1, and PLSCR1
H262Y were generated by designing and ordering large dsDNA gene blocks of PLSCR1 that
contained the desired mutations from IDT. These gene blocks were cloned into the PLSCR1-
SCRPSY vector [180] and confirmed by sequencing (see Supplementary Table 12 for sequences).

Lentivirus production

Lentivirus were generated in Lenti-X 293T cells by transfecting 200 ng VSV-G plasmid, 700
ng Gag-Pol plasmid, and 1100 ng plasmid of interest with lipofectamine 2000 in DMEM
supplemented with 5% FBS. Media was changed 4-6 hours later, and lentivirus harvested at 24
and 48 hours. Lentivirus from both timepoints was pooled, then filtered through a 0.45 µM filter
before aliquoting into 2 mL tubes and freezing at -80°C until use.

Cell transduction

0.3 million cells were transduced in suspension in a 12-well plate. Cells received 8 µg/mL
polybrene and 80 mM Hepes in addition to the lentivirus. Cells were then spinoculated at 37°C
for 1 hour at 1000 x g. The following day, cells were split into two, 6-well plates, then 24 hours
later one of the duplicates was treated with 2 µg/mL puromycin (when using SCRPSY-based
lentiviruses) to select for transduced cells. Further experiments were carried out using the cells
that had approximately 30% transduction before selection.

SARS-CoV-2 infection

In **Fig 8B**, Huh-7.5 and A549+ACE2 cells were plated at 6,000 cells/well and 3,000
cells/well, respectively, in 100 µL of media (DMEM, 10% FBS, 1X NEAA) in 96-well plates. The
following day, cells were treated with IFN (10pM for Huh-7.5 cells and 20pM for A549+ACE2 cells).
On the third day, the Huh-7.5 cells were infected with 0.1 µL of virus per well and the A549+ACE2
cells with 1 µL of virus per well, then spun at 200 g for 5 minutes and incubated at 37°C. Plates
were harvested the next day by fixing and staining as described above. For Figure 7A-E, Huh-7.5
cells were plated at 7,500 cells/well in 100µL of media (DMEM, 10% FBS, 1X NEAA) in 96-well
plates. The next day, cells were infected with quantities of virus that yielded comparable percent
infections in the WT cells, then spun at 200 g for 5 minutes and incubated at 33°C. The quantities
of virus used were as follows: 0.1 µL/well for parental, 0.05 µL/well for beta, 1 µL/well for delta,
0.5 µL/well for omicron, and 0.05 µL/well for kraken. The infected cells were fixed after 24 hours
and stained as described previously.

IF staining and imaging

In **Fig 8B**, cells were stained for IF as described in the *Unbiased arrayed CRISPR KO*
screening section, with different primary and secondary antibody solutions. The primary antibody
solution was a 1:3000 dilution of rabbit anti-SARS-CoV-2 nucleocapsid polyclonal antibody

(Genetex, cat. #GTX135357) in PBS-BGT, and the secondary antibody solution was 1:1000 goat
anti-rabbit 594 (ThermoFisher, cat. #A-11012) or 1:1000 goat anti-rabbit 647 (ThermoFisher, cat.
#A-21245), and 1:1000 Hoechst dye (ThermoFisher, cat. #62249) in PBS-BGT.
980

SARS-CoV-2 infection of human SV40-fibroblasts-ACE2
Generation of human SV40-fibroblasts ACE2 stable cell lines
ACE2 cDNA was inserted with In-Fusion cloning kit (Takara Bio) and using the XhoI and
BamHI restriction sites into linearized pTRIP-SFFV-CD271-P2A in accordance with the
manufacturers' instructions. We checked the entire sequence of the *ACE2* cDNA in the plasmid
by Sanger sequencing. Then, HEK293T cells were dispensed into a six-well plate at a density of
8×10^5 cells per well. On the next day, cells were transfected with pCMV-VSV-G (0.2 µg), pHXB2-
env (0.2 µg; NIH-AIDS Reagent Program; 1069), psPAX2 (1 µg; Addgene plasmid no. 12260) and
pTRIP-SFFV-CD271-P2A-ACE2 (1.6 µg) in Opti-MEM (Gibco; 300 µL) containing X-tremeGene-9
(Sigma Aldrich; 10 µL) according to the manufacturers' protocol. After transfection for 6 h, the
medium was replaced with 3 mL fresh culture medium, and the cells were incubated for a further
24 h for the production of lentiviral particles. The viral supernatant was collected and passed
through a syringe filter with 0.2 µm pores (Pall) to remove debris. Protamine sulfate (Sigma;
10 µg/mL) was added to the supernatant, which was then used immediately or stored at −80 °C
until use.
For the transduction of SV40-fibroblasts with ACE2, 5×10^5 cells per well were seeded in
six-well plates. Viral supernatant was added (500 µL per well). The cells were then further
incubated for 48 h at 37 °C. Cells were keep in culture and after 8 days, transduction efficiency
was evaluated by CD271 surface staining with CD271 AlexaFluor 647, 1:200 dilution (BD
Pharmigen 560326). MACS-sorting was performed with CD271 positive selection beads (Miltenyi
Biotec) if the proportion of CD271-positive cells was below 80% [7, 163].
Infection
5,000 cells per well were seeded in a 96-well plate and infected the next day with SARS-
CoV-2 at MOI = 0.05, using a titer determined on Vero E6 cells. Cells were fixed at 2 dpi, stained
and imaged as described in the *Unbiased arrayed CRISPR KO screen* section.
1006

Acknowledgments
We thank Georgia McClain for reading and editing this manuscript. We thank the staff at
the Laboratory of Virology and Infectious Disease: Ellen Castillo, Michela De Santis, Arnella Norris,
Aileen O'Connell, Santa Maria Pecoraro Di Vittorio, Glen Santiago, and Sonia Shirley. Ching-Wen
Chang and Lihong Liu (Columbia University, NY, USA), and Theodora Hatziioannou and Paul D.
Bieniasz (The Rockefeller University, NY, USA) generously provided plasmids and instructions to
generate SARS-CoV-2 spike-pseudotyped, single-cycle, replication-defective HIV-1 viruses [115].
Oded Danziger and Brad R. Rosenberg (Department of Microbiology at the Icahn School of
Medicine at Mount Sinai, NY, USA) kindly provided the A549-ACE2 cells [10] used in this study.
FACS was conducted at the Flow Cytometry Resource Center at Rockefeller University. mRNA-seq
was performed by the Genomics Resource Center at The Rockefeller University and by Novogene.

Confocal microscopy was performed in the Rockefeller University's Bio-Imaging Resource Center,
RRID:SCR_017791. We thank Ankit Patel, Sales Manager at Horizon Discovery.
Trim Galore was developed at The Babraham Institute by @FelixKrueger, now part of Altos
Labs. The Genotype-Tissue Expression (GTEx) Project [35] was supported by the Common Fund of
the Office of the Director of the National Institutes of Health, and by NCI, NHGRI, NHLBI, NIDA,
NIMH, and NINDS. The data used for the analyses described in this manuscript were obtained
from the EMBL-European Bioinformatics Institute portal,
"https://www.ebi.ac.uk/gxa/experiments/E-MTAB-5214/Results", on October 1st, 2020.
Work in the Laboratory of Virology and Infectious Disease was supported by NIH grants
P01AI138398-S1, 2U19AI111825, R01AI091707-10S1, and R01AI161444; a George Mason
University Fast Grant; the G. Harold and Leila Y. Mathers Charitable Foundation; the Meyer
Foundation; the Pilot Project Robertson Therapeutic Development Fund at The Rockefeller
University; and the Bawd Foundation. J.L.P. was supported by the Francois Wallace Monahan
Postdoctoral Fellowship at The Rockefeller University and the European Molecular Biology
Organization Long-Term Fellowship (ALTF 380-2018). G.P. was supported by the James H. Gilliam
Fellowship for Advanced Study from the Howard Hughes Medical Institute and the Graduate
Research Fellowship Program from the National Science Foundation (FAIN 1946429). M.B. was
supported by a Swiss National Science Foundation fellowship (P500PB_203007).

1036 Data and code availability

1037 To be determined.

1038

1039 Tables

1040 Table 1. PLSCR1 variants associated with severe COVID-19 in GWAS [23, 24]

1041

GWAS p-value	rsID	Genomic Coordinate (GRCh38)	Nucleotide Change	Gene	Functional Consequence
7.52E-07	rs116553931	chr3:146430956	C:T	PLSCR2	intron variant
1.08E-07	rs454645	chr3:146514682	C:T	PLSCR1	Downstream transcript variant
5.43E-08	rs343320	chr3:146517122	G:A	PLSCR1	His262Tyr
8.21E-08	rs343318	chr3:146518204	T:C	PLSCR1	intron variant
1.52E-07	rs343317	chr3:146518374	A:G	PLSCR1	intron variant
1.00E-07	rs186910	chr3:146520241	A:G	PLSCR1	intron variant
1.13E-07	rs173150	chr3:146520256	A:T	PLSCR1	intron variant

1.35E-07	rs71302408	chr3:146520389	T:C	PLSCR1	intron variant
7.06E-08	rs343316	chr3:146521151	A:G	PLSCR1	intron variant
4.64E-08	rs343314	chr3:146522652	C:T	PLSCR1	intron variant
7.46E-08	rs343312	chr3:146522970	G:A	PLSCR1	intron variant

1042

Figure Legends

Figure 1. Known ISGs restricting SARS-CoV-2 entry.

A schematic of SARS-CoV-2 entry and the sites where known ISG entry restriction factors function is shown. CD74 suppresses endolysosomal cathepsins, enzymes that process certain viral glycoproteins to make them fusion-competent [181, 182]. CH25H facilitates the sequestration of accessible cholesterol, which results in decreased virus-cell membrane fusion and viral entry [13, 59]. NCOA7 accelerates the acidification of the lysosome, leading to the degradation of viral antigens [116, 183]. LY6E and PLSCR1 restrict virus-cell membrane fusion at the endosome through unknown mechanisms, see [18, 157] and this study.

Figure 2. Unbiased arrayed CRISPR KO screens reveal IFN-dependent and independent genes influencing SARS-CoV-2 infection.

A. mRNA-seq comparison between Huh-7.5 and Calu-3 cells, focusing on a subset of 224 ISGs, in response to 24 h SARS-CoV-2 infection MOI 0.03. Red diamond, PLSCR1 RNA level. Viral RNA levels were comparable in both cell lines (not shown). ****, $p<0.0001$; two-tailed t-test.

B. Cells were treated with 0.5 nM IFN-α2a for 24 h. mRNA-seq analysis as in **(A)**.

C. Huh-7.5-Cas9 cells were pretreated with different amounts of IFN-α2a, then infected with SARS-CoV-2 for 24 h followed by IF staining for SARS-CoV-2 N protein; n = 6; error bars represent SEM. ****, $p<0.0001$; two-tailed t-test.

D. Diagram of the arrayed CRISPR KO screen method.

E. The virus level (percentage infected cells was determined by IF staining, then normalized and z-score was calculated) is plotted for 24 h 0 pM (y axis) or 1 pM (x axis) IFN-α2a pretreatment followed by 24 h infection (n ≥ 5). The genes were categorized as ISG or other based on mRNA-seq of IFN-α2a-treated cells as in **(B)**. ISGs were defined by a fold change ≥ 2 and padj ≤ 0.05 in the IFN-treatment versus untreated pairwise comparison.

F. Gene set enrichment analysis conducted on the arrayed CRISPR KO screens data represented in **(E)**. Description of the top pathways ranked by p-value for each quadrant. Databases: [1]Reactome; [2]WikiPathways; [3]Pathway Interaction Database; [4]KEGG; [5]Biocarta.

1073 Figure 3. Human genes significant in human genetics studies on COVID-19 patients and in
1074 functional genetic screens in cell culture
1075

Figure 4. PLSCR1 is a highly effective anti-SARS-CoV-2 effector ISG contributing to intrinsic
immunity in the absence of IFN.
A. Cells were pretreated with a JAK-STAT inhibitor (InSolution 1 µM) for 2 h, followed by IFN-
α2a (10 pM Huh-7.5 or 20 µM A549-ACE2) for 24 h and were infected with SARS-CoV-2 for
24 h followed by IF staining for viral N protein. Huh-7.5 infection using an MOI of 0.5 (titer
determined by focus forming assay on Huh-7.5 WT cells). A549-ACE2 infection using an
MOI of 0.01 (titer determined by focus forming assay on A549-ACE2 WT cells). The
percentage of SARS-CoV-2 positive cells is plotted.
B. Cells were reconstituted with the indicated proteins by stable transduction with
lentiviruses, then infected as in **(A)**.
C. Cells were co-cultured as indicated (50:50 mix), then infected as in **(A)** and the % infection
of each cell type was determined. n = 6; error bars represent sd. ****, $p < 0.0001$; two-
tailed t-test.
1089

Figure 5. PLSCR1 is not important for IFN signaling.
A. Huh-7.5 WT and PLSCR1 KO cells were infected with 17,000 FFU of CHIKV 181/25 mKate2
for 12 or 24 hours, then fixed, IF stained for nuclei, and the percentage of positive cells
determined by imaging for mKate2 reporter signal. n = 12 independent infections
(separate wells). Error bars represent sd. ns, non-significant; two-way ANOVA.
B. Huh-7.5 cells, PLSCR1 KO as indicated, were pretreated with different amounts of IFN-
α2a, then infected with 17,000 FFU of CHIKV-mKate for 12 h; n = 7; error bars represent
sd.
C-J. Cells were treated for 24 h by IFN, as indicated, followed by RT-qPCRs on ISGs.
1099

Figure 6. PLSCR1 restricts spike-mediated SARS-CoV-2 entry.
A. A549-ACE2 cells were IF stained using an anti-PLSCR1 antibody (white) and Hoechst
33342 nuclear staining (blue) and imaged at 63X magnification on a confocal microscope.
B. Focus forming assays: SARS-CoV-2 N IF (red) and Hoechst 33342 nuclear staining (blue)
on similarly infected WT or PLSCR1 KO Huh-7.5 and A549-ACE2 cells after 2 and 3 d,
respectively.
C. Quantification of **(B)**.
D. Huh-7.5 WT and PLSCR1 KO cells electroporated with SARS-CoV-2 replicon which
produces a secreted luciferase. Luciferase activity assayed 24 hours after electroporation.
n = 36 (separate wells from single electroporation). Error bars represent sd. Ns = non-
significant; two-tailed t-test.
E-F. Transduction of A549-ACE2 cells with an HIV-based replicon expressing the
nanoluciferase pseudotyped with VSV-G or SARS-CoV-2 spike, respectively. n = 5
independent infection (separate wells). Nanoluciferase signal measured 2 dpi. Error bars
represent sd. ns, non-significant, **, $p < 0.01$; two-tailed t-test.

G-I. A549 cells WT, expressing ACE2, or expressing ACE2-TMPRSS2 as indicated were transfected with PLSCR1 or non-template control (NTC) siRNAs as indicated for 3 d and infected with SARS-CoV-2 for 1 d. SARS-CoV-2 N was stained by IF and the percentage of positive cells was determined by imaging. n = 6 independent infection (separate wells). Error bars represent sd. ns, non-significant, **, p<0.01, ***, p<0.001; two-tailed t-test.

J-M. HEK293T cells expressing ACE2 or ACE2-TMPRSS2, as indicated, were transfected with siRNA knockdown of PLSCR1 or non-template control (NTC), as indicated, for 3 d and transduced with an HIV-based replicon expressing the nanoluciferase pseudotyped with VSV-G or SARS-CoV-2 spike, as indicated for 2 d. n = 3 independent infection (separate wells). Error bars represent sd. ns, non-significant, **, p<0.01; two-tailed t-test.

Figure 7. Newer variants of SARS-CoV-2 are less restricted by PLSCR1

A-E. Infection of Huh-7.5 cells with SARS-CoV-2 (parental) or its descendant variants, Beta, Delta, Omicron, and Kraken for 24 hours. SARS-CoV-2 N was stained by IF and the percentage of positive cells determined by imaging. n = 10 independent infection (separate wells). Error bars represent sd. ****, p < 0.0001; two-tailed t-test.

F. Ratio of WT/KO percent infection from A-E. Error bars represent sd. ns, non-significant. ***, p<0.001, ****, p<0.0001; one-way ANOVA.

G. Huh-7.5 WT and PLSCR1 KO cells were infected with 50 FFU of virus as titered on PLSCR1 KO cells in a focus-forming assay. Plaques were counted, and then a ratio of WT-to-KO plotted for each SARS-CoV-2 variant. n = 6 independent infection (separate wells). Error bars represent sd. **, p<0.01; one-way ANOVA.

Figure 8. PLSCR1 His262Tyr, which associates with severe COVID-19, leads to higher SARS-CoV-2 infection in cell culture.

A. Protein diagram of PLSCR1. Domain coordinates from UniProt.

B. Huh-7.5 cells, WT and PLSCR1 KO, stably expressing N-terminal FLAG-tagged Firefly Luciferase (Fluc), N-terminal FLAG-tagged PLSCR1, or N-terminal FLAG-tagged PLSCR1 H262Y mutant and infected with SARS-CoV-2 for 24 hours. SARS-CoV-2 N was stained by IF and the percentage of positive cells determined by imaging. n = 15 independent infection (separate wells). Error bars represent sd. ****, p<0.0001; two-tailed t-test.

C. SV40-Fibroblast-ACE2 cells, genotype as indicated, infected for two days with SARS-CoV-2. N = eight independent infection in separate wells. ns, non-significant, **, p<0.01, ***, p<0.001; two-tailed t-test.

Supplementary Figure Legends

Supplementary Figure 1.

A. Nuclei count (z-score) in arrayed genetic screen. Examples of VCP (essential gene) and PLSCR1 (SARS-CoV-2 antiviral hit) are plotted.

B. Volcano plot of Huh-7.5 cells mRNA-seq treated with 0.5 nM IFN-α2a for 24 h as in **Fig 2B**.

Supplementary Figure 2

    A. Occurrence of human genes interacting with SARS-CoV-2 drawn from a selection of 67 large-scale studies. The occurrence reflects the number of independent studies finding each gene as significant.

    B. Upset plot on data as in **(A)**, showing the overlap in significant genes in large-scale SARS-CoV-2 studies by category.

Supplementary Figure 3.

    A. Western blot on PLSCR1 KO cells against PLSCR1 (green) and ß-actin (red).

    B. Cells as indicated were seeded at similar density, treated or not with Blasticidin (used here as a control to decrease cell viability), and cultured for 4 days before resazurin cell viability assay. n = 4 independent wells.

    C. Western blot on PLSCR1 WT and KO A549-ACE2 cells against PLSCR1 (green) and ß-actin (red).

Supplementary Figure 4.

    A. Western blot against PLSCR1 and RPS11. Cas-9-expressing Huh-7.5 cells were transfected with 4-gRNA pools targeting PLSCR1 or non-template control (as indicated) and cells were in culture for 7 d.

    B. Quantification of bands intensity in **(A)**.

    C. mRNA-seq on cells as in **(B)**.

Supplementary Figure 5.

    A-I. Infection of A549-ACE2 cells with viruses as indicated. IF staining was used for IAV WSN, hCoV-OC43 and SARS-CoV-2, otherwise a fluorescent reporter was used. Percentage of virus positive cells was determined by imaging. N = six replicates (independent infections in separate wells), error bars represent SD.

    J-M. Infection of Huh-7.5 cells with viruses as indicated. siRNA knockdown of PLSCR1 vs non-template control (NTC). ****, p<0.0001; ns, non-significant; two-tailed t-test.

Supplementary Figure 6.

Western Blot against PLSCR1 and β-actin (loading control). Huh-7.5 PLSCR1 KO cells were stably transduced with FLAG-tagged Fluc, FLAG-tagged PLSCR1, or FLAG-tagged PLSCR1 H262Y.

Supplementary Figure 7.

Comparison between selected SARS-CoV-2 protein interactome studies [96-98, 100-105, 148].

Supplementary Figure 8.

Comparison between the relative mRNA levels of 97 hallmark IFN-α-stimulated genes and the remaining transcriptome in cell lines as indicated. Red diamond, *PLSCR1* RNA level. For Huh-7.5 cells mRNA-seq, cells treated with 0.5 nM IFN-α2a for 24 h as in **Fig 2B**. For primary human

1196 hepatocytes, cells were treated with 0.1 nM IFN-α2a for 24 h (full data and methods to be
1197 released elsewhere). For the human tissues, data from [35].
1198

Supplementary Material

1199 Supplementary Material
1200 Supplementary Table 1.
1201 mRNA-seq on SARS-CoV-2 infected cells, Normalized reads, related to **Fig 2A**.
1202

1203 Supplementary Table 2.
1204 mRNA-seq on SARS-CoV-2 infected cells, Differential Gene Expression.
1205

1206 Supplementary Table 3.
1207 mRNA-seq on IFN treated cells, Normalized reads, related to **Fig 2B**, and **Supp Fig 8.**
1208

1209 Supplementary Table 4.
1210 mRNA-seq on IFN treated cells, Differential Gene Expression, related to **Fig 2E** and **Supp Fig 1B**.
1211

1212 Supplementary Table 5.
1213 Arrayed CRISPR KO screen, raw data.
1214

1215 Supplementary Table 6.
1216 Arrayed CRISPR KO screen, analyzed data. Related to **Fig. 2E** and **Supp Fig 1A**.
1217

1218 Supplementary Table 7.
1219 Arrayed CRISPR KO screen, summary table.
1220

1221 Supplementary Table 8.
1222 GSEA on arrayed CRISPR KO screen. Related to **Fig. 2F**.
1223

1224 Supplementary Table 9.
1225 Consolidated list of human genes classified as hits in selected SARS-CoV-2 studies, full table.
1226 Related to **Fig 3** and **Supp Fig 2.**
1227

1228 Supplementary Table 10.
1229 Consolidated list of human genes classified as hits in selected SARS-CoV-2 studies, summary table.
1230 Related to **Fig 3** and **Supp Fig 2.**
1231

1232 Supplementary Table 11.
1233 Plasmids used in this study.
1234

1235 Supplementary Table 12.
1236 Gene fragments used in this study.
1237
1238 Supplementary Table 13.
1239 Primers used in this study.
1240

References

1.   Bastard P, Zhang Q, Zhang SY, Jouanguy E, Casanova JL. Type I interferons and SARS-CoV-
2: from cells to organisms. Curr Opin Immunol. 2022;74:172-82. Epub 20220125. doi:
10.1016/j.coi.2022.01.003. PubMed PMID: 35149239; PubMed Central PMCID:
PMCPMC8786610.
2.   Bastard P, Gervais A, Le Voyer T, Rosain J, Philippot Q, Manry J, et al. Autoantibodies
neutralizing type I IFNs are present in ~4% of uninfected individuals over 70 years old and
account for ~20% of COVID-19 deaths. Sci Immunol. 2021;6(62). Epub 2021/08/21. doi:
10.1126/sciimmunol.abl4340. PubMed PMID: 34413139.
3.   Lopez J, Mommert M, Mouton W, Pizzorno A, Brengel-Pesce K, Mezidi M, et al. Early
nasal type I IFN immunity against SARS-CoV-2 is compromised in patients with autoantibodies
against type I IFNs. J Exp Med. 2021;218(10). Epub 20210806. doi: 10.1084/jem.20211211.
PubMed PMID: 34357402; PubMed Central PMCID: PMCPMC8352718.
4.   Asano T, Boisson B, Onodi F, Matuozzo D, Moncada-Velez M, Maglorius Renkilaraj MRL,
et al. X-linked recessive TLR7 deficiency in ~1% of men under 60 years old with life-threatening
COVID-19. Sci Immunol. 2021;6(62). doi: 10.1126/sciimmunol.abl4348. PubMed PMID:
34413140; PubMed Central PMCID: PMCPMC8532080.
5.   Zhang Q, Bastard P, Liu Z, Le Pen J, Moncada-Velez M, Chen J, et al. Inborn errors of type
I IFN immunity in patients with life-threatening COVID-19. Science. 2020;370(6515). Epub
20200924. doi: 10.1126/science.abd4570. PubMed PMID: 32972995; PubMed Central PMCID:
PMCPMC7857407.
6.   Bastard P, Rosen LB, Zhang Q, Michailidis E, Hoffmann HH, Zhang Y, et al. Autoantibodies
against type I IFNs in patients with life-threatening COVID-19. Science. 2020;370(6515). Epub
20200924. doi: 10.1126/science.abd4585. PubMed PMID: 32972996; PubMed Central PMCID:
PMCPMC7857397.
7.   Zhang Q, Matuozzo D, Le Pen J, Lee D, Moens L, Asano T, et al. Recessive inborn errors of
type I IFN immunity in children with COVID-19 pneumonia. J Exp Med. 2022;219(8). Epub
20220616. doi: 10.1084/jem.20220131. PubMed PMID: 35708626; PubMed Central PMCID:
PMCPMC9206114.
8.   Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, et al. A pneumonia outbreak
associated with a new coronavirus of probable bat origin. Nature. 2020;579(7798):270-3. Epub
20200203. doi: 10.1038/s41586-020-2012-7. PubMed PMID: 32015507; PubMed Central
PMCID: PMCPMC7095418.
9.   Savan R, Gale M, Jr. Innate immunity and interferon in SARS-CoV-2 infection outcome.
Immunity. 2023;56(7):1443-50. doi: 10.1016/j.immuni.2023.06.018. PubMed PMID: 37437537;
PubMed Central PMCID: PMCPMC10361255.

10.   Danziger O, Patel RS, DeGrace EJ, Rosen MR, Rosenberg BR. Inducible CRISPR activation
screen for interferon-stimulated genes identifies OAS1 as a SARS-CoV-2 restriction factor. PLoS
Pathog. 2022;18(4):e1010464. Epub 20220414. doi: 10.1371/journal.ppat.1010464. PubMed
PMID: 35421191; PubMed Central PMCID: PMCPMC9041830.

11.   Martin-Sancho L, Lewinski MK, Pache L, Stoneham CA, Yin X, Becker ME, et al. Functional
landscape of SARS-CoV-2 cellular restriction. Mol Cell. 2021;81(12):2656-68 e8. Epub 20210413.
doi: 10.1016/j.molcel.2021.04.008. PubMed PMID: 33930332; PubMed Central PMCID:
PMCPMC8043580.

12.   Wickenhagen A, Sugrue E, Lytras S, Kuchi S, Noerenberg M, Turnbull ML, et al. A
prenylated dsRNA sensor protects against severe COVID-19. Science. 2021;374(6567):eabj3624.
Epub 20211029. doi: 10.1126/science.abj3624. PubMed PMID: 34581622.

13.   Zang R, Case JB, Yutuc E, Ma X, Shen S, Gomez Castro MF, et al. Cholesterol 25-
hydroxylase suppresses SARS-CoV-2 replication by blocking membrane fusion. Proc Natl Acad Sci
U S A. 2020;117(50):32105-13. Epub 20201125. doi: 10.1073/pnas.2012197117. PubMed PMID:
33239446; PubMed Central PMCID: PMCPMC7749331.

14.   Kaur R, Tada T, Landau NR. Restriction of SARS-CoV-2 replication by receptor transporter
protein 4 (RTP4). mBio. 2023;14(4):e0109023. Epub 20230629. doi: 10.1128/mbio.01090-23.
PubMed PMID: 37382452; PubMed Central PMCID: PMCPMC10470548.

15.   Schoggins JW, Wilson SJ, Panis M, Murphy MY, Jones CT, Bieniasz P, et al. A diverse range
of gene products are effectors of the type I interferon antiviral response. Nature.
2011;472(7344):481-5. Epub 20110410. doi: 10.1038/nature09907. PubMed PMID: 21478870;
PubMed Central PMCID: PMCPMC3409588.

16.   Schoggins JW. Interferon-Stimulated Genes: What Do They All Do? Annu Rev Virol.
2019;6(1):567-84. Epub 2019/07/10. doi: 10.1146/annurev-virology-092818-015756. PubMed
PMID: 31283436.

17.   Mac Kain A, Maarifi G, Aicher SM, Arhel N, Baidaliuk A, Munier S, et al. Identification of
DAXX as a restriction factor of SARS-CoV-2 through a CRISPR/Cas9 screen. Nat Commun.
2022;13(1):2442. Epub 20220504. doi: 10.1038/s41467-022-30134-9. PubMed PMID:
35508460; PubMed Central PMCID: PMCPMC9068693.

18.   Xu D, Jiang W, Wu L, Gaudet RG, Park ES, Su M, et al. PLSCR1 is a cell-autonomous
defence factor against SARS-CoV-2 infection. Nature. 2023;619(7971):819-27. Epub 20230712.
doi: 10.1038/s41586-023-06322-y. PubMed PMID: 37438530; PubMed Central PMCID:
PMCPMC10371867.

19.   Kim HS, Lee K, Kim SJ, Cho S, Shin HJ, Kim C, et al. Arrayed CRISPR screen with image-
based assay reliably uncovers host genes required for coxsackievirus infection. Genome Res.
2018;28(6):859-68. Epub 20180430. doi: 10.1101/gr.230250.117. PubMed PMID: 29712754;
PubMed Central PMCID: PMCPMC5991512.

20.   Kodigepalli KM, Bowers K, Sharp A, Nanjundan M. Roles and regulation of phospholipid
scramblases. FEBS Lett. 2015;589(1):3-14. Epub 20141203. doi: 10.1016/j.febslet.2014.11.036.
PubMed PMID: 25479087.

21.   Zhou Q, Zhao J, Stout JG, Luhm RA, Wiedmer T, Sims PJ. Molecular cloning of human
plasma membrane phospholipid scramblase. A protein mediating transbilayer movement of
plasma membrane phospholipids. J Biol Chem. 1997;272(29):18240-4. doi:
10.1074/jbc.272.29.18240. PubMed PMID: 9218461.

22. Dong B, Zhou Q, Zhao J, Zhou A, Harty RN, Bose S, et al. Phospholipid scramblase 1
potentiates the antiviral activity of interferon. J Virol. 2004;78(17):8983-93. doi:
10.1128/JVI.78.17.8983-8993.2004. PubMed PMID: 15308695; PubMed Central PMCID:
PMCPMC506946.
23. Pairo-Castineira E, Rawlik K, Bretherick AD, Qi T, Wu Y, Nassiri I, et al. GWAS and meta-
analysis identifies 49 genetic variants underlying critical COVID-19. Nature. 2023;617(7962):764-
8. Epub 20230517. doi: 10.1038/s41586-023-06034-3. PubMed PMID: 37198478; PubMed
Central PMCID: PMCPMC10208981.
24. Kousathanas A, Pairo-Castineira E, Rawlik K, Stuckey A, Odhams CA, Walker S, et al.
Whole-genome sequencing reveals host factors underlying critical COVID-19. Nature.
2022;607(7917):97-103. Epub 20220307. doi: 10.1038/s41586-022-04576-6. PubMed PMID:
35255492; PubMed Central PMCID: PMCPMC9259496.
25. Baggen J, Vanstreels E, Jansen S, Daelemans D. Cellular host factors for SARS-CoV-2
infection. Nat Microbiol. 2021;6(10):1219-32. Epub 20210901. doi: 10.1038/s41564-021-00958-
0. PubMed PMID: 34471255.
26. Hoffmann HH, Sanchez-Rivera FJ, Schneider WM, Luna JM, Soto-Feliciano YM, Ashbrook
AW, et al. Functional interrogation of a SARS-CoV-2 host protein interactome identifies unique
and shared coronavirus host factors. Cell Host Microbe. 2021;29(2):267-80 e5. Epub 20201216.
doi: 10.1016/j.chom.2020.12.009. PubMed PMID: 33357464; PubMed Central PMCID:
PMCPMC7833927.
27. Li Q, Wu J, Nie J, Zhang L, Hao H, Liu S, et al. The Impact of Mutations in SARS-CoV-2
Spike on Viral Infectivity and Antigenicity. Cell. 2020;182(5):1284-94 e9. Epub 20200717. doi:
10.1016/j.cell.2020.07.012. PubMed PMID: 32730807; PubMed Central PMCID:
PMCPMC7366990.
28. Riva L, Yuan S, Yin X, Martin-Sancho L, Matsunaga N, Pache L, et al. Discovery of SARS-
CoV-2 antiviral drugs through large-scale compound repurposing. Nature. 2020;586(7827):113-
9. Epub 20200724. doi: 10.1038/s41586-020-2577-1. PubMed PMID: 32707573; PubMed
Central PMCID: PMCPMC7603405.
29. Schneider WM, Luna JM, Hoffmann HH, Sanchez-Rivera FJ, Leal AA, Ashbrook AW, et al.
Genome-Scale Identification of SARS-CoV-2 and Pan-coronavirus Host Factor Networks. Cell.
2021;184(1):120-32 e14. Epub 20201209. doi: 10.1016/j.cell.2020.12.006. PubMed PMID:
33382968; PubMed Central PMCID: PMCPMC7796900.
30. Xia S, Lan Q, Su S, Wang X, Xu W, Liu Z, et al. The role of furin cleavage site in SARS-CoV-2
spike protein-mediated membrane fusion in the presence or absence of trypsin. Signal
Transduct Target Ther. 2020;5(1):92. Epub 20200612. doi: 10.1038/s41392-020-0184-0. PubMed
PMID: 32532959; PubMed Central PMCID: PMCPMC7289711.
31. Xia S, Liu M, Wang C, Xu W, Lan Q, Feng S, et al. Inhibition of SARS-CoV-2 (previously
2019-nCoV) infection by a highly potent pan-coronavirus fusion inhibitor targeting its spike
protein that harbors a high capacity to mediate membrane fusion. Cell Res. 2020;30(4):343-55.
Epub 20200330. doi: 10.1038/s41422-020-0305-x. PubMed PMID: 32231345; PubMed Central
PMCID: PMCPMC7104723.
32. Yan Y, Chang L, Wang L. Laboratory testing of SARS-CoV, MERS-CoV, and SARS-CoV-2
(2019-nCoV): Current status, challenges, and countermeasures. Rev Med Virol.

2020;30(3):e2106. Epub 20200417. doi: 10.1002/rmv.2106. PubMed PMID: 32302058; PubMed
Central PMCID: PMCPMC7235496.

33.   Barretina J, Caponigro G, Stransky N, Venkatesan K, Margolin AA, Kim S, et al. The Cancer
Cell Line Encyclopedia enables predictive modelling of anticancer drug sensitivity. Nature.
2012;483(7391):603-7. Epub 20120328. doi: 10.1038/nature11003. PubMed PMID: 22460905;
PubMed Central PMCID: PMCPMC3320027.

34.   Klijn C, Durinck S, Stawiski EW, Haverty PM, Jiang Z, Liu H, et al. A comprehensive
transcriptional portrait of human cancer cell lines. Nat Biotechnol. 2015;33(3):306-12. Epub
20141208. doi: 10.1038/nbt.3080. PubMed PMID: 25485619.

35.   Consortium GT. Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis:
multitissue gene regulation in humans. Science. 2015;348(6235):648-60. Epub 20150507. doi:
10.1126/science.1262110. PubMed PMID: 25954001; PubMed Central PMCID:
PMCPMC4547484.

36.   de Weerd NA, Samarajiwa SA, Hertzog PJ. Type I interferon receptors: biochemistry and
biological functions. J Biol Chem. 2007;282(28):20053-7. Epub 20070514. doi:
10.1074/jbc.R700006200. PubMed PMID: 17502368.

37.   de Weerd NA, Nguyen T. The interferons and their receptors--distribution and regulation.
Immunol Cell Biol. 2012;90(5):483-91. Epub 20120313. doi: 10.1038/icb.2012.9. PubMed PMID:
22410872; PubMed Central PMCID: PMCPMC7165917.

38.   Uze G, Schreiber G, Piehler J, Pellegrini S. The receptor of the type I interferon family.
Curr Top Microbiol Immunol. 2007;316:71-95. doi: 10.1007/978-3-540-71329-6_5. PubMed
PMID: 17969444.

39.   Horvath CM, Stark GR, Kerr IM, Darnell JE, Jr. Interactions between STAT and non-STAT
proteins in the interferon-stimulated gene factor 3 transcription complex. Mol Cell Biol.
1996;16(12):6957-64. doi: 10.1128/MCB.16.12.6957. PubMed PMID: 8943351; PubMed Central
PMCID: PMCPMC231699.

40.   Veals SA, Schindler C, Leonard D, Fu XY, Aebersold R, Darnell JE, Jr., et al. Subunit of an
alpha-interferon-responsive transcription factor is related to interferon regulatory factor and
Myb families of DNA-binding proteins. Mol Cell Biol. 1992;12(8):3315-24. doi:
10.1128/mcb.12.8.3315-3324.1992. PubMed PMID: 1630447; PubMed Central PMCID:
PMCPMC364572.

41.   Darnell JE, Jr., Kerr IM, Stark GR. Jak-STAT pathways and transcriptional activation in
response to IFNs and other extracellular signaling proteins. Science. 1994;264(5164):1415-21.
doi: 10.1126/science.8197455. PubMed PMID: 8197455.

42.   Schindler C, Strehlow I. Cytokines and STAT signaling. Adv Pharmacol. 2000;47:113-74.
doi: 10.1016/s1054-3589(08)60111-8. PubMed PMID: 10582086.

43.   Levy DE, Darnell JE, Jr. Stats: transcriptional control and biological impact. Nat Rev Mol
Cell Biol. 2002;3(9):651-62. doi: 10.1038/nrm909. PubMed PMID: 12209125.

44.   Perng YC, Lenschow DJ. ISG15 in antiviral immunity and beyond. Nat Rev Microbiol.
2018;16(7):423-39. doi: 10.1038/s41579-018-0020-5. PubMed PMID: 29769653; PubMed
Central PMCID: PMCPMC7097117.

45.   Speer SD, Li Z, Buta S, Payelle-Brogard B, Qian L, Vigant F, et al. ISG15 deficiency and
increased viral resistance in humans but not mice. Nat Commun. 2016;7:11496. Epub

20160519. doi: 10.1038/ncomms11496. PubMed PMID: 27193971; PubMed Central PMCID:
PMCPMC4873964.
46. Zhang X, Bogunovic D, Payelle-Brogard B, Francois-Newton V, Speer SD, Yuan C, et al.
Human intracellular ISG15 prevents interferon-alpha/beta over-amplification and auto-
inflammation. Nature. 2015;517(7532):89-93. Epub 20141012. doi: 10.1038/nature13801.
PubMed PMID: 25307056; PubMed Central PMCID: PMCPMC4303590.
47. Naka T, Narazaki M, Hirata M, Matsumoto T, Minamoto S, Aono A, et al. Structure and
function of a new STAT-induced STAT inhibitor. Nature. 1997;387(6636):924-9. doi:
10.1038/43219. PubMed PMID: 9202127.
48. Malakhova OA, Kim KI, Luo JK, Zou W, Kumar KG, Fuchs SY, et al. UBP43 is a novel
regulator of interferon signaling independent of its ISG15 isopeptidase activity. EMBO J.
2006;25(11):2358-67. Epub 20060518. doi: 10.1038/sj.emboj.7601149. PubMed PMID:
16710296; PubMed Central PMCID: PMCPMC1478183.
49. Francois-Newton V, Magno de Freitas Almeida G, Payelle-Brogard B, Monneron D,
Pichard-Garcia L, Piehler J, et al. USP18-based negative feedback control is induced by type I and
type III interferons and specifically inactivates interferon alpha response. PLoS One.
2011;6(7):e22200. Epub 20110714. doi: 10.1371/journal.pone.0022200. PubMed PMID:
21779393; PubMed Central PMCID: PMCPMC3136508.
50. Ziegler CGK, Allon SJ, Nyquist SK, Mbano IM, Miao VN, Tzouanas CN, et al. SARS-CoV-2
Receptor ACE2 Is an Interferon-Stimulated Gene in Human Airway Epithelial Cells and Is
Detected in Specific Cell Subsets across Tissues. Cell. 2020;181(5):1016-35 e19. Epub 20200427.
doi: 10.1016/j.cell.2020.04.035. PubMed PMID: 32413319; PubMed Central PMCID:
PMCPMC7252096.
51. Onabajo OO, Banday AR, Stanifer ML, Yan W, Obajemu A, Santer DM, et al. Interferons
and viruses induce a novel truncated ACE2 isoform and not the full-length SARS-CoV-2 receptor.
Nat Genet. 2020;52(12):1283-93. Epub 20201019. doi: 10.1038/s41588-020-00731-9. PubMed
PMID: 33077916; PubMed Central PMCID: PMCPMC9377523.
52. Zhao MM, Yang WL, Yang FY, Zhang L, Huang WJ, Hou W, et al. Cathepsin L plays a key
role in SARS-CoV-2 infection in humans and humanized mice and is a promising target for new
drug development. Signal Transduct Target Ther. 2021;6(1):134. Epub 20210327. doi:
10.1038/s41392-021-00558-8. PubMed PMID: 33774649; PubMed Central PMCID:
PMCPMC7997800.
53. Shang J, Wan Y, Luo C, Ye G, Geng Q, Auerbach A, et al. Cell entry mechanisms of SARS-
CoV-2. Proc Natl Acad Sci U S A. 2020;117(21):11727-34. Epub 20200506. doi:
10.1073/pnas.2003138117. PubMed PMID: 32376634; PubMed Central PMCID:
PMCPMC7260975.
54. Liu T, Luo S, Libby P, Shi GP. Cathepsin L-selective inhibitors: A potentially promising
treatment for COVID-19 patients. Pharmacol Ther. 2020;213:107587. Epub 20200526. doi:
10.1016/j.pharmthera.2020.107587. PubMed PMID: 32470470; PubMed Central PMCID:
PMCPMC7255230.
55. Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, et al. SARS-
CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease
Inhibitor. Cell. 2020;181(2):271-80 e8. Epub 20200305. doi: 10.1016/j.cell.2020.02.052. PubMed
PMID: 32142651; PubMed Central PMCID: PMCPMC7102627.

56. Jackson CB, Farzan M, Chen B, Choe H. Mechanisms of SARS-CoV-2 entry into cells. Nat
Rev Mol Cell Biol. 2022;23(1):3-20. Epub 20211005. doi: 10.1038/s41580-021-00418-x. PubMed
PMID: 34611326; PubMed Central PMCID: PMCPMC8491763.
57. Pellegrini S, John J, Shearer M, Kerr IM, Stark GR. Use of a selectable marker regulated
by alpha interferon to obtain mutations in the signaling pathway. Mol Cell Biol.
1989;9(11):4605-12. doi: 10.1128/mcb.9.11.4605-4612.1989. PubMed PMID: 2513475;
PubMed Central PMCID: PMCPMC363606.
58. Velazquez L, Fellous M, Stark GR, Pellegrini S. A protein tyrosine kinase in the interferon
alpha/beta signaling pathway. Cell. 1992;70(2):313-22. doi: 10.1016/0092-8674(92)90105-l.
PubMed PMID: 1386289.
59. Wang S, Li W, Hui H, Tiwari SK, Zhang Q, Croker BA, et al. Cholesterol 25-Hydroxylase
inhibits SARS-CoV-2 and other coronaviruses by depleting membrane cholesterol. EMBO J.
2020;39(21):e106057. Epub 20201005. doi: 10.15252/embj.2020106057. PubMed PMID:
32944968; PubMed Central PMCID: PMCPMC7537045.
60. Zhu Y, Feng F, Hu G, Wang Y, Yu Y, Zhu Y, et al. A genome-wide CRISPR screen identifies
host factors that regulate SARS-CoV-2 entry. Nat Commun. 2021;12(1):961. Epub 20210211. doi:
10.1038/s41467-021-21213-4. PubMed PMID: 33574281; PubMed Central PMCID:
PMCPMC7878750.
61. Wei LH, Sun Y, Guo JU. Genome-wide CRISPR screens identify noncanonical translation
factor eIF2A as an enhancer of SARS-CoV-2 programmed -1 ribosomal frameshifting. Cell Rep.
2023;42(8):112987. Epub 20230814. doi: 10.1016/j.celrep.2023.112987. PubMed PMID:
37581984.
62. Wei J, Alfajaro MM, DeWeirdt PC, Hanna RE, Lu-Culligan WJ, Cai WL, et al. Genome-wide
CRISPR Screens Reveal Host Factors Critical for SARS-CoV-2 Infection. Cell. 2021;184(1):76-91
e13. Epub 20201020. doi: 10.1016/j.cell.2020.10.028. PubMed PMID: 33147444; PubMed
Central PMCID: PMCPMC7574718.
63. Wang R, Simoneau CR, Kulsuptrakul J, Bouhaddou M, Travisano KA, Hayashi JM, et al.
Genetic Screens Identify Host Factors for SARS-CoV-2 and Common Cold Coronaviruses. Cell.
2021;184(1):106-19 e14. Epub 20201209. doi: 10.1016/j.cell.2020.12.004. PubMed PMID:
33333024; PubMed Central PMCID: PMCPMC7723770.
64. Song J, Chow RD, Pena-Hernandez MA, Zhang L, Loeb SA, So EY, et al. LRRC15 inhibits
SARS-CoV-2 cellular entry in trans. PLoS Biol. 2022;20(10):e3001805. Epub 20221013. doi:
10.1371/journal.pbio.3001805. PubMed PMID: 36228039; PubMed Central PMCID:
PMCPMC9595563.
65. Hossain MS, Kerkvliet JG, Hoppe AD. Whole genome CRISPR screening strategy to
identify genes contributing to SARS-CoV-2 spike and VSV-G mediated entry. J Med Virol.
2023;95(9):e29087. doi: 10.1002/jmv.29087. PubMed PMID: 37707319.
66. Rehfeld F, Eitson JL, Ohlson MB, Chang TC, Schoggins JW, Mendell JT. CRISPR screening
reveals a dependency on ribosome recycling for efficient SARS-CoV-2 programmed ribosomal
frameshifting and viral replication. Cell Rep. 2023;42(2):112076. Epub 20230130. doi:
10.1016/j.celrep.2023.112076. PubMed PMID: 36753415; PubMed Central PMCID:
PMCPMC9884621.
67. Rebendenne A, Roy P, Bonaventure B, Chaves VAL, Desmarets L, Rouille Y, et al.
Bidirectional genome-wide CRISPR screens reveal host factors regulating SARS-CoV-2, MERS-CoV

and seasonal HCoVs. Res Sq. 2021. Epub 20210527. doi: 10.21203/rs.3.rs-555275/v1. PubMed
PMID: 34075371; PubMed Central PMCID: PMCPMC8168385.

68.   Pahmeier F, Lavacca TM, Goellner S, Neufeldt CJ, Prasad V, Cerikan B, et al. Identification
of host dependency factors involved in SARS-CoV-2 replication organelle formation through
proteomics and ultrastructural analysis. J Virol. 2023:e0087823. Epub 20231031. doi:
10.1128/jvi.00878-23. PubMed PMID: 37905840.

69.   Loo L, Waller MA, Moreno CL, Cole AJ, Stella AO, Pop OT, et al. Fibroblast-expressed
LRRC15 is a receptor for SARS-CoV-2 spike and controls antiviral and antifibrotic transcriptional
programs. PLoS Biol. 2023;21(2):e3001967. Epub 20230209. doi: 10.1371/journal.pbio.3001967.
PubMed PMID: 36757924; PubMed Central PMCID: PMCPMC9910744.

70.   Israeli M, Finkel Y, Yahalom-Ronen Y, Paran N, Chitlaru T, Israeli O, et al. Genome-wide
CRISPR screens identify GATA6 as a proviral host factor for SARS-CoV-2 via modulation of ACE2.
Nat Commun. 2022;13(1):2237. Epub 20220425. doi: 10.1038/s41467-022-29896-z. PubMed
PMID: 35469023; PubMed Central PMCID: PMCPMC9039069.

71.   Grodzki M, Bluhm AP, Schaefer M, Tagmount A, Russo M, Sobh A, et al. Genome-scale
CRISPR screens identify host factors that promote human coronavirus infection. Genome Med.
2022;14(1):10. Epub 20220127. doi: 10.1186/s13073-022-01013-1. PubMed PMID: 35086559;
PubMed Central PMCID: PMCPMC8792531.

72.   Gordon DE, Hiatt J, Bouhaddou M, Rezelj VV, Ulferts S, Braberg H, et al. Comparative
host-coronavirus protein interaction networks reveal pan-viral disease mechanisms. Science.
2020;370(6521). Epub 20201015. doi: 10.1126/science.abe9403. PubMed PMID: 33060197;
PubMed Central PMCID: PMCPMC7808408.

73.   Daniloski Z, Jordan TX, Wessels HH, Hoagland DA, Kasela S, Legut M, et al. Identification
of Required Host Factors for SARS-CoV-2 Infection in Human Cells. Cell. 2021;184(1):92-105 e16.
Epub 20201024. doi: 10.1016/j.cell.2020.10.030. PubMed PMID: 33147445; PubMed Central
PMCID: PMCPMC7584921.

74.   Chan K, Farias AG, Lee H, Guvenc F, Mero P, Brown KR, et al. Survival-based CRISPR
genetic screens across a panel of permissive cell lines identify common and cell-specific SARS-
CoV-2 host factors. Heliyon. 2023;9(1):e12744. Epub 20221230. doi:
10.1016/j.heliyon.2022.e12744. PubMed PMID: 36597481; PubMed Central PMCID:
PMCPMC9800021.

75.   Biering SB, Sarnik SA, Wang E, Zengel JR, Leist SR, Schafer A, et al. Genome-wide
bidirectional CRISPR screens identify mucins as host factors modulating SARS-CoV-2 infection.
Nat Genet. 2022;54(8):1078-89. Epub 20220725. doi: 10.1038/s41588-022-01131-x. PubMed
PMID: 35879412; PubMed Central PMCID: PMCPMC9355872.

76.   Baggen J, Persoons L, Vanstreels E, Jansen S, Van Looveren D, Boeckx B, et al. Genome-
wide CRISPR screening identifies TMEM106B as a proviral host factor for SARS-CoV-2. Nat
Genet. 2021;53(4):435-44. Epub 20210308. doi: 10.1038/s41588-021-00805-2. PubMed PMID:
33686287.

77.   van der Made CI, Simons A, Schuurs-Hoeijmakers J, van den Heuvel G, Mantere T,
Kersten S, et al. Presence of Genetic Variants Among Young Men With Severe COVID-19. JAMA.
2020;324(7):663-73. doi: 10.1001/jama.2020.13719. PubMed PMID: 32706371; PubMed
Central PMCID: PMCPMC7382021.

78.   Shelton JF, Shastri AJ, Ye C, Weldon CH, Filshtein-Sonmez T, Coker D, et al. Trans-ancestry
analysis reveals genetic and nongenetic associations with COVID-19 susceptibility and severity.
Nat Genet. 2021;53(6):801-8. Epub 20210422. doi: 10.1038/s41588-021-00854-7. PubMed
PMID: 33888907.
79.   Roberts GHL, Partha R, Rhead B, Knight SC, Park DS, Coignet MV, et al. Expanded COVID-
19 phenotype definitions reveal distinct patterns of genetic association and protective effects.
Nat Genet. 2022;54(4):374-81. Epub 20220411. doi: 10.1038/s41588-022-01042-x. PubMed
PMID: 35410379.
80.   Pietzner M, Chua RL, Wheeler E, Jechow K, Willett JDS, Radbruch H, et al. ELF5 is a
potential respiratory epithelial cell-specific risk gene for severe COVID-19. Nat Commun.
2022;13(1):4484. Epub 20220815. doi: 10.1038/s41467-022-31999-6. PubMed PMID:
35970849; PubMed Central PMCID: PMCPMC9378714.
81.   Latini A, Agolini E, Novelli A, Borgiani P, Giannini R, Gravina P, et al. COVID-19 and
Genetic Variants of Protein Involved in the SARS-CoV-2 Entry into the Host Cells. Genes (Basel).
2020;11(9). Epub 20200827. doi: 10.3390/genes11091010. PubMed PMID: 32867305; PubMed
Central PMCID: PMCPMC7565048.
82.   Horowitz JE, Kosmicki JA, Damask A, Sharma D, Roberts GHL, Justice AE, et al. Genome-
wide analysis provides genetic evidence that ACE2 influences COVID-19 risk and yields risk
scores associated with severe disease. Nat Genet. 2022;54(4):382-92. Epub 20220303. doi:
10.1038/s41588-021-01006-7. PubMed PMID: 35241825; PubMed Central PMCID:
PMCPMC9005345.
83.   Garcia-Garcia A, Perez de Diego R, Flores C, Rinchai D, Sole-Violan J, Deya-Martinez A, et
al. Humans with inherited MyD88 and IRAK-4 deficiencies are predisposed to hypoxemic COVID-
19 pneumonia. J Exp Med. 2023;220(5). Epub 20230303. doi: 10.1084/jem.20220170. PubMed
PMID: 36880831; PubMed Central PMCID: PMCPMC9998661.
84.   Fallerini C, Daga S, Mantovani S, Benetti E, Picchiotti N, Francisci D, et al. Association of
Toll-like receptor 7 variants with life-threatening COVID-19 disease in males: findings from a
nested case-control study. Elife. 2021;10. Epub 20210302. doi: 10.7554/eLife.67569. PubMed
PMID: 33650967; PubMed Central PMCID: PMCPMC7987337.
85.   Severe Covid GG, Ellinghaus D, Degenhardt F, Bujanda L, Buti M, Albillos A, et al.
Genomewide Association Study of Severe Covid-19 with Respiratory Failure. N Engl J Med.
2020;383(16):1522-34. Epub 20200617. doi: 10.1056/NEJMoa2020283. PubMed PMID:
32558485; PubMed Central PMCID: PMCPMC7315890.
86.   Degenhardt F, Ellinghaus D, Juzenas S, Lerga-Jaso J, Wendorff M, Maya-Miles D, et al.
Detailed stratified GWAS analysis for severe COVID-19 in four European populations. Hum Mol
Genet. 2022;31(23):3945-66. doi: 10.1093/hmg/ddac158. PubMed PMID: 35848942; PubMed
Central PMCID: PMCPMC9703941.
87.   D'Antonio M, Nguyen JP, Arthur TD, Matsui H, Initiative C-HG, D'Antonio-Chronowska A,
et al. SARS-CoV-2 susceptibility and COVID-19 disease severity are associated with genetic
variants affecting gene expression in a variety of tissues. Cell Rep. 2021;37(7):110020. Epub
20211103. doi: 10.1016/j.celrep.2021.110020. PubMed PMID: 34762851; PubMed Central
PMCID: PMCPMC8563343.

88. Cruz R, Diz-de Almeida S, Lopez de Heredia M, Quintela I, Ceballos FC, Pita G, et al. Novel
genes and sex differences in COVID-19 severity. Hum Mol Genet. 2022;31(22):3789-806. doi:
10.1093/hmg/ddac132. PubMed PMID: 35708486; PubMed Central PMCID: PMCPMC9652109.
89. Initiative C-HG. A first update on mapping the human genetic architecture of COVID-19.
Nature. 2022;608(7921):E1-E10. Epub 20220803. doi: 10.1038/s41586-022-04826-7. PubMed
PMID: 35922517; PubMed Central PMCID: PMCPMC9352569.
90. Andolfo I, Russo R, Lasorsa VA, Cantalupo S, Rosato BE, Bonfiglio F, et al. Common
variants at 21q22.3 locus influence MX1 and TMPRSS2 gene expression and susceptibility to
severe COVID-19. iScience. 2021;24(4):102322. Epub 20210317. doi:
10.1016/j.isci.2021.102322. PubMed PMID: 33748697; PubMed Central PMCID:
PMCPMC7968217.
91. Zhou S, Butler-Laporte G, Nakanishi T, Morrison DR, Afilalo J, Afilalo M, et al. A
Neanderthal OAS1 isoform protects individuals of European ancestry against COVID-19
susceptibility and severity. Nat Med. 2021;27(4):659-67. Epub 20210225. doi: 10.1038/s41591-
021-01281-1. PubMed PMID: 33633408.
92. Huffman JE, Butler-Laporte G, Khan A, Pairo-Castineira E, Drivas TG, Peloso GM, et al.
Multi-ancestry fine mapping implicates OAS1 splicing in risk of severe COVID-19. Nat Genet.
2022;54(2):125-7. Epub 20220113. doi: 10.1038/s41588-021-00996-8. PubMed PMID:
35027740; PubMed Central PMCID: PMCPMC8837537.
93. Initiative C-HG. Mapping the human genetic architecture of COVID-19. Nature.
2021;600(7889):472-7. Epub 20210708. doi: 10.1038/s41586-021-03767-x. PubMed PMID:
34237774; PubMed Central PMCID: PMCPMC8674144.
94. Pairo-Castineira E, Clohisey S, Klaric L, Bretherick AD, Rawlik K, Pasko D, et al. Genetic
mechanisms of critical illness in COVID-19. Nature. 2021;591(7848):92-8. Epub 20201211. doi:
10.1038/s41586-020-03065-y. PubMed PMID: 33307546.
95. Banday AR, Stanifer ML, Florez-Vargas O, Onabajo OO, Papenberg BW, Zahoor MA, et al.
Genetic regulation of OAS1 nonsense-mediated decay underlies association with COVID-19
hospitalization in patients of European and African ancestries. Nat Genet. 2022;54(8):1103-16.
Epub 20220714. doi: 10.1038/s41588-022-01113-z. PubMed PMID: 35835913; PubMed Central
PMCID: PMCPMC9355882.
96. Payman S-T, Hala A, James DRK, Audrey A, Reuben S, Zhen-Yuan L, et al. A SARS-CoV-2 –
host proximity interactome. bioRxiv. 2020:2020.09.03.282103. doi:
10.1101/2020.09.03.282103.
97. Gordon DE, Jang GM, Bouhaddou M, Xu J, Obernier K, White KM, et al. A SARS-CoV-2
protein interaction map reveals targets for drug repurposing. Nature. 2020;583(7816):459-68.
Epub 20200430. doi: 10.1038/s41586-020-2286-9. PubMed PMID: 32353859; PubMed Central
PMCID: PMCPMC7431030.
98. Estelle MNL, Yorgos S, Anastassia K, Jean-Pascal G, Payman Samavarchi T, Dae-Kyum K, et
al. Global BioID-based SARS-CoV-2 proteins proximal interactome unveils novel ties between
viral polypeptides and host factors involved in multiple COVID19-associated mechanisms.
bioRxiv. 2020:2020.08.28.272955. doi: 10.1101/2020.08.28.272955.
99. St-Germain JR, Astori A, Samavarchi-Tehrani P, Abdouni H, Macwan V, Kim D-K, et al. A
SARS-CoV-2 BioID-based virus-host membrane protein interactome and virus peptide

compendium: new proteomics resources for COVID-19 research. bioRxiv.
2020:2020.08.28.269175. doi: 10.1101/2020.08.28.269175.

100.   Liu X, Huuskonen S, Laitinen T, Redchuk T, Bogacheva M, Salokas K, et al. SARS-CoV-2-
host proteome interactions for antiviral drug discovery. Mol Syst Biol. 2021;17(11):e10396. doi:
10.15252/msb.202110396. PubMed PMID: 34709727; PubMed Central PMCID:
PMCPMC8552907.

101.   May DG, Martin-Sancho L, Anschau V, Liu S, Chrisopulos RJ, Scott KL, et al. A BioID-
Derived Proximity Interactome for SARS-CoV-2 Proteins. Viruses. 2022;14(3). Epub 20220315.
doi: 10.3390/v14030611. PubMed PMID: 35337019; PubMed Central PMCID:
PMCPMC8951556.

102.   Zhou Y, Liu Y, Gupta S, Paramo MI, Hou Y, Mao C, et al. A comprehensive SARS-CoV-2-
human protein-protein interactome reveals COVID-19 pathobiology and potential host
therapeutic targets. Nat Biotechnol. 2023;41(1):128-39. Epub 20221010. doi: 10.1038/s41587-
022-01474-0. PubMed PMID: 36217030; PubMed Central PMCID: PMCPMC9851973.

103.   Stukalov A, Girault V, Grass V, Karayel O, Bergant V, Urban C, et al. Multilevel proteomics
reveals host perturbations by SARS-CoV-2 and SARS-CoV. Nature. 2021;594(7862):246-52. Epub
20210412. doi: 10.1038/s41586-021-03493-4. PubMed PMID: 33845483.

104.   Li J, Guo M, Tian X, Wang X, Yang X, Wu P, et al. Virus-Host Interactome and Proteomic
Survey Reveal Potential Virulence Factors Influencing SARS-CoV-2 Pathogenesis. Med.
2021;2(1):99-112 e7. Epub 20200721. doi: 10.1016/j.medj.2020.07.002. PubMed PMID:
32838362; PubMed Central PMCID: PMCPMC7373048.

105.   Davies JP, Almasy KM, McDonald EF, Plate L. Comparative Multiplexed Interactomics of
SARS-CoV-2 and Homologous Coronavirus Nonstructural Proteins Identifies Unique and Shared
Host-Cell Dependencies. ACS Infect Dis. 2020;6(12):3174-89. Epub 20201202. doi:
10.1021/acsinfecdis.0c00500. PubMed PMID: 33263384; PubMed Central PMCID:
PMCPMC7724760.

106.   Schmidt N, Ganskih S, Wei Y, Gabel A, Zielinski S, Keshishian H, et al. SND1 binds SARS-
CoV-2 negative-sense RNA and promotes viral RNA synthesis through NSP9. Cell.
2023;186(22):4834-50 e23. Epub 20231003. doi: 10.1016/j.cell.2023.09.002. PubMed PMID:
37794589; PubMed Central PMCID: PMCPMC10617981.

107.   Schmidt N, Lareau CA, Keshishian H, Ganskih S, Schneider C, Hennig T, et al. The SARS-
CoV-2 RNA-protein interactome in infected human cells. Nat Microbiol. 2021;6(3):339-53. Epub
20201221. doi: 10.1038/s41564-020-00846-z. PubMed PMID: 33349665; PubMed Central
PMCID: PMCPMC7906908.

108.   Labeau A, Fery-Simonian L, Lefevre-Utile A, Pourcelot M, Bonnet-Madin L, Soumelis V, et
al. Characterization and functional interrogation of the SARS-CoV-2 RNA interactome. Cell Rep.
2022;39(4):110744. doi: 10.1016/j.celrep.2022.110744. PubMed PMID: 35477000; PubMed
Central PMCID: PMCPMC9040432.

109.   Kamel W, Noerenberg M, Cerikan B, Chen H, Jarvelin AI, Kammoun M, et al. Global
analysis of protein-RNA interactions in SARS-CoV-2-infected cells reveals key regulators of
infection. Mol Cell. 2021;81(13):2851-67 e7. Epub 20210524. doi:
10.1016/j.molcel.2021.05.023. PubMed PMID: 34118193; PubMed Central PMCID:
PMCPMC8142890.

110.   Flynn RA, Belk JA, Qi Y, Yasumoto Y, Wei J, Alfajaro MM, et al. Discovery and functional
interrogation of SARS-CoV-2 RNA-host protein interactions. Cell. 2021;184(9):2394-411 e16.
Epub 20210311. doi: 10.1016/j.cell.2021.03.012. PubMed PMID: 33743211; PubMed Central
PMCID: PMCPMC7951565.

111.   Lee S, Lee YS, Choi Y, Son A, Park Y, Lee KM, et al. The SARS-CoV-2 RNA interactome. Mol
Cell. 2021;81(13):2838-50 e6. Epub 20210427. doi: 10.1016/j.molcel.2021.04.022. PubMed
PMID: 33989516; PubMed Central PMCID: PMCPMC8075806.

112.   Bouhaddou M, Memon D, Meyer B, White KM, Rezelj VV, Correa Marrero M, et al. The
Global Phosphorylation Landscape of SARS-CoV-2 Infection. Cell. 2020;182(3):685-712 e19.
Epub 20200628. doi: 10.1016/j.cell.2020.06.034. PubMed PMID: 32645325; PubMed Central
PMCID: PMCPMC7321036.

113.   Sanchez-Magraner L, Posada IM, Andraka N, Contreras FX, Viguera AR, Guerin DM, et al.
The C-terminal transmembrane domain of human phospholipid scramblase 1 is essential for the
protein flip-flop activity and Ca(2)(+)-binding. J Membr Biol. 2014;247(2):155-65. Epub
20131217. doi: 10.1007/s00232-013-9619-7. PubMed PMID: 24343571.

114.   Ricardo-Lax I, Luna JM, Thao TTN, Le Pen J, Yu Y, Hoffmann HH, et al. Replication and
single-cycle delivery of SARS-CoV-2 replicons. Science. 2021;374(6571):1099-106. Epub
20211014. doi: 10.1126/science.abj8430. PubMed PMID: 34648371; PubMed Central PMCID:
PMCPMC9007107.

115.   Schmidt F, Weisblum Y, Muecksch F, Hoffmann HH, Michailidis E, Lorenzi JCC, et al.
Measuring SARS-CoV-2 neutralizing antibody activity using pseudotyped and chimeric viruses. J
Exp Med. 2020;217(11). doi: 10.1084/jem.20201181. PubMed PMID: 32692348; PubMed
Central PMCID: PMCPMC7372514.

116.   Khan H, Winstone H, Jimenez-Guardeno JM, Graham C, Doores KJ, Goujon C, et al.
TMPRSS2 promotes SARS-CoV-2 evasion from NCOA7-mediated restriction. PLoS Pathog.
2021;17(11):e1009820. Epub 20211122. doi: 10.1371/journal.ppat.1009820. PubMed PMID:
34807954; PubMed Central PMCID: PMCPMC8648102.

117.   Thorne LG, Bouhaddou M, Reuschl AK, Zuliani-Alvarez L, Polacco B, Pelin A, et al.
Evolution of enhanced innate immune evasion by SARS-CoV-2. Nature. 2022;602(7897):487-95.
Epub 20211223. doi: 10.1038/s41586-021-04352-y. PubMed PMID: 34942634; PubMed Central
PMCID: PMCPMC8850198.

118.   Markov PV, Ghafari M, Beer M, Lythgoe K, Simmonds P, Stilianakis NI, et al. The evolution
of SARS-CoV-2. Nature Reviews Microbiology. 2023;21(6):361-79. doi: 10.1038/s41579-023-
00878-2.

119.   Carabelli AM, Peacock TP, Thorne LG, Harvey WT, Hughes J, de Silva TI, et al. SARS-CoV-2
variant biology: immune escape, transmission and fitness. Nature Reviews Microbiology.
2023;21(3):162-77. doi: 10.1038/s41579-022-00841-7.

120.   Huang P, Liao R, Chen X, Wu X, Li X, Wang Y, et al. Nuclear translocation of PLSCR1
activates STAT1 signaling in basal-like breast cancer. Theranostics. 2020;10(10):4644-58. Epub
20200325. doi: 10.7150/thno.43150. PubMed PMID: 32292520; PubMed Central PMCID:
PMCPMC7150476.

121.   Nanjundan M, Sun J, Zhao J, Zhou Q, Sims PJ, Wiedmer T. Plasma membrane
phospholipid scramblase 1 promotes EGF-dependent activation of c-Src through the epidermal

growth factor receptor. J Biol Chem. 2003;278(39):37413-8. Epub 20030718. doi:
10.1074/jbc.M306182200. PubMed PMID: 12871937.
122. Ben-Efraim I, Zhou Q, Wiedmer T, Gerace L, Sims PJ. Phospholipid scramblase 1 is
imported into the nucleus by a receptor-mediated pathway and interacts with DNA.
Biochemistry. 2004;43(12):3518-26. doi: 10.1021/bi0356911. PubMed PMID: 15035622.
123. Chen MH, Ben-Efraim I, Mitrousis G, Walker-Kopp N, Sims PJ, Cingolani G. Phospholipid
scramblase 1 contains a nonclassical nuclear localization signal with unique binding site in
importin alpha. J Biol Chem. 2005;280(11):10599-606. Epub 20041217. doi:
10.1074/jbc.M413194200. PubMed PMID: 15611084.
124. Wiedmer T, Zhao J, Nanjundan M, Sims PJ. Palmitoylation of phospholipid scramblase 1
controls its distribution between nucleus and plasma membrane. Biochemistry.
2003;42(5):1227-33. doi: 10.1021/bi026679w. PubMed PMID: 12564925.
125. Zhao J, Zhou Q, Wiedmer T, Sims PJ. Palmitoylation of phospholipid scramblase is
required for normal function in promoting Ca2+-activated transbilayer movement of membrane
phospholipids. Biochemistry. 1998;37(18):6361-6. doi: 10.1021/bi980218m. PubMed PMID:
9572851.
126. Dittmann M, Hoffmann HH, Scull MA, Gilmore RH, Bell KL, Ciancanelli M, et al. A serpin
shapes the extracellular environment to prevent influenza A virus maturation. Cell.
2015;160(4):631-43. doi: 10.1016/j.cell.2015.01.040. PubMed PMID: 25679759; PubMed
Central PMCID: PMCPMC4328142.
127. Burke JM, St Clair LA, Perera R, Parker R. SARS-CoV-2 infection triggers widespread host
mRNA decay leading to an mRNA export block. RNA. 2021;27(11):1318-29. Epub 20210727. doi:
10.1261/rna.078923.121. PubMed PMID: 34315815; PubMed Central PMCID:
PMCPMC8522697.
128. Zhang K, Miorin L, Makio T, Dehghan I, Gao S, Xie Y, et al. Nsp1 protein of SARS-CoV-2
disrupts the mRNA export machinery to inhibit host gene expression. Sci Adv. 2021;7(6). Epub
20210205. doi: 10.1126/sciadv.abe7386. PubMed PMID: 33547084; PubMed Central PMCID:
PMCPMC7864571.
129. Kimura I, Konno Y, Uriu K, Hopfensperger K, Sauter D, Nakagawa S, et al. Sarbecovirus
ORF6 proteins hamper induction of interferon signaling. Cell Rep. 2021;34(13):108916. Epub
20210312. doi: 10.1016/j.celrep.2021.108916. PubMed PMID: 33765414; PubMed Central
PMCID: PMCPMC7953434.
130. Setaro AC, Gaglia MM. All hands on deck: SARS-CoV-2 proteins that block early anti-viral
interferon responses. Curr Res Virol Sci. 2021;2:100015. Epub 20211112. doi:
10.1016/j.crviro.2021.100015. PubMed PMID: 34786565; PubMed Central PMCID:
PMCPMC8588586.
131. Ogunjimi B, Zhang SY, Sorensen KB, Skipper KA, Carter-Timofte M, Kerner G, et al. Inborn
errors in RNA polymerase III underlie severe varicella zoster virus infections. J Clin Invest.
2017;127(9):3543-56. Epub 20170807. doi: 10.1172/JCI92280. PubMed PMID: 28783042;
PubMed Central PMCID: PMCPMC5669568.
132. Zimmer MM, Kibe A, Rand U, Pekarek L, Ye L, Buck S, et al. The short isoform of the host
antiviral protein ZAP acts as an inhibitor of SARS-CoV-2 programmed ribosomal frameshifting.
Nat Commun. 2021;12(1):7193. Epub 20211210. doi: 10.1038/s41467-021-27431-0. PubMed
PMID: 34893599; PubMed Central PMCID: PMCPMC8664833.

133.   Klein S, Golani G, Lolicato F, Lahr C, Beyer D, Herrmann A, et al. IFITM3 blocks influenza
virus entry by sorting lipids and stabilizing hemifusion. Cell Host Microbe. 2023. Epub 20230329.
doi: 10.1016/j.chom.2023.03.005. PubMed PMID: 37003257.
134.   Guo X, Steinkuhler J, Marin M, Li X, Lu W, Dimova R, et al. Interferon-Induced
Transmembrane Protein 3 Blocks Fusion of Diverse Enveloped Viruses by Altering Mechanical
Properties of Cell Membranes. ACS Nano. 2021;15(5):8155-70. Epub 20210303. doi:
10.1021/acsnano.0c10567. PubMed PMID: 33656312; PubMed Central PMCID:
PMCPMC8159881.
135.   Suddala KC, Lee CC, Meraner P, Marin M, Markosyan RM, Desai TM, et al. Interferon-
induced transmembrane protein 3 blocks fusion of sensitive but not resistant viruses by
partitioning into virus-carrying endosomes. PLoS Pathog. 2019;15(1):e1007532. Epub 20190114.
doi: 10.1371/journal.ppat.1007532. PubMed PMID: 30640957; PubMed Central PMCID:
PMCPMC6347298.
136.   Desai TM, Marin M, Chin CR, Savidis G, Brass AL, Melikyan GB. IFITM3 restricts influenza
A virus entry by blocking the formation of fusion pores following virus-endosome hemifusion.
PLoS Pathog. 2014;10(4):e1004048. Epub 20140403. doi: 10.1371/journal.ppat.1004048.
PubMed PMID: 24699674; PubMed Central PMCID: PMCPMC3974867.
137.   Fadeel B, Gleiss B, Hogstrand K, Chandra J, Wiedmer T, Sims PJ, et al. Phosphatidylserine
exposure during apoptosis is a cell-type-specific event and does not correlate with plasma
membrane phospholipid scramblase expression. Biochem Biophys Res Commun.
1999;266(2):504-11. doi: 10.1006/bbrc.1999.1820. PubMed PMID: 10600532.
138.   Zhou Q, Zhao J, Al-Zoghaibi F, Zhou A, Wiedmer T, Silverman RH, et al. Transcriptional
control of the human plasma membrane phospholipid scramblase 1 gene is mediated by
interferon-alpha. Blood. 2000;95(8):2593-9. PubMed PMID: 10753839.
139.   Zhou Q, Zhao J, Wiedmer T, Sims PJ. Normal hemostasis but defective hematopoietic
response to growth factors in mice deficient in phospholipid scramblase 1. Blood.
2002;99(11):4030-8. doi: 10.1182/blood-2001-12-0271. PubMed PMID: 12010804.
140.   Kusano S, Eizuru Y. Human phospholipid scramblase 1 interacts with and regulates
transactivation of HTLV-1 Tax. Virology. 2012;432(2):343-52. Epub 20120711. doi:
10.1016/j.virol.2012.06.019. PubMed PMID: 22789739.
141.   Kusano S, Eizuru Y. Interaction of the phospholipid scramblase 1 with HIV-1 Tat results in
the repression of Tat-dependent transcription. Biochem Biophys Res Commun. 2013;433(4):438-
44. Epub 20130315. doi: 10.1016/j.bbrc.2013.02.098. PubMed PMID: 23501106.
142.   Sadanari H, Takemoto M, Ishida T, Otagiri H, Daikoku T, Murayama T, et al. The
Interferon-Inducible Human PLSCR1 Protein Is a Restriction Factor of Human Cytomegalovirus.
Microbiol Spectr. 2022;10(1):e0134221. Epub 2022/02/10. doi: 10.1128/spectrum.01342-21.
PubMed PMID: 35138119; PubMed Central PMCID: PMCPMC8826943.
143.   Metz P, Dazert E, Ruggieri A, Mazur J, Kaderali L, Kaul A, et al. Identification of type I and
type II interferon-induced effectors controlling hepatitis C virus replication. Hepatology.
2012;56(6):2082-93. Epub 20121014. doi: 10.1002/hep.25908. PubMed PMID: 22711689.
144.   Yang J, Zhu X, Liu J, Ding X, Han M, Hu W, et al. Inhibition of Hepatitis B virus replication
by phospholipid scramblase 1 in vitro and in vivo. Antiviral Res. 2012;94(1):9-17. Epub
20120209. doi: 10.1016/j.antiviral.2012.01.010. PubMed PMID: 22342889.

145.   Kusano S, Ikeda M. Interaction of phospholipid scramblase 1 with the Epstein-Barr virus
protein BZLF1 represses BZLF1-mediated lytic gene transcription. J Biol Chem.
2019;294(41):15104-16. Epub 2019/08/23. doi: 10.1074/jbc.RA119.008193. PubMed PMID:
31434743; PubMed Central PMCID: PMCPMC6791327.
146.   Luo W, Zhang J, Liang L, Wang G, Li Q, Zhu P, et al. Phospholipid scramblase 1 interacts
with influenza A virus NP, impairing its nuclear import and thereby suppressing virus replication.
PLoS Pathog. 2018;14(1):e1006851. Epub 20180119. doi: 10.1371/journal.ppat.1006851.
PubMed PMID: 29352288; PubMed Central PMCID: PMCPMC5792031.
147.   Dal Col J, Lamberti MJ, Nigro A, Casolaro V, Fratta E, Steffan A, et al. Phospholipid
scramblase 1: a protein with multiple functions via multiple molecular interactors. Cell Commun
Signal. 2022;20(1):78. Epub 2022/06/02. doi: 10.1186/s12964-022-00895-3. PubMed PMID:
35650588.
148.   Jonathan RS-G, Audrey A, Payman S-T, Hala A, Vinitha M, Dae-Kyum K, et al. A SARS-CoV-
2 BioID-based virus-host membrane protein interactome and virus peptide compendium: new
proteomics resources for COVID-19 research. bioRxiv. 2020:2020.08.28.269175. doi:
10.1101/2020.08.28.269175.
149.   Guo K, Barrett BS, Morrison JH, Mickens KL, Vladar EK, Hasenkrug KJ, et al. Interferon
resistance of emerging SARS-CoV-2 variants. Proc Natl Acad Sci U S A.
2022;119(32):e2203760119. Epub 20220722. doi: 10.1073/pnas.2203760119. PubMed PMID:
35867811; PubMed Central PMCID: PMCPMC9371743.
150.   Shalamova L, Felgenhauer U, Wilhelm J, Schaubmar AR, Buttner K, Schoen A, et al.
Omicron variant of SARS-CoV-2 exhibits an increased resilience to the antiviral type I interferon
response. PNAS Nexus. 2022;1(2):pgac067. Epub 20220523. doi: 10.1093/pnasnexus/pgac067.
PubMed PMID: 36713328; PubMed Central PMCID: PMCPMC9802332.
151.   Shi G, Li T, Lai KK, Johnson RF, Yewdell JW, Compton AA. Omicron Spike confers enhanced
infectivity and interferon resistance to SARS-CoV-2 in human nasal tissue. Nat Commun.
2024;15(1):889. Epub 20240130. doi: 10.1038/s41467-024-45075-8. PubMed PMID: 38291024;
PubMed Central PMCID: PMCPMC10828397.
152.   Willett BJ, Grove J, MacLean OA, Wilkie C, De Lorenzo G, Furnon W, et al. SARS-CoV-2
Omicron is an immune escape variant with an altered cell entry pathway. Nat Microbiol.
2022;7(8):1161-79. Epub 20220707. doi: 10.1038/s41564-022-01143-7. PubMed PMID:
35798890; PubMed Central PMCID: PMCPMC9352574.
153.   Peacock TP, Brown JC, Zhou J, Thakur N, Sukhova K, Newman J, et al. The altered entry
pathway and antigenic distance of the SARS-CoV-2 Omicron variant map to separate domains of
spike protein. bioRxiv. 2022:2021.12.31.474653. doi: 10.1101/2021.12.31.474653.
154.   Sette A, Crotty S. Adaptive immunity to SARS-CoV-2 and COVID-19. Cell.
2021;184(4):861-80. Epub 20210112. doi: 10.1016/j.cell.2021.01.007. PubMed PMID:
33497610; PubMed Central PMCID: PMCPMC7803150.
155.   Chen S, Francioli LC, Goodrich JK, Collins RL, Kanai M, Wang Q, et al. A genomic
mutational constraint map using variation in 76,156 human genomes. Nature.
2024;625(7993):92-100. doi: 10.1038/s41586-023-06045-0.
156.   Mar KB, Van Dyke MC, Lopez AH, Eitson JL, Fan W, Hanners NW, et al. LY6E protects mice
from pathogenic effects of murine coronavirus and SARS-CoV-2. bioRxiv. 2023. Epub 20230126.

doi: 10.1101/2023.01.25.525551. PubMed PMID: 36747632; PubMed Central PMCID:
PMCPMC9900800.
157.   Pfaender S, Mar KB, Michailidis E, Kratzel A, Boys IN, V'Kovski P, et al. LY6E impairs
coronavirus fusion and confers immune control of viral disease. Nat Microbiol. 2020;5(11):1330-
9. Epub 20200723. doi: 10.1038/s41564-020-0769-y. PubMed PMID: 32704094; PubMed
Central PMCID: PMCPMC7916999.
158.   Consortium GT. The Genotype-Tissue Expression (GTEx) project. Nat Genet.
2013;45(6):580-5. doi: 10.1038/ng.2653. PubMed PMID: 23715323; PubMed Central PMCID:
PMCPMC4010069.
159.   Blight KJ, McKeating JA, Rice CM. Highly permissive cell lines for subgenomic and
genomic hepatitis C virus RNA replication. J Virol. 2002;76(24):13001-14. doi:
10.1128/jvi.76.24.13001-13014.2002. PubMed PMID: 12438626; PubMed Central PMCID:
PMCPMC136668.
160.   Zhang L, Bukreyev A, Thompson CI, Watson B, Peeples ME, Collins PL, et al. Infection of
ciliated cells by human parainfluenza virus type 3 in an in vitro model of human airway
epithelium. J Virol. 2005;79(2):1113-24. doi: 10.1128/JVI.79.2.1113-1124.2005. PubMed PMID:
15613339; PubMed Central PMCID: PMCPMC538579.
161.   Hoffmann HH, Schneider WM, Rozen-Gagnon K, Miles LA, Schuster F, Razooky B, et al.
TMEM41B Is a Pan-flavivirus Host Factor. Cell. 2021;184(1):133-48 e20. Epub 20201209. doi:
10.1016/j.cell.2020.12.005. PubMed PMID: 33338421; PubMed Central PMCID:
PMCPMC7954666.
162.   Benboudjema L, Mulvey M, Gao Y, Pimplikar SW, Mohr I. Association of the herpes
simplex virus type 1 Us11 gene product with the cellular kinesin light-chain-related protein PAT1
results in the redistribution of both polypeptides. J Virol. 2003;77(17):9192-203. doi:
10.1128/jvi.77.17.9192-9203.2003. PubMed PMID: 12915535; PubMed Central PMCID:
PMCPMC187382.
163.   Lee D, Le Pen J, Yatim A, Dong B, Aquino Y, Ogishi M, et al. Inborn errors of OAS-RNase L
in SARS-CoV-2-related multisystem inflammatory syndrome in children. Science.
2023;379(6632):eabo3627. Epub 20230210. doi: 10.1126/science.abo3627. PubMed PMID:
36538032.
164.   Mendoza EJ, Manguiat K, Wood H, Drebot M. Two Detailed Plaque Assay Protocols for
the Quantification of Infectious SARS-CoV-2. Curr Protoc Microbiol. 2020;57(1):ecpmc105. doi:
10.1002/cpmc.105. PubMed PMID: 32475066; PubMed Central PMCID: PMCPMC7300432.
165.   Polo JM, Davis NL, Rice CM, Huang HV, Johnston RE. Molecular analysis of Sindbis virus
pathogenesis in neonatal mice by using virus recombinants constructed in vitro. J Virol.
1988;62(6):2124-33. doi: 10.1128/JVI.62.6.2124-2133.1988. PubMed PMID: 2835514; PubMed
Central PMCID: PMCPMC253309.
166.   Bick MJ, Carroll JW, Gao G, Goff SP, Rice CM, MacDonald MR. Expression of the zinc-
finger antiviral protein inhibits alphavirus replication. J Virol. 2003;77(21):11555-62. doi:
10.1128/jvi.77.21.11555-11562.2003. PubMed PMID: 14557641; PubMed Central PMCID:
PMCPMC229374.
167.   Atasheva S, Kim DY, Akhrymuk M, Morgan DG, Frolova EI, Frolov I. Pseudoinfectious
Venezuelan equine encephalitis virus: a new means of alphavirus attenuation. J Virol.

2013;87(4):2023-35. Epub 20121205. doi: 10.1128/JVI.02881-12. PubMed PMID: 23221545;
PubMed Central PMCID: PMCPMC3571482.
168.   Dalton KP, Rose JK. Vesicular stomatitis virus glycoprotein containing the entire green
fluorescent protein on its cytoplasmic domain is incorporated efficiently into virus particles.
Virology. 2001;279(2):414-21. doi: 10.1006/viro.2000.0736. PubMed PMID: 11162797.
169.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21. Epub 20121025. doi:
10.1093/bioinformatics/bts635. PubMed PMID: 23104886; PubMed Central PMCID:
PMCPMC3530905.
170.   Liao Y, Smyth GK, Shi W. The R package Rsubread is easier, faster, cheaper and better for
alignment and quantification of RNA sequencing reads. Nucleic Acids Res. 2019;47(8):e47. doi:
10.1093/nar/gkz114. PubMed PMID: 30783653; PubMed Central PMCID: PMCPMC6486549.
171.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol. 2014;15(12):550. doi: 10.1186/s13059-014-0550-8.
PubMed PMID: 25516281; PubMed Central PMCID: PMCPMC4302049.
172.   Korotkevich G, Sukhov V, Budin N, Shpak B, Artyomov MN, Sergushichev A. Fast gene set
enrichment analysis. bioRxiv. 2021:060012. doi: 10.1101/060012.
173.   Liberzon A, Subramanian A, Pinchback R, Thorvaldsdottir H, Tamayo P, Mesirov JP.
Molecular signatures database (MSigDB) 3.0. Bioinformatics. 2011;27(12):1739-40. Epub
20110505. doi: 10.1093/bioinformatics/btr260. PubMed PMID: 21546393; PubMed Central
PMCID: PMCPMC3106198.
174.   Fabregat A, Jupe S, Matthews L, Sidiropoulos K, Gillespie M, Garapati P, et al. The
Reactome Pathway Knowledgebase. Nucleic Acids Res. 2018;46(D1):D649-D55. doi:
10.1093/nar/gkx1132. PubMed PMID: 29145629; PubMed Central PMCID: PMCPMC5753187.
175.   Kanehisa M, Goto S. KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids Res.
2000;28(1):27-30. doi: 10.1093/nar/28.1.27. PubMed PMID: 10592173; PubMed Central
PMCID: PMCPMC102409.
176.   Agrawal A, Balci H, Hanspers K, Coort SL, Martens M, Slenter DN, et al. WikiPathways
2024: next generation pathway database. Nucleic Acids Res. 2024;52(D1):D679-D89. doi:
10.1093/nar/gkad960. PubMed PMID: 37941138; PubMed Central PMCID: PMCPMC10767877.
177.   Schaefer CF, Anthony K, Krupa S, Buchoff J, Day M, Hannay T, et al. PID: the Pathway
Interaction Database. Nucleic Acids Res. 2009;37(Database issue):D674-9. Epub 20081002. doi:
10.1093/nar/gkn653. PubMed PMID: 18832364; PubMed Central PMCID: PMCPMC2686461.
178.   Nishimura D. BioCarta. Biotech Software & Internet Report: The Computer Software
Journal for Scient. 2001;2(3):117-20.
179.   Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C(T) method.
Nat Protoc. 2008;3(6):1101-8. doi: 10.1038/nprot.2008.73. PubMed PMID: 18546601.
180.   Dittmann M, Hoffmann HH, Scull MA, Gilmore RH, Bell KL, Ciancanelli M, et al. A serpin
shapes the extracellular environment to prevent influenza A virus maturation. Cell.
2015;160(4):631-43. doi: 10.1016/j.cell.2015.01.040. PubMed PMID: 25679759; PubMed
Central PMCID: PMC4328142.
181.   Majdoul S, Compton AA. Lessons in self-defence: inhibition of virus entry by intrinsic
immunity. Nat Rev Immunol. 2022;22(6):339-52. Epub 20211013. doi: 10.1038/s41577-021-
00626-8. PubMed PMID: 34646033; PubMed Central PMCID: PMCPMC8511856.

182.   Bruchez A, Sha K, Johnson J, Chen L, Stefani C, McConnell H, et al. MHC class II
transactivator CIITA induces cell resistance to Ebola virus and SARS-like coronaviruses. Science.
2020;370(6513):241-7. Epub 20200827. doi: 10.1126/science.abb3753. PubMed PMID:
32855215; PubMed Central PMCID: PMCPMC7665841.
183.   Doyle T, Moncorge O, Bonaventure B, Pollpeter D, Lussignol M, Tauziet M, et al. The
interferon-inducible isoform of NCOA7 inhibits endosome-mediated viral entry. Nat Microbiol.
2018;3(12):1369-76. Epub 20181126. doi: 10.1038/s41564-018-0273-9. PubMed PMID:
30478388; PubMed Central PMCID: PMCPMC6329445.
1934

Figure 1

Figure 2

Figure 3

SARS-CoV-2 functional genetic screens (26 publications)

COVID-19 human genetics studies (24 publications)

2,603 genes

113 genes

28 genes

PROVIRAL

Virus entry
ACE2*   TMPRSS2

Cell adhesion and immune signaling
ICAM3   PCDH7

Intracellular trafficking and signaling
RAB2A

Guanyl-nucleotide exchange
ARHGEF38

IFN signaling
IRF1

Ion channels and membrane potential
KCNC3

Transcriptional regulation
FOXP4   ZBTB32

Cell metabolism and redox homeostasis
CAT*   FDX2*

ANTIVIRAL

Protective mucous barrier
MUC1*

Cell adhesion and immune signaling
ICAM1*

TLR7 signaling
TLR7
MYD88   UNC93B1

IFN signaling
IFNAR1*   IFNAR2*
JAK1*
STAT2*

IFN-stimulated genes
OAS1   OAS2   OAS3   PLSCR1*

Transcriptional regulation
KAT7*   ZBTB11*

RNA splicing
CLK2

* Hit in arrayed CRISPR KO screen.

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Supplementary figure 1

Supplementary figure 2

Supplementary figure 3

A

B

C

Supplementary figure 4

Supplementary figure 5

Supplementary figure 6

Actin

PLSCR1

WT background   KO background

Supplementary figure 7

A

B

Supplementary figure 8