Title: Foam rolling intervention improves lactate clearance after high-intensity exercise

Running head: Foam rolling improves lactate clearance

Kazuki Kasahara,[1] Keita Oneyama,[2] Takeru Ito,[2] Masatoshi Nakamura,[3] Genta Ochi[2,4]

[1] Sanyudo hospital, Fukuda Town, Yonezawa City, Yamagata, Japan

[2] Department of Health and Sports, Niigata University of Health and Welfare, Niigata, Japan

[3] Faculty of Rehabilitation Sciences, Nishi Kyushu University, 4490-9 Ozaki, Kanzaki, Saga, 842-

8585, Japan

[4] Institute for Human Movement and Medical Sciences, Niigata University of Health and Welfare,

Niigata, Japan

Corresponding Author:

Genta Ochi

Department of Health and Sports, Niigata University of Health and Welfare

Kita-ku, Shimami-cho 1398, Niigata, 950-3198

Phone: +81-025-257-4595

Email: ochi@nuhw.ac.jp

Statements and Declarations

Funding: This work was supported by the Japan Society for the Promotion of Science (JSPS)

Grant JP22K17739 (Genta Ochi) and the Grant-in-Aid for Scientific Research from the Niigata

University of Health and Welfare, 2023 (Genta Ochi).

Author Contributions: KK, MN, and GO designed the study. KK, KO, TI, and GO were

responsible for data acquisition. KK, MN, and GO performed data analysis and interpretation. All

authors were involved in drafting and critically revising the manuscript for intellectual content.

All authors have read and approved the final version of the manuscript.

Research involving human participants: This study was conducted in accordance with the

Declaration of Helsinki and approved by the Ethics Committee of the Niigata University of Health

and Welfare, Niigata, Japan (approval number 19170-231019). The participants were informed of

data confidentiality, the study details, and provided written informed consent to participate in the

study.

Conflict of Interest: The authors declare no conflicts of interest.

Data availability: All data generated or analyzed during this study are included in this article.

Abstract

Purpose: The effects foam rolling (FR) intervention on lactate clearance and the impaired executive function associated with fatigue after high-intensity exercise remain unclear. This study examined whether FR is an effective tool for fatigue recovery.

Methods: Eighteen healthy adults (10 males, 8 females; 21.8±1.7 years) without consistent exercise habits participated. The participants performed high-intensity exercises using an ergometer with a progressive load. Post-exercise FR intervention was compared with the control condition. Measurements included lactate, vigor/fatigue by the Profile of Mood States 2nd edition, cognitive function (cognitive task performance), and leg and body rating of perceived exertion (RPE) pre- and post-exercise, and post-intervention (5 min post-exercise).

Results: The blood lactate concentration post-FR intervention (-7.3±3.0 mmol/L) was significantly reduced. Increased lactate clearance by FR correlated with faster recovery of executive function for those with greater lactate clearance, although cognitive fatigue was not observed after high-intensity exercise ($P = .086$, $r = 0.41$). Lactate clearance was not significantly correlated with RPE in the FR condition, whereas RPE decreased with increased lactate clearance for those with greater lactate clearance in the control condition (leg; $r = 0.778$, body; $r = 0.669$).

Conclusions: These results indicated FR intervention may be an effective recovery tool for exhausting exercises.

Keywords: foam roller; fatigue; Profile of Mood States 2nd Edition (POMS2); recovery; cognitive function

INTRODUCTION

Many sports competitions involve high-intensity interval exercises involving
repeated sprints and jumps. Ball sports, such as basketball and handball, require high-
intensity interval exercises for extended periods with little time between halves or sets. High-
intensity interval exercise repeated with short rest periods, results in a noticeable decrease in
performance in the second half of matches. This resulting performance loss may include
muscle function as well as executive function.[1–3] Executive function is the foundation of
tactical understanding and execution, which includes the ability to make selective
decisions during sport activities,[4] and is a major determinant to athletic performance.
Cook and Beaven[5] reported that recovery to a good psychological state post-exercise is
important for maintaining performance, whereas recovery of both peripheral and central
functions post-exercise is an urgent issue. However, existing recovery methods, such as cold
water immersion[6] and full-body aerobic exercises, including a rowing ergometer,[7] are
difficult to implement during a game because of the time, location, and competition rules.
To maintain mental and physical functions until the latter half of a match, a new recovery
method that can be implemented in a limited space and short duration is required.

Muscle fatigue caused by high-intensity exercise is related to acidosis from
lactate production.[8] Therefore, many recovery protocols or methods target lactate
clearance.[9,10,11,12] Interestingly, increased peripheral blood lactate concentration induced
by strenuous exercise or intravenous infusion of lactate solution is inversely related to
attentional task performance,[13] suggesting that increased peripheral blood lactate
concentration may be a biomarker for cognitive fatigue because lactate crosses the blood-
brain barrier. Improving adequate lactate clearance may improve muscle fatigue and
executive dysfunction, thereby improving athletic performance; however none of the
previous recovery methods have focused on executive function.

Foam rolling (FR) is commonly used as a recovery method. Adamczyk et al.[14]
evaluated the effects of FR intervention after squat jumps at maximal effort for 1 min,
which significantly increased lactate clearance for the FR condition than the intervention
condition at 30 min post-exercise. In addition, Akinci et al.[15] implemented circuit-based
high-intensity exercise at 85% intensity, as calculated from the heart rate reserve method,
and measured lactate levels before exercise training, before the recovery period, and at 5
and 20 min into the recovery period to examine the effects of FR intervention on lactate
levels. They reported that FR intervention did not alter lactate concentrations. However,
these studies may not have adequately evaluated the effects of these exercises on

primarily active and fatigued muscles during exercise because of the long time
elapsed after the exercise and it is a whole-body exercise.

This study purported to determine whether FR improves lactate
clearance after exercise with fatigue and consequently improve impaired
executive function.

METHODS

Experimental Approach to the Problem
This study examined the immediate effects of FR, which can be performed in a confined
space for a limited period, on lactate clearance and cognitive function after high-intensity
exercise with progressive loading. In addition, this study assessed these effects separately
in males and females.

A randomized repeated-measures experimental design was used to compare the two
conditions: control and FR. Participants were instructed to visit the laboratory twice, with
a $\geq$48 h break. Measurements were conducted before the intervention (pre), after the
intervention (post), and 5 min post-intervention. The completed psychological scales
included the Profile of Mood States 2nd Edition (POMS2)[16] and participant ratings of
perceived exertion (RPE).[17] The parameters measured were executive function,
psychological scale scores, and blood lactate concentration.

Participants
A total of 22 right-handed Japanese-speaking young adults (12 males and 10 females) were
included in this study conducted between October 2023 and January 2024. The inclusion
criteria were: no regular exercise habits; no daily use of a foam roller; native Japanese
speakers; and no history of neurological, psychiatric, or respiratory diseases. All participants
were instructed to refrain from eating or drinking anything other than mineral water 2 h
before starting the experiment. In this study, analyses were conducted with data from 18
participants (10 males and 8 females); four participants were excluded because of their
inability to perform the FR intervention after high-intensity exercise. Table 1 presents the
participant characteristics. We performed a post-hoc sensitivity analysis using this sample
with 80% power and an $\alpha = 0.05$ demonstrated sufficient sensitivity to detect repeated-
measures effects exceeding $f = 0.31$ as computed using G*Power (3.1.9.2; The G*Power
Team).

This study was conducted in accordance with the Declaration of Helsinki and approved
by the Ethics Committee of the Niigata University of Health and Welfare, Niigata, Japan
(approval number 19170-231019). The participants were informed of data confidentiality
and the study details, and provided written informed consent to participate in the study.
131

Procedures

High-Intensity Exercise with a Progressive Load

The participants performed high-intensity exercises with progressive loading using an
ergometer (828E; Monark, Sweden). They performed an ergometer exercise for 3 min at a
load of 0.5 KP as a warm-up, followed by a 1-min rest period. After the break, males and
females performed ergometer exercises with a progressive increase in the load of 0.5 KP/min.
Participants performed the ergometer exercise and were instructed to pedal on time using a
metronome (Smart Metronome; Tomohiro Ihara, Japan) set at 60 bpm. The exercise was
stopped when the pedaling speed could no longer be sustained. During the exercise, strong
verbal encouragement was provided to elicit maximal effort. Exercise time and heart rate
(HR) pre- and post-intervention were measured during each session using a OH1 optical HR
sensor (Polar, Finland) worn on the arm.

FR Intervention

Figure 1 shows the FR intervention method using a foam roller (Stretch Roll SR-002,
Dream Factory, Umeda, Japan). A physical therapist instructed the participants prior to
the intervention. FR intervention was initiated with knee flexors, followed by knee
extensors. FR interventions were performed on the right and left sides. The FR area of the
knee flexor muscle group was defined as the area proximal from the knee fossa to the
sciatic tuberosity.[18] The knee extensor was defined as the superior anterior iliac spine at
the top of the patella.[19,20] One FR cycle was defined as one distal rolling movement
followed by one proximal rolling movement performed within 4 s.[21] The intervention
intensity was defined as the maximum intensity that the participants could tolerate (Kasahara
et al. 2022). The intervention time for each area was 60 s, and the participants were instructed
to prepare for the intervention for the next area during a 20-s interval. We used a metronome
(Smart Metronome; Tomohiro Ihara, Japan) operating at 60 bpm for control.

Psychological Measurements

POMS2 contains 35 items and evaluates seven mood states (anger-hostility, confusion-
bewilderment, depression-ejection, fatigue-inertia, tension-anxiety, vigor-activity, and
friendliness). This study used only a 10-item subset of vigor-activity and fatigue–inertia
to reduce the psychological burden on the participants.

We recorded the rate of perceived exertion (RPE) before and after exercise to assess
psychological exercise intensity in the whole body and lower extremities.

Blood Lactate Concentration
We used a Lactate Pro 2 (AKRAY, The Netherlands), a handheld point-of-care analyzer,
for enzymatic amperometric detection. It requires 0.3 μl of a whole blood sample and a
short time (15 s) to measure the lactate level. Lactate Pro 2 measurements range from 0.5
to 25.0 mmol/L. Therefore, only lactate values between 0.5 and 25 were included in this
study.

Cognitive task performance
We employed Schneider's spatial Stroop task to measure executive function.[22] This task
was created using web-building platforms (Lab.js v19.1.0).[23] In each experiment, the
participants made a spatially compatible key-press response to classify the spatial
meaning of a stimulus presented at an irrelevant position. The stimulus meaning and
position were either matched (congruent) or mismatched (incongruent) in the trial, with
16 congruent and 16 incongruent stimuli. Visual alert cues (squares) were briefly
presented at two potential stimulus positions 500 ms before stimulus onset. Although
previous studies used multiple stimulus types (arrows or words) and spatial dimensions
(horizontal or vertical), this study employed arrow stimulus-type tasks. In addition, since
preliminary experiments revealed that reaction times tended to be slower in vertical than
in horizontal spatial dimension tasks, and the load on the task was assumed to be higher
(fatigue was more likely to occur), vertical spatial dimension tasks were used in the
present study.

Participants pressed a key indicating up ("y") or down ("b") to classify the direction of
an up- or down-pointing arrow positioned above or below fixation (Figure 2). The correct
answer rates assigned to yes and no were 50%. Each stimulus was separated by an inter-
stimulus interval showing a fixation cross for 2 seconds to avoid prediction of the timing
of the subsequent trial. The stimulus remained on the screen until the patient responded
or for 1 s. This study adopted Stroop interference, a specifically defined cognitive process,
to elucidate the effects of an acute bout of exercise on executive function by calculating
the incongruent–congruent contrast, which was assumed to represent Stroop interference.

Data from 16 participants (eight males and eight females) were used only for the cognitive
task because data from two of the 18 participants could not be obtained.
194

Statistical Analyses

R software (4.3.2), Rstudio (2023.12.0 + 369), "EZR on R Commander" (12), and the

"anovaun" package were used for data analyses (The R Foundation, Vienna, Austria). A

three-way analysis of variance (ANOVA) was performed to compare conditions, times,

and sex. First, Mauchly's sphericity test was used to determine whether sphericity was

maintained. When this assumption was confirmed, a three-way ANOVA with

Greenhouse–Geisser's epsilon correction was performed; otherwise, only a three-way

ANOVA was performed. If the 3-way ANOVA was not significant, a 2-way ANOVA was

performed. The effect size classification was set where $\eta_p^2 < 0.01$ was considered small;

0.02–0.1, medium; and >0.1, large.[24] A corresponding t test with Holm's correction was

used to determine significant differences obtained from the ANOVA. Pearson's

correlation analyses were performed to clarify the relationships between the parameters

and executive performance. Statistical significance was set a priori at $P < .05$.

RESULTS

Physiological Parameters

Table 2 presents the results of the HR, RPE, total exercise time, and kp-max in the

control and FR conditions. No significant differences in the pre-test values were observed

between the two conditions for any of the measures. HR increased significantly ($P < .01$)

immediately post-exercise in both conditions (control: 101.9±14.4, FR: 97.7±22.5). RPE

(leg/body) was significantly ($P < .01$) higher at post-exercise (leg, control: 10.4±3.0, FR:

10.6±2.9; body, control: 9.1±2.6, FR: 8.9±3.5) and 5 min into the resting period (leg,

control: 6.7±3.8, FR: 6.2±3.1; body, control: 5.1±3.9, FR: 4.7±3.3) as compared to pre-

exercise in both conditions but significantly ($P < 0.01$) lower at 5 min into the resting

period when compared with post-exercise (leg, control: -3.7±2.0, FR: -4.4±2.4; body,

control: -4.0±2.3, FR: -4.2±3.0). In addition, sex differences were observed in the total

exercise time and post-exercise load, which were significantly ($P < .01$) higher in males

than in females. However, no significant differences were observed between the two

conditions.

Figure 3 presents the changes in lactate levels. A three-way ANOVA (conditions vs. time

vs. sex) showed no significant interactions ($F (2, 32) = 1.0047$, $P = 0.3774$, $\eta_p^2 = 0.0591$).

A two-way ANOVA (conditions vs. time) showed a significant interaction ($F (2,32) =$

4.0433, $P < .05$, $\eta_p^2 = 0.2017$) and a main effect of time ($F (2, 32) = 275.1504$, $P < .0001$,

$\eta_p^2 = 0.9450$). The post-hoc test results in the control condition indicated significantly

higher values at post-exercise (10.6±2.0 mmol/L) and 5 min into the resting period
(9.6±2.9 mmol/L) than those at pre-exercise ($P < .01$), although no significant difference
was observed between post-exercise and 5 min into the resting period (-1.0±2.1 mmol/L).
In the FR condition, post-exercise (16.3±2.0 mmol/L) and 5 min into the resting period
(9.0±2.5 mmol/L) values were significantly higher than pre-exercise ($P < .01$). In addition,
results at 5 min into the resting period (-7.3±3.0 mmol/L) were significantly lower than
that at post-exercise ($P < .01$).

Psychological Parameters
A three-way ANOVA (conditions vs. time vs. sex) showed no significant interaction between
vigor ($F(2, 32) = 1.6850$, $P = .2015$, $\eta_p^2 = 0.0953$) and fatigue ($F(2, 32) = 1.0022$, P
$= .3783$, $\eta_p^2 = 0.0589$) in POMS2. A two-way ANOVA (conditions vs. time) showed a main
effect of time on vigor ($F(2, 32) = 7.5235$, $P < .01$, $\eta_p^2 = 0.3198$) and fatigue ($F(2, 32) =$
163.5833, $P < .0001$, $\eta_p^2 = 0.9109$). A post-hoc test using Holm correction revealed that
vigor was significantly lower in post-exercise (control: -2.2±4.6, FR:-4.1±5.4) than in pre-
exercise conditions ($P < .01$). Fatigue was significantly higher at post-exercise (control:
12.7±4.1, FR: 12.4±3.7) and 5 min into the resting period (control: 8.1±4.4, FR: 8.4±4.8)
as compared to that at pre-exercise ($P < .0001$). In addition, post-exercise values (control: -
4.6±2.9, FR: -4.0±3.2) were significantly higher than those at 5 min into the resting period
($P < .0001$).

Relationship Between Parameters and Performance
Figure 4 displays the changes in lactate levels and executive function pre- and post-
intervention. A significant correlation was observed for the FR condition ($r = 0.41$, $P = .086$).
Sex-specific results showed a significant correlation for males ($r = 0.706$, $P < .05$), but not
for females ($r = 0.13$, $P = .747$). No significant correlation was found in the control group (r
$= -0.38$, $P = .118$).

Figure 5 presents the changes in lactate levels and RPE (leg/body) before and after the
intervention. No significant correlation was found in the FR condition; however, a
significant correlation was found in the control condition (leg: $r = 0.778$, $P < .01$; body: $r =$
0.669, $P < .01$).

DISCUSSION

We examined the acute effects of FR intervention on lactate clearance,
psychological parameters, and cognitive function after a progressively increasing load
exercise, resulting in exhaustion to propose FR as a new recovery method that athletes can
implement in a limited space. The results revealed that arterial blood lactate concentration,
which increased after exhaustive exercise, decreased in the FR condition regardless of sex,
indicating that FR intervention enhanced lactate clearance. Interestingly, cognitive function
improved after FR intervention as compared to those with greater lactate clearance, and this
effect was significantly greater in males than in females. This finding indicates that FR
intervention may be useful in improving exercise-induced cognitive decline, particularly in
males.

This study showed significant increases in HR at pre- and post-exercise in both the
FR and control conditions, without significant differences between them. In addition, the
Karvonen method[25] demonstrated that exercise intensity at post-exercise was $85.3 \pm 11.0\%$
in the FR condition and $83.6\% \pm 10.4\%$ in the control condition, with no significant
difference between the two conditions. Based on these results, we concluded that all
participants performed the same amount of exercise under both conditions.

First, we examined the effects of FR intervention on lactate clearance. In the present
study, ergometric exercise was performed with progressive loading by applying a strong load
to the lower extremities, and the FR intervention targeted the bilateral knee extensors and
flexors. The arterial blood lactate concentrations increased significantly immediately after
exercise under both conditions, and no difference was observed between the conditions.
Thereafter, no lactate concentration changes were observed in the control condition.
However, in the FR condition, the arterial blood lactate concentration significantly decreased
post-intervention. These results indicated that FR intervention immediately after exhaustive
exercise was useful in increasing lactate clearance. A previous study reported that increased
arterial blood lactate concentrations after high-intensity exercise were significantly lower in
active recovery groups that performed low- or high-intensity exercises than in passive
recovery groups that did not receive any intervention.[26,27] Interestingly, no significant
difference in the increasing effect of the ROM was observed between the FR and sham
rolling conditions, which simulated FR intervention movements.[28] These findings
demonstrated that active recovery, such as FR, may increase blood flow in active tissues and
lactate clearance.[29] In addition, FR intervention increases regional perfusion[30]) and,
therefore, may further increase lactate clearance.

Second, the effects of the FR intervention on executive function were
examined. In the present study, executive function task performance did not
decrease before or after exercise until exhaustion, and progressive-load exercises
leading to exhaustion did not induce cognitive fatigue. Previous study showed
progressive exercise to exhaustion caused impaired cognitive function, including
executive function[31]; however, its effects are not always clear. Whether laboratory
training impairs executive function varies greatly from person to person, and
further testing is required. Exercise in extreme environments, such as hypoxic
environments, decreases cognitive function.[2,3] To verify whether FR intervention
is effective in restoring exercise-induced cognitive decline, it should be investigated
under different exercise conditions such as a hypoxic environment.

This study demonstrated that the amount of lactate change was
significantly correlated with changes in executive function task performance
immediately after exercise and post-FR intervention in the FR condition. This result
indicates that executive function may recover in patients with increased lactate
clearance. Previous studies have reported that elevated blood lactate levels after
exhaustive exercise are associated with significantly worsened attentional
processes.[13] Therefore, increased lactate clearance due to FR may contribute to
improved cognitive performance due to increased lactate levels. Although we did
not observe a decline in executive function in this study, it may be possible to
propose a new effect of FR by testing it under experimental conditions in which
a decline in executive function occurs.

We examined the effects of the FR intervention on psychological
parameters. The results indicated that FR intervention did not affect the vigor or
fatigue of POMS2 and support the findings of a previous study[32] in which FR
intervention was administered to female basketball players and had no effect on
perceived recovery (total quality recovery) or fatigue level. However, a larger effect
size in professional male soccer players (Cohen's d = 1.08) was reported for total
perceived recovery in the FR intervention condition than in the control condition
after 24 h of training.[33] These differences in results may be due to the FR
intervention methods. Pernigoni et al. (2023) and Rey et al.[33] performed a 20-min
FR intervention on the entire lower extremity, including the quadriceps,
hamstrings, triceps, as well as abductor and adductor muscle groups. Conversely,
this study found that the FR intervention lasted 4 min for knee extensors and
knee flexors. A capacitance–response relationship has been found for the effect

of FR on the ROM increase.[34] The enhancement of psychological parameters through FR

may also be associated with the capacity–response relationship, necessitating further

research for a clearer understanding.

Regarding RPE, which represents perceived exertion, the results revealed a greater

decrease in lactate in the control condition was associated with a lower RPE. However, in

the FR condition, no significant correlation was observed between the changes in lactate and

RPE. These differences may be related to the FR intervention methods. A push-up bar was

used as the FR intervention method for the knee flexors (Figure 1A, B). Furthermore,

participants performed the intervention for the knee extensors in a plank posture (Figure 1C,

D). In this study, since we examined the intervention effects in participants who did not use

a foam roller daily, the FR intervention itself may have affected the RPE. Therefore,

although the lactate clearance increased in the FR condition, the effect of improved

perceived exertion might not have occurred. The on-field implementation of FR

interventions requires careful consideration of whether priority should be given to the

recovery of physiological parameters or subjective fatigue levels.

This study has a few limitations. First, the mechanisms underlying these results

remain unknown. The results indicated that FR intervention increased lactate clearance and

concomitantly affected executive function. However, the detailed mechanism by which this

occurs remains unclear. Future studies should examine the effects of FR intervention on

regional and cerebral perfusion and changes in lactate clearance and cognitive function.

Second limitation is the target population. As mentioned above, FR intervention for male

athletes showed fatigue recovery effects,[33] but not for female basketball players .[32] Therefore,

differences between those with exercise habits and older adults, as well as differences in

athletic specialties and sex among athletes, should be examined in the future. Finally, the

effect of FR intervention on exercise-induced cognitive decline (cognitive fatigue) was

discussed.

PRACTICAL APPLICATION

FR intervention may be a useful recovery method to improve muscle function and reduce

the decrease in cognitive performance. However, the FR intervention was not effective in

restoring RPE. Athletes and coaches should carefully consider whether the recovery of

physiological parameters or RPE should be prioritized when implementing FR interventions

in the field.

CONCLUSIONS

The results of the present study revealed that the acute effect of FR intervention after exhaustive exercise in healthy young adults was an increase in lactate clearance. Furthermore, the greater the effect, the greater the executive function recovery. However, FR intervention did not decrease RPE after exhaustive exercise. Although this study was conducted in young adults without exercise habits, further validation in athletes will allow us to propose a useful FR intervention program.

ACKNOWLEDGMENTS

The authors gratefully acknowledge all the participants involved in this study.

Funding: This work was supported by the Japan Society for the Promotion of Science (JSPS) Grant JP22K17739 (Genta Ochi) and the Grant-in-Aid for Scientific Research from the Niigata University of Health and Welfare, 2023 (Genta Ochi).

Research involving human participants: This study was conducted in accordance with the Declaration of Helsinki and approved by the Ethics Committee of the Niigata University of Health and Welfare, Niigata, Japan (approval number 19170-231019). The participants were informed of data confidentiality and the study details, and provided written informed consent to participate in the study.

Conflict of Interest: The authors declare no conflicts of interest.

Data availability: All data generated or analyzed during this study are included in this article.

REFERENCES

1. Marcora SM, Staiano W, Manning V. Mental fatigue impairs physical
performance in humans. *J Appl Physiol (1985)*. 2009;106(3):857-864.
doi:10.1152/japplphysiol.91324.2008

2. Ochi G, Yamada Y, Hyodo K, et al. Neural basis for reduced executive
performance with hypoxic exercise. *Neuroimage*. 2018;171:75-83.
doi:10.1016/j.neuroimage.2017.12.091

3. Ochi G, Kuwamizu R, Suwabe K, Fukuie T, Hyodo K, Soya H. Cognitive fatigue
due to exercise under normobaric hypoxia is related to hypoxemia during
exercise. *Sci Rep*. 2022;12(1):9835. doi:10.1038/s41598-022-14146-5

4. Vestberg T, Gustafson R, Maurex L, Ingvar M, Petrovic P. Executive functions
predict the success of top-soccer players. *PLoS One*. 2012;7(4):e34731.
doi:10.1371/journal.pone.0034731

5. Cook CJ, Beaven CM. Individual perception of recovery is related to subsequent
sprint performance. *Br J Sports Med*. 2013;47(11):705-709.
doi:10.1136/bjsports-2012-091647

6. Higgins TR, Greene DA, Baker MK. Effects of cold water immersion and
contrast water therapy for recovery from team sport: a systematic review and
meta-analysis. *J Strength Cond Res*. 2017;31(5):1443-1460.
doi:10.1519/JSC.0000000000001559

7. Raeder C, Wiewelhove T, Schneider C, et al. Effects of active recovery on muscle
function following high-intensity training sessions in elite Olympic weightlifters.
Advances in Skeletal Muscle Function Assessment. 2017;1:3–12.
doi:10.13154/294-7110

8. Robergs RA, Ghiasvand F, Parker D. Biochemistry of exercise-induced metabolic
acidosis. *Am J Physiol Regul Integr Comp Physiol*. 2004;287(3):R502-R516.
doi:10.1152/ajpregu.00114.2004

9. Fukuba Y, Walsh ML, Morton RH, Cameron BJ, Kenny CT, Banister EW. Effect
of endurance training on blood lactate clearance after maximal exercise. *J Sports
Sci*. 1999;17(3):239-248. doi:10.1080/026404199366145

10. Fleming N, Vaughan J, Feeback M. Ingestion of oxygenated water enhances
lactate clearance kinetics in trained runners. *J Int Soc Sports Nutr*. 2017;14:9.
doi:10.1186/s12970-017-0166-y

11. Warren CD, Szymanski DJ, Landers MR. Effects of three recovery protocols on range of motion, heart rate, rating of perceived exertion, and blood lactate in baseball pitchers during a simulated game. *J Strength Cond Res*. 2015;29(11):3016-3025. doi:10.1519/JSC.0000000000000487

12. Lucertini F, Gervasi M, D'Amen G, et al. Effect of water-based recovery on blood lactate removal after high-intensity exercise. *PLoS One*. 2017;12(9):e0184240. doi:10.1371/journal.pone.0184240

13. Coco M, Di Corrado D, Calogero RA, Perciavalle V, Maci T, Perciavalle V. Attentional processes and blood lactate levels. *Brain Res*. 2009;1302:205-211. doi:10.1016/j.brainres.2009.09.032

14. Adamczyk JG, Gryko K, Boguszewski D. Does the type of foam roller influence the recovery rate, thermal response and DOMS prevention?. *PLoS One*. 2020;15(6):e0235195. doi:10.1371/journal.pone.0235195

15. Akinci B, Zenginler Yazgan Y, Altinoluk T. The effectiveness of three different recovery methods on blood lactate, acute muscle performance, and delayed-onset muscle soreness: a randomized comparative study. *J Sports Med Phys Fitness*. 2020;60(3):345-354. doi:10.23736/S0022-4707.19.10142-9

16. Yokoyama K, Watanabe, K. *Japanese Translation of POMS 2: Profile of Mood States*. Tokyo: Kaneko Shobo; 2015.

17. Saenz C, Heitman E, Luna F, Litewka S, Goodman KW, Macklin R. Twelve years of Fogarty-funded bioethics training in Latin America and the Caribbean: achievements and challenges. *J Empir Res Hum Res Ethics*. 2014;9(2):80-91. doi:10.1525/jer.2014.9.2.80

18. Reiner MM, Tilp M, Guilhem G, Morales-Artacho A, Konrad A. Comparison of a single vibration foam rolling and static stretching exercise on the muscle function and mechanical properties of the hamstring muscles. *J Sports Sci Med*. 2022;21(2):287-297. doi:10.52082/jssm.2022.287

19. Kasahara K, Yoshida R, Yahata K, et al. Comparison of the acute effects of foam rolling with high and low vibration frequencies on eccentrically damaged muscle. *J Sports Sci Med*. 2022;21(1):112-119. doi:10.52082/jssm.2022.112

20. Nakamura M, Konrad A, Kasahara K, et al. The combined effect of static stretching and foam rolling with or without vibration on the range of motion, muscle performance, and tissue hardness of the knee extensor. *J Strength Cond Res*. 2023;37(2):322-327. doi:10.1519/JSC.0000000000004263

21. Behm DG, Alizadeh S, Hadjizadeh Anvar S, et al. Foam rolling prescription: a clinical commentary. *J Strength Cond Res*. 2020;34(11):3301-3308.

doi:10.1519/JSC.0000000000003765

22.   Schneider DW. Alertness and cognitive control: interactions in the spatial Stroop
task. *Atten Percept Psychophys*. 2020;82(5):2257-2270. doi:10.3758/s13414-
020-01993-5

23.   Shevchenko Y. Open Lab: a web application for running and sharing online
experiments. *Behav Res Methods*. 2022;54(6):3118-3125. doi:10.3758/s13428-
021-01776-2

24.   Cohen, J. Statistical Power Analysis for the Behavioral Sciences. New York:
Routledg; 1988.

25.   Karvonen J, Vuorimaa T. Heart rate and exercise intensity during sports activities.
Practical application. *Sports Med*. 1988;5(5):303-311. doi:10.2165/00007256-
198805050-00002

26.   Riganas CS, Papadopoulou Z, Psichas N, et al. The rate of lactate removal after
maximal exercise: the effect of intensity during active recovery. *J Sports Med*
Phys Fitness. 2015;55(10):1058-1063.

27.   Menzies P, Menzies C, McIntyre L, Paterson P, Wilson J, Kemi OJ. Blood lactate
clearance during active recovery after an intense running bout depends on the
intensity of the active recovery. *J Sports Sci*. 2010;28(9):975-982.
doi:10.1080/02640414.2010.481721

28.   Warneke K, Aragão-Santos JC, Alizadeh S, et al. Are acute effects of foam-rolling
attributed to dynamic warm up effects? A comparative study. *J Sports Sci Med*.
2023;22(2):180-188. doi:10.52082/jssm.2023.180

29.   Bogdanis GC, Nevill ME, Lakomy HK, Graham CM, Louis G. Effects of active
recovery on power output during repeated maximal sprint cycling. *Eur J Appl*
Physiol Occup Physiol. 1996;74(5):461-469. doi:10.1007/BF02337727

30.   Hotfiel T, Swoboda B, Krinner S, et al. Acute effects of lateral thigh foam rolling
on arterial tissue perfusion determined by spectral Doppler and power Doppler
ultrasound. *J Strength Cond Res*. 2017;31(4):893-900.
doi:10.1519/JSC.0000000000001641

31.   Chmura J, Nazar K, Kaciuba-Uściłko H. Choice reaction time during graded
exercise in relation to blood lactate and plasma catecholamine thresholds. *Int J*
Sports Med. 1994;15(4):172-176. doi:10.1055/s-2007-1021042

32.   Pernigoni M, Kreivytė R, Lukonaitienė I, Toper CR, Kamandulis S, Conte D. Is

foam rolling as effective as its popularity suggests? A randomised crossover study
exploring   post-match   recovery   in   female   basketball. *J   Sports   Sci*.
2023;41(18):1718-1725. doi:10.1080/02640414.2023.2296718
33. Rey E, Padrón-Cabo A, Costa PB, Barcala-Furelos R. Effects of foam rolling as
a   recovery   tool   in   professional   soccer   players. *J   Strength   Cond   Res*.
2019;33(8):2194-2201. doi:10.1519/JSC.0000000000002277
34. Sullivan KM, Silvey DB, Button DC, Behm DG. Roller-massager application to
the hamstrings increases sit-and-reach range of motion within five to ten seconds
without performance impairments. *Int J Sports Phys Ther*. 2013;8(3):228-236.
524
525

Figure Captions

Figure 1. Intervention method for knee flexors and knee extensors.
Panels A and B show the intervention method for the knee flexors. A push-up bar is used,
with the sole on the non-intervention side in contact with the ground. A: shows the distal
part of the knee flexors, and B: shows the proximal part. FR intervention is performed as
shown from panel A to B. Panels C and D: the intervention for the knee extensors in which
the non-intervention knee is instructed to touch the ground. C: the distal part of the knee
extensors, and D: the proximal part. C to D: FR intervention is performed bilaterally for
60 s at a speed of 1 repetition every 2 s in the range specified for each muscle.

Figure 2. Example of Schneider's spatial Stroop task.

In the Congruent task (A), the displayed position and the direction of the arrow match as

shown. In the Incongruent task (B), the position displayed on the screen and the direction of

the arrow are mismatched, as shown.

Figure 3. Changes in lactate at pre, post, and 5 min after each condition.

*: Significant difference in pre-exercise conditions ($P < .01$); †: Significant difference in

post-exercise conditions ($P < .01$). FR, foam rolling.

Figure 4. Correlation between the changes in lactate and Stroop interference in the FR
condition.
The dotted lines indicate the regression lines for sexes; blue for males and orange for
females. FR, foam rolling.
553

555 Figure 5. Correlations between the change in lactate for the RPE leg and whole body in
556 FR and control conditions. FR, foam rolling; RPE, rating of perceived exertion.
557

561 Table 1. Participant characteristics (mean±SD)

Characteristic	Males (N=10)	Females (N=8)
Height (cm)	172.1±5.0	158.4±5.0*
Weight (kg)	67.1±8.4	48.1±5.3*
Age (years)	22.4±1.3	21.0±2.0
Thigh circumference (Right) (cm)	53.1±4.9	46.7±6.0*
Thigh circumference (Left) (cm)	53.2±4.3	47.1±5.9*

562 *: Significant differences were found for females compared to males ($P < .05$). SD,
563 standard deviation.

565 Table 2. Physiological parameters of each condition before (pre), immediately after
566 exercise (post), and after intervention (5 min) for each sex (mean±SD)

Conditions and variables	Males	Females
Control		
HR pre (bpm)	76.9±10.4	75.5±6.4
HR post (bpm)	181.3±11.9*	174.4±12.8*
RPE pre: leg / body (point)	6.4±0.7 / 6.5±0.7	7.9±2.2 / 8.1±2.3
RPE post: leg / body (point)	18.3±1.5*/ 16.8±2.1*	16.4±1.8*/ 15.8±1.6*
RPE 5 min: leg / body (point)	14.7±2.9*† / 13.0±3.4*†	12.5±2.2*† / 11.5±2.7*†
Total exercise time (sec)	651.0±87.0	405±92.5**
kp max (a.u.)	3.4±0.4	2.3±0.4**
Foam Rolling		

HR pre (bpm)	77.1±12.2	81.6±6.6
HR post (bpm)	181.6±16.2*	170.9±25.2*
RPE pre: leg / body (point)	6.5±1.0 / 6.6±1.0	7.8±2.2 / 8.4±2.3
RPE post: leg / body (point)	18.6±1.4*/ 17.2±2.6*	16.5±2.0*/ 15.1±2.7*
RPE 5 min: leg / body (point)	14.4±2.4*†/ 12.7±3.2*†	11.9±2.0*†/ 11.4±2.1*†
Total exercise time (sec)	615.5±138.5	396.6±91.7**
kp max (a.u.)	3.2±0.6	2.3±0.4**

567   * Significant difference in pre-exercise conditions ($P < 0.01$); † significant difference in

568   post-exercise conditions ($P < .01$); ** significant difference in males ($P < .01$). HR, heart

569   rate; RPE, rating of perceived exertion; SD, standard deviation.

570