Effects of foam rolling intervention on lactate clearance and cognitive function after high-intensity

exercise with incremental load

Kazuki Kasahara[1], Keita Oneyama[2], Takeru Ito[2], Masatoshi Nakamura[3], Genta Ochi[1,2]

1) Institute for Human Movement and Medical Sciences, Niigata University of Health and Welfare, Niigata,

Japan

2) Department of Health and Sports, Niigata University of Health and Welfare, Niigata, Japan

3) Faculty of Rehabilitation Sciences, Nishi Kyushu University, 4490-9 Ozaki, Kanzaki, Saga, 842-8585,

Japan

Corresponding Author:

Genta Ochi

Department of Health and Sports, Niigata University of Health and Welfare, Kita-ku, Shimami-cho, 1398,

Niigata, 950-3198, Japan, Telephone: +81-025-257-4595

Email: ochi@nuhw.ac.jp

ORCID:

Kazuki Kasahara: 0009-0008-7054-9855

Masatoshi Nakamura: 0000-0002-8184-1121

Genta Ochi: 0000-0003-1739-2023

Abstract: Foam rolling (FR) intervention has attracted attention in the sports field. However, its effects on

lactate clearance, mental fatigue, and cognitive function associated with fatigue after high-intensity exercise are

unknown. Therefore, this study examined whether FR intervention is an effective fatigue recovery tool.

Eighteen healthy adults (10 males and 8 females; 21.8 ± 1.7 years) without exercise habits participated in the

study. Participants performed high-intensity exercise on an ergometer with a progressive load. The FR

intervention postexercise was compared with the control condition. Measurements were lactate,

vigor/fatigue by POMS2, cognitive function (cognitive task performance), and leg and body rating of

perceived exertion (RPE) pre- and postexercise and postintervention (i.e., 5 min postexercise). This study

showed a significant reduction in the blood lactate concentration post-FR intervention. In addition, greater

lactate clearance by FR was correlated with faster recovery of executive function in those with greater

lactate clearance, although cognitive fatigue was not identified after high-intensity exercise. Moreover,

lactate clearance was not significantly correlated with RPE in the FR condition, whereas RPE decreased

with greater lactate clearance in those with greater lactate clearance in the control condition. The study

results indicate that FR intervention may be effective as a recovery tool for exhausting exercises. However,

when implementing it in the field, whether priority should be given to the recovery of physiological

parameters or RPE should be carefully considered.

Keywords: Foam Roller, Fatigue, POMS2, recovery

Acknowledgments

40 This work was supported by the Japan Society for the Promotion of Science (JSPS) Grant JP22K17739

41 (Genta Ochi) and the Grant-in-Aid for Scientific Research from the Niigata University of Health and

42 Welfare, 2023 (Genta Ochi).

INTRODUCTION

Many sports competitions involve high-intensity interval exercises with repeated sprints and jumps. Ball sports, such as basketball and handball, require high-intensity interval exercise for extended periods, with little time between halves or sets. High-intensity interval exercise, repeated with such short rest periods, results in a noticeable decrease in performance (fatigue) in the second half of the match. Factors contributing to exercise-induced fatigue could involve functional impairment at the peripheral and central levels, such as acidosis associated with lactic acid accumulation (Robergs, Ghiasvand and Parker, 2004), mental fatigue (Marcora, Staiano and Manning, 2009), and cognitive fatigue (Ochi et al., 2018; Ochi et al., 2022). Cook and Beaven (2013) recommended that recovery to a good psychological state post-exercise is important for maintaining performance, and recovery of both peripheral and central functions post-exercise is an urgent issue. However, existing recovery methods, such as cold water immersion (Higgins, Greene and Baker, 2017) and full-body aerobic exercise, e.g., rowing ergometer (Raeder et al., 2017), are difficult to implement during a game due to the time, location, and rules of the competition. To maintain mental and physical functions until the latter half of the match, a new recovery method that can be implemented in a limited space and in a short time is required.

Foam rolling (FR) is commonly used as a recovery method in several cases. Adamczyk et al. evaluated the effects of FR intervention after squat jumps at maximal effort for 1 min (Adamczyk, Gryko and Boguszewski, 2020). They reported significantly ($p < 0.05$) greater lactate clearance in the FR than in the

intervention condition at 30 min post-exercise. In addition, Akinci et al. implemented circuit-based high-

intensity exercise at 85% intensity calculated from the heart rate reserve method and measured lactate levels

before the exercise training, before the recovery period, and 5 and 20 min of the recovery period to examine

the effects of FR intervention on lactate levels (Akinci, Zenginler Yazgan and Altinoluk, 2020). They reported

that FR intervention did not alter lactate concentrations. However, these studies may not have adequately

evaluated the effects on muscles that are primarily active and fatigued during exercise because of the long

time elapsed after the exercise and the fact that it is a whole-body exercise. Moreover, the acute effects of

recovery methods on lactate clearance, mental fatigue, and cognitive fatigue, which have been implicated in

poor performance, are unknown. In addition, only a few studies have compared FR lactate reduction rates

between men and women, and the physiological conditions and methodologies involved in lactate reduction

rates remain unclear. This study was designed to determine whether FR improves lactate clearance, mood

decline, and cognitive fatigue after exhaustive exercise and whether gender differences exist in these effects,

thereby providing a foothold for elucidating the mechanism of FR recovery.

METHODS

Experimental setup

This study examined the immediate effects of FR, which can be performed in a confined space

and for a limited time, on lactate clearance and cognitive function after high-intensity exercise with

progressive loading. In addition, this study examined these effects separately in males and females. A

randomized repeated measures experimental design was used to compare the two conditions: control and

FR. Participants were instructed to visit the laboratory twice with a ≥48 h break. The measurement was

conducted before the intervention (pre), after the intervention (post), and 5 min postintervention. The

psychological scales completed included the Profile of Mood States 2nd Edition (POMS2) (Yokoyama and

Watanabe, 2015) and participant ratings of perceived exertion (RPE) (Saenz et al., 2014). The parameters

measured were executive function, psychological scale scores, and blood lactate concentration, which were

assessed respectively to prevent the blood lactate concentration measure from affecting executive function

and psychological status.

Participants

A total of 22 right-handed Japanese-speaking young adults (12 males and 10 females) agreed to participate from

October 2023 to January 2024. The inclusion criteria were as follows: no regular exercise habits, no daily use

of a foam roller, native Japanese speaker, and no history of neurological, psychiatric, or respiratory diseases.

All participants were instructed to refrain from eating or drinking anything other than mineral water 2 h before

starting the experiment. In this study, data analysis was conducted with 18 participants (10 males and 8 females)

because four participants were excluded because of their inability to perform the FR intervention after a high-

intensity exercise. Table 1 shows the participants' characteristics. Post hoc sensitivity analysis performed

using this sample with 80% power and $\alpha = 0.05$ demonstrated sufficient sensitivity to detect repeated-

measures effects exceeding $f = 0.31$ as computed using G*Power (3.1.9.2; The G*Power Team).

High-intensity exercise with a progressive load

High-intensity exercise with progressive loading was performed using an ergometer (828E, Monark,

Sweden). Participants performed an ergometer exercise for 3 min at a load of 0.5 KP as a warm-up, followed

by a 1-min rest period. After the break, males and females performed ergometer exercises with a progressive

increase in load of 0.5 KP/min. The participants performed the ergometer exercise and were instructed to pedal

in time with a metronome (Smart Metronome; Tomohiro Ihara, Japan) set at 60 bpm. The exercise was stopped

when the pedaling speed could no longer be sustained. During the exercise, strong verbal encouragement was

provided to elicit maximum effort. Exercise time and heart rate (HR) pre- and postintervention were measured

during each session.

FR intervention

Figure 1 shows the FR intervention methods using a foam roller (Stretch Roll SR-002, Dream

Factory, Umeda, Japan). A physical therapist instructed the participants at preintervention. FR intervention

was initiated with knee flexors followed by knee extensors. FR intervention was performed on the right and

then on the left side. The FR area of the knee flexor muscle group was defined as the proximal knee fossa

to the sciatic tuberosity (Reiner et al., 2022). The knee extensor was defined as the superior anterior iliac

spine to the top of the patella (Kasahara et al. 2022, Nakamura et al., 2022). One FR cycle was defined as one

distal rolling movement followed by one proximal rolling movement performed in 4 s (Behm et al., 2020). The

intervention intensity was directed to be the maximum intensity that the participant could tolerate (Kasahara et

al., 2022). The intervention time for each area was 60 s, and the participants were instructed to prepare for the

intervention for the next area during the 20-s interval. A metronome (Smart Metronome; Tomohiro Ihara,

Japan) at 60 bpm was used as a control.

Psychological measurements

POMS2 contains 35 items and evaluates seven mood states (anger-hostility, confusion-

bewilderment, depression-ejection, fatigue–inertia, tension–anxiety, vigor-activity, and friendliness). This

study used only the 10-item subset on vigor-activity and fatigue–inertia to reduce the participant's

psychological burden.

RPE was recorded during pre- and postexercise intervention to assess psychological exercise

intensity.

Blood lactate concentration

Lactate Pro 2 (AKRAY Europe B.V. Prof. J.H Bavincklaan 51,183 AT, Amstelveen, the

Netherlands), a handheld point-of-care analyzer, was used for enzymatic amperometric detection. It

requires 0.3 µl of a whole blood sample and a short time (15 s) to measure the lactate level. Lactate Pro 2

measurement ranges from 0.5 to 25.0 mmol/L. In this study, only lactate values between 0.5 and 25 were

included.

Behavioral measurements

Schneider's (2020) (Schneider, 2020) spatial Stroop task was used to measure executive function.

This task was created using web-building platforms (Lab.js v19.1.0) (Shevchenko, 2022). In each

experiment, participants made a spatially compatible key-press response to classify the spatial meaning of

a stimulus presented at an irrelevant position. Stimulus meaning and position were either matched

(congruent) or mismatched (incongruent) in the trial: 16 congruent and 16 incongruent. Visual alerting cues

(squares) were briefly presented at the two potential stimulus positions 500 ms before the onset of the

stimulus. Although previous studies have used multiple stimulus types (arrows or words) and spatial

dimensions (horizontal or vertical), this study employed arrow stimulus-type tasks. In addition, because

preliminary experiments revealed that reaction times tended to be slower in vertical than in horizontal

spatial dimension tasks, and the load on the task was assumed to be higher (fatigue was more likely to

occur); therefore, vertical spatial dimension tasks were used in the present study.

Participants pressed a key indicating up ("y") or down ("b") to classify the direction of an up- or

down-pointing arrow positioned above or below fixation (Figure 2). The correct answer ratio assigned to

yes and no was 50%. Each stimulus was separated by an interstimulus interval showing a fixation cross for

2 s to avoid prediction of the subsequent trial's timing. The stimulus remained on the screen until the patient

responded or for 1 s. This study adopted Stroop interference, a specifically defined cognitive process, to

elucidate the effects of an acute bout of exercise on executive function by calculating the incongruent–

congruent contrast, which is assumed to represent Stroop interference.

Data from 16 participants (8 males and 8 females) were used only for the cognitive task because data

for 2 of the 18 participants could not be obtained.

Statistical analysis

The R software (4.3.2), Rstudio (2023.12.0 + 369), "EZR on R Commander (Kanda, 2013)," and

the "anovaun" package were used for all analyses. A three-way analysis of variance (ANOVA) was

performed to compare the conditions, times, and sex. First, Mauchly's sphericity test was used to determine

whether sphericity was maintained. When the sphericity assumption was confirmed, a three-way ANOVA

with Greenhouse–Geisser's epsilon correction was performed; otherwise, only a three-way ANOVA was

performed. The effect size classification was set where $\eta_p^2 < 0.01$ was considered small; 0.02–0.1, medium;

and >0.1, large (Cohen, 1988). The corresponding *t*-test with Holm's correction was used to determine the

significant differences obtained from ANOVA. To clarify the relationships between the parameters and

executive performance, Pearson's correlation analyses were performed. Statistical significance was set a

priori at $P < 0.05$ for all comparisons.

Ethics approval

This study was conducted following the Declaration of Helsinki and was approved by the ethics

committee of the Niigata University of Health and Welfare, Niigata, Japan (approval number 19170-

231019). Participants were informed about their data confidentiality and provided written informed consent

to participate in the study.

RESULTS

Physiological parameters

Table 2 shows the results of HR, RPE, total exercise time, and kp max in the control and FR

conditions. No significant differences in PRE values were observed between the two conditions for all measures.

HR increased significantly immediately postexercise in both conditions ($P < 0.01$). RPE (leg/body) was

significantly higher at POST and 5 min compared with PRE in both conditions ($P < 0.01$) but significantly

lower at 5 min compared with POST ($P < 0.01$). In addition, sex differences were observed in the total

exercise time and load at postexercise ($P < 0.01$), which were significantly higher in males than in females.

However, no significant differences were observed between the conditions.

189    Figure 3 shows the changes in lactate values. A three-way ANOVA (conditions vs. time vs. sex)

190  showed no significant interactions (F (2, 32) = 1.0047, P = 0.3774, η_p^2 = 0.0591). A two-way ANOVA

191  (conditions vs. time) showed a significant interaction (F (2 ,32) = 4.0433, P < 0.05, η_p^2 = 0.2017) and a

192  main effect of time (F (2, 32) = 275.1504, P < .0001, η_p^2 = 0.9450). The post hoc test results in the control

193  condition showed significantly higher values at POST and 5 min than those at PRE (P < 0.01), but no

194  significant difference was observed between POST and 5 min. In the FR condition, POST and 5 min values

195  were significantly higher than PRE (P < 0.01). In addition, 5 min was significantly lower than POST (P <

196  0.01).

198  ***Psychological parameters***

199    A three-way ANOVA (conditions vs. time vs. sex) showed no significant interaction between vigor

200  (F(2, 32) = 1.6850, P = 0.2015, η_p^2 = 0.0953) and fatigue (F(2, 32) = 1.0022, P = 0.3783, η_p^2 = 0.0589) in

201  POMS2. A two-way ANOVA (conditions vs. time) showed a main effect of time on vigor (F(2, 32) = 7.5235,

202  P < 0.01, η_p^2 = 0.3198) and fatigue (F(2, 32) = 163.5833, P < 0.0001, η_p^2 = 0.9109). A post hoc test using

203  Holm correction revealed that vigor was significantly lower in POST than in PRE (P < 0.01). Fatigue was

204  significantly higher at POST and 5 min compared with PRE (P < 0.0001). In addition, POST was significantly

205  higher than 5 min (P < 0.0001).

 Relationship between parameters and performance

Figure 4 shows lactate and executive function changes in pre- and postintervention. A significant

correlation trend was found in the FR condition ($r = 0.41$, $P = 0.086$). Sex-specific results showed a significant

correlation for males ($r = 0.706$, $P < 0.05$), but not for females ($r = 0.13$, $P = 0.747$). No significant correlation

was found in the control condition ($r = -0.38$, $P = 0.118$).

Figure 5 reveals lactate and RPE (leg/body) changes pre- and postintervention. No significant

correlation was found in the FR condition, but a significant correlation was found in the control (leg: $r = 0.778$,

$P < 0.01$, body: $r = 0.669$, $P < 0.01$).

DISCUSSION

The acute effects of FR intervention on lactate clearance, psychological parameters, and cognitive

function after progressively increasing load exercise were examined, resulting in exhaustion to propose FR as

a new recovery method that can be implemented in a limited space in athletes. The results revealed that,

regardless of sex, the arterial blood lactate concentration, which increased after exhaustive exercise, decreased

in the FR condition, indicating that the FR intervention enhanced the lactate clearance effect. Interestingly,

cognitive function improved after FR intervention compared with that in those with greater lactate clearance,

and this effect was significantly greater in males than in females. This finding indicates that FR intervention

may also be useful in improving exercise-induced cognitive decline (cognitive fatigue), particularly in males.

This study showed significant increases in HR at pre- and postexercise in both FR and control

conditions, without significant differences between them. In addition, the Karvonen method (Karvonen and

Vuorimaa, 1988) demonstrated that exercise intensity at postexercise was $85.3 \pm 11.0\%$ in the FR condition and

$83.6\% \pm 10.4\%$ in the control condition, without a significant difference between the two conditions. Based on

these results, we concluded that all participants could perform the same extent of exercise in both conditions.

First, we examined the effect of FR intervention on lactate clearance. In the present study, the

ergometer exercise was performed with progressive loading by applying a strong load to the lower extremities,

and the FR intervention targeted the bilateral knee extensors and flexors. In this study, arterial blood lactate

concentrations significantly increased immediately at postexercise in both conditions, and no difference was

observed between the conditions. Thereafter, no lactate concentration changes were observed in the control

condition, but in the FR condition, arterial blood lactate concentration significantly decreased postintervention.

These results indicate that FR intervention immediately after exhaustive exercise is useful for increasing lactate

clearance. A previous study reported that increased arterial blood lactate concentrations after high-intensity

exercise were significantly lower in active recovery groups that performed low- or high-intensity exercise than

in passive recovery groups that did not receive any intervention (Riganas et al., 2015; Menzies et al., 2010).

More interestingly, no significant difference in the increasing effect of ROM was observed between the FR

and sham rolling conditions, which simulated the FR intervention movements (Warneke et al., 2023). These

findings indicate that active recovery, such as FR, may again increase the blood flow in active tissues and

increase lactate clearance (Bogdanis et al., 1996). In addition, FR intervention increases regional perfusion

(Hotfiel et al., 2017), indicating that FR intervention may further increase lactate clearance.

Second, the effects of FR intervention on executive function were examined. In the present study,

executive function task performance did not decrease at pre- or postexercise to exhaustion, and the progressive

load exercise leading to exhaustion did not induce cognitive fatigue. Progressive exercise to exhaustion has

long been associated with impaired cognitive function, including executive function (Chmura, Nazar and

Kaciuba-Uściłko, 1994); however, the effects are not always clear. Recently, Sudo et al. reported that the type

of cognitive task was not associated with cognitive performance after a high-intensity exercise (Sudo et al.,

2022). Therefore, exercise conditions such as progressive load exercise and the spatial Stroop task possibly

failed to induce cognitive decline in the study participants. Conversely, exercise in extreme environments, such

as hypoxic environments, decreases cognitive function (Ochi et al., 2018; 2022). To verify whether FR

intervention is effective in restoring exercise-induced cognitive decline, it should be investigated under different

exercise conditions, such as a hypoxic environment.

This study demonstrated that the amount of lactate change was significantly correlated with changes

in executive function task performance immediately postexercise and post-FR intervention in the FR condition.

This result indicated that executive function may be recovered in patients with increased lactate clearance.

Previous studies have reported that elevated blood lactate levels after exhaustive exercise are associated

with a significantly worsened attentional process (Coco et al., 2009). Therefore, increasing lactate clearance

due to FR may contribute to improved cognitive decline (cognitive fatigue) due to increased lactate values.

In addition, previous studies have shown that low- and moderate-intensity exercises improve Stroop task

performance (Byun et al., 2014; Yanagisawa et al., 2010). The exercise-induced increase in body perfusion

(Mulser and Moreau, 2023) may increase nutrient (glucose and oxygen) supply to the brain. Although the

relationship between cognitive improvement and increased brain perfusion by exercise remains unclear, the

increase in whole-body perfusion due to FR intervention may contribute to improved executive function.

Finally, we examined the effects of FR intervention on psychological parameters. These results indicated that

FR intervention did not affect the vigor and fatigue of POMS2. The results support the findings of a previous

study (Pernigoni et al., 2023) in which FR intervention was administered to female basketball players and had

no effect on perceived recovery (TQR) or fatigue level. However, in professional male soccer players, a larger

effect size (Conhen's d = 1.08) was reported on TQR recovery in the FR intervention condition compared with

the control condition after 24 h of training (Rey et al., 2019). These differences in results may be due to the

methods of FR intervention. Pernigoni et al. (2023) and Ray et al. (2019) performed a 20-min FR

intervention on the entire lower extremity, including the quadriceps, hamstrings, triceps, and abductor and

adductor muscle groups. Conversely, this study found that the FR intervention was 4 min for knee extensors

and knee flexors. A capacitance–response relationship has been found for the effect of FR on ROM increase

(Sullivan et al., 2013). The enhancement of psychological parameters through FR may also be associated with

the capacity–response relationship, necessitating further research for a clearer understanding.

The results revealed that in terms of RPE, which represents the tightness to exercise, a greater

decrease in lactate in the control condition was associated with a lower RPE. However, in the FR condition, no

significant correlation was observed between changes in lactate and RPE. These differences may be related to

the FR intervention method. A push-up bar was used as the FR intervention method for the knee flexors (Figure

1A, 1B). Furthermore, intervention for the knee extensors was performed in a plank posture (Figure 1C, 1D).

In this study, the intervention effects were examined in participants who did not use the foam roller daily;

therefore, the FR intervention itself may have affected RPE. Therefore, while lactate clearance increased in the

FR condition, the effect of improved tightness to exercise may not have occurred. The on-field implementation

of FR interventions requires careful consideration of whether priority should be given to the recovery of

physiological parameters or the recovery of subjective fatigue levels.

There are limitations to this study. First, the mechanisms underlying the study results are unknown.

The results indicate that FR intervention increases lactate clearance and has a concomitant effect on executive

function; however, the detailed mechanism by which this occurs is unknown. Future studies should examine

the effects of FR intervention on regional and cerebral perfusion and changes in lactate clearance and cognitive

function. The second is the target population. As mentioned above, the FR intervention for male athletes showed

fatigue recovery effects (Rey et al. 2019), but not for female basketball players (Pernigoni et al., 2023).

Therefore, the differences between those who have exercise habits and older adults, as well as the differences

in athletic specialties and sex among athletes, should be examined in the future. Finally, the effect of FR

intervention on exercise-induced cognitive decline (cognitive fatigue) is discussed. In this study, a correlation

was observed between increased lactate clearance and improved cognitive task performance, but cognitive

fatigue was not observed in the participants. Using exercise conditions in which cognitive fatigue occurs and

accumulating verification, FR intervention can be proposed for the competitive scene as a contribution to the

improvement of not only peripheral but also central fatigue.

CONCLUSION

The results revealed that lactate clearance increased as an acute effect of FR intervention after

exhaustive exercise in healthy young adults. Furthermore, the greater the effect, the greater the recovery of

executive function. However, FR intervention does not decrease RPE by exhaustive exercise; therefore, careful

consideration should be given to whether recovery of physiological parameters or recovery of RPE should be

prioritized when implementing FR intervention in the field.

Conflict of Interest

The author declares that the author has no conflict of interest.

Data availability statement

The datasets generated and analyzed during the current study are not publicly available but are available

from the corresponding author, who was an organizer of the study.

316 Figure 1. Figures 1A and B show the intervention method for the knee flexors. A push-up bar was used and the sole of the non-intervention side was instructed to

317 contact the ground. 1A shows the distal part of the knee flexors, and 1B shows the proximal part. FR intervention was performed in the A-B range. Figures 1C and

318 1D show the intervention for the knee extensors, where the non-intervention knee was instructed to touch the ground. 1C shows the distal part of the knee extensors

319 and 1D shows the proximal part, and FR intervention was performed in the range of C-D. The FR intervention was performed bilaterally for 60 seconds at a speed of

320 1 repetition per 2 seconds in the range specified for each muscle.

(A) Congruent task (B) Incongruent task

Figure 2. This is an example of Schneider's spatial Stroop task. In the Congruent task, the displayed position and the direction of the arrow match as shown in A. On the

other hand, in the Incongruent task, the position displayed on the screen and the direction of the arrow are mismatched, as shown in B.

Figure 3. The changes in lactate at PRE, POST, and 5 min after each condition are shown.

*: significant difference with PRE ($P < .01$)

†:Significanct difference with POST ($P < .01$)

Figure 4. The correlation between the changes in lactate and Stroop interference in the FR condition are shown in the figure. The dotted lines
indicate the regression lines for sexes, blue for males and orange for females.

Figure 5. The correlations between the change in lactate and RPE leg and whole body in FR and control conditions are shown separately.

Table 1.

The following table shows the characteristics of the 18 participants (10 males and 8 females) in this

study (mean±SD).

Characteristic	Males (N=10)	Females (N=8)
Height (cm)	172.1±5.0	158.4±5.0*
Weight (kg)	67.1±8.4	48.1±5.3*
Age (years)	22.4±1.3	21.0±2.0
Thigh circumference (Right) (cm)	53.1±4.9	46.7±6.0†
Thigh circumference (Left) (cm)	53.2±4.3	47.1±5.9†

333   *: Significant difference with males ($P < .01$)

334   †: Significant difference with males ($P < .05$)

335 Table 2.

336 Physiological parameters of each condition before (PRE), immediately after exercise (POST), and

337 after intervention (5min) are shown for each sex (mean±SD).

Conditions and variables		males	females
Control			
	HR^a pre (bpm)	76.9±10.4	75.5±6.4
	HR^a post (bpm)	181.3±11.9*	174.4±12.8*
	RPE^b pre: leg / body (point)	6.4±0.7 / 6.5±0.7	7.9±2.2 / 8.1±2.3
	RPE^b post: leg / body (point)	18.3±1.5*/ 16.8±2.1*	16.4±1.8*/ 15.8±1.6*
	RPE^b 5min: leg / body (point)	14.7±2.9*† / 13.0±3.4*†	12.5±2.2*† / 11.5±2.7*†
	Total exercise time (sec)	651.0±87.0	405±92.5**
	kp max (a.u.)	3.4±0.4	2.3±0.4**
Foam Rolling			
	HR^a pre (bpm)	77.1±12.2	81.6±6.6
	HR^a post (bpm)	181.6±16.2*	170.9±25.2*
	RPE^b pre: leg / body (point)	6.5±1.0 / 6.6±1.0	7.8±2.2 / 8.4±2.3
	RPE^b post: leg / body (point)	18.6±1.4*/ 17.2±2.6*	16.5±2.0*/ 15.1±2.7*
	RPE^b 5min: leg / body (point)	14.4±2.4*† / 12.7±3.2*†	11.9±2.0*†/ 11.4±2.1*†
	Total exercise time (sec)	615.5±138.5	396.6±91.7**
	kp max (a.u.)	3.2±0.6	2.3±0.4**

338 ^{a}HR: heart rate

339 ^{b}RPE: Rating of Perceived Exertion

340 *: Significant difference with PRE ($P < .01$)

341 †: Significant difference with POST ($P < .01$)

342 **: Significant difference with males ($P < .01$)