Nuclear pore complexes undergo Nup221 exchange during blood stage asexual replication of *Plasmodium* parasites.

Authors:

James Blauwkamp[1]*, Sushma V Ambekar[2]*, Tahir Hussain[2], Gunnar R Mair[2,3#], Josh R. Beck[2#], Sabrina Absalon[1#]

Affiliations:

[1]Indiana University School of Medicine, Department of Pharmacology and Toxicology

[2]Iowa State University, Department of Biomedical Sciences

[3]School of Biological Sciences, Queen's University Belfast

*These authors contributed equally

#Co-corresponding Authors

Abstract

Plasmodium parasites, which are the causative agents of malaria, undergo closed mitosis without

breakdown of the nuclear envelope. Unlike the closed mitosis in yeast, *P. berghei* parasites

undergo multiple rounds of asynchronous nuclear divisions in a shared cytoplasm result in a

multinucleated (8-24) organism prior to formation of daughter cells within an infected red blood

cell. During this replication process, intact nuclear pore complexes (NPCs) and their component

nucleoporins are likely to play critical roles in parasite growth, facilitating selective bi-directional

nucleocytoplasmic transport and genome organization. Here we utilize ultrastructure expansion

microscopy (U-ExM) to investigate *P. berghei* Nup138, Nup221, and Nup313 at the single nucleus

level throughout the 24 hour blood-stage replication cycle. Our findings reveal that these Nups

are evenly distributed around the nuclei and organized in a rosette structure previously

undescribed around the centriolar plaque, which is responsible for intranuclear microtubule

nucleation during mitosis. We also detect an increased number of NPCs compared with previously

reported, highlighting the power of U-ExM. By adapting the recombination-induced tag exchange

(RITE) system to *P. berghei*, we provide evidence of NPC maintenance, demonstrating Nup221

turnover during parasite asexual replication. Our data shed light on the distribution of NPCs and

their homeostasis during the blood-stage replication of *P. berghei* parasites. Further studies into

the nuclear surface of these parasites will allow for a better understanding of parasites nuclear

mechanics and organization.

Introduction:

The survival and growth of eukaryotic organisms relies on the movement of macromolecules between the membrane bound nucleus and the cytosol. This nuclear-cytosolic transport is facilitated by nuclear pore complexes (NPCs) that are embedded into the nuclear envelope. Apart from being a channel for selective bidirectional nucleocytoplasmic transport, the NPC plays a key role in chromatin organization, gene regulation, DNA repair, and maintenance of epigenetic memory in many eukaryotes [1]. Typically, these NPCs are composed of multiple copies of more than 30 different nucleoporins (Nups) per NPC, with eukaryotic NPCs containing around 1000 proteins total [1-5]. Nups are widely conserved in most eukaryotes, including yeasts, humans, and various plant species, and have also been identified in parasitic protozoa such as *Trypanosoma brucei* and *Toxoplasma gondii* [6, 7]. Nups are separated into two distinct categories; structural Nups that form the pore and the intrinsically disordered phenylalanine-glycine repeat (FG) Nups which regulate nucleocytoplasmic transport (Figure 1a) [8, 9]. Structurally, the NPCs establish an inner channel called the inner ring, flanked by the outer rings on the cytosolic and nuclear side of the NPC. After the inner channel is constructed the cytoplasmic filament and the nuclear basket Nups attach to their appropriate side of the pore [8, 9]. The FG Nups, characterized by their FG repeats, occupy the space inside the inner ring and interact with the cargo to facilitate nuclear-cytosolic transport [8-10].

One of the most extensively studied organisms with respect to the dynamics and structural aspects of nuclear pore complexes is budding yeast. *Saccharomyces cerevisiae* undergoes closed mitosis during which the nuclear envelope remains intact during karyokinesis. In contrast to open mitosis, where NPC removal during prophase leads to nuclear envelope

breakdown, in closed mitosis it has been hypothesized that specific regions of nuclear pore/

complex disassembly cause small breaks in the nuclear envelope, facilitating nuclear envelope

fission and formation of two new nuclei [11, 12]. Like budding yeast, the parasites of the

Plasmodium genus, the causative agent of malaria, undergo a specialized form of closed mitosis.

In these parasites, multiple rounds of asynchronous nuclear division in a shared cytoplasm leads

to a multinucleated single-celled parasite [13]. While many Nups are conserved through

eukaryotes, *Plasmodium* Nups are very divergent, which makes their identification by

bioinformatics challenging. In a 2010 study aiming to identify Nups of different eukaryotes, only

two Nups were identified in *P. berghei* compared to 19 identified Nups in over 50 different

organisms [14]. Eleven Nups are currently known in the rodent malaria model *P. berghei*; Sec13.

Five FG Nups, and five novel Nups identified using Nup313 as bait in proximal labeling assays

(Figure 1a) [15-17]. Interestingly, only one of these proteins presents established structural

features of eukaryotic Nups [8].

The limited understanding of essential components of nuclear pore complexes in

Plasmodium is attributed to the absence of protein homology for Nup identification in

Plasmodium and the lack of suitable tools to investigate the dynamics of NPCs within relatively

small nuclei (a micrometer). While conventional fluorescence and electron microscopy

techniques have provided valuable insights into the perinuclear localization of identified Nups in

Plasmodium parasites, these approaches suffer from low resolution and the significant financial

and time investments required for electron microscopy [8, 11, 15, 18-22]. These challenges in

comprehending the dynamics and distribution of nuclear pore complexes (NPCs) in *Plasmodium*

parasites have spurred the exploration of innovative microscopy techniques to overcome these

limitations and gain deeper insights into these vital components. The adaptation of Ultrastructure

Expansion Microscopy (U-ExM) to work with parasites now allows for super-resolution imaging

of parasites in a cost-effective and high-throughput manner [23-26]. U-ExM achieves isotropic

expansion of biological samples, increasing their size up to 4.5 times by embedding denatured

samples into a hydrogel that expands when exposed to water. Notably, this technique has been

employed to achieve sub-nucleus resolution and has successfully visualized Nup313 at the

nuclear membrane in *P. falciparum* [23, 25].

In this study, we employed U-ExM to visualize individual NPCs using tagged versions of *P.*

berghei Nup138, Nup221, and Nup313. We successfully localized these Nups to the nuclear

envelope by employing NHS ester or BodipyFL Ceramide staining throughout the asexual life cycle

of the *P. berghei* parasite. Consistent with prior findings using "serial surface" imaging in focused

ion beam – scanning electron microscopy (FIB-SEM)[21], we observed an increase in NPC number

until the parasite's first mitosis, during which the NPCs were then distributed between the newly

developing nuclei. We proceeded to assess NPC assembly and protein turnover using the

Recombination Induced Tag Exchange (RITE) system marking the first time this has been

accomplished in *Plasmodium*. The RITE system uses a conditional, hormone activated Cre-

recombinase to generate permanent epitope tag switch in a target coding sequence. This method

has been previously utilized in human cells to observe the dynamic replacement of NPC

components in human neurons [27, 28].

Leveraging the enhanced resolution of U-ExM coupled with the RITE system, we were able

to monitor NPC assembly and maintenance dynamics within a single nucleus during *Plasmodium*

replication. In line with previous observations in mammalian cells, we found that NPCs in *P.*

berghei undergo protein maintenance after the NPCs have been produced. Remarkably, we also

confirmed the presence of Nup221 proteins of different ages in a single NPC by colocalization of

original and induced Nup221 signal. Our results expand upon existing data and showcase the new

insights achievable through the combination of U-ExM with genetic and biochemical systems.

Results:

U-ExM visualization of Nups around the nucleus.

Previous studies on NPCs in various organisms have predominantly relied on slide-based

immunofluorescence assays. For higher resolution, NPCs could be visualized by electron

microscopy, but this process is time consuming and costly. The emergence of expansion

microscopy in smaller organisms like yeast has significantly advanced NPC visualization and study

[29]. In our efforts to visualize nucleoporins (Nups) at single NPC resolution in *P. berghei*, we

introduced C-terminal endogenous epitope tags on three previously identified *P. berghei* FG Nups:

spaghetti monster Hemagglutinin (smHA) tags on Nup138 and Nup313 and a 3xHA/GFP tag on

Nup221 (Supplemental figure 2). In comparison to previous studies on unexpanded parasites [15,

17], and considering the expected size of the NPCs in yeast [29], expansion microscopy

successfully resolves individual NPCs around the nuclear periphery (Figure 1b). To confirm the

position of these Nups at the nuclear membrane, we employed different approaches, visualizing

the nuclear membrane using an endoplasmic reticulum (ER) antibody (PfBIP), employing

membrane stains such as Bodipy TR Ceramide and Bodipy FL Ceramide, and assessing membrane

localization through NHS Ester staining variations around the nucleus (Supplemental figure 1).

Among these methods, we determined that differences in NHS ester staining and high

concentrations of Bodipy FL Ceramide effectively delineated the nuclear envelope to confidently

assign the NPC signal to this structure. In summary, U-ExM allows us to observe individual NPCs

on the nuclear envelope.

***Plasmodium berghei* produces NPCs primarily before mitosis**.

We next wanted to quantify the number of NPCs surrounding each nucleus during various

stages of the parasite's intraerythrocytic development cycle. To achieve this, we performed U-

ExM on parasites that exhibited a range of nuclear numbers and were stained for either

Nup138::smHA or Nup313::smHA. In the early stages of the parasite's life cycle, before the

parasites begin their mitotic phase (1n, Ring), we observed Nup138 being uniformly distributed

around the nucleus (Figure 2a), while the Nup313 signal is concentrated at either end of the

nucleus (Supplemental figure 3). At this stage, we noticed a higher number of Nup313 foci

(approximately 40-50 NPCs) compared to Nup138 (around 25-35 NPCs) (Figure 2b). As the

parasites began mitosis (1n, Trophozoite premitotic and mitotic), there was a significant 5-fold

increase in both Nup138 and Nup313 numbers. The parasites produce over 250 NPCs containing

Nup313 and over 150 NPCs containing Nup138, distributed around the single nucleus (Figure 2b).

As more nuclei are formed (2-4n Schizont, 5-9n Schizont, 10+n Cytokinesis), the number of NPCs

around each nucleus decreased. Initially, approximately 250 NPCs surrounded a single nucleus,

which then reduced to around 90-100 NPCs around each of the 2 or 3 nuclei. Eventually, the count

further decreases to approximately 25-30 NPCs around 10-12 nuclei (Figure 2b). Our data

suggests that NPC production peaks before mitosis and NPCs are then distributed during each

karyokinesis.

147 *Plasmodium berghei* **NPCs are organized around the centriolar plaque.**

148   During and after the first mitosis we observe a uniform distribution of NPCs around

149 individual nuclei, as visualized by Nup138 and Nup313 (Figure 2). Interestingly, upon the first

150 round of mitosis we detect 4-10 foci of Nup138, Nup313, and Nup221, the homolog to yeast

151 Nup100, localizing around the centriolar plaque (CP), which is the nuclear microtubule

152 organization center responsible for intranuclear microtubule nucleation (Figure 3a)[25, 30, 31].

153 Quantification of NPCs around the centriolar plaque reveals a higher count of Nup313 compared

154 to the other two FG Nups used in this study (Figure 3b). It is important to highlight that after

155 parasite cytokinesis, the centriolar plaque disassembles [25], but NPCs retain a rosette formation

156 in the vicinity of the former plaque location (Figure 3c).

157 **The RITE system reveals NPC assembly and maintenance.**

158   NPCs in yeast undergo degradation and replacement of Nups and full NPCs [32-34], and

159 that Nup replacement occurs in non-dividing human cells [27]. Therefore, in order to assess the

160 turnover of Nups in *Plasmodium*, we adapted the Recombination Induced Tag Exchange (RITE)

161 system for the first time in these parasites. The RITE system uses a constitutively expressed Cre

162 recombinase fused to the human estrogen binding domain (EBD) which retains the protein in the

163 cytoplasm by association with HSP70. Upon the addition of β-estradiol, Cre-EBD translocates to

164 the nucleus where it triggers a permanent genetic switch through a recombination event between

165 two *LoxP* flanking the initial tag, removing this tag and bringing a distinct tag into frame [28]

166 (Supplemental figure 4a). Here we tagged *P. berghei* Nup221 with a RITE system cassette, which

167 includes a 3xHA/GFP tag, a 3xMyc/RFP tag, and the two *LoxP* sites for tag switching (Figure 4a).

We found that adding as little as 2 nM β-estradiol resulted in a partial switch from HA to Myc tag,

while 200 nM β-estradiol induces sufficient relocation into the nucleus for a complete switch from

HA to Myc tag in as little as 2 hours, confirmed by PCR analysis (Supplemental figure 4b, c).

Initial immunofluorescence imaging over the course of a 16-hour period for both control

and 200 nM β-estradiol treated parasites showed visible tag switching by eight hours post

induction with β-estradiol (Supplemental figure 5a). By the end of the 16-hour period we

observed individual parasites (merozoites) that expressed a mixture of Nup221 tagged with GFP

or RFP, both in a typical polar localization. By PCR genotyping, we can detect tag exchange within

two hours of estrogen addition. Notably, the continued detection of Nup221-HA through 16 hours

post induction indicates a stable population of Nup221 persists for at least eight hours during the

parasite life cycle (Supplemental figure 5). To validate complete turnover of Nup221 and ensure

no spurious excision was occurring, samples from control and induced conditions were allowed

to propagate for an additional 96 hours (four parasite development cycles). While Myc-tagged

Nup221 was never detected in control samples, HA-tagged Nup221 was no longer detectable in

β-estradiol treated parasites after 96 hours. These results demonstrate that the RITE system is an

effective tool for monitoring Nup221 turnover in *P. berghei.* Pairing this technique with U-ExM

will provide better resolution for determining NPC maintenance in the parasites.

***Plasmodium berghei* replaces Nups in post-mitotic NPCs.**

To investigate NPC assembly and maintenance during and after mitosis, we treated newly

invaded parasites (1N) with 200 nM β-estradiol and cultured for 20 hours. We then collected

processed samples for U-ExM at 0, 10, and 20 hours post induction with β-estradiol to determine

Nup221 presence in NPCs before mitosis occurs (10 hours post induction) and after mitosis,

during cytokinesis (20 hours post induction). We hypothesized that if Nup221 is being replaced in

the post-mitotic NPCs, we would observe a signal shift from the original HA signal (pink) to the

induced Myc signal (yellow), with a higher proportion of induced signal in the 20 hours post

induction samples.

At 0 hours post induction, early-stage parasites display only the original HA signal (Figure

4b). At 10 hours post induction, parasites mature and undergo the first rounds of mitosis resulting

in the formation of 2-3 nuclei. These parasites exhibit a mix of the HA and Myc signals, with 50%

of foci containing only HA signal, 35% of foci containing only Myc signal, and 15% of foci

containing both the HA and Myc signals (Figure 4b, c). The presence of co-localized HA and Myc

signals suggests multiple copies of Nup221 in individual NPCs. By 20 hours post induction with β-

estradiol the parasites have stopped producing new NPCs. At this time we found 35% of NPCs

contain only HA signal, while 50% of NPCs contain only Myc signal and 15% of NPCs contain both

HA and Myc signal (Figure 4b, c). This result suggests novel Nup221-Myc incorporation into

existing NPCs after the parasites have stopped producing them.

Subsequently, we wondered whether the Nups in the NPCs around the CP were subjected

to maintenance at the same rate as those around the nuclei. To test this, we specifically

monitored the turnover of Nup221 within the NPCs surrounding the CP. At 0 hours post induction,

when no CPs are visible, Nup221 is evenly distributed around the parasite nucleus. By 10 hours

post induction the CP is present, and now 50% of the NPCs surrounding it contained only the

original signal, mirroring observations around the rest of the nucleus. Around 20% of the NPCs

contain only the induced signal, a decrease of 15% compared to around the ones around the

nucleus, and 30% of the NPCs contain both the original and induced signal, an increase of 15%

compared to those scattered around the nucleus (Figure 4d). This suggests that the turnover of

the original Nup221 signal occurs at a slower pace in the NPCs surrounding the nucleus compared

to those scattered around it. This trend persists at 20 hours post induction, where 30% of NPCs

showed only the original signal, similar to what was seen around the rest of the nucleus. However,

45% of the NPCs contain only the induced signal (a decrease from 50%) and 25% of the NPCs

contain both the original and induced signal (an increase from 15%) (Figure 4d). Altogether, our

data indicate that proteins in the NPCs surrounding the CP exhibit greater stability than those in

NPCs scattered around the nucleus.

Discussion

In this study, we applied U-ExM and the RITE System to investigate the distribution and

homeostasis of NPCs in the asexual blood stages of the rodent malaria model, *P. berghei*. Previous

work on Nups in *Plasmodium* presented challenges due to both the divergent nature of the Nups

and the relatively small size of the parasite nucleus at approximately 1 μm. Our results

demonstrate the effectiveness of U-ExM in resolving individual NPCs around the parasite nucleus,

enabling us to address questions about their individual characteristics. Additionally, for the first

time we adapted the RITE system in *Plasmodium,* allowing us to dissect the mechanisms of NPC

maintenance. While the RITE system was employed here to monitor protein turnover by tag

exchange, it should also be amenable to other conditional mutagenesis strategies, providing an

alternative conditional excision strategy in addition to the established dimerizable Cre (DiCre)

system [35, 36], expanding the genetic toolkit in this important rodent malaria model.

This study defined the number of Nups surrounding the nucleus during the asexual life

cycle of *P. berghei*, unraveling a dual localization of NPCs around individual nuclei. Interestingly,

we observed a rosette structure around the centriolar plaque (CP) during nuclear replication and

this NPC organization persisted in merozoites even when the CP was no longer visible. Our

findings align with Ferreira et al, using CryoEM to examine microtubules in *P. falciparum* parasites,

revealing a similar ordered NPC structure around the CP during mitosis [37]. This circular

organization, previously unreported and not observed outside of *Plasmodium*, could suggest a

specific role of these NPCs in CP formation, duplication, maintenance, and function during

mitosis. For example, *S. cerevisiae* NPC protein Ndc1, a subunit of the transmembrane ring, is a

key protein in inserting the nuclear MTOC into the nuclear envelope [38, 39].

By quantifying individual Nup foci around the nuclei, we determined the number of NPCs

per nucleus throughout the life cycle. Our results indicate a rapid increase in NPCs just before the

first round of mitosis, consistent with previous observations *in P. falciparum* [21]. The higher NPC

count observed in our study (250 vs 60) might be attributed to greater sample numbers

achievable with U-ExM compared to FIB-SEM (our data was taken from 150+ nuclei from 60+ cells

over two biological samples) and improved NPC detection by immunostaining against the

endogenously tagged Nups.

Previous research in various cell types has shown that the composition of NPCs at specific

stages in the cell cycle can influence cell fate [40, 41]. In *Plasmodium* parasites it has been shown

that SEC13, a component of the nuclear pore complexes and the COPII coat, show a different

localization than FG Nups [15]. Our study delved into the heterogeneity of NPCs by examining

three FG-Nups: Nup138, Nup221 and Nup313. Notably, Nup313 showed a distinct localization in

the early ring stage of the parasite suggesting a potential specialized role during this stage of

development. Further investigations into the dynamic localization of different types of Nups

would be interesting to assess whether Nup313's temporal differential localization is unique or

shared with other Nups that compose the NPC.

Our data indicate that most NPCs contain a single version of Nup221, whether pre-mitotic

(Figure 4c – Nup221-HA, pink) or post-mitotic (Figure 4c – Nup221-Myc, yellow). However, some

NPCs exhibited both pre- and post-mitotic proteins, suggesting a gradual turnover in response to

protein degradation or damage (Figure 4c). These observations echoed findings using the RITE

system by Toyama et al., suggesting an active mechanism for NPC maintenance in mammalian

cells [27]. This could also point towards the octagonal rotational symmetry that has been seen in

other organisms [42, 43]. Interestingly, we note a slight increase in protein stability around the

CPs, illustrated by higher colocalization between Nup221-HA and Nup221-Myc proteins than

around the rest of the nucleus (Figure 4d), suggesting a higher longevity of this FG Nup in NPCs

rosette compared to those scattered across the nucleus.

With the identification of numerous Nups in the *Plasmodium* genome [15, 17], and

improved tools for visualizing NPCs, we are now poised to investigate the individual functions of

Nups. While the exact roles of specific Nups in the *Plasmodium* parasites remains unknown,

studies in other organisms have illuminated the involvement of FG Nups in various cellular

processes such as protein synthesis [44], NPC localization and composition [45, 46], as well as

RNA transport [6, 47]. While no studies of specific FG Nup function have been conducted in

Plasmodium parasites, deletions of Nup138 and Nup221 in genome wide screens and individual

knock-out attempts have shown all Nups to be essential [48-50]. Likewise, partial knockdown of

the Nup SEC13 in human malaria slows growth of the parasite in the blood [16], while a C-terminal

proline enriched sequence from SEC13 in mouse malaria models is required for parasite growth

[15]. Thus, it is likely that disruption of Nups can lead to growth delays and impaired cell viability

in malaria parasites. Our work, along with prior studies, lays the foundation for future research

into these divergent Nups as techniques for efficient knockdown continue to advance in malaria

parasites.

One of the hallmarks of nuclear envelope breakdown in mammalian open mitosis is the

disassembly of NPCs following mitotic kinase signaling [51, 52]. This disassembly is a decisive step

where multiple Nups are phosphorylated and sequestered to be used upon nuclear envelope

reassembly [53]. In contrast, *Plasmodium* parasites, like yeast, undergo closed mitosis, where the

nuclear envelope remains intact during nuclear division [13]. Recent work in fission yeast has

used confocal microscopy [11] and expansion microscopy [29] to look at the distribution and

dynamics of NPCs in a closed mitosis model. These studies identified NPCs at the bridge between

dividing nuclei before nuclear envelope fission. The authors hypothesize that through currently

unknown mechanisms, NPC disassembly in the bridge between dividing nuclei contributes to

nuclear envelope disassembly, enabling nuclear division. While we observed instances of dividing

nuclei with a cluster of NPCs along this bridge, further investigation is needed to explore this

mechanism in *Plasmodium* parasites. Identification and characterization of a cytoplasmic facing

Nup will allow for this hypothesis to be studied further, and potentially probe into the

mechanisms that determine where NPC disassembly occurs and its link to nuclear fission.

In conclusion, our study provides novel insights into the distribution, homeostasis, and

homogeneity of NPC complexes in *P. berghei* throughout the asexual life cycle. Understanding the

dynamics of these essential components is crucial, as NPCs play a vital role in parasite survival.

The methodologies employed here pave the way for further in-depth investigations into the

functions of these divergent complexes, facilitating a comprehensive understanding of their role

in *Plasmodium* biology. This study sets the stage for future investigations into the diverse Nups in

the *Plasmodium* genome and their individual functions. Disruption of Nups, whether structural

or FG Nups, likely impacts cell growth and viability in malaria parasites, as observed in related

studies. As techniques for knockdown efficiency improvement continue to evolve, exploring the

specific functions of Nups in *Plasmodium* is imminent.

Figures:

Figure 1: Divergent *P. berghei* NPCs can be analyzed by ultrastructure expansion microscopy (U-

ExM). **a.** Schematic representation and table of known nucleoporins (Nups) in yeast, *P.*

falciparum, and *P. berghei*. Nups highlighted in magenta indicate Nups visualized in this study,

Nups marked with an asterisk represent homologs, Nup637 predicted Nup. **b.** Schematic

representation of the *P. berghei* life cycle inside of host red blood cells, showing NPC distribution.

Parasites begin with a single nucleus that undergoes mitosis midway through the life cycle.

Multiple rounds of asynchronous mitosis lead to a multinucleated parasite with a single

cytoplasm before a shared cytokinetic event produces individual parasites. Also shown are

comparisons between unexpanded and U-ExM images of Nup138-smHA and Nup313-smHA

(magenta) through the life cycle showing differences in these microscopy techniques. Under the

U-ExM images lists the number of nuclei in the parasites shown. Scale bars 2 μm, number on

images indicates image depth in μm.

Figure 2: *P. berghei* produces NPCs until mitosis, distributing them among replicated nuclei.

a. U-ExM images showing Nup313-smHA at early stages and Nup138-smHA through the life cycle of *P. berghei*. Microtubules shown in yellow, Nup138-smHA shown in magenta, DNA shown in blue, and protein density shown in grayscale. Scale bars 2 μm, image depth in μm indicated. **b.** Number of NPCs per nucleus through the *P. berghei* life cycle calculated by counting Nup138-smHA and Nup313 signal. Green indicates Nup138 signal and magenta indicates Nup313-smHA signal.

Figure 3: P. berghei NPCs organize around the centriolar plaque (CP) during and after mitosis.
a. Representative images of mitotic nuclei containing CPs (indicated by black staining of NHS ester) and the organization of three respective FG Nups around the CP. Nups are shown in magenta, DNA is shown in blue, protein density is shown in grayscale. **b.** Quantitation of Nups around the CPs in Nup138-smHA, Nup221-3xHA/GFP, and Nup313-smHA stained NPCs. * indicates P-value > 0.05, ** indicates P-value >0.005. **c.** Representative images of post-mitotic CPs and the organization of three respective FG Nups around where the inner CP disassembles. Nups are shown in magenta, DNA is shown in blue, protein density is shown in grayscale Scale bars 2 μm, image depth in μm indicated on the image.

Figure 4: *P. berghei* replaces Nups in post-first mitotic NPCs.

a. Schematic showing endogenous tagging strategy to introduce the RITE system at the 3' end of the *Nup221* gene. Translocation of Cre-EBD to the nucleus following treatment with β-estradiol results in excision between loxP sites, removing the original 3xHA/GFP tag while bringing the 3xMyc/RFP tag into frame. **b.** Representative images of *P. berghei* parasites at 0 hours post, 10 hours post, and 20 hours post induction with β-estradiol. HA signal (Scale bars 2 μm, Image depth in μm indicated. White arrows indicate areas of colocalization between HA and Myc. **c.** Ratio Nup221-HA/Nup221-Myc foci in 0 hours-, 10 hours- and 20 hours post induction with β-estradiol. Green bars indicate number of foci that had both HA and Myc signal. **d.** Amount of HA or Myc signal found around the CP in 10 hour- and 20 hour post induction parasites determined by counting foci around the CP. Green bars indicate foci containing both HA and Myc signal.

Methods:

Parasite maintenance

P. berghei ANKA clone 2.34 and derivatives were maintained in Swiss Webster mice (Charles River). All experiments involving rodents were reviewed and approved by the Iowa State University Institutional Animal Care and Use Committee.

Genetic modification of *Plasmodium berghei*

Plasmid cloning was carried out using NEBuilder HIFI DNA assembly (NEB). Primer sequences are given in Table 1. To generate a smHA [54] fusion at the endogenous C terminus of *P. berghei* Nup138 the smHA coding sequence was amplified from plasmid pCAG_smFP_HA(Addgene #59759) using primers P4121/4122 and inserted between BamHI/BsiWI immediately downstream of the *Pbnup138* homology region in the plasmid pLIS0654 [17] resulting in the plasmid pSA026. To generate a smHA fusion at the endogenous C terminus of *P. berghei* Nup313, a homology region corresponding to the 3' coding sequence of *Pbnup313* up to but not including the stop codon was amplified with primers P4140/4126 from *P. berghei* genomic DNA and inserted between AvrII/BamHI in pSA026, replacing the *Pbnup138* homology region and resulting in the plasmid pSA028.

To generate parasites expressing Cre-EBD with the RITE cassette fused to the endogenous C terminus of *P. berghei* Nup221, the Cre-EBD coding sequence was amplified from the plasmid pTWO40 [55] (Addgene #64769) and the RITE cassette was amplified from the plasmid pKV016, which includes a 3xHA-GFP fusion in the pre-excised state and a 3xMYC-mRFP fusion in the post-excised state [56] (Addgene #64767). These sequences were assembled together with a *Pbnup221* homology region [17], placing Cre-EBD under the control of *P. berghei* EF1 alpha promoter and the hDHFR selectable marker under the control of the *P. berghei* EF1 delta promoter and positioning the hDHFR cassette between the two loxP-containing RITE tags to facilitate marker removed upon excision. The mRFP sequence in the post-excision RITE fusion was subsequently replaced with mRuby3 to yield the final plasmid pL857.

Plasmids pSA026, pSA028 and pL857 were linearized within the *Pbnup138, Pbnup313 or*
Pbnup221 homology region using HindIII, PacI or ScaI, respectively, and transfected into *P.*
berghei as described [57]. Transfected parasites were injected into naïve mice and selected with
0.07 mg/ml pyrimethamine provided *ad libitum* in drinking water beginning 24 hours post-
transfection. Parasites were genotyped to confirm integration at *Pbnup138, Pbnup313* or
Pbnup221 using PCR strategies described in Supplementary Figures 2 and 4.

Live imaging and immunofluorescence assays

For live imaging of parasites at regular intervals throughout the experiment, 100µl
parasite culture was spun down and resuspended in PBS containing 1µg/ml Hoechst 33342 to
stain nuclei. Following a brief incubation at 37℃, parasites were imaged with a 63x objective on
an Axio Observer 7 equipped with an Axiocam 702 mono camera and Zen 2.6 Pro software (Zeiss)
using the same exposure times for all images across sample groups and experimental
replicates.For immunofluorescence assays on unexpanded cells, thin blood smears were air dried
and fixed in ice-cold methanol/acetone (50:50) (for detection of smHA tagged Nup138, Nup313
or for detection of Cre). Fixed slides were blocked with Rockland blocking buffer (Rockland) and
probed with the primary and secondary antibodies indicated below. ,Slides were mounted in
Fluoromount-G with DAPI (Invitrogen).

Ultrastructure expansion microscopy

Ultrastructure expansion microscopy was performed as previously described on fixed
samples. Briefly, parasites were fixed in 4% PFA/PBS supplemented with 0.0075% glutaraldehyde
by the lab of Dr. Josh Beck at Iowa State University and were sent to the lab of Dr. Sabrina Absalon
at Indiana University School of Medicine. 12 mm round coverslips (Fisher Cat. No. NC1129240)
were treated with poly-D-lysine for 1 hour at 37 °C, washed three times with MilliQ water, and
placed in the wells of a 24-well dish. Fixed samples were resuspended at roughly 1.0 % hematocrit
in 1xPBS and 500 µL of parasite sample was added to the wells containing the coated coverslips
for 30 minutes at 37 °C. Following parasites settling, coverslips were washed 3 times in 1xPBS
before treating with 500 µL 1.4% v/v formaldehyde/2% v/v acylamide (FA/AA) in PBS. Samples

were incubated overnight at 37 °C. Gelation, denaturation, staining, and expansion of the gels

were performed as previously described [24]. Stained gels were imaged using a Zeiss LSM900

AxioObserver with an Airyscan 2 detector. Images were taken using a 63x Plan-Apochromat

objective lens with a numerical aperature of 1.4.

RITE experiment setup

Asynchronous Nup221::RITE *P. berghei* parasites were collected and allowed to develop

overnight in vitro into terminal schizonts. Schizonts were magnetically purified on a MACS LD

column (Miltenyi Biotec) and IV injected into naïve mice to establish a synchronous infection.

Infections were allowed to develop for three intraerythrocytic developmental cycles (IDCs) to

increase parasitemia, then collected at 72 hpi when parasites were predominantly ring stage and

passed through a MACS LD column to remove any trophozoites or schizonts by collecting the flow

through.

These synchronized ring-stage parasites were used to establish in vitro cultures in

complete RPMI containing 200nM β-estradiol or DMSO (vehicle control). Parasites were collected

every two hours for live imaging and genomic DNA isolation. After 24 hours, parasites were

washed to remove β-estradiol or DMSO and IV injected into naïve mice to monitor Nup turnover

in subsequent IDCs. Mice were not provided with pyrimethamine to avoid killing parasites that

had undergone excision, which also removes the hDHFR selection marker.

Antibodies and stains

The following antibodies were used for widefield IFA analysis: rabbit polyclonal anti-HA

(1:500, Thermofisher SG77), rabbit anti-Cre polyclonal (1:500, Abcam ab190177). The following

antibodies were used for the preparation of stained gels for expansion microscopy: Rat anti-HA

(3F10, 1:25, Roche 12158167001), Mouse IgG1 anti-alpha Tubulin Clone B-5-1-2 (1:500,

ThermoFisher 32-2500), Rabbit anti-Myc polyclonal (1:500, ThermoFisher 710007). Goat anti-

rabbit Alexa Fluor 594 (1:500, ThermoFisher A11012), Goat anti-rat Alexa Fluor 488 (1:500,

ThermoFisher A1106), Goat anti-mouse IgG(H+L) Alexa Fluor 555 (1:500, ThermoFisher A21422),

and Goat anti-rabbit Alexa Fluor 555 (1:500, ThermoFisher A32732) were used as secondary

antibodies. NHS Ester Alexa Fluor 405 in DMSO (8 μg/mL, ThermoFisher A30000) was used to

stain proteins and SYTOX™ Deep Red (1 μM, ThermoFisher S11380) was used as a DNA stain.

Image analysis

All image analysis was done using 3D Airyscan processing at moderate filter strength on

Zen Blue Version 3.1 (Zeiss, Oberkochen, Germany). Images shown in this manuscript are

maximum intensity projections of between 5 and 60 z-slices of the entire image, with the number

indicated as the total Z-depth of the image (Z depth = 0.13 μm * number of slices taken for

projection)

Statistical analysis

Graphical data and statistical analysis was performed on Graphpad Prism version 9.0.

Acknowledgements

We thank The Company of Biologists Travelling Fellowship grant JCSTF2202676 (S.V.A)

for initial travel funding to begin this project. This work was supported by NIH grant AI139579

(G.R.M. and J.R.B.) and was made possible by an award from the Indiana University School of

Medicine (S.A., BRG award #2286272). The content is solely the responsibility of the authors

and does not necessarily represent the official views of the Indiana University School of

Medicine.

Table S1. Sequences of primers used in this study.

Name	Sequence
P4121	GTTATTCTCACGAGCTTAGAAATATGAAGGGATCCATGTACCCTTATGATGTGCCCGATTATGC
P4122	GATAGCCAGCGTAGTCCGGGACGTCGTACGGGTAGCCACCCTTGTAGAGTTCGTC
P4140	GACGATAAAAATGAAAATACAAATCCTAGGGAAATTGAAATCGCTTGCACATATTTGAATC
P4126	CATCATAAGGGTACATGGATCCGTTTATGACATTTTGGCTAGTG
P5	GGAAAAAGCTTAACATTTAACTCTTCATTTTTTAAAACGAGCAC
P6	CCGGGACGTCGTACGGG

P7	CTTAGCCCATGCGAATGCATACTATTAGAAC
P8	GTCCATTAACGTCGCCATCCAACTCC
P3922	TTTAAATGATACAAAACACAGACG
P3915	ATACTAGTAGCGTAATCTGGAACG
P3888	TCAATGATTCATAAATAGTTGGACTTG
P3659	ATCAACAAATTATAAATGACAGAAC
P3004	AAAAGATCTATGGTTGGTTCGCTAAACTG
P3005	AAACAATTGTTAATCATTCTTCTCATATAC
P3491	GATAAATTCTCGAGCCCGGGTATCGATTATTTAGAGAATC
P3492	TCTTTACCTCCGCCGGATCCAATATTTTTAATAACATTTC
P4046	TTCCATGACCACCTTCATACGC
P3921	CGCATGAACTCCTTGATGACG

470 **Table S2: Plasmids used in this study.**

Genotype	Plasmid Name
nup138::smHA	pSA026
nup313::smHA	pSA028
nup221::RITE	pL857

Author Contributions:

Conceptualization: J.B., S.V.A., G.R.M., J.R.B., and S.A.; **Methodology:** J.B., S.V.A., G.R.M., J.R.B.,

and S.A.; **Investigation:** J.B., S.V.A., and T.H.; **Data Analysis:** J.B. **Writing – Original Draft:** J.B.;

Writing – Review and Editing: J.B., S.V.A., T.H., G.R.M., J.R.B., and S.A; **Funding Acquisition:**

S.V.A., G.R.M., J.R.B., and S.A.; **Supervision:** G.R.M., J.R.B. and S.A.

References:

1.   Lin, D.H. and A. Hoelz, *The Structure of the Nuclear Pore Complex (An Update).* Annu Rev
Biochem, 2019. **88**: p. 725-783.

2.   Hoelz, A., E.W. Debler, and G. Blobel, *The Structure of the Nuclear Pore Complex.* Annual Review
of Biochemistry, 2011. **80**(1): p. 613-643.

3.   Grossman, E., O. Medalia, and M. Zwerger, *Functional Architecture of the Nuclear Pore Complex.*
Annual Review of Biophysics, 2012. **41**(1): p. 557-584.

4.   Ori, A., et al., *Cell type-specific nuclear pores: a case in point for context-dependent stoichiometry*
of molecular machines. Mol Syst Biol, 2013. **9**: p. 648.

5.   Otsuka, S. and J. Ellenberg, *Mechanisms of nuclear pore complex assembly – two different ways*
of building one molecular machine. FEBS Letters, 2018. **592**(4): p. 475-488.

6.   Courjol, F., et al., *Characterization of a nuclear pore protein sheds light on the roles and*
composition of the Toxoplasma gondii nuclear pore complex. Cell Mol Life Sci, 2017. **74**(11): p.
2107-2125.

7.   Degrasse, J.A. and D. Devos, *A functional proteomic study of the Trypanosoma brucei nuclear*
pore complex: an informatic strategy. Methods Mol Biol, 2010. **673**: p. 231-8.

8.   Beck, M. and E. Hurt, *The nuclear pore complex: understanding its function through structural*
insight. Nature Reviews Molecular Cell Biology, 2017. **18**(2): p. 73-89.

9.   Devos, D., et al., *Simple fold composition and modular architecture of the nuclear pore complex.*
Proc Natl Acad Sci U S A, 2006. **103**(7): p. 2172-7.

10.   Strawn, L.A., et al., *Minimal nuclear pore complexes define FG repeat domains essential for*
transport. Nature Cell Biology, 2004. **6**(3): p. 197-206.

11.   Dey, G., et al., *Closed mitosis requires local disassembly of the nuclear envelope.* Nature, 2020.
585(7823): p. 119-123.

12.   Lénárt, P., et al., *Nuclear envelope breakdown in starfish oocytes proceeds by partial NPC*
disassembly followed by a rapidly spreading fenestration of nuclear membranes. J Cell Biol, 2003.
160(7): p. 1055-68.

13.   Gerald, N., B. Mahajan, and S. Kumar, *Mitosis in the human malaria parasite Plasmodium*
falciparum. Eukaryotic cell, 2011. **10**(4): p. 474-482.

14.   Neumann, N., D. Lundin, and A.M. Poole, *Comparative genomic evidence for a complete nuclear*
pore complex in the last eukaryotic common ancestor. PLoS One, 2010. **5**(10): p. e13241.

15.   Kehrer, J., et al., *Nuclear Pore Complex Components in the Malaria Parasite Plasmodium berghei.*
Scientific Reports, 2018. **8**(1): p. 11249.

16.   Dahan-Pasternak, N., et al., *PfSec13 is an unusual chromatin-associated nucleoporin of*
Plasmodium falciparum that is essential for parasite proliferation in human erythrocytes. Journal
of Cell Science, 2013. **126**(14): p. 3055-3069.

17.   Ambekar Sushma, V., R. Beck Josh, and R. Mair Gunnar, *TurboID Identification of Evolutionarily*
Divergent Components of the Nuclear Pore Complex in the Malaria Model Plasmodium berghei.
mBio, 2022. **13**(5): p. e01815-22.

18.   Bertiaux, E., et al., *Expansion microscopy provides new insights into the cytoskeleton of malaria*
parasites including the conservation of a conoid. PLoS Biol, 2021. **19**(3): p. e3001020.

19.   Colombi, P., et al., *The transmission of nuclear pore complexes to daughter cells requires a*
cytoplasmic pool of Nsp1. Journal of Cell Biology, 2013. **203**(2): p. 215-232.

20.   Doye, V., R. Wepf, and E.C. Hurt, *A novel nuclear pore protein Nup133p with distinct roles in*
poly(A)+ RNA transport and nuclear pore distribution. The EMBO Journal, 1994. **13**(24): p. 6062-
6075.

21.   Weiner, A., et al., *3D nuclear architecture reveals coupled cell cycle dynamics of chromatin and*
nuclear pores in the malaria parasite Plasmodium falciparum. Cellular Microbiology, 2011. **13**(7):
p. 967-977.

22.   Winey, M., et al., *Nuclear Pore Complex Number and Distribution throughout theSaccharomyces*
528   *cerevisiae Cell Cycle by Three-Dimensional Reconstruction from Electron Micrographs of Nuclear*
529   *Envelopes.* Molecular Biology of the Cell, 1997. **8**(11): p. 2119-2132.

530   23.   Caroline, S.S., et al., *An extended DNA-free intranuclear compartment organizes centrosome*
531   *microtubules in malaria parasites.* Life Science Alliance, 2021. **4**(11): p. e202101199.

532   24.   Liffner, B. and S. Absalon *Expansion Microscopy Reveals Plasmodium falciparum Blood-Stage*
533   *Parasites Undergo Anaphase with A Chromatin Bridge in the Absence of Mini-Chromosome*
534   *Maintenance Complex Binding Protein.* Microorganisms, 2021. **9**, DOI:
535   10.3390/microorganisms9112306.

536   25.   Liffner, B., et al., *Atlas of Plasmodium falciparum intraerythrocytic development using expansion*
537   *microscopy*. 2023, eLife Sciences Publications, Ltd.

538   26.   Gambarotto, D., et al., *Imaging cellular ultrastructures using expansion microscopy (U-ExM).* Nat
539   Methods, 2019. **16**(1): p. 71-74.

540   27.   Toyama, B.H., et al., *Visualization of long-lived proteins reveals age mosaicism within nuclei of*
541   *postmitotic cells.* Journal of Cell Biology, 2018. **218**(2): p. 433-444.

542   28.   Verzijlbergen, K.F., et al., *Recombination-induced tag exchange to track old and new proteins.*
543   Proceedings of the National Academy of Sciences, 2010. **107**(1): p. 64-68.

544   29.   Hinterndorfer, K., et al., *Ultrastructure expansion microscopy reveals the cellular architecture of*
545   *budding and fission yeast.* Journal of Cell Science, 2022. **135**(24): p. jcs260240.

546   30.   Arnot, D.E., E. Ronander, and D.C. Bengtsson, *The progression of the intra-erythrocytic cell cycle*
547   *of Plasmodium falciparum and the role of the centriolar plaques in asynchronous mitotic division*
548   *during schizogony.* Int J Parasitol, 2011. **41**(1): p. 71-80.

549   31.   Simon, C.S., et al., *An extended DNA-free intranuclear compartment organizes centrosome*
550   *microtubules in malaria parasites.* Life Sci Alliance, 2021. **4**(11).

551   32.   Allegretti, M., et al., *In-cell architecture of the nuclear pore and snapshots of its turnover.* Nature,
552   2020. **586**(7831): p. 796-800.

553   33.   Lee, C.-W., et al., *Selective autophagy degrades nuclear pore complexes.* Nature Cell Biology,
554   2020. **22**(2): p. 159-166.

555   34.   Tomioka, Y., et al., *TORC1 inactivation stimulates autophagy of nucleoporin and nuclear pore*
556   *complexes.* J Cell Biol, 2020. **219**(7).

557   35.   Fernandes, P., et al., *The dimerisable Cre recombinase allows conditional genome editing in the*
558   *mosquito stages of Plasmodium berghei.* PLoS One, 2020. **15**(10): p. e0236616.

559   36.   Kent, R.S., et al., *Inducible developmental reprogramming redefines commitment to sexual*
560   *development in the malaria parasite Plasmodium berghei.* Nat Microbiol, 2018. **3**(11): p. 1206-
561   1213.

562   37.   Ferreira, J.L., et al., *Variable microtubule architecture in the malaria parasite.* Nat Commun,
563   2023. **14**(1): p. 1216.

564   38.   Chial, H.J., et al., *Saccharomyces cerevisiae Ndc1p is a shared component of nuclear pore*
565   *complexes and spindle pole bodies.* J Cell Biol, 1998. **143**(7): p. 1789-800.

566   39.   Jaspersen, S.L. and S. Ghosh, *Nuclear envelope insertion of spindle pole bodies and nuclear pore*
567   *complexes.* Nucleus, 2012. **3**(3): p. 226-36.

568   40.   Gomar-Alba, M. and M. Mendoza, *Modulation of Cell Identity by Modification of Nuclear Pore*
569   *Complexes.* Frontiers in Genetics, 2020. **10**.

570   41.   Kane, M., et al., *Nuclear pore heterogeneity influences HIV-1 infection and the antiviral activity of*
571   *MX2.* eLife, 2018. **7**: p. e35738.

572   42.   Hinshaw, J.E. and R.A. Milligan, *Nuclear pore complexes exceeding eightfold rotational symmetry.*
573   Journal of Structural Biology, 2003. **141**(3): p. 259-268.

43.   Unwin, P.N. and R.A. Milligan, *A large particle associated with the perimeter of the nuclear pore*
complex. Journal of Cell Biology, 1982. **93**(1): p. 63-75.

44.   Pulianmackal, A.J., et al., *Misregulation of Nucleoporins 98 and 96 leads to defects in protein*
synthesis that promote hallmarks of tumorigenesis. Dis Model Mech, 2022. **15**(3).

45.   Ferreira, J., J.H. Stear, and H. Saumweber, *Nucleoporins NPP-10, NPP-13 and NPP-20 are required*
for HCP-4 nuclear import to establish correct centromere assembly. J Cell Sci, 2017. **130**(5): p.
963-974.

46.   Wu, X., et al., *Disruption of the FG nucleoporin NUP98 causes selective changes in nuclear pore*
complex stoichiometry and function. Proceedings of the National Academy of Sciences, 2001.
98(6): p. 3191-3196.

47.   Powers, M.A., et al., *The vertebrate GLFG nucleoporin, Nup98, is an essential component of*
multiple RNA export pathways. J Cell Biol, 1997. **136**(2): p. 241-50.

48.   Gomes, A.R., et al., *A genome-scale vector resource enables high-throughput reverse genetic*
screening in a malaria parasite. Cell Host Microbe, 2015. **17**(3): p. 404-413.

49.   Bushell, E., et al., *Functional Profiling of a Plasmodium Genome Reveals an Abundance of*
Essential Genes. Cell, 2017. **170**(2): p. 260-272.e8.

50.   Zhang, M., et al., *Uncovering the essential genes of the human malaria parasite Plasmodium*
falciparum by saturation mutagenesis. Science, 2018. **360**(6388).

51.   Dey, G. and B. Baum, *Nuclear envelope remodelling during mitosis.* Curr Opin Cell Biol, 2021. **70**:
p. 67-74.

52.   Linder, M.I., et al., *Mitotic Disassembly of Nuclear Pore Complexes Involves CDK1- and PLK1-*
Mediated Phosphorylation of Key Interconnecting Nucleoporins. Dev Cell, 2017. **43**(2): p. 141-
156.e7.

53.   Kutay, U., R. Jühlen, and W. Antonin, *Mitotic disassembly and reassembly of nuclear pore*
complexes. Trends in Cell Biology, 2021. **31**(12): p. 1019-1033.

54.   Viswanathan, S., et al., *High-performance probes for light and electron microscopy.* Nat Methods,
2015. **12**(6): p. 568-76.

55.   Terweij, M., et al., *Recombination-induced tag exchange (RITE) cassette series to monitor protein*
dynamics in Saccharomyces cerevisiae. G3 (Bethesda), 2013. **3**(8): p. 1261-72.

56.   Menendez-Benito, V., et al., *Spatiotemporal analysis of organelle and macromolecular complex*
inheritance. Proc Natl Acad Sci U S A, 2013. **110**(1): p. 175-80.

57.   Hussain, T., et al., *The PTEX Pore Component EXP2 Is Important for Intrahepatic Development*
during the Plasmodium Liver Stage. mBio, 2022. **13**(6): p. e0309622.

Supplemental Figures:

Supplemental figure 1: Nuclear envelope visualization in *P. berghei*. Staining with the

endoplasmic reticulum protein PfBip (top image, yellow), Bodipy TR ceramide and Bodipy FL

ceramide to stain for membranes (middle images, white), and differences in NHS ester staining

(grayscale) to visualize the nuclear envelope. Magenta indicates Nup138 or Nup313, scale bars 2

µm, image depth in µm indicated.

Supplemental figure 2: Genotyping of Nup138, Nup313 and Nup221 endogenously tagged
parasite lines. a. Schematic of general strategy for integration into the endogenous Nup138,
Nup313 and Nup221 loci to generate C-terminal tag fusions. The plasmid is linearized with the
homology flank targeting each gene (yellow/red box). Following transfection, homologous
recombination into the target locus (blue box) results in fusion of the tag to the 3' end of the
gene. Green and red lines show integration PCR products diagnostic for the successful
integration. sm, selectable marker. b,c. 5' integration PCRs for Nup138-smHA and Nup313-smHA
parasites. d. Schematic of the plasmid used to generate the Nup221-RITE parasites. Cre-mediated
recombination between loxP sequences removes the 3xHA-GFP tag and hDHFR selection marker
and brings the mRuby3-3xMyc tag into frame. e. Integration PCRs for the Nup221-RITE parasites.

630   Reactions 1 and 2 are diagnostic for successful 5' and 3' integration, respectively. Reactions 3 and

631   4 are controls amplifying an internal portion of the plasmid and an unrelated genomic locus,

632   respectively.

Supplemental figure 3: Nup313 through the P. berghei blood stage life cycle. U-ExM images showing Nup313 at early stages and Nup138 through the life cycle of *P. berghei*. DNA shown in blue, microtubules shown in yellow, Nup313 shown in magenta, protein density shown in grayscale. Scale bars 2 μm, image depth in μm indicated.

Supplemental figure 4: Cre-EBD enables inducible genomic excision in *P. berghei*. a.

Immunofluorescence of Cre-EBD expressing *P. berghei* parasites. The addition of β-estradiol to

the parasites induces the Cre-EBD to translocate to the nucleus. Scale bar 2 μm. **b.** PCR showing

genome shift upon addition of β-estradiol, inducing a tag switch. Primers used are indicated in

the diagrams and expected length of PCR products are listed. **c.** PCR showing concentration

dependence of the RITE system tested from 0.0002 μM to 200 μM β-estradiol. 2 nM β-estradiol

is sufficient to induce a partial tag switch, while 200 nM β-estradiol was sufficient to induce a

total tag switch after 2 hours of treatment. PCR primers used are listed in the diagram.

Supplemental figure 5: Time course of the RITE system in *P. berghei* of Nup221::RITE. a. Live

imaging of Nup221::RITE parasites showing control or β-estradiol treated parasites during 16

hours of treatment, samples taken every 2 hours. In addition, parasites were cultured an

additional 80 hours, where β-estradiol treated parasites only showed RFP signal and control

parasites still only showed GFP signal. Scale bars 2 μm. **b.** PCR gel showing the efficiency of Cre-

EBD system over 16 hours. PCR primers and expected band sizes are indicated in the diagram.

Supplemental figure 6: Uncropped images from Figure 4 showing Nup221::HA/GFP::Myc/RFP.

U-ExM images showing the uncropped parasites in Figure 4. Age of the parasites is listed, HA

signal represented in magenta, Myc signal represented in yellow, DNA represented in blue,

protein density represented in grayscale. Scale bar 2 µm, image depth indicated in µm.