1 **An explanation for the neighbour repulsion phenomenon in Patterson's f-**
2 **statistics**
3

4 Gözde Atağ and Mehmet Somel
5
6 Department of Biological Sciences, Middle East Technical University, 06800 Ankara,
7 Turkey

8
9
10 **Abstract**
11

Patterson's f-statistics are among the most heavily utilized tools for analysing genome-
wide allele frequency data for demographic inference. Beyond studying admixture, f_3 and
f_4 statistics are also used for clustering populations to identify groups with similar
histories. However, previous studies have noted an unexpected behaviour of f-statistics:
multiple populations from a certain region systematically show higher genetic affinity to
a more distant population than to their neighbours, a pattern that is mismatched with
alternative measures of genetic similarity as well as with spatial relationships. We call
this counter-intuitive pattern "neighbour repulsion". Here, we first present a novel
instance of neighbour repulsion, where genomes from Bronze Age East Anatolian sites
show higher affinity towards Bronze Age Greece rather than their neighbours. This is
observed both using f_3- and f_4-statistics, contrasts with archaeological/historical
expectation, and also contradicts genetic affinity patterns captured using PCA or MDS
on genetic distances. We then propose a simple demographic model to explain this
pattern, where the neighbouring populations (such as East Anatolia) receive independent
gene flow from an external source after their split from each other. Calculating f_3- and f_4-
statistics using simulated genetic data with varying population genetic parameters, we
confirm that low-level gene flow from an external source into populations from one
region, as well as unidirectional gene flow between the studied regions, can create
neighbour repulsion in f-statistics. Meanwhile, similar to our empirical observations, an
MDS analysis of genetic distances can still capture common demographic history
between neighbours. Overall, our results highlight the intricate effects of unaccounted
population dynamics when inferring shared demographic history using f-statistics.

Introduction

Over the last decade, Patterson's f-statistics (Patterson *et al.* 2012) have become one of the most commonly applied tools in population genetics studies involving the analysis of genome-wide data from recombining chromosomes. Various f-statistics (f_2, outgroup-f_3, admixture-f_3, and f_4) quantify divergence, shared genetic drift and treeness among populations by comparing allele frequencies. These measurements enable identification of population structure, differentiation and admixture events and help decipher the demographic history of the studied populations (Peter 2016; Harris and DeGiorgio 2017). The apparent robustness of f-statistics to heterogeneity of sample size/coverage and ascertainment has gained them high popularity.

The outgroup-f_3 statistic in the form of $f_3(Outgroup; Pop2, Pop3)$ measures shared drift between *Pop2* and *Pop3* relative to an outgroup, and its magnitude has been used as a measure of pairwise population affinity. The f_4-statistic measures treeness, such that $f_4(Outgroup, Pop1; Pop2, Pop3)$ is expected to be 0 if *Pop1* shares the same drift with *Pop2* and *Pop3*, i.e. *Pop1* is a real outgroup to the latter two populations. Significant deviation from 0 can be interpreted as evidence for admixture (Patterson *et al.* 2012), but the f_4-statistic can also be used for studying population structure and shared demographic histories. If *Pop2* and *Pop3* share more of their demographic history, i.e. shared drift and/or similar admixture events that *Pop1* did not experience, and if there is no post-split admixture, we expect $f_4(Outgroup, Pop1; Pop2, Pop3)$ to be non-significant, while $f_4(Outgroup, Pop2; Pop1, Pop3)$ and $f_4(Outgroup, Pop3; Pop1, Pop2)$ to both be positive. Gene flow from *Pop1* into *Pop3* in the same model will also cause $f_4(Outgroup, Pop1; Pop2, Pop3)$ to be positive. Thus, in addition to testing admixture, reciprocal affinity in f_4-tests has frequently been used as a tool for clustering populations based on allele frequency similarities, where clustering implies similar histories. The outgroup-f_3 statistic has similarly been used to evaluate shared history between populations.

Nevertheless, f-statistics are not always straightforward to interpret. Complications of diverse type and origin have been noted, such as possible misinterpretation of an outgroup as an admixture source when using admixture-f_3 (Patterson *et al.* 2012), or the sensitivity of f_4-statistics to SNP ascertainment schemes as showcased for demographic modelling of African human populations using SNPs identified in non-African groups (Bergström *et al.* 2020; Flegontov *et al.* 2023). The most widespread type of issue arguably involves the interpretation of f_4 -statistics under the assumption of a simple demographic model where the real history is substantially more complicated. For instance, unrecognized spatial structure can be misinterpreted as admixture events (Durand *et al.* 2011). "Ghost admixture", i.e. gene flow from an unknown source, can

2

79 lead to erroneous identification of donor/recipient populations of introgression events
80 (Rogers and Bohlender 2015; Tricou *et al.* 2022). A similar confounding effect can be
81 created by backward gene flow, as shown for Neanderthal admixture estimates varying
82 spuriously due to admixture between Eurasia and Africa within the last 45,000 years (Petr
83 *et al.* 2019a).
84
Previous work using f-statistics has noted another unexpected pattern that has yet
remained largely unexplored, to our knowledge. Using f_4-statistics, Kılınç et al. observed
higher affinity of all Anatolian Neolithic groups to a Neolithic genome from Greece than
to each other, a pattern that appeared difficult to interpret (Kılınç *et al.* 2017). Even more
surprising was the observation that modern-day North African populations showed
higher affinity to modern-day Sardinians than to each other using f_3- and f_4-statistics
(Rodríguez-Varela *et al.* 2017). This pattern was hypothesized to stem from different
degrees of gene flow from divergent sources, here sub-Saharan Africa, but the case was
not studied any further.
94
Here, we provide another case of this pattern, which we call "neighbour repulsion", by
comparing genomes from the Bronze Age (BA) East Mediterranean. We then propose
scenarios involving external and regional gene flow to explain neighbour repulsion and
recapitulate the observed f-statistics using simulations under these models, with various
conditions. Our results showcase how complex demographic histories can lead to
counter-intuitive patterns of f_3- and f_4-statistics when used in the context of studying
shared population history.
102
103
104 **Results and Discussion**
105
Our initial motivation was to investigate population structure and admixture patterns in
the East Mediterranean during the Bronze Ages, a period which saw population growth
and extensive interregional trade networks being established, as inferred from material
culture data (Şahoğlu 2005; Kouka 2016; Massa 2016). We studied genetic affinities
between published genomic data from n=6 populations from BA Greece (2400-1070
BCE) and n=5 populations from BA East Anatolia (3700-1303 BCE) (Lazaridis *et al.* 2017;
Skourtanioti *et al.* 2020; Lazaridis *et al.* 2022b) (Table S1). We restricted the analysis to
1240K capture data to avoid confounding by technical factors. We note that the results
described henceforth are also observed using Central and West Anatolian BA
populations, which we do not show here; we instead focus on East Anatolia as analyses
involving populations from this region are the most counter-intuitive.
117

First, we explored genetic clustering among populations with Principal Components
Analysis (PCA) and Multidimensional Scaling (MDS) analysis. In the PCA, where the
ancient genomes are projected on a PC space calculated from modern-day West
Eurasians, genomes from Greece group separately from those from East Anatolia (Figure
S1). Then, we calculated genetic distance between all pairs of populations using 1 -
f_3(Yoruba; Pop1, Pop2) to represent distance between populations Pop1 and Pop2.
Applying MDS analysis on the resulting pairwise distance matrix, we again observed
clear clustering in MDS space that mirrored geography (Figure 1A,B).
126

127
Figure 1: (A) Geographical distribution of the studied Bronze Age genomes from Greece
and East Anatolia. **(B)** Genetic clustering of the populations in line with geography
revealed by multidimensional scaling analysis (MDS), using 1 - outgroup-f_3 values as
genetic distances. **(C)** Intra-regional diversity of the populations measured as pairwise 1
- outgroup-f_3 values within regions. **(D)** f_3 statistics of the form f_3(Yoruba; East Anatolia,
X) where X corresponds to populations from Greece and East Anatolia. **(E)** f_4 statistics of
the form f_4(Yoruba, East Anatolia; East Anatolia, Greece), where positive values depict
higher affinity of East Anatolian populations towards Greek populations.
136

Intriguingly, however, and in contrast to the patterns observed in PCA and MDS, we
noticed that the values $f_3(Yoruba; EAnatoliaY, GreeceX)$ were consistently higher than the
values $f_3(Yoruba; EAnatoliaY, EAnatoliaZ)$, where *GreeceX* is one of the six populations
from Greece, while *EAnatoliaY* and *EAnatoliaZ* are any of the five populations from East
Anatolia (Figure 1C, Table S2). On face value, this would imply more similar histories
between any East Anatolian population with those from Greece than other neighbouring
East Anatolian populations.
144
We next calculated f_4-statistics of the form $f_4(Yoruba, AnatoliaY; AnatoliaZ, GreeceX)$. In
99% of 120 comparisons we observed positive values (42% significant at Z>3, without
correcting for multiple testing) (Figure 1D, Table S3). No f_4-statistic was significantly
negative, i.e. no East Anatolian group significantly shared more drift with one of its
neighbours over a population from Greece. This starkly contrasts with the expected
geographical clustering, while it does mirror the outgroup-f_3 results.
151
We hypothesized two explanations for this neighbour repulsion pattern. One may be
gene flow from Greece into each East Anatolian population, independently. Still, given
the thousands of kilometers of distance between Greece and East Anatolia, and that this
pattern was observed for all populations in the regions studied, this may not be a
parsimonious explanation. An alternative explanation and possibly more likely scenario
could involve admixture from an external and diverse source impacting East Anatolia,
similar to that hypothesized by Rodríguez-Varela and colleagues (2017).
159
The ancestral components of both Anatolia and Greek Early/Middle Bronze Age
genomes comprise Neolithic Anatolia and eastern-related ancestry from
Caucasus/Mesopotamia (Clemente *et al.* 2021; Lazaridis *et al.* 2022b; Skourtanioti *et al.*
2023). We hypothesized that additional external gene flow into East Anatolia (e.g. from
the Levant, Zagros, or Mesopotamia) that BA Greece did not receive could lead to
differentiation within the former region and might explain the observed neighbour
repulsion pattern. As a first test of this idea, we calculated intra-regional diversity within
each site, using pairwise 1- outgroup-f_3 among genomes within a site as the diversity
measure. This revealed significantly higher median diversity in BA East Anatolia than in
BA Greece (Wilcoxon rank sum test p=3e-09).
170
We then sought to replicate neighbour repulsion using population genetic simulations
under a simple demographic model with external admixture (Figure 2). We define A and
B as two sister populations, X as a more distant lineage, Z as the source of external gene
flow into A and B, and O as outgroup to all. Here, A, B and X as our focal populations; A
and B could represent two BA East Anatolian groups, X could be a population from BA

Greece, and Z would be an unknown source that admixes only with East Anatolia. The
ancestors of multiple populations are denoted by their merged names (i.e. the ancestor
population of A and B is called AB). Note that the demographic histories of A and B are
the same except for receiving independent gene flow from Z.
180

★ f4(O,A; B,X) < 0 : <u>neighbour attraction</u>   ★ f4(O,A; B,X) > 0 : <u>neighbour repulsion</u>

181 **Figure 2:** Demographic model used for the population genetic simulations. The ancestral
182 populations are named as combinations of the descending population names.
183 Population O is the outgroup used in the calculation of f-statistics. Dashed lines
184 represent equal rates of continuous migration from Z to A and B per generation. The blue
185 line corresponds to the parameter $\Delta T_{X-AB,A-B}$.
186

188 Using this general model we ran multiple simulations where we varied the parameters
189 $m_{Z\rightarrow A,B}$, Ne_Z, Ne_{AB}, Ne_{ABX}, T_{A-B}, and T_{X-AB} (Table 1). For simplicity, we refer to T_{A-B} and T_{X-AB}
190 as a single variable that shows T_{X-AB} - T_{A-B}, which we call $\Delta T_{X-AB,A-B}$. In each case we

191   sampled 25 genomes each from O, X, A, and B, and calculated various outgroup-f_3 and
192   f_4-statistics.
193
194   **Table 1:** Parameter values used for the population genetic simulations. Migration rates
195   are given per generation, divergence times are given in generations.
196

Parameter	Values	Definition
$m_{Z->A.B}$	0, 0.001, 0.005, 0.01, 0.1	Migration rate from Z to A and B
Ne_Z	5k, 15k, 35k	Effective population size of source Z
Ne_{AB}	5k, 15k, 35k	Effective population size of ancestral AB
Ne_{ABX}	1k, 3k, 5k	Effective population size of ancestral ABX
T_{A-B}	30, 60, 90	Divergence time of A and B
T_{X-AB}	100, 125, 150	Divergence time of X and AB
$\Delta T_{X-AB,A-B}$	10, 65, 120	Difference between T_{A-B} and T_{X-AB}
Ne_X*	1k, 3k, 5k	Effective population size of X
$m_{X->A,B}$*	0, 0.001, 0.005, 0.01, 0.1	Migration rate from X to A and B

197
198   * Given parameter values for Ne_X and $m_{X->A,B}$ were tested only for scenarios where Ne_{ABX}
199   = 1k and $\Delta T_{X-AB,A-B}$ = 65. At the remaining scenarios, Ne_X was fixed at 5k and $m_{X->A,B}$ at 0.
200
201
In this model (Figure 2) one would anticipate $f_4(O, A; B, X)$ values to be negative since
the populations A and B appear to share more similar histories with each other than with
X. Indeed, in the absence of gene flow from Z into A/B (m=0), f_4 statistics are mostly
negative, reflecting higher shared drift between A and B than with X (Figure 3, Figure S2).
However, f_4 values change as we allow migration from Z into A/B (from m=0 to m=0.1),
although in non-linear fashion. We observe a monotonic increase in $f_4(O, A; B, X)$ values
between 0 and 0.01, i.e. a stronger trend towards neighbour repulsion (Figure 3). Under
specific conditions, e.g. Ne_{ABX} = 1k, 0.001≤m≤0.1 and $\Delta T_{X-AB,A-B}$ ≤ 65, all $f_4(O, A; B, X)$
values are positive (Figure 3, S2). The same applies to $f_4(O, B; A, X)$. This neighbour
repulsion pattern is analogous to that observed in the empirical data where two East
Anatolian populations showed higher affinity to populations from Greece than to each
other (Figure 1D).

Figure 3: Change in f_4 statistics of the form $f_4(O, A; B, X)$ depending on the varying parameters in population genetic simulations, where negative values represent higher

217 affinity between A and B, and positive values higher affinity between A and X. The panels
218 for each parameter contain all values of the remaining studied parameters and are
219 explained in Table 1.
220
221
We note a number of conditions driving more positive $f_4(O, A; B, X)$ values under low-
level migration in our model (Figure 2; Table 1): (a) Low $\Delta T_{X\text{-}AB,A\text{-}B}$, which means A and B
share less drift relative to X (the phylogeny is more star-like), and also that A and B are
impacted by Z-sourced admixture for fewer generations. (b) Low Ne_{ABX}, which causes
stronger shared drift between A, B and X. (c) High Ne_{AB}, which leads to lower shared drift
between A and B. (d) High Ne_Z, whereby Z is more diverse and thus Z-sourced gene flow
differentiates A and B more effectively.
229
230 As opposed to results involving low-level migration (m=0.001 to m=0.01), at m=0.1, $f_4(O,$
231 $A; B, X)$ values shift back to negative values, i.e. neighbour attraction (Figure 3, S2). This
232 happens because high migration rates from Z leads to homogenization of A and B. The
233 drop in f_4 values is particularly strong if Ne_Z is low: e.g. at m=0.1 and $Ne_Z = 5k$ all results
234 are negative (Figure 3, S2).
235
We then asked whether in simulations where we detect neighbour repulsion between A
and B in f_4-statistics, we would still observe A and B clustering separately from X in the
MDS analysis, a pattern we had observed in the empirical data (Figure 1). For this, we
chose one random case from each scenario and calculated outgroup-f_3 statistics. We
then compared the mean distance between X to A and B [$\bar{d}_{mds}(X,A/B)$] versus the distance
between A and B [$d_{mds}(A,B)$] in MDS space. In 99% of all cases, populations A and B
clustered separately from X, i.e. $d_{mds}(A,B)$ was lower. This included scenarios where
100% of the f_4 values signified neighbour repulsion (Figure 4A,B). Meanwhile, under low-
level migration we also observed $f_3(O; A, B) < f_3(O; A/B, X)$, reflecting neighbour repulsion
(Figure 4C). Hence, similar to our observation with the East Mediterranean Bronze Age
genomes (Figure 1B,D), MDS-based clusterings in our simulations contradict pairwise f_3
statistics.

Figure 4: **(A)** MDS of simulated populations A, B, X and Z, using 1 - outgroup-f_3 values as genetic distances, for one sample scenario where $m_{Z\text{->}A.B}$: 0.01 ,Ne_Z : 15000, Ne_{AB} : 15000 ,Ne_{ABX} : 1000, and $\Delta T_{X\text{-}AB,A\text{-}B}$: 65. **(B)** MDS clusterings between A and B versus X and B, calculated by the Euclidean distances between the mean MDS coordinates of the corresponding populations, for all scenarios. $\bar{d}_{mds}(X,AB)$ is the mean distance of X to A and B. $d_{mds}(A,B)$ is the distance between A and B. Positive values for $\bar{d}_{mds}(X,AB)$ - $d_{mds}(A,B)$ represent higher distance of X to A and B, thus distinct clustering of X from A and B. **(C)** Change in f3 values with migration rate from Z for scenarios where Ne_{ABX} : 1000 and $\Delta T_{X\text{-}AB,A\text{-}B}$: 65. $\bar{d}_{f3}(X,A/B)$ is the mean 1-f3 distance of X to A or to B. $\bar{d}_{f3}(A,B)$ is the mean 1-f3 distance between A and B. Negative values for $\bar{d}_{f3}(X,A/B)$ - $\bar{d}_{f3}(A,B)$ show higher affinity between A or B and X, than between A and B. **(D)** Intra-population diversity differences between populations A,B and X in relation to migration rate from Z, for scenarios where Ne_{ABX} : 1000 and $\Delta T_{X\text{-}AB,A\text{-}B}$: 65. The diversities are measured by pairwise 1- outgroup f3 values per population. At migration rates 0 and 0.001, the values for X have higher deviation and overlap with A and B, thus have lower visibility.

10

In these models, we might expect regional differences in within-population diversity,
which could be a source of information with respect to admixture dynamics. Indeed, in
the Bronze Age dataset we had observed significantly higher diversity in genomes from
Anatolia versus Greece (Figure 1C). In our simulations that involved gene flow from Z,
we likewise observed elevated intra-population diversity in A and B versus X (Figure 4D).

We finally studied neighbour repulsion under two additional models of gene flow. In one
case, we asked how asymmetric Z-sourced gene flow, where only population B receives
migrants from Z, would impact f_4 statistics. In this case, as might be expected, $f_4(O, A;$
$B, X)$ tends to be positive, i.e. A shows neighbour repulsion, while $f_4(O, B; A, X)$ tends to
be negative, i.e. B is still attracted to its neighbour A (Figure S3).

We then asked whether gene flow from X into A and into B independently can also lead
to neighbour repulsion, without invoking the external source Z. We confirmed this was
the case using both f_3- and f_4-statistics (Figure S4A). We further found that patterns
observed both in the above simulations with external gene flow (Figure 3,4) and in the
empirical data (Figure 1), namely lower diversity in X versus A/B and separate MDS
clustering of X and A/B, can be replicated under gene flow from X, but only when the
effective population size of X (Ne$_X$) is low relative to Ne$_A$ and Ne$_B$ (Figure S4B,C).

The results above reveal how low-level gene flow from an external source (Z) can lead to
neighbour repulsion i.e. closely related pairs of populations (A and B) showing higher
affinity to a lineage that has split earlier from them (X). We also show that independent
regional gene flow from X into A and B can likewise create neighbour repulsion patterns,
especially when Ne$_X$ is relatively small.

We believe neighbour repulsion is an interesting case because in the scenarios we
studied, A and B share more similar demographic histories with each other than with X,
although this is not directly reflected in the f_3- and f_4-statistics. Whether post-split
admixture from X or an external population Z is more likely would need to be evaluated
using other sources of evidence, such as local ancestry inference or
archaeological/historical evidence for regional gene flow. In the empirical cases we
covered, such as neighbour repulsion in modern-day North Africa or Bronze Age East
Anatolia, external gene flow may be a more parsimonious explanation than the latter.

Methods

Ancient genomes

We used 79 published Bronze Age genomes from the East Mediterranean (Supplementary Table 1), from a publicly available dataset comprising the ancient individuals analyzed in Lazaridis et al., 2022, genotyped at the 1240K array positions (Lazaridis *et al.* 2022a). We only used Bronze Age genomes from East Anatolia and Greece produced with the 1240K enrichment method in order to avoid any technical bias that can arise when co-analyzing different data types, e.g. shotgun versus capture.

Principal Components Analysis (PCA)

For the principal components analysis, we used the Human Origins Dataset (Lazaridis *et al.* 2014, 2016), containing 2,583 present-day genomes genotyped at the Affymetrix Human Origins Array positions, merged with the Bronze Age genomes. PCA was performed using *smartpca v.18140*, with option *lsqproject:YES*, implemented in *EIGENSOFT v.7.2.0* (Patterson *et al.* 2006). We projected the ancient individuals onto the first two principal components calculated from present-day Western Eurasian diversity based on populations from the Human Origins panel.

f_3, f_4 - statistics and Multidimensional scaling (MDS)

We calculated outgroup f_3- and f_4-statistics using *admixr v0.9.1,* which runs *AdmixTools v.7.0.2,* with default parameters (Patterson *et al.* 2012; Petr *et al.* 2019b). Within-group genetic diversities were estimated by computing pairwise 1 - outgroup-f_3. All statistics were computed based on the 1240K dataset, using 3 modern-day Yoruba individuals as the outgroup. We estimated pairwise genetic distances with 1- outgroup f_3, computed a distance matrix of 1-f_3 values, and performed a classical multidimensional scaling analysis using the *cmdscale* algorithm implemented in the R package *stats* (R Core Team, 2023).

Simulations

We used *msprime v1.0* (Baumdicker *et al.* 2022) to simulate diploid sequences of 100Mbs, where we sampled 25 individuals from each population. The divergence time of the population used as outgroup (ABXZO) was given as 200 generations ago, followed by the divergence of ABXZ after 20 generations. The effective population size of each population other than the ones given in Table 1 were fixed as 5k, i.e. populations ABXZO,

O, ABXZ, X (except for scenarios where Ne_{ABX} = 1k and $\Delta T_{X\text{-}AB,A\text{-}B}$ = 65), A, and B. The
mutation rate and the recombination rate were chosen as 1e-8 and 1.1e-8 per
generation, respectively (Baumdicker *et al.* 2022). Each scenario had 100 replicates.
One-way migration rates from Z were set symmetric into both A and B, except for one
case (Ne_Z, Ne_{AB}, Ne_{ABX}, $\Delta T_{X\text{-}AB,A\text{-}B}$) where we tested the effect of asymmetric migration by
allowing gene flow (m=0.01) only into B. For scenarios where Ne_{ABX} = 1k and $\Delta T_{X\text{-}AB,A\text{-}B}$
= 65, one-way migration from X into A and B was also tested (Table 1). All combinations
of values for parameters $m_{Z\text{->}A.B}$, Ne_Z, Ne_{AB}, Ne_{ABX}, and $\Delta T_{X\text{-}AB,A\text{-}B}$ given in Table 1 were
tested. The outgroup-f_3 and f_4 statistics were calculated using *tskit v0.5.6*, which follows
the calculation in Patterson et al., 2012. The MDS analysis was performed the same way
as above. In order to compare the MDS distances between populations, we used the
mean MDS x and y coordinates of each individual, and calculated the mean coordinate
point per population. We then calculated Euclidean distances between the mean
coordinates of the corresponding populations.
364
365
366 **Data Availability**
367
368 The code and parameters for generating the simulated data are deposited under
369 https://github.com/goztag/msprime_fstats/tree/main.
370
371
372

373 **Acknowledgements**
374
We thank Emilia Huerta-Sanchez, Torsten Günther, Benjamin Vernot, Pavlos Pavlidis,
Dilek Koptekin, Yetkin Alıcı and members of the METU CompEvo Team for helpful
discussions. The work was supported by the H2020 ERC Consolidator grant (no. 772390
NEOGENE to M.S.), H2020-WIDESPREAD-05-2020 TWINNING grant (no. 952317
NEOMATRIX to M.S.) and Scientific and Technological Research Council of Turkey
(TÜBİTAK) through the 2210/A National Scholarship Programme for MSc Students (to
G.A.).

References

Baumdicker F., G. Bisschop, D. Goldstein, G. Gower, A. P. Ragsdale, *et al.*, 2022 Efficient

ancestry and mutation simulation with msprime 1.0. Genetics 220: iyab229.

https://doi.org/10.1093/genetics/iyab229

Bergström A., S. A. McCarthy, R. Hui, M. A. Almarri, Q. Ayub, *et al.*, 2020 Insights into human

genetic variation and population history from 929 diverse genomes. Science 367:

eaay5012. https://doi.org/10.1126/science.aay5012

Clemente F., M. Unterländer, O. Dolgova, C. E. G. Amorim, F. Coroado-Santos, *et al.*, 2021 The

genomic history of the Aegean palatial civilizations. Cell 184: 2565-2586.e21.

https://doi.org/10.1016/j.cell.2021.03.039

Durand E. Y., N. Patterson, D. Reich, and M. Slatkin, 2011 Testing for ancient admixture

between closely related populations. Mol Biol Evol 28: 2239–2252.

https://doi.org/10.1093/molbev/msr048

Flegontov P., U. Işıldak, R. Maier, E. Yüncü, P. Changmai, *et al.*, 2023 Modeling of African

population history using f-statistics is biased when applying all previously proposed SNP

ascertainment schemes. PLOS Genetics 19: e1010931.

https://doi.org/10.1371/journal.pgen.1010931

Harris A. M., and M. DeGiorgio, 2017 Admixture and Ancestry Inference from Ancient and

Modern Samples through Measures of Population Genetic Drift. Hum Biol 89: 21–46.

https://doi.org/10.13110/humanbiology.89.1.02

Kılınç G. M., D. Koptekin, Ç. Atakuman, A. P. Sümer, H. M. Dönertaş, *et al.*, 2017

Archaeogenomic analysis of the first steps of Neolithization in Anatolia and the Aegean.

Proceedings of the Royal Society B: Biological Sciences 284: 20172064.

https://doi.org/10.1098/rspb.2017.2064

Kouka O., 2016 The Built Environment and Cultural Connectivity in the Aegean Early Bronze

Age, in Molloy, B.P.C (ed.), Of Odysseys and Oddities, Sheffield Studies in Aegean

Archaeology, Oxbow Books, 203-222.

Lazaridis I., N. Patterson, A. Mittnik, G. Renaud, S. Mallick, *et al.*, 2014 Ancient human

genomes suggest three ancestral populations for present-day Europeans. Nature 513:

409–413. https://doi.org/10.1038/nature13673

Lazaridis I., D. Nadel, G. Rollefson, D. C. Merrett, N. Rohland, *et al.*, 2016 Genomic insights into

the origin of farming in the ancient Near East. Nature 536: 419–424.

https://doi.org/10.1038/nature19310

Lazaridis I., A. Mittnik, N. Patterson, S. Mallick, N. Rohland, *et al.*, 2017 Genetic origins of the

Minoans and Mycenaeans. Nature 548: 214–218. https://doi.org/10.1038/nature23310

Lazaridis I., S. Alpaslan-Roodenberg, R. Pinhasi, and D. Reich, 2022a The genetic history of

the Southern Arc: A bridge between West Asia and Europe

Lazaridis I., S. Alpaslan-Roodenberg, A. Acar, A. Açıkkol, A. Agelarakis, *et al.*, 2022b The

genetic history of the Southern Arc: A bridge between West Asia and Europe. Science

377: eabm4247. https://doi.org/10.1126/science.abm4247

Massa M., 2016 Networks before Empires: cultural transfers in western and central Anatolia

during the Early Bronze Age

Patterson N., A. L. Price, and D. Reich, 2006 Population Structure and Eigenanalysis. PLOS

Genetics 2: e190. https://doi.org/10.1371/journal.pgen.0020190

Patterson N., P. Moorjani, Y. Luo, S. Mallick, N. Rohland, *et al.*, 2012 Ancient Admixture in

Human History. Genetics 192: 1065–1093. https://doi.org/10.1534/genetics.112.145037

Peter B. M., 2016 Admixture, Population Structure, and F-Statistics. Genetics 202: 1485–1501.

https://doi.org/10.1534/genetics.115.183913

Petr M., S. Pääbo, J. Kelso, and B. Vernot, 2019a Limits of long-term selection against

Neandertal introgression. Proc. Natl. Acad. Sci. U.S.A. 116: 1639–1644.

https://doi.org/10.1073/pnas.1814338116

Petr M., B. Vernot, and J. Kelso, 2019b admixr—R package for reproducible analyses using

ADMIXTOOLS. Bioinformatics 35: 3194–3195.

https://doi.org/10.1093/bioinformatics/btz030

R Core Team, 2023 _R: A Language and Environment for Statistical   Computing_. R

Foundation for Statistical Computing, Vienna, Austria.   <https://www.R-

project.org/>.

Rodríguez-Varela R., T. Günther, M. Krzewińska, J. Storå, T. H. Gillingwater, *et al.*, 2017

Genomic Analyses of Pre-European Conquest Human Remains from the Canary Islands

Reveal Close Affinity to Modern North Africans. Current Biology 27: 3396-3402.e5.

https://doi.org/10.1016/j.cub.2017.09.059

Rogers A. R., and R. J. Bohlender, 2015 Bias in estimators of archaic admixture. Theoretical

Population Biology 100: 63–78. https://doi.org/10.1016/j.tpb.2014.12.006

Şahoğlu V., 2005 The Anatolian Trade Network and the Izmir region during the Early Bronze

Age. Oxford Journal of Archaeology 24: 339–361. https://doi.org/10.1111/j.1468-

0092.2005.00240.x

Skourtanioti E., Y. S. Erdal, M. Frangipane, F. Balossi Restelli, K. A. Yener, *et al.*, 2020

Genomic History of Neolithic to Bronze Age Anatolia, Northern Levant, and Southern

Caucasus. Cell 181: 1158-1175.e28. https://doi.org/10.1016/j.cell.2020.04.044

Skourtanioti E., H. Ringbauer, G. A. Gnecchi Ruscone, R. A. Bianco, M. Burri, *et al.*, 2023

Ancient DNA reveals admixture history and endogamy in the prehistoric Aegean. Nat

Ecol Evol 1–14. https://doi.org/10.1038/s41559-022-01952-3

Tricou T., E. Tannier, and D. M. de Vienne, 2022 Ghost Lineages Highly Influence the

Interpretation of Introgression Tests. Systematic Biology 71: 1147–1158.

https://doi.org/10.1093/sysbio/syac011