Cingulate cortex shapes early postnatal development of social vocalizations

Short title: Cortical development of vocal social behavior

Gurueswar Nagarajan[1], Denis Matrov[1], Anna C. Pearson[1], Cecil Yen[2],

Sean P. Bradley[3], Yogita Chudasama[1,3*]

[1]Section on Behavioral Neuroscience, National Institute of Mental Health,

[2]Reproductive Biology and Genetics Core, National Institute of Neurological Disorders and Stroke

[3]Rodent Behavioral Core, National Institute of Mental Health,

National Institutes of Health, Bethesda, MD 20892, USA.

Address for correspondence:
Yogita Chudasama, Ph.D.
National Institute of Mental Health
Building 35A, Room 2D-804
Bethesda, MD 20892
Tel: 301 594-4569
Email: yogita.chudasama@nih.gov

Abstract

The social dynamics of vocal behavior has major implications for social development in humans. We asked whether early life damage to the anterior cingulate cortex (ACC), which is closely associated with socioemotional regulation more broadly, impacts the normal development of vocal expression. The common marmoset provides a unique opportunity to study the developmental trajectory of vocal behavior, and to track the consequences of early brain damage on aspects of social vocalizations. We created ACC lesions in neonatal marmosets and compared their pattern of vocalization to that of age-matched controls throughout the first 6 weeks of life. We found that while early life ACC lesions had little influence on the production of vocal calls, developmental changes to the quality of social contact calls and their associated syntactical and acoustic characteristics were compromised. These animals made fewer social contact calls, and when they did, they were short, loud and monotonic. We further determined that damage to ACC in infancy results in a permanent alteration in downstream brain areas known to be involved in social vocalizations, such as the amygdala and periaqueductal gray. Namely, in the adult, these structures exhibited diminished GABA-immunoreactivity relative to control animals, likely reflecting disruption of the normal inhibitory balance following ACC deafferentation. Together, these data indicate that the normal development of social vocal behavior depends on the ACC and its interaction with other areas in the vocal network during early life.

Introduction

Vocal behavior is a critical mediator of social communication through different life stages of many animals, and particularly in social species such as primates[1]. The common marmoset is a small, arboreal monkey with an elaborate repertoire of acoustic calls. While the meaning and usage of most marmoset vocalizations are not well understood, research has shown that different call types convey information about their social organization, environment, and the presence of food or predators[2,1]. Moreover, these calls undergo developmental progression. During the first postnatal months, the acoustic properties and usage of marmoset infant vocalizations change markedly. For example, for different call types, parameters such as duration and frequency follow typical trajectories during the first months of life, transitioning from an immature babbling phase with a mixture of proto-calls to a more discrete and contingent usage of adult-like calls[3,4]. Recent evidence also suggests that parental or social interaction plays a significant role in the proper development of normal vocal behavior, raising the prospect that important aspects of marmoset vocal behavior are learned[5,6].

In this study we focus on the anterior cingulate cortex (ACC) and its contribution to vocal behavior and its development in early life. The ACC is a limbic cortical region known to contribute to vocal behaviors[7,8], and particularly those associated with emotional states[9–11]. Electrical stimulation of the most rostral segment of the ACC elicits vocalizations[12–14], whereas ACC ablations limit spontaneous vocalizations[15] and voluntary control of vocal behavior[16,17]. Its dense anatomical connections with the amygdala[18,19] underscore its role in shaping the affective component of vocalizations[20–23]. At the same time, its descending projections to the periaqueductal gray[24,25] endow the ACC direct control over activating the brainstem vocalization pathway[7,26,27]. Vocal production leads to expression of immediate early genes in the ACC[28], with early studies

reporting that infant ACC lesions abolish the characteristic cries that infants normally issue when separated from its mother[29]. These findings implicate the ACC in volitional and emotional control over vocal output.

Longitudinal monitoring of vocal behavior provides a tractable, high dimensional readout of the development of socio-affective circuits. It also provides a means to investigate how early life disruption to brain areas such as the ACC might affect the normal progression of social interaction. If the ACC contributes to the early-life maturation of vocal behavior, then neonatal ACC lesions should hamper the normal control of emotional vocal utterances. Here, we performed excitotoxic ACC lesions in neonatal marmosets and tracked their vocal behaviors, comparing them to age-matched controls throughout the first 6 weeks of life, and examined the impact of the early life lesion on interconnected brain regions in the vocal production network. We demonstrate that animals with neonatal damage to the ACC retained their capacity to issue calls. However, these animals showed a change in their vocal repertoire and an altered acoustic structure in their communicative "social" calls, as well as permanent anatomical changes in the amygdala (AMY) and periaqueductal gray (PAG).

Results

We studied the vocal behavior in 10 infant marmosets (five males and five females) from five different sets of unrelated parents. In five of the neonatal animals, we performed surgical excitotoxic lesions bilaterally to the rostral portion of the dorsal ACC (24a and 24b) (**Fig. 1A, B**, see **Methods**). Starting seven days before the surgery and continuing until six postnatal weeks, infant vocalization behavior was recorded in an isolated, temperature-controlled incubator in 5 min sessions, 2-3 times a week (**Fig. 1C, 1D**). In four of the animals, the estimate of ACC volume

from T2-weighted MR scans performed under anesthesia approximately eight months of age revealed a 60% decrease in ACC volume compared with four control animals (**Fig. 1E;** two animals were not scanned**).** Following sexual maturity, at approximately two years of age, the animals were euthanized and their brains were histologically visualized to verify the extent of the lesion. We also examined downstream effects of the lesion, including an evaluation of its effects on mature neurons (NeuN), inhibitory neurotransmitters (GABA and GAD67), glial cells (GFAP and Iba1) and fiber tracts (myelin; **Fig. 1F**). See Materials and Methods for details.

Verification of the ACC lesion and its impact on downstream vocal structures

The intended lesions and reconstructed ACC damage based on histological evaluation are shown in one hemisphere for four animals in **Figs 2A** and **2B**, respectively. The ACC lesion covered most of the target cytoarchitectonic areas 24a and 24b of Paxinos et al.[30], just above the corpus callosum. The rostral limit of the lesions was adjacent to the genu of the corpus callosum and the caudal limit just anterior of area 23a caudally. Dorsally, the lesions extended past 24b into motor area 6M. There was little if any encroachment into subgenual area 25. Apart from one case which showed some sparing of the lesion in the left hemisphere, there was extensive overlap in the placement of the ACC lesion.

Representative photomicrographs of the ACC lesion and a control are presented in **Fig. 2B,** which shows the distribution of myelinated fibers in the ACC region stained using a high-resolution Black-Gold II myelin stain (Histo-Chem Inc., Jefferson, AR). There was clear evidence that the lesion created a major disruption to the normal radial arrangement of the fibers in the ACC region caused by extensive demyelination of the axons (**Fig 2B**). The ACC lesion also impacted the integrity of white matter tracts local to the site of the lesion (data not shown), but the transverse diameter of major fiber tracts, namely the corpus callosum and the anterior commissure did not

differ between the groups. The loss of neurons, however, and the respective increase in glial cells at the lesioned site especially at the interface between the gray and white matter was clearly observed (**Fig. 2C**).

We examined the cellular and neurotransmitter composition of the amygdala (AMY) and periaqueductal gray (PAG), as these structures are downstream from the ACC and their natural development may be affected by the infant ACC lesions. We first investigated whether neurons in these structures were degenerated using Fluro-Jade C (Histo-Chem Inc.), which is as a marker for apoptotic, necrotic, and autophagic cells. There was no sign of neurodegeneration in these downstream brain regions two years following the infant lesion. We next examined the proportion of neurons in the AMY and PAG expressing GABA, since changes in the relative number of inhibitory interneurons could serve as a marker for downstream neuroplasticity in response to the ACC lesion[31]. We found a significant reduction of GABA positive neurons in two structures, the basomedial AMY and the dorsal portion of the PAG **(Fig 2D).** This reduction suggests a disruption of the normal inhibitory balance within the vocal network following the infant ACC lesions.

Vocal behavior persists immediately following neonatal ACC lesions

In the weeks following bilateral ACC lesions, infants tested in the isolated chamber continued to vocalize readily, which is somewhat surprising given the critical role of this structure in normal vocal behavior[12–14]. From (presurgical) postnatal week 2 to (postsurgical) postnatal week 6, we annotated 23,000 calls from the five lesioned and five control marmosets. Sample spectrographs from audio recordings of a twin pair before and after surgery are shown in **Fig. 3A-B.** While there was variability among individuals, the calls were complex and diversified from postnatal week 2, consisting of cries as well as immature versions of adult vocalizations including phee, twitter, and

trills, as well as complex calls when two calls merged such as trill-twitter or cry-phee, or any other combination (**Fig 3A**). By the sixth postnatal week, the call repertoire for both the lesioned animals and the controls had both evolved, with no conspicuous difference between groups (**Fig 3B**). Most notably, the relative reduction of the total rate and diversity of calls was similar between groups (**Fig. 3C,** $\chi2(2)=2.8464$, p=0.24), with the reduction matching the known maturational changes accompanying growth of the vocal apparatus and increased respiratory powers[32]. Further analysis showed that the ACC lesion had minimal effects on the rate of most call types during this period (phee (β= -0.07, 95% CI [-0.31, 0.17], p = 0.49); twitter (β= -0.07, 95% CI [-0.16, 0.01], p = 0.09); trill (β= -0.03, 95% CI [-0.11, 0.06], p = 0.49); cry (β= 0.13, 95% CI [-0.03, 0.29], p= 0.10) (**Fig. 3D**). The exception to this rule was an elevation in the rate of 'other' calls, which comprised tsik, egg, eck, chatter and seep calls. These calls were significantly elevated in animals after the ACC lesion (β= 0.11, 95% CI [0.03, 0.20], p = 0.018). This was driven mostly by an increase in the lesion group during postnatal week 4.

Two additional variables relatively unaffected by the ACC lesion were the call durations and inter-call intervals, acoustic features that have been used to track vocal development in previous studies[4,6,33]. Consistent with the overall decrease in vocalization rate with increasing age, there was an associated increase in inter-call intervals which was noted at late postnatal weeks (β= 0.33, 95% CI [0.22, 0.43], p < .001) which held true for both controls and lesioned group ($\chi2(2)=1.88$, p=0.39). None of the specific call types exhibited developmental changes in call type duration (phee ($\chi2(2)=1.08$, p=0.58); trill ($\chi2(2)=2.87$, p=0.24); twitter ($\chi2(2)=2.79$, p=0.25); cry ($\chi2(2)=0.057$, p=0.97), but there were slight changes in the duration of phee syllables exhibited only by animals with an ACC lesion which we discuss later.

Neonatal ACC lesions prevent the maturation of social contact calls

Whereas many of the basic vocal parameters evolved normally in the animals with the ACC lesions, one major difference related to their use of social contact calls. By 6 weeks of age, marmoset vocalizations are known to approach their mature state and become dominated by social contact calls, namely phees, trills, and twitters. The phee call is studied most extensively as a long-distance contact call. It is typically evoked when the animal is socially distanced or isolated, and it promotes vocal exchanges between marmosets located out of sight in far-away locations[34,35] to facilitate reunion with family groups[36]. Trills and twitters are short distance contact calls thought to monitor the presence of group members[37,38]. Since the ACC plays a major role in socioemotional cognition (for review, see Devinsky et al.[39]), we surmised that the ACC lesion might specifically influence the socioaffective content of the vocalization that is normally expressed through contact calls.

We thus grouped phee, twitter, and trill calls as social contact calls and compared them with non-contact calls (i.e., cries and other minor occurring calls like tsik, egg, ecks, chatter, and seep). At 6 weeks of age, social contact calls predominated the control animal's vocalization. However, in ACC-lesioned animals, this aspect of social vocal behavior was substantially reduced. This difference emerged gradually after the surgery, and was only evident at 6 weeks of age (**Fig 3E**; ($\chi 2(2)=7.58$, p=0.022**).** By the sixth week, the social vocal repertoire of the lesioned animals was altogether different from the control animals, with a much smaller proportion of social contact calls.

To further understand the effect of the ACC lesion on the normal distribution of calls, we investigated the call transition probabilities between contact calls, cries, and other calls (**Fig 3F**). In contrast to the control animals, whose repertoire was dominated by social contact calls, the ACC

lesion group showed frequent transitions mostly to other non-contact call types (u-index Wilcoxon test, p = 0.055). These data suggest, therefore, that the ACC mediates developmental changes within the first 6 weeks of life that lead to the dominant production of isolation-induced contact calls and the gradual reduction of cries and other calls.

Neonatal ACC lesions alter syntactical characteristics of social contact calls

We examined the syntax of vocal sequences to learn more about how early life ACC lesions might influence the acoustic signals that marmosets potentially relay to distantly located family members or other conspecifics when socially isolated. We focused on phee syllables, which are discrete elements or components of a call separated by very short intervals[39,40]. Thus, a sequence of phee calls comprises multiple syllables (**Fig 4A**). The functional significance of syllables is not clearly understood but a change in the number of syllables or their amplitude might feasibly alter the intensity of the message conveyed to a family that cannot be seen or heard. This is especially important to young infant monkeys that are naturally demanding of attention, and even more so if isolated. The number of syllables per phee call was highly variable amongst the animals, ranging from 1 to 8 syllables per phee.

The ACC lesion did not greatly affect the phee syllable count. Aside from a transient decline in the number of syllables in the week after the surgery (postnatal week 3: Wilcoxon test p=0.042), these animals showed the normal preferred range of 3-4 phee syllables at later postnatal weeks **(Fig 4B).** Even in the sixth postnatal week, when the proportion of phee and other contact calls was much lower in the ACC lesioned animals, the number of syllables in those phee calls that were issued was similar to the control group.

However, other phee call variables were affected by the lesion. For example, the duration of phee syllables was shortened in ACC-lesioned animals ($\chi2(5)$=13.27, p=0.021), particularly in the later syllables of a multi-syllablic phee. This effect emerged gradually and was most pronounced when the animals were 5-6 postnatal weeks (**Fig. 4C**). Likewise, the amplitude of phee syllables was also affected by the ACC lesion ($\chi2(5)$ = 48.178, p<0.0001), with lesioned animals making louder phee calls (**Fig 4D**). For each phee syllable, the amplitude difference between groups increased until postnatal week 5 and then disappeared at postnatal week 6 (postnatal week 4: β = 3017.75, 95% CI [1831.61, 4203.89] p < .001; postnatal week 5: β = 3719.00, 95% CI [2389.88, 5048.12], p < .001). Finally, we examined entropy of phee syllables as a measure of acoustic complexity. High entropy in multisyllabic phees would indicate that these vocalizations are diverse, variable and unpredictable. We found that animals with ACC lesions exhibited lower entropy in phee syllables relative to controls as early as postnatal week 3 ($\chi2(5)$ = 34.528, p < .0001) (**Fig 4E**) thereby expressing fixed, regular, and stereotyped phee calls. Together, our data suggest that the ACC lesion compromised the normal development of the phee signature for each monkey by making them shorter, louder, and monotonic.

Discussion

We found that early life ACC lesions led to rather specific alterations in the production of vocal calls, developmental changes to the quality of social contact calls, and their associated syntactical and acoustic characteristics were compromised. Contact calls that normally dominate the marmoset vocal repertoire at around 6 weeks were selectively diminished in the lesioned animals. When one common contact call, the phee call, was issued, its structural characteristics were unusual. The ACC lesions also led to permanent changes in remote brain areas known to be

involved in vocal behavior. Notably, the proportion of presumptive inhibitory interneurons was reduced in both the basomedial AMY and dorsal PAG. We can infer from these findings that an intact ACC in early life is integral to postnatal development of social vocalizations, and that its interactions with vocalization-eliciting sites from a very early age is fundamental to the normal vocal expression of social behavior.

Consistent with previous reports, the range of call types observed in both neonatal controls and neonatal ACC-lesioned animals within the first weeks of postnatal development were stereotyped and repetitive[33,41]. With increasing age, the call rate gradually declined such that by the time the animals were six weeks of age, the most common vocalizations were those that conveyed social distance. Such calls solicit attention from family members and may trigger a range of behaviors including search, approach, interaction, and caregiving in response to the need for social contact. This is especially so for infants whose well-being depends on social feedback and reciprocal interaction[4,5,35,42]. Our data suggest that the ability to effectively convey this social need was significantly altered in animals with early life ACC damage. At 6 weeks of age, these animals were not making social vocalizations at the same high rate as their age-matched controls. This reduction in social vocalizations does not appear to reflect a general slowing in vocal maturation, since other call types had advanced at the normal rate.

Since ACC lesions in humans cause social apathy[43,44], one possibility is that early life ACC removal altered the animals desire or motivation for social reinforcement; these infants appeared to make little effort in using vocalizations to solicit social contact when socially isolated. This change in call usage aligns with their social development period at around postnatal week 6 when infant marmosets transition from using fixed, stereotypical calls to a flexible and more

individualized call repertoire as they wean towards independence[33]. Our data suggest that this transition does not occur normally following an ACC lesion.

In addition, the neonatal ACC lesion altered the quality of the infants' long-distance phee calls; they were shorter in duration, louder in amplitude, and abnormal in their entropy such that the calls were blunted of variation, less diverse and more regular. This suggests that the social message conveyed by these infants over long distances to their families was deficient, limited, and/or indiscriminate because the calls were likely monotonic or stereotyped. The long-term behavioral implications of such imperfect vocalizations is currently unknown but could, ostensibly, affect their ability to use long distance social vocalizations to maintain intragroup functions such as warn of predators, strengthen family bonds, and maintain group cohesion more generally[2]. Since the ACC exerts regulation over autonomic responses[45–47], its ablation so early in life might feasibly blunt respiratory and vocalization responses in negative emotional environments such as social isolation. How these factors impact vocal behavior is a current topic of investigation (SheikhBahaei et al., *SfN* abstracts. 2023).

Although we found that an intact ACC in early life is integral to the postnatal maturation of social vocalizations, we also show that it is not critical for production of infant vocalizations more generally. A number of early observations reporting the loss of learned or spontaneous vocalizations following bilateral ACC lesions, left this question open, though it has been clear that vocal production in adults can withstand ACC damage[16,17,48–50]. Our findings indicate that innate vocal production in the earliest phases of life, as early as two weeks postnatally in the marmoset, is not critically dependent on the ACC. The infant 'babbling' behavior observed in marmosets and other primates[4,41,51] was largely preserved in the ACC-lesioned infant monkeys which, like the

control group, produced long sequences of concatenated calls composed of rudimentary features of mature adult-like calls.

While the ACC is not essential for infant vocal behavior, its absence affects not only the maturation of social vocal behavior, but also the anatomical compositions of structures with which it is interconnected. We noted a decrease in presumptive inhibitory interneurons in the dorsal PAG and basomedial AMY, two prominent ACC target regions involved in vocal behavior[7,20,52]. We can speculate that this reduction might stem from a prolonged deafferentation of cingulate inputs, gradually leading to a rebalancing of the excitatory/inhibitory elements in the local circuit.

One potentially related observation is that phee calls became louder in the weeks following the surgery. It is interesting to speculate that such amplitude increases might reflect a local decrease in inhibition in structures such as the AMY or PAG, whose activity is thought to tune the emotional characteristics of social vocalizations. The primate ACC receives dense projections from the basomedial AMY with notably minimal direct input from the lateral nucleus[53–55], and layer V pyramidal neurons project to the PAG[56,57] with a greater concentration directed to the dorsolateral column[58]. Thus, the ACC has the capacity to directly activate the brainstem vocalization pathway as early as the first few weeks of life. Both, the AMY and PAG are highly active during contexts in which threat related vocalizations would normally be triggered[59–61], and both regions elicit vocalizations through electrical stimulation or pharmacological disinhibition[12,62–64]. We cannot be sure when during postnatal development the ACC lesion altered GABA expression in the AMY and PAG, but from our results, we can infer that the appropriate modulation and coordination of social vocal behavior requires the normal postnatal development of the ACC.

Existing evidence in monkeys and humans demonstrate unequivocally the importance of the ACC in its contribution to emotional vocalization. In humans, as in monkeys, ACC lesions do not eliminate vocal behavior, but instead tend to remove the intonation and prosodic features of the vocalization characterized as expressionless[49]. This is consistent with the changes observed in the marmoset phee calls. In general, our data suggest that the ACC shapes the emotional structuring of social calls during the first few weeks of life in the marmoset. Given the many similarities to humans, and the strong contribution of socioemotional information to the vocal productions beginning in infancy, it is reasonable to speculate that similar mechanisms apply to the development of early life human vocal behavior. The ability to normalize brain circuits in early life would provide a major therapeutic advance for the remedial treatment of social deficits that plague disorders of mental health.

Materials and Methods

Subjects. All procedures accorded with the Guide for the Care and Use of Laboratory Animals and were approved by the Animal Care and Use Committee of the National Institute of Mental Health. A total of 10 marmosets (Callithrix jacchus), 5 males and 5 females, all born in captivity, were used in this study. Five infants received ACC lesions at 14-16 postnatal days old. Five others served as age-matched controls. The infants were raised by parents and siblings in family groups comprising 4-6 members and housed in temperature-controlled rooms (~27°C), 50-60% relative humidity under diurnal conditions (12h light:12h dark). Food and water were available ad libitum, supplemented with fresh fruit or vegetables. One animal showed sparing of the ACC lesion in one hemisphere. The final sample size for the behavioral data was n=5/per group. For the MRI and histological data, the final sample size was n=4/per group.

Surgery. We first obtained a reference MRI scan using a 14-day old ex-vivo sample **(Fig. 1A)**. A T2 weighted scan was obtained using 7T Bruker Biospin MRI platform with an eight-channel volume coil. Using ParaVision Acquisition 6.0.1, the following echo sequence was used to acquire a 3-dimensional volume of the infant marmoset brain: TR = 400, TE = 72ms, flip angle= 90 degrees, matrix size = 256x256x214, resolution = 0.15 mm isotropic, number of averages = 8, number of repetitions =1) and the total scan time was 3 hrs. The scan was aligned horizontally by rotating the image until the anterior and posterior commissures were positioned at the same height and water filled ear bars were used to obtain the interaural reference. We then used ITK-SNAP[65] (Yushkevich et al., 2006) to identify the anterior cingulate cortex (ACC) at the coronal planes before the genu of corpus callosum to the level of anterior commissure. The coordinates were calculated relative to the ear bars and midline references, both of which were visible on the scan. The resulting 14-day old marmoset scan served as a template atlas to calculate injection coordinates to target the rostral portion of the dorsal ACC (24a and 24b), bilaterally, in all marmosets. We calculated 5 injection coordinates for each hemisphere: (1) AP: 10.7 mm, ML: ±0.7 mm, DV: -2.9 mm; (2) AP: 10.7 mm, ML: ±0.7 mm, DV -4.5 mm; (3) AP: 9.5 mm, ML ±0.7 mm, DV -3.5 mm; (4) AP 8.5 mm, ML ±0.7 mm, DV -3.6 mm; (5) AP 7.5 mm, ML ±0.7 mm, DV -3.5 mm.

The entire surgical procedure was performed under aseptic conditions in infant monkeys that were 14-16 days old. During surgery, monkeys received isotonic fluids. Heart and respiration rates, body temperature, blood pressure, and expired CO_2 were monitored throughout the procedure. Pre- and post-operatively, monkeys received non-steroidal anti-inflammatory drugs, (meloxicam, 2mg/ml, s.c) to reduce swelling. The monkey was first immobilized with an anesthetic dose of alfaxalone (10mg/kg, i.m) combined with diazepam (5mg/kg, i.m). In this state,

the infant's head was shaved, and vital electrodes were secured on the infant's chest. Temperature was measured with a rectal probe. The infant's head was then secured in a small animal stereotaxic frame (Stoelting Company, Illinois) attached to a custom-built stage fitted with eye bars, ear bars and a pallet bar to accommodate the small head. Once the head was secured in the frame, anesthesia with isoflurane gas (1-2% to effect) was provided through the custom fitted mask **(Fig. 1B).** An integral part of the pallet bar was a gas hole (0.6 mm diameter) that ran along the length of the bar and connected, via tubing, to a vital monitor to measure small end tidal CO_2 volumes.

Following a midsagittal incision, the scalp was retracted, and a craniotomy was made above the target coordinates of the brain. A 5 µl syringe (33 gauge, Neurosyringe, Hamilton Company, Reno, NV) was used to administer bilateral injections of 0.12M NMDA (M3262, Sigma-Aldrich) dissolved in sterile filtered saline into the anterior cingulate cortex (0.5 µl per injection site). Each injection was made over 2 mins and the injector remained in place for an additional 4 mins for dispersion. When all injections were complete, the scalp was closed with intradermal absorbable sutures and the infant was allowed to recover in an intensive care unit that was void of extraneous sensory stimulation (e.g., excessive bright lights and loud noise). During recovery, marmosets received a combination of Esbalic and Enfamil (3:1 ratio) infant formula every 2-3 hours. When fully awake, each infant was returned to its family unit. A total of five marmosets received the neonatal cingulate lesion. Another five marmosets served as controls: two received saline injections (shams), one received a craniotomy only, and another two were unoperated.

Vocal recordings. Each infant was placed in a temperature-controlled incubator set to ~38°C (Thermocare, CA), and the emitted vocalization was recorded for 5 mins. Sound recordings were acquired using a cardioid microphone (Sennheiser ME 64, Sennheiser, Wedemark, Hanover, Germany) that was placed on the side of the incubator (Fig. 1C). The microphone was connected

to a computer and recorded sounds were digitized at a sampling frequency of 44 kHz using Raven Lite software (Cornell Lab of Ornithology, Ithaca, NY). Due to a variety of extraneous factors beyond our control including restrictions due to the COVID-19 pandemic, the exact day and number of recordings differed between monkeys. Therefore, recording sessions from each infant were grouped by week. All recording sessions were conducted without the presence of investigators in the recording room. The infant was then returned to its family unit.

Acoustic Analysis. The spectrograph of each audio file was obtained and visually inspected using Raven Pro 1.6 (Cornell Lab of Ornithology, Ithaca, NY). Spectrographs were generated with a Hann window of 512-sample points to filter the signal at 3dB bandwidth of 124 Hz (example, Fig. 2 A-B). The calls were manually classified by a defined classification system[37,51]. To identify call types in spectrograph of a recording, Raven software features, such as amplitude waveform, spectrograph and audio playbacks were used. Six major call types (phee, trill, trill-phee, twitter, cry, and complex calls) were identified from spectrographs, along with other minor call types (tsiks, chatter, egg, ock, seep). In some cases, when trill-phees and phees looked similar in spectrograph of a recording, acoustic parameters such as entropy were used to carefully classify calls. Complex calls, comprised vocalizations with elements from at least two different simple call types such as trill-twitter or twitter-phee, etc. From each recording, call types were manually annotated by 3 trained investigators (inter-rater reliability >80%). The spectrographs were used to obtain acoustic measurements such as peak frequency, RMS-amplitude and aggregated entropy and exported for further analysis.

Acoustic characteristics such as inter-call interval, syllable duration, number of phee syllables were quantified using a custom-written R script. Acoustic analyses were performed only for phees, which served as the major call type because of their abundance during postnatal weeks. In some

cases, vocalization quantity and amplitude was largely suppressed for several minutes after handling by the experimenters. Consequently, analysis of each recording began after 2 minutes had elapsed. Call counts, syllable count, temporal characteristics, and acoustics were quantified and analyzed.

Lesion Assessment with MRI. Gross ACC volume was measured from anesthetized MR scans performed at approximately 8 months of age. Anesthesia was induced with 5% isoflurane. The animals were then placed in an MR compatible cradle where its head was secured using ear bars. Isoflurane was maintained at 1.5 - 2.5% and vitals were monitored with V9004 Series Capnograph Monitor (San Clemente, CA). We obtained T2 weighted scan (n=4 Control and n=4 for ACC) using the MR procedure described above with the following echo sequence: TR = 30, TE = 48, matrix size = 144x144x128, resolution = 0.25 mm isotropic, number of averages = 8, number of repetitions =1. The rostro-caudal extent of the ACC was segmented and measured using ITK-SNAP 4.0. A representative sagittal view of the lesioned and non-lesioned ACC can be seen in Fig. 1E. Voxels containing ACC were carefully labeled from the anterior to posterior slice of the MR scan for each subject.

Histological preparation and quantification. At approximately 24 months of age, the marmosets were euthanized and perfused with 0.1 M PBS followed by 4% paraformaldehyde. The brains were extracted and cryoprotected in 0.1 M phosphate buffered sucrose (in steps of 10%, 20%, and 30% w/v). The brains were partitioned along the midline and right hemispheres were used for further histological processing after sectioning in coronal orientation on a sliding microtome and cryostat into 40 μm sections.

Immunohistochemistry. The immunohistochemistry (IHC) was performed on two series of free-floating sections. Initially, from each of the 8 brains, six sections each were collected around

three rostrocaudal planes at the following approximate locations (in reference to the interaural axis): +13.30 mm (target brain region: the anterior cingulate cortex); +9.20 mm (target brain region: the amygdala); and +2.05 mm (target brain region: the periaqueductal gray matter). Subsequently, these 18 sections were divided into two IHC series of 9 sections each, where three sections covered each target brain area, for separate processing. The first IHC series was processed to visualize major cell classes that are likely to be affected by a brain lesion (neurons: primary antibody against NeuN, astrocytes: primary antibody against the glial fibrillary acidic protein (GFAP), microglia/macrophages: primary antibody against the ionized calcium-binding adaptor molecule 1 (Iba1). The second IHC series was processed to visualize and assess the ratio of GABA-ergic neurons to all neurons: such changes in the relative number of inhibitory neurons could indicate local downstream neuroplasticity in reaction to the ACC lesion, as GABA-ergic neurons are primarily responsible for local inhibition. In the second IHC series we visualized the distribution of NeuN, as well as of neurotransmitter GABA and GAD67, a rate-limiting enzyme in GABA synthesis that produces more than 90% of GABA in the central nervous system.

The first IHC series was incubated in a cocktail of the following primary antibodies for 60 h at 4°C: NeuN (chicken, ABN91, Millipore Sigma), 1:500 dilution; GFAP (goat, SAB2500462, Millipore Sigma), 1:1,500 dilution; Iba1 (rabbit, 019-19741, Fujifilm Wako Chemicals), 1:1,500 dilution). Then after several washes the sections were incubated in a cocktail of the following secondary antibodies for 2 h at RT: 1:200 donkey anti-chicken Alexa Fluor 488 (703-545-155, Jackson ImmunoResearch), 1:500 donkey anti-goat Alexa Fluor 594 (A21207, Invitrogen), 1:400 donkey anti-rabbit Alexa Fluor 680 (711-625-152, Jackson ImmunoResearch). Additional antibodies used in the second IHC series were as follows. Primary: GABA (rabbit, A2052, Millipore Sigma), 1:600 dilution; GAD67 (mouse, MAB5406, Millipore Sigma), 1:500 dilution.

Secondary: 1:500 donkey anti-rabbit Alexa Fluor 594 (A21207, Invitrogen), 1:400 donkey anti-mouse Alexa Fluor 680 (A10038, Invitrogen). Finally, the sections were mounted on a glass slide, air-dried, and cover-slipped with DEPEX mounting media (13515, Electron Microscopy Sciences, Hatfield, PA).

Myelin staining Myelinated fibers were stained with aurohalophosphate-based Black-Gold II (Histo-Chem Inc., Jefferson, AR) as shown in Schmued et al.[66] according to the manufacturer's protocol and cover-slipped with Permount (SP15, Fisher Scientific).

Staining for degenerating neurons. Fluoro-Jade C (Histo-Chem Inc.) stain was used to visualize degenerating neurons according to the manufacturer's protocol in the target brain regions of ACC, amygdala, and PAG. There was no observable signal of neurodegeneration in studied regions.

Imaging and cell counting. Each histological section was digitized at 0.65 μm resolution (10✕ magnification) using a Zeiss Axioscan microscope slide scanner. Images were then split into a separate channel for each fluorophore. Cell detection and counting were done with an open source QuPath software 0.3.2.[67] . As each fluorescence channel was analyzed separately, the loci of immunofluorescence that were counted do not necessarily correspond to unique cells, especially for microglia and GABA channels where the fluorescent signal was more diffuse in appearance. Although the researcher performing cell counts was blinded to the marmosets' identity, it was possible to identify the site of the lesion and determine the animals group membership. Raw cell counts were transformed into cell densities per mm^2 to account for size differences in ROI areas.

Statistical analysis. All analyses were performed using R 4.2.3 (https://www.R-project.org/). For vocalization analysis, recording sessions were averaged by week and unless otherwise noted the data are represented as the mean values with mean ± confidence interval. Recording sessions

obtained before the surgery were grouped as postnatal week 2 (pre-surgery) and all the recordings after the surgery were binned into postnatal weeks 3 to 6 (post-surgery). Due to COVID-19 related restrictions, recordings for some infants could not be extended beyond postnatal week 3.

Packages in tidyverse library[68] were used for data processing and analyses. Linear mixed effect models (LMM) were used to analyze postnatal datasets, and these models were fitted with the lmer() function in lme4 package[69]. For estimation of coefficients, the maximum likelihood method was used. Models were fitted with postnatal weeks and experimental groups as fixed factors. To account for inter-individual variability, each monkey was modeled as a random effect. For multisyllabic phee analysis, syllables were nested within monkeys' random effect. Models with and without lesions were used to test the effect of ACC lesion and lesion effect is considered significant at an α of 0.05[70,71]. Model assumptions were tested using the check_model() function available in the performance package. Log transformation was performed on some dataset to meet LMM model assumptions. When normality assumption was violated, non-parametric test (Wilcoxon test) was also used. Graphs were created using the package ggplot2.

For analysis involving immunofluorescence there were some inhomogeneities in the spatial distribution of IHC signal. The raw immunopositive detections for specific markers were normalized by NeuN cell counts that were obtained in the same Qupath processing pipeline. The final measure for each antigen-specific immunopositivity count was the ratio computed from the number of antigen-positive detections divided by the sum of the antigen-positive detections and NeuN-positive cells. This transformation bounded the possible antigen-specific detections between 0 and 1 and allowed for parametric modeling with beta distribution. Cell counts were fitted using glmmTMB software package in R statistical computing environment. As there were 3 sections per animal for each region of interest, these were modeled as random effects. Widths of

major fiber tracts were modeled by a linear regression (function lm in base R). Final p values were

adjusted for multiple comparisons with the false discovery rate method for each antigen.

References

1. Eliades, S.J., and Miller, C.T. (2017). Marmoset vocal communication: Behavior and neurobiology. Dev. Neurobiol. *77*, 286–299. 10.1002/dneu.22464.

2. Oliveira, D.A.G., and Ades, C. (2004). Long-distance calls in Neotropical primates. An. Acad. Bras. Cienc. *76*, 393–398. 10.1590/s0001-37652004000200031.

3. Egnor, S.E.R., and Hauser, M.D. (2004). A paradox in the evolution of primate vocal learning. Trends Neurosci. *27*, 649–654. 10.1016/j.tins.2004.08.009.

4. Takahashi, D.Y., Fenley, A.R., Teramoto, Y., Narayanan, D.Z., Borjon, J.I., Holmes, P., and Ghazanfar, A.A. (2015). LANGUAGE DEVELOPMENT. The developmental dynamics of marmoset monkey vocal production. Science *349*, 734–738. 10.1126/science.aab1058.

5. Gultekin, Y.B., and Hage, S.R. (2017). Limiting parental feedback disrupts vocal development in marmoset monkeys. Nat. Commun. *8*, 14046. 10.1038/ncomms14046.

6. Gultekin, Y.B., and Hage, S.R. (2018). Limiting parental interaction during vocal development affects acoustic call structure in marmoset monkeys. Sci Adv *4*, eaar4012. 10.1126/sciadv.aar4012.

7. Jürgens, U. (2002). Neural pathways underlying vocal control. Neurosci. Biobehav. Rev. *26*, 235–258. 10.1016/s0149-7634(01)00068-9.

8. Miller, C.T., Iguina, C.G., and Hauser, M.D. (2005). Processing vocal signals for recognition during antiphonal calling in tamarins. Anim. Behav. *69*, 1387–1398. 10.1016/j.anbehav.2004.08.021.

9. Jürgens, U., and Pratt, R. (1979). The cingular vocalization pathway in the squirrel monkey. Exp. Brain Res. *34*, 499–510. 10.1007/BF00239145.

10. Sutton, D., Trachy, R.E., and Lindeman, R.C. (1981). Primate phonation: unilateral and bilateral cingulate lesion effects. Behav. Brain Res. *3*, 99–114. 10.1016/0166-4328(81)90031-0.

11. Kirzinger, A., and Jürgens, U. (1982). Cortical lesion effects and vocalization in the squirrel monkey. Brain Res. *233*, 299–315. 10.1016/0006-8993(82)91204-5.

12. Jürgens, U., and Ploog, D. (1970). Cerebral representation of vocalization in the squirrel

monkey. Exp. Brain Res. *10*, 532–554. 10.1007/BF00234269.

13. Müller-Preuss, P., Newman, J.D., and Jürgens, U. (1980). Anatomical and physiological evidence for a relationship between the "cingular" vocalization area and the auditory cortex in the squirrel monkey. Brain Res. *202*, 307–315. 10.1016/0006-8993(80)90143-2.

14. Jürgens, U. (1976). Reinforcing concomitants of electrically elicited vocalizations. Exp. Brain Res. *26*, 203–214. 10.1007/BF00238284.

15. Aitken, P.G. (1981). Cortical control of conditioned and spontaneous vocal behavior in rhesus monkeys. Brain Lang. *13*, 171–184. 10.1016/0093-934x(81)90137-1.

16. Sutton, D., Larson, C., and Lindeman, R.C. (1974). Neocortical and limbic lesion effects on primate phonation. Brain Res. *71*, 61–75. 10.1016/0006-8993(74)90191-7.

17. MacLean, P.D., and Newman, J.D. (1988). Role of midline frontolimbic cortex in production of the isolation call of squirrel monkeys. Brain Res. *450*, 111–123. 10.1016/0006-8993(88)91550-8.

18. Vogt, B.A., Pandya, D.N., and Rosene, D.L. (1987). Cingulate cortex of the rhesus monkey: I. Cytoarchitecture and thalamic afferents. J. Comp. Neurol. *262*, 256–270. 10.1002/cne.902620207.

19. Van Hoesen, G.W., Morecraft, R.J., and Vogt, B.A. (1993). Connections of the monkey cingulate cortex. In Neurobiology of Cingulate Cortex and Limbic Thalamus (Birkhäuser Boston), pp. 249–284. 10.1007/978-1-4899-6704-6_9.

20. Vogt, B.A., and Barbas, H. (1988). Structure and connections of the cingulate vocalization region in the rhesus monkey. In The Physiological Control of Mammalian Vocalization (Springer US), pp. 203–225. 10.1007/978-1-4613-1051-8_12.

21. Lloyd, R.L., and Kling, A.S. (1988). Amygdaloid electrical activity in response to conspecific calls in squirrel monkey (S. sciureus): Influence of environmental setting, cortical inputs, and recording site. In The Physiological Control of Mammalian Vocalization (Springer US), pp. 137–151. 10.1007/978-1-4613-1051-8_9.

22. Aggleton, J.P. (1993). The contribution of the amygdala to normal and abnormal emotional states. Trends Neurosci. *16*, 328–333. 10.1016/0166-2236(93)90110-8.

23. Gabriel, M. K., Foster, K., Orona, E., Saltwick, S. E., and Stanton, M., (1980). Neuronal activity of cingulate cortex, anteroventral thalamus, and hippocampal formation in discriminative conditioning: encoding and extraction of the significance of conditioned stimuli. In Progress in Psychobiology and Physiological Psychology. V. 9, J. M. Sprague and A. N. Epstein, eds. (Academic Press), pp. 126–223.

24. Müller-Preuss, P., and Jürgens, U. (1976). Projections from the "cingular" vocalization area in the squirrel monkey. Brain Res. *103*, 29–43. 10.1016/0006-8993(76)90684-3.

25. Hardy, S.G., and Leichnetz, G.R. (1981). Cortical projections to the periaqueductal gray in the monkey: a retrograde and orthograde horseradish peroxidase study. Neurosci. Lett. *22*, 97–101. 10.1016/0304-3940(81)90070-7.

26. Jürgens, U. (1994). The role of the periaqueductal grey in vocal behaviour. Behav. Brain Res. *62*, 107–117. 10.1016/0166-4328(94)90017-5.

27. An, X., Bandler, R., Ongür, D., and Price, J.L. (1998). Prefrontal cortical projections to longitudinal columns in the midbrain periaqueductal gray in macaque monkeys. J. Comp. Neurol. *401*, 455–479.

28. Simões, C.S., Vianney, P.V.R., de Moura, M.M., Freire, M.A.M., Mello, L.E., Sameshima, K., Araújo, J.F., Nicolelis, M.A.L., Mello, C.V., and Ribeiro, S. (2010). Activation of frontal neocortical areas by vocal production in marmosets. Front. Integr. Neurosci. *4*. 10.3389/fnint.2010.00123.

29. MacLean, P.D. (1985). Brain evolution relating to family, play, and the separation call. Arch. Gen. Psychiatry *42*, 405–417. 10.1001/archpsyc.1985.01790270095011.

30. Paxinos, G., Watson, C., Petrides, M., Rosa, M., and Tokuno, H. (2011). The Marmoset Brain in Stereotaxic Coordinates (Academic Press).

31. Atapour, N., Worthy, K.H., and Rosa, M.G.P. (2021). Neurochemical changes in the primate lateral geniculate nucleus following lesions of striate cortex in infancy and adulthood: implications for residual vision and blindsight. Brain Struct. Funct. *226*, 2763–2775. 10.1007/s00429-021-02257-0.

32. Zhang, Y.S., and Ghazanfar, A.A. (2018). Vocal development through morphological computation. PLoS Biol. *16*, e2003933. 10.1371/journal.pbio.2003933.

33. Gultekin, Y.B., Hildebrand, D.G.C., Hammerschmidt, K., and Hage, S.R. (2021). High plasticity in marmoset monkey vocal development from infancy to adulthood. Sci Adv *7*. 10.1126/sciadv.abf2938.

34. Ghazanfar, A.A., Flombaum, J.I., Miller, C.T., and Hauser, M.D. (2001). The units of perception in the antiphonal calling behavior of cotton-top tamarins (Saguinus oedipus): playback experiments with long calls. J. Comp. Physiol. A *187*, 27–35. 10.1007/s003590000173.

35. Miller, C.T., and Wang, X. (2006). Sensory-motor interactions modulate a primate vocal behavior: antiphonal calling in common marmosets. J. Comp. Physiol. A Neuroethol. Sens. Neural Behav. Physiol. *192*, 27–38. 10.1007/s00359-005-0043-z.

36. Bakker, J., and Langermans, J.A.M. (2018). Ultrasonic components of vocalizations in marmosets. In Handbook of Ultrasonic Vocalization - A Window into the Emotional Brain Handbook of behavioral neuroscience. (Elsevier), pp. 535–544. 10.1016/b978-0-12-809600-0.00049-4.

37. Bezerra, B.M., and Souto, A. (2008). Structure and Usage of the Vocal Repertoire of Callithrix jacchus. Int. J. Primatol. *29*, 671–701. 10.1007/s10764-008-9250-0.

38. Liao, D.A., Zhang, Y.S., Cai, L.X., and Ghazanfar, A.A. (2018). Internal states and extrinsic factors both determine monkey vocal production. Proc. Natl. Acad. Sci. U. S. A. *115*, 3978–3983. 10.1073/pnas.1722426115.

39. Devinsky, O., Morrell, M.J., and Vogt, B.A. (1995). Contributions of anterior cingulate cortex to behaviour. Brain *118 (Pt 1)*, 279–306. 10.1093/brain/118.1.279.

40. Agamaite, J.A., Chang, C.-J., Osmanski, M.S., and Wang, X. (2015). A quantitative acoustic analysis of the vocal repertoire of the common marmoset (Callithrix jacchus). J. Acoust. Soc. Am. *138*, 2906–2928. 10.1121/1.4934268.

41. Elowson, A.M., Snowdon, C., and Lazaro-Perea, C. (1998). Infant "babbling" in a nonhuman primate: Complex vocal sequences with repeated call types. Behaviour *135*, 643–664. 10.1163/156853998792897905.

42. Takahashi, D.Y., Liao, D.A., and Ghazanfar, A.A. (2017). Vocal Learning via Social Reinforcement by Infant Marmoset Monkeys. Curr. Biol. *27*, 1844–1852.e6. 10.1016/j.cub.2017.05.004.

43. Eslinger, P.J., and Damasio, A.R. (1985). Severe disturbance of higher cognition after bilateral frontal lobe ablation: patient EVR. Neurology *35*, 1731–1741. 10.1212/wnl.35.12.1731.

44. Damasio, A.R., Tranel, D., and Damasio, H. (1990). Individuals with sociopathic behavior caused by frontal damage fail to respond autonomically to social stimuli. Behav. Brain Res. *41*, 81–94. 10.1016/0166-4328(90)90144-4.

45. Robinson, B.W., and Mishkin, M. (1968). Alimentary responses to forebrain stimulation in monkeys. Exp. Brain Res. *4*, 330–366. 10.1007/BF00235700.

46. Neafsey, E.J. (1990). Prefrontal cortical control of the autonomic nervous system: anatomical and physiological observations. Prog. Brain Res. *85*, 147–165; discussion 165–166. 10.1016/s0079-6123(08)62679-5.

47. Buchanan, S.L., and Powell, D.A. (1993). Cingulothalamic and prefrontal control of autonomic function. In Neurobiology of Cingulate Cortex and Limbic Thalamus (Birkhäuser Boston), pp. 381–414. 10.1007/978-1-4899-6704-6_14.

48. Trachy, R.E., Sutton, D., and Lindeman, R.C. (1981). Primate phonation: Anterior cingulate lesion effects on response rate and acoustical structure. Am. J. Primatol. *1*, 43–55. 10.1002/ajp.1350010106.

49. von Cramon, D., and Jürgens, U. (1983). The anterior cingulate cortex and the phonatory control in monkey and man. Neurosci. Biobehav. Rev. *7*, 423–425. 10.1016/0149-7634(83)90049-0.

50. Jürgens, U. (1998). Neuronal control of mammalian vocalization, with special reference to the squirrel monkey. Naturwissenschaften *85*, 376–388. 10.1007/s001140050519.

51. Pistorio, A.L., Vintch, B., and Wang, X. (2006). Acoustic analysis of vocal development in a New World primate, the common marmoset (Callithrix jacchus). J. Acoust. Soc. Am. *120*, 1655–1670. 10.1121/1.2225899.

52. Jürgens, U. (2009). The neural control of vocalization in mammals: a review. J. Voice *23*, 1–10. 10.1016/j.jvoice.2007.07.005.

53. Amaral, D.G., and Price, J.L. (1984). Amygdalo-cortical projections in the monkey (Macaca fascicularis). J. Comp. Neurol. *230*, 465–496. 10.1002/cne.902300402.

54. Carmichael, S.T., and Price, J.L. (1996). Connectional networks within the orbital and medial prefrontal cortex of macaque monkeys. J. Comp. Neurol. *371*, 179–207. 10.1002/(SICI)1096-9861(19960722)371:2<179::AID-CNE1>3.0.CO;2-#.

55. Aggleton, J.P., Wright, N.F., Rosene, D.L., and Saunders, R.C. (2015). Complementary Patterns of Direct Amygdala and Hippocampal Projections to the Macaque Prefrontal Cortex. Cereb. Cortex *25*, 4351–4373. 10.1093/cercor/bhv019.

56. Mantyh, P.W. (1982). Forebrain projections to the periaqueductal gray in the monkey, with observations in the cat and rat. J. Comp. Neurol. *206*, 146–158. 10.1002/cne.902060205.

57. Morrell, J.I., Greenberger, L.M., and Pfaff, D.W. (1981). Hypothalamic, other diencephalic, and telencephalic neurons that project to the dorsal midbrain. J. Comp. Neurol. *201*, 589–620. 10.1002/cne.902010409.

58. Bandler, R. (1988). Brain mechanisms of aggression as revealed by electrical and chemical stimulation: Suggestion of a central role for the midbrain periaqueductal grey region. In Progress in Psychobiology and Physiological Psychology. V. 13, A. N. Epstein and A. R. Morrison, eds. (Academic Press), pp. 67–154.

59. Fecteau, S., Belin, P., Joanette, Y., and Armony, J.L. (2007). Amygdala responses to nonlinguistic emotional vocalizations. Neuroimage *36*, 480–487. 10.1016/j.neuroimage.2007.02.043.

60. Mobbs, D., Petrovic, P., Marchant, J.L., Hassabis, D., Weiskopf, N., Seymour, B., Dolan, R.J., and Frith, C.D. (2007). When fear is near: threat imminence elicits prefrontal-periaqueductal gray shifts in humans. Science *317*, 1079–1083. 10.1126/science.1144298.

61. Reis, F.M.C.V., Liu, J., Schuette, P.J., Lee, J.Y., Maesta-Pereira, S., Chakerian, M., Wang, W., Canteras, N.S., Kao, J.C., and Adhikari, A. (2021). Shared Dorsal Periaqueductal Gray Activation Patterns during Exposure to Innate and Conditioned Threats. J. Neurosci. *41*, 5399–5420. 10.1523/JNEUROSCI.2450-20.2021.

62. Robinson, B. (1967). Vocalization evoked from forebrain in Macaca mulatta☆. Physiol. Behav. *2*, 345–354. 10.1016/0031-9384(67)90050-9.

63. Jürgens, U., and Lu, C.L. (1993). The effects of periaqueductally injected transmitter antagonists on forebrain-elicited vocalization in the squirrel monkey. Eur. J. Neurosci. *5*, 735–741. 10.1111/j.1460-9568.1993.tb00537.x.

64. Forcelli, P.A., Waguespack, H.F., and Malkova, L. (2017). Defensive Vocalizations and Motor Asymmetry Triggered by Disinhibition of the Periaqueductal Gray in Non-human Primates. Front. Neurosci. *11*, 163. 10.3389/fnins.2017.00163.

65. Yushkevich, P.A., Piven, J., Hazlett, H.C., Smith, R.G., Ho, S., Gee, J.C., and Gerig, G. (2006). User-guided 3D active contour segmentation of anatomical structures: significantly improved efficiency and reliability. Neuroimage *31*, 1116–1128. 10.1016/j.neuroimage.2006.01.015.

66. Schmued, L., Bowyer, J., Cozart, M., Heard, D., Binienda, Z., and Paule, M. (2008). Introducing Black-Gold II, a highly soluble gold phosphate complex with several unique advantages for the histochemical localization of myelin. Brain Res. *1229*, 210–217. 10.1016/j.brainres.2008.06.129.

67. Bankhead, P., Loughrey, M.B., Fernández, J.A., Dombrowski, Y., McArt, D.G., Dunne, P.D., McQuaid, S., Gray, R.T., Murray, L.J., Coleman, H.G., et al. (2017). QuPath: Open source software for digital pathology image analysis. Sci. Rep. *7*, 16878. 10.1038/s41598-017-17204-5.

68. Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., Grolemund, G., Hayes, A., Henry, L., Hester, J., et al. (2019). Welcome to the tidyverse. J. Open Source Softw. *4*, 1686. 10.21105/joss.01686.

69. Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects models Usinglme4. J. Stat. Softw. *67*. 10.18637/jss.v067.i01.

70. Winter, B. (2013). Linear models and linear mixed effects models in R with linguistic applications. 10.48550/ARXIV.1308.5499.

71. Brown, V.A. (2021). An introduction to linear mixed-effects modeling in R. Adv. Methods Pract. Psychol. Sci. *4*, 251524592096035. 10.1177/2515245920960351.

Acknowledgements

This research was supported by the Intramural Research Program of the National Institute of Mental Health (ZIAMH002951 and ZICMH002952 to YC). We thank George Dold, William Bennett, and David Ide from the NIMH Section on Instrumentation for customization of the stereotaxic frame and surgical anesthesia gas mask. We would also like to thank the Veterinary

Declaration of interests

The authors declare that they have no competing interests.

Author Contributions (CRediT)

Conceptualization and methodology GN, YC; Software GN, DM, ACP, SPB; Formal analysis GN, DM; Investigation GN, DM, ACP; Resources YC; visualization GN, DM, YC; writing original draft GN, YC; Writing review and editing GN, DM, ACP, SPB, YC; Supervision DM, YC.

Data Availability

All data needed to evaluate the conclusions in the paper are present in the paper.

Figures

Fig. 1. Experimental Design and timeline.

(A) Reference MR scan was obtained using an ex-vivo sample of a 14-day old marmoset. Parasagittal view of the reference scan shows location of injection coordinates targeting the rostral portion of the dorsal ACC (24a and 24b), bilaterally (red). **(B)** Gas anesthesia was supplied

through a custom-made breathing system comprising a facemask fitted with a palate bar with a 0.6mm diameter hole. The palate bar was connected to a vital monitor to accurately detect small tidal end volumes during anesthesia while the animal was secured in the stereotaxic frame. **(C)** Five-minute vocalization recordings were obtained from infant placed in a softly padded temperature-controlled incubator. **(D)** Timeline of vocal recordings from postnatal week 2 to postnatal week 6. The ACC lesion was conducted at postnatal week 2 when animals were 14-16 days old. **(E)** Representative sagittal view of postoperative T2-weighted MR images of a control (left panel) and lesioned (right panel) infant to reveal extent of white hypersignal, which reflects edema due to injections of the excitotoxin and therefore approximate site of the ACC lesion. There was a significant reduction in total ACC volume in the ACC group relative to controls (n=4/per group; $F(1,6) = 82.78$, $p<0.0001$). A representative 3-dimensional view of area 24 is presented showing the reduced volume of the ACC (right panel) relative to the normal volume in the control (left panel). (F) Schematic illustration highlights the end point of the experiment involving histological processing and evaluation of cell markers. (G) Longitudinal timeline shows approximate age of animals following vocal recordings, MRI lesion assessments and histological processing.

Fig. 2. Lesion verification and impact of early life ACC lesion on vocal downstream

One lesioned animal showed sparing of the lesion in one hemisphere, so the lesion verification was performed in 4 animals only. (A) Left panel shows a sagittal section from the standard marmoset brain depicting the intended ACC lesion shaded in red. The right panel shows schematic lesion reconstructions superimposed on a sagittal and coronal marmoset brain section depicting the extent of the ACC lesion shaded in red. Regions that appear darker indicate greater overlap in

the damage present among different animals. (B) Left column shows the unilateral coronal section from the standard marmoset brain depicting area 24 of ACC lesion site. The right column shows magnified images of area 24 stained to visualize myelinated fibers in a representative control and ACC-lesioned animal. The normal radial arrangement of the myelinated fibers is disrupted following the ACC lesion. (C) Histological quantification of mature neurons and glia in representative ACC lesioned animal (top panel) and control (bottom panel). Magnified images show high levels of astrocytes (violet) and microglia/macrophages (white) surrounding the lesion site at the grey and white matter interface in the ACC-lesioned animal relative to the control. (D) Left panel shows grayscale images with anti-NeuN staining depicting divisions of AMY and PAG where relative distribution of GABA-positive immunoreactive expression were quantified. Right graphs show the proportion of GABA expression in each division depicted in the AMY and PAG. Each circle represents one animal (ACC lesioned animal is red, Control animal is blue). Mean expression is represented by black bar. Data for two animals in ACC group overlap for basomedial AMY quantification. GABA-positive immunoreactivity was significantly down in the basomedial AMY and dorsal PAG.

Fig 3. ACC in early life is integral to postnatal development of social contact calls

(A-B) Spectrograms show sample 30-second vocal recordings of a representative control and ACC-lesioned marmosets before (postnatal week 2) and after surgery (postnatal week 6). Before surgery, the infants 'babbled' by emitting a wide range of immature concatenated calls, each with its own spectrographic motif illustrated and labeled in boxes. After surgery, at postnatal week 6, calls show reduced variability separated by distinct gaps or inter-call intervals. (C) Both groups show a reduction in the relative call count with increasing age. (D) Animals in both groups were

able to emit calls of different call types. Those with ACC lesions made minor calls designated as 'other' more frequently than controls but all major call types were produced at equivalent rates. (E) Despite their ability to produce all call types, the proportion of social contact calls comprising phee, twitter and trills, was substantially reduced in animals with early life ACC lesions at postnatal week 6. (F) Chord diagrams show that at postnatal week 6, animals with ACC lesions show a high probability of transitions between all call types with lower frequency of transitions between social contact calls. The chord diagram represents weighted probabilities of transitions and their directionality from each group of call types. Weighted probabilities were used because of the variability in call counts. The size or thickness of arrows/links represents probabilities of call transition and the numbers around each chord diagram represents relative probability value for each call type transition.

Fig. 4. ACC lesion alters structural characteristics of long distance social phee calls

(A) Sample spectrograph with examples of three and five syllable phees. (B) The ACC lesion caused a reduction in average phee syllable counts immediately after the ACC lesion (red dots) at PW3 but then normalized to 3-4 syllables thereafter. (C) Phee syllable duration ACC-lesioned group became shorter for multisyllabic phees $\geq$ 3 especially with increasing age. (D) The effective amplitude for each phee syllable increased for the ACC group until postnatal week 6. (E) Animals with ACC lesions emitted low entropy phees for calls as low as 2-syllables and continued until postnatal week 6. Error bars are confidence intervals. Gray shaded lines or bars represent time of surgery.

Figure 1

A MRI guided surgery using a 14-day old ex-vivo sample

B Breathing system connected to stereotaxic frame

CO₂ Expiration

Isoflurane

C Temperature controlled incubator with microphone

D Timeline of vocal recordings

Postnatal age

Pre-Surgery

Surgery

Post-Surgery

week 2

week 3

week 4

week 5

week 6

E Lesion assessment with MRI

Volume (mm³)

Control   ACC lesion

Control   ACC lesion

F Evaluation of cell markers

G

~1.5 Month

Vocal Recording

~8 Months

T2 MRI scan

~24 months

Histological processing and quantification

Figure 2

A Intended lesion ACC lesion

Number of animals with overlapping lesions
1   4

B Myelinated fibres
CON   ACC lesion

C Mature neurons and glia
NeuN(green)-GFAP(violet)-Iba1(white)

Grey and white matter interface

D
Divisions of amygdala

Divisions of periaqueductal gray

dorsal
ventral

GABA-positive immunoreactivity in the amygdala and periaqueductal gray

Proportion of GABA-positive detections

Central amygdala   Basomedial amygdala   Basolateral amygdala   Lateral amygdala   Dorsal periaqueductal gray   Ventral periaqueductal gray

NS.   **   NS.   NS.   *   NS.

CON ACC   CON ACC   CON ACC   CON ACC   CON ACC   CON ACC

● CON ● ACC

Figure 3

Figure 4