Title: Extracellular vesicles alter trophoblast function in pregnancies complicated by COVID-19

One Sentence Summary: Circulating extracellular vesicles are altered by COVID-19 during

pregnancy and contribute to placental dysfunction.

Authors: Thea N. Golden[1,2,3*], Sneha Mani[1,2], Rebecca L. Linn[4], Rita Leite[1,2], Natalie A. Trigg[5],

Annette Wilson[6], Lauren Anton[1,2], Monica Mainigi[1,2], Colin C. Conine[2,5,7,8,9,10], Brett A.

Kaufman[6], Jerome F. Strauss III[1,2], Samuel Parry[1,2], and Rebecca A. Simmons[2,3,9,10]

Affiliations:

[1]Department of Obstetrics and Gynecology, Perelman School of Medicine at the University of

Pennsylvania; Philadelphia, USA.

[2]Center for Research on Reproduction and Women's Health, University of Pennsylvania;

Philadelphia, USA.

[3]Center for Excellence in Environmental Toxicology, University of Pennsylvania; Philadelphia,

USA.

[4]Department of Pathology and Laboratory Medicine, Children's Hospital of Philadelphia;

Philadelphia, USA.

[5]Epigenetics Institute, Perelman School of Medicine at the University of Pennsylvania;

Philadelphia, USA.

[6]Department of Medicine, University of Pittsburgh; Pittsburgh, USA.

[7]Institute for Regenerative Medicine, Perelman School of Medicine at the University of

Pennsylvania; Philadelphia, USA.

[8]Department of Genetics, Perelman School of Medicine at the University of Pennsylvania;

Philadelphia, USA.

[9]Department of Pediatrics, Perelman School of Medicine at the University of Pennsylvania;

Philadelphia, USA.

[10]Division of Neonatology, Children's Hospital of Philadelphia, Philadelphia, USA.

*Corresponding author. Email: goldent@pennmedicine.upenn.edu

Abstract:

Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection and resulting

coronavirus disease (COVID-19) causes placental dysfunction, which increases the risk of

adverse pregnancy outcomes. While abnormal placental pathology resulting from COVID-19 is

common, direct infection of the placenta is rare. This suggests that pathophysiology associated

with maternal COVID-19, rather than direct placental infection, is responsible for placental

dysfunction and alteration of the placental transcriptome. We hypothesized that maternal

circulating extracellular vesicles (EVs), altered by COVID-19 during pregnancy, contribute to

placental dysfunction. To examine this hypothesis, we characterized maternal circulating EVs

from pregnancies complicated by COVID-19 and tested their effects on trophoblast cell

physiology *in vitro*. We found that the gestational timing of COVID-19 is a major determinant

of circulating EV function and cargo. *In vitro* trophoblast exposure to EVs isolated from patients

with an active infection at the time of delivery, but not EVs isolated from Controls, altered key

trophoblast functions including hormone production and invasion. Thus, circulating EVs from

participants with an active infection, both symptomatic and asymptomatic cases, can disrupt vital

trophoblast functions. EV cargo differed between participants with COVID-19 and Controls,

which may contribute to the disruption of the placental transcriptome and morphology. Our

findings show that COVID-19 can have effects throughout pregnancy on circulating EVs and

circulating EVs are likely to participate in placental dysfunction induced by COVID-19.

INTRODUCTION

Maternal SARS-CoV-2 infection and resulting coronavirus disease (COVID-19) is

associated with an increased risk of pregnancy complications including preterm birth,

hypertensive disorders of pregnancy, fetal growth restriction, and pregnancy loss (*1, 2*).

Placental dysfunction is known to contribute to these complications, and placental pathology,

including vasculopathies and inflammation, is frequently reported following an acute or even

resolved infection during pregnancy (*3-6*). This suggests COVID-19 has a long-lasting effect on

pregnancy by altering placenta function. Despite extensive reports of placental abnormalities

following COVID-19, little is known about the underlying mechanisms contributing to placental

dysfunction and the related subsequent pregnancy complications. Direct infection of the

placenta is rare, which suggests that placental dysfunction is caused by the maternal response to

SARS-CoV-2 infection (*6-8*).

Circulating extracellular vesicles (EVs) are altered by SARS-CoV-2 infection and

contribute to COVID-19-induced organ damage (*9-13*). EVs are a means of cell-to-cell

communication resulting from their ability to carry and transfer bioactive cargo that elicits

signaling events in recipient cells. Compared to uninfected individuals, EV cargo composition in

patients with COVID-19 is significantly different, eliciting downstream systemic effects such as

coagulopathy (*14-17*) and inflammation (*10, 11, 14, 18*). For example, tissue factor protein

abundance in EVs is increased in COVID-19 and correlates with inflammation and disease

severity demonstrating the influence that circulating EVs have on the systemic response to

COVID-19, thereby leading to organ dysfunction (*9, 12, 13*).

During normal pregnancy, the placenta releases EVs into the maternal circulation (*19,*

20). Placental-derived EVs promote maternal adaptation to support a healthy pregnancy

including a shift in the maternal immune system to a tolerant state and the promotion of

angiogenesis (*21, 22*). Placental-derived EVs also affect trophoblast function through autocrine

and paracrine signaling. Trophoblasts are a specialized cell type of the placenta that are

responsible for invasion into maternal tissue to anchor the placenta, vascular remodeling for

adequate placental blood flow, nutrient transport, and hormone production for maternal and fetal

signaling. EV signaling impairs normal trophoblast function which is thought to contribute to

the placental dysfunction underlying pregnancy complications (*21, 23-26*).

Therefore, we hypothesized that COVID-19 alters circulating EV cargo which has a

functional consequence in the placenta. Similar to previous studies, we found that COVID-19

during pregnancy induced marked placental histologic and transcriptomic changes that were

dependent on the gestational timing of maternal SARS-CoV-2 infection. Importantly, we found

that COVID-19 altered EV cargo, and trophoblast exposure to these EVs resulted in reduced

trophoblast invasion and hormone production.

RESULTS

Pregnancy outcomes following active and resolved SARS-CoV-2 infection

Participants were enrolled at the time of delivery between July 2020-August 2022.

Controls had no known SARS-CoV-2 infection during pregnancy. COVID-19 cases were

divided based on the gestational timing of infection and either had a resolved infection that

occurred in the 1st, 2nd, or 3rd trimester (R1, R2, R3), or had an active infection at the time of

delivery (AI) (Table 1). All patients admitted to the labor and delivery unit underwent PCR

testing for SARS-CoV-2 at admission. There were no significant differences in maternal age

between groups, but there were significantly more Black participants with an infection in the

third trimester (resolved or active) compared to non-infected individuals (Controls) (R3 p=0.046,

AI p=0.011). Participants with an infection in the third trimester, resolved or active, also had an

earlier gestational age at delivery than Controls (R3 p<0.001, AI p=0.034) (Table 1). Not

surprisingly, the AI group, which underwent universal screening for SARS-CoV-2 at admission

for delivery, had a higher incidence of asymptomatic infection than those with resolved infection

(p<0.0001).

Multiple studies have shown that SARS-CoV-2 infection during pregnancy is associated

with adverse pregnancy outcomes. However, to date, no study has examined the relationship

between the timing of infection and pregnancy complications in a single cohort. Our study is

limited by relatively small numbers (n=15-32) in each participant group, but we found that the

type of adverse outcome differed depending on the gestational timing of infection. The

incidence of gestational hypertension (gHTN) was increased in R1 (p=0.016), whereas

spontaneous preterm birth (SPTB) was increased in R2 (p=0.008) compared to Controls.

Preeclampsia (PE) and medically indicated preterm birth (MPTB) were increased in R3 and AI

compared to Controls (PE: R3 p=0.032, AI p= 0.010, MPTB: R3 p=0.005, AI p=0.010) (Table

1). In our cohort, there was no increase in intrauterine growth restriction (IUGR) or intrauterine

fetal demise (IUFD) which have been reported in pregnancies complicated by COVID-19 (*27-*

29).

Placental pathology in COVID-19

Abnormal placental pathology is commonly reported in patients with active and resolved

SARS-CoV-2 infections (*3*). However, there has yet to be a comprehensive assessment of

placenta morphology following maternal infection at various gestational ages. Similar to

previous reports (*3-6, 30, 31*), we found maternal and fetal vascular malperfusion (MVM &

FVM) lesions were increased among patients with pregnancies complicated by COVID-19

(Table 1). Interestingly, high-grade MVM and perivillous fibrin deposition were increased in

participants with a resolved infection that occurred in the second or third trimester, but not the

first trimester or with an active infection. This suggests that it takes a substantial amount of time

for the placenta to recover from the impact of COVID-19. Alternatively, it is possible that the

first trimester placenta was more resilient. The lack of high-grade MVM and perivillous fibrin

deposition in placenta collected from participants with an active infection suggests these lesions

may take weeks to manifest.

Timing of COVID-19 impacts the placental transcriptome

To gain insight into potential novel pathways that might be affected by COVID-19 during

pregnancy, we performed RNA-seq on the placental transcriptome using biopsies from placenta

collected at delivery in COVID-19 cases and controls. Not surprisingly, active infection was

associated with significant differences in gene expression in the placenta. There were 72

upregulated genes and 384 downregulated genes in AI compared to Control placentas. (Figure 1,

Supplemental Table 1). Gene expression was also altered in recovered infections, but the

magnitude of change was smaller; 28, 2, and 56 differentially expressed genes (DEGs)

comparing R1, R2, and R3 to Controls, respectively. It's worth noting that in each of the 4

groups, there was differential expression of genes that regulate mitochondria activity, which

implies a common dysfunctional pathway resulting from COVID-19.

In addition, several other genes were differentially expressed in one or more COVID-19

groups. There were several transcripts that were increased in placenta from R1 and AI groups

compared to Controls. Several of these genes regulate fibrosis (*RORB*, *FN1*, *IGFB6*, *MMP12*,

and *AC027288.3*) (*32-36*) suggesting that this pathway plays an important role in placenta

pathology in COVID-19. Moreover, MMP12 regulates spiral artery remodeling and reduced

MMP12 activity contributes to the development of preeclampsia (*37*). Similarly, increased

expression of *FN1* slice variants containing Extra Domain A promotes inflammation via Toll-

like receptor 4 (TLR4) activation (*38, 39*) and is associated with an increased risk of

preeclampsia (*32, 38, 39*). Dysregulation of these pathways may contribute to the pathogenesis

of preeclampsia in pregnant individuals with COVID-19. Finally, *GREB1* was upregulated in R1

and AI placenta compared to Controls. GREB1 interacts with the progesterone (P4) receptor to

regulate P4 responsive genes (*40, 41*) and GREB1 promotes maternal tissue remodeling (*40*).

Several of the downregulated genes in AI compared to Controls were also

downregulated in R3 placenta compared to Controls (*PRELP*, *MEG9*, *VSTM4*, *CLIC2*, *C7*, and

CNN1). Interestingly, *CNN1*, which encodes calponin, is expressed by smooth muscle cells and

expression changes are associated with spiral artery remodeling (*42, 43*). Further, SARS-CoV-2

infection is known to disrupt complement pathways (*44*) and the gene encoding complement C7

was downregulated in R3 and AI placenta compared to Controls. The complement system plays

a dual role in pregnancy in that it protects the placenta from pathogen infection and participates

in spiral artery remodeling (*45*). While the effect of decreased *C7* in the placenta is unknown, it

may indicate that an imbalance in the complement system contributes to placental pathology

observed in R3 and AI placentas. *STAT4*, which encodes the signal transducer and activator of

transcription 4, is a key activator of immune regulating genes and was upregulated in R3 and AI

placentas compared to Controls. STAT4 mediated pathways are disrupted in preeclampsia and

circulating levels are elevated in patients with preeclampsia (*46, 47*).   This differential gene

expression was only seen in the placenta if the maternal infection was active or recently resolved,

suggesting a long period of time is necessary to attenuate these pathways.

162   Statistically significant disruptions in canonical pathways, as determined by Ingenuity

Pathway Analysis (IPA), include inflammation and fibrosis in the placentas from R1, R3, and AI

compared to Controls (Supplemental Table 2). This unbiased approach based on differential

gene expression supported our placental histopathology findings of inflammation and fibrosis in

these placentas. Thus, despite the long period of time following the resolution of maternal

SARS-CoV-2 infection, genes that regulate fibrosis and inflammation were altered, and

pathological evidence of fibrosis and inflammation were apparent in the placenta regardless of

the timing of infection.

170   We also used IPA to predict transcriptional regulators of genes differentially expressed in

171   the placenta from pregnancies complicated by COVID-19. Interestingly, genes that encode for

172   growth factors (IGF1, *IGF2*, *FGF3, FGF19, TGFB1*), immune-regulating proteins (*JUN, TNF,*

173   *IL1B, IL13, IL6, IL10, IL4, IFNG*) and hormone-regulating proteins (*PRLH, LEPR, ESR1, and*

174   *PGR*) were the top predicted transcriptional regulators (Supplemental Table 3).

175   **Sustained effects on circulating EVs following COVID-19 in early pregnancy**

176   As discussed above, SARS-CoV-2 rarely infects the placenta, implying a distal signal.

We hypothesized that maternal circulating EVs play a role in mediating placental dysfunction

associated with COVID-19. Therefore, we characterized EVs isolated from maternal plasma

collected at delivery to determine if COVID-19 altered the EV profile. We isolated large (LEV)

and small (SEV) EVs as they carry distinct cargo with distinct functional effects. We confirmed

the presence of large and small EVs in isolated particles by electron microscopy (Supplemental

Figure 1A). Large and small EVs had the expected size distribution of EVs (Supplemental

Figure 1B) and abundant expression of the EV-related tetraspanin CD9 (Supplemental Figure

1C).

EV characteristics, including concentration and size distribution, revealed long-lasting

alterations in patients with a resolved infection. The diameter of small EVs was significantly

increased in resolved infections compared to Controls (108.7nm vs. 117.2nm, p=0.023).

However, the difference in concentration was not significant (2.15×10^8 vs. 1.56×10^8 EVs/uL

plasma, p=0.074). When resolved infections were categorized by timing during gestation, we

found that small EVs isolated from R2 participants had an increased diameter and reduced

concentration, but small EVs isolated from R1 and R3 participants were not different from

Controls (Figure 2 D&E). The diameter of large EVs isolated from R2 patients was decreased

but there was no change in their number (Figure 2 A&B). COVID-19 in the second trimester was

uniquely associated with alterations in circulating EV concentration and size at the time of

delivery.

Altered cell of origin of EVs in patients with COVID-19 in early pregnancy

Characterizing the source of circulating EVs provides biological information about the

tissue and cell-type of origin and its functional state. We used flow cytometry to detect cell-

specific vesicle membrane protein expression and identified the relative contribution of each EV

tissue-cellular source. We identified EVs that originated from maternal endothelial cells (CD31+

CD34-), fetal endothelial cells (CD31+ CD34+), platelets (CD41a+), immune cells (CD45+), and

trophoblasts (PLAP+) (Figure 2 C&F).

Trophoblast-derived PLAP+ EVs comprised the largest proportion of circulating large

EVs (Figure 2C). The percentage of PLAP+ EVs was increased in the circulation of R2

compared to Controls suggesting the placenta secreted more large EVs into circulation.

Interestingly, we found a subset of endothelial-derived EVs that also express CD34, suggesting

that these EVs originated from fetal endothelial cells (48). Fetal endothelial cell-derived large

EVs (CD34+ CD31+) were also elevated in R1 and R2 compared to Controls. The percentage of

small EVs from the placenta was not altered by COVID-19 (Figure 2F). In fact, there was no

difference in the percentage of small EVs from any cell type measured. This suggests that

placenta-derived large, but not small EVs, were altered by COVID-19 in early pregnancy.

Circulating EVs from COVID-19 pregnant patients alter trophoblast function *in vitro*

The placenta is made up of three main functional cell types; 1) syncytiotrophoblast cells,

which are responsible for nutrient transport and hormone production; 2) cytotrophoblast cells,

the replicating precursors of the syncytiotrophoblast; and 3) extravillous trophoblasts (EVT),

which invade deep into the uterus to anchor the placenta and enable blood and nutrient flow to

the fetus. EV signaling is known to influence trophoblast function (49). Therefore, we tested the

capacity of circulating EVs isolated from participants with an active infection or Controls to alter

the function of trophoblast cell types. We focused our *in vitro* experiments on EVs isolated from

participants with an active infection compared to Controls because the changes in placental

pathology and transcriptome were the greatest in AI cases compared to Controls.

To assess the effect of AI EVs on EVT function, we used primary EVTs isolated from

first-trimester placenta and quantified EVT invasion through a collagen gel. EVT invasion,

which is vital for anchoring the placenta to the uterus and the remodeling of maternal uterine

arteries providing blood to the villous trophoblasts, was significantly reduced by exposure to AI

EVs compared to Control EVs (Figure 3A). Inadequate invasion and failure to completely

remodel maternal arteries increases the risk of preeclampsia, intrauterine growth restriction, and

fetal loss (*50, 51*).

To study syncytiotrophoblasts, we used the BeWo choriocarinoma cell line that is

commonly used to study this trophoblast lineage. After the addition of forskolin, BeWo cells

syncytialize forming cells that mimic the syncytiotrophoblast, including the production of

placental hormones (hCG and progesterone). Syncytiotrophoblast hormone production is

essential for the maintenance of pregnancy. The ratio of hCG to progesterone in the media of

syncytialized BeWo cells was significantly reduced following exposure to AI EVs compared to

Control EVs (Figure 3B). A reduction in the ratio of hCG to progesterone indicates that specific

pathways related to steroid hormone production were disrupted. These findings demonstrate that

EVs from the circulation of a pregnant individual with an active infection disrupt major

trophoblast functions including invasion and hormone production, which may have profound

effects on pregnancy maintenance.

To identify novel pathways that may contribute to trophoblast dysfunction, we analyzed

the BeWo transcriptome following EV exposure. AI EVs significantly altered gene expression

in BeWo cells compared to Control EVs. Multiple genes were dysregulated including genes that

encode for long non-coding RNA genes and histone proteins (Figure 3C). This suggests that

DNA packaging and transcription is disrupted in trophoblasts exposed to AI EVs compared to

Control EVs. Consistent with this, Biological Processes, determined by GO analysis, and top

canonical pathways, identified by IPA, were related to cellular transcription and DNA repair

(Figure 3D and Supplemental Table 4). The pathways disrupted by AI EVs suggest a

generalized effect on trophoblast gene expression that led to disrupted hormone production.

Increased mtDNA content in LEVs following COVID-19 in early pregnancy

Analysis of the placental transcriptome following COVID-19 identified differentially expressed genes indicative of mitochondrial dysfunction (Table 2). To determine if circulating EVs were enriched in mitochondrial cargo, we measured mitochondrial DNA (mtDNA) content (Figure 4A &C) and found that mtDNA was more abundant in large compared to small EVs. In contrast, nuclear DNA was not consistently measurable in all samples. Additionally, the abundance of mtDNA in large, but not small EVs, inversely correlated with the gestational timing of COVID-19 (Figure 4 B&D). The increase in mtDNA released in large EVs following COVID-19 during early pregnancy suggests that the mitochondrial function of cells producing large EVs was persistently disrupted.

COVID-19 during pregnancy alters EV RNA cargo

EVs contain small and larger (mRNAs) and long noncoding RNAs, however, small RNAs are the most commonly studied EV cargo (*52*). mRNAs encapsulated within EVs are transferred to recipient cells and translated into proteins, altering the behavior of the recipient cells (*53-56*). Therefore, we profiled the mRNA content of circulating EVs to determine if there were differences dependent on the gestational timing of COVID-19. We sequenced an average of 2,946 gene associated transcripts in large EVs and 1,947 in small EVs.

The most abundant mRNA transcripts in EVs were common to all groups. However, we also identified transcripts that were either uniquely expressed in COVID-19 groups (i.e. absent in Controls), or uniquely expressed in Controls, and absent in one or more COVID-19 groups. Multiple transcripts were uniquely expressed in large EVs from the COVID-19 groups including *YY1AP1*, *MOSPD1, RYBP*, and *H1-4* (Table 2). The proteins encoded by these mRNAs are related to transcription, except for *HI-4* which has an unknown cellular function. In small EVs,

MYL4, *C18orf32*, *CAPG*, and *CTSS* transcripts were uniquely present in EVs isolated from

COVID-19 groups (Table 2). These transcripts encode a motor (*MYL4* (myosin light chain 4))

and immune (*C18orf32* (Putative NF-Kappa-B-Activating Protein 200), *CAPG* (macrophage

capping protein), and *CTSS* (cathepsin S)) proteins. This suggests that COVID-19 alters

immune-related small EV cargo. Many other transcripts were uniquely detected in EVs isolated

from individual COVID-19 groups (Table 2). The unique transcriptome suggests that COVID-

19 increased expression of these genes making their transcripts more available for EV packaging

or increased specific transcript loading into EVs.

There were several interesting transcripts that were only present in large EVs from

Controls, including APBA3, MTSS, FCF1, PSG2, LOC100128233, PHOSPHO2, and THOC3

(Figure 5A). These transcripts encode proteins involved in various cellular functions, including

signal transduction, transcription, and proliferation. In contrast, only a few transcripts were

unique to Controls in small EVs. These included PYCR2, SCGB1C1, and CD300L4 (Figure

5B). PYCR2 encodes a cellular metabolism protein; CD300L4 encodes an immune-regulating

protein. The protein function of SCGB1C1 is unknown. Numerous other transcripts were

abundant in Controls but absent in individual COVID-19 groups. If present in the other COVID-

19 groups, their expression was decreased compared to Controls (Figure 5).

Transcripts carried by EVs reflect the activity of the secreting cell. While the individual

transcripts identified in EVs differ, their cellular functions often overlapped (Table 2, Figure 5).

For example, large and small EVs isolated from R2 carried transcripts that regulate cell

signaling, gene expression, immune regulation, and metabolism. Additional pathways include

proliferation, apoptosis, invasion, ubiquitination, platelet function, and vesicle formation.

Many transcripts identified in EVs are abundant in trophoblast cells and have been

previously reported to increase with gestation. Interestingly, *THOC3* mRNA, encoded by a

highly expressed gene in trophoblast cells, was abundant in Control large EVs but was low or

absent in the COVID-19 groups. In addition, large EVs carried a different highly expressed

trophoblast transcript, *RYBP*, in COVID-19 groups, but this transcript was absent in Controls.

Both proteins encoded by these genes are involved in transcriptional regulation (*57, 58*). Several

of the unique mRNAs in EVs isolated from COVID-19 groups have been previously reported to

be associated with adverse pregnancy outcomes, including gestational hypertension,

preeclampsia, preterm birth, and intrauterine growth restriction (Supplemental Table 5). The

number of these pregnancy complication-associated mRNAs was highest in AI EVs, but they

were also abundant in EVs isolated from participants with resolved infections. Interestingly, the

abundance of eleven transcripts that have been implicated in preeclampsia differed between AI

EVs and Control EVs. Of importance, participants with AI also had a higher incidence of

preeclampsia. Moreover, two transcripts associated with preterm birth were uniquely carried in

R2, but not Control EVs; R2 participants had a higher incidence of preterm birth. This suggests

that EV cargo may reflect etiological pathways leading to these pregnancy complications.

Circulating EVs carry transcriptional regulators of differentially expressed genes in the

placenta

Because EVs had a direct functional effect on trophoblasts *in vitro*, we investigated

whether they carried transcriptional regulators of genes whose expression was altered in placenta

of COVID-19 pregnant participants (Supplemental Table 3). Multiple mRNAs encoding

transcriptional regulators were differentially abundant in COVID-19 groups compared to

Controls. Several were contained in both small and large EVs (*JUN, FOS, LEPR, LGALS1,*

CD36) or only in small EVs *(PRKN)*. Moreover, Jun proto-oncogene B (*JUNB*) mRNA levels

were increased in large EVs from R1 compared to Controls (FC=2.25, p=0.01). This suggests

that *JUNB* carried in EVs could elicit the observed changes in transcription of its downstream

targets in the placenta. Similarly, mRNAs encoding 4 of the transcriptional regulators in R3

compared to Control placenta were found in large and small EVs (*IL1B, HIF1A, IGF2, PCBP1*);

1 was only in large EVs (*CXCR4*), and 3 were only in small EVs (*ESR1, AKT1, TP53, SPZ1,*

IRS2). Interestingly, interleukin 1 beta (*IL1B*) mRNA was decreased in large EVs from R3

compared to Controls (FC = -1.85, p=0.070), whereas estrogen receptor 1 (ESR1) mRNA levels

were lower in R3 small EVs compared to Controls (FC=-6.99, p=0.13). Importantly, expression

of genes controlled by these transcriptional regulators was altered in R3 placenta compared to

Controls. Similarly, many transcriptional regulators of genes with differential expression in AI

compared to Control placenta were found in both large and small EVs (*GRN* and *IL1B*), in large

EVs (*FAS*) or in small EVs (*JUN, STAT3, IFG1, IFNG, AGT*). These observations exemplify

potential EV-driven signaling leading to altered gene expression in the placenta that occurs in

COVID-19.

DISCUSSION

The long-term effects of COVID-19 during pregnancy have not yet been elucidated.

Previous studies and our findings reported here, show that the placenta is damaged, and the

likelihood of adverse pregnancy outcomes was increased in patients with a pregnancy

complicated by COVID-19. We have begun to elucidate the mechanisms underlying the

observed placental abnormalities associated with COVID-19. For the first time, we demonstrate

that circulating EVs from COVID-19 affected pregnancies 1) have a detrimental effect on

trophoblast function, including hormone production and invasion *in vitro*; 2) are altered after

SARS-CoV-2 infection; and 3) carry cargo that has been previously associated with adverse

pregnancy outcomes.

Our findings, in *in vitro* experiments, that trophoblast dysfunction following exposure to

EVs isolated from study participants with an active SARS-CoV-2 infection provides evidence

that circulating EVs contribute to the resulting placental pathology. We focused our *in vitro*

trophoblast experiments on the response to EVs isolated from participants with an active

infection because the magnitude of alteration in the placental transcriptome was greater

compared to those placentas from resolved infections. EVs from patients with an active

infection (AI) disrupted fundamental trophoblast functions that are crucial to maintain a healthy

pregnancy. Others have shown that trophoblast dysfunction, including failure to invade and

produce hormones, contributes to the development of preeclampsia, preterm birth, and

intrauterine growth restriction (*50, 51*). Therefore, the AI EV-induced reduction in EVT

invasion and syncytiotrophoblast hormone production may have contributed to the development

of these pregnancy complications following COVID-19.

Gestational age at the time of infection was a major determinant of COVID-19-induced

changes in the profile of circulating EVs. If participants were infected during the first or second

trimester of pregnancy, numbers of trophoblast and fetal endothelial cell large EV were

increased, and circulating large EVs carried more mtDNA. This suggests that COVID-19 during

early pregnancy disrupts mitochondrial function in the placenta. Mitochondrial dysfunction has

been reported in many organs following SARS-CoV-2 infection and is thought to contribute to

cell injury, cell death, and inflammation (*59-61*). Appelman et al. recently reported persistent

mitochondrial dysfunction in skeletal muscle long after the resolution of SARS-CoV-2 infection

(*62*). Further, elevated cell-free circulating mtDNA is commonly observed in COVID-19 and

363 correlates with severity and length of infection, reflecting significant mitochondria stress (*63-*

364 *66*). In support of a direct association of infection and mtDNA release in EVs, Faizan et al.

365 recently demonstrated that SARS-CoV-2 infection causes mitochondrial dysfunction and release

366 of EVs containing mtDNA in airway epithelial cells (*60*). Thus, our results suggest that

367 abnormal mitochondria may also play a role in the pathogenesis of placental dysfunction in

368 COVID-19.

369  Circulating EV cargo reflects the activity of the cells of origin. The transcripts carried by

370 EVs encode genes related to inflammation, vasculopathies, bioenergetics, and cell death,

371 processes and pathways that were present in the transcriptome and histopathology of the placenta

372 regardless of the timing of infection. EVs carry transcripts that are highly expressed by

373 trophoblasts, and have known functions in cellular metabolism, immune regulation, and

374 transcription. We also found that many of the transcripts in EVs from pregnancies complicated

375 by COVID-19 are encoded by genes that have been implicated in adverse pregnancy outcomes

376 including gestational hypertension, preeclampsia, preterm birth, and intrauterine growth

377 restriction. This points to shared pathways of placental dysfunction induced by a systemic

378 SARS-CoV-2 infection.

379  EV cargo can elicit a functional response when delivered to a recipient cell, as

380 demonstrated by our *in vitro* studies. While it is not known if mtDNA in EVs per se was

381 responsible for altering trophoblast function in our experiments, multiple studies have

382 demonstrated that mitochondria cargo can alter the recipient cell's mitochondrial function (*67-*

383 *69*). mRNA transcripts are also biologically active in recipient cells and we identified transcripts

384 in EVs that encode for multiple transcriptional regulators genes whose expression was altered in

385 placenta following COVID-19. Importantly, expression of several of these genes has been

previously reported to be altered in pregnancy complications. For example, JUN signaling was

disrupted in R1 placenta compared to Control placenta and JUNB mRNA was increased in

Control compared to R1 large EVs. JUN proteins are important for placentation, and a loss of

JUN signaling is implicated in preeclampsia (*70, 71*). Thus, low levels of JUNB in COVID-19

EVs may indicate placental dysfunction which in turn could contribute to the later development

of preeclampsia, which is observed at higher rates in pregnancies complicated by COVID-19

(*72*). In R3 compared to Controls, hormone receptor signaling was identified as a top canonical

pathway and differentially expressed genes were regulated by ESR1. *ESR1* mRNA was

abundant in small EVs isolated from Controls but not R3. ESR1 signaling is vital for placental

function and pregnancy maintenance because estrogen signaling is obligate for angiogenesis and

vasculature control (*73*). In fact, genetic variations in *ESR1* are associated with recurrent

pregnancy loss and preeclampsia, and both adverse pregnancy outcomes are increased in

maternal SARS-CoV-2 infection during pregnancy (*74, 75*). Thus, our findings suggest

placental dysfunction may in fact be a result of EV cargo delivery.

Our study is limited by the number of symptomatic patients with an active infection at the

time of delivery. Despite only 14% of pregnant participants experiencing COVID-19 related

symptoms, their placentas had significant pathology and an altered transcriptome. This was

associated with an increased incidence of preeclampsia and medically indicated preterm birth in

asymptomatic and symptomatic AI cases. It's worth noting that EVs obtained from

asymptomatic individuals have been found to exert significant impacts on trophoblast function

when tested in vitro. This discovery highlights the importance of exploring the potential

consequences of EV exposure in asymptomatic patients and may have important implications for

understanding the role of EVs in reproductive health.

Our study has provided significant insights into the profile and functional consequences

of circulating extracellular vesicles in mothers who were infected with SARS-CoV-2. This study

is the first to demonstrate the negative impact of maternal circulating vesicles on trophoblast

function in COVID-19. By comparing the placental transcriptome and EV cargo content, we

have identified shared pathways that are associated with pregnancy complications caused by

maternal COVID-19 and other pregnancy-related disorders that are not well understood.

MATERIALS AND METHODS

Patient cohort

The COMET study was conducted at the Hospital of the University of Pennsylvania (HUP) with

Institutional Review Board approval (IRB#843277). Study participants received a description of

the study and signed an informed consent before enrollment. Participants were enrolled at the

time of delivery in the COMET study between April 2020-June 2022. Participants were tested

for a SARS-CoV-2 infection by nasopharyngeal polymerase chain reaction (PCR) upon

admission to the labor and delivery unit at HUP. Participants who tested positive at the time of

delivery were enrolled in the active infection (AI) group. Those participants who tested negative

and had no known SARS-CoV-2 infection during their pregnancy were defined as Controls.

Participants with a negative test at delivery and a history of SARS-CoV-2 infection during their

pregnancy and greater than 14 days before enrollment, were defined as having a resolved

infection (R). All COVID-19 cases were unvaccinated against SARS-CoV-2. The gestational

age of SARS-CoV-2 infection was calculated, and participants were further divided into the

trimester of infection (resolved infection in the first trimester (R1), resolved infection in the

second trimester (R2) and resolved infection in the third trimester (R3)).

Clinical and demographic data collection

Clinical characteristics, such as maternal age, self-identified race, gestational age at infection, and pregnancy outcomes, were extracted from the medical record (Table 1). The severity of COVID-19 disease was categorized based on the National Institute of Health and Society for Maternal-Fetal Medicine definitions: Asymptomatic infection was defined as participants who tested positive but experienced no symptoms. Symptomatic participants included all levels of illness (mild-critical).

Sample collection

Placentas were collected at the time of delivery. All placentas were examined by the pathology department at the Hospital of the University of Pennsylvania (HUP). Placentas were assessed using a systematic protocol that includes recording the trimmed placental weight, membrane insertion site, gross appearance, dimensions of the placental disc, and umbilical cord insertion, length, and diameter. Full-thickness placental biopsies were collected from an area devoid of obvious pathology located equidistant between the placental cord insertion and the edge of the placenta. Tissue was fixed in 10% formalin for histological assessment. Macroscopic and microscopic lesions were identified and classified according to the Amsterdam Placental Workshop Group 2014 classification (*76-78*). Placental biopsies were also collected and stored in Trizol for RNA isolation.

Blood was collected at delivery in an EDTA tube and spun at 1,000G for 10 minutes at room temperature to isolate plasma, which was aliquoted and stored at -80°C.

Placenta and BeWo RNA isolation and sequencing

Total RNA was isolated from placental biopsy samples using Qiagen RNEasy Plus Mini

Kits (Cat# 74134 Qiagen, Hilden, Germany). Total RNA was isolated from syncytialized BeWo

cells with the PicoPure RNA Isolation Kit (Cat# KIT0204 Applied Biosystems, Waltham, MA).

Isolated RNA was sent to NovoGene for library preparation and sequencing.

RNA integrity and quantification were assessed using the RNA Nano 6000 Assay Kit of

the Bioanalyzer 2100 System (Agilent Technologies, CA, USA). RNA purity was determined

using a NanoPhotometer spectrophotometer (IMPLEN, CA, USA). A total of 1µg RNA per

sample was used as input material for the RNA sample preparation. Sequencing libraries were

generated using NEBNext Ultra RNA Library Prep Kit for Illumina (NEB, USA) following

manufacturer recommendations, and index codes were added to identify samples. Clustering of

the index-coded samples was performed on an Illumina Novaseq 6000 sequencer according to

the manufacturer's instructions. After cluster generation, libraries were sequenced, and pair-end

reads were generated. Raw data (raw reads) of FASTQ format were processed through fastp. and

clean data was obtained by removing reads containing adapter and poly-N sequences and reads

with low quality. Pair-end clean reads were aligned to the GRCh38/hg38 reference genome

using Spliced Transcripts Alignment to a Reference (STAR) software. FeatureCounts were used

to count the read number mapped to each gene. Then RPKM of each gene was calculated based

on the length of the gene and read count mapped to the gene. Differential gene expression

between COVID-19 groups and Controls was assessed by DESeq2. Differentially expressed

genes were determined based on their adjusted p-value (<0.05) and >1.5-fold change. Functional

analysis was conducted using Qiagen's ingenuity pathway analysis (IPA). The clusterProfiler R

package was used to perform a Gene Ontology enrichment analysis of genes that were

differentially expressed. Canonical pathways, transcriptional regulators, and GO terms were

considered significant if the p-value was less than 0.05.

EV isolation and characterization

Serial centrifugation was utilized to isolate EVs from plasma. One mL of plasma was

spun in an Eppendorf 5424 benchtop centrifuge at 2,000 x g for 10 minutes at 4°C. The

supernatant was then spun at 20,000 x g for 30 minutes at 4°C. The pellet was washed in 1mL

filtered PBS and spun again at 20,000 x g for 30 minutes at 4°C. The large EV pellet was

resuspended in 100uL filtered PBS. The supernatant was spun by the Beckman Ultracentrifuge

Optima Max TL using the TLA 120.2 rotor at 100,000 x g (48,000RPM) for 90 minutes at 4°C.

The pellet was washed with 1mL filtered PBS and spun again at 100,000 x g (48,000RPM) for

90 minutes at 4°C. The small EV pellet was resuspended in 100uL filtered PBS.

EV isolation was confirmed by transmission electron microscopy, nanoparticle tracking,

and protein measurement as recommended by the MISEV guidelines (79). Transmission

electron microscopy images were generated and resulting images reviewed for the presence of

EVs. EVs were analyzed by Particle Metrix Zetaview nanoparticle tracking. 11 fields were

captured using the following parameters: sensitivity 80, frame rate 30, shutter 100, minimum

brightness 1000, minimum area 10, trace length 15. Representative histograms and TEM images

for large and small EVs are included in Supplemental Figure 1. CD9 protein abundance was

determined by gel electrophoresis. Total protein was measured with a Qubit Protein Assay Kit

and EV suspension was evaporated by vacuum and resuspended in electrophoresis buffer. Three

large EV (5μg) and small EV (20μg) samples were loaded into BioRad Mini-Protean TGX Gel

4-20% polyacrylamide gels with Licor Chameleon Duo ladders (928-60000) and run at 20mA

for 2 hours. Proteins were transferred to nitrocellulose membrane via 200mA over 3 hours on

ice. The membrane was blocked in Licor Intercept Blocking Buffer for 1hour at room

temperature then incubated with CD9 antibody (HI9a Biolegend Cat 312112) at 1:5000

overnight at room temperature. The membrane was then incubated with Licor IRDye 800CW

streptavidin (926-32230) at 1:5000 for 2 hours at room temperature and the membrane was

imaged by Licor Odyssey.

Flow cytometry on EVs

EVs surface protein expression was determined by flow cytometry following the MISEV

guidelines (*79*). EVs were resuspended at $1 \times 10^{8}/10\mu L$ of filtered PBS. Di-8-ANEPPS

(Invitrogen Cat# D3167) was reconstituted in ethanol as per manufacturer instruction and further

diluted to 1:1000 in filtered PBS. Antibodies were spun at 20,000 x g for 30 minutes at 4°C

immediately before use. 10μL of EV suspension was incubated in ANEPPS (10μL) and

antibodies, 1.25μL CD45-Ry586 (Cat# BD568135), 1.25μL CD41a- PE/Cy7 (Cat#

BDB561424), 1.25μL PLAP- eFlour660 (Fisher Cat# 50-112-4573) , and 1.25μL CD34-

PE/CF594 (Cat# BDB562449) and 3μL CD31-AF700 (Biolegend Cat# 50-207-2950), for 30

minutes at room temperature. 470μL of filtered PBS was added before samples were measured

by BD Symphony A1 cytometer which has improved sensitivity for small particles. Negative

controls included: antibodies alone, EVs without Di-8-ANEPPS, and EVs treated with 1% triton.

Data was analyzed using FlowJo software. EVs were identified by Small Particle Side Scatter

(SP-SSC) and expression of Di-8-ANEPPS and the relative proportion that expresses cell-

specific surface proteins was determined by antibody detection.

***In vitro* trophoblast EV co-culture**

Extravillous trophoblasts (EVTs) were isolated from fresh first trimester placenta based

on an EVT outgrowth-based protocol established by Gram et al. (*80-84*). In brief, villous tissue

was finely minced and cultured at 37°C and 5% CO_2 in RPMI 1640 media with 20% FBS. After

attachment, EVT outgrowth occurs, and those cells were isolated. Isolated EVTs were confirmed

by staining for HLA-G and CK7. EVT invasion was measured using the MilliporeSigma

Chemicon QCM Collagen Cell Invasion Assay (Cat# ECM558). EVTs were added to the trans-

well invasion plate with EV-depleted media and large and small EVs at $1x10^6$/mL. Cells were

incubated at 37°C and 5% CO_2 for 48 hours. Cells that invaded through the collagen membrane

were quantified using a fluorescent plate reader (SpectraMax).

BeWo cells, subclone B30, were cultured in 75-cm^2 flasks (Fisher Scientific) at 37°C and

5% CO_2 in media (DMEM/F12, 10% FBS, 1% P/S, 1%L-alanyl-L-glutamine). EV-depleted

media was made with EV-depleted FBS (Gibco A2720801) and used for cell culture

experiments. Cells were plated at 250,000 cells/well in a 6 well plate and 1.5mL of EV-depleted

media was added. Cells adhered for 24 hours before adding 1μg/μL forskolin, a cAMP

producer, to promote syncytialization for an additional 24 hours. Large and small EVs were

resuspended in EV-depleted media at $1x10^6$/mL and added to BeWo cells for an additional 24

hours. At the time of harvest, cell media was collected, and cells released with 0.25% trypsin.

Cells were washed and collected as pellets for total DNA measurement and RNA isolation and

sequencing. Cell media was spun at 500 x g to clear cell debris and the supernatant was stored

for future hormone measurement. Hormones were measured by Penn Fertility Care using

Elecsys HCG+β (Cat# 03271749, Roche Diagnostics) and Elecsys Progesterone III (Cat#

07092539, Roche Diagnostics).

EV mtDNA measurement

mtDNA was isolated and quantified from large and small EVs by TaqMan-based

quantitative polymerase chain reaction (qPCR). We determined that $2x10^7$ large EV and $3x10^9$

small EVs were necessary to reliably and robustly measure mtDNA. As previously described,

we quantified mitochondrial-encoded human NADH: ubiquinone oxidoreductase core subunit 1

(ND1) as previously described (*85*). The qPCR reactions were performed in triplicates using a

QuantStudio 5 Real-time PCR System (Thermo Fisher) using the following thermocycling

conditions: 95 °C for 20 s followed by 40 cycles of 95 °C for 1 s, 63 °C for 20 s, and 60 °C for

20 s. Serial dilutions of pooled human placenta DNA quantified for copies of ND1 (copies/ μL)

by digital PCR (dPCR) were used as a standard curve. The mtDNA amount per EV was

determined by normalizing the resulting abundance by the number of EVs in starting material.

We calculated the Pearson correlation coefficient to determine the strength of the relationship

between gestational age at infection and abundance of EV mtDNA.

EV mRNA sequencing

Total RNA was isolated from large and small EVs isolated from 500μL of plasma.

Isolated EVs were treated with RNAseA (0.02μg/μL) (Invitrogen Cat# 12091021) for 20

minutes at 37°C to degrade extravesicular RNA. Enzyme activity was stopped by freezing

samples at -80°C for 5 minutes and immediate resuspension in Trizol. Nucleic acids were

isolated via BCP co-incubation, precipitated by isopropanol, and washed in ethanol. mRNA

libraries were prepared from total RNA using the SMART-Seq protocol (*86*). Briefly, RNA was

reverse transcribed using Superscript II (Invitrogen, Cat#18064014). The cDNA was amplified

with 20 cycles and cleaned up with AMPure XP beads (Beckman Coulter Cat#A63881). cDNA

was quantified with Qubit dsDNA HS Assay Kit (Life Technologies, Inc. Cat#Q32851), and 2ng

of each sample was used to construct a pool of uniquely indexed samples (Illumina Cat# FC-

131-1096). A second amplification was performed with 12 cycles and cleaned up with AMPure

XP beads. The final library was sequenced on a NextSeq 1000. Data were mapped against the

hg19 genome using RSEM and normalized to transcripts per million (tpm)(*87*).

To determine unique expression, we filtered genes to those that had greater than 5 tpm.

Unique genes had no expression (≤5 tpm) in all samples in the reference group and had

expression (>5 tpm) in the majority (≥ 50% of the samples) in the comparison group (data in

Table 2). Expression of unique genes in Controls, but not COVID-19 groups are shown in

Figure 4. These strict criteria identified genes in each group that were uniquely expressed, and

those genes were considered for subsequent analysis.

Statistical analysis

Statistical analysis was performed using GraphPad Prism. Differences in participant

demographics and outcomes were tested by a chi-squared test and considered statistically

different if $p<0.05$. Data was tested for normality and either parametric or non-parametric tests

were used to determine significance. Data points were identified as outliers and removed if they

exceeded two times the standard deviation from the mean. A one-way ANOVA tested for a

difference within all groups and subsequent post-hoc t-tests or Kruskal-Wallis determined the

significance of each COVID-19 group compared to Controls. Pearson's correlation was used to

determine a correlation between gestational age at infection and mtDNA abundance. A p-value

less than 0.05 was considered significant. Differential gene expression was determined to be

significant if the adjusted p-value was less than 0.05 and the fold change greater than 1.5.

List of Supplementary Materials

Supplemental Figure 1. EV Characterization

Supplemental Table 1. Placenta RNAseq DEG

Supplemental Table 2. Placenta RNAseq canonical pathways

Supplemental Table 3. Placental RNAseq transcriptional regulators

Supplemental Table 4. BeWo RNAseq pathway analysis

Supplemental Table 5. EV Transcripts Associated with Adverse Pregnancy Outcomes

References (*97-138*)

References and Notes

Acknowledgments: The authors would like to acknowledge the dedication and effort by the

Pregnancy and Perinatal Research Center, especially Meaghan McCabe, MPH, in

enrolling patients and collecting samples. The authors would also like to thank Dr. Luca

Musante at the University of Pennsylvania's Extracellular Vesicle Core for guidance in

EV-related methods. Finally, the authors would like to thank Dr. Jonni Moore and

Richard Schretzenmair at the University of Pennsylvania's Cytomics and Shared

Resource Laboratory for their guidance in conducting flow cytometry on EVs.

Funding:

Analyses of placentas collected at delivery (part of the COMET study) were supported (in part)

by a COVID-19 grant from the March of Dimes (SP, RAS)

National Institute of Health grant T32ES019851 (TNG)

National Institute of Health grant P30ES013508 (TNG)

Author contributions:

Experimental Design: TNG, SM, RLL, RL, LA, MM, CCC, BAK, JFS, SP, RAS

Data acquisition and analysis: TNG, SM, RLL, AW, LA, CCC, BAK

Writing- original draft: TNG, JFS, SP, RAS

Writing- review & editing: TNG, SM, RLL, RL, NAT, AW, LA, MM, CCC, BAK, JFS, SP,

RAS

Competing interests: The authors declare they have no competing interests.

Figure 1. Differential gene expression in the placentas from patients with COVID-19 during

pregnancy are represented by volcano plots. **(A-D)** The number and direction of differentially

expressed genes between Controls (n=5) and resolved infections in the first trimester (R1) (A),

second trimester (R2) (B), and third trimester (R3) (C), and active infection (AI) (D) is listed at

the top of each graph. The gene name for those transcripts with differential expression in more

than one COVID-19 group compared to Control is listed. (n=3-5/group)

Figure 2. Circulating EVs were persistently altered in participants who experienced COVID-19

in the second trimester. **(A)** The number of LEVs in the circulation at the time of delivery

(n=14-22/group). **(B)** The diameter of LEVs in the circulation (n=14-22/group). **(C)** Relative

frequency of LEVs derived from trophoblasts, endothelial cells, platelets, and immune cells

(n=11-16). **(D)** The number of small EVs in the circulation at the time of delivery (n=14-

22/group). **(E)** The diameter of small EVs in the circulation (n=14-22/group). **(F)** The relative

frequency of small EVs derived from endothelial cells, platelets, immune cells, and trophoblasts

(n=13-20/group). All data are presented as mean ± SD. All analyses were performed by one-

way ANOVA or the Kruskal-Wallis test, followed by post-hoc tests. Comparisons were made

between Controls and resolved infection in the first trimester (R1), second trimester (R2), third

trimester (R3), and active infection (AI).

Figure 3. Trophoblast function was disrupted by exposure to EVs isolated from patients with an

active infection (AI) compared to Controls. **(A)** Extravillous trophoblasts (EVTs) were isolated

from three placentas (identified by symbol shape) and exposed to Control or AI EVs (n=9-

10/group, 3 experiments). Invasion was calculated and normalized to invasion of EVTs derived

from the same placenta not exposed to EVs (noEVs). All data are presented as mean ± SD. **(B)**

Human chorionic gonadotropin (hCG) and progesterone were measured in the media of

forskolin-treated (syncytialized) BeWo cells. The ratio of hCG to progesterone was normalized

to hormone production by cells not exposed to EVs (noEVs). The results of two experiments

(identified by symbol shape) are reported in B. (n=6-7/group, 2 experiments). All analyses were

648 performed by one-way ANOVA or Kruskal-Wallis test, followed by post-hoc tests. **(C)**

649 Following EV exposure, BeWo cell transcriptome was measured, and the top differentially

650 expressed genes comparing the cellular response to AI EVs to Control EVs are listed in the heat

651 map (red represents increased and blue represents decreased expression). **(D)** The biological

652 processes altered by AI EVs compared to Control EVs were determined by Gene Ontology

653 enrichment analysis.

Figure 4. Large EV abundance of mtDNA is inversely correlated with gestational timing of

infection. **(A&C)** The amount of mtDNA in each large EV (A) and small EV (C) is reported for

each group (n=13-20/group). Data are presented as mean ± SD and tested by ANOVA followed

by post-hoc tests. Independent pairwise comparisons were made between Controls and resolved

infection in the 1st trimester (R1), second trimester (R2), third trimester (R3), or active infection

(AI). **(B&D)** The Pearson correlation between gestational age at infection and mtDNA content is

reported for large EV (B) and small EV (D).

Figure 5. Control EVs carried transcripts that were absent in COVID-19 groups. **(A&B)** mRNA

transcripts uniquely detected in EVs isolated from Controls but absent in EVs isolated from

COVID-19 groups (gray bars) and differential expression (log 2-fold change) in other COVID-

19 groups are listed in the heat map (increased expression is red and decreased expression is

blue). The general cellular function of each gene product is listed on the right. Large EV

transcripts are reported in (A) and small EV transcripts are reported in (B) (n=9-10/group).

Independent pairwise comparisons were made between Controls and resolved infection in the

first trimester (R1), second trimester (R2), third trimester (R3), or active infection (AI).

Transcripts known to be highly abundant in trophoblasts are marked by (T).

Demographics	Controls (n=32)	R1 (n=15)	R2 (n=20)	R3 (n=22)	AI (n=21)
Maternal age (years)	21 – 42 (Mean: 31.8)	25 - 43 (Mean: 31.7)	18 – 39 (Mean: 29.4)	21 – 42 (Mean: 30.5)	20 – 40 (Mean: 29.4)
Race: White	50%	46.7%	34%	27.3%	9.5%*
Race: Black	41%	46.7%	55%	68.2%*	76.2%*
Race: Asian	9.4%	0%	10%	0%	9.5%
Race: Other/Unknown	0%	6.7%	0%	4.5%	4.8%
GA at delivery (weeks)	37.3 – 41.1 (Mean: 39.3)	35.1 – 40.3 (Mean: 39.1)	33.7 – 41.3 (Mean: 38.4)	32.3 – 39.7 (Mean: 37.5) *	31 – 41 (Mean: 38.1) *
COVID severity	Controls (n=32)	R1 (n=15)	R2 (n=20)	R3 (n=22)	AI (n=21)
Symptomatic	N/A	73.3%	85%	81.8%	14.3%
Asymptomatic	N/A	26.7%	15%	18.2%	85.7%[+]
Hospitalized	N/A	13.3%	0%	31.8%	4.8%
Pregnancy outcomes	Controls (n=32)	R1 (n=15)	R2 (n=20)	R3 (n=22)	AI (n=21)
GHTN	18.8%	53.3%*[+]	20%	13.6%	9.5%
CHTN	3.1%	0%	5%	13.6%	4.8%
Preeclampsia	0%	0%	5%	18.2%*	19%*
Spontaneous PTB	0%	0%	20%*	4.5%	4.8%
Medically indicated PTB	0%	6.7%	10%	18.2%*	19%*
IUGR	6.3%	13.3%	0%	9.1%	4.8%
IUFD	0%	0%	0%	4.5%	0%
Placenta Pathology	Control (n=26)	R1 (n=9)	R2 (n=17)	R3 (n=19)	AI (n=19)
MVM or FVM	26.9%	100%*	71%*	63%*	82.6%*
MVM	15%	44%*	47%*	37%	53%*
High grade MVM	0%	11%	18%*	16%*	11%
FVM	15%	78%*	59%*	42%*	47%*
>10% perivillous fibrin deposition	0%	11%	18%*	21%*	5%

Table 1. Subject demographics, pregnancy outcomes, and placenta pathology

Participants enrolled in the COMET study formed five groups (controls, resolved infection in the 1st

trimester (R1), 2nd trimester (R2), and 3rd trimester (R3), and active infection (AI). Maternal

demographics including maternal age, race, and gestational age (GA) at birth are reported. The severity

of COVID-19 during their pregnancy, incidence of pregnancy complications (gestational hypertension

(gHTN), chronic hypertension (cHTN), preeclampsia (PE), spontaneous and medically indicated preterm

birth (PTB), intrauterine growth restriction (IUGR), and intrauterine demise (IUFD)) and placental

pathology (maternal vascular malperfusion (MVM), fetal vascular malperfusion (FVM) and perivillous

fibrin deposition) are reported * p<0.05 chi-squared test compared to Controls. [+] p<0.05 compared to

other COVID-19 groups

	Uniquely Expressed Transcripts in COVID EVs				
Gene Symbol	**Gene Name**	**Group(s) with Unique Expression**	**Cellular Function**	**High Trophoblast Expression**	**Increased in Pregnancy**
		Large EVs			
YY1AP1	YY1 Associated Protein 1	R1, R2	Transcription		
MOSPD1	Motile Sperm Domain Containing 1	R1, AI	Transcription		
RYBP	RING1 And YY1 Binding Protein	R1, AI	Transcription	Y	
H1-4	H1.4 linker histone, cluster member	R2, R3	Unknown		
A2M	Alpha-2-Macroglobulin	R1	Immune		
KIFAP3	Kinesin Associated Protein 3	R1	Chromosome Movement		Y(*88*)
MXD1	MAX Dimerization Protein 1	R1	Proliferation		
PEX19	Peroxisomal Biogenesis Factor 19	R1	Oxidative stress	Y	
TOB1	Transducer Of ERBB2, 1	R1	Proliferation		
HK1	Hexokinase 1	R2	Metabolism	Y	
LY9	Lymphocyte Antigen 9	R2	Immune		Y(*89*)
POLR3C	RNA Polymerase III Subunit C	R2	Nucleic acid binding activity		
SPNS3	SPNS Lysolipid Transporter 3, Sphingosine-1-Phosphate (Putative)	R2	Transporter Activity		
WDR46	WD Repeat Domain 46	R2	Nucleic acid binding activity		
CDC34	Cell Division Cycle 34, Ubiquitin Conjugating Enzyme	R3	Ubiquitination		
FUNDC2	FUN14 Domain Containing 2	R3	Metabolism		
LINC-PINT	Long Intergenic Non-Protein Coding RNA, P53 Induced Transcript	R3	Unknown		
R3HCC1	R3H Domain and Coiled-Coil Containing 1	R3	Nucleic acid binding activity		
ATPSCKMT	ATP Synthase C Subunit Lysine N-Methyltransferase	AI	Mitochondrial ATP synthesis		
DTWD1	DTW Domain Containing 1	AI	Translation		
FBXL4	F-Box and Leucine Rich Repeat Protein 4	AI	Ubiquitination		
FRA10AC1	FRA10A Associated CGG Repeat 1	AI	Transcription		
FOXP1	Forkhead Box P1	AI	Transcription		
GOLGA4	Golgi A4	AI	Protein and lipid transport	Y	Y(*88*)

GPBP1L1	GC-Rich Promoter Binding Protein 1 Like 1	AI	Transcription		
IL1R2	Interleukin 1 Receptor Type 2	AI	Immune		
KIAA1143	KIAA1143	AI	Unknown		
LINC01410	Long Intergenic Non-Protein Coding RNA 1410	AI	Unknown		
NEAT1	Nuclear Paraspeckle Assembly Transcript 1	AI	Transcription	Y	
WDR26	WD Repeat Domain 26	AI	Cell cycle progression and gene regulation	Y	
XRN1	5'-3' Exoribonuclease 1	AI	mRNA degradation		Y(*90*)
ZNF638	Zinc Finger Protein 638	AI	Transcription		
Small EVs					
MYL4	Myosin Light Chain 4	R1, AI	Motor Protein		Y(*91*)
C18orf32	Chromosome 18 Open Reading Frame 32	R1, AI	Immune	Y	
CAPG	Capping Actin Protein, Gelsolin Like	R2, AI	Motor Protein		
CTSS	Cathepsin S	R3, AI	Immune		Y(*92*)
CYREN	Cell Cycle Regulator Of NHEJ	R1	DNA Repair		
CCDC124	Coiled-Coil Domain Containing 124	R1	Transcription		
CD27-AS1	CD27 Antisense RNA 1	R1	Immune		
GNB2	G Protein Subunit Beta 2	R1	G Protein Signaling		
PHF5A	PHD Finger Protein 5A	R1	Immune	Y	
RBM8A	RNA Binding Motif Protein 8A	R1	Transcription	Y	
SIGMAR1	Sigma Non-Opioid Intracellular Receptor 1	R1	Calcium Signaling	Y	
CSF3R	Colony Stimulating Factor 3 Receptor	R2	Immune	Y	
ESD	Esterase D	R2	Metabolism		
GTF2IRD2	GTF2I Repeat Domain Containing 2	R2	Transcription		
SLC1A5	Solute Carrier Family 1 Member 5	R2	Metabolism	Y	Y(*93*)
AUP1	AUP1 Lipid Droplet Regulating VLDL Assembly Factor	R3	Ubiquitination	Y	
MFF	Mitochondrial Fission Factor	R3	Mitochondrial and Peroxisomal Fission		
ARCN1	Archain 1	AI	Vesicle	Y	
CKAP2	Cytoskeleton Associated Protein 2	AI	Proliferation		Y(*94*)

CSNK2B	Casein Kinase 2 Beta	AI	Metabolism		
CYP27C1	Cytochrome P450 Family 27 Subfamily C Member 1	AI	Metabolism		
GRHL1	Grainyhead Like Transcription Factor 1	AI	Epithelial Development	Y	
HSP90B1	Heat Shock Protein 90 Beta Family Member 1	AI	Molecular Chaperone	Y	Y(*95*)
KIF5B	Kinesin Family Member 5B	AI	Protein Binding Activity		
LINC01123	Long Intergenic Non-Protein Coding RNA 1123	AI	Unknown		
LOC728323	Unknown	AI	Unknown		
MANBAL	Mannosidase Beta Like	AI	Membrane Protein		Y(*96*)
MUC22	Mucin 22	AI	Membrane Protein		
MYLK	Myosin Light Chain Kinase	AI	Contractile Activity		
OR4F17	Olfactory Receptor Family 4 Subfamily F Member 17	AI	Vesicle		
PARP9	Poly (ADP-Ribose) Polymerase Family Member 9	AI	Immune		
RNF2	Ring Finger Protein 2	AI	Transcription		
SAR1A	Secretion Associated Ras Related GTPase 1A	AI	Vesicle	Y	
SHC4	SHC Adaptor Protein 4	AI	Proliferation		
SRSF8	Serine And Arginine Rich Splicing Factor 8	AI	Transcription	Y	
TMCC2	Transmembrane And Coiled-Coil Domain Family 2	AI	Metabolism	Y	
TRIM4	Tripartite Motif Containing 4	AI	Immune		
TWF2	Twinfilin Actin Binding Protein 2	AI	Actin and ATP Binding Site		
VTA1	Vesicle Trafficking 1	AI	Vesicle		
ZNF484	Zinc Finger Protein 484	AI	Transcription		

686   Table 2. Unique genes identified in EVs isolated from COVID-19 cases

687   Transcripts are carried by large EVs or small EVs isolated from COVID-19 cases that are absent in

688   Controls. The listed transcripts are not detected in EVs isolated from Controls but are present in the

689   identified COVID-19 group(s). The cellular function, expression in trophoblasts, and pregnancy

690   associated expression of each transcript is listed as well (reference listed).

References:

1. A. Conde-Agudelo, R. Romero, SARS-CoV-2 infection during pregnancy and risk of preeclampsia: a systematic review and meta-analysis. *Am J Obstet Gynecol* **226**, 68-89 e63 (2022).

2. E. R. Smith *et al.*, Adverse maternal, fetal, and newborn outcomes among pregnant women with SARS-CoV-2 infection: an individual participant data meta-analysis. *BMJ Glob Health* **8**, (2023).

3. C. M. Corbetta-Rastelli *et al.*, Analysis of placental pathology after COVID-19 by timing and severity of infection. *Am J Obstet Gynecol MFM* **5**, 100981 (2023).

4. F. M. Cribiu *et al.*, Severe SARS-CoV-2 placenta infection can impact neonatal outcome in the absence of vertical transmission. *J Clin Invest* **131**, (2021).

5. B. Joshi *et al.*, The placental pathology in Coronavirus disease 2019 infected mothers and its impact on pregnancy outcome. *Placenta* **127**, 1-7 (2022).

6. E. T. Patberg *et al.*, Coronavirus disease 2019 infection and placental histopathology in women delivering at term. *Am J Obstet Gynecol* **224**, 382 e381-382 e318 (2021).

7. A. G. Edlow *et al.*, Assessment of Maternal and Neonatal SARS-CoV-2 Viral Load, Transplacental Antibody Transfer, and Placental Pathology in Pregnancies During the COVID-19 Pandemic. *JAMA Netw Open* **3**, e2030455 (2020).

8. J. J. Mulvey, C. M. Magro, L. X. Ma, G. J. Nuovo, R. N. Baergen, Analysis of complement deposition and viral RNA in placentas of COVID-19 patients. *Ann Diagn Pathol* **46**, 151530 (2020).

9. C. Balbi *et al.*, Circulating extracellular vesicles are endowed with enhanced procoagulant activity in SARS-CoV-2 infection. *EBioMedicine* **67**, 103369 (2021).

10. D. Forte *et al.*, Circulating extracellular particles from severe COVID-19 patients show altered profiling and innate lymphoid cell-modulating ability. *Front Immunol* **14**, 1085610 (2023).

11. M. S. George *et al.*, Extracellular vesicles in COVID-19 convalescence can regulate T cell metabolism and function. *iScience* **26**, 107280 (2023).

12. A. Rosell *et al.*, Patients With COVID-19 Have Elevated Levels of Circulating Extracellular Vesicle Tissue Factor Activity That Is Associated With Severity and Mortality-Brief Report. *Arterioscler Thromb Vasc Biol* **41**, 878-882 (2021).

13.   L. Traby *et al.*, Extracellular Vesicles and Citrullinated Histone H3 in Coronavirus Disease 2019 Patients.

Thromb Haemost **122**, 113-122 (2022).

14.   E. Barberis *et al.*, Circulating Exosomes Are Strongly Involved in SARS-CoV-2 Infection. *Front Mol*

Biosci **8**, 632290 (2021).

15.   Y. Fujita *et al.*, Early prediction of COVID-19 severity using extracellular vesicle COPB2. *J Extracell*

Vesicles **10**, e12092 (2021).

16.   K. Mao *et al.*, Proteomics of extracellular vesicles in plasma reveals the characteristics and residual traces

of COVID-19 patients without underlying diseases after 3 months of recovery. *Cell Death Dis* **12**, 541

(2021).

17.   F. Puhm, L. Flamand, E. Boilard, Platelet extracellular vesicles in COVID-19: Potential markers and

makers. *J Leukoc Biol* **111**, 63-74 (2022).

18.   K. H. W. Yim, S. Borgoni, R. Chahwan, Serum extracellular vesicles profiling is associated with COVID-

19 progression and immune responses. *J Extracell Biol* **1**, e37 (2022).

19.   C. Salomon *et al.*, A gestational profile of placental exosomes in maternal plasma and their effects on

endothelial cell migration. *PLoS One* **9**, e98667 (2014).

20.   S. Sarker *et al.*, Placenta-derived exosomes continuously increase in maternal circulation over the first

trimester of pregnancy. *J Transl Med* **12**, 204 (2014).

21.   J. Ratajczak *et al.*, Paracrine proangiopoietic effects of human umbilical cord blood-derived purified

CD133+ cells--implications for stem cell therapies in regenerative medicine. *Stem Cells Dev* **22**, 422-430

(2013).

22.   K. Bai *et al.*, Placenta-Derived Exosomes as a Modulator in Maternal Immune Tolerance During

Pregnancy. *Front Immunol* **12**, 671093 (2021).

23.   X. Chang *et al.*, Exosomes From Women With Preeclampsia Induced Vascular Dysfunction by Delivering

sFlt (Soluble Fms-Like Tyrosine Kinase)-1 and sEng (Soluble Endoglin) to Endothelial Cells. *Hypertension*

72, 1381-1390 (2018).

24.   I. Hromadnikova, L. Dvorakova, K. Kotlabova, L. Krofta, The Prediction of Gestational Hypertension,

Preeclampsia and Fetal Growth Restriction via the First Trimester Screening of Plasma Exosomal C19MC

microRNAs. *Int J Mol Sci* **20**, (2019).

25.   P. Pillay, N. Maharaj, J. Moodley, I. Mackraj, Placental exosomes and pre-eclampsia: Maternal circulating
747   levels in normal pregnancies and, early and late onset pre-eclamptic pregnancies. *Placenta* **46**, 18-25
748   (2016).

749   26.   C. Salomon *et al.*, Gestational Diabetes Mellitus Is Associated With Changes in the Concentration and
750   Bioactivity of Placenta-Derived Exosomes in Maternal Circulation Across Gestation. *Diabetes* **65**, 598-609
751   (2016).

752   27.   M. Wilkinson, E. D. Johnstone, L. E. Simcox, J. E. Myers, The impact of COVID-19 on pregnancy
753   outcomes in a diverse cohort in England. *Sci Rep* **12**, 942 (2022).

754   28.   N. Alcover *et al.*, Systematic review and synthesis of stillbirths and late miscarriages following SARS-
755   CoV-2 infections. *Am J Obstet Gynecol* **229**, 118-128 (2023).

756   29.   K. Narang *et al.*, Impact of asymptomatic and mild COVID-19 infection on fetal growth during pregnancy.
757   *Eur J Obstet Gynecol Reprod Biol* **281**, 63-67 (2023).

758   30.   R. N. Baergen, D. S. Heller, J. A. Goldstein, Placental Pathology in COVID-19. *Am J Clin Pathol* **154**, 279
759   (2020).

760   31.   M. C. Sharps *et al.*, A structured review of placental morphology and histopathological lesions associated
761   with SARS-CoV-2 infection. *Placenta* **101**, 13-29 (2020).

762   32.   T. Ohmaru-Nakanishi *et al.*, Fibrosis in Preeclamptic Placentas Is Associated with Stromal Fibroblasts
763   Activated by the Transforming Growth Factor-beta1 Signaling Pathway. *Am J Pathol* **188**, 683-695 (2018).

764   33.   H. Li *et al.*, Human Placental Endothelial Cell and Trophoblast Heterogeneity and Differentiation Revealed
765   by Single-Cell RNA Sequencing. *Cells* **12**, (2022).

766   34.   A. G. Foote, Z. Wang, C. Kendziorski, S. L. Thibeault, Tissue specific human fibroblast differential
767   expression based on RNAsequencing analysis. *BMC Genomics* **20**, 308 (2019).

768   35.   F. Fang, Y. Yao, Z. Ma, Exploration of the Long Noncoding RNAs Involved in the Crosstalk between M2
769   Macrophages and Tumor Metabolism in Lung Cancer. *Genet Res (Camb)* **2023**, 4512820 (2023).

770   36.   A. Liso *et al.*, IGFBP-6: At the Crossroads of Immunity, Tissue Repair and Fibrosis. *Int J Mol Sci* **23**,
771   (2022).

772   37.   D. Chakraborty *et al.*, HIF-KDM3A-MMP12 regulatory circuit ensures trophoblast plasticity and placental
773   adaptations to hypoxia. *Proc Natl Acad Sci U S A* **113**, E7212-E7221 (2016).

38. H. Mogami *et al.*, Fetal fibronectin signaling induces matrix metalloproteases and cyclooxygenase-2

(COX-2) in amnion cells and preterm birth in mice. *J Biol Chem* **288**, 1953-1966 (2013).

39. Y. Okamura *et al.*, The extra domain A of fibronectin activates Toll-like receptor 4. *J Biol Chem* **276**,

10229-10233 (2001).

40. A. J. Camden *et al.*, Growth regulation by estrogen in breast cancer 1 (GREB1) is a novel progesterone-

responsive gene required for human endometrial stromal decidualization. *Mol Hum Reprod* **23**, 646-653

(2017).

41. S. B. Chadchan *et al.*, A GREB1-steroid receptor feedforward mechanism governs differential GREB1

action in endometrial function and endometriosis. *Nat Commun* **15**, 1947 (2024).

42. S. Biswas Shivhare, J. N. Bulmer, B. A. Innes, D. K. Hapangama, G. E. Lash, Altered vascular smooth

muscle cell differentiation in the endometrial vasculature in menorrhagia. *Hum Reprod* **29**, 1884-1894

(2014).

43. G. S. Whitley, J. E. Cartwright, Cellular and molecular regulation of spiral artery remodelling: lessons from

the cardiovascular field. *Placenta* **31**, 465-474 (2010).

44. B. Afzali, M. Noris, B. N. Lambrecht, C. Kemper, The state of complement in COVID-19. *Nat Rev*

Immunol **22**, 77-84 (2022).

45. C. B. Chighizola, P. A. Lonati, L. Trespidi, P. L. Meroni, F. Tedesco, The Complement System in the

Pathophysiology of Pregnancy and in Systemic Autoimmune Rheumatic Diseases During Pregnancy. *Front*

Immunol **11**, 2084 (2020).

46. L. Li *et al.*, Research progress on the STAT signaling pathway in pregnancy and pregnancy-associated

disorders. *Front Immunol* **14**, 1331964 (2023).

47. L. Zhang *et al.*, The diagnosis values of serum STAT4 and sEng in preeclampsia. *J Clin Lab Anal* **34**,

e23073 (2020).

48. O. Parant *et al.*, CD34+ cells in maternal placental blood are mainly fetal in origin and express endothelial

markers. *Lab Invest* **89**, 915-923 (2009).

49. C. Salomon *et al.*, Hypoxia-induced changes in the bioactivity of cytotrophoblast-derived exosomes. *PLoS*

One **8**, e79636 (2013).

50.   P. Kaufmann, S. Black, B. Huppertz, Endovascular trophoblast invasion: implications for the pathogenesis

802   of intrauterine growth retardation and preeclampsia. *Biol Reprod* **69**, 1-7 (2003).

803   51.   P. Tantbirojn, C. P. Crum, M. M. Parast, Pathophysiology of placenta creta: the role of decidua and

804   extravillous trophoblast. *Placenta* **29**, 639-645 (2008).

805   52.   M. Prieto-Vila, Y. Yoshioka, T. Ochiya, Biological Functions Driven by mRNAs Carried by Extracellular

806   Vesicles in Cancer. *Front Cell Dev Biol* **9**, 620498 (2021).

807   53.   M. C. Deregibus *et al.*, Endothelial progenitor cell derived microvesicles activate an angiogenic program in

808   endothelial cells by a horizontal transfer of mRNA. *Blood* **110**, 2440-2448 (2007).

809   54.   J. Ratajczak *et al.*, Embryonic stem cell-derived microvesicles reprogram hematopoietic progenitors:

810   evidence for horizontal transfer of mRNA and protein delivery. *Leukemia* **20**, 847-856 (2006).

811   55.   H. Valadi *et al.*, Exosome-mediated transfer of mRNAs and microRNAs is a novel mechanism of genetic

812   exchange between cells. *Nat Cell Biol* **9**, 654-659 (2007).

813   56.   Z. Wei *et al.*, Coding and noncoding landscape of extracellular RNA released by human glioma stem cells.

814   *Nat Commun* **8**, 1145 (2017).

815   57.   C. Shu, P. Xu, J. Han, S. Han, J. He, Upregulation of circRNA hsa_circ_0008726 in Pre-eclampsia Inhibits

816   Trophoblast Migration, Invasion, and EMT by Regulating miR-345-3p/RYBP Axis. *Reprod Sci* **29**, 2829-

817   2841 (2022).

818   58.   L. Wang *et al.*, The THO complex regulates pluripotency gene mRNA export and controls embryonic stem

819   cell self-renewal and somatic cell reprogramming. *Cell Stem Cell* **13**, 676-690 (2013).

820   59.   J. D. Domizio *et al.*, The cGAS-STING pathway drives type I IFN immunopathology in COVID-19.

821   *Nature* **603**, 145-151 (2022).

822   60.   M. I. Faizan *et al.*, NSP4 and ORF9b of SARS-CoV-2 Induce Pro-Inflammatory Mitochondrial DNA

823   Release in Inner Membrane-Derived Vesicles. *Cells* **11**, (2022).

824   61.   L. Gibellini *et al.*, Altered bioenergetics and mitochondrial dysfunction of monocytes in patients with

825   COVID-19 pneumonia. *EMBO Mol Med* **12**, e13001 (2020).

826   62.   B. Appelman *et al.*, Muscle abnormalities worsen after post-exertional malaise in long COVID. *Nat*

827   *Commun* **15**, 17 (2024).

828   63.   D. Scozzi *et al.*, Circulating mitochondrial DNA is an early indicator of severe illness and mortality from

829   COVID-19. *JCI Insight* **6**, (2021).

830   64.   S. Shoraka, S. R. Mohebbi, S. M. Hosseini, M. R. Zali, Comparison of plasma mitochondrial DNA copy

831   number in asymptomatic and symptomatic COVID-19 patients. *Front Microbiol* **14**, 1256042 (2023).

832   65.   J. J. Valdes-Aguayo *et al.*, Peripheral Blood Mitochondrial DNA Levels Were Modulated by SARS-CoV-2

833   Infection Severity and Its Lessening Was Associated With Mortality Among Hospitalized Patients With

834   COVID-19. *Front Cell Infect Microbiol* **11**, 754708 (2021).

835   66.   T. J. Costa *et al.*, Mitochondrial DNA and TLR9 activation contribute to SARS-CoV-2-induced endothelial

836   cell damage. *Vascul Pharmacol* **142**, 106946 (2022).

837   67.   L. Peruzzotti-Jametti *et al.*, Neural stem cells traffic functional mitochondria via extracellular vesicles.

838   *PLoS Biol* **19**, e3001166 (2021).

839   68.   D. G. Phinney *et al.*, Mesenchymal stem cells use extracellular vesicles to outsource mitophagy and shuttle

840   microRNAs. *Nat Commun* **6**, 8472 (2015).

841   69.   P. Sansone *et al.*, Packaging and transfer of mitochondrial DNA via exosomes regulate escape from

842   dormancy in hormonal therapy-resistant breast cancer. *Proc Natl Acad Sci U S A* **114**, E9066-E9075

843   (2017).

844   70.   M. Schorpp-Kistner, Z. Q. Wang, P. Angel, E. F. Wagner, JunB is essential for mammalian placentation.

845   *EMBO J* **18**, 934-948 (1999).

846   71.   A. M. Nuzzo *et al.*, JunB/cyclin-D1 imbalance in placental mesenchymal stromal cells derived from

847   preeclamptic pregnancies with fetal-placental compromise. *Placenta* **35**, 483-490 (2014).

848   72.   A. T. Papageorghiou *et al.*, Preeclampsia and COVID-19: results from the INTERCOVID prospective

849   longitudinal study. *Am J Obstet Gynecol* **225**, 289 e281-289 e217 (2021).

850   73.   N. Berkane *et al.*, From Pregnancy to Preeclampsia: A Key Role for Estrogens. *Endocr Rev* **38**, 123-144

851   (2017).

852   74.   W. Bahia *et al.*, Association of genetic variants in Estrogen receptor (ESR)1 and ESR2 with susceptibility

853   to recurrent pregnancy loss in Tunisian women: A case control study. *Gene* **736**, 144406 (2020).

854   75.   H. A. El-Beshbishy *et al.*, Estrogen Receptor Alpha (ESR1) Gene Polymorphisms in Pre-eclamptic Saudi

855   Patients. *Pak J Med Sci* **31**, 880-885 (2015).

76.   A. A. Freedman, L. S. Keenan-Devlin, A. Borders, G. E. Miller, L. M. Ernst, Formulating a Meaningful

and Comprehensive Placental Phenotypic Classification. *Pediatr Dev Pathol* **24**, 337-350 (2021).

77.   T. Y. Khong *et al.*, Sampling and Definitions of Placental Lesions: Amsterdam Placental Workshop Group

Consensus Statement. *Arch Pathol Lab Med* **140**, 698-713 (2016).

78.   R. W. Redline, S. Ravishankar, C. M. Bagby, S. T. Saab, S. Zarei, Four major patterns of placental injury: a

stepwise guide for understanding and implementing the 2016 Amsterdam consensus. *Mod Pathol* **34**, 1074-

1092 (2021).

79.   C. Thery *et al.*, Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position

statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines.

J Extracell Vesicles **7**, 1535750 (2018).

80.   L. Anton, A. G. Brown, S. Parry, M. A. Elovitz, Lipopolysaccharide induces cytokine production and

decreases extravillous trophoblast invasion through a mitogen-activated protein kinase-mediated pathway:

possible mechanisms of first trimester placental dysfunction. *Hum Reprod* **27**, 61-72 (2012).

81.   L. Anton *et al.*, HIF-1alpha Stabilization Increases miR-210 Eliciting First Trimester Extravillous

Trophoblast Mitochondrial Dysfunction. *Front Physiol* **10**, 699 (2019).

82.   S. Getsios, G. T. Chen, D. T. Huang, C. D. MacCalman, Regulated expression of cadherin-11 in human

extravillous cytotrophoblasts undergoing aggregation and fusion in response to transforming growth factor

beta 1. *J Reprod Fertil* **114**, 357-363 (1998).

83.   C. H. Graham, J. J. Lysiak, K. R. McCrae, P. K. Lala, Localization of transforming growth factor-beta at

the human fetal-maternal interface: role in trophoblast growth and differentiation. *Biol Reprod* **46**, 561-572

(1992).

84.   J. Y. Park *et al.*, A microphysiological model of human trophoblast invasion during implantation. *Nat

Commun* **13**, 1252 (2022).

85.   S. A. Ware *et al.*, An automated, high-throughput methodology optimized for quantitative cell-free

mitochondrial and nuclear DNA isolation from plasma. *J Biol Chem* **295**, 15677-15691 (2020).

86.   J. J. Trombetta *et al.*, Preparation of Single-Cell RNA-Seq Libraries for Next Generation Sequencing. *Curr

Protoc Mol Biol* **107**, 4 22 21-24 22 17 (2014).

87.   O. Yukselen, O. Turkyilmaz, A. R. Ozturk, M. Garber, A. Kucukural, DolphinNext: a distributed data

884   processing platform for high throughput genomics. *BMC Genomics* **21**, 310 (2020).

885   88.   K. S. Rehman, S. Yin, B. A. Mayhew, R. A. Word, W. E. Rainey, Human myometrial adaptation to

886   pregnancy: cDNA microarray gene expression profiling of myometrium from non-pregnant and pregnant

887   women. *Mol Hum Reprod* **9**, 681-700 (2003).

888   89.   C. M. Whittington *et al.*, Transcriptomic changes in the pre-implantation uterus highlight histotrophic

889   nutrition of the developing marsupial embryo. *Sci Rep* **8**, 2412 (2018).

890   90.   S. Peng *et al.*, Genetic regulation of the placental transcriptome underlies birth weight and risk of childhood

891   obesity. *PLoS Genet* **14**, e1007799 (2018).

892   91.   J. L. Maron *et al.*, Gene expression analysis in pregnant women and their infants identifies unique fetal

893   biomarkers that circulate in maternal blood. *J Clin Invest* **117**, 3007-3019 (2007).

894   92.   G. Song, F. W. Bazer, T. E. Spencer, Differential expression of cathepsins and cystatin C in ovine

895   uteroplacental tissues. *Placenta* **28**, 1091-1098 (2007).

896   93.   C. P. Chen *et al.*, Altered placental syncytin and its receptor ASCT2 expression in placental development

897   and pre-eclampsia. *BJOG* **113**, 152-158 (2006).

898   94.   K. L. Fulghum *et al.*, Metabolic signatures of pregnancy-induced cardiac growth. *Am J Physiol Heart Circ*

899   *Physiol* **323**, H146-H164 (2022).

900   95.   M. Prater *et al.*, RNA-Seq reveals changes in human placental metabolism, transport and endocrinology

901   across the first-second trimester transition. *Biol Open* **10**, (2021).

902   96.   M. B. Rabaglino, J. M. Sanchez, M. McDonald, E. O'Callaghan, P. Lonergan, Maternal blood

903   transcriptome as a sensor of foetal organ maturation at the end of organogenesis in cattle. *Biol Reprod*,

904   (2023).

905   97.   C. Y. Hung *et al.*, A defect in the inner kinetochore protein CENPT causes a new syndrome of severe

906   growth failure. *PLoS One* **12**, e0189324 (2017).

907   98.   W. Li *et al.*, Role of IGF2BP3 in trophoblast cell invasion and migration. *Cell Death Dis* **5**, e1025 (2014).

908   99.   H. Wen, L. Chen, J. He, J. Lin, MicroRNA expression profiles and networks in placentas complicated with

909   selective intrauterine growth restriction. *Mol Med Rep* **16**, 6650-6673 (2017).

100.   G. Wang *et al.*, Impact of intrauterine exposure to maternal diabetes on preterm birth: fetal DNA

methylation alteration is an important mediator. *Clin Epigenetics* **15**, 59 (2023).

101.   I. L. Aye, T. L. Powell, T. Jansson, Review: Adiponectin--the missing link between maternal adiposity,

placental transport and fetal growth? *Placenta* **34 Suppl**, S40-45 (2013).

102.   I. Hromadnikova *et al.*, Expression profile of C19MC microRNAs in placental tissue in pregnancy-related

complications. *DNA Cell Biol* **34**, 437-457 (2015).

103.   Y. Gu, S. A. Burlison, Y. Wang, PAF levels and PAF-AH activities in placentas from normal and

preeclamptic pregnancies. *Placenta* **27**, 744-749 (2006).

104.   T. Garrido-Gomez *et al.*, Severe pre-eclampsia is associated with alterations in cytotrophoblasts of the

smooth chorion. *Development* **144**, 767-777 (2017).

105.   R. Calicchio *et al.*, Preeclamptic plasma induces transcription modifications involving the AP-1

transcriptional regulator JDP2 in endothelial cells. *Am J Pathol* **183**, 1993-2006 (2013).

106.   W. Fan *et al.*, Upregulation of METTL14 contributes to trophoblast dysfunction by elevating FOXO3a

expression in an m(6)A-dependent manner. *Placenta* **124**, 18-27 (2022).

107.   J. Wang, F. Gao, X. Zhao, Y. Cai, H. Jin, Integrated analysis of the transcriptome-wide m6A methylome in

preeclampsia and healthy control placentas. *PeerJ* **8**, e9880 (2020).

108.   M. Diaz *et al.*, Placental and Cord Blood Methylation of Genes Involved in Energy Homeostasis:

Association With Fetal Growth and Neonatal Body Composition. *Diabetes* **66**, 779-784 (2017).

109.   P. Murthi, G. Rajaraman, J. Erwich, E. Dimitriadis, Decreased Placental FPR2 in Early Pregnancies That

Later Developed Small-For-Gestation Age: A Potential Role of FPR2 in the Regulation of Epithelial-

Mesenchymal Transition. *Cells* **9**, (2020).

110.   A. Chu *et al.*, Aldehyde dehydrogenase isoforms and inflammatory cell populations are differentially

expressed in term human placentas affected by intrauterine growth restriction. *Placenta* **81**, 9-17 (2019).

111.   A. Dobierzewska *et al.*, Impairment of Angiogenic Sphingosine Kinase-1/Sphingosine-1-Phosphate

Receptors Pathway in Preeclampsia. *PLoS One* **11**, e0157221 (2016).

112.   M. H. Fenstad *et al.*, Genetic and molecular functional characterization of variants within TNFSF13B, a

positional candidate preeclampsia susceptibility gene on 13q. *PLoS One* **5**, (2010).

113. J. G. Grudzinskas *et al.*, Identification of high-risk pregnancy by the routine measurement of pregnancy-

specific beta 1-glycoprotein. *Am J Obstet Gynecol* **147**, 10-12 (1983).

114. K. Pihl, T. Larsen, I. Laursen, L. Krebs, M. Christiansen, First trimester maternal serum pregnancy-specific

beta-1-glycoprotein (SP1) as a marker of adverse pregnancy outcome. *Prenat Diagn* **29**, 1256-1261 (2009).

115. R. M. Silver, K. D. Heyborne, K. K. Leslie, Pregnancy specific beta 1 glycoprotein (SP-1) in maternal

serum and amniotic fluid; pre-eclampsia, small for gestational age fetus and fetal distress. *Placenta* **14**,

583-589 (1993).

116. M. A. Mohamad *et al.*, A Review of Candidate Genes and Pathways in Preeclampsia-An Integrated

Bioinformatical Analysis. *Biology (Basel)* **9**, (2020).

117. B. Vora *et al.*, Meta-Analysis of Maternal and Fetal Transcriptomic Data Elucidates the Role of Adaptive

and Innate Immunity in Preterm Birth. *Front Immunol* **9**, 993 (2018).

118. M. N. Moufarrej *et al.*, Early prediction of preeclampsia in pregnancy with cell-free RNA. *Nature* **602**,

689-694 (2022).

119. J. Shang, L. Lin, X. Huang, L. Zhou, Q. Huang, Re-expression of circ_0043610 contributes to trophoblast

dysfunction through the miR-558/RYBP pathway in preeclampsia. *Endocr J* **69**, 1373-1385 (2022).

120. Y. Zhang, S. Fang, J. Wang, S. Chen, R. Xuan, Hsa_circ_0008726 regulates the proliferation, migration,

and invasion of trophoblast cells in preeclampsia through modulating the miR-1290-LHX6 signaling

pathway. *J Clin Lab Anal* **36**, e24540 (2022).

121. J. Wang *et al.*, Alpha-2-macroglobulin is involved in the occurrence of early-onset pre-eclampsia via its

negative impact on uterine spiral artery remodeling and placental angiogenesis. *BMC Med* **21**, 90 (2023).

122. J. A. Tronco *et al.*, Alpha-2-macroglobulin from circulating exosome-like vesicles is increased in women

with preterm pregnancies. *Sci Rep* **10**, 16961 (2020).

123. K. Leavey, S. L. Wilson, S. A. Bainbridge, W. P. Robinson, B. J. Cox, Epigenetic regulation of placental

gene expression in transcriptional subtypes of preeclampsia. *Clin Epigenetics* **10**, 28 (2018).

124. R. K. Yuen, M. S. Penaherrera, P. von Dadelszen, D. E. McFadden, W. P. Robinson, DNA methylation

profiling of human placentas reveals promoter hypomethylation of multiple genes in early-onset

preeclampsia. *Eur J Hum Genet* **18**, 1006-1012 (2010).

125.   J. Pan, X. Tian, H. Huang, N. Zhong, Proteomic Study of Fetal Membrane: Inflammation-Triggered

965   Proteolysis of Extracellular Matrix May Present a Pathogenic Pathway for Spontaneous Preterm Birth.

966   *Front Physiol* **11**, 800 (2020).

967   126.   T. Lekva *et al.*, Dysregulated non-coding telomerase RNA component and associated exonuclease XRN1

968   in leucocytes from women developing preeclampsia-possible link to enhanced senescence. *Sci Rep* **11**,

969   19735 (2021).

970   127.   W. Lai, L. Yu, Elevated MicroRNA 183 Impairs Trophoblast Migration and Invasiveness by

971   Downregulating FOXP1 Expression and Elevating GNG7 Expression during Preeclampsia. *Mol Cell Biol*

972   **41**, (2020).

973   128.   J. Chen *et al.*, Silencing of long non-coding RNA NEAT1 improves Treg/Th17 imbalance in preeclampsia

974   via the miR-485-5p/AIM2 axis. *Bioengineered* **12**, 8768-8777 (2021).

975   129.   S. Gremlich *et al.*, The long non-coding RNA NEAT1 is increased in IUGR placentas, leading to potential

976   new hypotheses of IUGR origin/development. *Placenta* **35**, 44-49 (2014).

977   130.   J. Schuster *et al.*, Protein Network Analysis of Whole Exome Sequencing of Severe Preeclampsia. *Front*

978   *Genet* **12**, 765985 (2021).

979   131.   Z. Miao, M. Chen, H. Wu, H. Ding, Z. Shi, Comparative proteomic profile of the human placenta in normal

980   and fetal growth restriction subjects. *Cell Physiol Biochem* **34**, 1701-1710 (2014).

981   132.   J. Zhao *et al.*, The m(6)A methyltransferase METTL3 promotes trophoblast cell invasion by regulating

982   MYLK expression. *Placenta* **129**, 1-6 (2022).

983   133.   J. Zadora *et al.*, Disturbed Placental Imprinting in Preeclampsia Leads to Altered Expression of DLX5, a

984   Human-Specific Early Trophoblast Marker. *Circulation* **136**, 1824-1839 (2017).

985   134.   H. Tiensuu *et al.*, Human placental proteomics and exon variant studies link AAT/SERPINA1 with

986   spontaneous preterm birth. *BMC Med* **20**, 141 (2022).

987   135.   F. Zhou, T. Cheng, Y. Xing, H. Ma, L. Yang, Network exploration of gene signatures underlying low birth

988   weight induced metabolic alterations. *Medicine (Baltimore)* **101**, e31489 (2022).

989   136.   M. A. Deyssenroth *et al.*, Whole-transcriptome analysis delineates the human placenta gene network and its

990   associations with fetal growth. *BMC Genomics* **18**, 520 (2017).

991   137.   D. A. Enquobahrie, C. Qiu, S. Y. Muhie, M. A. Williams, Maternal peripheral blood gene expression in

992   early pregnancy and preeclampsia. *Int J Mol Epidemiol Genet* **2**, 78-94 (2011).

993   138.   A. K. Edwards, K. A. Dunlap, T. E. Spencer, M. C. Satterfield, Identification of Pathways Associated with

994   Placental Adaptation to Maternal Nutrient Restriction in Sheep. *Genes (Basel)* **11**, (2020).

995