Exploring the Genomic Landscape of the GP63 family in *Trypanosoma cruzi*: Evolutionary Dynamics and Functional Peculiarities

Luisa Berná[1,2], María Laura Chiribao[1,3], Sebastian Pita[1,4], Fernando Alvarez-Valin[2], Adriana Parodi-Talice*[1,4]

[1] Laboratorio de Interacciones Hospedero–Patógeno—UBM, Institut Pasteur Montevideo, Montevideo 11400, Uruguay
[2] Laboratorio de Genómica Evolutiva, Facultad de Ciencias, Universidad de la República, Montevideo 11400, Uruguay
[3] Departamento de Bioquímica, Facultad de Medicina, Universidad de la República, Montevideo 11800, Uruguay
[4] Sección Genética Evolutiva, Instituto de Biología, Facultad de Ciencias, Universidad de la República, Montevideo 11400, Uruguay

*Corresponding author's email address apartal@fcien.edu.uy

Keywords: Zinc-metalloprotease GP63, Leishmanolysin, *Trypanosoma cruzi*, multigene family evolution

ABSTRACT

We analyzed the complete set of GP63 sequences from the parasitic protozoa *Trypanosoma cruzi*. Our analysis allowed us to refine annotation of sequences previously identified as functional and pseudogenes. Concerning the latter, we unified pseudogenic fragments derived from the same functional gene and excluded sequences incorrectly annotated as GP63 pseudogenes. We were able to identify eleven GP63 gene groups, which are sharply defined and have a high intra-group sequence identity. The sequences of each group showed a strong preference for genomic compartments. Some groups are located in the core and others in disruptive compartments of the *T. cruzi* genome. Groups located in the core compartment often contain tandem arrays of GP63 genes. On the contrary, genes from groups located in the disruptive compartment tend to be surrounded by genes encoding surface proteins such as MASP, mucins and trans-sialidases. Analysis of the immediate GP63 environments showed differences that may be the result of different genomic dynamics in these two compartments. Interestingly, each GP63 group showed a particular mRNA expression profile and some groups contain members that are differentially expressed between life cycle stages, being expressed at higher levels in trypomastigotes than in the replicative forms. This suggests that these groups of GP63 proteins may play a relevant role in the infective stage. The analysis of the M8 domain, that defines the GP63 protein family, allowed us to recognize that each group presented peculiarities in the conserved sites as well as in the presence of the predicted signal peptide and GPI anchor site. Phylogenetic analysis of the GP63 sequences, including other species of the genus *Trypanosoma* as well as other kinetoplastids, showed that ten of the 11 groups of *T. cruzi* not only are also present in the other *Trypanosoma* species but also are exclusive of genus, suggesting that the diversification of these subfamilies took place before speciation. However, each species then followed a different evolutionary path, amplifying specific groups in unique ways.

Data summary
The authors confirm all supporting data, code and protocols have been provided within the
article or through supplementary data files.

Impact statement

Our study contributes to the understanding of the GP63 gene family in *Trypanosoma*
cruzi, a crucial protein for the parasite's infectivity and evolution. We refined the
annotation of GP63 sequences, identifying eleven distinct gene groups with distinctive
preferences for genomic compartments -some in the core, others in the disruptive
compartment. This distribution hints at varied genomic dynamics and potential roles in
the parasite's life cycle, especially since some groups show enhanced expression in
infective stages, suggesting their importance in disease transmission.

Our exploration into the GP63 sequences' M8 domain revealed group-specific
peculiarities in conserved sites and structural motifs, emphasizing functional diversity.
Phylogenetic analysis across *Trypanosoma* species highlighted the evolutionary
uniqueness of these gene subfamilies within the genus, underscoring their role in the
species' distinct evolutionary paths and amplification patterns.

INTRODUCTION

Chagas disease, caused by the *Trypanosoma cruzi* parasite, affects approximately 7
million individuals worldwide and is classified as a neglected disease by the World
Health Organization (WHO Chagas disease). Primarily prevalent in rural regions of
Latin America, this disease holds a significant impact. The progression of Chagas
disease heavily relies on the interaction of parasitic virulence factors and the defensive
responses of the mammalian host. In this context, identifying parasite molecules that
engage with the host's immune or signaling systems could present novel targets crucial
for disease intervention.

The surface of the parasite, especially during its infective stages, serves as a crucial
interface for interaction with the vertebrate host. Surface proteins belonging to families
such as mucins, MASPs and trans-sialidases are prominently expressed in this
domain. While advancements have been achieved in understanding the surface
architecture of these parasites (1–3), the precise roles of individual components in
host-parasite interactions, as well as the dynamic expression patterns of all
constituents during the infection process, remain elusive.

Although different species exhibit distinct types of surface components, GP63
glycoproteins are the only surface protein components that are universally present
across all trypanosomatid species. The genes encoding GP63 proteins constitute a
family that exhibits considerable variability in both number and structure across these
species (4).

The GP63 protein is a metalloprotease that belongs to the group of zinc-dependent
endopeptidases. Sequences akin to GP63 are also found in other insect and plant
parasites (4). The diversification of trypanosomatid genomes has been accompanied
by the rapid evolution of their multigene families, including the GP63 family, which
underwent substantial modification after the emergence of parasitism. This is
particularly evident when comparing these genomes to those of free-living
kinetoplastids (5). Phylogenetic studies of the GP63 family showed that both *T. cruzi*

and *Leishmania* sp. have independently expanded their gene repertoires (5).
Therefore, this family has had a long evolutionary history, and it has been postulated
that co-evolutionary processes of parasites and their hosts have significantly influenced
the structure and functionality of its members (4,6).

The GP63 protein was first described in *Leishmania* (7), where it has been extensively
studied ever since. Some members are known to be expressed on the surface, others
are secreted, and others remain intracellular (8). It is the most abundant protein in the
parasite (500,000 molecules/parasite). GP63 in *Leishmania* plays several roles
relevant for infection such as: inhibition of complement-mediated lysis, facilitation of
phagocytosis, degradation of extracellular matrix components, alteration of host cell
signaling pathways and immune evasion (8–10). This multiplicity of functions suggests
a preponderant role for parasite biology. In contrast to the extensive evidence obtained
in *Leishmania*, very little is known about the functions of this family in *T. cruzi*, despite
its abundance in the genome.

Some studies showed that members of the GP63 family of *T. cruzi* may be involved in
infection. This is supported by the fact that pre-incubation with specific antibodies
partially blocked infection of Vero cells by trypomastigotes (11). Other studies showed
that GP63 RNAs are differentially expressed depending on the stage of the parasite life
cycle (12). A few studies on metalloproteinase activity detected on gelatin SDS-PAGE
gels have been reported for a group of GP63 proteins of 75-78 kDa (11,13)

The variability in the GP63 gene family across *Leishmania* species, ranging from 4 to
33 genes, contrasts starkly with the findings in *Trypanosoma cruzi*. The hybrid strain
CLBrener's genome, initially reported in 2005 by El-Sayed et al.(14), contained 425
GP63 copies. However, more recent genomic analyses using long-read, third-
generation sequencing techniques have provided a more accurate assembly and
annotation, revealing 378 GP63-encoding genes in the Dm28c strain of *T. cruzi* and
718 in the hybrid strain TCC (15). This sequencing approach not only offered a more
realistic gene count by resolving previously collapsed repetitive regions but also
unveiled a compartmentalized, or 'bipartite', genome structure with distinct GC content.
Specifically, we described the *T. cruzi* genome as comprising of a core compartment,
which contains genes conserved across trypanosomatids and encodes both known and
hypothetical proteins, and a disruptive compartment. This latter is composed by
genomic regions that are unique to *T. cruzi* and enriched with species-specific gene
families such as mucins, mucin-associated surface proteins (MASPs), and trans-
sialidases. Notably, GP63 family members are distributed across both genomic
compartments. The specific attributes of the genes within these compartments,
reflecting their unique evolutionary and functional dynamics, are yet to be fully
understood.

In *Leishmania* species, at least 60% of the amino acids in the GP63 protein, also
known as Leishmanolysin, are conserved. This includes the short zinc-binding motif
HEXXH and 19 cysteine residues distributed along the sequence. Following synthesis,
Leishmanolysin undergoes several post-translational modifications: the signal
sequence is cleaved in the endoplasmic reticulum, N-glycosylation is applied, and
roughly 25 C-terminal amino acids are substituted by a GPI anchor. The protein is
further processed by proteolytic cleavage of a propeptide, yielding the mature form.
The pro-peptide sequence harbors a cysteine residue conserved across all GP63
variants in different species (16). In *Leishmania major*, this cysteine residue is
implicated in a 'cysteine switch' mechanism, where it plays a role in inhibiting protease
activity by chelating the zinc atom at the active site. The mature protein in *L. major*

begins at Val100, features three N-glycosylation sites at Asn300, Asn407, and Asn534,
and concludes at Asn577, which is covalently attached via ethanolamine to the
carbohydrate portion of the GPI anchor.

The zinc-binding motif HEXXH located in the amino-terminal domain of GP63 is the
characteristic site of metalloproteases. Research demonstrates that the two histidines
and one glutamic acid within this motif are conserved across all GP63 sequences in the
trypanosomatids studied, playing a crucial role in proteolytic activity (17). Mutations in
any of these three residues result in a complete loss of catalytic function. Specifically, a
mutation from glutamic acid to aspartic acid at position E165 in the GP63 of *L. major*
leads to a loss of protease activity (18).

In this research, we aimed to extensively characterize the GP63 gene family in the
genome of *T. cruzi*, examining its subtypes, domains, genomic organization,
evolutionary dynamics, expression profiles, and structural characteristics. Through this
detailed analysis, we seek to contribute with enough knowledge necessary to elucidate
the unique role of these proteins to pathogenesis and immune system evasion.

158
159 RESULTS AND DISCUSSION
160
161 1 - The metalloprotease GP63 family contains multiple functional gene copies
162 and several pseudogenes in the genome of *Trypanosoma cruzi*

To analyze the GP63 family of metalloproteases, we utilized the annotated Dm28c
genome, sequenced using long-read technology (15). This approach is crucial for
including all family copies. In this genome, we identified 96 complete gene sequences
encoding GP63 proteins. Furthermore, the genome possesses annotations for 281
potential GP63 pseudogenes. The primary criterion for classifying a sequence as a
pseudogene was its length; specifically, sequences encoding polypeptides of 460 or
fewer amino acids were deemed pseudogenes.

Upon examining the length distribution of the entire sequence population, a broad
range of variation is evident, displaying a clear division between potential functional
sequences and pseudogenes (Figure 1A). Furthermore, sequences classified as
functional also exhibit significant variability. In fact, the distribution is bimodal, peaking
at lengths of approximately 560 and 700 amino acids (Figure 1B).

Figure 1: Distribution of GP63 amino acid sequence lengths in *T. cruzi* Dm28c. This figure displays two distinct length distributions: on the right, the combined length distribution of both GP63 genes and pseudogenes; on the left, solely GP63 functional genes.

Using length as the only criterion to categorize sequences as pseudogenes or functional, is incomplete to get a full grasp. These aspects led us to refine the annotation by incorporating additional criteria. Specifically, to classify sequences as functional, we now consider a sequence as such if it is longer than 460 amino acids, contains a full-length peptidase M8 domain and at least in the majority of the cases, presents a signal peptide (refer to Section 4 for further details).

Concerning pseudogenes, it is important to bear in mind that many sequences classified as pseudogenes may actually represent remnant fragments of the same former functional GP63 gene that underwent pseudogenization, as already reported for most trypanosomatid genes (19). In fact, among the 281 sequences previously classified as pseudogenes, 63 are observed in pairs or groups, either juxtaposed, overlapping, or separated by short distances. Analysis of these sequences shows that they are indeed sister fragments that originated by nonsense mutations or indels producing frameshifts of a previously intact coding sequence. Segments belonging to these groups are now classified as being parts of the same pseudogene. We also found that 29 pseudogenic fragments are not really related to GP63. In all likelihood, these fragments were originally incorrectly named as GP63 because of flawed transference of information based on homology. The revised list, now comprising 185 pseudogenes, is presented in Supplementary Table 1.

It is worth noting that many of these pseudogenic fragments retain high DNA sequence identity with complete (functional) genes, suggesting their status as "young" pseudogenes, namely nonsense mutations or indels causing pseudogenization occurred relatively recently. Moreover, comparative analysis of paired pseudogenic fragments, particularly at the indels sites, shows that several pairs (or groups) contain mutations at identical positions. This pattern suggests that they are derived from duplication events that have occurred after pseudogenization (Supplementary Figure 1).

2 - Eleven groups of GP63 genes are present in *Trypanosoma cruzi* genome

Previously, Cuevas et al. (11) conducted PCR analyses and identified *T. cruzi* sequences homologous to GP63 genes from *Leishmania*. They categorized GP63 genes in *T. cruzi* into three groups: Tcgp63-I and Tcgp63-II, and Tcgp63-III, consisting of pseudogenes. Subsequent re-analysis of the *T. cruzi* GP63 family genes led to their classification into four distinct groups (4). Within each group, the sequences exhibit 80% similarity, whereas between groups, the similarity drops to 30%. These authors identified a conserved segment of 100 amino acids following the active site, shared across all sequences, likely corresponding to the M8 domain, a feature ubiquitous to all GP63 sequences.

In this study, thanks to the development of third-generation technologies that allow sequencing of increasingly better-quality long reads, new *T. cruzi* genomes, with higher accuracy have become available. This advancement enabled us to conduct an exhaustive comparative analysis of the 96 annotated GP63 gene sequences, using both nucleotide and protein sequences (Supplementary Table 2). This approach led to the identification of 11 distinct groups within the GP63 gene family. Each group has a

characteristic average length of sequences, which in most groups has little variation
(Supplementary Figure 2). This classification is evident in a phylogenetic analysis
performed with the complete amino acid sequence (Figure 2A, Supplementary Figure
3). Furthermore, an amino acid identity matrix, created through all-against-all pairwise
comparisons, corroborated the observed clustering patterns (Figure 1B, Supplementary
Table 3). This matrix was visualized using Gephi, where each node represents a GP63
sequence and the color of each edge denotes the level of identity, and an edge weight
threshold of 65% was applied (Figure 2B).

Figure 2: Distinct Clustering of GP63 into 11 Groups. A) Phylogenetic analysis of amino acid
sequences of the 96 GP63 genes from *T. cruzi* Dm28c, illustrating the division into 11 unique groups. B)
Network visualization in Gephi of the pairwise identity distances among the 96 amino acid sequences of
GP63 genes, where nodes symbolize genes (color coding matches the phylogeny in A), and the color of
each edge indicates the level of identity. C) Identity matrix displaying average intra-group and inter-group
amino acid identities, highlighting the distinctiveness of each group. D) Network visualization similar to B,
with genes colored according to the genomic compartment to which they belong: green for the core
compartment and pink for the disruptive compartment.

Additionally, the identity matrix was condensed into a matrix representing average
intra-group and inter-group identity, as depicted in Figure 2C. The intra-group amino
acid identity levels are predominantly homogeneous, exceeding 90% in 8 out of 11
groups. In particular, three groups showed more variability, especially group G11,
which is the largest and exhibited the lowest homogeneity with 73% mean identity. In
contrast, amino acid identities between groups were notably lower, never exceeding
48%, indicating significant divergence among groups. This pronounced divergence
between groups, together with the strong homogeneity within most groups, justifies
their clustering into 11 distinct groups, as highlighted in Figure 2. Further expanding on
this classification, our BLAST analysis of *T. cruzi* genes and pseudogenes enabled us

to assign each pseudogene a parental group, thereby extending our clustering analysis
beyond genes to include pseudogenes. As revealed in Supplementary Table 2, a
significant proportion of these pseudogene sequences predominantly correspond to
groups G9 and G11, both of which are located in the disruptive compartment. This
inclusion of pseudogenes in our analysis underscores the comprehensive nature of the
classification, affirming the robustness of the 11-group clustering in capturing the
genetic diversity within *T. cruzi*.

259 3 - Non-random distribution of GP63 gene groups within the *T. cruzi* genome

The genome of *T. cruzi* is composed of two clearly defined compartments, the so-
called disruptive and core compartments (15). These genomic compartments are
recognized for their distinct base composition (% GC), gene content, and likely exhibit
differential regulation and functions within the parasite, alongside unique evolutionary
trajectories (20). The core compartment contains genes conserved between different *T.*
cruzi strains but also largely shares synteny conservation with other trypanosomatids.
On the other hand, the disruptive compartment is composed almost exclusively of
multigene families for surface proteins. While the majority of gene families encoding
surface proteins are situated within the disruptive compartment, GP63 does not follow
this pattern. Our findings reveal that the GP63 gene family is evenly distributed across
both compartments (Figure 2D and Supplementary Figure 4). This contrasts with
Leishmania major, which harbors a single tandem array with six GP63 genes in its
genome (21). Unlike *L. major*, *T. cruzi* shows a wide dispersion of GP63 genes
throughout its genome. Furthermore, the genomic positioning of the various GP63
groups shows a strong correlation with these specific genomic regions. Notably, genes
belonging to groups G1 to G8 are all found in the core compartment, whereas genes
from groups G9, G10 and G11 are present almost exclusively in the disruptive
compartment (Figure 2D and Supplementary Figure 4).

Figure 3: GP63 Gene Environment Composition. This figure illustrates the predominant genes found
adjacent to each GP63 gene, focusing on the three genes upstream and downstream. A) Analyzes the

environment of GP63 genes within the core compartment. B) Analyzes the environment of GP63 genes
within the disruptive compartment. C) Provides examples of GP63 gene environments in both
compartments, with visuals sourced from http://bioinformatica.fcien.edu.uy/cruzi/browser/dm28c.500.php.
For these examples, the contig and starting location of the depicted fragments are specified. Each region
presents a schematic dotplot, showing the organization of the tandem repeats.

We also performed a local gene environment analysis aiming to identify the most
frequent neighbors of GP63 genes. The three most abundant genes were identified at
both, 3' or 5' neighborhood of each GP63 coding sequence. By neighborhood we
understand the three closest neighbors. We found that depending on whether the
GP63 gene is located in the core or disruptive compartment, its gene environment is
quite different. While genes in the core compartment mostly have a sire-like repeat
sequence immediately upstream or downstream (Figure 3A), GP63 genes located in
the disruptive compartment are found together with MASP, RHS and TS genes (Figure
3B and 3C). In addition, many GP63 genes are distributed in tandem repeats, and this
is basically restricted to groups found in the core compartment (Figure 3C). In many
cases, these tandem arrays do not contain exclusively the GP63 sequences but also
the accompanying gene/s, meaning that the repetition unit can be somewhat larger.

The short interspersed repetitive element (SIRE)-like sequences, exclusive to *T. cruzi*,
are prevalent throughout its genome, numbering between 1,500 and 3,000 copies (22).
SIREs are often associated with protein-coding genes and are typically found in the 3'
untranslated regions (UTRs) of mRNAs, frequently contributing to polyadenylation sites
(23). Moreover, SIRE elements are components of VIPER, a distinctive retroelement
which is presumably active in *T. cruzi*. The presence of this type of sequences in the
UTRs of several of the GP63 genes may have implications in the regulation of their
expression. In fact, it is known that elements present in the UTRs influence their gene
expression (24). Furthermore, the observed enrichment of SIRE elements in
association with GP63 genes, especially within the core compartment, suggests their
potential involvement in the gene duplication mechanisms of GP63 genes. Conversely,
within the disruptive compartment, the genomic dynamics of GP63 genes are likely
influenced by the amplification of other multigenic sequences, including mucins, RHS,
trans-sialidases, and MASPs.

4 - Signal and domain recognition in predicted GP63 amino acid sequences

According to the MEROPS database, the peptidases of the M8 family are zinc-
metallopeptidases belonging to the MA clan of metallopeptidases, characterized by an
active site containing a HEXXH motif involved in the coordination of the zinc ion
required for catalytic activity. This clan includes several subfamilies and is known for its
functional diversity, including peptidases involved in extracellular protein degradation,
peptide signal processing, and other critical functions in both pathogens and their
hosts. M8 family proteases are notably prevalent across various protozoan organisms,
with *T. cruzi* boasting the largest collection of such proteins (298).

All identified sequences were found to contain a peptidase M8 domain (Leishmanolysin
family, clan MA(M), InterPro Acc IPR001577). The Metzincin clan of metalloproteinases
possess a consensus sequence that includes not only the two histidines that bind the
zinc ion and the general-base glutamate (HEXXH), but also an extended motif with and
additional strictly conserved glycine and a third zinc-binding histidine located further
along the sequence. Metzincins are also characterized by a conserved methionine-
containing 1,4-β-turn, known as the Met-turn (25). A distinctive aspect of
leishmanolysins is the presence of a large insertion between the glycine and the third

zinc-binding histidine. Our analysis focused on the presence of these conserved elements within the active site of the M8 domain. The zinc-binding site is preserved in most of the protein sequences, with extensive conservation observed beyond the second histidine, indicating a limited scope for variation in these positions, which consistently feature alanine and leucine (Figure 4). Intriguingly, two GP63 groups, G9 and G10, have a mutation in the highly conserved glutamic acid residue within the active site, by leucine and phenylalanine respectively. While the impact of this mutation on protein activity is unclear, it does not appear to influence gene expression. In fact, these genes exhibit RNAseq expression levels that are comparable to or exceed those of the other groups (see Figure 5). Furthermore, the variable number of residues between the conserved glycine in the active site and the third histidine is striking. While most of the groups have similar or exact spacing to the *Leishmania* sequences (62 amino acids), group G3 contains a significantly larger insertion of 150 to 270 residues. These insertions split the M8 domain into two segments, significantly increasing the distance between the third histidine and the preceding two histidines. The significance of this change in protein structure is not known but would merit further study in terms of its possible functional impact. In the case of group G1, this spacing is 71 residues. In addition, a Met-turn characterized by a highly conserved methionine is present in all groups except for this group G1. As illustrated in Figure 4, each group exhibits unique characteristics regarding these elements.

Figure 4: Structure of GP63 Group proteins. This figure provides a schematic overview of each GP63 group, depicting protein length, the location of the M8 domain, and the presence of signal peptides and GPI anchors. On the left side, each group is indicated and the number of sequences in each group is given in brackets. On the right side, the catalytic domain HEXXHXXG, the spacer to the subsequent histidine, and the conservation of the Met-turn are illustrated, along with all group-specific variations highlighted in red.

Mutations at H264, E265 and H268 of the active site of the *L. major* GP63 have been described to completely abrogate catalytic function (18). The presence of the essential active site elements in the majority of *T. cruzi* GP63 sequences suggests that these are proteins active in their protease function. In contrast, the absence of the highly conserved E residue of the active site of the sequences belonging to groups G9 and

G10 of *T. cruzi*, which has been considered essential in the *L. major* protein, suggests
that these two groups of sequences are not functional proteases. However, it should be
noted that these proteins could have other functions, as has been described for the
trans-sialidase family, a family that should be renamed because most of its members
have no enzymatic activity due to a mutation in the active site (26). However, these
proteins retain other important functions by facilitating interaction with host cell
receptors, promoting signaling and cell invasion of trypomastigote forms (27).

The structural analysis of GP63 amino acid sequences revealed a predicted N-terminal
signal sequence across almost all GP63 variants, ranging from 23 to 37 amino acids
depending on the sequence. Only in seven sequences a signal peptide could not be
predicted, due in almost all cases to N-terminal deletions. Of these, four sequences
corresponded to group G11, one to G9, one to G8 and one to G4. Conversely,
predictions for a GPI anchor site yielded inconsistent results. Most sequences from
groups G1, G3, G5, G6, G7, G9, G10, and G11 feature a GPI anchor site, whereas
sequences from groups G2, G4, and G8 appear to lack this site (Figure 4 and
Supplementary Table 1). Furthermore, our examination of putative protein features,
specifically the N-glycosylation sites across the 96 proteins, indicated a variability in the
number of N-glycosylation sites among the groups. Notably, group G9 exhibited the
highest count of glycosylation sites, as detailed in supplementary Figure 5.

The amino- and carboxyl-terminal ends of the GP63 sequences show the greatest
variability between the different members of each group, both in terms of length and
sequence composition. For leishmanolysin, as well as for several other families of
metzincin metalloproteinases, a "cysteine switch" mechanism has been described,
based on the presence of a cysteine residue in the pro-segment that prevents access
of substrates to the active site of the enzyme and thus keeps it inactive (18). With
proteolysis of the pro-segment, the enzyme would become active. A conserved motif
surrounding the cysteine residue has been described, which in the case of GP63 from
Leishmania sp. is HRCIHD or HRCVHD. Using the motif-based sequence analysis
tools of the MEME Suite, we examined the cysteine residue at the amino-terminal ends
of *T. cruzi* GP63 sequences. Our findings reveal that, except for nearly half the
sequences of group G11—where the cysteine is substituted by an arginine—the
cysteine residue remains conserved across all other groups where a similar motif was
detectable, despite significant variations in the adjacent sequence regions (see
Supplementary Table 4). An additional outlier is observed in one sequence from group
G2, characterized by a deletion in this specific region.

5 - Expression profiles of GP63 genes

Previous studies have demonstrated the upregulated transcriptional expression of
certain multigene families, such as TS (28) and MASP (2), during the trypomastigote
stage (a non-replicative, infective stage interacting with the host). In contrast, the
transcriptional profile of GP63 genes remains less defined. In fact, it has been
suggested that while one subset of GP63 genes is minimally expressed, another
exhibits increased expression in trypomastigotes, and a different subset shows
elevated expression in amastigotes (29). However, the reliability of these findings might
be compromised, as these analyses were based on collapsed genomes, potentially
failing to represent all gene copies accurately.

We have analyzed their expression profiles at the mRNA level with previously
generated transcriptomic data from three stages of the parasite (20), observing that

each group shows different expression levels and profiles and that several groups
and/or members show differential expression at the trypomastigote stage compared to
the epimastigote and amastigote stages. As can be seen in Figure 5, while groups G1,
G4, G6 and G10 do not show differential expression profiles, the rest of the groups
contain genes that are significantly expressed more in the trypomastigote stage
compared to the epimastigote and amastigote stages. Remarkably, insertions in the
active site of group G3 sequences appear to not affect their expression, as they are
distinctively upregulated during the trypomastigote stage, suggesting a functionality for
this group. Expression levels were also very different between the groups, with those of
group G2 and group G9 being the most highly expressed genes, reaching levels
between one and two orders of magnitude higher than the rest of the groups (Figure 5
and Supplementary Table 5). These differential expression profiles do not correlate
with the location of genes in the two genomic compartments. Groups showing
differential expression are located in both the core and disruptive compartments.
Moreover, the two groups with the highest expression at the RNA level are one in the
core and the other in the disruptive compartment. This suggests that differential
expression mechanisms are strongly linked to features associated with gene structure,
e.g. possible regulatory sequence elements present in the UTRs.

Figure 5: Expression profiles at the ARNm levels in the different GP63 groups. Each graph shows

the expression profiles of the genes in each GP63 group. On the y-axis, expression levels are expressed
in transcripts per million (tpm). Levels of expression were categorized as low, medium and high, and
coloured differently. On the x-axis are the three stages of *T. cruzi*: Epimastigotes (E), Amastigotes (A) and
Trypomastigotes (T). The black dots in each graph represent each individual gene. Statistical analysis of
differential expression between stages was performed by 2 way ANOVA. *** represents a p-value <0.001

6 - Patterns of divergence in regions embedding (surrounding) GP63 coding sequences is similar to that of proteins

To determine if the sequence identity observed within each GP63 group extends to
untranslated regions (UTRs) or is confined solely to the coding sequence, we
conducted a phylogenetic analysis paralleling the approach used for the proteins. This
involved analyzing 450 nucleotides upstream (5' UTR) and 700 nucleotides
downstream (3' UTR) of each GP63 gene, which represent the average length of the
UTRs present in the GP63 genes. As illustrated in Figure 6, the clustering patterns in
the 5' and 3' UTRs largely mirror those seen in the phylogenetic analysis of the
proteins. This suggests that GP63 genes within each group not only share sequence
identity in their coding regions but also in their UTRs, with the similarity being more
pronounced in the 3' UTR than in the 5' UTR. As previously discussed, tandem
duplication represents a form of gene amplification within this family, with conservation
in these regions indicating that duplication encompasses more than just the coding
sequence. Moreover, the untranslated regions (UTRs), particularly the 3'UTR, are
pivotal in gene expression regulation (30). Hence, the conserved clustering of these
sequences might suggest a differential expression regulation, potentially varying across
developmental stages or in response to different stress conditions. Notably, in the
3'UTR phylogeny, group G4 undergoes a division, with three sequences diverging from
this group (indicated by an arrow in Figure 6). These sequences are the 3'UTRs of
genes C4B63_170g41, C4B63_320g17, and C4B63_32g57, which represent the non-
expressed genes of G4 at any developmental stage (see Figure 5 and Supplementary
Table 5 for reference). Similarly, group G11 bifurcates, with 7 out of the 32 3'UTRs
clustering separately. The genes corresponding to these 3'UTRs show very low
expression levels during the trypomastigote stage, in contrast to the remaining genes,
which exhibit increased expression during this phase.

Figure 6: Phylogenetic Analysis of GP63 UTRs. Phylogenetic analysis of the 5'UTRs (450nt) on the left
and the 3'UTRs (700nt) on the right. Groups are differentiated by color coding, and labels include the
name of the respective gene and group. UTRs that do not cluster with their respective groups are
highlighted with a black arrow. Additionally, nodes are colored based on the genomic compartment to
which they belong.

Together, all these results suggest that this family of proteins with different levels of
mRNA expression may be strongly regulated at the post-transcriptional level by
elements present in the UTRs, as has been described for *Leishmania* genes (16)

7 - Amplification of GP63 genes is partially mirrored in other trypanosomes

The eleven groups described in section 2, most likely reflect functional diversification in
the GP63 family of *T. cruzi*. It is of great interest to determine whether this divergence
is observed in other species of the genus *Trypanosoma*, other trypanosomatids and
even outside the trypanosomatidae family. For this purpose, we conducted a
phylogenetic analysis including all completed GP63 sequences identified in eight
Trypanosoma species, covering the taxonomic diversity of the genus *(Trypanosoma*
cruzi, *T. cruzi marinkellei*, *Trypanosoma rangeli*, *Trypanosoma tehilieri*, *Trypanosoma*
grayi, and three african trypanosomes). We also included the genus *Strigomonas*
(*Blastocrithidia*), two *Leishmania* species and the free living kinetoplastid *Bodo saltans*.
The two phylogenetic trees presented in Figure 7 allow us to highlight different aspects
of this divergence. These two trees differ in that Figure 7A contains just one
representative copy of each group from the large *Trypanosoma* cluster, allowing an
easier identification of the divergence outside the genus. From Figure 7A it is evident
that ten of the eleven groups described above, represent *Trypanosoma* specific
subfamilies, whereas homologues to G1 are observed outside the genus. Interestingly,
members of G1 subfamily are found as single copy genes not only in *T. cruzi* but in all
species of *Trypanosoma* and *Strigomonas*. The gene is also present in *B. saltans* as
multicopy but absent in *Leishmania*. The phylogenetic location of G1, as the earliest
branching lineage, and its widespread presence in kinetoplastids is indicative of its
ancestrality. This, in turn, is reflected in two aspects of its biology that are worth
mentioning. First, it is the only group that lacks the conserved methionine of the Met-
turn motif (see Figure 4) and contains a spacing of 71 residues between the conserved
Glycine in the active site and the third histidine, which is unique to this group. Second,
in *T. cruzi,* the G1 group is transcribed at constant levels throughout the life cycle,
suggesting an absent or weak relationship with the infective phase, a feature that is
compatible with its presence in the non-infective kinetoplastid *B. saltans*. Analysis of
the conserved active site elements in sequences clustered with the *T. cruzi* G1 group
revealed the preservation of all elements, including the characteristic substitution of
methionine with leucine, which distinguishes this group from others. However, the
distance between the active site glycine and the third histidine exhibited some
variability, ranging from 69 to 74 amino acids across the sequences. A notable
deviation was observed in the *T. congolense* sequence, which contains a deletion
resulting in a shorter separation of 58 amino acids between these elements

Another interesting aspect from Figure 7A, is the pattern of amplification and
divergence of GP63 in both *L. major* and *L. braziliensis*. The interest in understanding
the divergence pattern in these two species and its comparison with *T. cruzi* is
fundamentally due to the fact that this is a classical and extensively studied *Leishmania*
protein (in fact is called leishmanolysin). Noteworthy, there are only three relatively

divergent gene copies, one of which forms a cluster that has 9 copies in *L. braziliensis*
and 4 in *L. major* (31). Similar to what is observed in *T. cruzi*, the degree of amino acid
divergence among copies within the same cluster is very modest, this being true for
both *Leishmania* species included in this study. Besides, similarities are higher within,
than among species. This may indicate that the amplifications emerged independently
(i.e. are species specific), or that there is a high rate of gene conversion. We note that
with this comparatively low copy number and differentiation within cluster in
Leishmania, this protease is able to exert different functions, facilitating adhesion and
invasion and cleaving or degrading several host cell proteins that play important roles
in the cellular response, such as transcription factors and components of signaling
cascades (9).

Figure 7: Global Phylogenetic Analysis of GP63 Genes Across Kinetoplastids. A) This panel shows a
phylogenetic analysis utilizing the Maximum Likelihood method based on the JTT model with gamma
distribution and 100 bootstrap pseudoreplicates. It focuses on the entire GP63 amino acid sequences from
representative genes of *T. cruzi, T. cruzi marinkellei, T. rangeli, T. grayi, T. theileri, T. vivax, T.*
congolense, T. brucei, Leishmania major, L. braziliensis, Strigomonas culicis, and *Bodo saltans*. Bootstrap
support values above 50 are indicated at the respective nodes. B) Presents a phylogenetic analysis of the
M8 domain using an enriched dataset of 282 sequences from a diverse array of *Trypanosoma* and
Leishmania species, including the free-living kinetoplastid *Bodo saltans*, highlighting the evolutionary
relationships within this gene family across kinetoplastids.

Figure 7B is basically an enlargement of part A that incorporates the intragroup
variability within the eight species of *Trypanosoma* genus, thus including all sequences

with a complete M8 domain (378). As previously noted, ten of the eleven *T. cruzi*
groups, defined in section 2, form part of a large clade. A relevant observation is that
members of these ten groups are also found in other *Trypanosoma* species, including
African ones. There are some exceptions though: group G3 is absent in African
trypanosomes, and groups G4 and G9 are not found in *T. grayi* and *T. theilieri*. The
copy number of each group exhibits considerable variation across different species,
with group G11 notably expanded in *T. cruzi*. *T. cruzi* exhibits very high intra-cluster
similarity, in contrast to what is observed in other species of *Trypanosoma* genus (and
Bodo). This pattern of *T. cruzi* is to some extent similar to that of *Leishmania.* Besides
the groups already described in *T. cruzi*, additional large clusters are present in other
species with no representation in *T. cruzi*. One of such clusters, designated LTGC1
(Large Theiliri Gp63 Cluster 1), includes over 35 copies in *T. theilieri*, with a few distant
relatives in *T. grayi*. A second notable expansion in *T. theilieri*, termed LTGC2,
comprises approximately 20 copies. Genes within these non-*T.cruzi* GP63 groups also
possess a complete M8 domain and signal peptide, suggesting their likely functionality.

557
558 **CONCLUSIONS**
559

In this study we address the amplification and diversification of GP63 genes in the
genome of *T. cruzi*, using as an initial reference our previous genomic studies on the
Dm28c strain. Our analysis identified the existence of eleven groups, each
distinguished by specific characteristics such as sequence length, unique features
within the conserved elements of the M8 motif and different expression profiles.
Notably, these groups exhibit variations in the presence of signals in the N- and C-
terminal regions as well as in the number of sites for N-glycosylation. While there is
remarkable homogeneity within each group, substantial divergence is observed when
comparing sequences across different groups.

Furthermore, we found a strong correlation between these GP63 groups and the core
and disruptive genomic compartments of the *T. cruzi*. Multi-copy tandem arrays of
GP63 genes are prevalent in groups situated in the core compartment, whereas those
in the disruptive compartment are typically flanked by genes from other multigene
families, such as mucins, MASPs, and trans-sialidases.

This study has also allowed us to elucidate the evolutionary expansion of the GP63
gene family in *Trypanosoma.* Phylogenetic analysis provided insights into the
evolutionary expansion of the GP63 gene family in *T. cruzi*, showing that this family has
followed different evolutionary paths, not only within the genus *Trypanosoma* but also
in other species. Ten of the 11 groups characterized in this study are shared only within
the genus *Trypanosoma* including African species, while only group G1 is also present
in other trypanosomatidae species and nonparasitic kinetoplastids, indicating that
group G1 represents the earliest branching lineage. These studies reveal on the one
hand, that the expansion of the remaining GP63 groups genes occurred prior to the
separation of *Trypanosoma* species. On the other hand, the expansion of GP63
observed in species of the genus *Leishmania* is independent of that of *Trypanosoma*,
and vastly more modest.

METHODS

GP63 Metalloprotease Sequences

We downloaded 378 sequences annotated as "surface_protease_GP63" within the *T. cruzi* Dm28c genome, from the database http://bioinformatica.fcien.edu.uy/cruzi/. Among them, 96 were identified as genes and 282 as pseudogenes.

To validate and refine these annotations, we focused on sequences that exceeded 460 amino acids in length and possessed a complete M8 peptidase domain, indicative of GP63 genes. This evaluation was performed using the InterPro predictor (32) and CDVist (33), a web server designed to maximize domain coverage in multidomain protein sequences, which enabled the identification of domains within the GP63 protein sequences. N-glycosilation site prediction were done using the NetNGlyc - 1.0 server that predicts N-Glycosylation sites proteins examining the sequence context of N-X-S/T sequences (https://services.healthtech.dtu.dk/services/NetNGlyc-1.0/).

The presence and location of signal peptide cleavage sites was predicted using SignalP 6.0 server (34) selecting Eukaria as an organism. The presence of a glycosylphosphatidylinositol (GPI) anchoring was predicted using two different web-servers: PredGPI (35), and NetGPI 1.1 (https://services.healthtech.dtu.dk/services/NetGPI-1.1/).

Reannotation of GP63 Pseudogenes

A homology search was performed using blastn, with functional GP63 sequences as a reference database. Additionally, the M8 domain was searched as previously described. Sequences located in close proximity or at short distances were meticulously analyzed using both blastn and blastp against GP63 genes, with YASS (36) facilitating dotplot visualizations. This approach allowed us to identify open reading frames (ORFs) as parts of genes that had undergone pseudogenization. Accordingly, we re-annotated these sequences, extending them from the first nucleotide (nt) of the initial sequence to the last nt of the final constituent sequence, marking the internal ORFs as derived fragments. The "new" parental pseudogene was re-annotated with an ID ending with a letter p.

Cluster analysis based on amino acid identity

Identity matrix of the GP63 sequences was generated by the pairwise sequence alignment tool Clustal Omega (37) (https://www.ebi.ac.uk/Tools/msa/clustalo/). Visualization of the clustering was achieved with Gephi, where each edge's color tone represents the degree of identity between sequences. An edge weight threshold of 65% was applied for visualization. Additionally, a condensed identity matrix, summarizing average intra-group and inter-group identity, was derived from mean group identity analyses. This condensed matrix was visualized using R.

Gene environment analysis

We analyzed the genomic context of each GP63 gene by identifying the three upstream (5') and three downstream (3') genes to assess the gene composition surrounding GP63 loci. For GP63 genes residing in both core and disruptive compartments, we determined the most prevalent gene types, utilizing the annotations of these neighboring genes with a custom script in Perl. The results were plotted using R.

Expression profile of GP63 genes

Expression profiling for each gene was derived from the work of (20). For each gene,

Transcript Per Million (TPM) values were calculated across replicates and
developmental stages (amastigote, epimastigote, and trypomastigote), with
differentially expressed genes listed in Supplementary Table 5. Rather than using
arbitrary TPM thresholds, expression levels were categorized based on quartile
distribution of TPM values within each stage: genes in the lowest quartile (<25%) were
classified as *low*, those between the 25th and 75th percentiles were deemed *medium*,
and genes in the highest quartile (>75%) were considered *high*.

Phylogenetic analyses
The peptide sequences of the 96 GP63 genes were aligned using MAFFT (38),
employing the '-einsi' option for enhanced accuracy in alignments involving sequences
with multiple conserved domains and gaps. Subsequent trimming was executed
utilizing the 'gappyout' option in TrimAl to remove poorly aligned regions and gaps.
Phylogenetic analysis was conducted using PhyML (39) with the following parameters:
'phyml -i 2024_M8.einsi.gappy.phy -b -1 -d aa'. Visualization of the resulting
phylogenetic tree was created using iTOL (40).

To enrich the taxonomic coverage of the dataset, 282 sequences from a variety of
Trypanosoma and *Leishmania* species, and the free living kinetoplastid *Bodo saltans*,
including *T. cruzi marinkellei* (17), *T. theileri* (103), *T. grayi* (33), *T. rangeli* (55), *T.*
congolense (11), *T. vivax* (12), *T. brucei* (23), *L. major* (6), *L. braziliensis* (10),
Strigomonas culicis (3), and *Bodo saltans* (9), were incorporated. The M8 domain was
identified in these sequences, and both alignment and phylogeny were specifically
performed on these extracted domains, as previously mentioned.

From this expanded set of 378 sequences (282 plus 96 from *T. cruzi*), we selected
representative sequences for each group of *T. cruzi* GP63 and all other clusters,
totaling 103 sequences. The alignment of the selected sequences was done using a
MAFFT (38), retaining all positions containing information in at least 80% of the
sequences. A phylogeny was then constructed using the Maximum likelihood method
using JTT model with gamma distribution and 100 bootstrap pseudoreplicates,
providing insights into the evolutionary relationships and diversity within the GP63 gene
family across these protozoan organisms.

UTR phylogenies were performed using 450 and 700 nucleotide positions upstream
and downstream of 5' and 3' ends, respectively. The sequences were retrieved from
tritrypdb.org (41), aligned with MAFFT (38) using -auto option, and the phylogenetic
tree was constructed with IQ-TREE (42) with default parameters.

SUPPLEMENTARY FIGURES

Supplementary Figure 1. Reassignment of pseudogene fragments, presenting two examples at the top. In these instances, alignments were conducted using the YASS tool, with the C4B63_1g558 gene serving as the reference. Below, the figure provides a visualization of the pairwise local alignments for four pseudogene pairs, demonstrating the consistent pattern observed in the examples above.

Supplementary Figure 2. Average length of the amino acid sequences for each GP63 group, including the standard deviation.

Supplementary Figure 3. Phylogenetic analysis of the 96 GP63 amino acid sequences from *T. cruzi* Dm28c. Groups are differentiated by color coding in labels. Nodes are colored based on the genomic compartment, core in green, disruptive in purple.

Supplementary Figure 4. Genomic distribution of GP63 gene sequences across *T. cruzi* groups. Green color corresponds to groups present in the core compartment, purple to groups in the disruptive compartment.

689 **Supplementary Figure 5.** Average number of predicted N-glycosylation sites in each group of GP63
690 sequences.

Funding information

693 This work was funded by Comisión Sectorial de Investigación Científica (CSIC)
694 Universidad de la República (project I+D 22520220100578UD) and Programa de
695 Desarrollo de las Ciencias Básicas (PEDECIBA).

696
Conflicts of interest
698 The authors declare that there are no conflicts of interest.

699
Consent for publication
701 All authors have approved the manuscript and agree with its submission.

702
703
REFERENCES

1.   Buscaglia CA, Campo VA, Frasch ACC, Di Noia JM. Trypanosoma cruzi surface
mucins: host-dependent coat diversity. Nat Rev Microbiol. 2006 Mar;4(3):229–36.

2.   Bartholomeu DC, Cerqueira GC, Leão ACA, daRocha WD, Pais FS, Macedo C, et
al. Genomic organization and expression profile of the mucin-associated surface
protein (masp) family of the human pathogen Trypanosoma cruzi. Nucleic Acids
Res. 2009 Jun;37(10):3407–17.

3.   Frevert U, Schenkman S, Nussenzweig V. Stage-specific expression and
intracellular shedding of the cell surface trans-sialidase of Trypanosoma cruzi.
Infect Immun. 1992 Jun;60(6):2349–60.

4.   Ma L, Chen K, Meng Q, Liu Q, Tang P, Hu S, et al. An evolutionary analysis of
trypanosomatid GP63 proteases. Parasitol Res. 2011 Oct;109(4):1075–84.

5.   Jackson AP, Otto TD, Aslett M, Armstrong SD, Bringaud F, Schlacht A, et al.
Kinetoplastid Phylogenomics Reveals the Evolutionary Innovations Associated
with the Origins of Parasitism. Curr Biol. 2016 Jan 25;26(2):161–72.

6.   Castro Neto AL, Brito ANALM, Rezende AM, Magalhães FB, de Melo Neto OP. In

silico characterization of multiple genes encoding the GP63 virulence protein from
Leishmania braziliensis: identification of sources of variation and putative roles in
immune evasion. BMC Genomics. 2019 Feb 7;20(1):118.

7.   Russell DG, Wilhelm H. The involvement of the major surface glycoprotein (gp63)
of Leishmania promastigotes in attachment to macrophages. J Immunol. 1986 Apr
1;136(7):2613–20.

8.   Yao C. Major surface protease of trypanosomatids: one size fits all? Infect Immun.
2010 Jan;78(1):22–31.

9.   Olivier M, Atayde VD, Isnard A, Hassani K, Shio MT. Leishmania virulence factors:
focus on the metalloprotease GP63. Microbes Infect. 2012 Dec;14(15):1377–89.

10.   Antonia AL, Gibbs KD, Trahair ED, Pittman KJ, Martin AT, Schott BH, et al.
Pathogen Evasion of Chemokine Response Through Suppression of CXCL10.
Front Cell Infect Microbiol. 2019 Aug 7;9:280.

11.   Cuevas IC, Cazzulo JJ, Sánchez DO. gp63 homologues in Trypanosoma cruzi:
surface antigens with metalloprotease activity and a possible role in host cell
infection. Infect Immun. 2003 Oct;71(10):5739–49.

12.   Grandgenett PM, Coughlin BC, Kirchhoff LV, Donelson JE. Differential expression
of GP63 genes in Trypanosoma cruzi. Mol Biochem Parasitol. 2000
Oct;110(2):409–15.

13.   Lowndes CM, Bonaldo MC, Thomaz N, Goldenberg S. Heterogeneity of
metalloprotease expression in Trypanosoma cruzi. Parasitology. 1996 Apr;112 (Pt
4):393–9.

14.   El-Sayed NM, Myler PJ, Bartholomeu DC, Nilsson D, Aggarwal G, Tran AN, et al.
The genome sequence of Trypanosoma cruzi, etiologic agent of Chagas disease.
Science. 2005 Jul 15;309(5733):409–15.

15.   Berná L, Rodriguez M, Chiribao ML, Parodi-Talice A, Pita S, Rijo G, et al.
Expanding an expanded genome: long-read sequencing of Trypanosoma cruzi.
Microb Genom [Internet]. 2018 May;4(5). Available from:
http://dx.doi.org/10.1099/mgen.0.000177

16.   Yao C, Donelson JE, Wilson ME. The major surface protease (MSP or GP63) of
Leishmania sp. Biosynthesis, regulation of expression, and function. Mol Biochem
Parasitol. 2003 Nov;132(1):1–16.

17.   McGwire BS, Chang KP. Posttranslational regulation of a Leishmania HEXXH
metalloprotease (gp63). The effects of site-specific mutagenesis of catalytic, zinc
binding, N-glycosylation, and glycosyl phosphatidylinositol addition sites on N-
terminal end cleavage, intracellular stability, and extracellular exit. J Biol Chem.
1996 Apr 5;271(14):7903–9.

18.   Macdonald MH, Morrison CJ, McMaster WR. Analysis of the active site and
activation mechanism of the Leishmania surface metalloproteinase GP63. Biochim
Biophys Acta. 1995 Dec 6;1253(2):199–207.

19.   Abrahim M, Machado E, Alvarez-Valín F, de Miranda AB, Catanho M. Uncovering
Pseudogenes and Intergenic Protein-coding Sequences in TriTryps' Genomes.
Genome Biol Evol [Internet]. 2022 Oct 7;14(10). Available from:
http://dx.doi.org/10.1093/gbe/evac142

20. Díaz-Viraqué F, Chiribao ML, Libisch MG, Robello C. Genome-wide chromatin
interaction map for Trypanosoma cruzi. Nat Microbiol. 2023 Nov;8(11):2103–14.

21. Button LL, Russell DG, Klein HL, Medina-Acosta E, Karess RE, McMaster WR.
Genes encoding the major surface glycoprotein in Leishmania are tandemly linked
at a single chromosomal locus and are constitutively transcribed. Mol Biochem
Parasitol. 1989 Jan 15;32(2-3):271–83.

22. Vázquez M, Lorenzi H, Schijman AG, Ben-Dov C, Levin MJ. Analysis of the
distribution of SIRE in the nuclear genome of Trypanosoma cruzi. Gene. 1999 Nov
1;239(2):207–16.

23. Vazquez M, Ben-Dov C, Lorenzi H, Moore T, Schijman A, Levin MJ. The short
interspersed repetitive element of Trypanosoma cruzi, SIRE, is part of VIPER, an
unusual retroelement related to long terminal repeat retrotransposons. Proc Natl
Acad Sci U S A. 2000 Feb 29;97(5):2128–33.

24. De Gaudenzi JG, Noé G, Campo VA, Frasch AC, Cassola A. Gene expression
regulation in trypanosomatids. Essays Biochem. 2011;51:31–46.

25. Gomis-Rüth FX. Structural aspects of the metzincin clan of
metalloendopeptidases. Mol Biotechnol. 2003 Jun;24(2):157–202.

26. Cremona ML, Sánchez DO, Frasch AC, Campetella O. A single tyrosine
differentiates active and inactive Trypanosoma cruzi trans-sialidases. Gene. 1995
Jul 4;160(1):123–8.

27. Pascuale CA, Burgos JM, Postan M, Lantos AB, Bertelli A, Campetella O, et al.
Inactive trans-Sialidase Expression in iTS-null Trypanosoma cruzi Generates
Virulent Trypomastigotes. Front Cell Infect Microbiol. 2017 Oct 4;7:430.

28. Jäger AV, Muiá RP, Campetella O. Stage-specific expression of Trypanosoma
cruzi trans-sialidase involves highly conserved 3' untranslated regions. FEMS
Microbiol Lett. 2008 Jun;283(2):182–8.

29. Berná L, Chiribao ML, Greif G, Rodriguez M, Alvarez-Valin F, Robello C.
Transcriptomic analysis reveals metabolic switches and surface remodeling as key
processes for stage transition in Trypanosoma cruzi. PeerJ. 2017 Mar 8;5:e3017.

30. Nozaki T, Cross GA. Effects of 3' untranslated and intergenic regions on gene
expression in Trypanosoma cruzi. Mol Biochem Parasitol. 1995 Dec;75(1):55–67.

31. Alvarez-Valin F, Tort JF, Bernardi G. Nonrandom spatial distribution of
synonymous substitutions in the GP63 gene from Leishmania. Genetics. 2000
Aug;155(4):1683–92.

32. Paysan-Lafosse T, Blum M, Chuguransky S, Grego T, Pinto BL, Salazar GA, et al.
InterPro in 2022. Nucleic Acids Res. 2023 Jan 6;51(D1):D418–27.

33. Adebali O, Ortega DR, Zhulin IB. CDvist: a webserver for identification and
visualization of conserved domains in protein sequences. Bioinformatics. 2015
May 1;31(9):1475–7.

34. Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos
KD, et al. SignalP 6.0 predicts all five types of signal peptides using protein
language models. Nat Biotechnol. 2022 Jul;40(7):1023–5.

35. Pierleoni A, Martelli PL, Casadio R. PredGPI: a GPI-anchor predictor. BMC
Bioinformatics. 2008 Sep 23;9:392.

36. Noé L, Kucherov G. YASS: enhancing the sensitivity of DNA similarity search.
Nucleic Acids Res. 2005 Jul 1;33(Web Server issue):W540–3.

37. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable
generation of high-quality protein multiple sequence alignments using Clustal
Omega. Mol Syst Biol. 2011 Oct 11;7:539.

38. Katoh K, Asimenos G, Toh H. Multiple alignment of DNA sequences with MAFFT.
Methods Mol Biol. 2009;537:39–64.

39. Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New
algorithms and methods to estimate maximum-likelihood phylogenies: assessing
the performance of PhyML 3.0. Syst Biol. 2010 May;59(3):307–21.

40. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic
tree display and annotation. Nucleic Acids Res. 2021 Jul 2;49(W1):W293–6.

41. Amos B, Aurrecoechea C, Barba M, Barreto A, Basenko EY, Bażant W, et al.
VEuPathDB: the eukaryotic pathogen, vector and host bioinformatics resource
center. Nucleic Acids Res. 2022 Jan 7;50(D1):D898–911.

42. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler
A, et al. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference
in the Genomic Era. Mol Biol Evol. 2020 May 1;37(5):1530–4.