POMC neurons control fertility through differential signaling of MC4R in Kisspeptin neurons

Authors:

Rajae Talbi[1,2*], Todd L. Stincic[4,8*], Kaitlin Ferrari[2], Choi Ji Hae[2], Karol Walec[2], Elizabeth Medve[2], Achi Gerutshang[2], Silvia León[1,2], Elizabeth A. McCarthy[1,2], Oline K. Rønnekleiv[3,4], Martin J. Kelly[3,4,6] and Víctor M. Navarro[1,2,5,6]

[1]Harvard Medical School, Boston, MA, USA; [2]Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA; [3]Department of Chemical Physiology and Biochemistry, Oregon Health & Science University, Portland, OR, USA; [4]Division of Neuroscience, Oregon National Primate Research Center, Beaverton, OR, USA; [5]Harvard Program in Neuroscience, Boston, MA, USA

*These authors contributed equally
[6] Corresponding authors
[8]Present address: Department of Biology, Appalachian State University, Boone, NC 28608.

Correspondence:

Victor M. Navarro, PhD, Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, 221 Longwood Avenue, Room 219, Boston, Massachusetts 02115. E-mail: vnavarro@bwh.harvard.edu.

Martin J. Kelly, PhD, Department of Chemical Physiology and Biochemistry, Oregon Health & Science University, 3181 SW Sam Jackson Park Road, Portland, OR 97239. E-mail: kellym@ohsu.edu .

Keywords:

Kiss1 neurons, POMC neurons, MC4R, Fertility, Ovulation, Obesity, Estradiol.

Abstract

Inactivating mutations in the melanocortin 4 receptor (*MC4R*) gene cause monogenic obesity.

Interestingly, female patients also display various degrees of reproductive disorders, in line with

the subfertile phenotype of MC4RKO female mice. However, the cellular mechanisms by which

MC4R regulates reproduction are unknown. Kiss1 neurons directly stimulate gonadotropin-

releasing hormone (GnRH) release through two distinct populations; the Kiss1[ARH] neurons,

controlling GnRH pulses, and the sexually dimorphic Kiss1[AVPV/PeN] neurons controlling the

preovulatory LH surge. Here, we show that *Mc4r* expressed in Kiss1 neurons regulates fertility in

females. *In vivo*, deletion of *Mc4r* from Kiss1 neurons in female mice replicates the reproductive

impairments of MC4RKO mice without inducing obesity. Conversely, reinsertion of *Mc4r* in Kiss1

neurons of MC4R null mice restores estrous cyclicity and LH pulsatility without reducing their

obese phenotype. *In vitro*, we dissect the specific action of MC4R on Kiss1[ARH] vs Kiss1[AVPV/PeN]

neurons and show that MC4R activation excites Kiss1[ARH] neurons through direct synaptic actions.

In contrast, Kiss1[AVPV/PeN] neurons are normally inhibited by MC4R activation except under

elevated estradiol levels, thus facilitating the activation of Kiss1[AVPV/PeN] neurons to induce the LH

surge driving ovulation in females. Our findings demonstrate that POMC[ARH] neurons acting

through MC4R, directly regulate reproductive function in females by stimulating the "pulse

generator" activity of Kiss1[ARH] neurons and restricting the activation of Kiss1[AVPV/PeN] neurons to

the time of the estradiol-dependent LH surge, and thus unveil a novel pathway of the metabolic

regulation of fertility by the melanocortin system.

Introduction

Obesity rates have skyrocketed in Western societies in the last decades, resulting in an alarming rise in comorbidities that place a significant burden on healthcare systems. The increase in obesity correlates with a decrease in fertility rates, leading to conception challenges that are experienced by approximately 15% of couples in the United States currently (1).

The melanocortin 4 receptor (MC4R) binds i) α-melanocyte stimulating hormone (αMSH), an agonist product of the pro-opiomelanocortin (*Pomc*) gene, and ii) the inverse agonist, agouti-related peptide (AgRP), to regulate food intake and energy expenditure (2, 3). While the role of MC4R on food intake is largely mediated by neurons located in the paraventricular nucleus of the hypothalamus (PVN) (4), its expression in the brain is widespread (5) with the specific function of the different MC4R-expressing neurons in areas beyond the PVN remaining to be fully explored. Inactivating mutations in *MC4R* are a leading cause of monogenic obesity in humans (6) and cause excessive obesity and hyperphagia in mice (7). Scant evidence in humans shows an association between *MC4R* mutations and higher incidence of hypogonadotropic hypogonadism (HH) (8), alterations in the timing of puberty onset (9), and polycystic ovary syndrome (PCOS) (10); however, conflicting reports exist (6). Therefore, the role of MC4R signaling in reproductive function in humans remains controversial despite the clear association found in mice. Supporting this role of MC4R, *Mc4r* null mice display an array of reproductive abnormalities that largely affects females, characterized by irregular estrous cycles, disrupted luteinizing hormone (LH) secretion, reduced corpora lutea and reduced fertility (11-13). Further evidence from mice demonstrates that MC4R agonists robustly increase LH release in a kisspeptin-dependent manner (14).

Kisspeptin is the most potent gonadotropin releasing hormone (GnRH) secretagogue known to date and it is mainly produced in two distinct neuronal populations. Kiss1 neurons of the arcuate nucleus of the hypothalamus (Kiss1[ARH]), present in both sexes, control the pulsatile (tonic) release of GnRH, and sex steroids attenuate their release of kisspeptin. Kiss1 neurons of the

anteroventral periventricular continuum area (Kiss1[AVPV/PeN]), are predominantly present in

females, and are responsible for generating the preovulatory GnRH/LH surge essential for

ovulation (15). Despite these critical roles of both populations of Kiss1 neurons for reproduction,

the underlying mechanisms that determine how each Kiss1 population responds differently to sex

steroids to regulate the GnRH tonic vs surge release remains unresolved.

In rodents, compelling evidence indicates a close interaction between Kiss1 neurons and the

melanocortin system: 1) fibers from POMC neurons in the arcuate nucleus (POMC[ARH]) project to

and juxtapose Kiss1[ARH] neurons (14); 2) the melanocortin signaling through MC4R contributes to

the permissive role of leptin on puberty onset (14, 16, 17); and 3) MC4R expression on both

Kiss1[ARH] (18-20) and Kiss1[AVPV/PeN] (18, 21). Altogether, this evidence suggests a clear role for

MC4R in Kiss1 neurons, in the control of reproductive maturation and fertility.

In this study, we investigated the contribution of MC4R signaling versus obesity *per se* in the

etiology of the reproductive impairments observed in *Mc4r* null mice, which could explain similar

impairments observed in humans. Using genetic mouse models with specific deletion or

reinsertion of *Mc4r* in Kiss1 neurons, we show that MC4R action in Kiss1 neurons is necessary

for normal reproductive function in female mice. Whole cell voltage clamp recordings evidenced

an excitatory action of MC4R on Kiss1[ARH] neurons and an estradiol-dependent inhibitory action

on Kiss1[AVPV/PeN] neurons, with important implications for the timing of the preovulatory LH surge.

Results

Mc4r **within Kiss1 neurons determines the timing of puberty onset in females.**

The expression of *Mc4r* within Kiss1[ARH] and Kiss1[AVPV/PeN] neurons has already been described

elsewhere (18-21), suggesting a role for MC4R in the regulation of fertility. To further investigate

the specific role of MC4R in Kiss1 neurons, we generated a Kiss1-specific MC4R knockout mouse

model (Kiss1-MC4RKO). The specific deletion of *Mc4r* from Kiss1 neurons, as well as the

absence of global recombination, were confirmed through RNAscope. While Kiss1 neurons of

Kiss1-MC4RKO mice lack *Mc4r* transcript compared to their control littermates (**Figure S1A**),

Mc4r was detectable in the PVN of all mice, which is a major site of *Mc4r* expression in the brain

in the regulation of metabolism, therefore supporting the specific deletion of *Mc4r* only within Kiss1

neurons (**Figure S1B**).

Puberty onset was assessed daily from weaning age through the monitoring of vaginal opening

(VO) and first estrus (FE). Kiss1-MC4RKO females showed a significant advancement in the age

of VO (P=0.0150, Kiss1-MC4RKO: 24.67± 0.39 vs controls: 26.32±0.53) and FE (P=0.0341,

Kiss1-MC4RKO: 32.63±1.4 vs controls: 37.84±1.90) compared to their littermate controls (**Figure**

1A, B). Body weight of Kiss1-MC4RKO females at the age of puberty onset was similar between

groups (P=0.2596, Kiss1-MC4RKO: 12.41±0.30 vs controls: 12.89±0.26) (**Figure 1C**). These data

suggest that melanocortin signaling on Kiss1 neurons participates in the timing of puberty onset

in females. Since the role of the melanocortin system on puberty onset is largely unexplored, we

evaluated the expression profile of the main components of this system in the hypothalami of WT

female mice at postnatal days (PND) 10, 15, 22 and 30. Interestingly, expressions of *Agrp*, *Mc3r*

and *Mc4r* were significantly lower at the time of puberty onset (PND30) compared to earlier

developmental ages (**Figure 1D**). This supports our *in vivo* data and suggests that a decrease in

the hypothalamic melanocortin tone drives puberty onset, in line with the advancement in the age

of VO and FE observed in Kiss1-MC4RKO female mice (**Figure 1A, B**).

MC4R in Kiss1 neurons is required for female reproduction.

Because Kiss1-MC4RKO females showed altered puberty onset, we further investigated their

reproductive phenotype. Interestingly, Kiss1-MC4RKO female mice displayed normal BW

throughout the time of the study (up to PND150) **(Figure 2A)**; however, they presented with

irregular estrous cycles with predominantly more time spent in diestrus (P=0.0001) and less time

spent in estrus (P<0.0001) than their control littermates (**Figure 2B, C**). The assessment of the

pulsatile release of LH every 10 min over 180 min revealed no change in the total number of

pulses between groups but a significant increase in basal LH release (P=0.04) in Kiss1-MC4RKO

female mice (0.42±0.03) compared to controls (0.34±0.02) (**Figure 2D-H**). The analysis of the

gene expression of the "KNDy" systems in the ARH, which control the GnRH pulse generator

(15), revealed normal expression levels of *Kiss1*, *Tac2* and *Tacr3*, but significantly lower

expression of *Pdyn* (P=0.0067, Kiss1-MC4RKO: 0.78±0.038 vs controls: 1.00±0.019) (**Figure 2I-**

L). The conserved expression of *Kiss1*, *Tac2* and *Tacr3* correlates with the preserved LH pulse

frequency and amplitude in Kiss1-MC4RKO mice, while the lower inhibitory tone of dynorphin

(**Figure 2I**) correlates with the higher basal LH levels (**Figure 2E**). However, the mRNA

expression of these genes, alone, does not necessarily reflect changes of their activities. To

assess the contribution of MC4R signaling in Kiss1 neurons to the induction of ovulation through

the preovulatory LH surge, Kiss1-MC4RKO females and control littermates were submitted to an

LH surge induction protocol that showed the LH surge was significantly blunted in the Kiss1-

MC4RKO females (P=0.0091), while the protocol clearly evoked the expected afternoon rise of

LH in control mice (**Figure 2M**). In line with these findings, the ovaries of Kiss1-MC4RKO females

displayed fewer corpora lutea, markers of recent ovulation (P=0.0054, Kiss1-MC4RKO: 0.80±0.37

vs controls: 2.80±0.37), in addition to increased cystic follicles (P=0.0337, Kiss1-MC4RKO:

1.60±0.40 vs controls: 0.40±0.24) (**Figure 2N-P**), which correlate with decreased fertility as

observed by the extended time to deliver pups (*i.e.* longer time to get pregnant) (P=0.0030, Kiss1-

MC4RKO: 30.30±4.60 vs controls: 20.88±0.22), and fewer pups per litter (P=0.0095, Kiss1-

MC4RKO: 6.66±0.55 vs controls: 8.44±0.24) (**Figure 2S, T**).

The increase in serum LH levels and decreased ovulation observed in the Kiss1-MC4RKO

females is reminiscent of polycystic ovary syndrome (PCOS) mouse models (22, 23). Thus, we

investigated whether Kiss1-MC4RKO females display a PCOS-like phenotype. We analyzed

circulating levels of testosterone (T) and anti-müllerian hormone (AMH), which are frequently

elevated in PCOS models (24). The Kiss1-MC4RKO females expressed normal T and AMH levels

compared to control mice in diestrus (**Figure 2*Q, R***). Thus, we can exclude a PCOS-like

reproductive phenotype mediated by the lack of melanocortin signaling on Kiss1 neurons.

Re-insertion of MC4R in Kiss1 neurons of MC4RKO mice improves reproductive function.

Kiss1-MC4RKO females displayed reproductive abnormalities resembling those described in

MC4RKO females (11-13). Thus, we hypothesized that the reproductive defects described for the

MC4RKO mice would be, at least in part, due to the absence of MC4R signaling in Kiss1 neurons.

To further investigate this hypothesis, we generated mice that do not express *Mc4r* anywhere

(MC4R-LoxTB, *i.e.* MC4RKO) or that express *Mc4r* only in Kiss1 neurons (*Kiss1-cre:MC4R-*

loxTB). The specific re-insertion of *Mc4r* within Kiss1 neurons in the *Kiss1-cre:MC4R-loxTB* mice

was confirmed through RNAscope (**Figure S2*A*).** *Mc4r* expression was not detected in the PVN

of these mice (**Figure S2*B***), and it was only detected in Kiss1 neurons in the *Kiss1-cre:MC4R-*

loxTB mice. Puberty onset was assessed daily from weaning age through the monitoring of

vaginal opening (VO) and first estrus (FE). MC4R-LoxTB and *Kiss1-cre:MC4R-loxTB* female mice

displayed normal timing of puberty onset compared to their control littermates, as assessed by

VO (P=0.104, MC4R-LoxTB: 30.86±1.90 vs *Kiss1-cre:MC4R-loxTB: 30.09*±1.00 vs controls:

27.57±0.81), and FE (P=0.8472, MC4R-LoxTB: 35.14±2.46 vs *Kiss1-cre:MC4R-loxTB:*

34.55±1.64 vs controls: 35.80±0.84) (**Figure 3*A***), despite displaying significantly higher body

weight at the time of puberty onset (P=0.0001, MC4R-LoxTB: 15.49±0.50 vs *Kiss1-cre:MC4R-*

loxTB: 15.54±0.40 vs Controls: 13.29±0.30) (**Figure 3*B***). As expected, MC4R-LoxTB females

(MC4RKO) displayed increased body weight (**Figure 3C**) and an array of reproductive

impairments that included: irregular estrous cycles with significantly more days in diestrus

(P=0.0016) and fewer days in estrus (P=0.0317) compared to controls (**Figure 3*D, E***),

significantly higher serum LH levels characterized by higher basal (P=0.0882, MC4R-LoxTB:

0.48±0.02 vs Controls: 0.37±0.06) and amplitude levels per LH pulse (P=0.0102, MC4R-LoxTB:

0.730±0.009 vs Controls: 0.58±0.07) (**Figure 3*F-J***), and fewer corporal lutea (P=0.0366, MC4R-

LoxTB: 0.80±0.80 vs Controls: 4.20±1.15) (**Figure 3*K, L***), recapitulating the same reproductive

phenotype observed in Kiss1-MC4RKO mice despite the differences in BW. Re-introduction of

MC4R into Kiss1 neurons in *Kiss1-cre:MC4R-loxTB* mice completely recovered estrous cyclicity,

which was similar to controls (diestrus: P=0.1932, proestrus: P=0.8262, estrus: P=0.0547,

compared to controls) (**Figure 3*D, E***) despite *Kiss1-cre:MC4R-loxTB* mice showing the same

degree of obesity as MC4R-LoxTB mice (**Figure 3*C***), indicating that obesity *per se* was not

mediating the irregular estrous cycles in MC4RKO mice. As indicated above, MC4R-LoxTB mice

display higher overall circulating LH levels, and this feature was also recovered by the re-insertion

of MC4R in Kiss1 neurons of *Kiss1-cre:MC4R-loxTB* mice (basal LH: 0.32±0.04, LH amplitude:

0.48±0.04, LH pulses/180min: 2.40±0.67 and AUC: 70.62±7.15 compared to controls) (**Figure

3*G-J***), further confirming that the melanocortin action on Kiss1 neurons is required for the normal

control of gonadotropin release. Despite these significant improvements in reproductive function,

Kiss1-cre:MC4R-loxTB mice presented fewer corpora lutea than controls (0.80±0.80), similar to

MC4R-LoxTB mice, suggesting that an ovulatory impairment still persists (**Figure 3*K, L***). While

tonic LH release and estrous cyclicity are predominantly controlled by Kiss1^{ARH} neurons, ovulation

is mediated by the induction of the preovulatory LH surge by Kiss1$^{AVPV/PeN}$ neurons. It is possible

that the metabolic signals derived from their obese phenotype, and/or the absence of the direct

action of MC4R in GnRH neurons (16, 25) prevents the complete recovery of ovulation in *Kiss1-*

cre:MC4R-loxTB mice. Both genetic models displayed significantly lower T (MC4R-LoxTB:

28.50±6.20 vs *Kiss1-cre:MC4R-loxTB:* 31.22±3.90 vs Controls: 62.84±16.53) and AMH (MC4R-

LoxTB: 150.6±22.0 vs *Kiss1-cre:MC4R-loxTB:* 146.9±10.8 vs Controls: 321.8±15.7) levels than

control mice in diestrus (**Figure 3*M, N***). Therefore, we can exclude, once again, a PCOS-like

reproductive phenotype mediated by the lack of melanocortin signaling.

Kiss1^{ARH} neurons are excited by MC4R agonists.

Based on the expression of *Mc4r* in Kiss1 neurons and the reproductive impairment found after

Mc4r deletion within Kiss1 neurons (**Figure 2**), we hypothesized that Kiss1^{ARH} neurons are direct

targets of melanocortins and would respond to MC4R activation. Initially we did whole-cell, voltage

clamp recordings in Kiss1^{ARH} neurons (**Figure 4A**). *Kiss1-Cre* x Ai32 or *Kiss1-Cre* AAV-injected

mice underwent ovariectomies (OVX) and were given estradiol (E2) replacement (*see Methods*).

We targeted fluorescent cells for recording which were isolated synaptically by bathing the slices

in tetrodotoxin (TTX, 1 μM). Focal application of the high-affinity melanocortin receptor agonist

melanotan II (MTII, ~250 nM) elicited a small inward current in half of the isolated Kiss1^{ARH} cells

(10/21) (**Figure 4B)**. Next, we examined if the E2 state affected MTII response by recording from

Kiss1^{ARH} neurons from OVX females and found two thirds (6/9 cells) responded, but there was no

significant difference in the average current (**Figure 4C**). Finally, it must be noted that at this

concentration MTII is a non-selective MCR agonist, so we followed up using a perfusion of the

MC4R-selective agonist THIQ (100 nM) to determine if activation of MC4Rs alone was sufficient

to invoke a postsynaptic response in Kiss1^{ARH} neurons from OVX+E2 females. Similar to MTII,

THIQ was able to elicit an excitatory inward current in Kiss1^{ARH} neurons (4/5 cells, -7.4±0.5 pA),

indicating MC4R activation is sufficient (**Figure 4D**).

POMC neurons synapse directly with Kiss1^{ARH} neurons.

Based on the pharmacological activation of MCRs, we wanted to address whether POMC^{ARH}

neurons are the source of melanocortins to excite Kiss1^{ARH} neurons directly. To answer this

question, we injected an AAV-EF1α-DIO-ChR2:mCherry vector into the ARH of adult *Pomc-Cre*

female mice (**Figure 4E**) (24). After 2-4 weeks we did whole cell recordings from putative Kiss1^{ARH}

neurons and looked for a response to high frequency optogenetic stimulation of POMC fibers in

OVX + E2-treated females (**Figure 4F**). We were able to confirm that these were Kiss1 neurons

based on the presence of a persistent sodium current and/or *post hoc* identification by scRT-PCR

(40/77) cells, **Figure 4G**). While this current is more prevalent in the AVPV Kiss1 population,

Kiss1 neurons are the only ARH neurons to display this electrophysiological "fingerprint" of a

pronounced INaP paired with a high capacitance and a low input resistance (26). Additionally,

while recording from putative Kiss1[ARH] neurons, we optogenetically stimulated POMC fibers at

low frequency (1 Hz, 5 ms, 470 nm light) and recorded fast postsynaptic inward currents. The fast

kinetics of the EPSC are also a tell-tale sign of an ionotropic glutamatergic response (27).

Previously, CNQX was sufficient to block similar excitatory ESPCs in other postsynaptic targets

of POMC neurons (28). While the consistent latency from optogenetic stimulus and current

response was indicative of a single synaptic delay, we wanted to establish that it was a direct

synaptic connection, as we have previously shown for output of POMC neurons (28) and Kiss1

neurons (29-31). First, we abrogated the optogenetic response with the addition of TTX (1 μM) to

the bath, we were able to rescue the postsynaptic glutamate response with the addition of the K^+

channel blockers 4-aminopyridine (4-AP; 0.5 mM) and tetraethyl ammonium (TEA; 7.5 mM)

(**Figure 4*H***). K+ channel block enables the calcium influx from ChR2 activation in the presynaptic

terminals to be sufficient to restore vesicle fusion. Therefore, there appears to be no intervening

synapses between POMC and the downstream Kiss1[ARH] neurons.

Previously we found that E2 increases glutamate release from POMC neurons by increasing

Vglut2 (28). Therefore, we compared the postsynaptic glutamatergic responses following two

optogenetic stimuli (50 ms interval between light flashes) in Kiss1[ARH] neurons from OVX + vehicle

versus OVX + E2-treated female mice (**Figure 4*I***). As anticipated, we found that treatment with

E2 increased the probability of glutamate release from POMC neurons onto Kiss1[ARH] neurons

based on the significant decrease in the paired pulse ratio of the two stimuli (32) (**Figure 4*J***).

Although in the present study we used an *in vivo* treatment paradigm, we know from previous

studies that this augmentation of glutamate release can happen quite rapidly after a brief

exposure to E2 *in vitro* (within 15 min) (28). Therefore, Kiss1[ARH] neurons are excited by the

glutamatergic input from POMC neurons in an E2-dependent manner.

Next, we tested for a melanocortin response using a high frequency stimulation protocol known

to elicit peptide release (33). Only cells displaying a glutamatergic response were tested. Indeed,

high frequency stimulation evoked an excitatory inward current (**Figure 4*K***) that was blocked by

the selective MC3/4R antagonist SHU 9119 (**Figure 4*L***). However, the number of high frequency

responses seen in Kiss1 neurons was unexpectedly infrequent and small (4/16, 25%, mean

inward current: -1.4±3.1 pA) compared to the efficacy of pharmacological MCR activation. E2-

treatment enhances *Pomc* expression and β-endorphin labeling (34) (35), but the effect, if any,

on αMSH release or the MCRs in Kiss1 neurons is unknown. We suspected that the high

frequency stimulation was eliciting co-release of αMSH and β-endorphin which would exert

oppositive effects on a postsynaptic cell. In order to isolate melanocortin signaling we pretreated

slices with the non-selective opioid antagonist naloxone (1 μM) and found an increase in the

likelihood and magnitude of a response to high frequency stimulation (9/23, 40%, mean inward

current: -7.6±2.9 pA) (**Figure 4*M***).

Together these optogenetic findings reinforced our pharmacological results showing that Kiss1^{ARH}

neurons are excited by MCR agonists. Additional studies would need to be done to identify the

cation conductance that is affected by the MC4R signaling cascade in Kiss1^{ARH} neurons, but

clearly we have established that there are excitatory glutamatergic and peptidergic inputs from

POMC to Kiss1^{ARH} neurons.

Kiss1$^{AVPV/PeN}$ neurons are inhibited by MC4R agonists.

Having established a direct, excitatory projection from POMC to Kiss1^{ARH} neurons, we next

examined POMC inputs to Kiss1^{AVPVPeN} neurons. First, we used immunocytochemistry to label

fibers expressing αMSH to demonstrate that POMC fibers densely innervated the AVPV/PeN area

(**Figure 5*A***). Next, we recorded from AVPV neurons in slices taken from OVX+E2, POMC-Cre

mice injected with AAV1-DIO-YFP:ChR2 into the ARH (**Figure *5B***). We targeted cells along the

ventricle in the AVPV and PeN, surrounded by YFP terminals, and Kiss1$^{AVPV/PeN}$ neurons were

identified based on the expression of several endogenous conductances (I_{NaP}, T-type calcium

current, h-current) that are unique to these neurons (26, 36). We recorded a direct glutamatergic

synaptic response following optogenetic stimulation in six neurons, which was further verified

through pharmacological "rescue" of the synaptic response in the majority (5/6) neurons tested

(**Figure 5C**), indicating that POMC neurons make direct monosynaptic connections with

Kiss1$^{AVPV/PeN}$ neurons.

We next hypothesized that Kiss1$^{AVPV/PeN}$ neurons would be excited by exogenous application of

melanocortin agonists because the receptor is typically Gs-coupled (37, 38). We did whole-cell,

voltage clamp recordings from Kiss1$^{AVPV/PeN}$-Cre::Ai32 neurons, which were isolated synaptically

by bathing the neurons in TTX (1 µM), from OVX females. For these experiments, we tested the

response to the high affinity melanocortin receptor agonist MTII (500 nM). Surprisingly, we

consistently measured an outward (inhibitory) current that could be reversed on washout of MTII

(**Figure 5D**). This outward current was associated with an increase in a K$^+$ conductance based

on the current-voltage plot *(i.e.,* the outward current reversed at ~E_{K+}, **Figure 5E**). Also, the

inwardly-rectifying I/V could indicate that MC4R is coupled to activation of inwardly-rectifying K$^+$

channels as has been previously reported for MC4R signaling in pre-autonomic parasympathetic

neurons in the brainstem (39). We hypothesized that the coupling could be modulated by E2 via

a Gα$_q$-coupled membrane estrogen receptor (Gq-mER) as we have previously demonstrated in

POMC neurons (40-42). For these experiments, we again utilized OVX females and targeted

Kiss1$^{AVPV/PeN}$-Cre::Ai32 neurons. Once in the whole-cell voltage-clamp configuration, we perfused

the slices with STX (10 nM), a selective ligand for the putative Gq-mER (42), and tested the

response to the MTII. Indeed, the outward current (inhibitory) response to MTII was completely

abrogated and even reversed by STX (**Figure 5F, G**). The short-term (bath) treatment with STX

ensured that there was no desensitization of the Gq-mER with longer-term (*in vivo*) treatment with

E2. Therefore, estrogen receptor activation can rapidly uncouple (*i.e.,* desensitize) the MC4R

inhibitory response in Kiss1$^{AVPV/PeN}$ neurons. We would predict that the intracellular signaling

cascade for the heterologous desensitization is similar to what we have elucidated in POMC

neurons (42), but this will need to be determined in future experiments. We next investigated the

response in vehicle-treated, OVX females. Indeed, Kiss1$^{AVPV/PeN}$ neurons were even more

inhibited by the same MTII exposure (**Figure 5*G***). Therefore, in contrast to Kiss1^{ARH} neurons, the

MC4R appears to be coupled to opening of K$^+$ channels in Kiss1$^{AVPV/PeN}$ neurons. To further

establish whether the MTII was having pre- or postsynaptic effects, we measured the

glutamatergic mEPSCs before and after melanocortin receptor activation. We found there was no

significant effect on the frequency (**Figure 5H**), but there was an effect on the amplitude (**Figure**

5I). This further supports a postsynaptic locus of STX's effects. However, *posthoc* comparisons

found STX increased amplitude compared to both OVX and OVX+E. This result hints at a

membrane delimited effect as STX is a much more potent agonist for Gq-mER than E2 (43).

Discussion

Our findings show that melanocortins act directly on Kiss1 neurons to contribute to the metabolic

regulation of fertility, in line with previous reports showing that αMSH stimulates LH release in a

kisspeptin-dependent manner (14) and that Kiss1 neurons express the melanocortin receptor

MC4R (18-21). Several studies in humans have linked *Mc4r* mutations to reproductive

abnormalities, including precocious puberty (9), PCOS (10), and hypogonadism (8). However, a

direct association between *MC4R* mutations and reproductive function, independent from the

obese condition of these patients, has not been identified (6). In the current study, we show that

the reproductive impairments observed in MC4R deficient mice, which replicate many of the

conditions described in humans, are largely mediated by the direct action of melanocortins via

MC4R on Kiss1 neurons and not to their obese phenotype. This is because the ablation of MC4R

from Kiss1 neurons largely replicated the reproductive impairments observed in MC4RKO female

mice without inducing obesity, and the selective reinsertion of MC4R into Kiss1 neurons of

MC4RKO mice significantly improved their reproductive function without changing their obese

phenotype. Strikingly, puberty onset was advanced in Kiss1-MC4RKO females. Our findings

revealed a low melanocortin tone in the hypothalamus of WT females during pubertal

development, characterized by lower levels of *Agrp, Mc4r* and *Mc3r* expression at the time of

puberty onset (~PND30). This developmental decline in the expression of melanocortin receptors

is in line with the advanced timing of puberty onset in females when MC4R is congenitally ablated

from Kiss1 neurons. A recent study using whole body MC3RKO mice showed a trend to delayed

puberty onset (19), suggesting a larger role of melanocortins in the timing of puberty onset. Here,

we show that in females the combination of MC3R and MC4R action may be necessary for

pubertal development. Interestingly, the MC4R-LoxTB and *Kiss1-cre:MC4R-LoxTB* mice do not

show an advancement in puberty onset as we observed in Kiss1-MC4RKO mice; however, these

mice are obese at the time of puberty onset already, which may affect the timing of sexual

maturation independently of MC4R action in Kiss1 neurons. Of note, the findings presented in

this study do not rule out the existence of additional sites of action of MC4R in other neuronal

populations to regulate reproduction, *e.g.,* GnRH neurons, which also express MC4R (16, 25).

Indeed, the lack of full recovery of the reproductive function in *Kiss1-cre:MC4R-LoxTB* mice

suggests that MC4R in Kiss1 neurons is necessary but not sufficient to achieve full reproductive

capabilities.

In addition to early puberty onset, Kiss1-MC4RKO females displayed an impaired preovulatory

LH surge driving ovulation and had fewer corpora lutea in their ovaries compared to control

littermates, further supporting a role of MC4R in regulating ovulation. In adulthood, the

reproductive phenotype observed in Kiss1-MC4RKO and MC4R-loxTB female mice (increased

LH, irregular estrous cycles, oligo-ovulation, increased cystic follicles) correlates with the

phenotype observed in PCOS mouse models (22, 23). In fact, an association between *MC4R*

mutations and PCOS has been reported (10). However, our mouse models failed to display higher

levels of circulating androgens or AMH, two of the hallmarks of PCOS (24), suggesting that

melanocortin signaling is unlikely to be a main contributing factor in the development of this

syndrome.

Our data revealing early puberty onset and impaired LH pulse and surge driving ovulation support

a direct role of MC4R on the two primary Kiss1 populations regulating these functions (ARH and

AVPV/PeN). We further investigated this and found that MC4R signaling excites Kiss1[ARH] neurons

and inhibits Kiss1[AVPV/PeN] neurons in an estradiol-dependent manner. In Kiss1[ARH] neurons, bath

applied or optogenetically-evoked release of melanocortin agonists induce a direct excitatory

inward current. Our studies using whole cell voltage clamp in female mice were able to detect a

melanocortin-mediated inward current in Kiss1[ARH] neurons that would increase excitability.

Previously, an MCR-activation effect was not detected in cell-attached recordings of diestrus

females (14, 44). However, cell-attached recordings do not measure a direct response in isolated

Kiss1[ARH] neurons but rather the summation of multiple synaptic inputs to Kiss1[ARH] neurons. Most

importantly, we did not see a difference in the excitatory response in synaptically isolated Kiss1[ARH]

neurons from E2-treated versus vehicle-treated, ovariectomized females. This emphasizes the

importance of doing whole-cell recording from isolated Kiss1 neurons. The modulatory actions of

POMC[ARH] neuronal projections to Kiss1[ARH] neurons can also occur through a) the excitatory action

of glutamate in a process that is facilitated in the presence of estradiol, in line with previous

publications (33, 45), and b) the inhibitory action of β-endorphin, suggesting that POMC[ARH]

neurons are able to exert a precise regulatory role of the GnRH pulse generator, *i.e.* Kiss1[ARH]

neurons, in response to metabolic challenges. In the AVPV/PeN, MC4R agonists inhibited

Kiss1[AVPV/PeN] neurons in an estradiol-dependent manner, which utilized a $G\alpha_q$-coupled membrane

estrogen receptor (STX receptor) to attenuate the inhibitory tone of MC4R in these neurons. Rapid

estrogen signaling may act to ease transitions between states. Membrane delimited E2 actions

can quickly attenuate or enhance coupling between receptors and signaling cascades. These

effects will precede E2-driven changes in gene expression that produce more stable alterations

in signaling. This combination of mechanisms will reduce any lag between rises in serum E2 and

physiological effects. Considering the abbreviated mouse reproductive cycle, parallel

mechanisms acting on different timescales are particularly important.

Our pharmacology data clearly demonstrates the estradiol-dependent action of MC4R in

Kiss1[AVPV/PeN] neurons, suggesting that similar to Kiss1[ARH] neurons, this population is regulated by

melanocortins. The effect of MC4R agonists on Kiss1 neurons in the presence of TTX, suggests

a direct synaptic effect without the need of channelrhodopsin-2(ChR2)-assisted circuit mapping.

Nonetheless, to further demonstrate this interaction, we provide evidence of profound innervation

of αMSH fibers and the presence of synaptic contact between POMC and Kiss1[AVPV/PeN] neurons

through the optogenetic stimulation of a glutamatergic response in Kiss1[AVPV/PeN] neurons after low

frequency stimulation of POMC terminals. We intentionally avoided high frequency stimulation of

POMC terminals to prevent the co-release of β-endorphins (e.g., **Figure 4M**) and their possible

additive effect on the inhibitory action of MC4R. Future studies will be aimed at the

characterization of the endogenous opioid pathways in the induction of the LH surge.

To date, two instances of MC4R-mediated inhibition have been described in the central nervous

system: MC4R can Gi,o couple to open K+ (Kir6.2) channels in parasympathetic preganglionic

neurons in the brainstem (39), and MC4R can directly couple to K+ (Kv7.1) channels in the

hypothalamic paraventricular nucleus neurons (46). Interestingly, our findings describe an

inhibitory role of MC4R in the reproductive neuroendocrine axis. This steroid state dependent

effect is particularly relevant because in the context of reproduction, the two populations of Kiss1

neurons are strikingly different in their response to estradiol, where peptide expression in Kiss1[ARH]

and Kiss1[AVPV/PeN] neurons are inhibited or stimulated, respectively, in order to mount the negative

versus positive feedback of sex steroids (15). This differential regulation allows for the episodic

versus surge release of GnRH; however, the cellular mechanisms underlying these opposing

roles of Kiss1 neurons to the same stimulus, *i.e.* circulating E2 levels, remain unknown. Although

there were no differences in the magnitude of the electrophysiological responses to MTII in

Kiss1[ARH] neurons in vehicle- versus E2-treated, ovariectomized females, we cannot rule out that

E2 affects other systems activated by the MC4R signaling cascade as recently reported in MC4R

expressing neurons of the ventromedial nucleus of the hypothalamus (VMH) (47). Interestingly,

puberty onset was advanced in Kiss1-MC4RKO females. This data suggests that the inhibitory

tone of MC4R signaling on Kiss1$^{AVPV/PeN}$ neurons might also be involved in the timing of puberty

onset, which in turn suggests a role for this female-specific population of Kiss1 neurons in sexual

maturation.. Furthermore, our findings suggest that metabolic cues, through the regulation of the

melanocortin output onto Kiss1$^{AVPV/PeN}$ neurons, are essential for the timing and magnitude of the

GnRH/LH surge. This study suggests that the melanocortin-kisspeptin pathway is one of the

multiple pathways essential for the metabolic regulation of the HPG axis. Future studies are

warranted to fully elucidate synaptic input and postsynaptic cascades in the POMC-Kiss1$^{AVPV/PeN}$

circuit.

Altogether, our data reveal a differential regulatory action of MC4R in the neural control of

GnRH/LH release, participating in both the surge (E2-treated, ovariectomized female) and the

pulse-like (ovariectomized female) modes of LH release through the cellular regulation of Kiss1

neurons, which translate into our findings on the conditional knockout of *Mc4r* from Kiss1 neurons

(**Figure 6**). These findings are important because the reproductive abnormalities often attributed

to obesity in MC4R deficient patients may not be caused by the excess in body weight but, at

least in part, by a deficiency in MC4R signaling directly at the level of Kiss1 neurons that affects

predominantly the ability to mount a preovulatory LH surge.

Materials and methods:

In-vivo experimental procedures:

Generation of Kiss1-MC4RKO, Kiss1-cre:MC4R-LoxTB and MC4R-LoxTB Transgenic Mice

Kiss1-MC4RKO mice were generated by crossing *Kiss1-cre* knock-in mice and *MC4R$^{lox/lox}$*

mice. *Kiss1-cre* mice (RRID:MGI:6278139) were obtained from Dr. Richard Palmiter (University

of Washington, Seattle, WA) (49) and *Mc4r$^{lox/lox}$* (RRID:IMSR_JAX:023720) were a gift from Dr.

Brad Lowell (Beth Israel Deaconess Hospital, Boston, MA) (4). These mice were crossed to

generate Kiss1-MC4RKO mice lacking *Mc4r* expression selectively from Kiss1 neurons (*Kiss1-*

cre$^{+/-}$; *MC4R$^{lox/lox}$*) and their control littermates (MC4R$^{lox/lox}$). To generate *Kiss1-cre:MC4R-LoxTB*

and *MC4R-LoxTB* mice, *MC4R-LoxTB* mice were purchased from The Jackson Laboratory

(MC4R-LoxTB; catalog no. 006414). These mice present with a loxp-flanked transcriptional

blocking (LoxTB) sequence preventing normal endogenous gene transcription and translation

from the endogenous locus. Homozygous MC4R-loxTB/loxTB mice are devoid of functional

MC4R mRNA (MC4RKO mice), while the presence of Cre recombinase on the Kiss1 promotor in

Kiss1-cre mice will result in the removal of the transcription blocker and subsequent expression

of MC4R in tissue-specific sites (*i.e.* Kiss1 neurons) therefore resulting in the generation of *Kiss1-*

cre:MC4R-LoxTB (with *Mc4r* expression restored in Kiss1 neurons), their obese control

littermates *MC4R-LoxTB* mice (MC4RKO), and their WT controls (MC4R$^{+/+}$). To rule out early

embryonic recombination of the MC4R$^{loxTB/loxTB}$ or MC4R$^{lox/lox}$ alleles, we ran PCR assays on tail

DNA designed to detect wild-type allele (MC4R$^{+/+}$), undeleted lox (MC4R$^{lox/lox}$) or loxTB alleles

(MC4R$^{loxTB/loxTB}$). Genotyping was confirmed by sending tissue to Transnetyx, Inc., for testing by

real-time polymerase chain reaction. Mice were housed in Harvard Medical School Animal

Resources facilities where they were fed standard mouse chow (Teklad F6 Rodent Diet 8664)

and were given *ad libitum* access to tap water under constant conditions of temperature (22–

24˚C) and light (12 hr light [07:00]/dark [19:00] cycle).

RNAscope in situ hybridization

To validate the Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and MC4R-LoxTB (MC4RKO) mouse

models and investigate the co-expression of *Kiss1* and *Mc4r* mRNA in these mice, dual

fluorescence ISH was performed using RNAscope (ACD, Multiplex Fluorescent v.2) according to

the manufacturer's protocol using the following probes: *Mc4r* (319181-C2) and *Kiss1* (500141-

C1). Brains (n=4/group) from WT OVX (for expression in the ARH) and OVX+E2 (for expression

in the AVPV/PeN) mice and OVX Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and MC4RKO were

removed fresh frozen on dry ice, and then stored at -80˚C until sectioned. Five sets of 20 µm

sections in the coronal plane were cut on a cryostat, from the diagonal band of Broca to the

mammillary bodies, thaw mounted onto SuperFrost Plus slides (VWR Scientific) and stored at -

80˚C until use. A single set was used for ISH experiment (adjacent sections 100 mm apart).

Images were taken at 20x magnification of the sections containing AVPV, PeN, and the three

rostro-to-caudal levels of the ARH, and Kiss1 neurons expressing (*Kiss1-cre*:MC4R-LoxTB mice)

or lacking (Kiss1-MC4RKO mice) *Mc4r* were identified using ImageJ.

Reproductive maturation of Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and MC4R-LoxTB

mice

To assess the reproductive phenotype of mice with selective reinsertion of *Mc4r* in Kiss1 neurons

(*Kiss1-cre*:MC4R-LoxTB, n=11), selective deletion of *Mc4r* from Kiss1 neurons (Kiss1-MC4RKO,

n=24), global deletion of *Mc4r* (MC4R-LoxTB, n=7); and their control MC4R$^{lox/lox}$ littermates

(n=14), mice were weaned at post-natal day (PND) 21 and were monitored daily for puberty onset.

Females were monitored daily for vaginal opening (VO, indicative of the complete canalization of

the vaginal cavity) and for first estrus (first day with cornified cells determined by daily morning

vaginal cytology) after the day of VO. Body weight (BW) was measured at the day of puberty

onset to determine if changes in puberty onset could be due to differences in BW. Estrous cyclicity

was monitored in females by daily vaginal cytology, for a period of 15 days, in 6-month-old mice

and their respective control littermates (n=5/group). Cytology samples were obtained every

morning (9 am), placed on a glass slide and stained with hematoxylin and eosin for determination

of the estrous cycle stage under the microscope.

Fecundity test of Kiss1-MC4RKO females

Adult 6-month-old Kiss1-MC4RKO and control littermate female mice (n=9/group) were placed

with adult WT males proven to father litters for 90 days and time to deliver pups and number of

pups per litter were monitored.

Characterization of the estradiol-induced luteinizing hormone surge

Kiss1-MC4RKO (n=6) and control MC4R$^{lox/lox}$ littermate (n=5) adult female mice were subjected

to bilateral ovariectomy (OVX) via abdominal incision under light isoflurane anesthesia.

Immediately after OVX, capsules filled with E2 (1 ug/20g BW) were implanted subcutaneously via

a small mid- scapular incision on the back. Five days later, mice were subcutaneously injected in

the morning with estradiol benzoate (1 ug/20g BW) to produce elevated proestrus-like E2 levels

(preovulatory LH surge) on the following day (50). Blood samples were collected at 8am and 7pm;

LH levels were stored at -80˚C until measured via LH ELISA.

Ovarian histology and hormone measurements

Bilateral ovariectomy of 6-month-old Kiss1-MC4RKO (n=4), *Kiss1-cre*:MC4R-LoxTB, MC4R-

LoxTB and their control littermates (n=5/group) was performed under light isoflurane anesthesia.

Briefly, the ventral skin was shaved and cleaned, and one small abdominal incision was made.

Once the ovaries were identified and excised, the muscle incision was sutured, and the skin was

closed with surgical clips. Ovaries were stored in Bouin's fixative, sectioned, and stained with

hematoxylin and eosin at the Harvard Histopathology Core. Corporal lutea were counted in the

middle section of each ovary. Serum samples were also collected for analysis of testosterone and

anti-müllerian hormone (AMH) levels in these mice. These hormones levels were measured at

the University of Virginia Ligand Assay core with the Mouse & Rat Testosterone ELISA assay

(reportable average range 10-1600 ng/dL; sensitivity of 10 ng/dL); AMH ELISA assay (reportable

average range 0.2 – 15 ng/mL; sensitivity of 0.2 ng/mL).

LH pulsatile secretion profile in gonad intact Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and

MC4R-LoxTB female mice

To assess the profile of LH pulses secretion, adult 6 months old gonad intact Kiss1-MC4RKO

females (n=6), their control Mc4r$^{lox/lox}$ littermates (n=7); and *Kiss1-cre*:MC4R-LoxTB (n=5/group),

MC4R-LoxTB and their control WT females (n=4/group) were handled daily for 3 weeks to allow

acclimation to sampling conditions prior to the experiment. Pulsatile measurements of LH

secretion were assessed in diestrus by repeated blood collection through a single incision at the

tip of the tail. The tail was cleaned with saline and 4 µL of blood was taken at each time point from

the cut tail with a pipette. We collected sequential blood samples every 10 minutes over a 180-

minute sampling period. Samples were immediately frozen on dry ice and stored at −80 °C until

analyzed with LH ELISA as previously described (51). The functional sensitivity of the ELISA

assay was 0.0039 ng/ml with a CV% of 3.3%.

LH pulses analysis

LH pulses in mice were analyzed using a custom-made MATLAB-based algorithm. The MATLAB

code includes a loop that determines LH pulses as any LH peak: (i) whose height is 20% greater

than the heights of the 2 previous values; (ii) 10% greater than the height of the following value;

and (iii) the peak at the second time interval needs to be 20% greater than the single value that

comes before it to be considered a pulse, as we previously described (52).

LH pulsatility was assessed by measuring: (1) the total secretory mass, assessed by area under

the curve (AUC); (2) the LH pulse amplitude, calculated by averaging the 4 highest LH values in

the samples collection period for each animal; (3) the basal LH, calculated by averaging the 4

lowest LH values in the samples collection period for each animal; and (4) the total number of

pulses throughout the 180 minutes sampling period.

Immunohistochemistry

Animals and Treatment: Coronal brain blocks (2 mm each) from adult female C57BL/6 mice (n=4)

were fixed by immersion in 4% paraformaldehyde for ~ 8 hours, cryoprotected in 30% sucrose

solution, frozen in isopentane at –55°C, sectioned coronally on a cryostat at 20 µm, and thaw-

mounted onto Superfrost Plus slides (Fisher Scientific, Pittsburgh, PA). The 20 µm sections were

stored at –20°C until used for immunocytochemistry.

Immunocytochemistry: The *s*ections were rinsed in PB (0.1M phosphate buffer, pH 7.4) for at

least 30 min. Next, sections were incubated with normal serum corresponding to the host for the

secondary antiserum (5 % normal serum with 0.3% Triton-X100 in PBS for 30 min), rinsed in PB

and then incubated for ~45 h at 4º C with a rabbit polyclonal antiserum against α-MSH (1:2,500).

The specificity of this antiserum has been documented (53). After rinsing, sections were first

incubated for 2-3 hours at room temperature with biotinylated donkey anti rabbit gamma globulin

(IgG; 1:500) and next with streptavidin-Alexa 488 (1:2500). Both the primary and secondary

antibodies were diluted in tris-(hydroxymethyl) aminomethane (0.5 %, Jackson ImmunoResearch,

Philadelphia, PA) in PB containing 0.7 % seaweed gelatin (Jackson ImmunoResearch,

Philadelphia, PA) and 0.5 % Triton X-100 and 3% bovine serum albumin (BSA; Jackson

ImmunoResearch, Philadelphia, PA) adjusted to pH 7.6. Following a final rinse overnight, slides

were coverslipped with gelvatol containing the anti-fading agent, 1,4-diazabicyclo(2,2)octane

(DABCO; Cold Spring Harbor Protocols, 2006).

Imaging: Photomicrographs of labeling were initially acquired using a Nikon E800 fluorescent

microscope (Eclipse E800; Nikon Instruments, Melville, NY) equipped with a fiber illuminator

(Intensilight C-HGFI; Nikon Instruments) and a high-definition digital microscope camera head

(DS-Fi1; Nikon Instruments) interfaced with a PC-based camera controller (DS-U3; Nikon

Instruments).

Real time quantitative PCR

i) To investigate the changes in the expression of the melanocortin genes *Agrp*, *Pomc, Mc3r* and

Mc4r in the MBH during development in prepubertal and pubertal WT females at ages P10 (n=6),

P15 (n=6), P22 (n=6), and P30 (n=5), and ii) evaluate the gene expression profile of *Pdyn*, *Kiss1*,

Tac2, *Tacr3*, in the ARH of adult intact (in diestrus) female Kiss1-MC4RKO (n=6) and their control

MC4R$^{lox/lox}$ littermates (n=3), the brains were removed and rapidly embedded in Tissue-Tek,

frozen in −30°C isopentane solution and stored at −80°C until use. Frozen tissue punches were

recovered through MBH with a 1 mm diameter canula (54). These tissue punches encompassed

the whole MBH from the WT females (figure 1D) and the ARH from the Kiss1-MC4RKO and their

control females (Figure 2I). The tissues were homogenized, and total RNA was isolated using

TRIzol reagent (Invitrogen) followed by chloroform/isopropanol extraction. RNA purity and

concentration were measured via an absorbance spectrophotometer (260/280 nm > 1.8;

NanoDrop 1000, Thermo Fisher Scientific). Total RNA (1 µg) was reverse transcribed to cDNA

using random hexamers (High-Capacity cDNA Synthesis Kit, Life Technologies). Quantitative

real-time PCR assays were performed using SYBR Green RT-qPCR master mix (Applied

Biosystems) and analyzed using ABI Prism 7000 SDS software (Applied Biosystems). The cycling

conditions were: 2 minutes incubation at 95°C (hot start), 45 amplification cycles (95°C for 30

seconds, 60°C for 30 seconds, and 45 seconds at 75°C, with fluorescence detection at the end

of each cycle), followed by melting curve of the amplified products obtained by ramped increase

of the temperature from 55°C to 95°C to confirm the presence of single amplification product per

reaction. PCR specificity was verified by melting curve analysis and agarose gel electrophoresis.

Each sample was run in duplicate to obtain an average cycle threshold (CT) value and relative

expression of each target gene was determined using the comparative Ct method (55). The data

were normalized to Hypoxanthine Guanine Phosphoribosyltransferase (*Hprt)* expression levels in

each sample. Results were expressed as fold differences in relative gene expression with respect

to i) P10 for melanocortin genes expression analysis during development in WT mice, and ii)

controls for the KNDy genes expression in female Kiss1-MC4RKO mice. The primers used are

listed in Table 1.

Gene	Gene length (pb)	Accession #	Primers	Location (nt)	Sequence
Hprt	352	NM_013556.2	Hprt-F	704-728	CCTGCTGGATTACATTAAAGCGCTG
			Hprt-R	377-401	GTCAAGGGCATATCCAACAACAAAC
Agrp	136	NM_001271806.1	Agrp-F	466-488	GCCTCAAGAAGACAACTGCAGAC
			Agrp-R	580-601	AAGCAGGACTCGTGCAGCCTTA
Pomc	138	NM_001278584.1	Pomc-F	187-208	CCATAGATGTGTGGAGCTGGTG
			Pomc-R	303-324	CACCTCCGTTGCCAGGAAACAC
Mc4r	101	NM_016977.4	Mc4r-F	552-570	CCCGGACGGAGGATGCTAT
			Mc4r-R	632-652	TCGCCACGATCACTAGAATGT
Pdyn	200	NM_018863.4	Pdyn-F	45-64	ACAGGGGGAGACTCTCATCT
			Pdyn-R	223-244	GGGGATGAATGACCTGCTTACT
Kiss1	129	AF472576.1	Kiss1-F	147-166	GCTGCTGCTTCTCCTCTGTG
			Kiss1-R	256-275	TCTGCATACCGCGATTCCTT
Tac2	234	NM_001199971.1	Tac2- F	238-257	GCTCCACAGCTTTGTCCTTC
			Tac2- R	452-471	GCTAGCCTTGCTCAGCACTT
Tacr3	159	NM_021382.6	Tacr3-F	759-779	GCCATTGCAGTGGACAGGTAT
			Tacr3-R	898-917	ACGGCCTGGCATGACTTTTA

Data analysis

Statistical data are expressed as means ± SEM, where n represents the number of animals in

each study group. The significance of differences between groups was evaluated using unpaired

two-tailed Student's t test, or a one- or two-way ANOVA test (with *post hoc* comparisons).

Significance level was set at $p<0.05$. All analyses were performed with GraphPad Prism Software,

Inc (San Diego, CA).

In-vitro experimental procedures:

Animals:

All animal procedures described in the electrophysiology studies were performed in accordance

with institutional guidelines based on National Institutes of Health standards and approved by the

Institutional Animal Care and Use Committees at Oregon Health and Science University and

Appalachian State University. *Kiss1-cre* (v2) mice (49) were crossed with Ai32 (56) or C57B6J

mice. *Pomc-Cre* mice (RRID:IMSR_JAX:005965) (57) were crossed with wildtype C57B6J

(RRID:IMSR_JAX:000664) mice. The Ai32 cross was not used with *Pomc-Cre* mice because the

gene is transiently expressed during development in some cells fated to be Kiss1 or AgRP cells

(58) and early recombinant events lead to persistent expression of the ChR2-mCh fusion protein

in non-POMC cells. However, we have previously shown that AAV-driven expression in adult

Pomc-cre animals is restricted to β-endorphin labeled cells in the ARH (45), avoiding this non-

specificity issue. All colonies were maintained onsite under controlled temperature (21-23 ºC) and

photoperiod (12:12-h light-dark cycle 0600 to 1800) while receiving ad libitum food (5L0D;

LabDiet, St. Louis, MO) and water access. Following surgeries, mice received a *s.c.* dose of 4-5

mg/kg carprofen (Rimadyl; Pfizer Animal Health, New York, NY) and given at least 1 week of

recovery.

Ovariectomies:

Seven to ten days prior to each experiment, ovaries were removed as described previously while

under isoflurane anesthesia (45). Two days before experiments females received either an

injection of sesame oil (50 µl, *sc*; Sigma-Aldrich, St. Louis, MO) or a priming dose (0.25 µg /50 µl

sesame oil, *sc*) of E2 benzoate (Sigma-Aldrich, St. Louis, MO) in the morning. On the following

day, oil or a high (1.5 µg) dose of E2 benzoate, which generates a LH surge, was administered

(59). Circulating levels of E2 were verified by the uterine weights (<25 mg for OVX and >95 mg

for OVX E2-treated) at the time of euthanasia (59).

AAV Delivery:

Bilateral ARH injections of AAV1-Ef1a-DIO-ChR2:mCherry (RRID:Addgene_20297) or AAV1-

Ef1a-DIO-ChR2:YFP (RRID:Addgene_20298) were performed on adult *Kiss1-cre* mice or *Pomc-*

cre mice on a stereotaxic frame under isoflurane anesthesia. ARH injection coordinates were

anteroposterior (AP): −1.10 mm, mediolateral (ML): ± 0.30 mm, dorsoventral (DL): −5.80 mm

(surface of brain z = 0.0 mm); 400 nl of the AAV (2.0 x 10^{12} particles/ml) was injected (100 nl/min)

into each position. Mice were given carprofen for analgesia and allowed to recover for at least

two weeks before euthanasia.

Visualized whole-cell patch recordings:

Hypothalamic coronal brain slices (240 um) were made from female mice, using a Leica VT1000S

vibratome, in ice cold cutting solution bubbled with O_2/CO_2 (95%/5%). Slices were then transferred

to a holding chamber with artificial cerebrospinal fluid bubbled with the same gas mix and allowed

to recover for at least 1 hour. For recordings, slices were placed in a perfusion chamber and

visualized with an Olympus BX51W1 using either differential infrared contrast or oblique

illumination. Kiss1 neurons in the ARH or AVPV/PeN were targeted for electrophysiological

recordings as done previously and described below (60, 61). Final concentration was calculated

based on the drug in the known volume of the bath. Perfused drugs (TTX, naloxone, and STX)

were constantly circulated and given at least 10 minutes to reach maximal effect. Focal application

of drugs (MTII and THIQ) was done with the pump stopped (10-20 minutes). 0.3 to 1 µl was

directly added to the known volume of the bath to achieve the desired final concentration. This

approach enables precise timing of recordings and reduces the risk of receptor desensitization.

Solutions/drugs:

Standard vibratome slicing, external, and internal recording solutions were utilized as previously

described (60, 61). Tetrodotoxin was purchased from Alomone Labs (Jerusalem, Israel),

Melanotan II and αMSH from Tocris (Minneapolis, MN). TEA, 4-AP, 17β-estradiol benzoate, and

Naloxone were purchased from Millipore-Sigma. STX was produced by AAPharmaSyn, LLC (Ann

Arbor, MI) under contract.

Electrophysiology data analysis:

Electrophysiological data were analyzed using Clampfit 10/11 (Molecular Devices) and Prism 7/10

(Dotmatics). All values are expressed as Mean ± SEM. Comparisons between two groups were

made using un-paired Student's t-test or between multiple groups using an ANOVA (with *post hoc*

comparisons) with p-values < 0.05 considered significant. When variances differed significantly,

Mann-Whitney U test was used instead.

Targeting of Kiss1 neurons for electrophysiological recordings.

For non-optogenetic experiments, brain slices were taken from AAV injected Kiss1-Cre AAV or

Kiss1xAi32 female mice. Ai32 mice (RRID:IMSR_JAX:024109, C57BL/6 background) carry the

floxed ChR2 (H134R)-EYFP gene in their Gt(ROSA)26Sor Locus (56), allowing its expression in

a Cre-dependent manner. Due to concerns of nonspecific expression (62), we previously

validated this model using single cell RT-PCR and documented that *Kiss1* mRNA was detectable

in 99% of individually harvested eYFP cells (n=126). In addition, we have used both AAV injected

Kiss1-Cre AAV or Kiss1xAi32 female mice and found no differences in electrophysiological results

(60). We used AAV injected POMC-Cre female mice and avoided small soma, high input

resistance (>800 MΩ), ventrally located cells that are typically NPY/AgRP neurons. Instead, we

targeted larger, more dorsomedial neurons in the ARH while avoiding fluorescent (POMC) cells.

Unlike ARH Kiss1 neurons, POMC neurons do not make reciprocal projections, so uninfected

POMC neurons do not display monosynaptic EPSCs in response to optogenetic stimulation. Next,

we used a ramp IV protocol to probe for the presence of a persistent sodium current (I_{NaP}). While

this current is more prevalent in the AVPV Kiss1 population, Kiss1 neurons are the only ARH

neurons to display this electrophysiological "fingerprint" of a pronounced I_{NaP} paired with a high

capacitance and low input resistance (26). Finally, we also harvested the cytosol at the end of all

recordings and measured the expression of Kiss1. Only cells that expressed a persistent sodium

current and/or expressed *Kiss1* mRNA were included in final analysis (n = 40 ARH & 6 AVPV).

Acknowledgments:

The authors would like to recognize the excellent technical expertise of Ms. Martha A. Bosch

(tissue preparation, immunohistochemical procedure and single cell RT-PCR of recorded

neurons). In addition, we thank Dr. Rona Carroll for her assistance at Harvard Medical School

and Mr. Cole Martinson, a student worker in the Ronnekleiv/ Kelly laboratories, for his assistance

with genotyping and care of mouse colonies at OHSU. The University of Virginia Center for

Research in Reproduction Ligand Assay and Analysis Core is supported by the Eunice Kennedy

Shriver NICHD/NIH (NCTRI) Grant P50-HD28934. This work was supported by PHS MPI grant

DK68098 to MJK and OKR; HD090151, HD099084 and DK133760 to VMN, and The Charles A.

King Trust Postdoctoral Research Fellowship Award, The Lalor Foundation Postdoctoral

Fellowship Award, the Women's brain initiative Fellowship Award, the ROSA SCORE pilot grant

(supported by NIH Research Grant U54 AG062322 funded by The National Institute on Aging

(NIA) and Office of Research on Women's Health (ORWH)) and the IBRO-ISN Research

Fellowship Award to RT, and startup funds from Appalachian State University to TLS.

Authors contribution:

Declaration of interests:

735   The authors declare they have no competing interest.

Figure legends:

Figure 1. *Mc4r* expressed in Kiss1 neurons determines the timing of puberty onset.

Kiss1-MC4RKO females display advanced puberty onset, assessed by daily monitoring of vaginal opening **(A)** and first estrus **(B)**, as documented by cumulative percent and mean age of animals at vaginal opening **(A)** and first estrus **(B)** in Kiss1-MC4RKO females (n=24) compared to WT littermates (n=19). *p<0.05 by Student's t-test. Data presented as the mean± SEM. **(C)** Female Kiss1-MC4RKO (n=19) have normal body weight at the time of puberty onset compared to their WT littermates (n=15). **(D)** Ontogeny expression of melanocortin genes (*Agrp*, *Pomc*, *Mc4r and Mc3r*) in the ARH of WT female mice at different postnatal pre-pubertal and pubertal ages: P10, P15, P22 and P30, normalized to the housekeeping gene *Hprt*. (*n values*: females at P10 (n=6), P15 (n=6), P22 (n=6), and P30 (n=5)). Groups with different letters are significantly different

(p<0.05), as determined by one-way ANOVA followed by the Student-Newman-Keuls test. Data

presented as the mean± SEM.

Figure 2. Deletion of MC4R from Kiss1 neurons impairs fertility in Kiss1-MC4RKO females.

(A) Female Kiss1-MC4RKO (n=19) have normal body weight from weaning and until post-natal

day (PND) 150 compared to their WT littermates (n=15). **(B)** Representative examples of estrous

cycles in Kiss1-MC4RKO and control females (n=5/group) assessed by daily vaginal cytology for

15 days. **(C)** Kiss1-MC4RKO females displayed irregular estrous cycles with a longer time in

diestrus and a shorter time in estrus compared to control females. *** P < 0.001 Two-way ANOVA.

(D) Pattern of LH pulsatility was analyzed in gonad intact Kiss1-MC4RKO (n=6) and control (n=7)

females. # Represents LH pulses. **(E)** LH total secretory mass assessed by area under the curve

(AUC), **(F)** LH pulse amplitude, **(G)** Basal LH and **(H)** Total number of pulses /180 min were

analyzed. *p<0.05 by Student's t-test. The expression of the KNDy genes *Pdyn* **(I)**, *Kiss1* **(J)**,

Tac2 **(K)** and *Tacr3* **(L)**, was assessed in the ARH of adult Kiss1-MC4RKO (n=6) and control

(n=3) females. **p<0.01 by Student's t-test. Data presented as the mean± SEM. **(M)** Kiss1-

MC4RKO (n=6) and control (n=5) females were subjected to an LH surge induction protocol. LH

samples were collected in the morning (AM [8 a.m.]) and evening (PM [7 p.m.]) after lights off. **P

< 0.01, Two-way ANOVA followed by Bonferroni *post hoc* test. **(N)** Ovarian histology shows a

decrease in the number of corpora lutea (CL, **O**) and increase in the number of cystic follicles

(CF, **P**) of Kiss1-MC4RKO compared to controls (n=5/group). # represents cystic follicles. Serum

levels of Testosterone **(Q)** and AMH **(R)** in adult gonad intact Kiss1-MC4RKO and control females

(n=5/group). Student t test for unpaired samples. Data presented as the mean± SEM. Kiss1-

MC4RKO females display impaired fertility characterized by increased time to deliver pups **(S)**

and decreased number of pups per litter **(T)**, (n=9/group). *p<0.05 and **p<0.01 by Student's t-

test. Data presented as the mean± SEM.

Figure 3. Re-insertion of MC4R in Kiss1 neurons restores estrous cyclicity and LH pulsatility in Kiss1-cre:MC4R-LoxTB females.

(A) *Kiss1-cre*:MC4R-LoxTB (n=11) and MC4R-loxTB (MC4RKO) (n=7) displayed normal puberty onset as compared to their control littermates (n=14), as documented by cumulative percent and mean age of animals at vaginal opening and first estrus. **(B)** *Kiss1-cre*:MC4R-LoxTB and MC4R-loxTB both had significantly higher body weights at the time of puberty onset compared to their controls. **P < 0.01, *** P < 0.001 by one way ANOVA. **(C)** MC4R-loxTB (n=7) and *Kiss1-cre*:MC4R-LoxTB (n=11) females displayed significantly higher body weight than their littermates (n=10) from post-natal day 30 onwards, *p<0.05 and **P < 0.01. Data presented as the mean± (SEM). **(D)** Representative examples of estrous cycles of MC4R-loxTB (n=4), *Kiss1-cre*:MC4R-LoxTB (n=5) and control littermate mice, (n=4) assessed by daily monitoring of vaginal smears for 15 days. (D/M: diestrus/metestrus, P: proestrus, E: estrus). **(E)** MC4R-loxTB females displayed irregular estrous cycles, presenting longer time in diestrus and shorter time in estrus compared to control females, while *Kiss1-cre*:MC4R-LoxTB females displayed regular estrous cyclicity, similar to controls. *p<0.05, **P < 0.01, *** P < 0.001, ****P < 0.0001, Two-way ANOVA. Data presented as the mean± SEM. **(F)** Pattern of LH pulsatility analyzed in gonad intact MC4R-loxTB (n=4), *Kiss1-cre*:MC4R-LoxTB (n=5) and control littermates females (n=5). LH samples were collected every 10 min for 180 minutes; # represents LH pulses. **(G)** LH total secretory mass, **(H)** LH pulse amplitude, **(I)** basal LH, and **(J)** total number of pulses /180 min were assessed. *p<0.05, **P < 0.01 by one way ANOVA. **(K)** Representative samples of ovarian histology from MC4R-loxTB, *Kiss1-cre*:MC4R-LoxTB and control females (n=5/group); CL: Corpora lutea. Data are presented as the mean± SEM. **(L)** Ovarian histology showed a significant decrease in the number of corpora lutea in the MC4RKO and *Kiss1-cre*:MC4R-LoxTB compared to controls. Groups with different letters are significantly different. Serum levels of **(M)** Testosterone and **(N)** AMH in adult gonad intact females MC4R-loxTB, *Kiss1-cre*:MC4R-LoxTB and their control littermates. *p<0.05, ****P < 0.0001, One Way ANOVA. Data are presented as the mean± SEM.

Figure 4. Kiss1^{ARH} neurons are activated by melanocortin agonists and respond to optogenetic stimulation of POMC neurons.

(A) Slices were taken from *Kiss1-cre* injected (confocal image) brains to target fluorescent cells (arrowhead) for recording (white outline over electrode). **(B)** Whole cell voltage clamp recording of Kiss1^{ARH} neurons following the direct application of the high affinity melanocortin receptor agonist melanotan II (MTII, 250 nM) was added directly to the bath and the response was compared between slices from OVX and OVX+E females. **(C)** While the average inward current was slightly higher in the OVX+E state, this was not a significant difference (Student's t-test for unpaired samples, p>0.05). **(D)** The MC4R-selective agonist THIQ (100 nM) was perfused, and excitatory inward currents were generated in Kiss1^{ARH} neurons. **(E)** Images are of AAV-driven labeling of POMC cells as seen through the confocal. **(F)** Using the AAV-driven expression of channelrhodopsin in adult *Pomc-cre* mice, high frequency stimulation elicited a slow inward current in Kiss1 neurons. **(G)** The identity of cells was confirmed through the presence of a persistent sodium current (see (26)) and/or with RT-PCR of harvested cytoplasm showing *Kiss1* expression. (Gel: MM=molecular marker, TC=tissue controls). **(H)** A direct synaptic projection from POMC to Kiss1^{ARH} neuron (*posthoc* identified) was confirmed using the "rescue" protocol: 1, baseline glutamatergic responses were initially generated (black trace); 2, then action potentials were eliminated by blocking voltage-gated sodium channels with TTX and the postsynaptic response (red trace); 3, blockade of potassium channels facilitated calcium entry into the terminal through ChR2 to release synaptic vesicles, "rescuing" the postsynaptic glutamate response (green trace). **I)** Glutamatergic responses were also often observed, particularly in the ventral ARH. As seen when targeting low input resistance neurons (i.e., Kiss1^{ARH}) with low frequency optogenetic stimulation (5 ms pulse, 50 ms inter-spike interval), the first response in OVX + vehicle females was larger relative to the second response in OVX + E2- treated female mice. Representative traces are the average of 30 sweeps. **(J)** The averaged paired-pulse ratio was

lower in recordings from estradiol-treated female mice, indicating an increased release probability

(Students t-test p<0.05,). **(K)** High frequency optogenetic stimulation elicited a small inward

current. **(L)** In a different cell, a high frequency inward current was noted before constant perfusion

of SHU9119 for 15 minutes (break between traces). When the stimulation protocol was repeated,

no inward current was elicited. **(M)** The majority of high frequency responses were inward and

only twice was an inhibitory outward current recorded in identified Kiss1^{ARH} neurons. Perfusion of

the non-selective opioid receptor antagonist naloxone reversed the current, eliminating MCR-

activation as the mechanism.

Figure 5. Kiss1$^{AVPV/PeN}$ neurons are inhibited by MC4R agonists in an E2-dependent manner.

(A) Immunohistochemistry showing robust labeling of α-MSH fibers in the AVPV/PeN region in

OVX+E2 WT female mice. **(B)** Low and high power bright-field images taken during

electrophysiology recording in OVX+E2 POMC-Cre mice expressing YFP:ChR2 in the ARH

following AAV injection. (left) Low power image shows the location of recorded cells. (right) Higher

power image of area from white box inset displaying POMC fibers innervating the area. **(C)** Fibers

surrounded putative Kiss1-AVPV neurons (expressing I$_{NaP}$, IT and I$_h$) and optogenetic stimuli was

able to elicit postsynaptic currents that were eliminated with tetrodotoxin (1 μM), but "rescued"

with addition of K$^+$ channel blockers (4-AP and TEA) which indicates a monosynaptic connection

between POMC^{ARH} and Kiss1^{AVPV} neurons. **(D)** Whole-cell voltage clamp recordings were made

in ChR2-YFP positive cells in brain slices taken from *Kiss1-cre*:Ai32 female mice. Bath application

of melanotan II (MTII, 500 nM) generated an inhibitory outward current in a Kiss1$^{AVPV/PeN}$ neuron

from OVX female. As the cell was synaptically isolated using bath-applied tetrodotoxin (TTX, 1

μM) this represents a direct effect. Washout of MTII while still in TTX quickly led to a return to

baseline RMP. **(E)** An IV relationship was plotted using voltage steps before and after MTII

administration. The crossing at -80 mV (~EK$^+$) indicates that the opening of K+ channels underlie

the MC4R inhibition of Kiss1^{AVPV} neurons. **(F)** In a subset of recordings from OVX brains slices,

the selective membrane estrogen receptor (Gq-mER) agonist STX (10 nM) was added to the bath

for ~10 minutes prior to addition of MTII. STX pretreatment resulted in either a greatly attenuated

outward current or even an inward current as shown in this example. **(G)** The mean outward

current was significantly higher when recording in Kiss1$^{AVPV/Pen}$ neurons from brain slices from

vehicle-treated, OVX females compared to E2-treated, OVX females or acute STX-treated brain

slices from OVX females (One-way ANOVA $F_{(2,19)}$=12.32, p<0.001; Holm-Sidak *posthoc*

comparisons found significant differences between all groups; * p<0.05, *** p<0.001). **(H)** There

was no difference in the frequency of mEPSCs after MTII, calculated as a percent of baseline

between groups. **(I)** However, there was a significant difference in the mEPSC amplitude

(calculated as a percent of baseline) between all groups: main effect $F_{(2,19)}$ =10.58, p<0.001.

Posthoc comparisons using Holm-Sidak found OVX+STX to be different from both OVX (p<0.001)

and OVX+E (p<0.05).

Figure 6. Working model of the neuronal input from POMC ARH to Kiss1 ARH and AVPV

populations.

POMC^{ARH} neurons excite Kiss1^{ARH} neurons via glutamate and αMSH release, which activate

glutamatergic ionotropic receptors and MC4R, respectively. In low E2 state, αMSH fibers project

to the AVPV/PeN, and MC4R inhibit Kiss1$^{AVPV/PeN}$ neurons via opening a K$^+$ channel. However,

in high E2 state, E2 enhances the glutamatergic excitation of both POMC and Kiss1 ARH neurons.

E2-treatment rapidly abrogates (uncouples) the inhibitory MC4R signaling in Kiss1$^{AVPV/PeN}$

neurons, allowing greater excitatory actions of Kiss1^{ARH} neurons via enhanced glutamate release

(26).

References:

1. Health U. Infertility. 2020 [Available from: http://obgyn.ucla.edu/infertility

2. Andermann ML, Lowell BB. Toward a Wiring Diagram Understanding of Appetite Control. Neuron. 2017;95(4):757-78.

3. Cone RD. Studies on the physiological functions of the melanocortin system. Endocr Rev. 2006;27(7):736-49.

4. Shah BP, Vong L, Olson DP, Koda S, Krashes MJ, Ye C, et al. MC4R-expressing glutamatergic neurons in the paraventricular hypothalamus regulate feeding and are synaptically connected to the parabrachial nucleus. Proceedings of the National Academy of Sciences of the United States of America. 2014;111(36):13193-8.

5. Wang K, Mao W, Zhang X, Zhao Y, Fan K, Pan D, et al. Neuroanatomy of melanocortin-4 receptor pathway in the mouse brain. Open Life Sci. 2020;15(1):580-7.

6. Farooqi IS, Keogh JM, Yeo GS, Lank EJ, Cheetham T, O'Rahilly S. Clinical spectrum of obesity and mutations in the melanocortin 4 receptor gene. The New England journal of medicine. 2003;348(12):1085-95.

7. Balthasar N, Dalgaard LT, Lee CE, Yu J, Funahashi H, Williams T, et al. Divergence of melanocortin pathways in the control of food intake and energy expenditure. Cell. 2005;123(3):493-505.

8. Hainerová IA, Zamrazilová H, Sedláčková D, Hainer V. Hypogonadotropic hypogonadism in a homozygous MC4R mutation carrier and the effect of sibutramine treatment on body weight and obesity-related health risks. Obes Facts. 2011;4(4):324-8.

9. Doulla M, McIntyre AD, Hegele RA, Gallego PH. A novel MC4R mutation associated with childhood-onset obesity: A case report. Paediatr Child Health. 2014;19(10):515-8.

10. Batarfi AA, Filimban N, Bajouh OS, Dallol A, Chaudhary AG, Bakhashab S. MC4R variants rs12970134 and rs17782313 are associated with obese polycystic ovary syndrome patients in the Western region of Saudi Arabia. BMC Med Genet. 2019;20(1):144.

11. Chen X, Huang L, Tan HY, Li H, Wan Y, Cowley M, et al. Deficient melanocortin-4 receptor causes abnormal reproductive neuroendocrine profile in female mice. Reproduction (Cambridge, England). 2017;153(3):267-76.

12. Sandrock M, Schulz A, Merkwitz C, Schoneberg T, Spanel-Borowski K, Ricken A. Reduction in corpora lutea number in obese melanocortin-4-receptor-deficient mice. Reprod Biol Endocrinol. 2009;7:24.

13. Cui L, Tan C, Huang L, Wang W, Huang Z, Geng F, et al. Dapagliflozin partially restores reproductive function in MC4R KO obese female mice. The Journal of endocrinology. 2022;254(2):65-76.

14. Manfredi-Lozano M, Roa J, Ruiz-Pino F, Piet R, Garcia-Galiano D, Pineda R, et al. Defining a novel leptin-melanocortin-kisspeptin pathway involved in the metabolic control of puberty. Mol Metab. 2016;5(10):844-57.

15. Goodman RL, Herbison AE, Lehman MN, Navarro VM. Neuroendocrine control of gonadotropin-releasing hormone: Pulsatile and surge modes of secretion. Journal of neuroendocrinology. 2022;34(5):e13094.

16. Israel DD, Sheffer-Babila S, de Luca C, Jo YH, Liu SM, Xia Q, et al. Effects of leptin and
melanocortin signaling interactions on pubertal development and reproduction. Endocrinology.
2012;153(5):2408-19.

17. Manfredi-Lozano M, Roa J, Tena-Sempere M. Connecting metabolism and gonadal
function: Novel central neuropeptide pathways involved in the metabolic control of puberty and
fertility. Front Neuroendocrinol. 2018;48:37-49.

18. Cravo RM, Margatho LO, Osborne-Lawrence S, Donato J, Jr., Atkin S, Bookout AL, et al.
Characterization of Kiss1 neurons using transgenic mouse models. Neuroscience. 2011;173:37-
56.

19. Lam BYH, Williamson A, Finer S, Day FR, Tadross JA, Gonçalves Soares A, et al. MC3R links
nutritional state to childhood growth and the timing of puberty. Nature. 2021;599(7885):436-41.

20. Villa PA, Ruggiero-Ruff RE, Jamieson BB, Campbell RE, Coss D. Obesity alters POMC and
kisspeptin neuron crosstalk leading to reduced luteinizing hormone in male mice. The Journal of
Neuroscience. 2024:e0222242024.

21. Stephens SBZ, Kauffman AS. Estrogen Regulation of the Molecular Phenotype and Active
Translatome of AVPV Kisspeptin Neurons. Endocrinology. 2021;162(9).

22. Moore AM, Prescott M, Campbell RE. Estradiol negative and positive feedback in a
prenatal androgen-induced mouse model of polycystic ovarian syndrome. Endocrinology.
2013;154(2):796-806.

23. McCarthy EA, Dischino D, Maguire C, Leon S, Talbi R, Cheung E, et al. Inhibiting Kiss1
Neurons With Kappa Opioid Receptor Agonists to Treat Polycystic Ovary Syndrome and
Vasomotor Symptoms. J Clin Endocrinol Metab. 2022;107(1):e328-e47.

24. Dewailly D, Barbotin AL, Dumont A, Catteau-Jonard S, Robin G. Role of Anti-Müllerian
Hormone in the Pathogenesis of Polycystic Ovary Syndrome. Frontiers in endocrinology.
2020;11:641.

25. Roa J, Herbison AE. Direct regulation of GnRH neuron excitability by arcuate nucleus
POMC and NPY neuron neuropeptides in female mice. Endocrinology. 2012;153(11):5587-99.

26. Zhang C, Bosch MA, Qiu J, Ronnekleiv OK, Kelly MJ. 17beta-Estradiol increases persistent
Na(+) current and excitability of AVPV/PeN Kiss1 neurons in female mice. Mol Endocrinol.
2015;29(4):518-27.

27. Clements JD, Westbrook GL. Activation kinetics reveal the number of glutamate and
glycine binding sites on the N-methyl-D-aspartate receptor. Neuron. 1991;7(4):605-13.

28. Stincic TL, Grachev P, Bosch MA, Rønnekleiv OK, Kelly MJ. Estradiol drives the
anorexigenic activity of proopiomelanocortin neurons in female mice. eNeuro. 2018;5(4):e0103-
18.2018.

29. Qiu J, Nestor CC, Zhang C, Padilla SL, Palmiter RD, Kelly MJ, et al. High-frequency
stimulation-induced peptide release synchronizes arcuate kisspeptin neurons and excited GnRH
neurons. eLife. 2016;5:e16246.

30. Qiu J, Rivera HM, Bosch MA, Padilla SL, Stincic TL, Palmiter RD, et al. Estrogenic-dependent
glutamatergic neurotransmission from kisspeptin neurons governs feeding circuits in females.
eLife. 2018;7:e35656.

31. Stincic TL, Qiu J, Connors AM, Kelly MJ, Rønnekleiv OK. Arcuate and Preoptic Kisspeptin
neurons exhibit differential projections to hypothalamic nuclei and exert opposite postsynaptic

effects on hypothalamic paraventricular and dorsomedial nuclei in the female mouse. eneuro.
2021:ENEURO.0093-21.2021.
32.   Herman MA, Ackermann F, Trimbuch T, Rosenmund C. Vesicular glutamate transporter
expression level affects synaptic vesicle release probability at hippocampal synapses in culture.
The Journal of Neuroscience. 2014;34(35):11781-91.
33.   Qiu J, Rivera HM, Bosch MA, Padilla SL, Stincic TL, Palmiter RD, et al. Estrogenic-dependent
glutamatergic neurotransmission from kisspeptin neurons governs feeding circuits in females.
Elife. 2018;7.
34.   Thornton JE, Loose MD, Kelly MJ, Rønnekleiv OK. Effects of estrogen on the number of
neurons expressing β-endorphin in the medial basal hypothalamus of the female guinea pig. The
Journal of comparative neurology. 1994;341:68-77.
35.   Petersen SL, Keller ML, Carder SA, McCrone S. Differential Effects of Estrogen and
Progesterone on Levels of POMC mRNA Levels in the Arcuate Nucleus: Relationship to the Timing
of LH Surge Release. Journal of Neuroendocrinology. 1993;5(6):643-8.
36.   Zhang C, Tonsfeldt KJ, Qiu J, Bosch MA, Kobayashi K, Steiner RA, et al. Molecular
mechanisms that drive estradiol-dependent burst firing of Kiss1 neurons in the rostral
periventricular preoptic area. American journal of physiology Endocrinology and metabolism.
2013;305(11):E1384-97.
37.   Mountjoy KG, Mortrud MT, Low MJ, Simerly RB, Cone RD. Localization of the
melanocortin-4 receptor (MC4-R) in neuroendocrine and autonomic control circuits in the brain.
Molecular Endocrinology. 1994;8(10):1298-308.
38.   Yu J, Gimenez LE, Hernandez CC, Wu Y, Wein AH, Han GW, et al. Determination of the
melanocortin-4 receptor structure identifies Ca^{2+} as a cofactor for ligand binding.
Science. 2020;368(6489):428-33.
39.   Sohn JW, Harris LE, Berglund ED, Liu T, Vong L, Lowell BB, et al. Melanocortin 4 receptors
reciprocally regulate sympathetic and parasympathetic preganglionic neurons. Cell.
2013;152(3):612-9.
40.   Kelly MJ, Loose MD, Ronnekleiv OK. Estrogen suppresses mu-opioid- and GABAB-
mediated hyperpolarization of hypothalamic arcuate neurons. The Journal of neuroscience : the
official journal of the Society for Neuroscience. 1992;12(7):2745-50.
41.   Lagrange AH, Ronnekleiv OK, Kelly MJ. The potency of mu-opioid hyperpolarization of
hypothalamic arcuate neurons is rapidly attenuated by 17 beta-estradiol. The Journal of
neuroscience : the official journal of the Society for Neuroscience. 1994;14(10):6196-204.
42.   Qiu J, Bosch MA, Tobias SC, Grandy DK, Scanlan TS, Ronnekleiv OK, et al. Rapid signaling
of estrogen in hypothalamic neurons involves a novel G-protein-coupled estrogen receptor that
activates protein kinase C. The Journal of neuroscience : the official journal of the Society for
Neuroscience. 2003;23(29):9529-40.
43.   Qiu J, Bosch MA, Tobias SC, Krust A, Graham SM, Murphy SJ, et al. A G-protein-coupled
estrogen receptor is involved in hypothalamic control of energy homeostasis. The Journal of
neuroscience : the official journal of the Society for Neuroscience. 2006;26(21):5649-55.
44.   Sabine Hessler XL, Allan E. Herbison. Direct inhibition of arcuate kisspeptin neurones by
neuropeptide Y in the male and female mouse. Journal of Neuroendocrinology. 2020.
45.   Stincic TL, Grachev P, Bosch MA, Ronnekleiv OK, Kelly MJ. Estradiol Drives the
Anorexigenic Activity of Proopiomelanocortin Neurons in Female Mice. eNeuro. 2018;5(4).

46.   Ghamari-Langroudi M, Digby GJ, Sebag JA, Millhauser GL, Palomino R, Matthews R, et al.
G-protein-independent coupling of MC4R to Kir7.1 in hypothalamic neurons. Nature.
2015;520(7545):94-8.
47.   Krause WC, Rodriguez R, Gegenhuber B, Matharu N, Rodriguez AN, Padilla-Roger AM, et
al. Oestrogen engages brain MC4R signalling to drive physical activity in female mice. Nature.
2021;599(7883):131-5.
48.   Brix N, Ernst A, Lauridsen LLB, Parner E, Stovring H, Olsen J, et al. Timing of puberty in
boys and girls: A population-based study. Paediatr Perinat Epidemiol. 2019;33(1):70-8.
49.   Padilla SL, Johnson CW, Barker FD, Patterson MA, Palmiter RD. A Neural Circuit Underlying
the Generation of Hot Flushes. Cell Rep. 2018;24(2):271-7.
50.   Dror T, Franks J, Kauffman AS. Analysis of multiple positive feedback paradigms
demonstrates a complete absence of LH surges and GnRH activation in mice lacking kisspeptin
signaling. Biology of reproduction. 2013;88(6):146.
51.   Steyn FJ, Wan Y, Clarkson J, Veldhuis JD, Herbison AE, Chen C. Development of a
methodology for and assessment of pulsatile luteinizing hormone secretion in juvenile and adult
male mice. Endocrinology. 2013;154(12):4939-45.
52.   Talbi R, Ferrari K, Choi JH, Gerutshang A, McCarthy EA, Dischino D, et al. Characterization
of the Action of Tachykinin Signaling on Pulsatile LH Secretion in Male Mice. Endocrinology.
2021;162(8).
53.   Dave JR, Rubinstein N, Eskay RL. Evidence that b-endorphin binds to specific receptors in
rat peripheral tissues and stimulates the adenylate cyclase-adenosine 3',5'-monophosphate
system. Endocrinology. 1985;117:1389-96.
54.   Naulé L, Robert V, Parmentier C, Martini M, Keller M, Cohen-Solal M, et al. Delayed
pubertal onset and prepubertal Kiss1 expression in female mice lacking central oestrogen
receptor beta. Hum Mol Genet. 2015;24(25):7326-38.
55.   Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR.
Nucleic Acids Res. 2001;29(9):e45.
56.   Madisen L, Mao T, Koch H, Zhuo JM, Berenyi A, Fujisawa S, et al. A toolbox of Cre-
dependent optogenetic transgenic mice for light-induced activation and silencing. Nature
Neuroscience. 2012;15(5):793-802.
57.   Balthasar N, Coppari R, McMinn J, Liu SM, Lee CE, Tang V, et al. Leptin receptor signaling
in POMC neurons is required for normal body weight homeostasis. Neuron. 2004;42(6):983-91.
58.   Padilla SL, Carmody JS, Zeltser LM. Pomc-expressing progenitors give rise to antagonistic
neuronal populations in hypothalamic feeding circuits. Nature Medicine. 2010;16(4):403-5.
59.   Bosch MA, Tonsfeldt KJ, Ronnekleiv OK. mRNA expression of ion channels in GnRH
neurons: subtype-specific regulation by 17beta-estradiol. Molecular and cellular endocrinology.
2013;367(1-2):85-97.
60.   Qiu J, Nestor CC, Zhang C, Padilla SL, Palmiter RD, Kelly MJ, et al. High-frequency
stimulation-induced peptide release synchronizes arcuate kisspeptin neurons and excites GnRH
neurons. Elife. 2016;5.
61.   Stincic TL, Qiu J, Connors AM, Kelly MJ, Ronnekleiv OK. Arcuate and Preoptic Kisspeptin
Neurons Exhibit Differential Projections to Hypothalamic Nuclei and Exert Opposite Postsynaptic
Effects on Hypothalamic Paraventricular and Dorsomedial Nuclei in the Female Mouse. eNeuro.
2021;8(4).

62.   Qiu J, Stincic TL, Bosch MA, Connors AM, Kaech Petrie S, Ronnekleiv OK, et al. Deletion of
Stim1 in Hypothalamic Arcuate Nucleus Kiss1 Neurons Potentiates Synchronous GCaMP Activity
and Protects against Diet-Induced Obesity. The Journal of neuroscience : the official journal of
the Society for Neuroscience. 2021;41(47):9688-701.

1056

A. Kiss1 neuron

B. MT II

20 pA
20 s

C.
MTII Current (pA)
OVX   OVX+E

D. THIQ

10 pA
1 m

E. Pomc^Cre Injected
3V
200μm

F.
470 nm
POMC → Kiss1 ARH

G.
Recorded Cells   TC
MM 1 2 3 4 5   - +
400
200
100
Kiss1 120 bp

H.
5 ms   2) TTX

10 pA
20 ms
3) TTX+4AP +TEA
1) Baseline

I.
5 ms
5 pA   OVX
5 pA   OVX+E

J.
Paired Pulse Ratio
OVX   OVX+E

K.
10 s, 20 Hz
20 pA

L.
10 s, 20 Hz   SHU 9119 Pretreatment
20 pA

M.
10 s, 20 Hz   Naloxone
20 pA   1 m