POMC neurons control fertility through differential signaling of MC4R in Kisspeptin neurons in the female mouse

Authors:
Rajae Talbi[1,2*], Todd L. Stincic[4,*#], Kaitlin Ferrari[2], Choi Ji Hae[2], Karol Walec[2,] Elizabeth Medve[2], Achi Gerutshang[2], Silvia León[1,2], Elizabeth A. McCarthy[1,2], Oline K. Rønnekleiv [4,5], Martin J. Kelly[4,5] and Víctor M. Navarro[1,2,6]

[1]Harvard Medical School, Boston, MA, USA; [2]Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA; [4]Department of Chemical Physiology and Biochemistry, Oregon Health & Science University, Portland, OR, USA; [5]Division of Neuroscience, Oregon National Primate Research Center, Beaverton, OR, USA; [6]Harvard Program in Neuroscience, Boston, MA, USA

*First authors contributed equally
#Present address: Department of Biology, Appalachian State University, Boone, NC 28608.

Correspondence:
Victor M. Navarro, PhD, Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, 221 Longwood Avenue, Room 219, Boston, Massachusetts 02115. E-mail: vnavarro@bwh.harvard.edu.

Martin J. Kelly, PhD, Department of Chemical Physiology and Biochemistry, Oregon Health & Science University, 3181 SW Sam Jackson Park Road, Portland, OR 97239. E-mail: kellym@ohsu.edu .

Authors contribution: RT, TLS, MJK and VMN conceived and designed the study; RT, TLS, KF, CJH, EM, AG, KW, SL, EAM, OKR performed the experiments. RT, TLS, OKR, MJK and VMN analyzed data. RT, TLS and VMN wrote the manuscript with input from all the authors.

Competing Interests Statement: The authors declare they have no competing interest.

Data and Materials Availability: All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials.

Keywords: Kiss1 neurons, MC4R, POMC, Fertility, Obesity

Abstract

Inactivating mutations in the melanocortin 4 receptor (*MC4R*) gene cause monogenic obesity. Interestingly, female patients also display various degrees of reproductive disorders, in line with the subfertile phenotype of MC4RKO female mice. However, the cellular mechanisms by which MC4R regulates reproduction are unknown. Kiss1 neurons directly stimulate gonadotropin-releasing hormone (GnRH) release through two distinct populations; the Kiss1^{ARH} neurons, controlling GnRH pulses, and the sexually dimorphic Kiss1$^{AVPV/PeN}$ neurons controlling the preovulatory LH surge driving ovulation. In the present study, we show that *Mc4r* is expressed within Kiss1^{ARH} and Kiss1$^{AVPV/PeN}$ neurons. *In vivo*, deletion of MC4R from Kiss1 neurons in female mice replicates the reproductive impairments of MC4RKO mice without inducing obesity. Conversely, reinsertion of MC4R in Kiss1 neurons of MC4R null mice restores estrous cyclicity and LH pulsatility without reducing their obese phenotype. *In vitro*, we show that MC4R activation excites Kiss1^{ARH} neurons through direct synaptic actions. In contrast, Kiss1$^{AVPV/PeN}$ neurons are normally inhibited by MC4R activation except under elevated estradiol levels, thus facilitating the activation of Kiss1$^{AVPV/PeN}$ neurons to induce the preovulatory LH surge driving ovulation in females. Our findings demonstrate that POMC^{ARH} neurons acting through MC4R, directly regulate reproductive function in females by stimulating the "pulse generator" activity of Kiss1^{ARH} neurons and restricting the activation of Kiss1$^{AVPV/PeN}$ neurons to the time of the estradiol-dependent LH surge, and thus unveil a novel pathway of metabolic regulation of fertility.

Significance:

Women with inactivating mutations in proopiomelanocortin (*POMC*) or melanocortin receptor 4 (*MC4R*) genes are severely obese and display various degrees of reproductive disorders; however, the mechanisms by which MC4R regulates reproduction remain unknown. Here, we show that the melanocortins, known for their fundamental role in the control of energy balance, also play a direct role in reproduction by modulating the tonic and surge-like release of GnRH/LH via direct actions on MC4R of Kiss1 neurons. These findings identify a direct link between the neural regulation of metabolism and fertility and offer a new platform for the development of novel approaches to treat reproductive deficiencies derived from metabolic impairments.

Introduction

Obesity rates have skyrocketed in Western societies in the last decades, resulting in an alarming rise in comorbidities that place a significant burden on healthcare systems. The increase in obesity correlates with a decrease in fertility rates, leading to conception challenges that are experienced by approximately 15% of couples in the United States currently (1).

The melanocortin receptor 4 (MC4R) binds α-melanocyte stimulating hormone (αMSH), an agonist product of the pro-opiomelanocortin (*Pomc*) gene, and the inverse agonist of the agouti-related peptide (AgRP) to regulate food intake and energy expenditure (2, 3). While the role of MC4R on food intake is largely mediated by neurons located in the paraventricular nucleus of the hypothalamus (PVN) (4), its expression in the brain is widespread (5) with the specific function of the different MC4R-expressing neurons in areas beyond the PVN remaining to be fully explored. Inactivating mutations in *MC4R* are a leading cause of monogenic obesity in humans and mice (6). Scant evidence in humans shows an association between *MC4R* mutations and higher incidence of hypogonadotropic hypogonadism (HH) (7), alterations in the timing of puberty onset (8), and polycystic ovary syndrome (PCOS) (9); however, conflicting reports exist (6). Therefore, the role of MC4R signaling in reproductive function in humans remains controversial despite the clear association found in mice. Supporting this role of MC4R, *Mc4r* null mice display an array of reproductive abnormalities that largely affects females, characterized by irregular estrous cycles, disrupted luteinizing hormone (LH) secretion, reduced corpora lutea and reduced fertility (10-12). Further evidence from mice demonstrates that MC4R agonists robustly increase LH release in a kisspeptin-dependent manner (13).

Kisspeptin is the most potent gonadotropin releasing hormone (GnRH) secretagogue known to date and it is mainly produced in two distinct neuronal populations. Kiss1 neurons of the arcuate nucleus (Kiss1^{ARH}) neurons control the pulsatile (tonic) release of GnRH (present in both sexes), and sex steroids attenuate their release of kisspeptin. However, while testosterone inhibits

Kiss1[ARH] glutamate release (14), estradiol enhances the vesicular-glutamate transporter 2

(*Vglut2)* expression and glutamate release probability which could reflect changing roles of

Kiss1[ARH] neurons across the estrous cycle (15). Kiss1 neurons of the anteroventral periventricular

continuum area (Kiss1[AVPV/PeN]), which are predominantly present in females, are responsible for

generating the preovulatory GnRH/LH surge (16). Despite these critical roles of both populations

of Kiss1 neurons for reproduction, the underlying mechanisms that determine how each Kiss1

population responds differently to sex steroids remains unresolved. Fibers from POMC neurons

in the arcuate nucleus (POMC[ARH]) project to and juxtapose Kiss1[ARH] neurons (13); and

melanocortin signaling through MC4R contributes to the permissive role of leptin on puberty onset

(13, 17, 18). Altogether, the existing data in rodents suggests a clear role for MC4R, likely on

Kiss1 neurons, in the control of reproductive maturation and fertility.

In this study, we investigated the contribution of MC4R signaling versus obesity per se in the

etiology of the reproductive impairments observed in *Mc4r* null mice, which could explain similar

impairments observed in humans.

Results

***Mc4r* is expressed within Kiss1 neurons and determines the timing of puberty onset in**

females.

As the expression of *Mc4r* in Kiss1 neurons has been inconclusive in previous studies, we began

by using the highly sensitive RNAscope *in situ* hybridization method. Dual fluorescence labeling

of *Mc4r* and *Kiss1* probes showed that 84% of Kiss1[ARH] neurons and 74% of Kiss1[AVPV/PeN] neurons

expressed *Mcr4* (**Fig. 1*A, B***). To further investigate the specific role of MC4R in Kiss1 neurons,

we generated a Kiss1-specific MC4R knockout model (Kiss1-MC4RKO). The specific deletion of

Mc4r from Kiss1 neurons, as well as the absence of global recombination, were confirmed through

RNAscope. While Kiss1 neurons of Kiss1-MC4RKO mice lack *Mc4r* transcript compared to their

control littermates (**Fig. S1A**), *Mc4r* was detectable in the PVN of all mice, which is a major site

of *Mc4r* expression in the brain, therefore supporting the specific deletion of *Mc4r* only within

Kiss1 neurons (**Fig. S1B**).

Puberty onset was assessed daily from weaning age through the monitoring of vaginal opening

(VO) and first estrus (FE). Kiss1-MC4RKO females showed a significant advancement in the age

of VO (P=0.0150, Kiss1-MC4RKO: 24.67± 0.3978 vs controls: 26.32±0.5301) and FE (P=0.0341,

Kiss1-MC4RKO: 32.63±1.489 vs controls: 37.84±1.903) compared to their littermate controls

(**Fig. 1C, D**). Body weight of Kiss1-MC4RKO females at the age of puberty onset was similar

between groups (P=0.2596, Kiss1-MC4RKO: 12.41±0.3041 vs controls: 12.89±0.2631) (**Fig. 1E**).

These data suggest that melanocortin signaling on Kiss1 neurons participates in the timing of

puberty onset in females. Since the role of the melanocortin system on puberty onset is largely

unexplored, we evaluated the expression profile of the main components of this system in the

hypothalami of female mice at postnatal days (PND) 10, 15, 22 and 30. Interestingly, expressions

of *Agrp*, *Mc3r* and *Mc4r* were significantly lower at the time of puberty onset (PND30) in Kiss1-

MC4RKO females compared to earlier developmental ages (**Fig. 1F**). This suggests that a

decrease in the hypothalamic melanocortin tone drives puberty onset, in line with the

advancement in the age of VO and FE observed in Kiss1-MC4RKO female mice (**Fig. 1C, D**).

MC4R in Kiss1 neurons is required for female reproduction.

Because Kiss1-MC4RKO females showed altered puberty onset, we further investigated their

reproductive phenotype. Interestingly, Kiss1-MC4RKO female mice displayed normal BW

throughout the time of the study (up to PND150) **(Fig. 2A)**; however, they presented with irregular

estrous cycles with predominantly more time spent in diestrus (P=0.0001) and less time spent in

estrus (P<0.0001) than their control littermates (**Fig. 2B, C**). The assessment of the pulsatile

release of LH every 10 min over 180 min revealed no change in the total number of pulses

between groups but a significant increase in basal LH release (P=0.0368) in Kiss1-MC4RKO

female mice (0.426±0.036) compared to controls (0.339±0.026) (**Fig. 2D-H**). The analysis of the

gene expression of the "KNDy" systems in the ARH, which control the GnRH pulse generator

(16), revealed normal expression levels of *Kiss1*, *Tac2* and *Tacr3*, but significantly lower

expression of *Pdyn* (P=0.0067, Kiss1-MC4RKO: 0.781±0.0384 vs controls: 1.000±0.0190) (**Fig.**

2I-L). The conserved expression of *Kiss1*, *Tac2* and *Tacr3* correlates with the preserved LH pulse

frequency and amplitude in Kiss1-MC4RKO mice, while the lower inhibitory tone of dynorphin

(**Fig. 2I**) correlates with the higher basal LH levels (**Fig. 2G**). To assess the contribution of MC4R

signaling in Kiss1 neurons to the induction of the preovulatory LH surge, Kiss1-MC4RKO females

and control littermates were submitted to an LH surge induction protocol that showed the LH

surge was significantly blunted in the Kiss1-MC4RKO females (P=0.0091), while the protocol

clearly evoked the expected afternoon rise of LH in control mice (**Fig. 2M**). In line with these

findings, the ovaries of Kiss1-MC4RKO females displayed fewer corpora lutea (P=0.0054, Kiss1-

MC4RKO: 0.800±0.374 vs controls: 2.80±0.374) and increased cystic follicles (P=0.0337, Kiss1-

MC4RKO: 1.60±0.400 vs controls: 0.400±0.244) (**Fig. 2N-P**), which correlate with decreased

fertility as observed by the extended time to deliver pups (*i.e.* longer time to get pregnant)

(P=0.0030, Kiss1-MC4RKO: 30.30±4.600 vs controls: 20.88±0.2266), and fewer pups per litter

(P=0.0095, Kiss1-MC4RKO: 6.667±0.5528 vs controls: 8.444±0.242) (**Fig. 2S, T**).

The increase in serum LH levels and decreased ovulation observed in the Kiss1-MC4RKO

females is reminiscent of polycystic ovary syndrome (PCOS) mouse models (19, 20). Thus, we

analyzed circulating levels of testosterone (T) and anti-müllerian hormone (AMH), which are

frequently elevated in PCOS models (21). The Kiss1-MC4RKO females expressed normal T and

AMH levels compared to control mice in diestrus (**Fig. 2Q, R**). Thus, we can exclude a PCOS-like

reproductive phenotype mediated by the lack of melanocortin signaling on Kiss1 neurons.

Re-insertion of MC4R in Kiss1 neurons of MC4RKO mice improves reproductive function.

Kiss1-MC4RKO females displayed reproductive abnormalities resembling those described in MC4RKO females (10-12). Thus, we hypothesized that the reproductive defects described for the MC4RKO mice would be, at least in part, due to the absence of MC4R signaling in Kiss1 neurons. To further investigate this hypothesis, we generated mice that do not express *Mc4r* anywhere (MC4R-LoxTB, *i.e.* MC4RKO) or that express *Mc4r* only in Kiss1 neurons (*Kiss1-cre:MC4R-loxTB*). The re-insertion of *Mc4r* within Kiss1 neurons in the *Kiss1-cre:MC4R-loxTB* mice was confirmed through RNAscope (**Fig. S2A).** *Mc4r* expression was not detected in the PVN of these mice (**Fig. S2B**), and it was only detected in Kiss1 neurons in the *Kiss1-cre:MC4R-loxTB* mice. Puberty onset was assessed daily from weaning age through the monitoring of vaginal opening (VO) and first estrus (FE). MC4R-LoxTB and *Kiss1-cre:MC4R-loxTB* female mice displayed normal timing of puberty onset compared to their control littermates, as assessed by VO (P=0.104, MC4R-LoxTB: 30.86±1.908 vs *Kiss1-cre:MC4R-loxTB: 30.09*±1.004 vs controls: 27.57±0.8168), and FE (P=0.8472, MC4R-LoxTB: 35.14±2.463 vs *Kiss1-cre:MC4R-loxTB: 34.55*±1.648 vs controls: 35.80±0.8406) (**Fig. 3A**), despite displaying significantly higher body weight at the time of puberty onset (P=0.0001, MC4R-LoxTB: 15.49±0.5078 vs *Kiss1-cre:MC4R-loxTB:* 15.54±0.40 vs Controls: 13.29±0.30) (**Fig. 3B**). As expected, MC4R-LoxTB (MC4RKO) displayed increased body weight (**Fig. 3C**) and an array of reproductive impairments that included: irregular estrous cycles with significantly more days in diestrus (P=0.0016) and fewer days in estrus (P=0.0317) compared to controls (**Fig. 3D, E**), significantly higher serum LH levels characterized by higher basal (P=0.0882, MC4R-LoxTB: 0.48±0.02 vs Controls: 0.37±0.06) and amplitude levels per LH pulse (P=0.0102, MC4R-LoxTB: 0.73±0.009 vs Controls: 0.58±0.07) despite unaltered number of LH pulses/180min (P=0.2285, MC4R-LoxTB: 2.00±0.40 vs Controls: 3.50±0.50) (**Fig. 3F-J**), and fewer corporal lutea (P=0.0366, MC4R-LoxTB: 0.80±0.80 vs Controls: 4.20±1.15) (**Fig. 3K, L**), recapitulating the same reproductive phenotype observed in Kiss1-MC4RKO mice despite the differences in BW. Re-introduction of MC4R into Kiss1 neurons in *Kiss1-cre:MC4R-loxTB* mice

completely recovered estrous cyclicity, which was similar to controls (diestrus: P=0.1932,

proestrus: P=0.8262, estrus: P=0.0547, compared to controls) (**Fig. 3D, E)** despite *Kiss1-*

cre:MC4R-loxTB mice showing the same degree of obesity as MC4R-LoxTB mice (**Fig. 3C**),

indicating that obesity *per se* was not mediating the irregular estrous cycles in MC4RKO mice. As

indicated above, MC4R-LoxTB mice display higher overall circulating LH levels, and this feature

was also recovered by the insertion of MC4R in Kiss1 neurons of *Kiss1-cre:MC4R-loxTB* mice

(basal LH: 0.32±0.044, LH amplitude: 0.48±0.04, LH pulses/180min: 2.40±0.67 and AUC:

70.62±7.156 compared to controls: 80.36±9.058 compared to controls) (**Fig. 3G-I**), further

confirming that melanocortin action on Kiss1 neurons is required for the normal control of

gonadotropin release. Despite these significant improvements in reproductive function, *Kiss1-*

cre:MC4R-loxTB mice presented fewer corpora lutea than controls (0.80±0.80), similar to MC4R-

LoxTB mice, suggesting that an ovulatory impairment still persists (**Fig. 3K, L**). While tonic LH

release and estrous cyclicity are predominantly controlled by Kiss1^{ARH} neurons, ovulation is

mediated by the induction of the preovulatory LH surge by Kiss1$^{AVPV/PeN}$ neurons. It is possible

that the metabolic signals derived from their obese phenotype, and/or the absence of the direct

action of MC4R in GnRH neurons (17, 22) prevents the complete recovery of ovulation in *Kiss1-*

cre:MC4R-loxTB mice. Both genetic models displayed significantly lower T (MC4R-LoxTB:

28.50±6.20 vs *Kiss1-cre:MC4R-loxTB:* 31.22±3.90 vs Controls: 62.84±16.53) and AMH (MC4R-

LoxTB: 150.6±22.00 vs *Kiss1-cre:MC4R-loxTB:* 146.9±10.83 vs Controls: 321.8±15.70) levels

than control mice in diestrus (**Fig. 3M, N**). Therefore, we can exclude a PCOS-like reproductive

phenotype mediated by the lack of melanocortin signaling.

Kiss1^{ARH} neurons are excited by MC4R agonists.

Based on the RNAscope and reproductive phenotype data, we hypothesized that Kiss1^{ARH}

neurons are direct targets of melanocortins and would respond to MC4R activation. Initially we

did whole-cell, voltage clamp recordings in Kiss1^{ARH} neurons (**Fig. 4A**). *Kiss1-Cre* x Ai32 or *Kiss1-*

Cre AAV-injected mice underwent OVX and were given E2 replacement (see Methods). We

targeted fluorescent cells for recording which were isolated synaptically by bathing the slices in

tetrodotoxin (TTX, 1 µM). Focal application of the high-affinity melanocortin receptor agonist

melanotan II (MTII, ~250 nM) elicited a small inward current in half of the isolated Kiss1^{ARH} cells

(10/21) (**Fig. 4B**). Next, we examined if the E2 state affected MTII response by recording from

Kiss1^{ARH} neurons from OVX females and found two thirds (6/9 cells) responded, but there was no

significant difference in the average current (**Fig. 4C**). Finally, it must be noted that at this

concentration MTII is a non-selective MCR agonist, so we followed up using a perfusion of the

MC4R-selective agonist THIQ (100 nM) to determine if activation of MC4Rs alone was sufficient

to invoke a postsynaptic response in Kiss1^{ARH} neurons from OVX+E2 females. Similar to MTII,

THIQ was able to elicit an excitatory inward current in Kiss1^{ARH} neurons (4/5 cells, -7.4±0.5 pA),

indicating MC4R activation is sufficient (**Fig. 4D**).

POMC neurons synapse directly with Kiss1^{ARH} neurons.

Based on the pharmacological activation of MCRs, we wanted to address whether POMC^{ARH}

neurons are the source of melanocortins to excite Kiss1^{ARH} neurons directly. To answer this

question, we injected an AAV-EF1α-DIO-ChR2:mCherry vector into the ARH of adult *Pomc-Cre*

female mice (**Fig. 4E**) (24). After 2-4 weeks we did whole cell recordings from putative Kiss1^{ARH}

neurons and looked for a response to high frequency optogenetic stimulation of POMC fibers in

OVX + E2-treated females (**Fig. 4F**). We were able to confirm that these were Kiss1 neurons

based on the presence of a persistent sodium current and/or *post hoc* identification by scRT-PCR

(41/69 cells, **Fig. 4G**). While this current is more prevalent in the AVPV Kiss1 population, Kiss1

neurons are the only ARH neurons to display this electrophysiological "fingerprint" of a

pronounced INaP paired with a high capacitance and low input resistance (23). Additionally, while

recording from putative Kiss1^{ARH} neurons, we optogenetically stimulated POMC fibers at low

frequency (1 Hz, 5 ms, 470 nm light) and recorded fast postsynaptic inward currents. The fast

kinetics of the EPSC are also a tell-tale sign of an ionotropic glutamatergic response (24).
Previously, CNQX was sufficient to block similar excitatory ESPCs in other postsynaptic targets
of POMC neurons (25). While the consistent latency from optogenetic stimulus and current
response was indicative of a single synaptic delay, we wanted to establish that it was a direct
synaptic connection, as we have previously shown for output of POMC neurons (25) and Kiss1
neurons (26-28). First, we abrogated the optogenetic response with the addition of TTX (1 µM)
to the bath, we were able to rescue the postsynaptic glutamate response with the addition of the
K^+ channel blockers 4-aminopyridine (4-AP; 0.5 mM) and tetraethyl ammonium (TEA; 7.5 mM)
(**Fig. 4H**). K+ channel block enables the calcium inflow from ChR2 in the presynaptic terminals
to be sufficient to restore vesicle fusion. Therefore, there appears to be no intervening synapses
between POMC and the downstream Kiss1^{ARH} neurons.
Previously we found that E2 increases glutamate release from POMC neurons by increasing
Vglut2 (25). Therefore, we compared the postsynaptic glutamatergic responses following two
optogenetic stimuli (50 ms interval between light flashes) in Kiss1^{ARH} neurons from OVX + vehicle
versus OVX + E2-treated female mice (**Fig. 4I**). As anticipated, we found that treatment with E2
increased the probability of glutamate release from POMC neurons onto Kiss1^{ARH} neurons based
on the significant decrease in the paired pulse ratio of the two stimuli (29) (**Fig. 4J**). Although in
the present study we used an *in vivo* treatment paradigm, we know from previous studies that this
augmentation of glutamate release can happen quite rapidly after a brief exposure to E2 *in vitro*
(within 15 min) (25). Therefore, Kiss1^{ARH} neurons are excited by the glutamatergic input from
POMC neurons in an E2-dependent manner.
Finally, given that POMC neurons project directly onto Kiss1^{ARH} neurons, we wanted to show that
the evoked slow inward current with high frequency stimulation was mediated by αMSH. Indeed,
high frequency stimulation evoked an excitatory inward current (**Fig. 4K**) that was blocked by the
selective MC3/4R antagonist SHU 9119 (**Fig. 4L**). The number of high frequency (peptidergic)
responses seen in Kiss1 neurons that displayed low frequency postsynaptic (glutamatergic)

currents was low (4/16, 25%, mean inward current: -1.4±3.1 pA). However, since these recordings

were made in brain slices from E2-treated, OVX females and β-endorphin expression is enhanced

by E2 treatment (30), we were concerned that co-release of the opioid peptide could obscure

melanocortin postsynaptic effects. Indeed, there was a notable increase in slow inward currents

generated by high frequency stimulation of POMC fibers in slices that were perfused with the non-

selective opioid antagonist naloxone (1 µM) (9/23, 40%, mean inward current: -7.6±2.9 pA) (**Fig.**

4M).

Together these optogenetic findings reinforced our pharmacological results showing that Kiss1[ARH]

neurons are excited by MTII, and THIQ. Additional studies would need to be done to identify the

cation conductance that is affected by the MC4R signaling cascade in Kiss1[ARH] neurons, but

clearly we have established that there are excitatory glutamatergic and peptidergic inputs from

POMC to Kiss1[ARH] neurons.

Kiss1[AVPV/PeN] neurons are inhibited by MC4R agonists.

Having established a direct, excitatory projection from POMC to Kiss1[ARH] neurons, we next

examined POMC inputs to Kiss1[AVPVPeN] neurons. First, we used immunocytochemistry to label

fibers expressing αMSH to demonstrate that POMC fibers densely innervated the AVPV/ PeN

area (**Fig. 5A**). We hypothesized that Kiss1[AVPV/PeN] neurons would be excited by exogenous

application of melanocortins because the receptor is typically Gs-coupled (31, 32). We did whole-

cell, voltage clamp recordings from Kiss1[AVPV/PeN]-Cre::Ai32 neurons, which were isolated

synaptically by bathing the neurons in TTX (1 µM), from OVX females. For these experiments,

we tested the response to the high affinity melanocortin receptor agonist MTII (500 nM).

Surprisingly, we consistently measured an outward (inhibitory) current that could be reversed on

washout of MTII (**Fig. 5B**). This outward current was associated with an increase in a K$^+$

conductance based on the current-voltage plot (*i.e.,* the outward current reversed at ~E$_{K+}$, **Fig.**

5C). Also, the inwardly-rectifying I/V could indicate that MC4R is coupled to activation of inwardly-

rectifying K$^+$ channels as has been previously reported for MC4R signaling in pre-autonomic

parasympathetic neurons in the brainstem (33). We hypothesized that the coupling could be

modulated by E2 via a Gα_q-coupled membrane estrogen receptor (Gq-mER) as we have

previously demonstrated in POMC neurons (34-36). For these experiments, we again utilized

OVX females and targeted Kiss1$^{AVPV/PeN}$-Cre::Ai32 neurons. Once in the whole-cell voltage-clamp

configuration, we perfused the slices with STX (10 nM), a selective ligand for the putative Gq-

mER (36), and tested the response to the MTII. Indeed, the outward current (inhibitory) response

to MTII was completely abrogated and even reversed by STX (**Fig. 5D, E**). The short-term (bath)

treatment with STX ensured that there was no desensitization of the Gq-mER with longer-term (*in*

vivo) treatment with E2. Therefore, estrogen receptor activation can rapidly uncouple (*i.e.,*

desensitize) the MC4R inhibitory response in Kiss1$^{AVPV/PeN}$ neurons. We would predict that the

intracellular signaling cascade for the heterologous desensitization is similar to what we have

elucidated in POMC neurons (36), but this will need to be determined in future experiments. We

next investigated the response in vehicle-treated, OVX females. Indeed, Kiss1$^{AVPV/PeN}$ neurons

were even more inhibited by the same MTII exposure (**Fig. 5E**). Therefore, in contrast to Kiss1^{ARH}

neurons, the MC4R appears to be coupled to opening of K$^+$ channels in Kiss1$^{AVPV/PeN}$ neurons. To

further establish whether the MTII was having pre- or postsynaptic effects, we measured the

glutamatergic mEPSCs before and after melanocortin receptor activation. We found there was no

significant effect on the frequency (**Fig. 5F**), but there was an effect on the amplitude (**Fig. 5G**).

This further supports a postsynaptic locus of STX's effects. However, *posthoc* comparisons found

STX increased amplitude compared to both OVX and OVX+E. This result hints at a membrane

delimited effect as STX is a much more potent agonist for Gq-mER than E2 (37).

Discussion

Our findings show that melanocortins act directly on Kiss1 neurons to contribute to the metabolic

regulation of fertility, in line with previous reports showing that αMSH stimulates LH release in a

kisspeptin-dependent manner (13). Several studies in humans have linked *MC4R* mutations to

reproductive abnormalities, including precocious puberty (8), PCOS (9), and hypogonadism (7).

However, a direct association between *MC4R* mutations and reproductive function, independent

from the obese condition of these patients, has not been identified (6). In the current study, we

demonstrate that the majority of Kiss1 neurons express the *Mc4r* transcript, and we show that the

reproductive impairments observed in MC4R deficient mice, which replicate many of the

conditions described in humans, are largely mediated by the direct action of melanocortins via

MC4R on Kiss1 neurons and not to their obese phenotype. This is because the ablation of MC4R

from Kiss1 neurons largely replicated the reproductive impairments observed in MC4RKO female

mice without inducing obesity, and the selective reinsertion of MC4R into Kiss1 neurons of

MC4RKO mice significantly improved their reproductive function without changing their obese

phenotype. Strikingly, puberty onset was advanced in Kiss1-MC4RKO females. Our findings

revealed a low melanocortin tone in the hypothalamus of females during pubertal development,

characterized by lower levels of *Agrp, Mc4r* and *Mc3r* expression at the time of puberty onset

(~PND30). This developmental decline in the expression of melanocortin receptors is in line with

the advanced timing of puberty onset in females when MC4R is congenitally ablated from Kiss1

neurons. A recent study using whole body MC3RKO mice showed a trend to delayed puberty

onset in both sexes (38), suggesting that MC3R, which is also expressed in Kiss1 neurons, may

be more critical for the timing of puberty onset in males. However, in females the combination of

MC3R and MC4R action may be necessary for pubertal development. Interestingly, the Kiss1-

MC4RKO and *Kiss1-cre:MC4R-LoxTB* mice do not show an advancement in puberty onset as we

observed in Kiss1-MC4RKO mice; however, these mice show a significant increase in body

weight at the time of puberty onset already. We can therefore speculate that the increased

adiposity is masking the specific effect of MC4R ablation from Kiss1 neurons on puberty, that we

see in the non-obese Kiss1-MC4RKO female mice. Our findings presented in this study do not

rule out the existence of additional sites of action of MC4R in other neuronal populations to

regulate reproduction, *e.g.,* GnRH neurons, which also express *Mc4r* (17, 22). Indeed, the lack of

full recovery of the reproductive function in *Kiss1-cre:MC4R-LoxTB* mice suggests that MC4R in

Kiss1 neurons is necessary but not sufficient to achieve full reproductive capabilities.

In addition to early puberty onset, Kiss1-MC4RKO females displayed an impaired preovulatory

LH surge driving ovulation and had fewer corpora lutea in their ovaries compared to control

littermates, further supporting a role of MC4R in regulating ovulation. In adulthood, the

reproductive phenotype observed in Kiss1-MC4RKO and MC4R-loxTB female mice (increased

LH, irregular estrous cycles, oligo-ovulation, increased cystic follicles) correlates with the

phenotype observed in PCOS mouse models (19, 20). In fact, an association between *MC4R*

mutations and PCOS has been reported in patients (9). However, our mouse models failed to

display higher levels of circulating androgens or AMH, two of the hallmarks of PCOS (21),

suggesting that melanocortin signaling is unlikely to be a main contributing factor in the

development of this syndrome.

Our data revealing early puberty onset and impaired LH pulse and surge driving ovulation support

a direct role of MC4R on the two primary Kiss1 populations regulating these functions (ARH and

AVPV/PeN). We further investigated this and found that MC4R signaling excites Kiss1^{ARH} neurons

and inhibits Kiss1$^{AVPV/PeN}$ neurons in an estradiol-dependent manner. In Kiss1^{ARH} neurons, bath

applied or optogenetically-evoked release of melanocortin agonists induce a direct excitatory

inward current. Our studies using whole cell voltage clamp in female mice were able to detect a

melanocortin-mediated inward current in Kiss1^{ARH} neurons that would increase excitability.

Previously, an MCR-activation effect was not detected in cell-attached recordings of diestrus

females (13). However, cell-attached recordings do not measure a direct response in isolated

Kiss1^{ARH} neurons but rather the summation of multiple synaptic inputs to Kiss1^{ARH} neurons. Most

importantly, we did not see a difference in the excitatory response in synaptically isolated Kiss1^{ARH}

neurons from E2-treated versus vehicle-treated females. This emphasizes the importance of

doing whole-cell recording from isolated Kiss1 neurons. The modulatory actions of POMC^{ARH}

neuronal projections to Kiss1^{ARH} neurons can also occur through a) the excitatory action of

glutamate in a process that is facilitated in the presence of estradiol, in line with previous

publications (15, 39), and b) the inhibitory action of β-endorphin, suggesting that POMC^{ARH}

neurons are able to exert a precise regulatory role of the GnRH pulse generator, *i.e.* Kiss1^{ARH}

neurons, in response to metabolic challenges. In the AVPV/PeN, MC4R agonists inhibited

Kiss1$^{AVPV/PeN}$ neurons in an estradiol-dependent manner, which utilized a Gα$_q$-coupled membrane

estrogen receptor (STX receptor) to attenuate the inhibitory tone of MC4R in these neurons. To

date, two instances of MC4R-mediated inhibition have been described in the central nervous

system: MC4R can Gi,o couple to open K+ (Kir6.2) channels in parasympathetic preganglionic

neurons in the brain stem (33), and MC4R can directly couple to K+ (Kv7.1) channels in the

hypothalamic paraventricular nucleus neurons (40). Based on the robust αMSH labeling, we

believe that POMC projections represent a significant input to Kiss1$^{AVPV/PeN}$ neurons. Interestingly,

our findings describe an inhibitory role of MC4R in the reproductive neuroendocrine axis. This

steroid state dependent effect is particularly relevant because in the context of reproduction, the

two populations of Kiss1 neurons are strikingly different in their response to estradiol, where

Kiss1^{ARH} and Kiss1$^{AVPV/PeN}$ neurons are inhibited or stimulated, respectively, in order to mount the

negative versus positive feedback of sex steroids (16). This differential regulation allows for the

episodic versus surge release of GnRH; however, the cellular mechanisms underlying these

opposing roles of Kiss1 neurons to the same stimulus, *i.e.* circulating E2 levels, remain unknown.

Although there were no differences in the magnitude of the electrophysiological responses to

MTII in Kiss1^{ARH} neurons in vehicle- versus E2-treated, ovariectomized females, we cannot rule

out that E2 affects other systems activated by the MC4R signaling cascade as recently reported

in MC4R expressing neurons of the ventromedial nucleus (VMH) (41). Interestingly, puberty onset

was advanced in Kiss1-MC4RKO females. This data suggests that the inhibitory tone of MC4R

signaling on Kiss1$^{AVPV/PeN}$ neurons might also be involved in the timing of puberty onset, which in

turn suggests a role for this female-specific population of Kiss1 neurons in sexual maturation. This

phenomenon could explain the earlier age of puberty onset frequently seen in females of most

mammalian species compared to their male counterparts (42). Furthermore, our findings suggest

that metabolic cues, through the regulation of the melanocortin output onto Kiss1$^{AVPV/PeN}$ neurons,

are essential for the timing and magnitude of the GnRH/LH surge. Future studies are warranted

to fully elucidate synaptic input and postsynaptic cascades in the POMC-Kiss1$^{AVPV/PeN}$ circuit.

Altogether, our data reveal a differential regulatory action of MC4R in the neural control of GnRH

release, participating in both the surge (E2-treated, ovariectomized female) and the pulse-like

(ovariectomized female) modes of LH release through the cellular regulation of Kiss1 neurons

(**Fig. 6**), which translate into our findings on the conditional knockout of *Mc4r* from Kiss1 neurons.

These findings are important because the reproductive abnormalities often attributed to obesity

in MC4R deficient patients may not be caused by the excess in body weight but, at least in part,

by a deficiency in MC4R signaling directly at the level of Kiss1 neurons that affects predominantly

the ability to mount a preovulatory LH surge.

Materials and Methods:

Detailed description of the materials and methods used in this study are given in ***SI Appendix***.

Figure legends:

Fig. 1. *Mc4r* is expressed within Kiss1 neurons and determines the timing of puberty onset.

(A) *In situ* hybridization using RNAscope technology showed co-expression of *Kiss1* (green) and

Mc4r (red) mRNA transcripts within ARH neurons cell bodies of ovariectomized female mice

(n=4). *Kiss1* and *Mc4r mRNAs* were co-expressed above background threshold levels within

84.01% of ARH cell bodies. Scale bar is 100 μm. **(B)** RNAscope in the AVPV/PeN of OVX+ E2

treated females (n=4/group) shows co-expression of *Kiss1* (green) and *Mc4r* (red) mRNA

transcripts within AVPV/PeN neurons. *Kiss1* and *Mc4r* mRNAs are co-expressed above

background threshold levels within 74.06% AVPV/PeN cell bodies. Scale bar is 100 μm.

Kiss1-MC4RKO females display advanced puberty onset, assessed by daily monitoring of vaginal

opening **(C)** and first estrus **(D)**, as documented by cumulative percent and mean age of animals

at vaginal opening **(C)** and first estrus **(D)** in Kiss1-MC4RKO females (n=24) compared to WT

littermates (n=19). *p<0.05 by Student's t-test. Data presented as the mean± SEM. **(E)** Female

Kiss1-MC4RKO (n=19) have normal body weight at the time of puberty onset compared to their

WT littermates (n=15). **(F)** Ontogeny expression of melanocortin genes (*Agrp*, *Pomc*, *Mc4r and*

Mc3r) in the ARH of WT female mice at different postnatal pre-pubertal and pubertal ages: P10,

P15, P22 and P30, normalized to the housekeeping gene *Hprt*. (*n values*: females at P10 (n=6),

P15 (n=6), P22 (n=6), and P30 (n=5)). Groups with different letters are significantly different

(p<0.05), as determined by one-way ANOVA followed by the Student-Newman-Keuls test. Data

presented as the mean± SEM.

Fig. 2. Deletion of MC4R from Kiss1 neurons impairs fertility in Kiss1-MC4RKO females.

(A) Female Kiss1-MC4RKO (n=19) have normal body weight from weaning and until post-natal

day (PND) 150 compared to their WT littermates (n=15). **(B)** Representative examples of estrous

cycles in Kiss1-MC4RKO and control females (n=5/group) assessed by daily vaginal cytology for

15 days. **(C)** Kiss1-MC4RKO females displayed irregular estrous cycles with a longer time in

diestrus and a shorter time in estrus compared to control females. *** P < 0.001 Two-way ANOVA.

(D) Pattern of LH pulsatility was analyzed in gonad intact Kiss1-MC4RKO (n=6) and control (n=7)

females. # Represents LH pulses. **(E)** LH total secretory mass assessed by area under the curve

(AUC), **(F)** LH pulse amplitude, **(G)** Basal LH and **(H)** total number of pulses /180 min were

analyzed. *p<0.05 by Student's t-test. The expression of the KNDy genes *Pdyn* **(I)**, *Kiss1* **(J)**,

Tac2 **(K)** and *Tacr3* **(L)**, was assessed in the ARH of adult Kiss1-MC4RKO (n=6) and control

(n=3) females. **p<0.01 by Student's t-test. Data presented as the mean± SEM. **(M)** Kiss1-

MC4RKO (n=6) and control (n=5) females were subjected to an LH surge induction protocol. LH samples were collected in the morning (AM [8 a.m.]) and evening (PM [7 p.m.]) after lights off. **P < 0.01, Two-way ANOVA followed by Bonferroni *post hoc* test. **(N)** Ovarian histology shows a decrease in the number of corpora lutea (CL, **O**) and increase in the number of cystic follicles (CF, **P**) of Kiss1-MC4RKO compared to controls(n=5/group). # represents cystic follicles. Serum levels of Testosterone **(Q)** and AMH **(R)** in adult gonad intact Kiss1-MC4RKO and control females (n=5/group). Student t test for unpaired samples. Data presented as the mean± SEM. Kiss1-MC4RKO females display impaired fertility characterized by increased time to deliver pups **(S)** and decreased number of pups per litter **(T)**, (n=9/group). *p<0.05 by Student's t-test. Data presented as the mean± SEM.

Fig. 3. Re-insertion of MC4R in Kiss1 neurons restores estrous cyclicity and LH pulsatility in Kiss1-cre:MC4R-LoxTB females.

(A) *Kiss1-cre*:MC4R-LoxTB (n=11) and MC4R-loxTB (MC4RKO) (n=7) displayed normal puberty onset as compared to their control littermates (n=14), as documented by cumulative percent and mean age of animals at vaginal opening and first estrus. **(B)** *Kiss1-cre*:MC4R-LoxTB and MC4R-loxTB both had significantly higher body weights at the time of puberty onset compared to their controls. **P < 0.01, *** P < 0.001 by one way ANOVA. **(C)** MC4R-loxTB (n=7) and *Kiss1-cre*:MC4R-LoxTB (n=11) females displayed significantly higher body weight than their littermates (n=10) from post-natal day 30 onwards, *p<0.05 and **P < 0.01. Data presented as the mean± (SEM). **(D)** Representative examples of estrous cycles of MC4R-loxTB (n=4), *Kiss1-cre*:MC4R-LoxTB (n=5) and control littermate mice, (n=4) assessed by daily monitoring of vaginal smears for 15 days. (D/M: diestrus/metestrus, P: proestrus, E: estrus). **(E)** MC4R-loxTB females displayed irregular estrous cycles, presenting longer time in diestrus and shorter time in estrus compared to control females, while *Kiss1-cre*:MC4R-LoxTB females displayed regular estrous cyclicity, similar to controls. *p<0.05, **P < 0.01, *** P < 0.001, ****P < 0.0001, Two-way ANOVA. Data

presented as the mean± SEM. **(F)** Pattern of LH pulsatility analyzed in gonad intact MC4R-loxTB

(n=4), *Kiss1-cre*:MC4R-LoxTB (n=5) and control littermates females (n=5). LH samples were

collected every 10 min for 180 minutes; # represents LH pulses. **(G)** LH total secretory mass, **(H)**

LH pulse amplitude, **(I)** basal LH, and **(J)** total number of pulses /180 min were assessed. *p<0.05,

P < 0.01 by one way ANOVA. **(K) Representative samples of ovarian histology from MC4R-

loxTB, *Kiss1-cre*:MC4R-LoxTB and control females (n=5/group); CL: Corpora lutea. Data are

presented as the mean± SEM. **(L)** Ovarian histology showed a significant decrease in the number

of corpora lutea in the MC4RKO and *Kiss1-cre*:MC4R-LoxTB compared to controls. Groups with

different letters are significantly different. Serum levels of **(M)** Testosterone and **(N)** AMH in adult

gonad intact females MC4R-loxTB, *Kiss1-cre*:MC4R-LoxTB and their control littermates. *p<0.05,

****P < 0.0001, One Way ANOVA. Data are presented as the mean± SEM.

Fig. 4. Kiss1^{ARH} neurons are activated by melanocortin agonists and respond to

optogenetic stimulation of POMC neurons.

(A) Slices were taken from *Kiss1-cre* injected (confocal image) brains to target fluorescent cells

(arrowhead) for recording (white outline over electrode). **(B)** Whole cell voltage clamp recording

of Kiss1^{ARH} neurons following the direct application of the high affinity melanocortin receptor

agonist melanotan II (MTII, 250 nM) was added directly to the bath and the response was

compared between slices from OVX and OVX+E females. **(C)** While the average inward current

was slightly higher in the OVX+E state, this was not a significant difference (Student's t-test for

unpaired samples, p>0.05). **(D)** The MC4R-selective agonist THIQ (100 nM) was perfused, and

excitatory inward currents were generated in Kiss1^{ARH} neurons. **(E)** Images are of AAV-driven

labeling of POMC cells as seen through the confocal. **(F)** Using the AAV-driven expression of

channelrhodopsin in adult *Pomc-cre* mice, high frequency stimulation elicited a slow inward

current in Kiss1 neurons. **(G)** The identity of cells was confirmed through the presence of a

persistent sodium current (see (23)) and/or with RT-PCR of harvested cytoplasm showing *Kiss1*

expression. (Gel: MM=molecular marker, TC=tissue controls). **(H)** A direct synaptic projection

from POMC to Kiss1^ARH neuron (*posthoc* identified) was confirmed using the "rescue" protocol: 1,

baseline glutamatergic responses were initially generated (black trace); 2, then action potentials

were eliminated by blocking voltage-gated sodium channels with TTX and the postsynaptic

response (red trace); 3, blockade of potassium channels facilitated calcium entry into the terminal

through ChR2 to release synaptic vesicles, "rescuing" the postsynaptic glutamate response

(green trace). **I)** Glutamatergic responses were also often observed, particularly in the ventral

ARH. As seen when targeting low input resistance neurons (i.e., Kiss1^ARH) with low frequency

optogenetic stimulation (5 ms pulse, 50 ms inter-spike interval), the first response in OVX +

vehicle females was larger relative to the second response in OVX + E2- treated female mice.

Representative traces are the average of 30 sweeps. **(J)** The averaged paired-pulse ratio was

lower in recordings from estradiol-treated female mice, indicating an increased release probability

(Students t-test $p<0.05$,). **(K)** High frequency optogenetic stimulation elicited a small inward

current. **(L)** In a different cell, a high frequency inward current was noted before constant perfusion

of SHU9119 for 15 minutes (break between traces). When the stimulation protocol was repeated,

no inward current was elicited. **(M)** The majority of high frequency responses were inward and

only twice was an inhibitory outward current recorded in identified Kiss1^ARH neurons. Perfusion of

the non-selective opioid receptor antagonist naloxone reversed the current, eliminating MCR-

activation as the mechanism.

Fig. 5. Kiss1^AVPV/PeN neurons are inhibited by MC4R agonists in an E2-dependent manner.

(A) Immunohistochemistry showing typical labeling of α-MSH fibers in the AVPV/PeN region in

OVX+E2 WT female mice. **(B)** Whole-cell voltage clamp recordings were made in ChR2-YFP

positive cells in brain slices taken from *Kiss1-cre*:Ai32 female mice. Bath application of melanotan

II (MTII, 500 nM) generated an inhibitory outward current in a Kiss1^AVPV/PeN neuron from OVX

female. As the cell was synaptically isolated using bath-applied tetrodotoxin (TTX, 1 µM) this

represents a direct effect. Washout of MTII while still in TTX quickly led to a return to baseline

RMP. **C)** An IV relationship was plotted using voltage steps before and after MTII administration.

The crossing near -85 mV (~EK^+) suggested the opening of a K+ channel. **D)** In a subset of

recordings from OVX brains slices, the selective membrane estrogen receptor (Gq-mER) agonist

STX (10 nM) was added to the bath for ~10 minutes prior to addition of MTII. STX pretreatment

resulted in either a greatly attenuated outward current or even an inward current as shown in this

example. **(E)** The mean outward current was significantly higher when recording in Kiss1[AVPV/Pen]

neurons from brain slices from vehicle-treated, OVX females compared to E2-treated, OVX

females or acute STX-treated brain slices from OVX females (One-way ANOVA $F_{(2,19)}$=12.32,

p<0.001; Holm-Sidak *posthoc* comparisons found significant differences between all groups; *

p<0.05, *** p<0.001). **(F)** There was no difference in the frequency of mEPSCs after MTII,

calculated as a percent of baseline between groups. **(G)** However, there was a significant

difference in the mEPSC amplitude (calculated as a percent of baseline) between all groups: main

effect $F_{(2,19)}$ =10.58, p<0.001. *Posthoc* comparisons using Holm-Sidak found OVX+STX to be

different from both OVX (p<0.001) and OVX+E (p<0.05).

Fig. 6. Working model of POMC to Kiss1 neuronal circuitry.

In ovariectomized females, POMC neurons excite Kiss1[ARH] neurons via glutamate and αMSH

release, which activate glutamatergic ionotropic and MC4R, respectively. E2-treatment enhances

the glutamatergic excitation of these neurons. In contrast in ovariectomized females, αMSH fibers

project to the AVPV/PeN, and MC4R agonists (*i.e.*, MTII) inhibit Kiss1[AVPV/PeN] neurons via opening

a K^+ channel.   However, E2-treatment rapidly abrogates (uncouples) the inhibitory MC4R

signaling in Kiss1[AVPV/PeN] neurons, allowing greater excitatory actions of Kiss1[ARH] neurons via

enhanced glutamate release (26).

Fig. S1. Validation of Kiss-MC4RKO mouse model, related to figure 5.

(A) Representative images depicting co-expression of *Kiss1* and *Mc4r* mRNA, using RNAscope, in OVX WT female mice (upper left panel), and OVX Kiss1-MC4RKO female mice (upper right panel). As expected, *Mc4r* was not detected within Kiss1 neurons in Kiss1-MC4RKO females. **(B)** Representative images depicting comparable *Mc4r* mRNA expression in the paraventricular hypothalamus between OVX WT and Kiss1-MC4RKO females. Scale bar is 100 µm.

Fig. S2. Validation of Kiss1-cre:MC4R-LoxTB mouse model, related to figure 4.

(A) Representative images depicting co-expression of *Kiss1* and *Mc4r* mRNA, using RNAscope, in OVX *Kiss1-cre:MC4R-LoxTB* female mice in the ARH (upper right panel). As expected, Mc4r is detected within Kiss1 neurons in *Kiss1-cre:MC4R-LoxTB* females. **(B)** Representative images depicting the absence of *Mc4r* mRNA expression in the paraventricular hypothalamus of *Kiss1-cre*:MC4R-LoxTB female, compared to their WT control littermates. Scale bar is 100 µm.

Acknowledgments:

The authors would like to recognize the excellent technical expertise of Ms. Martha A. Bosch (tissue preparation, immunohistochemical procedure and single cell RT-PCR of recorded neurons). In addition, we thank Dr. Rona Carroll for her assistance at Harvard Medical School and Mr. Cole Martinson, a student worker in the Ronnekleiv/ Kelly laboratories, for his assistance with genotyping and care of mouse colonies at OHSU. The University of Virginia Center for Research in Reproduction Ligand Assay and Analysis Core is supported by the Eunice Kennedy Shriver NICHD/NIH (NCTRI) Grant P50-HD28934. This work was supported by PHS MPI grant DK68098 to MJK and OKR; HD090151, HD099084 and DK133760 to VMN, and The Charles A. King Trust Postdoctoral Research Fellowship Award, The Lalor Foundation Postdoctoral Fellowship Award, the Women's brain initiative Fellowship Award, the ROSA SCORE pilot grant

630   (supported by NIH Research Grant U54 AG062322 funded by The National Institute on Aging

631   (NIA) and Office of Research on Women's Health (ORWH)) and the IBRO-ISN Research

632   Fellowship Award to RT, and startup funds from Appalachian State University to TLS.

633
634
References:
636
1.   U. Health (2020) Infertility.
2.   M. L. Andermann, B. B. Lowell, Toward a Wiring Diagram Understanding of
Appetite Control. *Neuron* **95**, 757-778 (2017).
3.   R. D. Cone, Studies on the physiological functions of the melanocortin system.
Endocr Rev **27**, 736-749 (2006).
4.   B. P. Shah *et al.*, MC4R-expressing glutamatergic neurons in the paraventricular
hypothalamus regulate feeding and are synaptically connected to the parabrachial
nucleus. *Proceedings of the National Academy of Sciences of the United States of
America* **111**, 13193-13198 (2014).
5.   K. Wang *et al.*, Neuroanatomy of melanocortin-4 receptor pathway in the mouse
brain. *Open Life Sci* **15**, 580-587 (2020).
6.   I. S. Farooqi *et al.*, Clinical spectrum of obesity and mutations in the melanocortin
4 receptor gene. *The New England journal of medicine* **348**, 1085-1095 (2003).
7.   I. A. Hainerová, H. Zamrazilová, D. Sedláčková, V. Hainer, Hypogonadotropic
hypogonadism in a homozygous MC4R mutation carrier and the effect of
sibutramine treatment on body weight and obesity-related health risks. *Obes Facts*
4, 324-328 (2011).
8.   M. Doulla, A. D. McIntyre, R. A. Hegele, P. H. Gallego, A novel MC4R mutation
associated with childhood-onset obesity: A case report. *Paediatr Child Health* **19**,
515-518 (2014).
9.   A. A. Batarfi *et al.*, MC4R variants rs12970134 and rs17782313 are associated
with obese polycystic ovary syndrome patients in the Western region of Saudi
Arabia. *BMC Med Genet* **20**, 144 (2019).
10.   X. Chen *et al.*, Deficient melanocortin-4 receptor causes abnormal reproductive
neuroendocrine profile in female mice. *Reproduction (Cambridge, England)* **153**,
267-276 (2017).
11.   M. Sandrock *et al.*, Reduction in corpora lutea number in obese melanocortin-4-
receptor-deficient mice. *Reprod Biol Endocrinol* **7**, 24 (2009).
12.   L. Cui *et al.*, Dapagliflozin partially restores reproductive function in MC4R KO
obese female mice. *The Journal of endocrinology* **254**, 65-76 (2022).
13.   M. Manfredi-Lozano *et al.*, Defining a novel leptin-melanocortin-kisspeptin
pathway involved in the metabolic control of puberty. *Mol Metab* **5**, 844-857 (2016).
14.   C. C. Nestor *et al.*, Optogenetic Stimulation of Arcuate Nucleus Kiss1 Neurons
Reveals a Steroid-Dependent Glutamatergic Input to POMC and AgRP Neurons
in Male Mice. *Mol Endocrinol* **30**, 630-644 (2016).
15.   J. Qiu *et al.*, Estrogenic-dependent glutamatergic neurotransmission from
kisspeptin neurons governs feeding circuits in females. *Elife* **7** (2018).

16. R. L. Goodman, A. E. Herbison, M. N. Lehman, V. M. Navarro, Neuroendocrine
control of gonadotropin-releasing hormone: Pulsatile and surge modes of
secretion. *Journal of neuroendocrinology* **34**, e13094 (2022).
17. D. D. Israel *et al.*, Effects of leptin and melanocortin signaling interactions on
pubertal development and reproduction. *Endocrinology* **153**, 2408-2419 (2012).
18. M. Manfredi-Lozano, J. Roa, M. Tena-Sempere, Connecting metabolism and
gonadal function: Novel central neuropeptide pathways involved in the metabolic
control of puberty and fertility. *Front Neuroendocrinol* **48**, 37-49 (2018).
19. A. M. Moore, M. Prescott, R. E. Campbell, Estradiol negative and positive feedback
in a prenatal androgen-induced mouse model of polycystic ovarian syndrome.
Endocrinology **154**, 796-806 (2013).
20. E. A. McCarthy *et al.*, Inhibiting Kiss1 Neurons With Kappa Opioid Receptor
Agonists to Treat Polycystic Ovary Syndrome and Vasomotor Symptoms. *J Clin
Endocrinol Metab* **107**, e328-e347 (2022).
21. D. Dewailly, A. L. Barbotin, A. Dumont, S. Catteau-Jonard, G. Robin, Role of Anti-
Müllerian Hormone in the Pathogenesis of Polycystic Ovary Syndrome. *Frontiers
in endocrinology* **11**, 641 (2020).
22. J. Roa, A. E. Herbison, Direct regulation of GnRH neuron excitability by arcuate
nucleus POMC and NPY neuron neuropeptides in female mice. *Endocrinology*
153, 5587-5599 (2012).
23. C. Zhang, M. A. Bosch, J. Qiu, O. K. Ronnekleiv, M. J. Kelly, 17beta-Estradiol
increases persistent Na(+) current and excitability of AVPV/PeN Kiss1 neurons in
female mice. *Mol Endocrinol* **29**, 518-527 (2015).
24. J. D. Clements, G. L. Westbrook, Activation kinetics reveal the number of
glutamate and glycine binding sites on the N-methyl-D-aspartate receptor. *Neuron*
7, 605-613 (1991).
25. T. L. Stincic, P. Grachev, M. A. Bosch, O. K. Rønnekleiv, M. J. Kelly, Estradiol
drives the anorexigenic activity of proopiomelanocortin neurons in female mice.
eNeuro **5**, e0103-0118.2018 (2018).
26. J. Qiu *et al.*, High-frequency stimulation-induced peptide release synchronizes
arcuate kisspeptin neurons and excited GnRH neurons. *eLife* **5**, e16246 (2016).
27. J. Qiu *et al.*, Estrogenic-dependent glutamatergic neurotransmission from
kisspeptin neurons governs feeding circuits in females. *eLife* **7**, e35656 (2018).
28. T. L. Stincic, J. Qiu, A. M. Connors, M. J. Kelly, O. K. Rønnekleiv, Arcuate and
Preoptic Kisspeptin neurons exhibit differential projections to hypothalamic nuclei
and exert opposite postsynaptic effects on hypothalamic paraventricular and
dorsomedial nuclei in the female mouse. *eneuro* 10.1523/eneuro.0093-21.2021,
ENEURO.0093-0021.2021 (2021).
29. M. A. Herman, F. Ackermann, T. Trimbuch, C. Rosenmund, Vesicular glutamate
transporter expression level affects synaptic vesicle release probability at
hippocampal synapses in culture. *The Journal of Neuroscience* **34**, 11781-11791
(2014).
30. J. E. Thornton, M. D. Loose, M. J. Kelly, O. K. Rønnekleiv, Effects of estrogen on
the number of neurons expressing β-endorphin in the medial basal hypothalamus
of the female guinea pig. *The Journal of comparative neurology* **341**, 68-77 (1994).

31.   K. G. Mountjoy, M. T. Mortrud, M. J. Low, R. B. Simerly, R. D. Cone, Localization
720   of the melanocortin-4 receptor (MC4-R) in neuroendocrine and autonomic control
721   circuits in the brain. *Molecular Endocrinology* **8**, 1298-1308 (1994).

722   32.   J. Yu *et al.*, Determination of the melanocortin-4 receptor structure identifies
723   Ca^{2+} as a cofactor for ligand binding. *Science* **368**, 428-433 (2020).

724   33.   J. W. Sohn *et al.*, Melanocortin 4 receptors reciprocally regulate sympathetic and
725   parasympathetic preganglionic neurons. *Cell* **152**, 612-619 (2013).

726   34.   M. J. Kelly, M. D. Loose, O. K. Ronnekleiv, Estrogen suppresses mu-opioid- and
727   GABAB-mediated hyperpolarization of hypothalamic arcuate neurons. *The Journal*
728   *of neuroscience : the official journal of the Society for Neuroscience* **12**, 2745-2750
729   (1992).

730   35.   A. H. Lagrange, O. K. Ronnekleiv, M. J. Kelly, The potency of mu-opioid
731   hyperpolarization of hypothalamic arcuate neurons is rapidly attenuated by 17
732   beta-estradiol. *The Journal of neuroscience : the official journal of the Society for*
733   *Neuroscience* **14**, 6196-6204 (1994).

734   36.   J. Qiu *et al.*, Rapid signaling of estrogen in hypothalamic neurons involves a novel
735   G-protein-coupled estrogen receptor that activates protein kinase C. *The Journal*
736   *of neuroscience : the official journal of the Society for Neuroscience* **23**, 9529-9540
737   (2003).

738   37.   J. Qiu *et al.*, A G-protein-coupled estrogen receptor is involved in hypothalamic
739   control of energy homeostasis. *The Journal of neuroscience : the official journal of*
740   *the Society for Neuroscience* **26**, 5649-5655 (2006).

741   38.   B. Y. H. Lam *et al.*, MC3R links nutritional state to childhood growth and the timing
742   of puberty. *Nature* **599**, 436-441 (2021).

743   39.   T. L. Stincic, P. Grachev, M. A. Bosch, O. K. Ronnekleiv, M. J. Kelly, Estradiol
744   Drives the Anorexigenic Activity of Proopiomelanocortin Neurons in Female Mice.
745   *eNeuro* **5** (2018).

746   40.   M. Ghamari-Langroudi *et al.*, G-protein-independent coupling of MC4R to Kir7.1 in
747   hypothalamic neurons. *Nature* **520**, 94-98 (2015).

748   41.   W. C. Krause *et al.*, Oestrogen engages brain MC4R signalling to drive physical
749   activity in female mice. *Nature* **599**, 131-135 (2021).

750   42.   N. Brix *et al.*, Timing of puberty in boys and girls: A population-based study.
751   *Paediatr Perinat Epidemiol* **33**, 70-78 (2019).

A. anti-αMSH 100 μm

B. MTII Washout 20 pA 30 s

C. Baseline MTII Current (pA) Voltage (mV)

D. 10 nM STX 250 nM MTII 10 pA 40 ms

E. MTII Current (pA) — OVX (8), OVX+E (8), OVX+STX (7); *** * *

F. % change mEPSC frequcy — OVX (7), OVX+E (8), OVX+STX (7); n.s

G. % change mEPSC amplitude — OVX (7), OVX+E (8), OVX+STX (7); *** * n.s

Ovariectomized females

Kiss1^{AVPV}

Glu

α-MSH

+

−

Kiss1^{ARH}

+

POMC

E2-treated, ovariectomized females

Kiss1^{AVPV}

Glu

α-MSH

+

+

+

Kiss1^{ARH}

+

POMC