POMC neurons control fertility through differential signaling of MC4R in Kisspeptin neurons

Authors:
Rajae Talbi[1,2*], Todd L. Stincic[4,8*], Kaitlin Ferrari[2], Choi Ji Hae[2], Karol Walec[2], Elizabeth Medve[2], Achi Gerutshang[2], Silvia León[1,2], Elizabeth A. McCarthy[1,2], Oline K. Rønnekleiv[3,4], Martin J. Kelly[3,4,6] and Víctor M. Navarro[1,2,5,6]

[1]Harvard Medical School, Boston, MA, USA; [2]Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA; [3]Department of Chemical Physiology and Biochemistry, Oregon Health & Science University, Portland, OR, USA; [4]Division of Neuroscience, Oregon National Primate Research Center, Beaverton, OR, USA; [5]Harvard Program in Neuroscience, Boston, MA, USA

*These authors contributed equally
[6] Corresponding authors
[8]Present address: Department of Biology, Appalachian State University, Boone, NC 28608.

Correspondence:
Victor M. Navarro, PhD, Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, 221 Longwood Avenue, Room 219, Boston, Massachusetts 02115. E-mail: vnavarro@bwh.harvard.edu.

Martin J. Kelly, PhD, Department of Chemical Physiology and Biochemistry, Oregon Health & Science University, 3181 SW Sam Jackson Park Road, Portland, OR 97239. E-mail: kellym@ohsu.edu .

Keywords:
Kiss1 neurons, POMC neurons, MC4R, Fertility, Ovulation, Obesity, Estradiol.

Abstract

Inactivating mutations in the melanocortin 4 receptor (*MC4R*) gene cause monogenic obesity. Interestingly, female patients also display various degrees of reproductive disorders, in line with the subfertile phenotype of MC4RKO female mice. However, the cellular mechanisms by which MC4R regulates reproduction are unknown. Kiss1 neurons directly stimulate gonadotropin-releasing hormone (GnRH) release through two distinct populations; the Kiss1[ARH] neurons, controlling GnRH pulses, and the sexually dimorphic Kiss1[AVPV/PeN] neurons controlling the preovulatory LH surge. Here, we show that *Mc4r* expressed in Kiss1 neurons regulates fertility in females. *In vivo*, deletion of *Mc4r* from Kiss1 neurons in female mice replicates the reproductive impairments of MC4RKO mice without inducing obesity. Conversely, reinsertion of *Mc4r* in Kiss1 neurons of MC4R null mice restores estrous cyclicity and LH pulsatility without reducing their obese phenotype. *In vitro*, we dissect the specific action of MC4R on Kiss1[ARH] vs Kiss1[AVPV/PeN] neurons and show that MC4R activation excites Kiss1[ARH] neurons through direct synaptic actions. In contrast, Kiss1[AVPV/PeN] neurons are normally inhibited by MC4R activation except under elevated estradiol levels, thus facilitating the activation of Kiss1[AVPV/PeN] neurons to induce the LH surge driving ovulation in females. Our findings demonstrate that POMC[ARH] neurons acting through MC4R, directly regulate reproductive function in females by stimulating the "pulse generator" activity of Kiss1[ARH] neurons and restricting the activation of Kiss1[AVPV/PeN] neurons to the time of the estradiol-dependent LH surge, and thus unveil a novel pathway of the metabolic regulation of fertility by the melanocortin system.

Introduction

Obesity rates have skyrocketed in Western societies in the last decades, resulting in an alarming rise in comorbidities that place a significant burden on healthcare systems. The increase in obesity correlates with a decrease in fertility rates, leading to conception challenges that are experienced by approximately 15% of couples in the United States currently (1).

The melanocortin 4 receptor (MC4R) binds i) α-melanocyte stimulating hormone (αMSH), an agonist product of the pro-opiomelanocortin (*Pomc*) gene, and ii) the inverse agonist, agouti-related peptide (AgRP), to regulate food intake and energy expenditure (2, 3). While the role of MC4R on food intake is largely mediated by neurons located in the paraventricular nucleus of the hypothalamus (PVN) (4), its expression in the brain is widespread (5) with the specific function of the different MC4R-expressing neurons in areas beyond the PVN remaining to be fully explored. Inactivating mutations in *MC4R* are a leading cause of monogenic obesity in humans (6) and cause excessive obesity and hyperphagia in mice (7). Scant evidence in humans shows an association between *MC4R* mutations and higher incidence of hypogonadotropic hypogonadism (HH) (8), alterations in the timing of puberty onset (9), and polycystic ovary syndrome (PCOS) (10); however, other studies have shown no association between MC4R and reproductive disorders (6). Therefore, the role of MC4R signaling in reproductive function in humans remains controversial despite the clear association found in mice. Supporting this role of MC4R, *Mc4r* null mice display an array of reproductive abnormalities that largely affects females, characterized by irregular estrous cycles, disrupted luteinizing hormone (LH) secretion, reduced corpora lutea and reduced fertility (11-13). Further evidence from mice demonstrates that MC4R agonists robustly increase LH release in a kisspeptin-dependent manner (14).

Kisspeptin is the most potent gonadotropin releasing hormone (GnRH) secretagogue known to date and it is mainly produced in two distinct neuronal populations. Kiss1 neurons of the arcuate

nucleus of the hypothalamus (Kiss1^{ARH}), present in both sexes, control the pulsatile (tonic)

release of GnRH, and sex steroids attenuate their release of kisspeptin. Kiss1 neurons of the

anteroventral periventricular continuum area (Kiss1$^{AVPV/PeN}$), are predominantly present in

females, and are responsible for generating the preovulatory GnRH/LH surge essential for

ovulation (15). Despite these critical roles of both populations of Kiss1 neurons for reproduction,

the underlying mechanisms that determine how each Kiss1 population responds differently to

sex steroids to regulate the GnRH tonic vs surge release remains unresolved.

In rodents, compelling evidence indicates a close interaction between Kiss1 neurons and the

melanocortin system: 1) fibers from POMC neurons in the arcuate nucleus (POMC^{ARH}) project to

and juxtapose Kiss1^{ARH} neurons (14); 2) the melanocortin signaling through MC4R contributes

to the permissive role of leptin on puberty onset (14, 16, 17); and 3) MC4R expression on both

Kiss1^{ARH} (18-20) and Kiss1$^{AVPV/PeN}$ (18, 21). Altogether, this evidence suggests a clear role for

MC4R in Kiss1 neurons, in the control of reproductive maturation and fertility.

In this study, we investigated the contribution of MC4R signaling versus obesity *per se* in the

etiology of the reproductive impairments observed in *Mc4r* null mice, which could explain similar

impairments observed in humans. Using genetic mouse models with specific deletion or

reinsertion of *Mc4r* in Kiss1 neurons, we show that MC4R action in Kiss1 neurons is necessary

for normal reproductive function in female mice. Whole cell voltage clamp recordings evidenced

an excitatory action of MC4R on Kiss1^{ARH} neurons and an estradiol-dependent inhibitory action

on Kiss1$^{AVPV/PeN}$ neurons, with important implications for the timing of the preovulatory LH surge.

Results

***Mc4r* within Kiss1 neurons determines the timing of puberty onset in females.**

The expression of *Mc4r* within Kiss1^{ARH} and Kiss1$^{AVPV/PeN}$ neurons has already been described

elsewhere (18-21), suggesting a role for MC4R in the regulation of fertility. To further investigate

the specific role of MC4R in Kiss1 neurons, we generated a Kiss1-specific MC4R knockout

mouse model (Kiss1-MC4RKO). The specific deletion of *Mc4r* from Kiss1 neurons, as well as

the absence of global recombination, were confirmed through RNAscope. While Kiss1 neurons

of Kiss1-MC4RKO mice lack *Mc4r* transcript compared to their control littermates (**Figure S1A**),

Mc4r was detectable in the PVN of all mice, which is a major site of *Mc4r* expression in the brain

in the regulation of metabolism, therefore supporting the specific deletion of *Mc4r* only within

Kiss1 neurons (**Figure S1B**).

Puberty onset was assessed daily from weaning age through the monitoring of vaginal opening

(VO) and first estrus (FE). Kiss1-MC4RKO females showed a significant advancement in the

age of VO (P=0.0150, Kiss1-MC4RKO: 24.67± 0.39 vs controls: 26.32±0.53) and FE (P=0.0341,

Kiss1-MC4RKO: 32.63±1.4 vs controls: 37.84±1.90) compared to their littermate controls

(**Figure 1A, B**). Body weight of Kiss1-MC4RKO females at the age of puberty onset was similar

between groups (P=0.2596, Kiss1-MC4RKO: 12.41±0.30 vs controls: 12.89±0.26) (**Figure 1C**).

These data suggest that melanocortin signaling on Kiss1 neurons participates in the timing of

puberty onset in females. Since the role of the melanocortin system on puberty onset is largely

unexplored, we evaluated the expression profile of the main components of this system in the

hypothalami of WT female mice at postnatal days (PND) 10, 15, 22 and 30. Interestingly,

expressions of *Agrp*, *Mc3r* and *Mc4r* were significantly lower at the time of puberty onset

(PND30) compared to earlier developmental ages (**Figure 1D**). This supports our *in vivo* data

and suggests that a decrease in the hypothalamic melanocortin signaling drives puberty onset,

in line with the advancement in the age of VO and FE observed in Kiss1-MC4RKO female mice

(**Figure 1A, B**).

MC4R in Kiss1 neurons is required for female reproduction.

Because Kiss1-MC4RKO females showed altered puberty onset, we further investigated their

reproductive phenotype. Interestingly, Kiss1-MC4RKO female mice displayed normal BW

throughout the time of the study (up to PND150) **(Figure 2*A*)**; however, they presented with

irregular estrous cycles with predominantly more time spent in diestrus (P=0.0028) and less

time spent in estrus (P=0.0067) than their control littermates (**Figure 2*B, S3a, b***). The

assessment of the pulsatile release of LH every 10 min over 180 min revealed no change in the

total number of pulses between groups but a significant increase in basal LH release (P=0.04)

in Kiss1-MC4RKO female mice (0.42±0.03) compared to controls (0.34±0.02) (**Figure 2*C-H***).

The analysis of the gene expression of the "KNDy" systems in the ARH, which control the GnRH

pulse generator (15), revealed normal expression levels of *Kiss1*, *Tac2* and *Tacr3*, but

significantly lower expression of *Pdyn* (P=0.0067, Kiss1-MC4RKO: 0.78±0.038 vs controls:

1.00±0.019) (**Figure 2*I-L***). The conserved expression of *Kiss1*, *Tac2* and *Tacr3* correlates with

the preserved LH pulse frequency and amplitude in Kiss1-MC4RKO mice, while the lower

inhibitory tone of dynorphin (**Figure 2*I***) correlates with the higher basal LH levels (**Figure 2*G***).

However, the mRNA expression of these genes, alone, does not necessarily reflect changes of

their activities. To assess the contribution of MC4R signaling in Kiss1 neurons to the induction

of ovulation through the preovulatory LH surge, Kiss1-MC4RKO females and control littermates

were submitted to an LH surge induction protocol that showed the induced LH surge was

significantly blunted in the Kiss1-MC4RKO females (P=0.0091), while the protocol clearly

evoked the expected afternoon rise of LH in control mice (**Figure 2*M***). In line with these

findings, the ovaries of Kiss1-MC4RKO females displayed fewer corpora lutea, markers of

recent ovulation (P=0.0054, Kiss1-MC4RKO: 0.80±0.37 vs controls: 2.80±0.37), in addition to

increased cystic follicles (P=0.0337, Kiss1-MC4RKO: 1.60±0.40 vs controls: 0.40±0.24) (**Figure

2*N-P***), which correlate with decreased fertility as observed by the extended time to deliver pups

(*i.e.* longer time to get pregnant) (P=0.0030, Kiss1-MC4RKO: 30.30±4.60 vs controls:

20.88±0.22), and fewer pups per litter (P=0.0095, Kiss1-MC4RKO: 6.66±0.55 vs controls:

8.44±0.24) (**Figure 2*S, T***).

The increase in serum LH levels and decreased ovulation observed in the Kiss1-MC4RKO

females is reminiscent of polycystic ovary syndrome (PCOS) mouse models (22, 23). Thus, we

investigated whether Kiss1-MC4RKO females display a PCOS-like phenotype. We analyzed

circulating levels of testosterone (T) and anti-müllerian hormone (AMH), which are frequently

elevated in PCOS models (24). The Kiss1-MC4RKO females expressed normal T and AMH

levels compared to control mice in diestrus (**Figure 2*Q, R***). Thus, we can exclude a PCOS-like

reproductive phenotype mediated by the lack of melanocortin signaling on Kiss1 neurons.

Re-insertion of MC4R in Kiss1 neurons of MC4RKO mice improves reproductive function.

Kiss1-MC4RKO females displayed reproductive abnormalities resembling those described in

MC4RKO females (11-13). Thus, we hypothesized that the reproductive defects described for

the MC4RKO mice would be, at least in part, due to the absence of MC4R signaling in Kiss1

neurons. To further investigate this hypothesis, we generated mice that do not express *Mc4r*

anywhere (MC4R-LoxTB, *i.e.* MC4RKO) or that express *Mc4r* only in Kiss1 neurons (*Kiss1-*

cre:MC4R-loxTB). The specific re-insertion of *Mc4r* within Kiss1 neurons in the *Kiss1-*

cre:MC4R-loxTB mice was confirmed through RNAscope (**Figure S2*A***). *Mc4r* expression was

not detected in the PVN of these mice (**Figure S2*B***), and it was only detected in Kiss1 neurons

in the *Kiss1-cre:MC4R-loxTB* mice. Puberty onset was assessed daily from weaning age

through the monitoring of vaginal opening (VO) and first estrus (FE). MC4R-LoxTB and *Kiss1-*

cre:MC4R-loxTB female mice displayed normal timing of puberty onset compared to their

control littermates, as assessed by VO (P=0.104, MC4R-LoxTB: 30.86±1.90 vs *Kiss1-*

cre:MC4R-loxTB: 30.09±1.00 vs controls: 27.57±0.81), and FE (P=0.8472, MC4R-LoxTB:

35.14±2.46 vs *Kiss1-cre:MC4R-loxTB: 34.55*±1.64 vs controls: 35.80±0.84) (**Figure 3*A, B***),

despite displaying significantly higher body weight at the time of puberty onset (P=0.0001,

MC4R-LoxTB: 15.49±0.50 vs *Kiss1-cre:MC4R-loxTB:* 15.54±0.40 vs Controls: 13.29±0.30)

(**Figure 3*C***). As expected, MC4R-LoxTB females (MC4RKO) displayed increased body weight

(**Figure 3D**) and an array of reproductive impairments that included: irregular estrous cycles

with significantly more days in diestrus (P=0.0016) and fewer days in estrus (P=0.0317)

compared to controls (**Figure 3*E, S3c, e***), significantly higher AUC (MC4R-LoxTB: 109.6±2.39

vs Controls: 80.36±9.05) serum LH levels characterized by higher basal (MC4R-LoxTB:

0.48±0.02 vs Controls: 0.37±0.06) and amplitude levels per LH pulse (MC4R-LoxTB:

0.730±0.009 vs Controls: 0.58±0.07) (**Figure 3*F-J***), and fewer corporal lutea (P=0.0366, MC4R-

LoxTB: 0.80±0.80 vs Controls: 4.20±1.15) (**Figure 3*K, L***), recapitulating the same reproductive

phenotype observed in Kiss1-MC4RKO mice despite the differences in BW. Re-introduction of

MC4R into Kiss1 neurons in *Kiss1-cre:MC4R-loxTB* mice completely recovered estrous

cyclicity, which was similar to controls (diestrus: P=0.1932, proestrus: P=0.8262, estrus:

P=0.0547, compared to controls) (**Figure 3*E, S3c, d***) despite *Kiss1-cre:MC4R-loxTB* mice

showing the same degree of obesity as MC4R-LoxTB mice (**Figure 3*D***), indicating that obesity

per se was not mediating the irregular estrous cycles in MC4RKO mice. As indicated above,

MC4R-LoxTB mice display higher overall circulating LH levels, and this feature was also

recovered by the re-insertion of MC4R in Kiss1 neurons of *Kiss1-cre:MC4R-loxTB* mice (basal

LH: 0.32±0.04, LH amplitude: 0.48±0.04, LH pulses/180min: 2.40±0.67 and AUC: 70.62±7.15

compared to controls) (**Figure 3*G-J***), further confirming that the melanocortin action on Kiss1

neurons is required for the normal control of gonadotropin release. Despite these significant

improvements in reproductive function, *Kiss1-cre:MC4R-loxTB* mice presented fewer corpora

lutea than controls (0.80±0.80), similar to MC4R-LoxTB mice, suggesting that an ovulatory

impairment still persists (**Figure 3*K, L***). While tonic LH release and estrous cyclicity are

predominantly controlled by Kiss1[ARH] neurons, ovulation is mediated by the induction of the

preovulatory LH surge by Kiss1[AVPV/PeN] neurons. It is possible that the metabolic signals derived

from their obese phenotype, and/or the absence of the direct action of MC4R in GnRH neurons

(16, 25) prevents the complete recovery of ovulation in *Kiss1-cre:MC4R-loxTB* mice. Both

genetic models displayed significantly lower T (MC4R-LoxTB: 28.50±6.20 vs *Kiss1-cre:MC4R-*

loxTB: 31.22±3.90 vs Controls: 62.84±16.53) and AMH (MC4R-LoxTB: 150.6±22.0 vs *Kiss1-*

cre:MC4R-loxTB: 146.9±10.8 vs Controls: 321.8±15.7) levels than control mice in diestrus

(**Figure 3*M, N***). Therefore, we can exclude, once again, a PCOS-like reproductive phenotype

mediated by the lack of melanocortin signaling.

Kiss1^{ARH} neurons are excited by MC4R agonists.

Based on the expression of *Mc4r* in Kiss1 neurons and the reproductive impairment found after

Mc4r deletion within Kiss1 neurons (**Figure 2**), we hypothesized that Kiss1^{ARH} neurons are

direct targets of melanocortins and would respond to MC4R activation. Initially we did whole-

cell, voltage clamp recordings in Kiss1^{ARH} neurons (**Figure 4*A***). *Kiss1-Cre* x Ai32 or *Kiss1-Cre*

AAV-injected mice underwent ovariectomy (OVX) and were given estradiol (E2) replacement

(*see Methods*). We targeted fluorescent cells for recording which were isolated synaptically by

bathing the slices in tetrodotoxin (TTX, 1 µM). Focal application of the high-affinity melanocortin

receptor agonist melanotan II (MTII, ~250 nM) elicited a small inward current in half of the

isolated Kiss1^{ARH} cells (10/21) (**Figure 4*B***). Next, we examined if the E2 state affected MTII

response by recording from Kiss1^{ARH} neurons from OVX females and found two thirds (6/9 cells)

responded, but there was no significant difference in the average current (**Figure 4*C***). Finally, it

must be noted that at this concentration MTII is a non-selective MCR agonist, so we followed up

using a perfusion of the MC4R-selective agonist THIQ (100 nM) to determine if activation of

MC4Rs alone was sufficient to invoke a postsynaptic response in Kiss1^{ARH} neurons from

OVX+E2 females. Similar to MTII, THIQ was able to elicit an excitatory inward current in

Kiss1^{ARH} neurons (4/5 cells, -7.4±0.5 pA), indicating MC4R activation is sufficient (**Figure 4*D***).

POMC neurons synapse directly with Kiss1^{ARH} neurons.

Based on the pharmacological activation of MCRs, we wanted to address whether POMC[ARH]

neurons are the source of melanocortins to excite Kiss1[ARH] neurons directly. To answer this

question, we injected an AAV-EF1α-DIO-ChR2:mCherry vector into the ARH of adult *Pomc-Cre*

female mice (**Figure 4E**) (24). After 2-4 weeks we did whole cell recordings from putative

Kiss1[ARH] neurons and looked for a response to high frequency optogenetic stimulation of POMC

fibers in OVX + E2-treated females (**Figure 4F**). We were able to confirm that these were Kiss1

neurons based on the presence of a persistent sodium current and/or *post hoc* identification by

scRT-PCR (40/77) cells, **Figure 4G**). While this current is more prevalent in the AVPV Kiss1

population, Kiss1 neurons are the only ARH neurons to display this electrophysiological

"fingerprint" of a pronounced I_{NaP} paired with a high capacitance and a low input resistance (26).

Additionally, while recording from putative Kiss1[ARH] neurons, we optogenetically stimulated

POMC fibers at low frequency (1 Hz, 5 ms, 470 nm light) and recorded fast postsynaptic inward

currents. The fast kinetics of the EPSC are also a tell-tale sign of an ionotropic glutamatergic

response (27). Previously, CNQX was sufficient to block similar excitatory ESPCs in other

postsynaptic targets of POMC neurons (28). While the consistent latency from optogenetic

stimulus and current response was indicative of a single synaptic delay, we wanted to establish

that it was a direct synaptic connection, as we have previously shown for output of POMC

neurons (28) and Kiss1 neurons (29-31). First, we abrogated the optogenetic response with the

addition of TTX (1 µM) to the bath, and then we were able to rescue the postsynaptic glutamate

response with the addition of the K^+ channel blockers 4-aminopyridine (4-AP; 0.5 mM) and

tetraethyl ammonium (TEA; 7.5 mM) (**Figure 4H**). K+ channel block enables the calcium influx

from ChR2 activation in the presynaptic terminals to be sufficient to restore vesicle fusion.

Therefore, there appears to be no intervening synapses between POMC and the downstream

Kiss1[ARH] neurons.

Previously we found that E2 increases glutamate release from POMC neurons by increasing

Vglut2 (28). Therefore, we compared the postsynaptic glutamatergic responses following two

optogenetic stimuli (50 ms interval between light flashes) in Kiss1^{ARH} neurons from OVX +

vehicle versus OVX + E2-treated female mice (**Figure 4*I***). As anticipated, we found that

treatment with E2 increased the probability of glutamate release from POMC neurons onto

Kiss1^{ARH} neurons based on the significant decrease in the paired pulse ratio of the two stimuli

(32) (**Figure 4*J***). Although in the present study we used an *in vivo* treatment paradigm, we

know from previous studies that this augmentation of glutamate release can happen quite

rapidly after a brief exposure to E2 *in vitro* (within 15 min) (28). Therefore, Kiss1^{ARH} neurons are

excited by the glutamatergic input from POMC neurons in an E2-dependent manner.

Next, we tested for a melanocortin response using a high frequency stimulation protocol known

to elicit peptide release (33). Only cells displaying a glutamatergic response were tested.

Indeed, high frequency stimulation evoked an excitatory inward current (**Figure 4*K***) that was

blocked by the selective MC3/4R antagonist SHU 9119 (**Figure 4*L***). However, the number of

high frequency responses seen in Kiss1 neurons was unexpectedly infrequent and relatively

small (4/16, 25%, mean inward current: -1.4±3.1 pA) compared to the efficacy of

pharmacological MCR activation. E2-treatment enhances *Pomc* expression and β-endorphin

labeling (34) (35), but the effect, if any, on αMSH release or the MCRs in Kiss1 neurons is

unknown. We suspected that the high frequency stimulation was eliciting co-release of αMSH

and β-endorphin, which would exert oppositive effects on a postsynaptic cell. In order to isolate

melanocortin signaling we pretreated slices with the non-selective opioid antagonist naloxone (1

µM) and found an increase in the likelihood and magnitude of a response to high frequency

stimulation (9/23, 40%, mean inward current: -7.6±2.9 pA) (**Figure 4*M***).

Together these optogenetic findings reinforced our pharmacological results showing that

Kiss1^{ARH} neurons are excited by MCR agonists. Additional studies would need to be done to

identify the cation conductance that is affected by the MC4R signaling cascade in Kiss1^{ARH}

neurons, but clearly we have established that there are excitatory glutamatergic and peptidergic

inputs from POMC to Kiss1[ARH] neurons.

Kiss1[AVPV/PeN] neurons are inhibited by MC4R agonists.

Having established a direct, excitatory projection from POMC to Kiss1[ARH] neurons, we next

examined POMC inputs to Kiss1[AVPVPeN] neurons. First, we used immunocytochemistry to label

fibers expressing αMSH to demonstrate that POMC fibers densely innervated the AVPV/PeN

area (**Figure 5*A***). Next, we recorded from AVPV neurons in slices taken from OVX+E2, POMC-

Cre mice injected with AAV1-DIO-YFP:ChR2 into the ARH (**Figure *5B***). We targeted cells along

the ventricle in the AVPV and PeN, surrounded by YFP terminals, and Kiss1[AVPV/PeN] neurons

were identified based on the expression of several endogenous conductances (I_{NaP}, T-type

calcium current, h-current) that are unique to these neurons (26, 36). We recorded a direct

glutamatergic synaptic response following optogenetic stimulation in six neurons, which was

further verified through pharmacological "rescue" of the synaptic response in the majority (5/6)

neurons tested (**Figure *5C***), indicating that POMC neurons make direct monosynaptic

connections with Kiss1[AVPV/PeN] neurons.

We next hypothesized that Kiss1[AVPV/PeN] neurons would be excited by exogenous application of

melanocortin agonists because the receptor is typically Gs-coupled (37, 38). We did whole-cell,

voltage clamp recordings from Kiss1[AVPV/PeN]-Cre::Ai32 neurons, which were isolated synaptically

by bathing the neurons in TTX (1 μM), from OVX females. For these experiments, we tested the

response to the high affinity melanocortin receptor agonist MTII (500 nM). Surprisingly, we

consistently measured an outward (inhibitory) current that could be reversed on washout of MTII

(**Figure 5*D***). This outward current was associated with an increase in a K^+ conductance based

on the current-voltage plot (*i.e.,* the outward current reversed at ~E_{K+}, **Figure 5E**). Also, the

inwardly-rectifying I/V could indicate that MC4R is coupled to activation of inwardly-rectifying K^+

channels as has been previously reported for MC4R signaling in pre-autonomic

parasympathetic neurons in the brainstem (39). We hypothesized that the coupling could be

modulated by E2 via a $G\alpha_q$-coupled membrane estrogen receptor (Gq-mER) as we have

previously demonstrated in POMC neurons (40-42). For these experiments, we again utilized

OVX females and targeted Kiss1[AVPV/PeN]-Cre::Ai32 neurons. Once in the whole-cell voltage-

clamp configuration, we perfused the slices with STX (10 nM), a selective ligand for the putative

Gq-mER (42), and tested the response to the MTII. Indeed, the outward current (inhibitory)

response to MTII was completely abrogated and even reversed by STX (**Figure 5*F, G***). The

short-term (bath) treatment with STX ensured that there was no desensitization of the Gq-mER

with longer-term (*in vivo*) treatment with E2. Therefore, estrogen receptor activation can rapidly

uncouple (*i.e.,* desensitize) the MC4R inhibitory response in Kiss1[AVPV/PeN] neurons. We would

predict that the intracellular signaling cascade for the heterologous desensitization is similar to

what we have elucidated in POMC neurons (42), but this will need to be determined in future

experiments. We next investigated the response in vehicle-treated, OVX females. Indeed,

Kiss1[AVPV/PeN] neurons were even more inhibited by the same MTII exposure (**Figure 5*G***).

Therefore, in contrast to Kiss1[ARH] neurons, the MC4R appears to be coupled to opening of K^+

channels in Kiss1[AVPV/PeN] neurons. To further establish whether the MTII was having pre- or

postsynaptic effects, we measured the glutamatergic mEPSCs before and after melanocortin

receptor activation. We found there was no significant effect on the frequency (**Figure 5H**), but

there was an effect on the amplitude (**Figure 5I**). This further supports a postsynaptic locus of

STX's effects. However, *posthoc* comparisons found STX increased amplitude compared to

both OVX and OVX+E. This result hints at a membrane delimited effect as STX is a much more

potent agonist for Gq-mER than E2 (43).

Discussion

Our findings show that melanocortins act directly on Kiss1 neurons to contribute to the

metabolic regulation of fertility, in line with previous reports showing that αMSH stimulates LH

release in a kisspeptin-dependent manner (14) and that Kiss1 neurons express the

melanocortin receptor MC4R (18-21). Several studies in humans have linked *Mc4r* mutations to

reproductive abnormalities, including precocious puberty (9), PCOS (10), and hypogonadism

(8). However, a direct association between *MC4R* mutations and reproductive function,

independent from the obese condition of these patients, has not been identified (6). In the

current study, we show that the reproductive impairments observed in MC4R deficient mice,

which replicate many of the conditions described in humans, are largely mediated by the direct

action of melanocortins via MC4R on Kiss1 neurons and not to their obese phenotype. This is

because the ablation of MC4R from Kiss1 neurons largely replicated the reproductive

impairments observed in MC4RKO female mice without inducing obesity, and the selective

reinsertion of MC4R into Kiss1 neurons of MC4RKO mice significantly improved their

reproductive function without changing their obese phenotype. Strikingly, puberty onset was

advanced in Kiss1-MC4RKO females. Our findings revealed a low melanocortin expression in

the hypothalamus of WT females during pubertal development, characterized by lower levels of

Agrp, Mc4r and *Mc3r* expression at the time of puberty onset (~PND30). This developmental

decline in the expression of melanocortin receptors is in line with the advanced timing of puberty

onset in females when MC4R is congenitally ablated from Kiss1 neurons. A recent study using

whole body MC3RKO mice showed a trend to delayed puberty onset (19), suggesting a larger

role of melanocortins in the timing of puberty onset. Here, we show that in females the

combination of MC3R and MC4R action may be necessary for pubertal development.

Interestingly, the MC4R-LoxTB and *Kiss1-cre:MC4R-LoxTB* mice do not show an advancement

in puberty onset as we observed in Kiss1-MC4RKO mice; however, these mice are obese at the

time of puberty onset already, which may affect the timing of sexual maturation independently of

MC4R action in Kiss1 neurons. Of note, the findings presented in this study do not rule out the

existence of additional sites of action of MC4R in other neuronal populations to regulate

reproduction, *e.g.,* GnRH neurons, which also express MC4R (16, 25). Indeed, the lack of full

recovery of the reproductive function in *Kiss1-cre:MC4R-LoxTB* mice suggests that MC4R in

Kiss1 neurons is necessary but not sufficient to achieve full reproductive capabilities.

In addition to early puberty onset, Kiss1-MC4RKO females displayed an impaired preovulatory

LH surge driving ovulation and had fewer corpora lutea in their ovaries compared to control

littermates, further supporting a role of MC4R in regulating ovulation. While MC4R has been

documented in the ovary (11), our data of i) impaired LH surge observed in the Kiss1-MC4RKO

females (**Figure 2M**), and ii) direct action of αMSH on Kiss1 ARC and AVPV/PeN neuronal

populations through MC4R (**Figures 4, 5**) support a CNS role for MC4R in the regulation of

fertility. However, an additional effect of MC4R at the level of the ovary cannot be ruled. One

possibility is that the genetic deletion of *Mc4r* from *Kiss1* gene may have affected the ovary.

However, to test this hypothesis, the co-expression of *Mc4r* and *Kiss1* in the ovary needs to be

investigated. Another possibility is that given the partial fertility of these mice, follicles may have

developed at different rates, potentially due to irregular secretion of gonadotropins. As a result,

fewer follicles would develop and ovulate. In adulthood, the reproductive phenotype observed in

Kiss1-MC4RKO and MC4R-loxTB female mice (increased LH, irregular estrous cycles, oligo-

ovulation, increased cystic follicles) correlates with the phenotype observed in PCOS mouse

models (22, 23). In fact, an association between *MC4R* mutations and PCOS has been reported

(10). However, our mouse models failed to display higher levels of circulating androgens or

AMH, two of the hallmarks of PCOS (24), suggesting that melanocortin signaling is unlikely to

be a main contributing factor in the development of this syndrome.

Our data revealing early puberty onset, augmented LH pulse and impaired LH surge driving

ovulation support a direct role of MC4R on the two primary Kiss1 populations regulating these

functions (ARH and AVPV/PeN). We further investigated this and found that MC4R signaling

excites Kiss1^{ARH} neurons and inhibits Kiss1$^{AVPV/PeN}$ neurons in an estradiol-dependent manner.

In Kiss1^{ARH} neurons, bath applied or optogenetically-evoked release of melanocortin agonists

induce a direct excitatory inward current. Our studies using whole cell voltage clamp in female

mice were able to detect a melanocortin-mediated inward current in Kiss1[ARH] neurons that

would increase excitability. Previously, an MCR-activation effect was not detected in cell-

attached recordings of diestrus females (14, 44). However, cell-attached recordings do not

measure a direct response in isolated Kiss1[ARH] neurons but rather the summation of multiple

synaptic inputs to Kiss1[ARH] neurons. Most importantly, we did not see a difference in the

excitatory response in synaptically isolated Kiss1[ARH] neurons from E2-treated versus vehicle-

treated, ovariectomized females. This emphasizes the importance of doing whole-cell recording

from isolated Kiss1 neurons. The modulatory actions of POMC[ARH] neuronal projections to

Kiss1[ARH] neurons can also occur through a) the excitatory action of glutamate in a process that

is facilitated in the presence of estradiol, in line with previous publications (33, 45), and b) the

inhibitory action of β-endorphin, suggesting that POMC[ARH] neurons are able to exert a precise

regulatory role of the GnRH pulse generator, *i.e.* Kiss1[ARH] neurons, in response to metabolic

challenges. In the AVPV/PeN, MC4R agonists inhibited Kiss1[AVPV/PeN] neurons in an estradiol-

dependent manner, which utilized a $G\alpha_q$-coupled membrane estrogen receptor (STX receptor)

to attenuate the inhibitory tone of MC4R in these neurons. Rapid estrogen signaling may act to

ease transitions between states. Membrane delimited E2 actions can quickly attenuate or

enhance coupling between receptors and signaling cascades. These effects will precede E2-

driven changes in gene expression that produce more stable alterations in signaling. This

combination of mechanisms will reduce any lag between rises in serum E2 and physiological

effects. Considering the abbreviated mouse reproductive cycle, parallel mechanisms acting on

different timescales are particularly important.

Our pharmacology data clearly demonstrates the estradiol-dependent action of MC4R in

Kiss1[AVPV/PeN] neurons, suggesting that similar to Kiss1[ARH] neurons, this population is regulated

by melanocortins. The effect of MC4R agonists on Kiss1 neurons in the presence of TTX,

suggests a direct synaptic effect without the need of channelrhodopsin-2(ChR2)-assisted circuit

mapping. Nonetheless, to further demonstrate this interaction, we provide evidence of profound

innervation of αMSH fibers and the presence of synaptic contact between POMC and

Kiss1$^{AVPV/PeN}$ neurons through the optogenetic stimulation of a glutamatergic response in

Kiss1$^{AVPV/PeN}$ neurons after low frequency stimulation of POMC terminals. We intentionally

avoided high frequency stimulation of POMC terminals to prevent the co-release of β-

endorphins (e.g., **Figure 4M**) and their possible additive effect on the inhibitory action of MC4R.

Future studies will be aimed at the characterization of the endogenous opioid pathways in the

induction of the LH surge.

To date, two instances of MC4R-mediated inhibition have been described in the central nervous

system: MC4R can Gi,o couple to open K+ (Kir6.2) channels in parasympathetic preganglionic

neurons in the brainstem (39), and MC4R can directly couple to K+ (Kv7.1) channels in the

hypothalamic paraventricular nucleus neurons (46). Interestingly, our findings describe an

inhibitory role of MC4R in the reproductive neuroendocrine axis. This steroid state dependent

effect is particularly relevant because in the context of reproduction, the two populations of

Kiss1 neurons are strikingly different in their response to estradiol, where peptide expression in

Kiss1^{ARH} and Kiss1$^{AVPV/PeN}$ neurons are inhibited or stimulated, respectively, in order to mount

the negative versus positive feedback of sex steroids (15). This differential regulation allows for

the episodic versus surge release of GnRH; however, the cellular mechanisms underlying these

opposing roles of Kiss1 neurons to the same stimulus, *i.e.* circulating E2 levels, remain

unknown. Although there were no differences in the magnitude of the electrophysiological

responses to MTII in Kiss1^{ARH} neurons in vehicle- versus E2-treated, ovariectomized females,

we cannot rule out that E2 affects other systems activated by the MC4R signaling cascade as

recently reported in MC4R expressing neurons of the ventromedial nucleus of the hypothalamus

(VMH) (47). Interestingly, puberty onset was advanced in Kiss1-MC4RKO females. This data

suggests that the inhibitory tone of MC4R signaling on Kiss1$^{AVPV/PeN}$ neurons might also be

involved in the timing of puberty onset, which in turn suggests a role for this female-specific

population of Kiss1 neurons in sexual maturation. Furthermore, our findings suggest that

metabolic cues, through the regulation of the melanocortin output onto Kiss1$^{AVPV/PeN}$ neurons,

are essential for the timing and magnitude of the GnRH/LH surge. This study suggests that the

melanocortin-kisspeptin pathway is one of the multiple pathways essential for the metabolic

regulation of the HPG axis. Future studies are warranted to fully elucidate synaptic input and

postsynaptic cascades in the POMC-Kiss1$^{AVPV/PeN}$ circuit.

Altogether, our data reveal a differential regulatory action of MC4R in the neural control of

GnRH/LH release, participating in both the surge (E2-treated, ovariectomized female) and the

pulse-like (ovariectomized female) modes of LH release through the cellular regulation of Kiss1

neurons, which translate into our findings on the conditional knockout of *Mc4r* from Kiss1

neurons (**Figure 6**). These findings are important because the reproductive abnormalities often

attributed to obesity in MC4R deficient patients may not be caused by the excess in body weight

but, at least in part, by a deficiency in MC4R signaling directly at the level of Kiss1 neurons that

affects predominantly the ability to mount a preovulatory LH surge.

Materials and methods:

In-vivo experimental procedures:

Generation of Kiss1-MC4RKO, Kiss1-cre:MC4R-LoxTB and MC4R-LoxTB Transgenic Mice

Kiss1-MC4RKO mice were generated by crossing *Kiss1-cre* knock-in mice and *MC4R$^{lox/lox}$* mice.

Kiss1-cre mice (RRID:MGI:6278139) were obtained from Dr. Richard Palmiter (University of

Washington, Seattle, WA) (48) and *Mc4r$^{lox/lox}$* (RRID:IMSR_JAX:023720) were a gift from Dr.

Brad Lowell (Beth Israel Deaconess Hospital, Boston, MA) (4). These mice were crossed to

generate Kiss1-MC4RKO mice lacking *Mc4r* expression selectively from Kiss1 neurons (*Kiss1-*

cre$^{+/-}$; *MC4R$^{lox/lox}$*) and their control littermates (MC4R$^{lox/lox}$). To generate *Kiss1-cre:MC4R-LoxTB*

and *MC4R-LoxTB* mice, *MC4R-LoxTB* mice were purchased from The Jackson Laboratory

(MC4R-LoxTB; catalog no. 006414). These mice present with a loxp-flanked transcriptional

blocking (LoxTB) sequence preventing normal endogenous gene transcription and translation

from the endogenous locus. Homozygous MC4R-loxTB/loxTB mice are devoid of functional

MC4R mRNA (MC4RKO mice), while the presence of Cre recombinase on the Kiss1 promotor

in *Kiss1-cre* mice will result in the removal of the transcription blocker and subsequent

expression of MC4R in tissue-specific sites (*i.e.* Kiss1 neurons) therefore resulting in the

generation of *Kiss1-cre*:MC4R-LoxTB (with *Mc4r* expression restored in Kiss1 neurons), their

obese control littermates *MC4R-LoxTB* mice (MC4RKO), and their WT controls (MC4R$^{+/+}$). To

rule out early embryonic recombination of the MC4R$^{loxTB/loxTB}$ or MC4R$^{lox/lox}$ alleles, we ran PCR

assays on tail DNA designed to detect wild-type allele (MC4R$^{+/+}$), undeleted lox (MC4R$^{lox/lox}$) or

loxTB alleles (MC4R$^{loxTB/loxTB}$). Genotyping was confirmed by sending tissue to Transnetyx, Inc.,

for testing by real-time polymerase chain reaction. Mice were housed in Harvard Medical School

Animal Resources facilities where they were fed standard mouse chow (Teklad F6 Rodent Diet

8664) and were given *ad libitum* access to tap water under constant conditions of temperature

(22–24°C) and light (12 hr light [07:00]/dark [19:00] cycle).

RNAscope in situ hybridization

To validate the Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and MC4R-LoxTB (MC4RKO) mouse

models and investigate the co-expression of *Kiss1* and *Mc4r* mRNA in these mice, dual

fluorescence ISH was performed using RNAscope (ACD, Multiplex Fluorescent v.2) according

to the manufacturer's protocol using the following probes: *Mc4r* (319181-C2) and *Kiss1*

(500141-C1). Brains (n=4/group) from WT OVX (for expression in the ARH) and OVX+E2 (for

expression in the AVPV/PeN) mice and OVX Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and

MC4RKO were removed fresh frozen on dry ice, and then stored at -80°C until sectioned. Five

sets of 20 μm sections in the coronal plane were cut on a cryostat, from the diagonal band of

Broca to the mammillary bodies, thaw mounted onto SuperFrost Plus slides (VWR Scientific)

and stored at -80°C until use. A single set was used for ISH experiment (adjacent sections 100

mm apart). Images were taken at 20x magnification of the sections containing AVPV, PeN, and

the three rostro-to-caudal levels of the ARH, and Kiss1 neurons expressing (*Kiss1-cre*:MC4R-

LoxTB mice) or lacking (Kiss1-MC4RKO mice) *Mc4r* were identified using ImageJ.

Reproductive maturation of Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and MC4R-LoxTB

mice

To assess the reproductive phenotype of mice with selective reinsertion of *Mc4r* in Kiss1

neurons (*Kiss1-cre*:MC4R-LoxTB, n=11), selective deletion of *Mc4r* from Kiss1 neurons (Kiss1-

MC4RKO, n=24), global deletion of *Mc4r* (MC4R-LoxTB, n=7); and their control MC4R$^{lox/lox}$

littermates (n=14), mice were weaned at post-natal day (PND) 21 and were monitored daily for

puberty onset. Females were monitored daily for vaginal opening (VO, indicative of the

complete canalization of the vaginal cavity) and for first estrus (first day with cornified cells

determined by daily morning vaginal cytology) after the day of VO. Body weight (BW) was

measured at the day of puberty onset to determine if changes in puberty onset could be due to

differences in BW. Estrous cyclicity was monitored in females by daily vaginal cytology, for a

period of 15 days, in 6-month-old mice and their respective control littermates (n=5/group).

Cytology samples were obtained every morning (9 am), placed on a glass slide and stained with

hematoxylin and eosin for determination of the estrous cycle stage under the microscope.

Fecundity test of Kiss1-MC4RKO females

Adult 6-month-old Kiss1-MC4RKO and control littermate female mice (n=9/group) were placed

with adult WT males proven to father litters for 90 days and time to deliver pups and number of

pups per litter were monitored.

Characterization of the estradiol-induced luteinizing hormone surge

Kiss1-MC4RKO (n=6) and control MC4R$^{lox/lox}$ littermate (n=5) adult female mice were subjected to bilateral ovariectomy (OVX) via abdominal incision under light isoflurane anesthesia. Immediately after OVX, capsules filled with E2 (1 ug/20g BW) were implanted subcutaneously via a small mid- scapular incision on the back. Five days later, mice were subcutaneously injected in the morning with estradiol benzoate (1 ug/20g BW) to produce elevated proestrus-like E2 levels (induced LH surge) on the following day (49). Blood samples were collected at 8am and 7pm; LH levels were stored at -80°C until measured via LH ELISA.

Ovarian histology and hormone measurements

Bilateral ovariectomy of 6-month-old Kiss1-MC4RKO (n=4), *Kiss1-cre*:MC4R-LoxTB, MC4R-LoxTB and their control littermates (n=5/group) was performed under light isoflurane anesthesia. Briefly, the ventral skin was shaved and cleaned, and one small abdominal incision was made. Once the ovaries were identified and excised, the muscle incision was sutured, and the skin was closed with surgical clips. Ovaries were stored in Bouin's fixative, sectioned, and stained with hematoxylin and eosin at the Harvard Histopathology Core. Corporal lutea were counted in the middle section of each ovary. Serum samples were also collected for analysis of testosterone and anti-müllerian hormone (AMH) levels in these mice. These hormones levels were measured at the University of Virginia Ligand Assay core with the Mouse & Rat Testosterone ELISA assay (reportable average range 10-1600 ng/dL; sensitivity of 10 ng/dL); AMH ELISA assay (reportable average range 0.2 – 15 ng/mL; sensitivity of 0.2 ng/mL).

LH pulsatile secretion profile in gonad intact Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and MC4R-LoxTB female mice

To assess the profile of LH pulses secretion, adult 6 months old gonad intact Kiss1-MC4RKO females (n=6), their control Mc4r$^{lox/lox}$ littermates (n=7); and *Kiss1-cre*:MC4R-LoxTB (n=5/group),

MC4R-LoxTB and their control WT females (n=4/group) were handled daily for 3 weeks to allow

acclimation to sampling conditions prior to the experiment. Pulsatile measurements of LH

secretion were assessed in diestrus by repeated blood collection through a single incision at the

tip of the tail. The tail was cleaned with saline and 4 µL of blood was taken at each time point

from the cut tail with a pipette. We collected sequential blood samples every 10 minutes over a

180-minute sampling period. Samples were immediately frozen on dry ice and stored at –80 °C

until analyzed with LH ELISA as previously described (50). The functional sensitivity of the

ELISA assay was 0.0039 ng/ml with a CV% of 3.3%.

LH pulses analysis

LH pulses in mice were analyzed using a custom-made MATLAB-based algorithm. The

MATLAB code includes a loop that determines LH pulses as any LH peak: (i) whose height is

20% greater than the heights of the 2 previous values; (ii) 10% greater than the height of the

following value; and (iii) the peak at the second time interval needs to be 20% greater than the

single value that comes before it to be considered a pulse, as we previously described (51).

LH pulsatility was assessed by measuring: (1) the total secretory mass, assessed by area under

the curve (AUC); (2) the LH pulse amplitude, calculated by averaging the 4 highest LH values in

the samples collection period for each animal; (3) the basal LH, calculated by averaging the 4

lowest LH values in the samples collection period for each animal; and (4) the total number of

pulses throughout the 180 minutes sampling period.

Immunohistochemistry

Animals and Treatment: Coronal brain blocks (2 mm each) from adult female C57BL/6 mice

(n=4) were fixed by immersion in 4% paraformaldehyde for ~ 8 hours, cryoprotected in 30%

sucrose solution, frozen in isopentane at –55°C, sectioned coronally on a cryostat at 20 µm, and

thaw-mounted onto Superfrost Plus slides (Fisher Scientific, Pittsburgh, PA).   The 20 µm

sections were stored at –20°C until used for immunocytochemistry.

Immunocytochemistry: The *s*ections were rinsed in PB (0.1M phosphate buffer, pH 7.4) for at

least 30 min. Next, sections were incubated with normal serum corresponding to the host for

the secondary antiserum (5 % normal serum with 0.3% Triton-X100 in PBS for 30 min), rinsed in

PB and then incubated for ~45 h at 4º C with a rabbit polyclonal antiserum against α-MSH

(1:2,500). The specificity of this antiserum has been documented (52). After rinsing, sections

were first incubated for 2-3 hours at room temperature with biotinylated donkey anti rabbit

gamma globulin (IgG; 1:500) and next with streptavidin-Alexa 488 (1:2500). Both the primary

and secondary antibodies were diluted in tris-(hydroxymethyl) aminomethane (0.5 %, Jackson

ImmunoResearch, Philadelphia, PA) in PB containing 0.7 % seaweed gelatin (Jackson

ImmunoResearch, Philadelphia, PA) and 0.5 % Triton X-100 and 3% bovine serum albumin

(BSA; Jackson ImmunoResearch, Philadelphia, PA) adjusted to pH 7.6. Following a final rinse

overnight, slides were coverslipped with gelvatol containing the anti-fading agent, 1,4-

diazabicyclo(2,2)octane (DABCO; Cold Spring Harbor Protocols, 2006).

Imaging: Photomicrographs of labeling were initially acquired using a Nikon E800 fluorescent

microscope (Eclipse E800; Nikon Instruments, Melville, NY) equipped with a fiber illuminator

(Intensilight C-HGFI; Nikon Instruments) and a high-definition digital microscope camera head

(DS-Fi1; Nikon Instruments) interfaced with a PC-based camera controller (DS-U3; Nikon

Instruments).

Real time quantitative PCR

i) To investigate the changes in the expression of the melanocortin genes *Agrp*, *Pomc, Mc3r*

and *Mc4r* in the MBH during development in prepubertal and pubertal WT females at ages P10

(n=6), P15 (n=6), P22 (n=6), and P30 (n=5), and ii) evaluate the gene expression profile of

Pdyn, *Kiss1*, *Tac2*, *Tacr3*, in the ARH of adult intact (in diestrus) female Kiss1-MC4RKO (n=6)

and their control MC4R$^{lox/lox}$ littermates (n=3), the brains were removed and rapidly embedded in

Tissue-Tek, frozen in –30°C isopentane solution and stored at –80°C until use. Frozen tissue

punches were recovered through MBH with a 1 mm diameter canula (53). These tissue punches

encompassed the whole MBH from the WT females (figure 1D) and the ARH from the Kiss1-

MC4RKO and their control females (Figure 2I). The tissues were homogenized, and total RNA

was isolated using TRIzol reagent (Invitrogen) followed by chloroform/isopropanol extraction.

RNA purity and concentration were measured via an absorbance spectrophotometer (260/280

nm > 1.8; NanoDrop 1000, Thermo Fisher Scientific). Total RNA (1 μg) was reverse transcribed

to cDNA using random hexamers (High-Capacity cDNA Synthesis Kit, Life Technologies).

Quantitative real-time PCR assays were performed using SYBR Green RT-qPCR master mix

(Applied Biosystems) and analyzed using ABI Prism 7000 SDS software (Applied Biosystems).

The cycling conditions were: 2 minutes incubation at 95°C (hot start), 45 amplification cycles

(95°C for 30 seconds, 60°C for 30 seconds, and 45 seconds at 75°C, with fluorescence

detection at the end of each cycle), followed by melting curve of the amplified products obtained

by ramped increase of the temperature from 55°C to 95°C to confirm the presence of single

amplification product per reaction. PCR specificity was verified by melting curve analysis and

agarose gel electrophoresis. Each sample was run in duplicate to obtain an average cycle

threshold (CT) value and relative expression of each target gene was determined using the

comparative Ct method (54). The data were normalized to Hypoxanthine Guanine

Phosphoribosyltransferase (*Hprt)* expression levels in each sample. Results were expressed as

fold differences in relative gene expression with respect to i) P10 for melanocortin genes

expression analysis during development in WT mice, and ii) controls for the KNDy genes

expression in female Kiss1-MC4RKO mice. The primers used are listed in Table 1.

Gene	Gene length (pb)	Accession #	Primers	Location (nt)	Sequence

Hprt	352	NM_013556.2	Hprt-F	704-728	CCTGCTGGATTACATTAAAGCGCTG
			Hprt-R	377-401	GTCAAGGGCATATCCAACAACAAAC
Agrp	136	NM_001271806.1	Agrp-F	466-488	GCCTCAAGAAGACAACTGCAGAC
			Agrp-R	580-601	AAGCAGGACTCGTGCAGCCTTA
Pomc	138	NM_001278584.1	Pomc-F	187-208	CCATAGATGTGTGGAGCTGGTG
			Pomc-R	303-324	CACCTCCGTTGCCAGGAAACAC
Mc4r	101	NM_016977.4	Mc4r-F	552-570	CCCGGACGGAGGATGCTAT
			Mc4r-R	632-652	TCGCCACGATCACTAGAATGT
Pdyn	200	NM_018863.4	Pdyn-F	45-64	ACAGGGGGAGACTCTCATCT
			Pdyn-R	223-244	GGGGATGAATGACCTGCTTACT
Kiss1	129	AF472576.1	Kiss1-F	147-166	GCTGCTGCTTCTCCTCTGTG
			Kiss1-R	256-275	TCTGCATACCGCGATTCCTT
Tac2	234	NM_001199971.1	Tac2- F	238-257	GCTCCACAGCTTTGTCCTTC
			Tac2- R	452-471	GCTAGCCTTGCTCAGCACTT
Tacr3	159	NM_021382.6	Tacr3-F	759-779	GCCATTGCAGTGGACAGGTAT
			Tacr3-R	898-917	ACGGCCTGGCATGACTTTTA

Data analysis

Statistical data are expressed as means ± SEM, where n represents the number of animals in each study group. The significance of differences between groups was evaluated using unpaired two-tailed Student's t test, or a one- or two-way ANOVA test (with *post hoc* comparisons). Significance level was set at $p<0.05$. All analyses were performed with GraphPad Prism Software, Inc (San Diego, CA).

In-vitro experimental procedures:

Animals:

All animal procedures described in the electrophysiology studies were performed in accordance

with institutional guidelines based on National Institutes of Health standards and approved by

the Institutional Animal Care and Use Committees at Oregon Health and Science University and

Appalachian State University. *Kiss1-cre* (v2) mice (48) were crossed with Ai32 (55) or C57B6J

mice. *Pomc-Cre* mice (RRID:IMSR_JAX:005965) (56) were crossed with wildtype C57B6J

(RRID:IMSR_JAX:000664) mice. The Ai32 cross was not used with *Pomc-Cre* mice because

the gene is transiently expressed during development in some cells fated to be Kiss1 or AgRP

cells (57) and early recombinant events lead to persistent expression of the ChR2-mCh fusion

protein in non-POMC cells. However, we have previously shown that AAV-driven expression in

adult *Pomc-cre* animals is restricted to β-endorphin labeled cells in the ARH (45), avoiding this

non-specificity issue. All colonies were maintained onsite under controlled temperature (21-23

ºC) and photoperiod (12:12-h light-dark cycle 0600 to 1800) while receiving ad libitum food

(5L0D; LabDiet, St. Louis, MO) and water access. Following surgeries, mice received a *s.c.*

dose of 4-5 mg/kg carprofen (Rimadyl; Pfizer Animal Health, New York, NY) and given at least 1

week of recovery.

Ovariectomies:

Seven to ten days prior to each experiment, ovaries were removed as described previously

while under isoflurane anesthesia (45). Two days before experiments females received either

an injection of sesame oil (50 µl, *sc*; Sigma-Aldrich, St. Louis, MO) or a priming dose (0.25 µg

/50 µl sesame oil, *sc*) of E2 benzoate (Sigma-Aldrich, St. Louis, MO) in the morning. On the

following day, oil or a high (1.5 µg) dose of E2 benzoate, which generates an induced LH surge,

was administered (58). Circulating levels of E2 were verified by the uterine weights (<25 mg for

OVX and >95 mg for OVX E2-treated) at the time of euthanasia (58).

AAV Delivery:

Bilateral ARH injections of AAV1-Ef1a-DIO-ChR2:mCherry (RRID:Addgene_20297) or AAV1-

Ef1a-DIO-ChR2:YFP (RRID:Addgene_20298) were performed on adult *Kiss1-cre* mice or

Pomc-cre mice on a stereotaxic frame under isoflurane anesthesia. ARH injection coordinates

were anteroposterior (AP): −1.10 mm, mediolateral (ML): ± 0.30 mm, dorsoventral (DL): −5.80

mm (surface of brain z = 0.0 mm); 400 nl of the AAV (2.0×10^{12} particles/ml) was injected (100

nl/min) into each position. Mice were given carprofen for analgesia and allowed to recover for at

least two weeks before euthanasia.

Visualized whole-cell patch recordings:

Hypothalamic coronal brain slices (240 um) were made from female mice, using a Leica

VT1000S vibratome, in ice cold cutting solution bubbled with O_2/CO_2 (95%/5%). Slices were

then transferred to a holding chamber with artificial cerebrospinal fluid bubbled with the same

gas mix and allowed to recover for at least 1 hour. For recordings, slices were placed in a

perfusion chamber and visualized with an Olympus BX51W1 using either differential infrared

contrast or oblique illumination. Kiss1 neurons in the ARH or AVPV/PeN were targeted for

electrophysiological recordings as done previously and described below (59, 60). Final

concentration was calculated based on the drug in the known volume of the bath. Perfused

drugs (TTX, naloxone, and STX) were constantly circulated and given at least 10 minutes to

reach maximal effect. Focal application of drugs (MTII and THIQ) was done with the pump

stopped (10-20 minutes). 0.3 to 1 μl was directly added to the known volume of the bath to

achieve the desired final concentration. This approach enables precise timing of recordings and

reduces the risk of receptor desensitization.

Solutions/drugs:

696   Standard vibratome slicing, external, and internal recording solutions were utilized as previously

697   described (59, 60). Tetrodotoxin was purchased from Alomone Labs (Jerusalem, Israel),

698   Melanotan II and αMSH from Tocris (Minneapolis, MN). TEA, 4-AP, 17β-estradiol benzoate, and

699   Naloxone were purchased from Millipore-Sigma. STX was produced by AAPharmaSyn, LLC

700   (Ann Arbor, MI) under contract.

Electrophysiology data analysis:

703   Electrophysiological data were analyzed using Clampfit 10/11 (Molecular Devices) and Prism

704   7/10 (Dotmatics). All values are expressed as Mean ± SEM. Comparisons between two groups

705   were made using un-paired Student's t-test or between multiple groups using an ANOVA (with

706   *post hoc* comparisons) with p-values < 0.05 considered significant. When variances differed

707   significantly, Mann-Whitney U test was used instead.

Targeting of Kiss1 neurons for electrophysiological recordings.

For non-optogenetic experiments, brain slices were taken from AAV injected Kiss1-Cre AAV or

Kiss1xAi32 female mice. Ai32 mice (RRID:IMSR_JAX:024109, C57BL/6 background) carry the

floxed ChR2 (H134R)-EYFP gene in their Gt(ROSA)26Sor Locus (55), allowing its expression

in a Cre-dependent manner. Due to concerns of nonspecific expression (61), we previously

validated this model using single cell RT-PCR and documented that *Kiss1* mRNA was

detectable in 99% of individually harvested eYFP cells (n=126). In addition, we have used both

AAV injected Kiss1-Cre AAV or Kiss1xAi32 female mice and found no differences in

electrophysiological results (59). We used AAV injected POMC-Cre female mice and avoided

small soma, high input resistance (>800 MΩ), ventrally located cells that are typically

NPY/AgRP neurons. Instead, we targeted larger, more dorsomedial neurons in the ARH while

avoiding fluorescent (POMC) cells. Unlike ARH Kiss1 neurons, POMC neurons do not make

reciprocal projections, so uninfected POMC neurons do not display monosynaptic EPSCs in

response to optogenetic stimulation. Next, we used a ramp IV protocol to probe for the presence of a persistent sodium current (I_{NaP}). While this current is more prevalent in the AVPV Kiss1 population, Kiss1 neurons are the only ARH neurons to display this electrophysiological "fingerprint" of a pronounced I_{NaP} paired with a high capacitance and low input resistance (26). Finally, we also harvested the cytosol at the end of all recordings and measured the expression of Kiss1. Only cells that expressed a persistent sodium current and/or expressed *Kiss1* mRNA were included in final analysis (n = 40 ARH & 6 AVPV).

Acknowledgments:

The authors would like to recognize the excellent technical expertise of Ms. Martha A. Bosch (tissue preparation, immunohistochemical procedure and single cell RT-PCR of recorded neurons). In addition, we thank Dr. Rona Carroll for her assistance at Harvard Medical School and Mr. Cole Martinson, a student worker in the Ronnekleiv/ Kelly laboratories, for his assistance with genotyping and care of mouse colonies at OHSU. The University of Virginia Center for Research in Reproduction Ligand Assay and Analysis Core is supported by the Eunice Kennedy Shriver NICHD/NIH (NCTRI) Grant P50-HD28934. This work was supported by PHS MPI grant DK68098 to MJK and OKR; HD090151, HD099084 and DK133760 to VMN, and The Charles A. King Trust Postdoctoral Research Fellowship Award, The Lalor Foundation Postdoctoral Fellowship Award, the Women's brain initiative Fellowship Award, the ROSA SCORE pilot grant (supported by NIH Research Grant U54 AG062322 funded by The National Institute on Aging (NIA) and Office of Research on Women's Health (ORWH)) and the IBRO-ISN Research Fellowship Award to RT, and startup funds from Appalachian State University to TLS.

Authors contribution:

RT, TLS, MJK and VMN conceived and designed the study; RT, TLS, KF, CJH, EM, AG, KW, SL, EAM, OKR performed the experiments. RT, TLS, OKR, MJK and VMN analyzed data. RT, TLS and VMN wrote the manuscript with input from all the authors.

Declaration of interests:

The authors declare they have no competing interest.

Figure legends:

Figure 1. *Mc4r* expressed in Kiss1 neurons determines the timing of puberty onset.

Kiss1-MC4RKO females display advanced puberty onset, assessed by daily monitoring of vaginal opening **(A)** and first estrus **(B)**, as documented by cumulative percent and mean age of animals at vaginal opening **(A)** and first estrus **(B)** in Kiss1-MC4RKO females (n=24) compared to WT littermates (n=19). *p<0.05 by Student's t-test. Data presented as the mean± SEM. **(C)** Female Kiss1-MC4RKO (n=19) have normal body weight at the time of puberty onset compared to their WT littermates (n=15). **(D)** Ontogeny expression of melanocortin genes (*Agrp*, *Pomc*, *Mc4r and Mc3r*) in the ARH of WT female mice at different postnatal pre-pubertal and pubertal ages: P10, P15, P22 and P30, normalized to the housekeeping gene *Hprt*. (*n values*: females at P10 (n=6), P15 (n=6), P22 (n=6), and P30 (n=5)). Groups with different letters are significantly different (p<0.05), as determined by one-way ANOVA followed by the Student-Newman-Keuls test. Data presented as the mean± SEM.

Figure 2. Deletion of MC4R from Kiss1 neurons impairs fertility in Kiss1-MC4RKO females.

(A) Female Kiss1-MC4RKO (n=19) have normal body weight from weaning and until post-natal day (PND) 150 compared to their WT littermates (n=15). **(B)** Kiss1-MC4RKO females displayed irregular estrous cycles with a longer time in diestrus and a shorter time in estrus compared to control females. ** $P < 0.01$ Two-way ANOVA. **(C, D)** Pattern of LH pulsatility was analyzed in gonad intact Kiss1-MC4RKO (n=6) and control (n=7) females. # Represents LH pulses. **(E)** LH total secretory mass assessed by area under the curve (AUC), **(F)** LH pulse amplitude, **(G)** Basal LH and **(H)** Total number of pulses /180 min were analyzed. *p<0.05 by Student's t-test. The expression of the KNDy genes *Pdyn* **(I)**, *Kiss1* **(J)**, *Tac2* **(K)** and *Tacr3* **(L)**, was assessed in the ARH of adult Kiss1-MC4RKO (n=6) and control (n=3) females. **p<0.01 by Student's t-test. Data presented as the mean± SEM. **(M)** Kiss1-MC4RKO (n=6) and control (n=5) females were subjected to an LH surge induction protocol. LH samples were collected in the morning (AM [8 a.m.]) and evening (PM [7 p.m.]) after lights off. ** $P < 0.01$, Two-way ANOVA followed by Bonferroni *post hoc* test. **(N)** Ovarian histology shows a decrease in the number of corpora lutea (CL, **O**) and increase in the number of cystic follicles (CF, **P**) of Kiss1-MC4RKO compared to controls (n=5/group). # represents cystic follicles. Serum levels of Testosterone **(Q)** and AMH **(R)** in adult gonad intact Kiss1-MC4RKO and control females (n=5/group). Student t test for unpaired samples. Data presented as the mean± SEM. Kiss1-MC4RKO females display impaired fertility characterized by increased time to deliver pups **(S)** and decreased number of pups per litter **(T)**, (n=9/group). *p<0.05 and **p<0.01 by Student's t-test. Data presented as the mean± SEM.

Figure 3. Re-insertion of MC4R in Kiss1 neurons restores estrous cyclicity and LH pulsatility in Kiss1-cre:MC4R-LoxTB females.

(A, B) *Kiss1-cre*:MC4R-LoxTB (n=11) and MC4R-loxTB (MC4RKO) (n=7) displayed normal puberty onset as compared to their control littermates (n=14), as documented by cumulative

percent and mean age of animals at vaginal opening and first estrus. **(C)** *Kiss1-cre*:MC4R-

LoxTB and MC4R-loxTB both had significantly higher body weights at the time of puberty onset

compared to their controls. **P < 0.01, *** P < 0.001 by one way ANOVA. **(D)** MC4R-loxTB

(n=7) and *Kiss1-cre*:MC4R-LoxTB (n=11) females displayed significantly higher body weight

than their littermates (n=10) from post-natal day 30 onwards, *p<0.05 and **P < 0.01. Data

presented as the mean± (SEM). **(E)** MC4R-loxTB females displayed irregular estrous cycles,

presenting longer time in diestrus and shorter time in estrus compared to control females, while

Kiss1-cre:MC4R-LoxTB females displayed regular estrous cyclicity, similar to controls. *p<0.05,

**P < 0.01, *** P < 0.001, ****P < 0.0001, Two-way ANOVA. Data presented as the mean±

SEM. **(F)** Pattern of LH pulsatility analyzed in gonad intact MC4R-loxTB (n=4), *Kiss1-cre*:MC4R-

LoxTB (n=5) and control littermates females (n=5). LH samples were collected every 10 min for

180 minutes; # represents LH pulses. **(G)** LH total secretory mass, **(H)** LH pulse amplitude, **(I)**

basal LH, and **(J)** total number of pulses /180 min were assessed. *p<0.05, **P < 0.01 by one

way ANOVA. **(K)** Representative samples of ovarian histology from MC4R-loxTB, *Kiss1-*

cre:MC4R-LoxTB and control females (n=5/group); CL: Corpora lutea. Data are presented as

the mean± SEM. **(L)** Ovarian histology showed a significant decrease in the number of corpora

lutea in the MC4RKO and *Kiss1-cre*:MC4R-LoxTB compared to controls. Groups with different

letters are significantly different. Serum levels of **(M)** Testosterone and **(N)** AMH in adult gonad

intact females MC4R-loxTB, *Kiss1-cre*:MC4R-LoxTB and their control littermates. *p<0.05, ****P

< 0.0001, One Way ANOVA. Data are presented as the mean± SEM.

Figure 4. Kiss1[ARH] neurons are activated by melanocortin agonists and respond to

optogenetic stimulation of POMC neurons.

(A) Slices were taken from *Kiss1-cre* injected (confocal image) brains to target fluorescent cells

(arrowhead) for recording (white outline over electrode). **(B)** Whole cell voltage clamp recording

of Kiss1[ARH] neurons following the direct application of the high affinity melanocortin receptor

agonist melanotan II (MTII, 250 nM) was added directly to the bath and the response was

compared between slices from OVX and OVX+E females. **(C)** While the average inward current

was slightly higher in the OVX+E state, this was not a significant difference (Student's t-test for

unpaired samples, p>0.05). **(D)** The MC4R-selective agonist THIQ (100 nM) was perfused, and

excitatory inward currents were generated in Kiss1^{ARH} neurons. **(E)** Images are of AAV-driven

labeling of POMC cells as seen through the confocal. **(F)** Using the AAV-driven expression of

channelrhodopsin in adult *Pomc-cre* mice, high frequency stimulation elicited a slow inward

current in Kiss1 neurons. **(G)** The identity of cells was confirmed through the presence of a

persistent sodium current (see (26)) and/or with RT-PCR of harvested cytoplasm showing *Kiss1*

expression. (Gel: MM=molecular marker, TC=tissue controls). **(H)** A direct synaptic projection

from POMC to Kiss1^{ARH} neuron (*posthoc* identified) was confirmed using the "rescue" protocol:

1, baseline glutamatergic responses were initially generated (black trace); 2, then action

potentials were eliminated by blocking voltage-gated sodium channels with TTX and the

postsynaptic response (red trace); 3, blockade of potassium channels facilitated calcium entry

into the terminal through ChR2 to release synaptic vesicles, "rescuing" the postsynaptic

glutamate response (green trace). **I)** Glutamatergic responses were also often observed,

particularly in the ventral ARH. As seen when targeting low input resistance neurons (i.e.,

Kiss1^{ARH}) with low frequency optogenetic stimulation (5 ms pulse, 50 ms inter-spike interval), the

first response in OVX + vehicle females was larger relative to the second response in OVX +

E2- treated female mice. Representative traces are the average of 30 sweeps. **(J)** The

averaged paired-pulse ratio was lower in recordings from estradiol-treated female mice,

indicating an increased release probability (Students t-test p<0.05,). **(K)** High frequency

optogenetic stimulation elicited a small inward current. **(L)** In a different cell, a high frequency

inward current was noted before constant perfusion of SHU9119 for 15 minutes (break between

traces). When the stimulation protocol was repeated, no inward current was elicited. **(M)** The

majority of high frequency responses were inward and only twice was an inhibitory outward

current recorded in identified Kiss1^{ARH} neurons. Perfusion of the non-selective opioid receptor

antagonist naloxone reversed the current, eliminating MCR-activation as the mechanism.

Figure 5. Kiss1$^{AVPV/PeN}$ neurons are inhibited by MC4R agonists in an E2-dependent

manner.

(A) Immunohistochemistry showing robust labeling of α-MSH fibers in the AVPV/PeN region in

OVX+E2 WT female mice. **(B)** Low and high power bright-field images taken during

electrophysiology recording in OVX+E2 POMC-Cre mice expressing YFP:ChR2 in the ARH

following AAV injection. (left) Low power image shows the location of recorded cells. (right)

Higher power image of area from white box inset displaying POMC fibers innervating the area.

(C) Fibers surrounded putative Kiss1-AVPV neurons (expressing I$_{NaP}$, IT and I$_h$) and optogenetic

stimuli was able to elicit postsynaptic currents that were eliminated with tetrodotoxin (1 μM), but

"rescued" with addition of K$^+$ channel blockers (4-AP and TEA) which indicates a monosynaptic

connection between POMC^{ARH} and Kiss1^{AVPV} neurons. **(D)** Whole-cell voltage clamp recordings

were made in ChR2-YFP positive cells in brain slices taken from *Kiss1-cre*:Ai32 female mice.

Bath application of melanotan II (MTII, 500 nM) generated an inhibitory outward current in a

Kiss1$^{AVPV/PeN}$ neuron from OVX female. As the cell was synaptically isolated using bath-applied

tetrodotoxin (TTX, 1 μM) this represents a direct effect. Washout of MTII while still in TTX

quickly led to a return to baseline RMP. **(E)** An IV relationship was plotted using voltage steps

before and after MTII administration. The crossing at -80 mV (~EK$^+$) indicates that the opening

of K+ channels underlie the MC4R inhibition of Kiss1^{AVPV} neurons. **(F)** In a subset of recordings

from OVX brains slices, the selective membrane estrogen receptor (Gq-mER) agonist STX (10

nM) was added to the bath for ~10 minutes prior to addition of MTII. STX pretreatment resulted

in either a greatly attenuated outward current or even an inward current as shown in this

example. **(G)** The mean outward current was significantly higher when recording in Kiss1$^{AVPV/Pen}$

neurons from brain slices from vehicle-treated, OVX females compared to E2-treated, OVX

females or acute STX-treated brain slices from OVX females (One-way ANOVA $F_{(2,19)}$=12.32,

p<0.001; Holm-Sidak *posthoc* comparisons found significant differences between all groups; *

p<0.05, *** p<0.001). **(H)** There was no difference in the frequency of mEPSCs after MTII,

calculated as a percent of baseline between groups. **(I)** However, there was a significant

difference in the mEPSC amplitude (calculated as a percent of baseline) between all groups:

main effect $F_{(2,19)}$ =10.58, p<0.001. *Posthoc* comparisons using Holm-Sidak found OVX+STX to

be different from both OVX (p<0.001) and OVX+E (p<0.05).

Figure 6. Working model of the neuronal input from POMC ARH to Kiss1 ARH and AVPV

populations.

POMC^{ARH} neurons excite Kiss1^{ARH} neurons via glutamate and αMSH release, which activate

glutamatergic ionotropic receptors and MC4R, respectively. In low E2 state, αMSH fibers project

to the AVPV/PeN, and MC4R inhibit Kiss1$^{AVPV/PeN}$ neurons via opening a K$^+$ channel. However,

in high E2 state, E2 enhances the glutamatergic excitation of both POMC and Kiss1 ARH

neurons.   E2-treatment rapidly abrogates (uncouples) the inhibitory MC4R signaling in

Kiss1$^{AVPV/PeN}$ neurons, allowing greater excitatory actions of Kiss1^{ARH} neurons via enhanced

glutamate release (26).

References:
897

1.   Health U. Infertility. 2020 [Available from: http://obgyn.ucla.edu/infertility
2.   Andermann ML, Lowell BB. Toward a Wiring Diagram Understanding of Appetite
Control. Neuron. 2017;95(4):757-78.
3.   Cone RD. Studies on the physiological functions of the melanocortin system. Endocr Rev.
2006;27(7):736-49.
4.   Shah BP, Vong L, Olson DP, Koda S, Krashes MJ, Ye C, et al. MC4R-expressing
glutamatergic neurons in the paraventricular hypothalamus regulate feeding and are
synaptically connected to the parabrachial nucleus. Proceedings of the National Academy of
Sciences of the United States of America. 2014;111(36):13193-8.

5.   Wang K, Mao W, Zhang X, Zhao Y, Fan K, Pan D, et al. Neuroanatomy of melanocortin-4
receptor pathway in the mouse brain. Open Life Sci. 2020;15(1):580-7.
6.   Farooqi IS, Keogh JM, Yeo GS, Lank EJ, Cheetham T, O'Rahilly S. Clinical spectrum of
obesity and mutations in the melanocortin 4 receptor gene. The New England journal of
medicine. 2003;348(12):1085-95.
7.   Balthasar N, Dalgaard LT, Lee CE, Yu J, Funahashi H, Williams T, et al. Divergence of
melanocortin pathways in the control of food intake and energy expenditure. Cell.
2005;123(3):493-505.
8.   Hainerová IA, Zamrazilová H, Sedláčková D, Hainer V. Hypogonadotropic hypogonadism
in a homozygous MC4R mutation carrier and the effect of sibutramine treatment on body
weight and obesity-related health risks. Obes Facts. 2011;4(4):324-8.
9.   Doulla M, McIntyre AD, Hegele RA, Gallego PH. A novel MC4R mutation associated with
childhood-onset obesity: A case report. Paediatr Child Health. 2014;19(10):515-8.
10.   Batarfi AA, Filimban N, Bajouh OS, Dallol A, Chaudhary AG, Bakhashab S. MC4R variants
rs12970134 and rs17782313 are associated with obese polycystic ovary syndrome patients in
the Western region of Saudi Arabia. BMC Med Genet. 2019;20(1):144.
11.   Chen X, Huang L, Tan HY, Li H, Wan Y, Cowley M, et al. Deficient melanocortin-4
receptor causes abnormal reproductive neuroendocrine profile in female mice. Reproduction
(Cambridge, England). 2017;153(3):267-76.
12.   Sandrock M, Schulz A, Merkwitz C, Schoneberg T, Spanel-Borowski K, Ricken A.
Reduction in corpora lutea number in obese melanocortin-4-receptor-deficient mice. Reprod
Biol Endocrinol. 2009;7:24.
13.   Cui L, Tan C, Huang L, Wang W, Huang Z, Geng F, et al. Dapagliflozin partially restores
reproductive function in MC4R KO obese female mice. The Journal of endocrinology.
2022;254(2):65-76.
14.   Manfredi-Lozano M, Roa J, Ruiz-Pino F, Piet R, Garcia-Galiano D, Pineda R, et al. Defining
a novel leptin-melanocortin-kisspeptin pathway involved in the metabolic control of puberty.
Mol Metab. 2016;5(10):844-57.
15.   Goodman RL, Herbison AE, Lehman MN, Navarro VM. Neuroendocrine control of
gonadotropin-releasing hormone: Pulsatile and surge modes of secretion. Journal of
neuroendocrinology. 2022;34(5):e13094.
16.   Israel DD, Sheffer-Babila S, de Luca C, Jo YH, Liu SM, Xia Q, et al. Effects of leptin and
melanocortin signaling interactions on pubertal development and reproduction. Endocrinology.
2012;153(5):2408-19.
17.   Manfredi-Lozano M, Roa J, Tena-Sempere M. Connecting metabolism and gonadal
function: Novel central neuropeptide pathways involved in the metabolic control of puberty
and fertility. Front Neuroendocrinol. 2018;48:37-49.
18.   Cravo RM, Margatho LO, Osborne-Lawrence S, Donato J, Jr., Atkin S, Bookout AL, et al.
Characterization of Kiss1 neurons using transgenic mouse models. Neuroscience. 2011;173:37-
56.
19.   Lam BYH, Williamson A, Finer S, Day FR, Tadross JA, Gonçalves Soares A, et al. MC3R
links nutritional state to childhood growth and the timing of puberty. Nature.
2021;599(7885):436-41.

20.   Villa PA, Ruggiero-Ruff RE, Jamieson BB, Campbell RE, Coss D. Obesity alters POMC and
kisspeptin neuron crosstalk leading to reduced luteinizing hormone in male mice. The Journal of
Neuroscience. 2024:e0222242024.

21.   Stephens SBZ, Kauffman AS. Estrogen Regulation of the Molecular Phenotype and Active
Translatome of AVPV Kisspeptin Neurons. Endocrinology. 2021;162(9).

22.   Moore AM, Prescott M, Campbell RE. Estradiol negative and positive feedback in a
prenatal androgen-induced mouse model of polycystic ovarian syndrome. Endocrinology.
2013;154(2):796-806.

23.   McCarthy EA, Dischino D, Maguire C, Leon S, Talbi R, Cheung E, et al. Inhibiting Kiss1
Neurons With Kappa Opioid Receptor Agonists to Treat Polycystic Ovary Syndrome and
Vasomotor Symptoms. J Clin Endocrinol Metab. 2022;107(1):e328-e47.

24.   Dewailly D, Barbotin AL, Dumont A, Catteau-Jonard S, Robin G. Role of Anti-Müllerian
Hormone in the Pathogenesis of Polycystic Ovary Syndrome. Frontiers in endocrinology.
2020;11:641.

25.   Roa J, Herbison AE. Direct regulation of GnRH neuron excitability by arcuate nucleus
POMC and NPY neuron neuropeptides in female mice. Endocrinology. 2012;153(11):5587-99.

26.   Zhang C, Bosch MA, Qiu J, Ronnekleiv OK, Kelly MJ. 17beta-Estradiol increases persistent
Na(+) current and excitability of AVPV/PeN Kiss1 neurons in female mice. Mol Endocrinol.
2015;29(4):518-27.

27.   Clements JD, Westbrook GL. Activation kinetics reveal the number of glutamate and
glycine binding sites on the N-methyl-D-aspartate receptor. Neuron. 1991;7(4):605-13.

28.   Stincic TL, Grachev P, Bosch MA, Rønnekleiv OK, Kelly MJ. Estradiol drives the
anorexigenic   activity   of   proopiomelanocortin   neurons   in   female   mice.   eNeuro.
2018;5(4):e0103-18.2018.

29.   Qiu J, Nestor CC, Zhang C, Padilla SL, Palmiter RD, Kelly MJ, et al. High-frequency
stimulation-induced peptide release synchronizes arcuate kisspeptin neurons and excited GnRH
neurons. eLife. 2016;5:e16246.

30.   Qiu J, Rivera HM, Bosch MA, Padilla SL, Stincic TL, Palmiter RD, et al. Estrogenic-
dependent glutamatergic neurotransmission from kisspeptin neurons governs feeding circuits
in females. eLife. 2018;7:e35656.

31.   Stincic TL, Qiu J, Connors AM, Kelly MJ, Rønnekleiv OK. Arcuate and Preoptic Kisspeptin
neurons exhibit differential projections to hypothalamic nuclei and exert opposite postsynaptic
effects on hypothalamic paraventricular and dorsomedial nuclei in the female mouse. eneuro.
2021:ENEURO.0093-21.2021.

32.   Herman MA, Ackermann F, Trimbuch T, Rosenmund C. Vesicular glutamate transporter
expression level affects synaptic vesicle release probability at hippocampal synapses in culture.
The Journal of Neuroscience. 2014;34(35):11781-91.

33.   Qiu J, Rivera HM, Bosch MA, Padilla SL, Stincic TL, Palmiter RD, et al. Estrogenic-
dependent glutamatergic neurotransmission from kisspeptin neurons governs feeding circuits
in females. Elife. 2018;7.

34.   Thornton JE, Loose MD, Kelly MJ, Rønnekleiv OK. Effects of estrogen on the number of
neurons expressing β-endorphin in the medial basal hypothalamus of the female guinea pig.
The Journal of comparative neurology. 1994;341:68-77.

35.   Petersen SL, Keller ML, Carder SA, McCrone S. Differential Effects of Estrogen and
Progesterone on Levels of POMC mRNA Levels in the Arcuate Nucleus: Relationship to the
Timing of LH Surge Release. Journal of Neuroendocrinology. 1993;5(6):643-8.

36.   Zhang C, Tonsfeldt KJ, Qiu J, Bosch MA, Kobayashi K, Steiner RA, et al. Molecular
mechanisms that drive estradiol-dependent burst firing of Kiss1 neurons in the rostral
periventricular preoptic area. American journal of physiology Endocrinology and metabolism.
2013;305(11):E1384-97.

37.   Mountjoy KG, Mortrud MT, Low MJ, Simerly RB, Cone RD. Localization of the
melanocortin-4 receptor (MC4-R) in neuroendocrine and autonomic control circuits in the
brain. Molecular Endocrinology. 1994;8(10):1298-308.

38.   Yu J, Gimenez LE, Hernandez CC, Wu Y, Wein AH, Han GW, et al. Determination of the
melanocortin-4 receptor structure identifies Ca^{2+} as a cofactor for ligand binding. Science.
2020;368(6489):428-33.

39.   Sohn JW, Harris LE, Berglund ED, Liu T, Vong L, Lowell BB, et al. Melanocortin 4
receptors reciprocally regulate sympathetic and parasympathetic preganglionic neurons. Cell.
2013;152(3):612-9.

40.   Kelly MJ, Loose MD, Ronnekleiv OK. Estrogen suppresses mu-opioid- and GABAB-
mediated hyperpolarization of hypothalamic arcuate neurons. The Journal of neuroscience : the
official journal of the Society for Neuroscience. 1992;12(7):2745-50.

41.   Lagrange AH, Ronnekleiv OK, Kelly MJ. The potency of mu-opioid hyperpolarization of
hypothalamic arcuate neurons is rapidly attenuated by 17 beta-estradiol. The Journal of
neuroscience : the official journal of the Society for Neuroscience. 1994;14(10):6196-204.

42.   Qiu J, Bosch MA, Tobias SC, Grandy DK, Scanlan TS, Ronnekleiv OK, et al. Rapid signaling
of estrogen in hypothalamic neurons involves a novel G-protein-coupled estrogen receptor that
activates protein kinase C. The Journal of neuroscience : the official journal of the Society for
Neuroscience. 2003;23(29):9529-40.

43.   Qiu J, Bosch MA, Tobias SC, Krust A, Graham SM, Murphy SJ, et al. A G-protein-coupled
estrogen receptor is involved in hypothalamic control of energy homeostasis. The Journal of
neuroscience : the official journal of the Society for Neuroscience. 2006;26(21):5649-55.

44.   Sabine Hessler XL, Allan E. Herbison. Direct inhibition of arcuate kisspeptin neurones by
neuropeptide Y in the male and female mouse. Journal of Neuroendocrinology. 2020.

45.   Stincic TL, Grachev P, Bosch MA, Ronnekleiv OK, Kelly MJ. Estradiol Drives the
Anorexigenic Activity of Proopiomelanocortin Neurons in Female Mice. eNeuro. 2018;5(4).

46.   Ghamari-Langroudi M, Digby GJ, Sebag JA, Millhauser GL, Palomino R, Matthews R, et al.
G-protein-independent coupling of MC4R to Kir7.1 in hypothalamic neurons. Nature.
2015;520(7545):94-8.

47.   Krause WC, Rodriguez R, Gegenhuber B, Matharu N, Rodriguez AN, Padilla-Roger AM, et
al. Oestrogen engages brain MC4R signalling to drive physical activity in female mice. Nature.
2021;599(7883):131-5.

48.   Padilla SL, Johnson CW, Barker FD, Patterson MA, Palmiter RD. A Neural Circuit
Underlying the Generation of Hot Flushes. Cell Rep. 2018;24(2):271-7.

49.   Dror T, Franks J, Kauffman AS. Analysis of multiple positive feedback paradigms
demonstrates a complete absence of LH surges and GnRH activation in mice lacking kisspeptin
signaling. Biology of reproduction. 2013;88(6):146.

50.   Steyn FJ, Wan Y, Clarkson J, Veldhuis JD, Herbison AE, Chen C. Development of a
methodology for and assessment of pulsatile luteinizing hormone secretion in juvenile and
adult male mice. Endocrinology. 2013;154(12):4939-45.

51.   Talbi R, Ferrari K, Choi JH, Gerutshang A, McCarthy EA, Dischino D, et al.
Characterization of the Action of Tachykinin Signaling on Pulsatile LH Secretion in Male Mice.
Endocrinology. 2021;162(8).

52.   Dave JR, Rubinstein N, Eskay RL. Evidence that b-endorphin binds to specific receptors in
rat peripheral tissues and stimulates the adenylate cyclase-adenosine 3',5'-monophosphate
system. Endocrinology. 1985;117:1389-96.

53.   Naulé L, Robert V, Parmentier C, Martini M, Keller M, Cohen-Solal M, et al. Delayed
pubertal onset and prepubertal Kiss1 expression in female mice lacking central oestrogen
receptor beta. Hum Mol Genet. 2015;24(25):7326-38.

54.   Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR.
Nucleic Acids Res. 2001;29(9):e45.

55.   Madisen L, Mao T, Koch H, Zhuo JM, Berenyi A, Fujisawa S, et al. A toolbox of Cre-
dependent optogenetic transgenic mice for light-induced activation and silencing. Nature
Neuroscience. 2012;15(5):793-802.

56.   Balthasar N, Coppari R, McMinn J, Liu SM, Lee CE, Tang V, et al. Leptin receptor signaling
in POMC neurons is required for normal body weight homeostasis. Neuron. 2004;42(6):983-91.

57.   Padilla SL, Carmody JS, Zeltser LM. Pomc-expressing progenitors give rise to antagonistic
neuronal populations in hypothalamic feeding circuits. Nature Medicine. 2010;16(4):403-5.

58.   Bosch MA, Tonsfeldt KJ, Ronnekleiv OK. mRNA expression of ion channels in GnRH
neurons: subtype-specific regulation by 17beta-estradiol. Molecular and cellular endocrinology.
2013;367(1-2):85-97.

59.   Qiu J, Nestor CC, Zhang C, Padilla SL, Palmiter RD, Kelly MJ, et al. High-frequency
stimulation-induced peptide release synchronizes arcuate kisspeptin neurons and excites GnRH
neurons. Elife. 2016;5.

60.   Stincic TL, Qiu J, Connors AM, Kelly MJ, Ronnekleiv OK. Arcuate and Preoptic Kisspeptin
Neurons Exhibit Differential Projections to Hypothalamic Nuclei and Exert Opposite
Postsynaptic Effects on Hypothalamic Paraventricular and Dorsomedial Nuclei in the Female
Mouse. eNeuro. 2021;8(4).

61.   Qiu J, Stincic TL, Bosch MA, Connors AM, Kaech Petrie S, Ronnekleiv OK, et al. Deletion
of Stim1 in Hypothalamic Arcuate Nucleus Kiss1 Neurons Potentiates Synchronous GCaMP
Activity and Protects against Diet-Induced Obesity. The Journal of neuroscience : the official
journal of the Society for Neuroscience. 2021;41(47):9688-701.

1072