POMC neurons control fertility through differential signaling of MC4R in Kisspeptin neurons

Authors:

Rajae Talbi[1,2*], Todd L. Stincic[4,8*], Kaitlin Ferrari[2], Choi Ji Hae[2], Karol Walec[2], Elizabeth Medve[2], Achi Gerutshang[2], Silvia León[1,2], Elizabeth A. McCarthy[1,2], Oline K. Rønnekleiv[3,4], Martin J. Kelly[3,4,6] and Víctor M. Navarro[1,2,5,6]

[1]Harvard Medical School, Boston, MA, USA; [2]Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital, Boston, MA, USA; [3]Department of Chemical Physiology and Biochemistry, Oregon Health & Science University, Portland, OR, USA; [4]Division of Neuroscience, Oregon National Primate Research Center, Beaverton, OR, USA; [5]Harvard Program in Neuroscience, Boston, MA, USA

*These authors contributed equally
[6] Corresponding authors
[8]Present address: Department of Biology, Appalachian State University, Boone, NC 28608.

Correspondence:

Victor M. Navarro, PhD, Division of Endocrinology, Diabetes and Hypertension, Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, 221 Longwood Avenue, Room 219, Boston, Massachusetts 02115. E-mail: vnavarro@bwh.harvard.edu.

Martin J. Kelly, PhD, Department of Chemical Physiology and Biochemistry, Oregon Health & Science University, 3181 SW Sam Jackson Park Road, Portland, OR 97239. E-mail: kellym@ohsu.edu .

Keywords:

Kiss1 neurons, POMC neurons, MC4R, Fertility, Ovulation, Obesity, Estradiol.

Abstract

Inactivating mutations in the melanocortin 4 receptor (*MC4R*) gene cause monogenic obesity.

Interestingly, female patients also display various degrees of reproductive disorders, in line with

the subfertile phenotype of MC4RKO female mice. However, the cellular mechanisms by which

MC4R regulates reproduction are unknown. Kiss1 neurons directly stimulate gonadotropin-

releasing hormone (GnRH) release through two distinct populations; the Kiss1^{ARH} neurons,

controlling GnRH pulses, and the sexually dimorphic Kiss1$^{AVPV/PeN}$ neurons controlling the

preovulatory LH surge. Here, we show that *Mc4r* expressed in Kiss1 neurons is required for

fertility in females. *In vivo*, deletion of *Mc4r* from Kiss1 neurons in female mice replicates the

reproductive impairments of MC4RKO mice without inducing obesity. Conversely, reinsertion of

Mc4r in Kiss1 neurons of MC4R null mice restores estrous cyclicity and LH pulsatility without

reducing their obese phenotype. *In vitro*, we dissect the specific action of MC4R on Kiss1^{ARH} vs

Kiss1$^{AVPV/PeN}$ neurons and show that MC4R activation excites Kiss1^{ARH} neurons through direct

synaptic actions. In contrast, Kiss1$^{AVPV/PeN}$ neurons are normally inhibited by MC4R activation

except under elevated estradiol levels, thus facilitating the activation of Kiss1$^{AVPV/PeN}$ neurons to

induce the LH surge driving ovulation in females. Our findings demonstrate that POMC^{ARH}

neurons acting through MC4R, directly regulate reproductive function in females by stimulating

the "pulse generator" activity of Kiss1^{ARH} neurons and restricting the activation of Kiss1$^{AVPV/PeN}$

neurons to the time of the estradiol-dependent LH surge, and thus unveil a novel pathway of the

metabolic regulation of fertility by the melanocortin system.

Introduction

Obesity rates have skyrocketed in Western societies in the last decades, resulting in an alarming rise in comorbidities that place a significant burden on healthcare systems. The increase in obesity correlates with a decrease in fertility rates, leading to conception challenges that are experienced by approximately 15% of couples in the United States currently [1].

The melanocortin 4 receptor (MC4R) binds α-melanocyte stimulating hormone (αMSH), an agonist product of the pro-opiomelanocortin (*Pomc*) gene, and the inverse agonist of the agouti-related peptide (AgRP) to regulate food intake and energy expenditure [2,3]. While the role of MC4R on food intake is largely mediated by neurons located in the paraventricular nucleus of the hypothalamus (PVN) [4], its expression in the brain is widespread [5] with the specific function of the different MC4R-expressing neurons in areas beyond the PVN remaining to be fully explored. Inactivating mutations in *MC4R* are a leading cause of monogenic obesity in humans and mice [6]. Scant evidence in humans shows an association between *MC4R* mutations and higher incidence of hypogonadotropic hypogonadism (HH) [7], alterations in the timing of puberty onset [8], and polycystic ovary syndrome (PCOS) [9]; however, conflicting reports exist [6]. Therefore, the role of MC4R signaling in reproductive function in humans remains controversial despite the clear association found in mice. Supporting this role of MC4R, *Mc4r* null mice display an array of reproductive abnormalities that largely affects females, characterized by irregular estrous cycles, disrupted luteinizing hormone (LH) secretion, reduced corpora lutea and reduced fertility [10-12]. Further evidence from mice demonstrates that MC4R agonists robustly increase LH release in a kisspeptin-dependent manner [13].

Kisspeptin is the most potent gonadotropin releasing hormone (GnRH) secretagogue known to date and it is mainly produced in two distinct neuronal populations. Kiss1 neurons of the arcuate nucleus of the hypothalamus (Kiss1[ARH]), present in both sexes, control the pulsatile (tonic) release of GnRH, and sex steroids attenuate their release of kisspeptin. Kiss1 neurons of the

anteroventral periventricular continuum area (Kiss1$^{AVPV/PeN}$), are predominantly present in

females, and are responsible for generating the preovulatory GnRH/LH surge essential for

ovulation [14]. Despite these critical roles of both populations of Kiss1 neurons for reproduction,

the underlying mechanisms that determine how each Kiss1 population responds differently to

sex steroids to regulate the GnRH tonic vs surge release remains unresolved.

In rodents, compelling evidence indicates a close interaction between Kiss1 neurons and the

melanocortin system: 1) fibers from POMC neurons in the arcuate nucleus (POMC^{ARH}) project to

and juxtapose Kiss1^{ARH} neurons [13]; 2) the melanocortin signaling through MC4R contributes to

the permissive role of leptin on puberty onset [13,15,16]; and 3) MC4R expression on both Kiss1^{ARH}

[17-19] and Kiss1$^{AVPV/PeN}$ [17,20]. Altogether, this evidence suggests a clear role for MC4R in Kiss1

neurons, in the control of reproductive maturation and fertility.

In this study, we investigated the contribution of MC4R signaling versus obesity *per se* in the

etiology of the reproductive impairments observed in *Mc4r* null mice, which could explain similar

impairments observed in humans. Using genetic mouse models with specific deletion or

reinsertion of *Mc4r* in Kiss1 neurons, we show that MC4R action in Kiss1 neurons is necessary

for normal reproductive function in female mice. Whole cell voltage clamp recordings evidenced

an excitatory action of MC4R on Kiss1^{ARH} neurons and an estradiol-dependent inhibitory action

on Kiss1$^{AVPV/PeN}$ neurons, with important implications for the timing of the preovulatory LH surge.

Results

***Mc4r* within Kiss1 neurons determines the timing of puberty onset in females.**

The expression of *Mc4r* within Kiss1^{ARH} and Kiss1$^{AVPV/PeN}$ neurons has already been described

elsewhere [17-20], suggesting a role for MC4R in the regulation of fertility. To further investigate the

specific role of MC4R in Kiss1 neurons, we generated a Kiss1-specific MC4R knockout mouse

model (Kiss1-MC4RKO). The specific deletion of *Mc4r* from Kiss1 neurons, as well as the

absence of global recombination, were confirmed through RNAscope. While Kiss1 neurons of

Kiss1-MC4RKO mice lack *Mc4r* transcript compared to their control littermates (**Figure S1A**),

Mc4r was detectable in the PVN of all mice, which is a major site of *Mc4r* expression in the brain

in the regulation of metabolism, therefore supporting the specific deletion of *Mc4r* only within

Kiss1 neurons (**Figure S1B**).

Puberty onset was assessed daily from weaning age through the monitoring of vaginal opening

(VO) and first estrus (FE). Kiss1-MC4RKO females showed a significant advancement in the

age of VO (P=0.0150, Kiss1-MC4RKO: 24.67± 0.3978 vs controls: 26.32±0.5301) and FE

(P=0.0341, Kiss1-MC4RKO: 32.63±1.489 vs controls: 37.84±1.903) compared to their littermate

controls (**Figure 1A, B**). Body weight of Kiss1-MC4RKO females at the age of puberty onset

was similar between groups (P=0.2596, Kiss1-MC4RKO: 12.41±0.3041 vs controls:

12.89±0.2631) (**Figure 1C**). These data suggest that melanocortin signaling on Kiss1 neurons

participates in the timing of puberty onset in females. Since the role of the melanocortin system

on puberty onset is largely unexplored, we evaluated the expression profile of the main

components of this system in the hypothalami of WT female mice at postnatal days (PND) 10,

15, 22 and 30. Interestingly, expressions of *Agrp*, *Mc3r* and *Mc4r* were significantly lower at the

time of puberty onset (PND30) compared to earlier developmental ages (**Figure 1D**). This

suggests that a decrease in the hypothalamic melanocortin tone drives puberty onset, in line

with the advancement in the age of VO and FE observed in Kiss1-MC4RKO female mice

(**Figure 1A, B**).

MC4R in Kiss1 neurons is required for female reproduction.

Because Kiss1-MC4RKO females showed altered puberty onset, we further investigated their

reproductive phenotype. Interestingly, Kiss1-MC4RKO female mice displayed normal BW

throughout the time of the study (up to PND150) **(Figure 2A)**; however, they presented with

irregular estrous cycles with predominantly more time spent in diestrus (P=0.0001) and less

time spent in estrus (P<0.0001) than their control littermates (**Figure 2*B, C***). The assessment of

the pulsatile release of LH every 10 min over 180 min revealed no change in the total number of

pulses between groups but a significant increase in basal LH release (P=0.0368) in Kiss1-

MC4RKO female mice (0.426±0.036) compared to controls (0.339±0.026) (**Figure 2*D-H***). The

analysis of the gene expression of the "KNDy" systems in the ARH, which control the GnRH

pulse generator [14], revealed normal expression levels of *Kiss1*, *Tac2* and *Tacr3*, but significantly

lower expression of *Pdyn* (P=0.0067, Kiss1-MC4RKO: 0.781±0.0384 vs controls: 1.000±0.0190)

(**Figure 2*I-L***). The conserved expression of *Kiss1*, *Tac2* and *Tacr3* correlates with the

preserved LH pulse frequency and amplitude in Kiss1-MC4RKO mice, while the lower inhibitory

tone of dynorphin (**Figure 2*I***) correlates with the higher basal LH levels (**Figure 2*E***). To assess

the contribution of MC4R signaling in Kiss1 neurons to the induction of ovulation through the

preovulatory LH surge, Kiss1-MC4RKO females and control littermates were submitted to an LH

surge induction protocol that showed the LH surge was significantly blunted in the Kiss1-

MC4RKO females (P=0.0091), while the protocol clearly evoked the expected afternoon rise of

LH in control mice (**Figure 2*M***). In line with these findings, the ovaries of Kiss1-MC4RKO

females displayed fewer corpora lutea, markers of recent ovulation (P=0.0054, Kiss1-MC4RKO:

0.800±0.374 vs controls: 2.80±0.374), in addition to increased cystic follicles (P=0.0337, Kiss1-

MC4RKO: 1.60±0.400 vs controls: 0.400±0.244) (**Figure 2*N-P***), which correlate with decreased

fertility as observed by the extended time to deliver pups (*i.e.* longer time to get pregnant)

(P=0.0030, Kiss1-MC4RKO: 30.30±4.600 vs controls: 20.88±0.2266), and fewer pups per litter

(P=0.0095, Kiss1-MC4RKO: 6.667±0.5528 vs controls: 8.444±0.242) (**Figure 2*S, T***).

The increase in serum LH levels and decreased ovulation observed in the Kiss1-MC4RKO

females is reminiscent of polycystic ovary syndrome (PCOS) mouse models [21,22]. Thus, we

investigated whether Kiss1-MC4RKO females display a PCOS-like phenotype. We analyzed

circulating levels of testosterone (T) and anti-müllerian hormone (AMH), which are frequently

elevated in PCOS models [23]. The Kiss1-MC4RKO females expressed normal T and AMH levels

compared to control mice in diestrus (**Figure 2*Q, R***). Thus, we can exclude a PCOS-like

reproductive phenotype mediated by the lack of melanocortin signaling on Kiss1 neurons.

Re-insertion of MC4R in Kiss1 neurons of MC4RKO mice improves reproductive function.

Kiss1-MC4RKO females displayed reproductive abnormalities resembling those described in

MC4RKO females [10-12]. Thus, we hypothesized that the reproductive defects described for the

MC4RKO mice would be, at least in part, due to the absence of MC4R signaling in Kiss1

neurons. To further investigate this hypothesis, we generated mice that do not express *Mc4r*

anywhere (MC4R-LoxTB, *i.e.* MC4RKO) or that express *Mc4r* only in Kiss1 neurons (*Kiss1-*

cre:MC4R-loxTB). The specific re-insertion of *Mc4r* within Kiss1 neurons in the *Kiss1-*

cre:MC4R-loxTB mice was confirmed through RNAscope (**Figure S2*A*).** *Mc4r* expression was

not detected in the PVN of these mice (**Figure S2*B***), and it was only detected in Kiss1 neurons

in the *Kiss1-cre:MC4R-loxTB* mice. Puberty onset was assessed daily from weaning age

through the monitoring of vaginal opening (VO) and first estrus (FE). MC4R-LoxTB and *Kiss1-*

cre:MC4R-loxTB female mice displayed normal timing of puberty onset compared to their

control littermates, as assessed by VO (P=0.104, MC4R-LoxTB: 30.86±1.908 vs *Kiss1-*

cre:MC4R-loxTB: 30.09±1.004 vs controls: 27.57±0.8168), and FE (P=0.8472, MC4R-LoxTB:

35.14±2.463 vs *Kiss1-cre:MC4R-loxTB: 34.55*±1.648 vs controls: 35.80±0.8406) (**Figure 3*A***),

despite displaying significantly higher body weight at the time of puberty onset (P=0.0001,

MC4R-LoxTB: 15.49±0.5078 vs *Kiss1-cre:MC4R-loxTB:* 15.54±0.40 vs Controls: 13.29±0.30)

(**Figure 3*B***). As expected, MC4R-LoxTB females (MC4RKO) displayed increased body weight

(**Figure 3C**) and an array of reproductive impairments that included: irregular estrous cycles

with significantly more days in diestrus (P=0.0016) and fewer days in estrus (P=0.0317)

compared to controls (**Figure 3*D, E***), significantly higher serum LH levels characterized by

higher basal (P=0.0882, MC4R-LoxTB: 0.48±0.02 vs Controls: 0.37±0.06) and amplitude levels

per LH pulse (P=0.0102, MC4R-LoxTB: 0.73±0.009 vs Controls: 0.58±0.07) (**Figure 3*F-J***), and

fewer corporal lutea (P=0.0366, MC4R-LoxTB: 0.80±0.80 vs Controls: 4.20±1.15) (**Figure 3*K,***

L), recapitulating the same reproductive phenotype observed in Kiss1-MC4RKO mice despite

the differences in BW. Re-introduction of MC4R into Kiss1 neurons in *Kiss1-cre:MC4R-loxTB*

mice completely recovered estrous cyclicity, which was similar to controls (diestrus: P=0.1932,

proestrus: P=0.8262, estrus: P=0.0547, compared to controls) (**Figure 3*D, E*)** despite *Kiss1-*

cre:MC4R-loxTB mice showing the same degree of obesity as MC4R-LoxTB mice (**Figure 3*C***),

indicating that obesity *per se* was not mediating the irregular estrous cycles in MC4RKO mice.

As indicated above, MC4R-LoxTB mice display higher overall circulating LH levels, and this

feature was also recovered by the re-insertion of MC4R in Kiss1 neurons of *Kiss1-cre:MC4R-*

loxTB mice (basal LH: 0.32±0.044, LH amplitude: 0.48±0.04, LH pulses/180min: 2.75±0.67 and

AUC: 70.62±7.156 compared to controls: 80.36±9.058 compared to controls) (**Figure 3*G-J***),

further confirming that the melanocortin action on Kiss1 neurons is required for the normal

control of gonadotropin release. Despite these significant improvements in reproductive

function, *Kiss1-cre:MC4R-loxTB* mice presented fewer corpora lutea than controls (0.80±0.80),

similar to MC4R-LoxTB mice, suggesting that an ovulatory impairment still persists (**Fig. 3*K, L***).

While tonic LH release and estrous cyclicity are predominantly controlled by Kiss1^{ARH} neurons,

ovulation is mediated by the induction of the preovulatory LH surge by Kiss1$^{AVPV/PeN}$ neurons. It

is possible that the metabolic signals derived from their obese phenotype, and/or the absence of

the direct action of MC4R in GnRH neurons [15,24] prevents the complete recovery of ovulation in

Kiss1-cre:MC4R-loxTB mice. Both genetic models displayed significantly lower T (MC4R-

LoxTB: 28.50±6.20 vs *Kiss1-cre:MC4R-loxTB:* 31.22±3.90 vs Controls: 62.84±16.53) and AMH

(MC4R-LoxTB: 150.6±22.00 vs *Kiss1-cre:MC4R-loxTB:* 146.9±10.83 vs Controls: 321.8±15.70)

levels than control mice in diestrus (**Fig. 3*M, N***). Therefore, we can exclude, once again, a

PCOS-like reproductive phenotype mediated by the lack of melanocortin signaling.

Kiss1^{ARH} neurons are excited by MC4R agonists.

Based on the expression of *Mc4r* in Kiss1 neurons and the reproductive impairment found after

Mc4r deletion within Kiss1 neurons (**Figure 2**), we hypothesized that Kiss1^{ARH} neurons are

direct targets of melanocortins and would respond to MC4R activation. Initially we did whole-

cell, voltage clamp recordings in Kiss1^{ARH} neurons (**Figure 4A**). *Kiss1-Cre* x Ai32 or *Kiss1-Cre*

AAV-injected mice underwent ovariectomies (OVX) and were given estradiol (E2) replacement

(*see Methods*). We targeted fluorescent cells for recording which were isolated synaptically by

bathing the slices in tetrodotoxin (TTX, 1 µM). Focal application of the high-affinity melanocortin

receptor agonist melanotan II (MTII, ~250 nM) elicited a small inward current in half of the

isolated Kiss1^{ARH} cells (10/21) (**Figure 4B**). Next, we examined if the E2 state affected MTII

response by recording from Kiss1^{ARH} neurons from OVX females and found two thirds (6/9 cells)

responded, but there was no significant difference in the average current (**Figure 4C**). Finally, it

must be noted that at this concentration MTII is a non-selective MCR agonist, so we followed up

using a perfusion of the MC4R-selective agonist THIQ (100 nM) to determine if activation of

MC4Rs alone was sufficient to invoke a postsynaptic response in Kiss1^{ARH} neurons from

OVX+E2 females. Similar to MTII, THIQ was able to elicit an excitatory inward current in

Kiss1^{ARH} neurons (4/5 cells, -7.4±0.5 pA), indicating MC4R activation is sufficient (**Figure 4D**).

POMC neurons synapse directly with Kiss1^{ARH} neurons.

Based on the pharmacological activation of MCRs, we wanted to address whether POMC^{ARH}

neurons are the source of melanocortins to excite Kiss1^{ARH} neurons directly. To answer this

question, we injected an AAV-EF1α-DIO-ChR2:mCherry vector into the ARH of adult *Pomc-Cre*

female mice (**Figure 4E**) (24). After 2-4 weeks we did whole cell recordings from putative

Kiss1^{ARH} neurons and looked for a response to high frequency optogenetic stimulation of POMC

fibers in OVX + E2-treated females (**Figure 4F**). We were able to confirm that these were Kiss1

neurons based on the presence of a persistent sodium current and/or *post hoc* identification by

scRT-PCR (41/69 cells, **Figure 4*G***). While this current is more prevalent in the AVPV Kiss1

population, Kiss1 neurons are the only ARH neurons to display this electrophysiological

"fingerprint" of a pronounced INaP paired with a high capacitance and a low input resistance [25].

Additionally, while recording from putative Kiss1[ARH] neurons, we optogenetically stimulated

POMC fibers at low frequency (1 Hz, 5 ms, 470 nm light) and recorded fast postsynaptic inward

currents. The fast kinetics of the EPSC are also a tell-tale sign of an ionotropic glutamatergic

response [26]. Previously, CNQX was sufficient to block similar excitatory ESPCs in other

postsynaptic targets of POMC neurons [27]. While the consistent latency from optogenetic

stimulus and current response was indicative of a single synaptic delay, we wanted to establish

that it was a direct synaptic connection, as we have previously shown for output of POMC

neurons [27] and Kiss1 neurons [28-30]. First, we abrogated the optogenetic response with the

addition of TTX (1 µM) to the bath, we were able to rescue the postsynaptic glutamate response

with the addition of the K^+ channel blockers 4-aminopyridine (4-AP; 0.5 mM) and tetraethyl

ammonium (TEA; 7.5 mM) (**Figure 4*H***). K+ channel block enables the calcium influx from

ChR2 activation in the presynaptic terminals to be sufficient to restore vesicle fusion. Therefore,

there appears to be no intervening synapses between POMC and the downstream Kiss1[ARH]

neurons.

Previously we found that E2 increases glutamate release from POMC neurons by increasing

Vglut2 [27]. Therefore, we compared the postsynaptic glutamatergic responses following two

optogenetic stimuli (50 ms interval between light flashes) in Kiss1[ARH] neurons from OVX +

vehicle versus OVX + E2-treated female mice (**Figure 4*I***). As anticipated, we found that

treatment with E2 increased the probability of glutamate release from POMC neurons onto

Kiss1[ARH] neurons based on the significant decrease in the paired pulse ratio of the two stimuli [31]

(**Figure 4*J***). Although in the present study we used an *in vivo* treatment paradigm, we know

from previous studies that this augmentation of glutamate release can happen quite rapidly after

a brief exposure to E2 *in vitro* (within 15 min) [27]. Therefore, Kiss1[ARH] neurons are excited by the

glutamatergic input from POMC neurons in an E2-dependent manner.

Finally, given that POMC neurons project directly onto Kiss1[ARH] neurons, we wanted to show

that the evoked slow inward current with high frequency stimulation was mediated by αMSH.

Indeed, high frequency stimulation evoked an excitatory inward current (**Figure 4K**) that was

blocked by the selective MC3/4R antagonist SHU 9119 (**Figure 4L**). The number of high

frequency (peptidergic) responses seen in Kiss1 neurons that displayed low frequency

postsynaptic (glutamatergic) currents was low (4/16, 25%, mean inward current: -1.4±3.1 pA).

However, since these recordings were made in brain slices from E2-treated OVX females and

β-endorphin (another product of POMC neurons) expression is enhanced by E2 treatment [32], we

were concerned that co-release of this opioid peptide could obscure melanocortin postsynaptic

effects. Indeed, there was a notable increase in slow inward currents generated by high

frequency stimulation of POMC fibers in slices that were perfused with the non-selective opioid

antagonist naloxone (1 µM) (9/23, 40%, mean inward current: -7.6±2.9 pA) (**Figure 4M**).

Together these optogenetic findings reinforced our pharmacological results showing that

Kiss1[ARH] neurons are excited by MTII, and THIQ. Additional studies would need to be done to

identify the cation conductance that is affected by the MC4R signaling cascade in Kiss1[ARH]

neurons, but clearly we have established that there are excitatory glutamatergic and peptidergic

inputs from POMC to Kiss1[ARH] neurons.

Kiss1[AVPV/PeN] neurons are inhibited by MC4R agonists.

Having established a direct, excitatory projection from POMC to Kiss1[ARH] neurons, we next

examined POMC inputs to Kiss1[AVPVPeN] neurons. First, we used immunocytochemistry to label

fibers expressing αMSH to demonstrate that POMC fibers densely innervated the AVPV/PeN

area (**Figure 5A**). Next, we recorded from AVPV neurons in slices taken from OVX+E2, POMC-

Cre mice injected with AAV1-DIO-YFP:ChR2 into the ARH (**Figure 5B**). We targeted cells along

305 the ventricle in the AVPV and PeN, surrounded by YFP terminals, and Kiss1$^{AVPV/PeN}$ neurons

306 were identified based on the expression of several endogenous conductances (I_{NaP}, T-type

307 calcium current, h-current) that are unique to these neurons [25,33]. We recorded a direct

308 glutamatergic synaptic response following optogenetic stimulation in 5 neurons, which was

309 further verified through pharmacological "rescue" of the synaptic response in 3 out of the 4

310 neurons tested (**Figure 5C**), indicating that POMC neurons make direct monosynaptic

311 connections with Kiss1$^{AVPV/PeN}$ neurons. We hypothesized that Kiss1$^{AVPV/PeN}$ neurons would be

312 excited by exogenous application of melanocortins because the receptor is typically Gs-coupled

313 [34,35]. We did whole-cell, voltage clamp recordings from Kiss1$^{AVPV/PeN}$-Cre::Ai32 neurons, which

314 were isolated synaptically by bathing the neurons in TTX (1 µM), from OVX females. For these

315 experiments, we tested the response to the high affinity melanocortin receptor agonist MTII (500

316 nM). Surprisingly, we consistently measured an outward (inhibitory) current that could be

317 reversed on washout of MTII (**Figure 5D**). This outward current was associated with an increase

318 in a K$^+$ conductance based on the current-voltage plot (*i.e.,* the outward current reversed at

319 ~E_{K+}, **Figure 5E**). Also, the inwardly-rectifying I/V could indicate that MC4R is coupled to

320 activation of inwardly-rectifying K$^+$ channels as has been previously reported for MC4R signaling

321 in pre-autonomic parasympathetic neurons in the brainstem [36]. We hypothesized that the

322 coupling could be modulated by E2 via a $G\alpha_q$-coupled membrane estrogen receptor (Gq-mER)

323 as we have previously demonstrated in POMC neurons [37-39]. For these experiments, we again

324 utilized OVX females and targeted Kiss1$^{AVPV/PeN}$-Cre::Ai32 neurons. Once in the whole-cell

325 voltage-clamp configuration, we perfused the slices with STX (10 nM), a selective ligand for the

326 putative Gq-mER [39], and tested the response to the MTII. Indeed, the outward current

327 (inhibitory) response to MTII was completely abrogated and even reversed by STX (**Figure 5F,**

328 **G**). The short-term (bath) treatment with STX ensured that there was no desensitization of the

329 Gq-mER with longer-term (*in vivo*) treatment with E2. Therefore, estrogen receptor activation

330 can rapidly uncouple (*i.e.,* desensitize) the MC4R inhibitory response in Kiss1$^{AVPV/PeN}$ neurons.

We would predict that the intracellular signaling cascade for the heterologous desensitization is

similar to what we have elucidated in POMC neurons [39], but this will need to be determined in

future experiments.   We next investigated the response in vehicle-treated, OVX females.

Indeed, Kiss1[AVPV/PeN] neurons were even more inhibited by the same MTII exposure (**Figure**

5G). Therefore, in contrast to Kiss1[ARH] neurons, the MC4R appears to be coupled to opening of

K^+ channels in Kiss1[AVPV/PeN] neurons. To further establish whether the MTII was having pre- or

postsynaptic effects, we measured the glutamatergic mEPSCs before and after melanocortin

receptor activation. We found there was no significant effect on the frequency (**Figure 5H**), but

there was an effect on the amplitude (**Figure 5I**). This further supports a postsynaptic locus of

STX's effects. However, *posthoc* comparisons found STX increased amplitude compared to

both OVX and OVX+E. This result hints at a membrane delimited effect as STX is a much more

potent agonist for Gq-mER than E2 [40].

Discussion

Our findings show that melanocortins act directly on Kiss1 neurons to contribute to the

metabolic regulation of fertility, in line with previous reports showing that αMSH stimulates LH

release in a kisspeptin-dependent manner [13] and that Kiss1 neurons express the melanocortin

receptor MC4R [17-20]. Several studies in humans have linked *MC4R* mutations to reproductive

abnormalities, including precocious puberty [8], PCOS [9], and hypogonadism [7]. However, a direct

association between *MC4R* mutations and reproductive function, independent from the obese

condition of these patients, has not been identified [6]. In the current study, we show that the

reproductive impairments observed in MC4R deficient mice, which replicate many of the

conditions described in humans, are largely mediated by the direct action of melanocortins via

MC4R on Kiss1 neurons and not to their obese phenotype. This is because the ablation of

MC4R from Kiss1 neurons largely replicated the reproductive impairments observed in

MC4RKO female mice without inducing obesity, and the selective reinsertion of MC4R into

Kiss1 neurons of MC4RKO mice significantly improved their reproductive function without

changing their obese phenotype. Strikingly, puberty onset was advanced in Kiss1-MC4RKO

females. Our findings revealed a low melanocortin tone in the hypothalamus of WT females

during pubertal development, characterized by lower levels of *Agrp, Mc4r* and *Mc3r* expression

at the time of puberty onset (~PND30). This developmental decline in the expression of

melanocortin receptors is in line with the advanced timing of puberty onset in females when

MC4R is congenitally ablated from Kiss1 neurons. A recent study using whole body MC3RKO

mice showed a trend to delayed puberty onset [18], suggesting a larger role of melanocortins in

the timing of puberty onset. Here, we show that in females the combination of MC3R and MC4R

action may be necessary for pubertal development. Interestingly, the MC4R-LoxTB and *Kiss1-*

cre:MC4R-LoxTB mice do not show an advancement in puberty onset as we observed in Kiss1-

MC4RKO mice; however, these mice are obese at the time of puberty onset already, which may

affect the timing of sexual maturation independently of MC4R action in Kiss1 neurons. Of note,

the findings presented in this study do not rule out the existence of additional sites of action of

MC4R in other neuronal populations to regulate reproduction, *e.g.,* GnRH neurons, which also

express MC4R [15,24]. Indeed, the lack of full recovery of the reproductive function in *Kiss1-*

cre:MC4R-LoxTB mice suggests that MC4R in Kiss1 neurons is necessary but not sufficient to

achieve full reproductive capabilities.

In addition to early puberty onset, Kiss1-MC4RKO females displayed an impaired preovulatory

LH surge driving ovulation and had fewer corpora lutea in their ovaries compared to control

littermates, further supporting a role of MC4R in regulating ovulation. In adulthood, the

reproductive phenotype observed in Kiss1-MC4RKO and MC4R-loxTB female mice (increased

LH, irregular estrous cycles, oligo-ovulation, increased cystic follicles) correlates with the

phenotype observed in PCOS mouse models [21,22]. In fact, an association between *MC4R*

mutations and PCOS has been reported [9]. However, our mouse models failed to display higher

levels of circulating androgens or AMH, two of the hallmarks of PCOS [23], suggesting that

melanocortin signaling is unlikely to be a main contributing factor in the development of this

syndrome.

Our data revealing early puberty onset and impaired LH pulse and surge driving ovulation

support a direct role of MC4R on the two primary Kiss1 populations regulating these functions

(ARH and AVPV/PeN). We further investigated this and found that MC4R signaling excites

Kiss1^{ARH} neurons and inhibits Kiss1$^{AVPV/PeN}$ neurons in an estradiol-dependent manner. In

Kiss1^{ARH} neurons, bath applied or optogenetically-evoked release of melanocortin agonists

induce a direct excitatory inward current. Our studies using whole cell voltage clamp in female

mice were able to detect a melanocortin-mediated inward current in Kiss1^{ARH} neurons that

would increase excitability. Previously, an MCR-activation effect was not detected in cell-

attached recordings of diestrus females [13]. However, cell-attached recordings do not measure a

direct response in isolated Kiss1^{ARH} neurons but rather the summation of multiple synaptic

inputs to Kiss1^{ARH} neurons. Most importantly, we did not see a difference in the excitatory

response in synaptically isolated Kiss1^{ARH} neurons from E2-treated versus vehicle-treated,

ovariectomized females. This emphasizes the importance of doing whole-cell recording from

isolated Kiss1 neurons. The modulatory actions of POMC^{ARH} neuronal projections to Kiss1^{ARH}

neurons can also occur through a) the excitatory action of glutamate in a process that is

facilitated in the presence of estradiol, in line with previous publications [41,42], and b) the

inhibitory action of β-endorphin, suggesting that POMC^{ARH} neurons are able to exert a precise

regulatory role of the GnRH pulse generator, *i.e.* Kiss1^{ARH} neurons, in response to metabolic

challenges. In the AVPV/PeN, MC4R agonists inhibited Kiss1$^{AVPV/PeN}$ neurons in an estradiol-

dependent manner, which utilized a Gα_q-coupled membrane estrogen receptor (STX receptor)

to attenuate the inhibitory tone of MC4R in these neurons.

Our pharmacology data clearly demonstrates the estradiol-dependent action of MC4R in

Kiss1$^{AVPV/PeN}$ neurons, suggesting that similar to Kiss1^{ARH} neurons, this population is tightly

controlled by melanocortins. The effect of MC4R agonists on Kiss1 neurons in the presence of

TTX, suggests a direct synaptic effect without the need of channelrhodopsin-2(ChR2)-assisted

circuit mapping. Nonetheless, to further demonstrate this interaction, we provide evidence of

profound innervation of αMSH fibers and the presence of synaptic contact between POMC and

Kiss1[AVPV/PeN] neurons through the optogenetic stimulation of a glutamatergic response in

Kiss1[AVPV/PeN] neurons after low frequency stimulation of POMC terminals. We intentionally

avoided high frequency stimulation of POMC terminals to prevent the co-release of β-

endorphins (e.g., **Figure 4M**) and their possible additive effect on the inhibitory action of MC4R.

Future studies will be aimed at the characterization of the endogenous opioid pathways in the

induction of the LH surge.

To date, two instances of MC4R-mediated inhibition have been described in the central nervous

system: MC4R can Gi,o couple to open K+ (Kir6.2) channels in parasympathetic preganglionic

neurons in the brainstem [36], and MC4R can directly couple to K+ (Kv7.1) channels in the

hypothalamic paraventricular nucleus neurons [43]. Interestingly, our findings describe an

inhibitory role of MC4R in the reproductive neuroendocrine axis. This steroid state dependent

effect is particularly relevant because in the context of reproduction, the two populations of

Kiss1 neurons are strikingly different in their response to estradiol, where peptide expression in

Kiss1[ARH] and Kiss1[AVPV/PeN] neurons are inhibited or stimulated, respectively, in order to mount

the negative versus positive feedback of sex steroids [14]. This differential regulation allows for

the episodic versus surge release of GnRH; however, the cellular mechanisms underlying these

opposing roles of Kiss1 neurons to the same stimulus, *i.e.* circulating E2 levels, remain

unknown. Although there were no differences in the magnitude of the electrophysiological

responses to MTII in Kiss1[ARH] neurons in vehicle- versus E2-treated, ovariectomized females,

we cannot rule out that E2 affects other systems activated by the MC4R signaling cascade as

recently reported in MC4R expressing neurons of the ventromedial nucleus of the hypothalamus

(VMH) [44]. Interestingly, puberty onset was advanced in Kiss1-MC4RKO females. This data

suggests that the inhibitory tone of MC4R signaling on Kiss1[AVPV/PeN] neurons might also be

involved in the timing of puberty onset, which in turn suggests a role for this female-specific population of Kiss1 neurons in sexual maturation. This phenomenon could explain the earlier age of puberty onset frequently seen in females of most mammalian species compared to their male counterparts [45]. Furthermore, our findings suggest that metabolic cues, through the regulation of the melanocortin output onto Kiss1$^{AVPV/PeN}$ neurons, are essential for the timing and magnitude of the GnRH/LH surge. Future studies are warranted to fully elucidate synaptic input and postsynaptic cascades in the POMC-Kiss1$^{AVPV/PeN}$ circuit.

Altogether, our data reveal a differential regulatory action of MC4R in the neural control of GnRH/LH release, participating in both the surge (E2-treated, ovariectomized female) and the pulse-like (ovariectomized female) modes of LH release through the cellular regulation of Kiss1 neurons, which translate into our findings on the conditional knockout of *Mc4r* from Kiss1 neurons (**Figure 6**). These findings are important because the reproductive abnormalities often attributed to obesity in MC4R deficient patients may not be caused by the excess in body weight but, at least in part, by a deficiency in MC4R signaling directly at the level of Kiss1 neurons that affects predominantly the ability to mount a preovulatory LH surge.

Materials and methods:

In-vivo experimental procedures:

Generation of Kiss1-MC4RKO, Kiss1-cre:MC4R-LoxTB and MC4R-LoxTB Transgenic Mice

Kiss1-MC4RKO mice were generated by crossing *Kiss1-cre* knock-in mice and *MC4R$^{lox/lox}$* mice. *Kiss1-cre* mice (RRID:MGI:6278139) were obtained from Dr. Richard Palmiter (University of Washington, Seattle, WA) [46] and *Mc4r$^{lox/lox}$* (RRID:IMSR_JAX:023720) were a gift from Dr. Brad Lowell (Beth Israel Deaconess Hospital, Boston, MA) [4]. These mice were crossed to generate Kiss1-MC4RKO mice lacking *Mc4r* expression selectively from Kiss1 neurons (*Kiss1-cre$^{+/-}$*; *MC4R$^{lox/lox}$*) and their control littermates (MC4R$^{lox/lox}$). To generate *Kiss1-cre:MC4R-LoxTB* and

MC4R-LoxTB mice, *MC4R-LoxTB* mice were purchased from The Jackson Laboratory (MC4R-

LoxTB; catalog no. 006414). These mice present with a loxp-flanked transcriptional blocking

(LoxTB) sequence preventing normal endogenous gene transcription and translation from the

endogenous locus. Homozygous MC4R-loxTB/loxTB mice are devoid of functional MC4R

mRNA (MC4RKO mice), while the presence of Cre recombinase on the Kiss1 promotor in

Kiss1-cre mice will result in the removal of the transcription blocker and subsequent expression

of MC4R in tissue-specific sites (*i.e.* Kiss1 neurons) therefore resulting in the generation of

Kiss1-cre:MC4R-LoxTB (with *Mc4r* expression restored in Kiss1 neurons), their obese control

littermates *MC4R-LoxTB* mice (MC4RKO), and their WT controls (MC4R$^{+/+}$). To rule out early

embryonic recombination of the MC4R$^{loxTB/loxTB}$ or MC4R$^{lox/lox}$ alleles, we ran PCR assays on tail

DNA designed to detect wild-type allele (MC4R$^{+/+}$), undeleted lox (MC4R$^{lox/lox}$) or loxTB alleles

(MC4R$^{loxTB/loxTB}$). Genotyping was confirmed by sending tissue to Transnetyx, Inc., for testing by

real-time polymerase chain reaction. Mice were housed in Harvard Medical School Animal

Resources facilities where they were fed standard mouse chow (Teklad F6 Rodent Diet 8664)

and were given *ad libitum* access to tap water under constant conditions of temperature (22–

24˚C) and light (12 hr light [07:00]/dark [19:00] cycle).

RNAscope in situ hybridization

To validate the Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and MC4R-LoxTB (MC4RKO) mouse

models and investigate the co-expression of *Kiss1* and *Mc4r* mRNA in these mice, dual

fluorescence ISH was performed using RNAscope (ACD, Multiplex Fluorescent v.2) according

to the manufacturer's protocol using the following probes: *Mc4r* (319181-C2) and *Kiss1*

(500141-C1). Brains (n=4/group) from WT OVX (for expression in the ARH) and OVX+E2 (for

expression in the AVPV/PeN) mice and OVX Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and

MC4RKO were removed fresh frozen on dry ice, and then stored at -80˚C until sectioned. Five

sets of 20 μm sections in the coronal plane were cut on a cryostat, from the diagonal band of

Broca to the mammillary bodies, thaw mounted onto SuperFrost Plus slides (VWR Scientific)

and stored at -80°C until use. A single set was used for ISH experiment (adjacent sections 100

mm apart). Images were taken at 20x magnification of the sections containing AVPV, PeN, and

the three rostro-to-caudal levels of the ARH, and Kiss1 neurons expressing (*Kiss1-cre*:MC4R-

LoxTB mice) or lacking (Kiss1-MC4RKO mice) *Mc4r* were identified using ImageJ.

Reproductive maturation of Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and MC4R-LoxTB

mice

To assess the reproductive phenotype of mice with selective reinsertion of *Mc4r* on Kiss1

neurons (*Kiss1-cre*:MC4R-LoxTB, n=11), selective deletion of *Mc4r* from Kiss1 neurons (Kiss1-

MC4RKO, n=24), global deletion of *Mc4r* (MC4R-LoxTB, n=7); and their control MC4R$^{lox/lox}$

littermates (n=14), mice were weaned at post-natal day (PND) 21 and were monitored daily for

puberty onset. Females were monitored daily for vaginal opening (VO, indicative of the

complete canalization of the vaginal cavity) and for first estrus (first day with cornified cells

determined by daily morning vaginal cytology) after the day of VO. Body weight (BW) was

measured at the day of puberty onset to determine if changes in puberty onset could be due to

differences in BW. Estrous cyclicity was monitored in females by daily vaginal cytology, for a

period of 15 days, in 6-month-old mice and their respective control littermates (n=5/group).

Cytology samples were obtained every morning (9 am), placed on a glass slide and stained with

hematoxylin and eosin for determination of the estrous cycle stage under the microscope.

Fecundity test of Kiss1-MC4RKO females

Adult 6-month-old Kiss1-MC4RKO and control littermate female mice (n=9/group) were placed

with adult WT males proven to father litters for 90 days and time to deliver pups and number of

pups per litter were monitored.

Characterization of the estradiol-induced luteinizing hormone surge

Kiss1-MC4RKO (n=6) and control MC4R[lox/lox] littermate (n=5) adult female mice were subjected

to bilateral ovariectomy (OVX) via abdominal incision under light isoflurane anesthesia.

Immediately after OVX, capsules filled with E2 (1 ug/20g BW) were implanted subcutaneously

via a small mid- scapular incision on the back. Five days later, mice were subcutaneously

injected in the morning with estradiol benzoate (1 ug/20g BW) to produce elevated proestrus-

like E2 levels (preovulatory LH surge) on the following day [47]. Blood samples were collected at

8am and 7pm; LH levels were stored at -80˚C until measured via LH ELISA.

Ovarian histology and hormone measurements

Bilateral ovariectomy from 6-month-old Kiss1-MC4RKO (n=4), *Kiss1-cre*:MC4R-LoxTB, MC4R-

LoxTB and their control littermates (n=5/group) was performed under light isoflurane

anesthesia. Briefly, the ventral skin was shaved and cleaned, and one small abdominal incision

was made. Once the ovaries were identified and excised, the muscle incision was sutured, and

the skin was closed with surgical clips. Ovaries were stored in Bouin's fixative, sectioned, and

stained with hematoxylin and eosin at the Harvard Histopathology Core. Corporal lutea were

counted in the middle section of each ovary. Serum samples were also collected for analysis of

testosterone and anti-müllerian hormone (AMH) levels in these mice. These hormones levels

were measured at the University of Virginia Ligand Assay core with the Mouse & Rat

Testosterone ELISA assay (reportable average range 10-1600 ng/dL; sensitivity of 10 ng/dL);

AMH ELISA assay (reportable average range 0.2 – 15 ng/mL; sensitivity of 0.2 ng/mL).

LH pulsatile secretion profile in gonad intact Kiss1-MC4RKO, *Kiss1-cre*:MC4R-LoxTB and

MC4R-LoxTB female mice

To assess the profile of LH pulses secretion, adult 6 months old gonad intact Kiss1-MC4RKO

females (n=6), their control Mc4r$^{lox/lox}$ littermates (n=7); and *Kiss1-cre*:MC4R-LoxTB (n=5/group),

MC4R-LoxTB and their control WT females (n=4/group) were handled daily for 3 weeks to allow

acclimation to sampling conditions prior to the experiment. Pulsatile measurements of LH

secretion were assessed by repeated blood collection through a single incision at the tip of the

tail. The tail was cleaned with saline and 4 µL of blood was taken at each time point from the cut

tail with a pipette. We collected sequential blood samples every 10 minutes over a 180-minute

sampling period. Samples were immediately frozen on dry ice and stored at –80 °C until

analyzed with LH ELISA as previously described [48]. The functional sensitivity of the ELISA

assay was 0.0039 ng/ml with a CV% of 3.3%.

LH pulses analysis

LH pulses in mice were analyzed using a custom-made MATLAB-based algorithm. The

MATLAB code includes a loop that determines LH pulses as any LH peak: (i) whose height is

20% greater than the heights of the 2 previous values; (ii) 10% greater than the height of the

following value; and (iii) the peak at the second time interval needs to be 20% greater than the

single value that comes before it to be considered a pulse, as we previously described [49].

LH pulsatility was assessed by measuring: (1) the total secretory mass, assessed by area under

the curve (AUC); (2) the LH pulse amplitude, calculated by averaging the 4 highest LH values in

the samples collection period for each animal; (3) the basal LH, calculated by averaging the 4

lowest LH values in the samples collection period for each animal; and (4) the total number of

pulses throughout the 180 minutes sampling period.

Immunohistochemistry

Animals and Treatment: Coronal brain blocks (2 mm each) from adult female C57BL/6 mice

(n=4) were fixed by immersion in 4% paraformaldehyde for ~ 8 hours, cryoprotected in 30%

sucrose solution, frozen in isopentane at –55°C, sectioned coronally on a cryostat at 20 μm, and

thaw-mounted onto Superfrost Plus slides (Fisher Scientific, Pittsburgh, PA).   The 20 μm

sections were stored at –20°C until used for immunocytochemistry.

Immunocytochemistry: The _s_ections were rinsed in PB (0.1M phosphate buffer, pH 7.4) for at

least 30 min. Next, sections were incubated with normal serum corresponding to the host for

the secondary antiserum (5 % normal serum with 0.3% Triton-X100 in PBS for 30 min), rinsed in

PB and then incubated for ~45 h at 4º C with a rabbit polyclonal antiserum against α-MSH

(1:2,500). The specificity of this antiserum has been documented [50]. After rinsing, sections were

first incubated for 2-3 hours at room temperature with biotinylated donkey anti rabbit gamma

globulin (IgG; 1:500) and next with streptavidin-Alexa 488 (1:2500). Both the primary and

secondary antibodies were diluted in tris-(hydroxymethyl) aminomethane (0.5 %, Jackson

ImmunoResearch, Philadelphia, PA) in PB containing 0.7 % seaweed gelatin (Jackson

ImmunoResearch, Philadelphia, PA) and 0.5 % Triton X-100 and 3% bovine serum albumin

(BSA; Jackson ImmunoResearch, Philadelphia, PA) adjusted to pH 7.6. Following a final rinse

overnight, slides were coverslipped with gelvatol containing the anti-fading agent, 1,4-

diazabicyclo(2,2)octane (DABCO; Cold Spring Harbor Protocols, 2006).

Imaging: Photomicrographs of labeling were initially acquired using a Nikon E800 fluorescent

microscope (Eclipse E800; Nikon Instruments, Melville, NY) equipped with a fiber illuminator

(Intensilight C-HGFI; Nikon Instruments) and a high-definition digital microscope camera head

(DS-Fi1; Nikon Instruments) interfaced with a PC-based camera controller (DS-U3; Nikon

Instruments).

Real time quantitative PCR

i) To investigate the changes in the expression of the melanocortin genes _Agrp_, _Pomc, Mc3r_

and _Mc4r_ in the MBH during development in prepubertal and pubertal WT females at ages P10

(n=6), P15 (n=6), P22 (n=6), and P30 (n=5), and ii) evaluate the gene expression profile of

Pdyn, *Kiss1*, *Tac2*, *Tacr3*, in the ARH of adult intact (in diestrus) female Kiss1-MC4RKO (n=6)

and their control MC4R^lox/lox littermates (n=3), the brains were removed and rapidly embedded in

Tissue-Tek, frozen in –30°C isopentane solution and stored at –80°C until use. Frozen tissue

punches were recovered through MBH with a 1 mm diameter canula [51]. These tissue punches

encompassed the whole MBH from the WT females and the ARH from the Kiss1-MC4RKO and

their control females. The tissues were homogenized, and total RNA was isolated using TRIzol

reagent (Invitrogen) followed by chloroform/isopropanol extraction. RNA purity and

concentration were measured via an absorbance spectrophotometer (260/280 nm > 1.8;

NanoDrop 1000, Thermo Fisher Scientific). Total RNA (1 μg) was reverse transcribed to cDNA

using random hexamers (High-Capacity cDNA Synthesis Kit, Life Technologies). Quantitative

real-time PCR assays were performed using SYBR Green RT-qPCR master mix (Applied

Biosystems) and analyzed using ABI Prism 7000 SDS software (Applied Biosystems). The

cycling conditions were: 2 minutes incubation at 95°C (hot start), 45 amplification cycles (95°C

for 30 seconds, 60°C for 30 seconds, and 45 seconds at 75°C, with fluorescence detection at

the end of each cycle), followed by melting curve of the amplified products obtained by ramped

increase of the temperature from 55°C to 95°C to confirm the presence of single amplification

product per reaction. PCR specificity was verified by melting curve analysis and agarose gel

electrophoresis. Each sample was run in duplicate to obtain an average cycle threshold (CT)

value and relative expression of each target gene was determined using the comparative Ct

method [52]. The data were normalized to Hypoxanthine Guanine Phosphoribosyltransferase

(*Hprt)* expression levels in each sample. Results were expressed as fold differences in relative

gene expression with respect to i) P10 for melanocortin genes expression analysis during

development in WT mice, and ii) controls for the KNDy genes expression in female Kiss1-

MC4RKO mice. The primers used are listed in Table 1.

Gene	Gene length (pb)	Accession #	Primers	Location (nt)	Sequence
Hprt	352	NM_013556.2	Hprt-F	704-728	CCTGCTGGATTACATTAAAGCGCTG
			Hprt-R	377-401	GTCAAGGGCATATCCAACAACAAAC
Agrp	136	NM_001271806 .1	Agrp-F	466-488	GCCTCAAGAAGACAACTGCAGAC
			Agrp-R	580-601	AAGCAGGACTCGTGCAGCCTTA
Pomc	138	NM_001278584 .1	Pomc-F	187-208	CCATAGATGTGTGGAGCTGGTG
			Pomc-R	303-324	CACCTCCGTTGCCAGGAAACAC
Mc4r	101	NM_016977.4	Mc4r-F	552-570	CCCGGACGGAGGATGCTAT
			Mc4r-R	632-652	TCGCCACGATCACTAGAATGT
Pdyn	200	NM_018863.4	Pdyn-F	45-64	ACAGGGGGAGACTCTCATCT
			Pdyn-R	223-244	GGGGATGAATGACCTGCTTACT
Kiss1	129	AF472576.1	Kiss1-F	147-166	GCTGCTGCTTCTCCTCTGTG
			Kiss1-R	256-275	TCTGCATACCGCGATTCCTT
Tac2	234	NM_001199971 .1	Tac2- F	238-257	GCTCCACAGCTTTGTCCTTC
			Tac2- R	452-471	GCTAGCCTTGCTCAGCACTT
Tacr3	159	NM_021382.6	Tacr3-F	759-779	GCCATTGCAGTGGACAGGTAT
			Tacr3-R	898-917	ACGGCCTGGCATGACTTTTA

Data analysis

Statistical data are expressed as means ± SEM, where n represents the number of animals in
each study group. The significance of differences between groups was evaluated using
unpaired two-tailed Student's t test, or a one- or two-way ANOVA test (with *post hoc*
comparisons). Significance level was set at $p<0.05$. All analyses were performed with GraphPad
Prism Software, Inc (San Diego, CA).

<u>In-vitro experimental procedures:</u>

Animals:

All animal procedures described in the electrophysiology studies were performed in accordance

with institutional guidelines based on National Institutes of Health standards and approved by

the Institutional Animal Care and Use Committees at Oregon Health and Science University and

Appalachian State University. *Pomc-Cre* mice (RRID:IMSR_JAX:005965) [53] were crossed with

wildtype C57B6J (RRID:IMSR_JAX:000664) mice. *Kiss1-cre*(v2) mice [46] were crossed with Ai32

[54] or C57B6J mice. All colonies were maintained onsite under controlled temperature (21-23 ºC)

and photoperiod (12:12-h light-dark cycle 0600 to 1800) while receiving ad libitum food (5L0D;

LabDiet, St. Louis, MO) and water access. Following surgeries, mice received a *s.c.* dose of 4-5

mg/kg carprofen (Rimadyl; Pfizer Animal Health, New York, NY) and given at least 1 week of

recovery.

Ovariectomies:

At least 7 days prior to each experiment, ovaries were removed as described previously while

under isoflurane anesthesia [42]. Two days before experiments females received either an

injection of sesame oil (50 µl, *sc*; Sigma-Aldrich, St. Louis, MO) or a priming dose (0.25 µg /50

µl sesame oil, *sc*) of E2 benzoate (Sigma-Aldrich, St. Louis, MO) in the morning. On the

following day, oil or a high (1.5 µg) dose of E2 benzoate, which generates a LH surge, was

administered [55]. Circulating levels of E2 were verified by the uterine weights (<25 mg for OVX

and >95 mg for OVX E2-treated) at the time of euthanasia [55].

AAV Delivery:

Bilateral ARH injections of AAV1-Ef1a-DIO-ChR2:mCherry (RRID:Addgene_20297) or AAV1-

Ef1a-DIO-ChR2:YFP (RRID:Addgene_20298) were performed on adult *Kiss1-cre* mice or

Pomc-cre mice on a stereotaxic frame under isoflurane anesthesia. ARH injection coordinates

were anteroposterior (AP): –1.10 mm, mediolateral (ML): ± 0.30 mm, dorsoventral (DL): –5.80

mm (surface of brain z = 0.0 mm); 400 nl of the AAV (2.0×10^{12} particles/ml) was injected (100

nl/min) into each position. Mice were given carprofen for analgesia and allowed to recover for at

least two weeks before euthanasia.

Visualized whole-cell patch recordings:

Hypothalamic coronal brain slices (240 um) were made from female mice, using a Leica

VT1000S vibratome, in ice cold cutting solution bubbled with O_2/CO_2 (95%/5%). Slices were

then transferred to a holding chamber with artificial cerebrospinal fluid bubbled with the same

gas mix and allowed to recover for at least 1 hour. For recordings, slices were placed in a

perfusion chamber and visualized with a Olympus BX51W1 using either differential infrared

contrast or oblique illumination. Kiss1 neurons in the ARH or AVPV/PeN were targeted for

electrophysiological recordings as done previously and described below [56,57].

Solutions/drugs:

Standard vibratome slicing, external, and internal recording solutions were utilized as previously

described [56,57]. Tetrodotoxin was purchased from Alomone Labs (Jerusalem, Israel), Melanotan

II and αMSH from Tocris (Minneapolis, MN). TEA, 4-AP, 17β-estradiol benzoate, and Naloxone

were purchased from Millipore-Sigma. STX was produced by AAPharmaSyn, LLC (Ann Arbor,

MI) under contract.

Electrophysiology data analysis:

Electrophysiological data were analyzed using Clampfit 10/11 (Molecular Devices) and Prism

7/10 (Dotmatics). All values are expressed as Mean ± SEM. Comparisons between two groups

were made using un-paired Student's t-test or between multiple groups using an ANOVA (with

post hoc comparisons) with p-values < 0.05 considered significant. When variances differed

significantly, Mann-Whitney U test was used instead.

Targeting of Kiss1 neurons for electrophysiological recordings.

For non-optogenetic experiments, brain slices were taken from AAV injected Kiss1-Cre AAV or

Kiss1xAi32 female mice. Ai32 mice (RRID:IMSR_JAX:024109, C57BL/6 background) carry the

floxed ChR2 (H134R)-EYFP gene in their Gt(ROSA)26Sor Locus [54], allowing its expression in a

Cre-dependent manner. Due to concerns of nonspecific expression [58], we previously validated

this model using single cell RT-PCR and documented that *Kiss1* mRNA was detectable in 99%

of individually harvested eYFP cells (n=126). In addition, we have used both AAV injected

Kiss1-Cre AAV or Kiss1xAi32 female mice and found no differences in electrophysiological

results [56]. We used AAV injected POMC-Cre female mice and avoided small soma, high input

resistance (>800 MΩ), ventrally located cells that are typically NPY/AgRP neurons. Instead, we

targeted larger, more dorsomedial neurons in the ARH while avoiding fluorescent (POMC) cells.

Unlike ARH Kiss1 neurons, POMC neurons do not make reciprocal projections, so uninfected

POMC neurons do not display monosynaptic EPSCs in response to optogenetic stimulation.

Next, we used a ramp IV protocol to probe for the presence of a persistent sodium current (I_{NaP}).

While this current is more prevalent in the AVPV Kiss1 population, Kiss1 neurons are the only

ARH neurons to display this electrophysiological "fingerprint" of a pronounced I_{NaP} paired with a

high capacitance and low input resistance [25]. Finally, we also harvested the cytosol at the end of

all recordings and measured the expression of Kiss1. Only cells that expressed a persistent

sodium current and/or expressed *Kiss1* mRNA were included in final analysis (n = 44).

Acknowledgments:

The authors would like to recognize the excellent technical expertise of Ms. Martha A. Bosch

(tissue preparation, immunohistochemical procedure and single cell RT-PCR of recorded

neurons). In addition, we thank Dr. Rona Carroll for her assistance at Harvard Medical School

and Mr. Cole Martinson, a student worker in the Ronnekleiv/ Kelly laboratories, for his

assistance with genotyping and care of mouse colonies at OHSU. The University of Virginia

Center for Research in Reproduction Ligand Assay and Analysis Core is supported by the

Eunice Kennedy Shriver NICHD/NIH (NCTRI) Grant P50-HD28934. This work was supported by

PHS MPI grant DK68098 to MJK and OKR; HD090151, HD099084 and DK133760 to VMN, and

The Charles A. King Trust Postdoctoral Research Fellowship Award, The Lalor Foundation

Postdoctoral Fellowship Award, the Women's brain initiative Fellowship Award, the ROSA

SCORE pilot grant (supported by NIH Research Grant U54 AG062322 funded by The National

Institute on Aging (NIA) and Office of Research on Women's Health (ORWH)) and the IBRO-

ISN Research Fellowship Award to RT, and startup funds from Appalachian State University to

TLS.

Authors contribution:

RT, TLS, MJK and VMN conceived and designed the study; RT, TLS, KF, CJH, EM, AG, KW,

SL, EAM, OKR performed the experiments. RT, TLS, OKR, MJK and VMN analyzed data. RT,

TLS and VMN wrote the manuscript with input from all the authors.

Declaration of interests:

The authors declare they have no competing interest.

Figure legends:

Figure 1. *Mc4r* expressed in Kiss1 neurons determines the timing of puberty onset.

Kiss1-MC4RKO females display advanced puberty onset, assessed by daily monitoring of vaginal opening **(A)** and first estrus **(B)**, as documented by cumulative percent and mean age of animals at vaginal opening **(A)** and first estrus **(B)** in Kiss1-MC4RKO females (n=24) compared to WT littermates (n=19). *p<0.05 by Student's t-test. Data presented as the mean± SEM. **(C)** Female Kiss1-MC4RKO (n=19) have normal body weight at the time of puberty onset compared to their WT littermates (n=15). **(D)** Ontogeny expression of melanocortin genes (*Agrp*, *Pomc*, *Mc4r and Mc3r*) in the ARH of WT female mice at different postnatal pre-pubertal and pubertal ages: P10, P15, P22 and P30, normalized to the housekeeping gene *Hprt*. (*n values*: females at P10 (n=6), P15 (n=6), P22 (n=6), and P30 (n=5)). Groups with different letters are significantly different (p<0.05), as determined by one-way ANOVA followed by the Student-Newman-Keuls test. Data presented as the mean± SEM.

Figure 2. Deletion of MC4R from Kiss1 neurons impairs fertility in Kiss1-MC4RKO females.

(A) Female Kiss1-MC4RKO (n=19) have normal body weight from weaning and until post-natal day (PND) 150 compared to their WT littermates (n=15). **(B)** Representative examples of estrous cycles in Kiss1-MC4RKO and control females (n=5/group) assessed by daily vaginal cytology for 15 days. **(C)** Kiss1-MC4RKO females displayed irregular estrous cycles with a longer time in diestrus and a shorter time in estrus compared to control females. *** P < 0.001 Two-way ANOVA. **(D)** Pattern of LH pulsatility was analyzed in gonad intact Kiss1-MC4RKO (n=6) and control (n=7) females. # Represents LH pulses. **(E)** Basal LH, **(F)** LH total secretory mass assessed by area under the curve (AUC), **(G)** LH pulse amplitude and **(H)** Total number of

pulses /180 min were analyzed. *p<0.05 by Student's t-test. The expression of the KNDy genes

Pdyn **(I)**, *Kiss1* **(J)**, *Tac2* **(K)** and *Tacr3* **(L)**, was assessed in the ARH of adult Kiss1-MC4RKO

(n=6) and control (n=3) females. **p<0.01 by Student's t-test. Data presented as the mean±

SEM. **(M)** Kiss1-MC4RKO (n=6) and control (n=5) females were subjected to an LH surge

induction protocol. LH samples were collected in the morning (AM [8 a.m.]) and evening (PM [7

p.m.]) after lights off. **P < 0.01, Two-way ANOVA followed by Bonferroni *post hoc* test. **(N)**

Ovarian histology shows a decrease in the number of corpora lutea (CL, **O**) and increase in the

number of cystic follicles (CF, **P**) of Kiss1-MC4RKO compared to controls(n=5/group). #

represents cystic follicles. Serum levels of Testosterone **(Q)** and AMH **(R)** in adult gonad intact

Kiss1-MC4RKO and control females (n=5/group). Student t test for unpaired samples. Data

presented as the mean± SEM. Kiss1-MC4RKO females display impaired fertility characterized

by increased time to deliver pups **(S)** and decreased number of pups per litter **(T)**, (n=9/group).

*p<0.05 by Student's t-test. Data presented as the mean± SEM.

Figure 3. Re-insertion of MC4R in Kiss1 neurons restores estrous cyclicity and LH

pulsatility in Kiss1-cre:MC4R-LoxTB females.

(A) *Kiss1-cre*:MC4R-LoxTB (n=11) and MC4R-loxTB (MC4RKO) (n=7) displayed normal

puberty onset as compared to their control littermates (n=14), as documented by cumulative

percent and mean age of animals at vaginal opening and first estrus. **(B)** *Kiss1-cre*:MC4R-

LoxTB and MC4R-loxTB both had significantly higher body weights at the time of puberty onset

compared to their controls. **P < 0.01, *** P < 0.001 by one way ANOVA. **(C)** MC4R-loxTB

(n=7) and *Kiss1-cre*:MC4R-LoxTB (n=11) females displayed significantly higher body weight

than their littermates (n=10) from post-natal day 30 onwards, *p<0.05 and **P < 0.01. Data

presented as the mean± (SEM). **(D)** Representative examples of estrous cycles of MC4R-loxTB

(n=4), *Kiss1-cre*:MC4R-LoxTB (n=5) and control littermate mice, (n=4) assessed by daily

monitoring of vaginal smears for 15 days. (D/M: diestrus/metestrus, P: proestrus, E: estrus). **(E)**

MC4R-loxTB females displayed irregular estrous cycles, presenting longer time in diestrus and

shorter time in estrus compared to control females, while *Kiss1-cre*:MC4R-LoxTB females

displayed regular estrous cyclicity, similar to controls. *p<0.05, **P < 0.01, *** P < 0.001, ****P <

0.0001, Two-way ANOVA. Data presented as the mean± SEM. **(F)** Pattern of LH pulsatility

analyzed in gonad intact MC4R-loxTB (n=4), *Kiss1-cre*:MC4R-LoxTB (n=5) and control

littermates females (n=5). LH samples were collected every 10 min for 180 minutes; #

represents LH pulses. **(G)** LH total secretory mass, **(H)** LH pulse amplitude, **(I)** basal LH, and **(J)**

total number of pulses /180 min were assessed. *p<0.05, **P < 0.01 by one way ANOVA. **(K)**

Representative samples of ovarian histology from MC4R-loxTB, *Kiss1-cre*:MC4R-LoxTB and

control females (n=5/group); CL: Corpora lutea. Data are presented as the mean± SEM. **(L)**

Ovarian histology showed a significant decrease in the number of corpora lutea in the MC4RKO

and *Kiss1-cre*:MC4R-LoxTB compared to controls. Groups with different letters are significantly

different. Serum levels of **(M)** Testosterone and **(N)** AMH in adult gonad intact females MC4R-

loxTB, *Kiss1-cre*:MC4R-LoxTB and their control littermates. *p<0.05, ****P < 0.0001, One Way

ANOVA. Data are presented as the mean± SEM.

Figure 4. Kiss1^{ARH} neurons are activated by melanocortin agonists and respond to

optogenetic stimulation of POMC neurons.

(A) Slices were taken from *Kiss1-cre* injected (confocal image) brains to target fluorescent cells

(arrowhead) for recording (white outline over electrode). **(B)** Whole cell voltage clamp recording

of Kiss1^{ARH} neurons following the direct application of the high affinity melanocortin receptor

agonist melanotan II (MTII, 250 nM) was added directly to the bath and the response was

compared between slices from OVX and OVX+E females. **(C)** While the average inward current

was slightly higher in the OVX+E state, this was not a significant difference (Student's t-test for

unpaired samples, p>0.05). **(D)** The MC4R-selective agonist THIQ (100 nM) was perfused, and

excitatory inward currents were generated in Kiss1^{ARH} neurons. **(E)** Images are of AAV-driven

labeling of POMC cells as seen through the confocal. **(F)** Using the AAV-driven expression of

channelrhodopsin in adult *Pomc-cre* mice, high frequency stimulation elicited a slow inward

current in Kiss1 neurons. **(G)** The identity of cells was confirmed through the presence of a

persistent sodium current (see [25]) and/or with RT-PCR of harvested cytoplasm showing *Kiss1*

expression. (Gel: MM=molecular marker, TC=tissue controls). **(H)** A direct synaptic projection

from POMC to Kiss1^{ARH} neuron (*posthoc* identified) was confirmed using the "rescue" protocol:

1, baseline glutamatergic responses were initially generated (black trace); 2, then action

potentials were eliminated by blocking voltage-gated sodium channels with TTX and the

postsynaptic response (red trace); 3, blockade of potassium channels facilitated calcium entry

into the terminal through ChR2 to release synaptic vesicles, "rescuing" the postsynaptic

glutamate response (green trace). **I)** Glutamatergic responses were also often observed,

particularly in the ventral ARH. As seen when targeting low input resistance neurons (i.e.,

Kiss1^{ARH}) with low frequency optogenetic stimulation (5 ms pulse, 50 ms inter-spike interval), the

first response in OVX + vehicle females was larger relative to the second response in OVX +

E2- treated female mice. Representative traces are the average of 30 sweeps. **(J)** The

averaged paired-pulse ratio was lower in recordings from estradiol-treated female mice,

indicating an increased release probability (Students t-test $p < 0.05$,). **(K)** High frequency

optogenetic stimulation elicited a small inward current. **(L)** In a different cell, a high frequency

inward current was noted before constant perfusion of SHU9119 for 15 minutes (break between

traces). When the stimulation protocol was repeated, no inward current was elicited. **(M)** The

majority of high frequency responses were inward and only twice was an inhibitory outward

current recorded in identified Kiss1^{ARH} neurons. Perfusion of the non-selective opioid receptor

antagonist naloxone reversed the current, eliminating MCR-activation as the mechanism.

Figure 5. Kiss1$^{AVPV/PeN}$ neurons are inhibited by MC4R agonists in an E2-dependent manner.

(A) Immunohistochemistry showing robust labeling of α-MSH fibers in the AVPV/PeN region in OVX+E2 WT female mice. **(B)** Low and high power bright-field images taken during electrophysiology recording in OVX+E2 POMC-Cre mice expressing YFP:ChR2 in the ARH following AAV injection. (left) Low power image shows the location of recorded cells. (right) Higher power image of area from white box inset displaying POMC fibers innervating the area. **(C)** Fibers surrounded putative Kiss1-AVPV neurons (expressing I_{NaP}, IT and I_h) and optogenetic stimuli was able to elicit postsynaptic currents that were eliminated with tetrodotoxin (1 μM), but "rescued" with addition of K$^+$ channel blockers (4-AP and TEA) which indicates a monosynaptic connection between POMC^{ARH} and Kiss1^{AVPV} neurons. **(D)** Whole-cell voltage clamp recordings were made in ChR2-YFP positive cells in brain slices taken from *Kiss1-cre*:Ai32 female mice. Bath application of melanotan II (MTII, 500 nM) generated an inhibitory outward current in a Kiss1$^{AVPV/PeN}$ neuron from OVX female. As the cell was synaptically isolated using bath-applied tetrodotoxin (TTX, 1 μM) this represents a direct effect. Washout of MTII while still in TTX quickly led to a return to baseline RMP. **(E)** An IV relationship was plotted using voltage steps before and after MTII administration. The crossing at -80 mV (~EK$^+$) indicates that the opening of K+ channels underlie the MC4R inhibition of Kiss1^{AVPV} neurons. **(F)** In a subset of recordings from OVX brains slices, the selective membrane estrogen receptor (Gq-mER) agonist STX (10 nM) was added to the bath for ~10 minutes prior to addition of MTII. STX pretreatment resulted in either a greatly attenuated outward current or even an inward current as shown in this example. **(G)** The mean outward current was significantly higher when recording in Kiss1$^{AVPV/Pen}$ neurons from brain slices from vehicle-treated, OVX females compared to E2-treated, OVX females or acute STX-treated brain slices from OVX females (One-way ANOVA $F_{(2,19)}=12.32$, $p<0.001$; Holm-Sidak *posthoc* comparisons found significant differences between all groups; * $p<0.05$, *** $p<0.001$). **(H)** There was no difference in the frequency of mEPSCs after MTII,

calculated as a percent of baseline between groups. **(I)** However, there was a significant

difference in the mEPSC amplitude (calculated as a percent of baseline) between all groups:

main effect $F_{(2,19)}$ =10.58, p<0.001. *Posthoc* comparisons using Holm-Sidak found OVX+STX to

be different from both OVX (p<0.001) and OVX+E (p<0.05).

References:

1. Health, U. (2020). Infertility. http://obgyn.ucla.edu/infertility
2. Andermann, M.L., and Lowell, B.B. (2017). Toward a Wiring Diagram Understanding of
Appetite Control. Neuron *95*, 757-778. 10.1016/j.neuron.2017.06.014.
3. Cone, R.D. (2006). Studies on the physiological functions of the melanocortin system.
Endocr Rev *27*, 736-749. 10.1210/er.2006-0034.
4. Shah, B.P., Vong, L., Olson, D.P., Koda, S., Krashes, M.J., Ye, C., Yang, Z., Fuller,
P.M., Elmquist, J.K., and Lowell, B.B. (2014). MC4R-expressing glutamatergic neurons
in the paraventricular hypothalamus regulate feeding and are synaptically connected to
the parabrachial nucleus. Proceedings of the National Academy of Sciences of the
United States of America *111*, 13193-13198. 10.1073/pnas.1407843111.
5. Wang, K., Mao, W., Zhang, X., Zhao, Y., Fan, K., Pan, D., Liu, H., Li, P., Hai, R., and Du,
C. (2020). Neuroanatomy of melanocortin-4 receptor pathway in the mouse brain. Open
Life Sci *15*, 580-587. 10.1515/biol-2020-0063.
6. Farooqi, I.S., Keogh, J.M., Yeo, G.S., Lank, E.J., Cheetham, T., and O'Rahilly, S. (2003).
Clinical spectrum of obesity and mutations in the melanocortin 4 receptor gene. The
New England journal of medicine *348*, 1085-1095. 10.1056/NEJMoa022050.
7. Hainerová, I.A., Zamrazilová, H., Sedláčková, D., and Hainer, V. (2011).
Hypogonadotropic hypogonadism in a homozygous MC4R mutation carrier and the
effect of sibutramine treatment on body weight and obesity-related health risks. Obes
Facts *4*, 324-328. 10.1159/000330763.
8. Doulla, M., McIntyre, A.D., Hegele, R.A., and Gallego, P.H. (2014). A novel MC4R
mutation associated with childhood-onset obesity: A case report. Paediatr Child Health
19, 515-518. 10.1093/pch/19.10.515.
9. Batarfi, A.A., Filimban, N., Bajouh, O.S., Dallol, A., Chaudhary, A.G., and Bakhashab, S.
(2019). MC4R variants rs12970134 and rs17782313 are associated with obese
polycystic ovary syndrome patients in the Western region of Saudi Arabia. BMC Med
Genet *20*, 144. 10.1186/s12881-019-0876-x.
10. Chen, X., Huang, L., Tan, H.Y., Li, H., Wan, Y., Cowley, M., Veldhuis, J.D., and Chen, C.
(2017). Deficient melanocortin-4 receptor causes abnormal reproductive neuroendocrine
profile in female mice. Reproduction (Cambridge, England) *153*, 267-276. 10.1530/REP-
16-0341.
11. Sandrock, M., Schulz, A., Merkwitz, C., Schoneberg, T., Spanel-Borowski, K., and
Ricken, A. (2009). Reduction in corpora lutea number in obese melanocortin-4-receptor-
deficient mice. Reprod Biol Endocrinol *7*, 24. 10.1186/1477-7827-7-24.
12. Cui, L., Tan, C., Huang, L., Wang, W., Huang, Z., Geng, F., Wu, M., Chen, X., Cowley,
M., Roelfsema, F., and Chen, C. (2022). Dapagliflozin partially restores reproductive
function in MC4R KO obese female mice. The Journal of endocrinology *254*, 65-76.
10.1530/JOE-21-0449.

899 13. Manfredi-Lozano, M., Roa, J., Ruiz-Pino, F., Piet, R., Garcia-Galiano, D., Pineda, R.,
900   Zamora, A., Leon, S., Sanchez-Garrido, M.A., Romero-Ruiz, A., et al. (2016). Defining a
901   novel leptin-melanocortin-kisspeptin pathway involved in the metabolic control of
902   puberty. Mol Metab *5*, 844-857. 10.1016/j.molmet.2016.08.003.

903 14. Goodman, R.L., Herbison, A.E., Lehman, M.N., and Navarro, V.M. (2022).
904   Neuroendocrine control of gonadotropin-releasing hormone: Pulsatile and surge modes
905   of secretion. Journal of neuroendocrinology *34*, e13094. 10.1111/jne.13094.

906 15. Israel, D.D., Sheffer-Babila, S., de Luca, C., Jo, Y.H., Liu, S.M., Xia, Q., Spergel, D.J.,
907   Dun, S.L., Dun, N.J., and Chua, S.C., Jr. (2012). Effects of leptin and melanocortin
908   signaling interactions on pubertal development and reproduction. Endocrinology *153*,
909   2408-2419. 10.1210/en.2011-1822.

910 16. Manfredi-Lozano, M., Roa, J., and Tena-Sempere, M. (2018). Connecting metabolism
911   and gonadal function: Novel central neuropeptide pathways involved in the metabolic
912   control of puberty and fertility. Front Neuroendocrinol *48*, 37-49.
913   10.1016/j.yfrne.2017.07.008.

914 17. Cravo, R.M., Margatho, L.O., Osborne-Lawrence, S., Donato, J., Jr., Atkin, S., Bookout,
915   A.L., Rovinsky, S., Frazao, R., Lee, C.E., Gautron, L., et al. (2011). Characterization of
916   Kiss1 neurons using transgenic mouse models. Neuroscience *173*, 37-56.
917   10.1016/j.neuroscience.2010.11.022.

918 18. Lam, B.Y.H., Williamson, A., Finer, S., Day, F.R., Tadross, J.A., Gonçalves Soares, A.,
919   Wade, K., Sweeney, P., Bedenbaugh, M.N., Porter, D.T., et al. (2021). MC3R links
920   nutritional state to childhood growth and the timing of puberty. Nature *599*, 436-441.
921   10.1038/s41586-021-04088-9.

922 19. Villa, P.A., Ruggiero-Ruff, R.E., Jamieson, B.B., Campbell, R.E., and Coss, D. (2024).
923   Obesity alters POMC and kisspeptin neuron crosstalk leading to reduced luteinizing
924   hormone in male mice. The Journal of Neuroscience, e0222242024.
925   10.1523/jneurosci.0222-24.2024.

926 20. Stephens, S.B.Z., and Kauffman, A.S. (2021). Estrogen Regulation of the Molecular
927   Phenotype and Active Translatome of AVPV Kisspeptin Neurons. Endocrinology *162*.
928   10.1210/endocr/bqab080.

929 21. Moore, A.M., Prescott, M., and Campbell, R.E. (2013). Estradiol negative and positive
930   feedback in a prenatal androgen-induced mouse model of polycystic ovarian syndrome.
931   Endocrinology *154*, 796-806. 10.1210/en.2012-1954.

932 22. McCarthy, E.A., Dischino, D., Maguire, C., Leon, S., Talbi, R., Cheung, E., Schteingart,
933   C.D., Riviere, P.J.M., Reed, S.D., Steiner, R.A., and Navarro, V.M. (2022). Inhibiting
934   Kiss1 Neurons With Kappa Opioid Receptor Agonists to Treat Polycystic Ovary
935   Syndrome and Vasomotor Symptoms. J Clin Endocrinol Metab *107*, e328-e347.
936   10.1210/clinem/dgab602.

937 23. Dewailly, D., Barbotin, A.L., Dumont, A., Catteau-Jonard, S., and Robin, G. (2020). Role
938   of Anti-Müllerian Hormone in the Pathogenesis of Polycystic Ovary Syndrome. Frontiers
939   in endocrinology *11*, 641. 10.3389/fendo.2020.00641.

940 24. Roa, J., and Herbison, A.E. (2012). Direct regulation of GnRH neuron excitability by
941   arcuate nucleus POMC and NPY neuron neuropeptides in female mice. Endocrinology
942   *153*, 5587-5599. 10.1210/en.2012-1470.

943 25. Zhang, C., Bosch, M.A., Qiu, J., Ronnekleiv, O.K., and Kelly, M.J. (2015). 17beta-
944   Estradiol increases persistent Na(+) current and excitability of AVPV/PeN Kiss1 neurons
945   in female mice. Mol Endocrinol *29*, 518-527. 10.1210/me.2014-1392.

946 26. Clements, J.D., and Westbrook, G.L. (1991). Activation kinetics reveal the number of
947   glutamate and glycine binding sites on the N-methyl-D-aspartate receptor. Neuron *7*,
948   605-613. 10.1016/0896-6273(91)90373-8.

27.   Stincic, T.L., Grachev, P., Bosch, M.A., Rønnekleiv, O.K., and Kelly, M.J. (2018).
Estradiol drives the anorexigenic activity of proopiomelanocortin neurons in female mice.
eNeuro *5*, e0103-0118.2018. 10.1523/eneuro.0103-18.2018.
28.   Qiu, J., Nestor, C.C., Zhang, C., Padilla, S.L., Palmiter, R.D., Kelly, M.J., and
Rønnekleiv, O.K. (2016). High-frequency stimulation-induced peptide release
synchronizes arcuate kisspeptin neurons and excited GnRH neurons. eLife *5*, e16246,
e16246. 10.7554/eLife.16246.
29.   Qiu, J., Rivera, H.M., Bosch, M.A., Padilla, S.L., Stincic, T.L., Palmiter, R.D., Kelly, M.J.,
and Rønnekleiv, O.K. (2018). Estrogenic-dependent glutamatergic neurotransmission
from kisspeptin neurons governs feeding circuits in females. eLife *7*, e35656, e35656.
10.7554/eLife.35656.
30.   Stincic, T.L., Qiu, J., Connors, A.M., Kelly, M.J., and Rønnekleiv, O.K. (2021). Arcuate
and Preoptic Kisspeptin neurons exhibit differential projections to hypothalamic nuclei
and exert opposite postsynaptic effects on hypothalamic paraventricular and
dorsomedial nuclei in the female mouse. eneuro, ENEURO.0093-0021.2021.
10.1523/eneuro.0093-21.2021.
31.   Herman, M.A., Ackermann, F., Trimbuch, T., and Rosenmund, C. (2014). Vesicular
glutamate transporter expression level affects synaptic vesicle release probability at
hippocampal synapses in culture. The Journal of Neuroscience *34*, 11781-11791.
10.1523/JNEUROSCI.1444-14.2014.
32.   Thornton, J.E., Loose, M.D., Kelly, M.J., and Rønnekleiv, O.K. (1994). Effects of
estrogen on the number of neurons expressing β-endorphin in the medial basal
hypothalamus of the female guinea pig. The Journal of comparative neurology *341*, 68-
77. 10.1002/cne.903410107.
33.   Zhang, C., Tonsfeldt, K.J., Qiu, J., Bosch, M.A., Kobayashi, K., Steiner, R.A., Kelly, M.J.,
and Ronnekleiv, O.K. (2013). Molecular mechanisms that drive estradiol-dependent
burst firing of Kiss1 neurons in the rostral periventricular preoptic area. American journal
of physiology. Endocrinology and metabolism *305*, E1384-1397.
10.1152/ajpendo.00406.2013.
34.   Mountjoy, K.G., Mortrud, M.T., Low, M.J., Simerly, R.B., and Cone, R.D. (1994).
Localization of the melanocortin-4 receptor (MC4-R) in neuroendocrine and autonomic
control circuits in the brain. Molecular Endocrinology *8*, 1298-1308.
10.1210/mend.8.10.7854347.
35.   Yu, J., Gimenez, L.E., Hernandez, C.C., Wu, Y., Wein, A.H., Han, G.W., McClary, K.,
Mittal, S.R., Burdsall, K., Stauch, B., et al. (2020). Determination of the melanocortin-4
receptor structure identifies Ca^{2+} as a cofactor for ligand binding. Science *368*, 428-433.
doi:10.1126/science.aaz8995.
36.   Sohn, J.W., Harris, L.E., Berglund, E.D., Liu, T., Vong, L., Lowell, B.B., Balthasar, N.,
Williams, K.W., and Elmquist, J.K. (2013). Melanocortin 4 receptors reciprocally regulate
sympathetic and parasympathetic preganglionic neurons. Cell *152*, 612-619.
10.1016/j.cell.2012.12.022.
37.   Kelly, M.J., Loose, M.D., and Ronnekleiv, O.K. (1992). Estrogen suppresses mu-opioid-
and GABAB-mediated hyperpolarization of hypothalamic arcuate neurons. The Journal
of neuroscience : the official journal of the Society for Neuroscience *12*, 2745-2750.
10.1523/JNEUROSCI.12-07-02745.1992.
38.   Lagrange, A.H., Ronnekleiv, O.K., and Kelly, M.J. (1994). The potency of mu-opioid
hyperpolarization of hypothalamic arcuate neurons is rapidly attenuated by 17 beta-
estradiol. The Journal of neuroscience : the official journal of the Society for
Neuroscience *14*, 6196-6204. 10.1523/JNEUROSCI.14-10-06196.1994.
39.   Qiu, J., Bosch, M.A., Tobias, S.C., Grandy, D.K., Scanlan, T.S., Ronnekleiv, O.K., and
Kelly, M.J. (2003). Rapid signaling of estrogen in hypothalamic neurons involves a novel

G-protein-coupled estrogen receptor that activates protein kinase C. The Journal of neuroscience : the official journal of the Society for Neuroscience *23*, 9529-9540. 10.1523/JNEUROSCI.23-29-09529.2003.

40. Qiu, J., Bosch, M.A., Tobias, S.C., Krust, A., Graham, S.M., Murphy, S.J., Korach, K.S., Chambon, P., Scanlan, T.S., Rønnekleiv, O.K., and Kelly, M.J. (2006). A G-protein-coupled estrogen receptor is involved in hypothalamic control of energy homeostasis. The Journal of neuroscience : the official journal of the Society for Neuroscience *26*, 5649-5655. 10.1523/jneurosci.0327-06.2006.

41. Qiu, J., Rivera, H.M., Bosch, M.A., Padilla, S.L., Stincic, T.L., Palmiter, R.D., Kelly, M.J., and Ronnekleiv, O.K. (2018). Estrogenic-dependent glutamatergic neurotransmission from kisspeptin neurons governs feeding circuits in females. Elife *7*. 10.7554/eLife.35656.

42. Stincic, T.L., Grachev, P., Bosch, M.A., Ronnekleiv, O.K., and Kelly, M.J. (2018). Estradiol Drives the Anorexigenic Activity of Proopiomelanocortin Neurons in Female Mice. eNeuro *5*. 10.1523/ENEURO.0103-18.2018.

43. Ghamari-Langroudi, M., Digby, G.J., Sebag, J.A., Millhauser, G.L., Palomino, R., Matthews, R., Gillyard, T., Panaro, B.L., Tough, I.R., Cox, H.M., et al. (2015). G-protein-independent coupling of MC4R to Kir7.1 in hypothalamic neurons. Nature *520*, 94-98. 10.1038/nature14051.

44. Krause, W.C., Rodriguez, R., Gegenhuber, B., Matharu, N., Rodriguez, A.N., Padilla-Roger, A.M., Toma, K., Herber, C.B., Correa, S.M., Duan, X., et al. (2021). Oestrogen engages brain MC4R signalling to drive physical activity in female mice. Nature *599*, 131-135. 10.1038/s41586-021-04010-3.

45. Brix, N., Ernst, A., Lauridsen, L.L.B., Parner, E., Stovring, H., Olsen, J., Henriksen, T.B., and Ramlau-Hansen, C.H. (2019). Timing of puberty in boys and girls: A population-based study. Paediatr Perinat Epidemiol *33*, 70-78. 10.1111/ppe.12507.

46. Padilla, S.L., Johnson, C.W., Barker, F.D., Patterson, M.A., and Palmiter, R.D. (2018). A Neural Circuit Underlying the Generation of Hot Flushes. Cell Rep *24*, 271-277. 10.1016/j.celrep.2018.06.037.

47. Dror, T., Franks, J., and Kauffman, A.S. (2013). Analysis of multiple positive feedback paradigms demonstrates a complete absence of LH surges and GnRH activation in mice lacking kisspeptin signaling. Biology of reproduction *88*, 146. 10.1095/biolreprod.113.108555.

48. Steyn, F.J., Wan, Y., Clarkson, J., Veldhuis, J.D., Herbison, A.E., and Chen, C. (2013). Development of a methodology for and assessment of pulsatile luteinizing hormone secretion in juvenile and adult male mice. Endocrinology *154*, 4939-4945. 10.1210/en.2013-1502.

49. Talbi, R., Ferrari, K., Choi, J.H., Gerutshang, A., McCarthy, E.A., Dischino, D., Leon, S., and Navarro, V.M. (2021). Characterization of the Action of Tachykinin Signaling on Pulsatile LH Secretion in Male Mice. Endocrinology *162*. 10.1210/endocr/bqab074.

50. Dave, J.R., Rubinstein, N., and Eskay, R.L. (1985). Evidence that b-endorphin binds to specific receptors in rat peripheral tissues and stimulates the adenylate cyclase-adenosine 3',5'-monophosphate system. Endocrinology *117*, 1389-1396.

51. Naulé, L., Robert, V., Parmentier, C., Martini, M., Keller, M., Cohen-Solal, M., Hardin-Pouzet, H., Grange-Messent, V., Franceschini, I., and Mhaouty-Kodja, S. (2015). Delayed pubertal onset and prepubertal Kiss1 expression in female mice lacking central oestrogen receptor beta. Hum Mol Genet *24*, 7326-7338. 10.1093/hmg/ddv430.

52. Pfaffl, M.W. (2001). A new mathematical model for relative quantification in real-time RT-PCR. Nucleic Acids Res *29*, e45. 10.1093/nar/29.9.e45.

53. Balthasar, N., Coppari, R., McMinn, J., Liu, S.M., Lee, C.E., Tang, V., Kenny, C.D., McGovern, R.A., Chua, S.C., Jr., Elmquist, J.K., and Lowell, B.B. (2004). Leptin receptor

signaling in POMC neurons is required for normal body weight homeostasis. Neuron *42*,
983-991. 10.1016/j.neuron.2004.06.004.
54. Madisen, L., Mao, T., Koch, H., Zhuo, J.M., Berenyi, A., Fujisawa, S., Hsu, Y.W., Garcia,
A.J., 3rd, Gu, X., Zanella, S., et al. (2012). A toolbox of Cre-dependent optogenetic
transgenic mice for light-induced activation and silencing. Nature Neuroscience *15*, 793-
802. 10.1038/nn.3078.
55. Bosch, M.A., Tonsfeldt, K.J., and Ronnekleiv, O.K. (2013). mRNA expression of ion
channels in GnRH neurons: subtype-specific regulation by 17beta-estradiol. Molecular
and cellular endocrinology *367*, 85-97. 10.1016/j.mce.2012.12.021.
56. Qiu, J., Nestor, C.C., Zhang, C., Padilla, S.L., Palmiter, R.D., Kelly, M.J., and
Ronnekleiv, O.K. (2016). High-frequency stimulation-induced peptide release
synchronizes arcuate kisspeptin neurons and excites GnRH neurons. Elife *5*.
10.7554/eLife.16246.
57. Stincic, T.L., Qiu, J., Connors, A.M., Kelly, M.J., and Ronnekleiv, O.K. (2021). Arcuate
and Preoptic Kisspeptin Neurons Exhibit Differential Projections to Hypothalamic Nuclei
and Exert Opposite Postsynaptic Effects on Hypothalamic Paraventricular and
Dorsomedial Nuclei in the Female Mouse. eNeuro *8*. 10.1523/ENEURO.0093-21.2021.
58. Qiu, J., Stincic, T.L., Bosch, M.A., Connors, A.M., Kaech Petrie, S., Ronnekleiv, O.K.,
and Kelly, M.J. (2021). Deletion of Stim1 in Hypothalamic Arcuate Nucleus Kiss1
Neurons Potentiates Synchronous GCaMP Activity and Protects against Diet-Induced
Obesity. The Journal of neuroscience : the official journal of the Society for
Neuroscience *41*, 9688-9701. 10.1523/JNEUROSCI.0622-21.2021.
1073

A. Kiss1 neuron

B. MT II

C.

D. THIQ

E. Pomc^Cre Injected · 3V · 200μm

F. 470 nm · POMC · Kiss1 ARH

G. MM · Recorded Cells · 1 2 3 4 5 · TC · + − · 400− 200− 100− · Kiss1 120 bp

H. 5 ms · 2) TTX · 10 pA · 20 ms · 3) TTX+4AP +TEA · 1) Baseline

I. 5 ms · 5 pA · OVX · 5 pA · OVX+E

J.

K. 10 s, 20 Hz · 20 pA

L. 10 s, 20 Hz · SHU 9119 Pretreatment · 20 pA

M. 10 s, 20 Hz · Naloxone · 20 pA · 1 m