Polyamine-mediated sensitization of *Klebsiella pneumoniae* to macrolides through a dual

mode of action

Joshua M. E. Adams[1], Peri B. Moulding[1] and Omar M. El-Halfawy[1,2*]

[1] Department of Chemistry and Biochemistry, Faculty of Science, University of Regina, Regina,

SK, S4S 0A2, Canada

[2] Department of Microbiology and Immunology, Faculty of Pharmacy, Alexandria University,

Alexandria, 21521, Egypt

[*] **Correspondence: omar.el-halfawy@uregina.ca (O.M.E.)**

Running title: Uncovering new polyamine macrolide antibiotic adjuvants

Abstract

The chemical landscape bacteria encounter at the infection site could shape their stress and

antibiotic responses; these altered responses are not typically detected in standard lab conditions.

Polyamines are small molecules typically overproduced by the host during infection and have

been shown to alter bacterial stress responses. We sought to determine the effect of polyamines

on the antibiotic response of *Klebsiella pneumoniae*, a Gram-negative priority pathogen.

Interestingly, putrescine and other natural polyamines sensitized *K. pneumoniae,* including pan-

drug resistant and hypervirulent isolates, to azithromycin, a macrolide protein translation

inhibitor typically used for Gram-positive bacteria. The synergy was further potentiated in the

physiological buffer, bicarbonate. Chemical genomic screens suggested a dual mechanism

whereby putrescine acts at the membrane and ribosome levels. Putrescine permeabilized both the

outer and inner membranes of *K. pneumoniae* in fluorometric (NPN) and colorimetric (β-

lactamase and β-galactosidase) assays. Further, mutants with chemically and genetically

perturbed membranes exhibited a loss of putrescine-azithromycin synergy. Putrescine also

inhibited protein synthesis in an *Escherichia coli*-derived cell-free protein expression assay

simultaneously monitoring transcription and translation. Profiling the putrescine-azithromycin

synergy against a combinatorial array of antibiotics targeting various ribosomal sites suggested

that putrescine acts as tetracyclines targeting the 30S ribosomal acceptor site. Next, we exploited

the natural polyamine-azithromycin synergy to uncover azithromycin adjuvants by screening a

polyamine analog library. We discovered four azithromycin synergists with activity from the low

micromolar range, sharing modes of action similar to putrescine. This work sheds light on the

bacterial antibiotic responses under conditions more reflective of those at the infection site and

provides a new strategy to extend the macrolide spectrum to drug-resistant *K. pneumoniae*.

Keywords: Polyamines; macrolides; putrescine; Gram-negative bacteria; antibiotic adjuvants

Introduction

Antimicrobial resistance (AMR) is an increasingly rising problem gravely impacting today's society. In 2019, there was an estimated 4.95 million deaths worldwide associated with bacterial AMR [1]. The AMR public health crisis prompted the World Health Organization (WHO) to issue a list of priority pathogens posing the most serious threat that requires urgent action; Gram-negative bacteria make up the majority of this list due to the limited therapeutic options available against these bacteria[2-4]. Unlike Gram-positive bacteria, the Gram-negative ones possess an outer membrane, an integral component of which are lipopolysaccharides (LPS), which constitutes a formidable barrier that prevents antibiotic permeation [5]. *Klebsiella pneumoniae* is one of the Gram-negative bacteria on the WHO priority pathogens list, gaining international concern due to the high prevalence of hypervirulent and drug resistant strains [4]. With the constant rise in AMR and the inadequate antibiotic drug discovery pipeline [6], there is a dire need for new strategies to combat Gram-negative resistant strains[1].

Bacteria encounter a myriad of chemicals at the infection site that may alter their antibiotic susceptibility and virulence [7, 8]. The potential effects of such chemicals are not typically captured in standard antibiotic susceptibility assays, whose conditions follow the guidelines issued by standardizing agencies such as the Clinical and Laboratory Standards institute (CLSI). Polyamines constitute an important class of chemicals present at the infection sites, especially since they are overproduced by the host during infection [9]. Natural polyamines, such as spermine, spermidine, and putrescine, are small polycationic molecules produced by almost all living organisms and are critical for their viability [10]. During infection, polyamine biosynthesis is upregulated in regenerating tissues of the host at the inflammatory and infection site; for example, polyamines accumulate in the lungs during pneumonia infections, suggesting a

role in suppressing infection [11-13]. Additionally, polyamines play an immunomodulatory role,

where spermine exhibits macrophage suppression and anti-inflammatory activity [12]. The

polyamine backbone structure consists of regularly spaced positive charges interrupted by

hydrophobic hydrocarbon bridges; this unique structure allows polyamines to interact with

various macromolecules [14]. Polyamines can stabilize nucleic acids and thus influence

transcription, chromosomal structure, apoptosis, and protein synthesis [15-20]; they are involved in

cell growth, differentiation, cell membrane stability and immune responses in humans [14, 15].

Polyamines are similarly involved in cellular homeostasis and stress response in bacteria

[21]. They can also alter antibiotic susceptibility in a wide range of Gram-positive and Gram-

negative bacteria [22-27]. For example, polyamines increased resistance of *Pseudomonas*

aeruginosa and *Burkholderia cenocepacia* to antibiotics, including polymyxins and

fluoroquinolones [23-25], and conversely increased the susceptibility of *Staphylococcus aureus* and

enteric bacteria, such as *Escherichia coli*, to β-lactam antibiotics [26].

Macrolides are a class of antibiotics that target and inhibit protein synthesis via binding to

the 23S portion of the 50S bacterial ribosome subunits[28]. Antibiotics in this class are highly

prescribed due to their potency, low toxicity, and favourable pharmacokinetic properties,

particularly their oral bioavailability [29-31]. Newer classes of macrolides, such as the azalide

azithromycin, are clinically advantageous due to their extensive and rapid distribution into

tissues and intracellular compartments yet minimal accumulation in fat and muscle [30]. Typically,

macrolides are used to treat Gram-positive infections with less application in Gram-negative

ones due to their poor penetration of the outer membrane [32]. However, macrolides show potency

against Gram-negative bacteria albeit under certain infection-relevant conditions. For example,

E. coli exhibit heightened sensitivity to macrolides in the presence of the physiologically relevant

buffer bicarbonate, which interferes with the proton motive force increasing transport of

macrolides through the inner membrane [33]. Understanding the effects of macrolides against

priority Gram-negative bacteria, such as *K. pneumoniae*, especially under host relevant

conditions, may expand their spectrum of activity to this group of bacteria where the clinical

need is high.

In this study, we show that polyamines sensitize *K. pneumoniae* to the macrolide

azithromycin. We determined the mechanism of this synergy, revealing both a membrane and

target-dependent effects. These observations prompted us to screen a collection of synthetic

polyamine analogs for macrolide adjuvants against *K. pneumoniae*, identifying micromolar

azithromycin synergists. Together, this work offers potential new antimicrobial strategies against

multi-drug resistant Gram-negative bacteria.

Results and discussion

Natural polyamines synergize with azithromycin.

We conducted checkerboard assays to test the effect of the polyamine putrescine on

azithromycin susceptibility in *K. pneumoniae* MKP103, a derivative of the clinical outbreak

isolate KPNIH1 lacking its carbapenemase-encoding gene [34]. Putrescine synergized with

azithromycin with a fractional inhibitory concentration index (FICI) of 0.47±0.01 (Fig 1A). FICI

≤ 0.5 were interpreted as synergy, FICI > 4.0 as antagonism, and 0.5 < FICI ≤4.0 as additive,

indifference, or no interaction. This synergy matches previously reported polyamine-induced

macrolide sensitivity in another Gram-negative organism, *E. coli* [26]; the mechanism of such

interaction is unknown. The putrescine-azithromycin synergy reversed macrolide resistance of *K.*

pneumoniae MKP103; putrescine at 20 mM lowered the azithromycin MIC to below the CLSI

clinical breakpoint (Fig 1A). A bacterial strain is deemed sensitive to an antibiotic at MIC values

less than the clinical breakpoint, thus may become clinically used[35]. We also observed

azithromycin synergy with other natural polyamines (spermine and spermidine, Fig 1B-C).

Polyamine physiological levels vary but are generally within the millimolar range and are further

increased at the infection site upon infection [36, 37]; hence polyamines at the site of infection may

be sufficient to drive azithromycin synergy similar to what we observed *in vitro*.

The polyamine-azithromycin synergy was maintained across diverse *K. pneumoniae*

clinical isolates. We conducted putrescine-azithromycin checkerboard assays against the *K.*

pneumoniae diversity panel, comprising 100 clinical isolates collected from 63 facilities in 19

countries with an array of genotypic and phenotypic differences that span pan-sensitive, pan-

resistant, and hypervirulent lineages (Supplementary Table 1) [38]. We observed the putrescine-

azithromycin synergy in 78 isolates (Fig 1D), which lowered azithromycin MIC below the CLSI

breakpoint in 77 isolates (Supplementary Table 1; examples in Fig 1E). Notably, such synergy

was conserved across pan-sensitive to pan-drug-resistant isolates, including those encoding

carbapenemase, OXA, NDM, and CTX β-lactamases, Rmt(B,F,H) methyltransferases, MCR-1,

Tet(X), and ErmB, which drive resistance against carbapenems, β-lactams, aminoglycosides,

colistin, tetracyclines, and erythromycin, respectively (supplementary Table 1). On the other

hand, the putrescine-azithromycin combination showed additive effects in 22 isolates, which

harbour genes encoding the macrolide phosphotransferases Mph(A), Mph(B), and Mph(E)

(Supplementary Table 1).

Figure 1. **Natural polyamines synergize with azithromycin against K. pneumoniae rendering it susceptible to the macrolide antibiotic.** A-C) Representative checkerboard assays of K. pneumoniae MKP103 of azithromycin combined with putrescine, spermine, and spermidine. D) A scatter plot representing the distribution of FICI values across the K. pneumoniae diversity panel. E) Representative checkerboard assays of select clinical isolates from the diversity panel. F) A checkerboard assay of putrescine and azithromycin against K. pneumoniae MKP103 in the presence of 25 mM bicarbonate buffer. G) Chemical structures of the polyamines used in this study and azithromycin. Dark blue regions represent high cell density determined by $OD_{600.}$ Red dotted lines refer to the clinical breakpoint as set by CLSI. Mean of the FICI and SEM (standard error of the mean) are displayed on each representative checkerboard, n=3.

Next, we sought to test whether polyamines further sensitize *K. pneumoniae* to azithromycin in the presence of bicarbonate, the physiological buffer, in which Gram-negative bacteria are less resistant to macrolides [33]. We show that exposure to the physiological bicarbonate concentration (25 mM) resulted in 4-fold reduction in azithromycin MIC against *K. pneumoniae* (Fig 1F compared to 1A), as previously reported [33]. Putrescine further sensitized *K. pneumoniae* to azithromycin (further 16-fold reduction in MIC) in the presence of bicarbonate

(Fig 1F). These observations suggest that multiple physiologically-relevant factors may act in

tandem to alter bacterial responses to antibiotics.

Genomic screens identify a dual mechanism of the polyamine-azithromycin synergy.

To identify the mechanism of polyamine-azithromycin synergy, we hypothesized that a

mutant in the gene encoding the determinant of synergy will show increased azithromycin

susceptibility and loss in synergy with polyamines. We used an arrayed sequence-defined

transposon mutant library covering the non-essential genome of *K. pneumoniae* MKP103,

comprising one or multiple non-redundant mutants for each non-essential gene, with an average

of 2.5 mutants per gene [34]. First, we screened the library at two azithromycin sub-inhibitory

concentrations, 32 and 64 µg/mL, to identify mutants susceptible to azithromycin (Fig 2A-B and

Supplementary Fig 1). The screen revealed 71 mutants representing 67 genes showing partial or

complete growth inhibition at the sub-inhibitory azithromycin concentrations (Fig 2A-B and

Supplementary Fig 1). These genes include those whose products are involved in stress induced

mutagenesis (*recC* and *recB*), ribosomal function (*rpsO, rpsR, ybeY,* and *rsmB*), and membrane

integrity (*waaD, waaC, waaE, dsbA, acrB,* and *bamB*); knockouts of genes involved in these

functions have been previously identified in either *K. pneumoniae* or *E. coli* or both as more

susceptible to macrolides and other protein translation inhibitors [39-41].

Figure 2. **A chemogenomic screen identifies putrescine-azithromycin synergy determinants.** A) A flow chart of the chemogenomic screen process that led to the identification of genetic determinants of azithromycin-putrescine synergy B) An index plot of the primary screen of *K. pneumoniae* MKP103 mutants grown in 32 μg/mL of azithromycin shown as % growth relative to that of the untreated mutants. Data points below a cut-off (the red line) of 1.5x the standard deviation of all datapoints below the interquartile mean are considered putative hits. C-F) Heat maps of representative putrescine-azithromycin checkerboards of the *K. pneumoniae* MKP103 mutants identified from the chemogenomic screen. Dark blue represents maximal growth determined by OD_{600}. Mean of the FICI and SEM are displayed on each representative checkerboard, n=3.

Next, we conducted putrescine-azithromycin checkerboard assays against the 71 mutants

(Supplementary Fig 2 and Fig 2C-F). Azithromycin sensitivity (MIC ≤ 32μg/mL) and lack of

synergy (FICI >0.5), suggested mutants of genetic determinants of the putrescine-azithromycin

synergy; mutants with disruption in four genes (*waaD, waaC*, *rpsO,* and *rpsR*) fit these criteria

(Fig 2C-F). The disruptions in these mutants can be categorized into membrane (*waaD*::Tn and

waaC::Tn) and 30S ribosomal subunit (*rpsO*::Tn and *rpsR*::Tn) related disruptions.

Knockouts of *waaD* and *waaC* lead to the same deep rough LPS phenotype [42-44]. WaaD is

necessary for the production of ADP-L-glycero-D-mannoheptose, which WaaC incorporates into

the LPS structure; loss of these enzymes results in a truncated LPS without an outer core and

most of the inner core exposing Kdo2-lipid A (Supplementary Fig 3A-B)[42]. Notably, the

waaD::Tn was hypersensitive to novobiocin (Supplementary Fig 4), as expected for LPS deep

rough mutants [42, 45]. The azithromycin hypersensitivity of these *K. pneumoniae* mutants is in

agreement with previous reports of *E. coli* with a compromised LPS structure being more

sensitive to macrolides [39, 46]. Setting the azithromycin susceptibility cutoff to MIC $\leq$64µg/mL led

to the inclusion of *dsbA*::Tn, *bamB*::Tn, and KPNIH1_25560::Tn (predicted *waaE* [44]), which

exhibited partial loss of synergy (Supplementary Fig 2A-C and Supplementary Fig 3C). Loss of

DsbA leads to accumulation of inactive LptD (a lipopolysaccharide assembly protein), BamB is

a lipoprotein essential for assembly and insertion of β-barrel proteins into the outer membrane,

and WaaE is responsible for the transfer of the branch β-D-glucopyranose (Glc) to L-glycero-D-

manno-heptose I (Heptose I) [44, 47, 48]. Loss of *dsbA* and *waaE* has been previously shown to

exhibit macrolide sensitivity in *E. coli* and *K. pneumoniae,* respectively [41, 44], whereas knockouts

of *bamB* exhibit loss of a membrane barrier effect in *E. coli,* becoming sensitive to novobiocin [48].

We tested other mutants involved in LPS biosynthesis to determine the extent of LPS

truncation necessary for the loss of putrescine-azithromycin synergy. WaaF and WaaQ are

enzymes responsible for the sequential addition of heptose II and heptose III, respectively, to

complete the LPS inner core [44]. Knockouts of genes encoding both enzymes showed no change

in azithromycin MIC and retained putrescine-azithromycin synergy (Supplementary Fig 3D-E).

Together, a deep rough truncation of the LPS core leads to loss of putrescine-azithromycin

synergy, suggesting that a similar polyamine-mediated membrane permeabilization induces the

observed synergy (Supplementary Fig 3). For subsequent follow up assays, we used *waaD*::Tn as

a representative of the deep rough mutants.

The ribosomal proteins RpsO, RpsR, and RpsF create a complex within the platform

domain of the 30S ribosomal subunit, where RpsR and RpsF form heterodimers [49]. Both

rpsO::Tn and *rpsR*::Tn mutants exhibited an azithromycin MIC of 32 µg/mL and an FICI value

of 0.7 ± 0.04 and 0.6 ± 0.04, respectively (Fig 2G-H). RpsF is not represented in the *K.*

pneumoniae MKP103 library. The deletion of *rpsO* has been reported to influence the functional

capacity of 30S subunits and decrease the amount of 70S complexes [50]. RpsR is hypothesized to

be involved in the binding of aminoacyl-tRNA to the P site of the ribosome [51]. Genes encoding

other 30S ribosomal subunit proteins are essential except for *rpsA,* whose mutant yielded no

change in azithromycin susceptibility (data not shown). RpsA is weakly associated with the 30S

ribosome and even lacks association to the ribosome in some *E. coli* strains [52]. Notably, the

checkerboard of *infB*::Tn, a mutant of the translation initiation factor for *rpsO*, exhibited a partial

reduction in synergy and a 4-fold decrease in azithromycin MIC compared to that of the wild

type (Supplementary Fig 2D).

Together, our results suggest that polyamines synergize with azithromycin via a dual

mechanism of action: (1) increasing membrane permeability allowing azithromycin to cross this

barrier, and (2) acting on the 30S ribosomal subunit compounding the effects of azithromycin in

suppressing protein synthesis.

Putrescine permeabilizes bacterial outer and inner membranes.

Gram-negative bacteria, including *K. pneumoniae*, possess outer and inner membranes. We studied the effects of polyamines on both membranes. Starting with the outer membrane, we showed, in kinetic 1-*N*-phenylnaphthylamine (NPN) fluorometric assays, that putrescine disrupted the outer membrane of *K. pneumoniae* shown as an increase in NPN uptake (Fig 3A). A disruption of the outer membrane occurs at 10 mM and 20 mM of putrescine (P < 0.0001), which is within the synergistic range of the putrescine-azithromycin interaction. Such putrescine-mediated outer membrane perturbation was similar to that displayed by the deep rough LPS mutant *waaD*::Tn, which showed a similar increase in NPN uptake relative to the wild type (P < 0.005, Fig 3B). *Salmonella enterica* serovar Typhimurium and *E. coli waaD* deletion mutants also showed a similar (2 to 5-fold) increase in NPN uptake in comparison to the respective parent strains [53, 54]. Using colorimetric quantification of the hydrolysis of nitrocefin, a β-lactam probe, by a periplasmic β-lactamase constitutively expressed from pBR322 as another proxy for outer membrane perturbation, we further confirmed the increased outer membrane permeability of the *waaD*::Tn mutant in comparison to its parent strain (Fig 3C), in agreement with previous reports [55].

Given that outer membrane permeabilization of the deep rough mutant (Fig 3B-C) is sufficient to enable macrolides to cross that barrier (Fig 1A, Fig 2C-D, [39]), further membrane perturbation may not lead to enhancement of the macrolide activity. Thus, the lack of putrescine-azithromycin synergy in *waaD*::Tn further suggests that the polyamine-mediated outer membrane perturbation contributes to the synergy. As such, we reasoned that chemically or genetically perturbing the membrane should exhibit increased susceptibility to azithromycin and a loss of synergy with putrescine, similar to *waaD*::Tn. Chemically perturbing *K. pneumoniae* membrane by 1 µg/mL of polymyxin B nonapeptide (PMBN), a cationic peptide that disrupts the

outer membrane with little to no antimicrobial activity [56], led to a 4-fold decrease in

azithromycin MIC (Fig 3 D-E), consistent with previous reports [5, 56, 57], and a loss of putrescine-

azithromycin synergy (FICI 0.83 ± 0.10) (Fig 3E). Similarly, an *E. coli* with a compromised

outer membrane genetically-induced by the deletion of 9 efflux pump transmembrane proteins

(Δ9 mutant) [58], exhibited an 8-fold decrease in azithromycin MIC and a loss in synergy (FICI =

1.00 ± 0.00) relative to its parent strain (M6394), which showed a synergistic phenotype (FICI =

0.41 ± 0.07) (Supplementary Fig 5A-B). The *E. coli* wild type and Δ9 mutant showed NPN

profiles (Supplementary Fig 5C) similar to those of *K. pneumoniae* MKP103 wild type and

waaD::Tn (Fig 3B).

Figure 3. **Putrescine permeabilizes the outer and inner membranes of *K. pneumoniae* to azithromycin.** A-B)
Fluorometric kinetic assays displaying % NPN uptake in wild type *K. pneumoniae* MKP103 treated with different
concentrations of putrescine (A) and in *K. pneumoniae waaD*::Tn relative to its parent MKP103 strain (B). C) A
kinetic absorbance assay displaying nitrocefin hydrolysis by a periplasmic β-lactamase in *K. pneumoniae* wild type
and *K. pneumoniae waaD*::Tn. D-E) Representative checkerboards of putrescine-azithromycin combination against
K. pneumoniae MKP103 in the absence (D) and presence (E) of 1 µg/mL polymyxin B nonapeptide; dark blue
regions represent high cell density determined by OD_{600}. F) A kinetic absorbance assay of ONPG hydrolysis in *E.
coli* ML-35 as a proxy for inner membrane perturbation. All kinetic assays are the combined data of three

independent assays each done in triplicate. All error bars denote SEM. Mean of the FICI and SEM are displayed on
each representative checkerboard, n=3.
265

Notably, spermidine and spermine at the micromolar range have been reported to confer

resistance to polymyxin B in *P. aeruginosa* by localizing at the outer membrane. Molecular

dynamics simulations of model bacterial outer membranes have shown that polyamines engage

in electrostatic interactions with negatively charged phosphate groups, stabilizing the outer

membrane with the largest stabilizing effects in the order of spermidine, spermine, then

putrescine [59]. The activity of putrescine at the outer membrane in *K. pneumoniae* at the

physiologically relevant millimolar range has not been previously reported. At low putrescine

concentrations, we confirmed that putrescine can stabilize the outer membrane shown as reduced

NPN uptake relative to the untreated cells (Supplementary Fig 6). As such, it appears that

putrescine displays a dose-dependent activity on the outer membrane, ranging from stabilization

at low concentrations to disruption at higher concentrations.

The inner membrane of Gram-negative bacteria may constitute a second physical barrier

for azithromycin. We sought to test if putrescine also affects the inner membrane by measuring

the β-galactosidase-mediated o-nitrophenyl-β-D-galactopyranoside (ONPG) hydrolysis in a

lactose permease-deficient *E. coli* that constitutively expresses a cytoplasmic β-galactosidase (*E.*

coli ML-35). Putrescine at 40 mM significantly increased ONPG hydrolysis (P < 0.0001, Fig

3F), suggesting that polyamines can also disrupt the inner membrane, potentially contributing to

increasing intracellular azithromycin concentration. Taken together, we have shown evidence

that putrescine permeabilizes both the outer and inner membranes, contributing to its synergy

with azithromycin.

Polyamines act on the ribosome compounding the protein synthesis inhibitory effect of

azithromycin.

The lack of azithromycin-putrescine synergy in the 30S ribosomal subunit mutants

(*rpsO*::Tn and *rpsR*::Tn, Fig 2E-F) suggests the synergy may be driven, at least in part, by an

interaction between both molecules at the ribosomal level. Hence, we sought to test the effect of

putrescine on protein translation. To this end, we used a cell-free protein synthesis system, an

extract-based transcription/translation system derived from *E. coli* designed to synthesize

proteins under the control of T7 RNA Polymerase. We quantified the levels of a plasmid-

encoded destabilized enhanced Green Fluorescent Protein (deGFP) fluorometrically in the

presence and absence of putrescine (Fig 4 and Supplementary Fig 7). Simultaneously, we

measured the effects on transcriptional activity in real-time (Fig 4 and Supplementary Fig 7A-C),

since the plasmid also included a malachite green mRNA aptamer allowing for such

measurement fluorometrically via binding of malachite green dye [60]. At the higher tested

concentration of putrescine, mRNA levels were reduced but protein synthesis was further

inhibited (Supplementary Fig 7A-F). Similar reduction in gene expression has been previously

observed in cell-free *in vitro* gene expression models at high spermine and spermidine

concentrations [61]. Thus, to determine the effect of putrescine on protein synthesis due to a

ribosomal effect, the deGFP signal was normalized by the mRNA signal (Fig 4). The lower

putrescine concentrations exhibited increased protein synthesis (in the first ~90 minutes of

exposure, Fig 4), which is in agreement with previous reports that low concentrations of

polyamines promote protein synthesis in cell free systems [61]. A majority of intracellular

polyamines exist as polyamine-RNA and polyamine-ribosome complexes with evidence

suggesting they contribute to transcription and translation, playing an essential role in ribosome

biogenesis, tRNA positioning on the ribosome, and translation elongation speed [62-65]. Cross-

linking studies showed that ribosomes have >500 stably linked polyamines per ribosome, and it

has been suggested polyamine binding happens in a stochastic manner [65, 66]. On the other hand,

our results show that 80 mM of putrescine marked a significant decrease in protein synthesis,

suggesting a decreased ribosomal function of the cell-free protein expression system (P=0.0001,

Fig 4). The addition of spermine to ribosome complexes was previously shown to slow down

protein expression, inhibit peptidyl transferase activity, and possibly alter ribosome conformation

[67, 68].

Figure 4. **Putrescine inhibits protein synthesis**. Protein (deGFP) and mRNA levels were simultaneously monitored
fluorometrically (433/475 and 610/650 nm, respectively) in black 384 well plates using the NEBExpress™ cell-free
protein expression system in 200 mM HEPES buffer with an engineered plasmid encoding deGFP with a malachite
green mRNA aptamer. Results shown are normalized by dividing each well's RFU deGFP value by the malachite
green RFU value and converted into a percentage of the maximal signal observed in the untreated condition. Raw
RFU data found in supplementary Fig 7A-F. All error bars denote SEM. A one-way ANOVA analysis followed by a
Brown-Forsythe and Welch test assessed the significance of the difference as indicated on the plot.

Next, we sought to determine the potential target site of putrescine within the ribosome

through a phenotypic combinatorial approach using antibiotics targeting different ribosomal sites

and a control (rifampicin) acting outside the ribosome. We hypothesize that, in the wild type,

antibiotics targeting the same site within the ribosome as that of putrescine will show similar

macrolide synergy as that of putrescine-azithromycin combination. Additionally, we hypothesize

that a compound that mirrors the phenotype of putrescine-azithromycin will show additive

effects when in combination with putrescine. We tested eight antibiotics, including azithromycin,

in combination with one another and with putrescine to compare their FICI values (Fig 5A).

These antibiotics targeted various sites within the ribosome: Paromomycin and gentamicin bind

to the aminoacyl site within the 30S ribosomal subunit leading to misreading of genetic code.

Chloramphenicol binds to the aminoacyl site of the 50S ribosomal subunit and prevents peptidyl

transferase. Azithromycin and erythromycin both block the 50S ribosomal subunit nascent

peptide tunnel inhibiting transpeptidation. Tetracycline and doxycycline target the 30S ribosomal

acceptor site preventing the association of aminoacyl-tRNA with the ribosome complex and

prevent mRNA binding [69]. We used rifampicin, an RNA polymerase inhibitor, as a control. Out

of the seven antibiotics tested, only tetracyclines (tetracycline and doxycycline) synergized with

azithromycin as putrescine did (Fig 5A). Synergy between the tetracycline doxycycline and

azithromycin has been observed before in *P. aeruginosa*, albeit the underlying mechanism has

not been resolved [70]. Checkerboard assays of putrescine-azithromycin combinations were

phenocopied by tetracycline-azithromycin combinations in the *K. pneumoniae* wild type strain

(Fig 5B-C) and tetracycline also showed additive effects with putrescine (Fig 5D). Similarly, the

additive effects of the putrescine-azithromycin combination against the ribosomal mutants

rpsO::Tn and *rpsR*::Tn was also observed in the tetracycline-azithromycin combination (Fig 5E-

H). Additionally, the combination of tetracycline and azithromycin, in checkerboards of PMBN

treated *K. pneumoniae* (FICI 0.50 ± 0.00) and the *waaD*::Tn mutant still exhibited significant

synergy (FICI 0.41 ± 0.07) (Supplementary Fig 8A-C); under these conditions, while the outer

membrane is perturbed chemically or genetically, synergy may still occur at the ribosomal level.

Previously, *E. coli* serially passaged in sub-lethal putrescine concentrations evolved

putrescine-resistant mutants that had truncations of the S7 30S ribosomal protein, a protein that

organizes the folding of the 3' major domain of 16S rRNA and crosslinks to the anticodon loop

of tRNAs [71, 72]. These mutants exhibited increased growth rates relative to the wild type when

grown in the presence of putrescine [72]. Indeed, adaptive resistance is likely to occur at the target

for a compound, supporting our findings that putrescine may act on the 30S ribosomal subunit.

Polyamine cross-linking in the 30S subunit leads to 16S rRNA structural changes [66]; similarly,

tetracycline has been shown to alter the structure of 16S rRNA [73]. Moreover, 30S subunits

lacking RpsO exhibit subtle changes in 16S rRNA nucleotide reactivity [50, 74]. Thus, it is probable

that the mutants of RpsO and RpsR lead to an altered 16S rRNA structure similar to that

occurring as a result of polyamine and tetracycline binding to the 30S subunit, which may

explain the lack of synergy of these drugs with azithromycin in *rpsO*::Tn and *rpsR*::Tn mutant

backgrounds. Together, our results provide support that putrescine acts on the 30S ribosomal

subunit similar to tetracycline and that it mediates the macrolide synergy via a dual mechanism

(at the membrane and ribosomal levels).

Figure 5. **Putrescine may inhibit the ribosomal 30S subunit in a similar manner as tetracyclines.** A) Heat map
showing the average FICI value of duplicate checkerboard assays against *K. pneumoniae* MKP103 with various
drug combinations. B-D) Checkerboard assays against *K. pneumoniae* (MKP103) treated with B) putrescine-
azithromycin, C) tetracycline-azithromycin, and D) putrescine-tetracycline combinations. E-F) Checkerboard assays
against *K. pneumoniae rpsO*::Tn treated with E) putrescine-azithromycin and F) tetracycline-azithromycin. G-H)
Checkerboard assays against *K. pneumoniae rpsR*::Tn treated with G) putrescine-azithromycin and H) tetracycline-
azithromycin. Dark blue regions represent high cell density determined by OD_{600}. FICI mean and SEM, derived
from 3 independent experiments, are displayed on each representative checkerboard.

A screen of a polyamine analog library uncovers azithromycin synergists.

We sought to exploit the polyamine-azithromycin synergy and the understanding of its mode of action to uncover an antibiotic adjuvant that may expand the spectrum of azithromycin to *K. pneumoniae*. We hypothesized that a synthetic polyamine analog may lead to synergy with azithromycin similar to that mediated by natural polyamines, serving as a potential macrolide adjuvant. To this end, we screened a library of 86 polyamine analogs against MKP103 alone and in combination with a sub-inhibitory concentration (1/8th MIC) of azithromycin (Fig 6A). Six compounds displayed growth inhibition alone and in combination with azithromycin. Follow up checkerboard assays of the six primary hits revealed that OES2-017, OES2-046, OES2-045, and OES2-008 synergized with azithromycin at concentrations starting at the low micromolar range with FICI values of 0.54 ± 0.11, 0.32 ± 0.05, 0.37 ± 0.03, and 0.29 ± 0.05, respectively (Fig 6B-E). Similar to putrescine, the identified polyamine analogs showed growth inhibitory effects against *K. pneumoniae* MKP103; their MIC values (31.25 μM, 2.5, 2.5, and 5 mM for OES2-017, -045, -046, and -008, respectively, Fig 6B-E) were substantially lower than that of putrescine (40 mM).

Next, we sought to identify the mechanism of azithromycin synergy by these four compounds, first by checking if they acted through a similar mechanism as that of putrescine. Therefore, we conducted checkerboard assays of each compound with azithromycin against the *rpsO*::Tn, *rpsR*::Tn, and *waaD*::Tn *K. pneumoniae* mutants, against which putrescine lost its synergy with the macrolide antibiotic, and tested their outer and inner membrane permeabilization effects. Both OES2-017 and OES2-008 showed reduced or complete loss of azithromycin synergy against all three mutants (Supplementary Fig 9A-F). Conversely, OES2-045 and OES2-046 exhibited no loss of synergy against the mutants, except that the extent of the

synergy against *waaD*::Tn (mean FICI= 0.419 and 0.489, respectively) was reduced relative to

that against the wild type (mean FICI= 0.325 and 0.375, respectively) (Supplementary Fig 9G-

L). We showed in NPN fluorometric kinetic assays that OES2-017, OES2-008, OES2-045, and

OES2-046, in descending order of magnitude, disrupted the outer membrane of *K. pneumoniae*

in a dose dependant manner across the synergistic range of each compound (Supplementary Fig

10A-D). Further, all four analogs exhibited inner membrane activity, in colorimetric ONPG

hydrolysis assays against *E. coli* ML-35, in a dose dependant manner across the synergistic range

of each compound (Supplementary Fig 11A-D). Taken together, these results suggest that OES2-

017 and OES2-008 synergize with azithromycin via mechanisms similar to those of putrescine at

the ribosome and outer and inner membranes levels, whereas OES2-045 and OES2-046 may

exert their azithromycin synergy through membrane activity only. Importantly, these analogs

synergize with azithromycin at substantially lower concentrations than putrescine, serving as

potential macrolide adjuvants.

Figure 6. **Screening for azithromycin adjuvants identifies polyamine analogs with more potent azithromycin synergy and growth inhibitory activities.** A) A replica plot displaying the OD readings of *K. pneumoniae* MKP103 in the presence of the polyamine analogs at 2.5 mM alone (x-axis) and in combination with 64 μg/mL azithromycin (y-axis). B-E) Representative checkerboard assays of *K. pneumoniae* MKP103 treated with the prioritized polyamine analog hits in combination with azithromycin. All FICI values are shown as the average of at least 3 independent repeats and the SEM. Dark blue regions represent high cell density determined by OD_{600}.

Conclusions

In this study, we uncovered a sensitization of *K. pneumoniae* to the macrolide azithromycin in the presence of potentially physiological levels of polyamines. We showed that the putrescine-azithromycin synergy occurs in pan-drug-resistant and hypervirulent *K. pneumoniae* lineages, bringing azithromycin susceptibility in 78 of 100 clinical isolates below the CLSI breakpoint. Further, we have provided evidence towards a dual mechanism of action underlying the putrescine-azithromycin synergy, whereby the polyamine: 1. permeabilized the outer and inner membranes to azithromycin and 2. compounded the protein synthesis inhibiting effects of azithromycin by acting at the ribosome, likely at the 30S subunit.

Further, we discovered four synthetic polyamine analogs that displayed growth inhibitory

effects and synergy with azithromycin against *K. pneumoniae*, with greater potency compared to

natural polyamines. Synergy with currently available Gram-positive specific antibiotics, such as

macrolides, including by perturbing the Gram-negative outer membrane, represents a potential

strategy to combat drug-resistant Gram-negative bacteria [57]. As such, the polyamine analogs

discovered herein may broaden the spectrum of azithromycin, serving as a potential strategy to

combat antimicrobial resistant *K. pneumoniae*.

Methods

Strains, reagents, and culture conditions. Table 1 lists bacteria and plasmids used in this work.

Bacteria were grown in cation-adjusted Mueller Hinton medium (CAMHB, herein referred to as

MHB) at 37 °C buffered with 100 mM Tris-HCl pH7.4 buffer (to offset the alkalinity of higher

polyamine concentrations), unless otherwise indicated. Antibiotics and reagents were obtained

from Sigma unless otherwise indicated.

Table 1. Bacterial strains and plasmids used in this study.

Bacterial Strain	Description	Reference
K. pneumoniae		
MKP103	A derivative of the clinical outbreak isolate KPNIH1 lacking its carbapenemase-encoding gene	34
MKP103 Library	An arrayed sequence-defined transposon mutant library covering the non-essential genome of *K. pneumoniae* MKP103	34
K. pneumoniae Diversity Panel	A *K. pneumoniae* diversity panel, comprising 100 clinical isolates – Received from BEI resources	38
MKP103 pBR322	MKP103 transformed with pBR322	This study
MKP103 *waaD*::Tn pBR322	MKP103 *waaD*::Tn transformed with pBR322	This study

E. coli		
M6394	Wild-type GC4468 Δ(argF-lac)169 λ− IN(rrnD-rrnE)1 rpsL179 (Strr)	58
M6394 Δ9	M6394 Δ*acrB* Δ*acrD* Δ*acrEF*::spc Δ*emrB* Δ*emrY* Δ*entS*::cam Δ*macB* Δ*mdtC* Δ*mdtF*	58
ML-35 pBR322	lacZ$^+$Y$^-$I$^-$, *E. coli* with constitutive expression of β-galactosidase but lacking the lactose permease ML-35 with periplasmic β-lactamase	75
Bacterial Plasmids		
pBR322	Plasmid encoding a periplasmic β-lactamase	76
PT7-deGFP-Mgapt	pIVEX2.3d with a transcription unit of PT7:deGFP-Mgapt:T7term	60

Antimicrobial susceptibility testing. All MIC and checkerboard assays were conducted

following the CLSI method for MIC testing by broth microdilution [77]. The assays were

conducted in polystyrene 96-well plates with final volume of 100 μL/well and inoculum size

adjusted as optical density at 600 nm (OD_{600}) of 0.001. Plates were incubated while shaking at

600 RPM and 37°C; growth of the bacterial culture was determined by OD_{600} measurement at 24

hours. To assess the combined antimicrobial activity, fractional inhibitory concentration indices

(FICI) were calculated as FICI = $A/MIC_A + B/MIC_B$. A and B represent the MIC of the drugs

while in combination. MIC_A and MIC_B represent the individual MIC of each drug alone. FICI

values less than or equal to 0.5 were considered synergism, values greater than 4 were

considered antagonism, and values between 0.5 and 4 were indicated indifference, additive, or no

interaction.

Arrayed transposon *Klebsiella pneumoniae* MKP103 library screen. Overnight cultures of

the *K. pneumoniae* MKP103 library[34] were performed using the BioMatrix benchtop robot (S&P

robotics) into 384 well-density plates containing MHB with 150 μg/mL of chloramphenicol and

incubated at 37°C for 24 hours. Subsequently, the entire library was inoculated into MHB with

azithromycin at 32 μg/mL and 64 μg/mL and without azithromycin. The plates were incubated at

37°C, and OD_{600} was read after 24 hours.

NPN assays. The outer membrane permeability of *K. pneumoniae* MKP103 and *E. coli* M6394

wild type and mutants was measured using NPN uptake assays as previously described [78].

Briefly, bacteria were subcultured in MHB until mid to late log phase, washed, and resuspended

in either 100 mM Tris-HCl (pH 7.4) with 20 mM glucose or 5 mM HEPES (pH7.4) with 20 mM

glucose. The latter buffer was used only when low putrescine concentrations, causing minimal

pH changes, were tested. NPN was added to the bacterial suspension at a final concentration of

20 μM and dispensed into black 96 well plates pre-seeded with a concentration gradient of

putrescine to a total final volume of 200 μL. We tested untreated bacteria suspended in the same

buffer adjusted to the final pH reached upon addition of each tested putrescine concentration as a

negative control (untreated pH control) used to correct for any pH related signal. Fluorescence

was read kinetically (excitation λ=355 nm, emission λ=420 nm) every 5 minutes for 215

minutes. The %NPN uptake was calculated using the following equation: %NPN uptake = (F_{obs}-

F_0)/(F_{100}-F_0) x 100, where F_{obs} is the observed fluorescence at a given concentration of

putrescine, F_0 is the fluorescence in non-treated cells (untreated pH control), and F_{100} is the

fluorescence of NPN in 8 μg/mL colistin-treated cells (used as a positive control).

β-lactamase assay. The outer membrane integrity of *K. pneumoniae* wild type and *waaD*::Tn

mutant were tested by measuring β-lactamase activity with nitrocefin hydrolysis as previously

described [79]. First, we extracted pBR322 from an *E. coli* strain carrying the plasmid (Monarch

Mini-Prep kit, NEB) and transformed it into electrocompetent *K. pneumoniae* strains of interest

by electroporation[34] Overnight cultures of *K. pneumoniae* pBR322 and *K. pneumoniae waaD*::Tn

pBR322 were diluted 100-fold into TSB and incubated at 37°C until late log phase (OD_{600} of

~1.8). Aliquots of the sub-cultures were centrifuged and washed twice with PBS then

resuspended in 100 mM Tris-HCl buffer (pH 7.4) at a final OD_{600} of 0.01. The bacterial

suspensions mixed with 30 μM nitrocefin were dispensed into 96 well plates pre-seeded with the

tested drugs with a total final volume of 100 μL/well. The plates were incubated at 37°C while

measuring the absorbance kinetically at 492 nm every 10 minutes.

β-galactosidase assay. The inner membrane integrity of *E. coli* ML-35 pBR322 was tested by

measuring β-galactosidase activity with ONPG hydrolysis as previously described [79]. An

overnight culture of *E. coli* ML-35 pBR322 was diluted 100-fold into TSB and incubated at

37°C until late log phase (OD_{600} of ~1.8). Aliquots of the sub-culture were centrifuged and

washed twice with PBS then resuspended in 100 mM Tris-HCl buffer (pH 7.4) at a final OD_{600}

of 0.01. The bacterial suspensions mixed with 1.5 mM ONPG were dispensed into 96 well plates

pre-seeded with the tested drugs with a total final volume of 100 μL/well. The plates were

incubated at 37°C while measuring the absorbance kinetically at 405 nm every 2 minutes.

Cell free protein expression. An NEB Express Cell-free *E. coli* protein synthesis system (New

England Biolabs) was used for *in vitro* protein expression. Transcription and translation were

measured fluorometrically (excitation/emission wavelengths at 610/650 nm MGapt, 485/525 nm

deGFP) using an engineered plasmid that encodes a deGFP protein and a malachite green RNA

aptamer (PT7-deGFP-MGapt was a gift from Richard Murray, via Addgene plasmid # 67741;

http://n2t.net/addgene:67741) [60]. The cell-free protein synthesis system reaction mixture was

prepared following the manufacturer's protocol and dispensed as 5 μL aliquots into black 384

well plates. Putrescine, dH_2O, plasmid (124 ng/μL final concentration), and Malachite green dye

(10 μM final concentration) top up the reaction mixture reaching a combined final volume of 11

μL.

Polyamine analog library screen. *K. pneumoniae* MKP103 was inoculated in MHB with 100

mM Tris-HCl buffer (pH 7.4) at a 0.001 OD_{600} and dispensed into 96 well plates at a total final

volume of 100 μL (including polyamine analogs, azithromycin, vehicle, or combinations

thereof). Polyamine analogs were screened at 2.5 mM alone and in combination with 64 μg/mL

azithromycin. Plates were incubated at 37°C and 600 RPM; growth of the bacterial culture was

determined by OD_{600} readings at 24 hours.

Statistical analysis. All statistical analyses were calculated using GraphPad Prism 9.

Acknowledgements.

This work was funded by a Natural Sciences and Engineering Research Council of Canada

discovery grant, a Saskatchewan Health Research Foundation establishment grant, and startup

funds from the Faculty of Science at the University of Regina. O.M.E. holds a Canada Research

Chair in Chemogenomics and Antimicrobial Research. P.B.M. was supported by a CIHR CGS-

M award.

References

(1) Antimicrobial Resistance, C. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. *Lancet* **2022**, *399* (10325), 629-655. DOI: 10.1016/S0140-6736(21)02724-0.

(2) Nikaido, H. Molecular basis of bacterial outer membrane permeability revisited. *Microbiol Mol Biol Rev* **2003**, *67* (4), 593-656. DOI: 10.1128/MMBR.67.4.593-656.2003.

(3) Ak, O.; Batirel, A.; Ozer, S.; Çolakoğlu, S. Nosocomial infections and risk factors in the intensive care unit of a teaching and research hospital: a prospective cohort study. *Med Sci Monit* **2011**, *17* (5), Ph29-34. DOI: 10.12659/msm.881750.

(4) WHO. Global Priority List of Antibiotic-Resistant Bacteria to Guide Research, Discovery, and Development of New Antibiotics. world health organization: 2017.

(5) Delcour, A. H. Outer membrane permeability and antibiotic resistance. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics* **2009**, *1794* (5), 808-816. DOI: 10.1016/j.bbapap.2008.11.005.

(6) WHO. *Antibacterial agents in clinical development: an analysis of the antibacterial clinical development pipeline*; 2019.

(7) El-Halfawy, O. M.; Valvano, M. A. Chemical Communication of Antibiotic Resistance by a Highly Resistant Subpopulation of Bacterial Cells. *Plos One* **2013**, *8* (7). DOI: 10.1371/journal.pone.0068874.

(8) Perry, E. K.; Meirelles, L. A.; Newman, D. K. From the soil to the clinic: the impact of microbial secondary metabolites on antibiotic tolerance and resistance. *Nat Rev Microbiol* **2022**, *20* (3), 129-142. DOI: 10.1038/s41579-021-00620-w.

(9) Zhang, M.; Wang, H.; Tracey, K. J. Regulation of macrophage activation and inflammation by spermine: a new chapter in an old story. *Crit Care Med* **2000**, *28* (4 Suppl), N60-66.

(10) Agostinelli, E.; Arancia, G.; Vedova, L. D.; Belli, F.; Marra, M.; Salvi, M.; Toninello, A. The biological functions of polyamine oxidation products by amine oxidases: perspectives of clinical applications. *Amino Acids* **2004**, *27* (3-4), 347-358. DOI: 10.1007/s00726-004-0114-4.

(11) Bjelakovic, G.; Stojanovic, I.; Jevtovic Stoimenov, T.; Pavlovic, D.; Kocic, G.; Rossi, S.; Tabolacci, C.; Nikolic, J.; Sokolovic, D.; Bjelakovic, L. Metabolic correlations of glucocorticoids and polyamines in inflammation and apoptosis. *Amino Acids* **2010**, *39* (1), 29-43.

(12) Zhang, M.; Wang, H.; Tracey, K. J. Regulation of macrophage activation and inflammation by spermine: A new chapter in an old story. *Critical Care Medicine* **2000**, *28* (4), N60-N66.

(13) Liao, C. P.; Lasbury, M. E.; Wang, S. H.; Zhang, C.; Durant, P. J.; Tschang, D.; Lee, C. H. Inflammatory cells are sources of polyamines that induce alveolar macrophage to undergo apoptosis during Pneumocystis pneumonia. *J Eukaryot Microbiol* **2006**, *53 Suppl 1*, S134-135. DOI: 10.1111/j.1550-7408.2006.00203.x.

(14) Gugliucci, A. Polyamines as clinical laboratory tools. *Clin Chim Acta* **2004**, *344* (1-2), 23-35. DOI: 10.1016/j.cccn.2004.02.022.

(15) Larque, E.; Sabater-Molina, M.; Zamora, S. Biological significance of dietary polyamines. *Nutrition* **2007**, *23* (1), 87-95.

(16) Feuerstein, B. G.; Pattabiraman, N.; Marton, L. J. Spermine-DNA interactions: a theoretical study. *Proc Natl Acad Sci U S A* **1986**, *83* (16), 5948-5952. DOI: 10.1073/pnas.83.16.5948.

(17) Feuerstein, B. G.; Pattabiraman, N.; Marton, L. J. Molecular dynamics of spermine-DNA interactions: sequence specificity and DNA bending for a simple ligand. *Nucleic Acids Res* **1989**, *17* (17), 6883-6892. DOI: 10.1093/nar/17.17.6883.

(18) Feuerstein, B. G.; Pattabiraman, N.; Marton, L. J. Molecular mechanics of the interactions of spermine with DNA: DNA bending as a result of ligand binding. *Nucleic Acids Res* **1990**, *18* (5), 1271-1282. DOI: 10.1093/nar/18.5.1271.

(19) Davis, R. H.; Morris, D. R.; Coffino, P. Sequestered end products and enzyme regulation: the case of ornithine decarboxylase. *Microbiol Rev* **1992**, *56* (2), 280-290. DOI: 10.1128/mr.56.2.280-290.1992.

(20) Miyamoto, S.; Kashiwagi, K.; Ito, K.; Watanabe, S.; Igarashi, K. Estimation of polyamine distribution and polyamine stimulation of protein synthesis in *Escherichia coli*. *Arch Biochem Biophys* **1993**, *300* (1), 63-68. DOI: 10.1006/abbi.1993.1009.

(21) Wortham, B. W.; Oliveira, M. A.; Patel, C. N. Polyamines in Bacteria: Pleiotropic Effects yet Specific Mechanisms. Springer New York, 2007; pp 106-115.

(22) Hasan, C. M.; Pottenger, S.; Green, A. E.; Cox, A. A.; White, J. S.; Jones, T.; Winstanley, C.; Kadioglu, A.; Wright, M. H.; Neill, D. R.; Fothergill, J. L. *Pseudomonas aeruginosa* utilizes the host-derived polyamine spermidine to facilitate antimicrobial tolerance. *JCI Insight* **2022**, *7* (22). DOI: 10.1172/jci.insight.158879.

(23) El-Halfawy, O. M.; Valvano, M. A. Chemical communication of antibiotic resistance by a highly resistant subpopulation of bacterial cells. *PLOS One* **2013**, *8* (7), e68874. DOI: 10.1371/journal.pone.0068874.

(24) El-Halfawy, O. M.; Valvano, M. A. Putrescine reduces antibiotic-induced oxidative stress as a mechanism of modulation of antibiotic resistance in *Burkholderia cenocepacia*. *Antimicrob Agents Chemother* **2014**, *58* (7), 4162-4171. DOI: AAC.02649-14.

(25) Kwon, D. H.; Lu, C. D. Polyamines induce resistance to cationic peptide, aminoglycoside, and quinolone antibiotics in *Pseudomonas aeruginosa* PAO1. *Antimicrob Agents Chemother* **2006**, *50* (5), 1615-1622.

(26) Kwon, D. H.; Lu, C. D. Polyamine effects on antibiotic susceptibility in bacteria. *Antimicrob Agents Chemother* **2007**, *51* (6), 2070-2077.

(27) El-Halfawy, O. M.; Valvano, M. A. Non-genetic mechanisms communicating antibiotic resistance: Rethinking strategies for antimicrobial drug design. *Expert Opinion On Drug Discovery* **2012**, *7* (10), 923-933.

(28) Vázquez-Laslop, N.; Mankin, A. S. How Macrolide Antibiotics Work. *Trends Biochem Sci* **2018**, *43* (9), 668-684. DOI: 10.1016/j.tibs.2018.06.011.

(29) Lalak, N. J.; Morris, D. L. Azithromycin clinical pharmacokinetics. *Clin Pharmacokinet* **1993**, *25* (5), 370-374. DOI: 10.2165/00003088-199325050-00003.

(30) Kohno, Y. Chapter 6 - Pharmacokinetics and Metabolism of Macrolides. In *Macrolide Antibiotics (Second Edition)*, Ōmura, S. Ed.; Academic Press, 2003; pp 327-361.

(31) Magill, S. S.; O'Leary, E.; Ray, S. M.; Kainer, M. A.; Evans, C.; Bamberg, W. M.; Johnston, H.; Janelle, S. J.; Oyewumi, T.; Lynfield, R.; et al. Antimicrobial Use in US Hospitals: Comparison of Results From Emerging Infections Program Prevalence Surveys, 2015 and 2011. *Clin Infect Dis* **2021**, *72* (10), 1784-1792. DOI: 10.1093/cid/ciaa373.

(32) Wilson, D. N. The A–Z of bacterial translation inhibitors. *Critical Reviews in Biochemistry and Molecular Biology* **2009**, *44* (6), 393-433. DOI: 10.3109/10409230903307311.

(33) Farha, M. A.; French, S.; Stokes, J. M.; Brown, E. D. Bicarbonate Alters Bacterial Susceptibility to Antibiotics by Targeting the Proton Motive Force. *ACS Infectious Diseases* **2018**, *4* (3), 382-390. DOI: 10.1021/acsinfecdis.7b00194.

(34) Ramage, B.; Erolin, R.; Held, K.; Gasper, J.; Weiss, E.; Brittnacher, M.; Gallagher, L.; Manoil, C. Comprehensive Arrayed Transposon Mutant Library of Klebsiella pneumoniae Outbreak Strain KPNIH1. *J Bacteriol* **2017**, *199* (20). DOI: 10.1128/JB.00352-17 From NLM PubMed-not-MEDLINE.

(35) CLSI. Performance Standards for Antimicrobial Susceptibility Testing. *CLSI supplement M100* **2020**, *30th ed.* .

(36) Thurlow, L. R.; Joshi, G. S.; Clark, J. R.; Spontak, J. S.; Neely, C. J.; Maile, R.; Richardson, A. R. Functional modularity of the arginine catabolic mobile element contributes to the success of USA300 methicillin-resistant *Staphylococcus aureus*. *Cell Host Microbe* **2013**, *13* (1), 100-107. DOI: 10.1016/j.chom.2012.11.012.

(37) Sagar, N. A.; Tarafdar, S.; Agarwal, S.; Tarafdar, A.; Sharma, S. Polyamines: Functions, Metabolism,
and Role in Human Disease Management. *Med Sci (Basel)* **2021**, *9* (2). DOI: 10.3390/medsci9020044.

(38) Martin, M. J.; Stribling, W.; Ong, A. C.; Maybank, R.; Kwak, Y. I.; Rosado-Mendez, J. A.; Preston, L. N.;
Lane, K. F.; Julius, M.; Jones, A. R.; et al. A panel of diverse *Klebsiella pneumoniae* clinical isolates for
research and development. *Microb Genom* **2023**, *9* (5). DOI: 10.1099/mgen.0.000967.

(39) Liu, A.; Tran, L.; Becket, E.; Lee, K.; Chinn, L.; Park, E.; Tran, K.; Miller, J. H. Antibiotic sensitivity
profiles determined with an *Escherichia coli* gene knockout collection: generating an antibiotic bar code.
Antimicrob Agents Chemother **2010**, *54* (4), 1393-1403. DOI: 10.1128/AAC.00906-09.

(40) Izquierdo, L.; Coderch, N.; Pique, N.; Bedini, E.; Corsaro, M. M.; Merino, S.; Fresno, S.; Tomas, J. M.;
Regue, M. The *Klebsiella pneumoniae wabG* gene: role in biosynthesis of the core lipopolysaccharide and
virulence. *J Bacteriol* **2003**, *185* (24), 7213-7221. DOI: 10.1128/JB.185.24.7213-7221.2003.

(41) Furniss, R. C. D.; Kaderabkova, N.; Barker, D.; Bernal, P.; Maslova, E.; Antwi, A. A. A.; McNeil, H. E.;
Pugh, H. L.; Dortet, L.; Blair, J. M. A.; et al. Breaking antimicrobial resistance by disrupting
extracytoplasmic protein folding. *Elife* **2022**, *11*. DOI: 10.7554/eLife.57974.

(42) Wang, J.; Ma, W.; Wang, Z.; Li, Y.; Wang, X. Construction and Characterization of an *Escherichia coli*
Mutant Producing Kdo2-Lipid A. *Marine Drugs* **2014**, *12* (3), 1495-1511. DOI: 10.3390/md12031495.

(43) Kneidinger, B.; Marolda, C.; Graninger, M.; Zamyatina, A.; McArthur, F.; Kosma, P.; Valvano, M. A.;
Messner, P. Biosynthesis Pathway of ADP-L-glycero-beta-D-manno-heptose in *Escherichia coli*. *Journal of*
Bacteriology **2002**, *184* (2), 363-369. DOI: 10.1128/jb.184.2.363-369.2002.

(44) Regué, M.; Climent, N. R.; Abitiu, N.; Coderch, N. R.; Merino, S.; Izquierdo, L.; Altarriba, M.; TomáS,
J. M. Genetic Characterization of the *Klebsiella pneumoniae waa* Gene Cluster, Involved in Core
Lipopolysaccharide Biosynthesis. *Journal of Bacteriology* **2001**, *183* (12), 3564-3573. DOI:
10.1128/jb.183.12.3564-3573.2001.

(45) Yethon, J. A.; Heinrichs, D. E.; Monteiro, M. A.; Perry, M. B.; Whitfield, C. Involvement of *waaY*,
waaQ, and *waaP* in the modification of *Escherichia coli* lipopolysaccharide and their role in the
formation of a stable outer membrane. *J Biol Chem* **1998**, *273* (41), 26310-26316. DOI:
10.1074/jbc.273.41.26310.

(46) Vaara, M. Agents that increase the permeability of the outer membrane. *Microbiological Reviews*
1992, *56* (3), 395-411. DOI: doi:10.1128/mr.56.3.395-411.1992.

(47) Meehan, B. M.; Landeta, C.; Boyd, D.; Beckwith, J. The Disulfide Bond Formation Pathway Is
Essential for Anaerobic Growth of *Escherichia coli*. *J Bacteriol* **2017**, *199* (16). DOI: 10.1128/jb.00120-17.

(48) Onufryk, C.; Crouch, M. L.; Fang, F. C.; Gross, C. A. Characterization of six lipoproteins in the sigmaE
regulon. *J Bacteriol* **2005**, *187* (13), 4552-4561. DOI: 10.1128/jb.187.13.4552-4561.2005.

(49) Sykes, M. T.; Williamson, J. R. A complex assembly landscape for the 30S ribosomal subunit. *Annu*
Rev Biophys **2009**, *38*, 197-215. DOI: 10.1146/annurev.biophys.050708.133615.

(50) Bubunenko, M.; Korepanov, A.; Court, D. L.; Jagannathan, I.; Dickinson, D.; Chaudhuri, B. R.; Garber,
M. B.; Culver, G. M. 30S ribosomal subunits can be assembled in vivo without primary binding ribosomal
protein S15. *RNA* **2006**, *12* (7), 1229-1239. DOI: 10.1261/rna.2262106.

(51) Ginzburg, I.; Miskin, R.; Zamir, A. N-ethyl maleimide as a probe for the study of functional sites and
conformations of 30 S ribosomal subunits. *Journal of Molecular Biology* **1973**, *79* (3), 481-494. DOI:
https://doi.org/10.1016/0022-2836(73)90400-2.

(52) Delvillani, F.; Papiani, G.; Deho, G.; Briani, F. S1 ribosomal protein and the interplay between
translation and mRNA decay. *Nucleic Acids Res* **2011**, *39* (17), 7702-7715. DOI: 10.1093/nar/gkr417.

(53) Soh, S. M.; Jang, H.; Mitchell, R. J. Loss of the lipopolysaccharide (LPS) inner core increases the
electrocompetence of *Escherichia coli*. *Appl Microbiol Biotechnol* **2020**, *104* (17), 7427-7435. DOI:
10.1007/s00253-020-10779-6.

(54) Sporing, I.; Felgner, S.; Preusse, M.; Eckweiler, D.; Rohde, M.; Haussler, S.; Weiss, S.; Erhardt, M.
Regulation of Flagellum Biosynthesis in Response to Cell Envelope Stress in *Salmonella enterica Serovar*
Typhimurium. *mBio* **2018**, *9* (3). DOI: 10.1128/mBio.00736-17.
(55) Frirdich, E.; Bouwman, C.; Vinogradov, E.; Whitfield, C. The Role of Galacturonic Acid in Outer
Membrane Stability in *Klebsiella pneumoniae*. *Journal of Biological Chemistry* **2005**, *280* (30), 27604-
27612. DOI: 10.1074/jbc.m504987200.
(56) Al-Marzooq, F.; Ghazawi, A.; Tariq, S.; Daoud, L.; Collyns, T. Discerning the role of polymyxin B
nonapeptide in restoring the antibacterial activity of azithromycin against antibiotic-resistant *Escherichia*
coli. *Front Microbiol* **2022**, *13*, 998671. DOI: 10.3389/fmicb.2022.998671.
(57) Wesseling, C. M. J.; Martin, N. I. Synergy by Perturbing the Gram-Negative Outer Membrane:
Opening the Door for Gram-Positive Specific Antibiotics. *ACS Infectious Diseases* **2022**, *8* (9), 1731-1757.
DOI: 10.1021/acsinfecdis.2c00193.
(58) Rosner, J. L.; Martin, R. G. Reduction of cellular stress by TolC-dependent efflux pumps in
Escherichia coli indicated by BaeSR and CpxARP activation of spy in efflux mutants. *J Bacteriol* **2013**, *195*
(5), 1042-1050. DOI: 10.1128/JB.01996-12.
(59) Li, J.; Beuerman, R.; Verma, C. S. Mechanism of polyamine induced colistin resistance through
electrostatic networks on bacterial outer membranes. *Biochim Biophys Acta Biomembr* **2020**, *1862* (9),
183297. DOI: 10.1016/j.bbamem.2020.183297.
(60) Siegal-Gaskins, D.; Tuza, Z. A.; Kim, J.; Noireaux, V.; Murray, R. M. Gene circuit performance
characterization and resource usage in a cell-free "breadboard". *ACS Synth Biol* **2014**, *3* (6), 416-425.
DOI: 10.1021/sb400203p.
(61) Kanemura, A.; Yoshikawa, Y.; Fukuda, W.; Tsumoto, K.; Kenmotsu, T.; Yoshikawa, K. Opposite effect
of polyamines on In vitro gene expression: Enhancement at low concentrations but inhibition at high
concentrations. *PLoS One* **2018**, *13* (3), e0193595. DOI: 10.1371/journal.pone.0193595.
(62) Igarashi, K.; Kashiwagi, K. Effects of polyamines on protein synthesis and growth of *Escherichia coli*.
Journal of Biological Chemistry **2018**, *293* (48), 18702-18709. DOI: 10.1074/jbc.tm118.003465.
(63) Agrawal, R. K.; Penczek, P.; Grassucci, R. A.; Burkhardt, N.; Nierhaus, K. H.; Frank, J. Effect of buffer
conditions on the position of tRNA on the 70 S ribosome as visualized by cryoelectron microscopy. *J Biol*
Chem **1999**, *274* (13), 8723-8729. DOI: 10.1074/jbc.274.13.8723.
(64) Winther, K. S.; Sorensen, M. A.; Svenningsen, S. L. Polyamines are Required for tRNA Anticodon
Modification in *Escherichia coli*. *J Mol Biol* **2021**, *433* (15), 167073. DOI: 10.1016/j.jmb.2021.167073.
(65) Dever, T. E.; Ivanov, I. P. Roles of polyamines in translation. *Journal of Biological Chemistry* **2018**,
293 (48), 18719-18729. DOI: 10.1074/jbc.tm118.003338.
(66) Amarantos, I.; Zarkadis, I. K.; Kalpaxis, D. L. The identification of spermine binding sites in 16S rRNA
allows interpretation of the spermine effect on ribosomal 30S subunit functions. *Nucleic Acids Res* **2002**,
30 (13), 2832-2843. DOI: 10.1093/nar/gkf404.
(67) Kalpaxis, D. L.; Drainas, D. Inhibitory effect of spermine on ribosomal peptidyltransferase. *Arch*
Biochem Biophys **1993**, *300* (2), 629-634. DOI: 10.1006/abbi.1993.1087.
(68) Teraoka, H.; Tanaka, K. Effect of Spermine on the Binding of Erythromycin to *Escherichia coli*
Ribosomes and the Peptidyl-Transfer Reaction. *European Journal of Biochemistry* **1973**, *33* (3), 578-583.
DOI: 10.1111/j.1432-1033.1973.tb02718.x.
(69) Chopra, I.; Roberts, M. Tetracycline Antibiotics: Mode of Action, Applications, Molecular Biology,
and Epidemiology of Bacterial Resistance. *Microbiology and Molecular Biology Reviews* **2001**, *65* (2),
232-260. DOI: 10.1128/mmbr.65.2.232-260.2001.
(70) Saiman, L.; Chen, Y.; Gabriel, P. S.; Knirsch, C. Synergistic activities of macrolide antibiotics against
Pseudomonas aeruginosa, Burkholderia cepacia, Stenotrophomonas maltophilia, and *Alcaligenes*
xylosoxidans isolated from patients with cystic fibrosis. *Antimicrob Agents Chemother* **2002**, *46* (4),
1105-1107.

(71) Wimberly, B. T.; White, S. W.; Ramakrishnan, V. The structure of ribosomal protein S7 at 1.9 A
resolution reveals a beta-hairpin motif that binds double-stranded nucleic acids. *Structure* **1997**, *5* (9),
1187-1198. DOI: 10.1016/s0969-2126(97)00269-4.
(72) Lennen, R. M.; Lim, H. G.; Jensen, K.; Mohammed, E. T.; Phaneuf, P. V.; Noh, M. H.; Malla, S.;
Borner, R. A.; Chekina, K.; Ozdemir, E.; et al. Laboratory evolution reveals general and specific tolerance
mechanisms for commodity chemicals. *Metab Eng* **2023**, *76*, 179-192. DOI:
10.1016/j.ymben.2023.01.012.
(73) Noah, J. W.; Dolan, M. A.; Babin, P.; Wollenzien, P. Effects of Tetracycline and Spectinomycin on the
Tertiary Structure of Ribosomal RNA in the *Escherichia coli* 30 S Ribosomal Subunit. *Journal of Biological*
Chemistry **1999**, *274* (23), 16576-16581. DOI: 10.1074/jbc.274.23.16576.
(74) Svensson, P.; Changchien, L. M.; Craven, G. R.; Noller, H. F. Interaction of ribosomal proteins, S6, S8,
S15 and S18 with the central domain of 16 S ribosomal RNA. *J Mol Biol* **1988**, *200* (2), 301-308. DOI:
10.1016/0022-2836(88)90242-2.
(75) Buttin, G.; Cohen, G. N.; Monod, J.; Rickenberg, H. V. Galactoside-permease of *Escherichia coli*. *Ann*
Inst Pasteur (Paris) **1956**, *91* (6), 829-857.
(76) Bolivar, F.; Rodriguez, R. L.; Greene, P. J.; Betlach, M. C.; Heyneker, H. L.; Boyer, H. W.; Crosa, J. H.;
Falkow, S. Construction and characterization of new cloning vehicles. II. A multipurpose cloning system.
Gene **1977**, *2* (2), 95-113. DOI: 0378-1119(77)90000-2 [pii].
(77) CLSI. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria That Grow Aerobically;*
Approved Standard—Ninth Edition. CLSI document M07-A9.; Clinical and Laboratory Standards Institute,
2012.
(78) Hancock, R. E.; Wong, P. G. Compounds which increase the permeability of the *Pseudomonas*
aeruginosa outer membrane. *Antimicrob Agents Chemother* **1984**, *26* (1), 48-52.
(79) Farha, M. A.; El-Halfawy, O. M.; Gale, R. T.; MacNair, C. R.; Carfrae, L. A.; Zhang, X.; Jentsch, N. G.;
Magolan, J.; Brown, E. D. Uncovering the hidden antibiotic potential of Cannabis. *ACS Infect Dis* **2020**, *6*
(3), 338-346. DOI: 10.1021/acsinfecdis.9b00419.