Mating-compatibility genes employed as diagnostic markers to identify novel incursions of the myrtle rust pathogen *Austropuccinia psidii*

Jinghang Feng[1], Austin Bird[1], Zhenyan Luo[1], Rita Tam[1], Luc Shepherd[1,2], Lydia Murphy[1], Lavi Singh[1], Abigail Graetz[1], Mareike Moeller[1], Lilian Amorim[3], Nelson Sidnei Massola Júnior[3], M. Asaduzzaman Prodhan[2], Louise Shuey[4], Douglas Beattie[5], Alejandro Trujillo Gonzalez[6], Peri A. Tobias[7], Amanda Padovan[1], Rohan Kimber[8], Alistair McTaggart[9], Monica Kehoe[2], Benjamin Schwessinger[1,†,*], Thaís R. Boufleur[1,3,*]

[1] Research School of Biology, The Australian National University, Canberra, ACT 2601, Australia.

[2] DPIRD Diagnostics and Laboratory Services, Department of Primary Industries and Regional Development; 3 Baron-Hay Court, South Perth, WA 6151, Australia.

[3] Department of Plant Pathology and Nematology, Luiz de Queiroz College of Agriculture, University of São Paulo, Piracicaba, SP 13418-900, Brazil.

[4] Queensland Department of Agriculture and Fisheries, Ecosciences Precinct, 41 Boggo Rd, Dutton Park QLD 4102, Australia.

[5] Department of Health, Australian Government.

[6] Centre for Conservation Ecology and Genomics, Institute for Applied Ecology, University of Canberra, Bruce, Australian Capital Territory, Australia.

[7] School of Life and Environmental Sciences, Faculty of Science, The University of Sydney, Camperdown, New South Wales 2006, Australia.

[8] Crop Sciences, South Australian Research and Development Institute (SARDI), Waite Research Precinct, Urrbrae, SA 5064, Australia

[9] Centre for Horticultural Science, Queensland Alliance for Agriculture and Food Innovation, The

University of Queensland, Ecosciences Precinct, Dutton Park, Queensland, Australia

J. Feng, A. Bird, and Z. Luo contributed equally to this manuscript.

*B. Schwessinger and T. R. Boufleur share the last authorship.

[†]Corresponding author: B. Schwessinger; E-mail: benjamin.schwessinger@anu.edu.au

ABSTRACT

Austropuccinia psidii is the causal agent of myrtle rust in over 480 species within the

family Myrtaceae. Lineages of *A. psidii* are structured by host in its native range, and

some have success on new-encounter hosts. For example, the pandemic biotype has

spread beyond South America, and proliferation of other lineages is an additional risk to

biodiversity and industries. Efforts to manage *A. psidii* incursions, including lineage

differentiation, relies on variable microsatellite markers. Testing these markers is time-

consuming and complex, particularly on a large scale. We designed a novel diagnostic

approach targeting the fungal mating-type *HD* (homeodomain) transcription factor locus

to address these limitations. The *HD* locus (*bW1/2*-HD1 and *bE1/2-HD2)* is highly

polymorphic, facilitating clear biological predictions about its inheritance from founding

populations. To be considered the same lineage, all four *HD* alleles must be identical.

Our lineage diagnostics relies on PCR amplification of the *HD* locus in different

genotypes of *A. psidii* followed by amplicon sequencing using Oxford Nanopore

Technologies (ONT) and comparative analysis. The lineage-specific assay was

validated on four isolates with existing genomes, uncharacterized isolates, and directly

from infected leaf material. We reconstructed *HD* alleles from amplicons and confirmed

their sequence identity relative to their reference. Genealogies using *HD* alleles

confirmed the variations at the *HD* loci among lineages/isolates. Our study establishes a

robust diagnostic tool, for differentiating known lineages of *A. psidii* based biological

predictions. This tool holds promise for detecting new pathogen incursions and can be

refined for broader applications, including air-sample detection and mixed-isolate

infections.

Keywords: Myrtaceae, Diagnostics, Oxford Nanopore Technologies, mating-type,

Homeodomain genes.

Background

Austropuccinia psidii, the causal agent of myrtle rust in over 480 host species

within the Myrtaceae family (Soewarto et al. 2019; Carnegie and Giblin 2021), is among

the world's top ten priority fungal species for biosecurity (Hyde et al. 2018). Its high

virulence and rapid adaptability to new environments is a threat to biodiversity and

industries (Chock 2020), especially in regions like Australia and New Zealand, where

species of the Myrtaceae family prevail (Hyde et al. 2018). In eucalypts, for example,

losses in volume due to rust severity can vary from 23% to 35% (Santos et al. 2020).

Initially described in Brazil (Winter 1884), *A. psidii* remained limited to the

Americas for many decades before spreading to all continents, except Europe and

Antarctica (Simpson et al. 2006). In Australia, where the pathogen was first detected in

2010 (Carnegie et al. 2010), only the pandemic biotype (group of organisms with

identical genetic constitution) has been reported, and isolates belonging to exotic

lineages of *A. psidii* are considered a threat to Australian natural environments and

commercial native forests (DAFF 2023; Makinson et al. 2020). Disease symptoms and

spore morphology are highly similar across isolates of *A. psidii* belonging to different

lineages, even in cases with strong host associations (Morales et al. 2023; Boufleur et

al. 2023; Ferrarezi et al. 2022). The disease is predominantly caused by the clonal

stage of the pathogen and started by urediniospores. It is characterized by the initial

appearance of small chlorotic spots that develop into bright orange pustules that

sporulate to generate re-invective urediniospores under most conditions (Glen et al.,

2007).

Early detection and diagnosis are crucial for tracking, and potentially limiting rust

fungi incursions (Hussain et al. 2020). Microsatellite markers have been used to

differentiate lineages of *A. psidii* (Stewart et al. 2017; Graça et al. 2013), but their

application can be time-consuming and complex, particularly on a large scale. The

current internationally approved assay to diagnose *A. psidii* is a species-specific qPCR

(Quantitative Polymerase Chain Reaction) (IPPC 2018; Baskarathevan et al. 2016),

however, the choice of gene lacks the variability needed to differentiate among

pathogen lineages (Beenken 2017; Boufleur et al. 2023; Bini 2016). Therefore, there is

a need to identify novel target regions that precisely diagnose different lineages of *A.*

psidii for faster and precise action in a biosecurity response.

Mating in fungi is controlled through genes expressed at mating-type (*MAT*) loci

(Wilson et al. 2015). In rust fungi, these are two unlinked loci, one contains pheromone

precursors and receptors (*P/R*) and the other contains homeodomain (*HD*) transcription

factors that are closely linked via a short DNA sequence. The *HD* locus encodes *bW-*

HD1 and *bE-HD2* genes, which are highly multiallelic in rust fungi (Luo et al. 2024) and

many Basidiomycota (Coelho et al. 2017). These transcription factors form

heterodimeric complexes between alleles and regulate cellular development during

mating and the fungal life cycle (Coelho et al. 2017; Cuomo et al. 2017; Holden et al.

2023; Wilson et al. 2015). The analysis of *A. psidii* genomes confirmed physically

unlinked, heterozygous *P/R* and *HD* loci, supporting that mate compatibility in this

pathogen is governed by two multiallelic *HD* genes (*bW-HD1* and *bE-HD2*) and a

biallelic *P/R* gene (Ferrarezi et al. 2022).

The aim of this study was to develop a highly sensitive assay for the detection

and identification of *A. psidii* lineages distinct from the pandemic biotype. This assay

can be used for monitoring existing incursions/outbreaks, and to help prevent and limit

further incursions of exotic lineages. Here we introduce novel primers designed to target

the mating-type *HD* locus of *A. psidii*. These primers are designed to be used in

combination with long-read sequencing such as those facilitated by Oxford Nanopore

Technologies (ONT).□□□

Material and Methods

HD locus identification and primer design

The *HD* locus of three *A. psidii* lineages was identified on complete dikaryotic

genome assemblies, including Brazilian isolates belonging to two different lineages MF-

1 (from *Eucalyptus grandis*) (PRJNA215767, GCA_000469055.2) and LFNJM1

(unpublished data) from *Syzygium jambos* (Boufleur et al. 2023), and APG1 from

Psidium guajava 2/28/2024 8:02:00 AM (unpublished data), along with the Au3 isolate

that belongs to the pandemic biotype (Au3_v2) (PRJNA810573, GCA_023105745.1,

GCA_023105775.1) (Edwards et al. 2022), and the South African isolate Apsidii_AM,

that belongs to the South African Biotype (PRJNA480390, GCA_003724095.1)

(McTaggart et al. 2018). The *HD* loci containing regions were identified with BLASTx

(v.2.15.0) in combination with annotated *bW-HD1* and *bE-HD2 A. psidii* genes, as

described by Ferrarezi et al., (2022). As expected for the *HD* locus in dikaryotic genome

assemblies, two alleles of each *bW-HD1* and *bE-HD2* gene were retrieved. The alleles

of each gene (*bW-HD1* and *bE-HD2*) were aligned separately with MACSE (v.2.07)

(Ranwez et al. 2018) and two Bayesian inference genealogy trees were generated with

Mr. Bayes v.3.2.6 (Huelsenbeck and Ronquist 2001).

Primers were designed manually based on a multiple sequence alignment of

contigs containing the *HD* alleles of the pandemic Au3_v2 isolate, the South African

isolate Apsidii_AM, and the Brazilian isolate MF-1. Two pairs of degenerate primers

were designed to amplify *bW-HD1* and *bE-HD2* individually, and the combination of the

most forward and the most reverse primers was used to amplify the full-length *HD* locus

(Table 1). Regions that were fully or nearly fully conserved having a maximum of two

single nucleotide differences between all *HD* loci alleles were selected to allow

amplification of all alleles at the same time. The expected amplicon size was ~1600 bp

and ~1400 bp for each *bW-HD* and *bE-HD,* and ~3000 bp for the complete *HD* locus.

DNA extraction and HD locus amplification

The designed primers were tested with diverse samples, including four positive

controls of *A. psidii*, ten field samples and three non-target rust species (Table 2).

Genomic DNA (gDNA) was extracted directly from *A. psidii* urediniospores, from

infected leaf material or from urediniospores of non-target rust species (Table 2) with

the DNeasy Plant mini kit (Qiagen) according to the manufacturer's instructions. The

integrity and quality of the DNA was measured with a Nanodrop spectrophotometer

(Thermo Fisher Scientific) and checked by agarose gel (0.8%) electrophoresis stained

with SYBR Safe (Thermo Fisher Scientific). The DNA concentration was determinate

using the Qubit 4 (Thermo Fisher Scientific), and all samples were adjusted to 25 ng/µL

for further studies.

In the first round of PCR tests, the aim was to evaluate the amplification of (non-)

target sequences by the designed primers followed by sequence analysis using *A. psidii*

and three non-target rust isolates (Table 2, Figure 1A-B). PCR was performed on a

Mastecycler nexus X2 thermal cycler (Eppendorf). The reaction mixture, with a final

volume of 25 µL, included 5 µL of 5X reaction buffer, 0.5 µL of dNTPs [10 mM], 1.25 µL

of each primer [10 µM], 0.25 µL of Q5 High-Fidelity DNA Polymerase (NEB), 14.75 uL

of Nuclease Free Water (NFW) and 2 µL (up to 50 ng) of template DNA. The PCR

amplification had an initial denaturation step at 98 °C for 30 s, followed by 35 cycles of

denaturation at 98 °C for 30 s, annealing at 58 °C for 30 s and extension at 72 °C for 30

s, with a final extension step at a 72 °C for 2 min. Specificity tests were run in duplicate,

and PCR products were visualised on 2% agarose gel stained with SYBR safe (Thermo

Fisher Scientific).

In the second round of PCR tests, the performance of the primers was assessed

against single-spore isolates and field samples (urediniospores and infected leaves)

collected in Brazil and Australia (Table 2). Identical reaction conditions, as described

above, were used to amplify the *bW-HD* and *bE-HD* loci individually or the full-length

HD locus (Table 2).

ONT sequencing

For Oxford Nanopore sequencing, libraries were generated following

manufacturer instructions for V14 Ligation Sequencing of amplicons (Native Barcoding

Kit V14 96 - SQK-NBD114.96) with modifications as follows. An initial bead clean was

performed using 1.2 x of 2 % Sera-Mag beads to purify the PCR product and 200 fmol

of clean DNA carried through to End-prep reaction. The End-prep reaction was

incubated at 20 °C and then 65 °C for 15 min to maximize yield. One µL of the end-

prepped DNA was amplicons were barcoded with 1 µL of Nanopore Native Barcode

using 5 µL Blunt/TA ligase master mix (NEB) in total reaction volume of 10 µL for 20

min at 20 °C. The reaction was stopped by adding 1 µL of EDTA to each ligation

reaction. The individually barcoded PCR amplicons were pooled, bead cleaned with 0.6

x volume 2% Sera-Mag beads and two washes of 70% ethanol. The pool of barcoded

PCR amplicons was eluted in 21 μL of nuclease-free water. Library preparation was

completed according to the manufacturers protocol. Twenty fmol of the barcoded library

was loaded on a MinION R10.4.1 Flowcell (FLO-MIN114) and sequencing was ran

using a MinION. Basecalling was performed with Guppy v. 6.4.2 Super High Accuracy

mode.   All   long-read   amplicon   datasets   were   deposited   to   Zenodo

(https://doi.org/10.5281/zenodo.10656657).

De novo reconstruction of HD loci and genealogies

A two-step filtering process was implemented on the base-called sequences. In

the initial step, sequencing reads were filtered based on their Phred quality scores, with

reads having a mean quality score below 15 being removed. This ensured that the

remaining reads had an average per base accuracy of ≥ 97%. The second filtering step

involved selecting sequencing reads with lengths around the expected amplicon length

for each *HD* gene region: 1500-1800 bp for *bW-HD1* and 1300-1500 bp for *bE-HD2*.

The VSEARCH clustering algorithm (Rognes et al. 2016) was applied for quality control

of sequencing reads of each sample, using global similarity during clustering. It is

important to note that the VSEARCH clustering algorithm could not recognize

sequences of the same gene in the opposite direction. Hence, we obtained two clusters

for each *bW-HD1* and *bE-HD2* alleles in the sample along with two consensus

sequences - one forward and the other reverse. The analysis code is available on

Github   (https://github.com/TheRainInSpain/Lineage-Specific-Marker.git).   And   the

dataset is available on (https://doi.org/10.5281/zenodo.10656657). Geneious software

(v.2023.2.1) was used to visualize the forward read consensus sequences.

The reconstructed amplicon and reference sequences were aligned (Table 2)

with MUSCLE (v.5.1) (Edgar 2004), and simple genealogical trees were reconstructed

using the Geneious tree builder (v.2023.2.1).

Results

Specificity of the diagnostic assay achieved through HD loci amplification and

ONT sequencing

During the initial phase of primer testing, *HD* PCRs were performed on positive

controls of *A. psidii* (Au3, MF-1, APG1 and LFNJM1) and negative controls with non-

target rust species (*Miyagia psudosphaeria, Puccinia striiformis* f. sp. *tritici, P. graminis*,

P. triticina, Thekopsora minima). In addition, we focused on the individual PCR

amplicons (*bW-HD* and *bE-HD*) because the amplification of the full-length locus was

not robust enough across samples and technical replicates. The sizes of the amplicons

were as expected in the positive control samples, being of ~1600 bp for *bW-HD*, ~1400

bp for *bE-HD*, and ~3000 kb for the full HD locus (Table 2). In addition, we observed

bands of variable sizes in some of our technical repeats of non-target species. All

samples were sequenced with our ONT amplicon sequencing workflow because our

assay does not rely exclusively on the PCR amplification product but requires that the

amplified sequences match the *A. psidii HD* sequences in a genealogical framework.

None of the non-target species amplicon sequences matched the full-length *A.*

psidii HD sequences. For each *A. psidii* isolate, four *HD* alleles (2x *bW-HD1* and 2x *bE-*

HD2) were reconstructed based on ONT sequencing results, as expected for dikaryotic

organisms. Isolates were considered identical if their four *HD* alleles were identical or

carried minor non-functional variation (for example synonymous variation, variation

outside the variable domain or within introns). The applicability of the lineage diagnostic

test was confirmed by comparing the *de novo* reconstructed consensus sequences

derived from ONT amplicon sequencing with the *HD* amplicon sequences derived from

the reference genomes (APG1, MF-1, LFNJM1, Au3 and Apsidii_AM). All *de novo*

reconstructed ONT amplicon sequences clearly grouped with their respective *HD* alleles

obtained from reference genomes (Table 2, Figure 1A-B). Moreover, the reference

trees for *bW-HD1* and *bE-HD2* revealed that the Brazilian isolates and the South African

isolate carry clearly distinct alleles when compared against the pandemic isolate (Figure

1A-B), corroborating variations previously described using microsatellite markers

(Stewart et al. 2017; Roux et al. 2016; Graça et al. 2013).

Primers targeting the *HD* region distinguished different lineages of *A. psidii*

The designed primers successfully amplified individual *HD* loci of DNA extracted

from different sources of field samples, including urediniospores and infected leaf

material. As observed previously, the full *HD* locus amplification was not possible for the

LFNJRM1 isolate and the field samples (Table 2). The reconstructed amplicon and

reference sequences were aligned and revealed a pairwise identity of 67.9% to 100%

for *bW-HD1* and 76.9% and to 100% for *bE-HD2* alleles across the analyzed sample

isolates. Our results revealed that an isolate collected from field samples in Australia in

2022 (SYD) belong to the same lineage as the pandemic lineage because all four *HD*

alleles clearly grouped with those derived from the pandemic reference isolate, while

isolates collected from field samples from Brazil, belong to different lineages having at

least two different *HD* alleles (Figure 1C-D).

Discussion

Traditionally, *A. psidii* lineages have been identified using microsatellite markers

(Stewart et al. 2017; Graça et al. 2013; Roux et al. 2016; Sandhu et al. 2016). In this

study, a diagnostic assay targeting the *HD* locus of *A. psidii*, coupled with ONT

sequencing was developed. This method successfully amplified diverse copies of the

HD locus from *A. psidii* isolates collected in Australia and Brazil and is predicted to work

for the South African isolate.

The genealogical tree of *de novo* reconstructed amplicons confirmed the

biological expectation of lineage specific variation among isolates originating from single

spores, field samples from different hosts, and distinct geographic locations. The isolate

collected from field samples in Australia had *HD* alleles that matched the pandemic

lineage, whereas in Brazil, the center of origin of the disease, there was a strong

association of *HD* allele status with their original host species, as previously observed

(Morales et al. 2023; Graça et al. 2013; Stewart et al. 2017). Overall, the high allelic

diversity at the *HD* locus in our samples is in line with the previously identified diversity

observed at this locus in rust fungi including *A. psidii* and *Puccinia* spp. (Ferrarezi et al.

2022; Holden et al. 2023; Luo et al., 2024).

Individual *HD* genes were amplified in field samples for which the entire *HD* locus

could not be amplified by PCR. In the future, our primers and/or amplification conditions

can be further improved to amplify the entire locus of natural *A. psidii* populations within

its center of diversity, as it is easier in a Biosecurity response to amplify one long region

than two. In addition, we observed off target PCR amplification in some of our technical

replicates of non-target species, yet none of these sequences match the *A. psidii HD*

sequence. Improved primers or nested PCRs could improve PCR specificity increasing

sensitivity and throughput when applied under routine testing conditions.

To date, the non-pandemic isolates shared at most one out of four *HD* alleles

with the pandemic isolate (Au_3). This finding shows non-pandemic isolates can be

differentiated from the pandemic lineage based on their *HD* alleles. This protocol serves

as a valuable tool in detecting new incursions of the pathogen in regions where a single

lineage is present as for example in Australia where only a single incursion of the

pandemic biotype has been reported to date. Further, comparing *HD* allele identity with

known reference sequences could link novel incursions with related populations in

source regions. This could help in identifying risks in import pathways of this exotic

pathogen and improve risk mitigation strategies. In addition, the assay could potentially

detect recombination between populations if purified single pustule isolates were

analyzed. In the future, we anticipate that the diagnostic test can be refined to detect

urediniospores of *A. psidii* from complex samples derived from air-sampling or mixed

infections to enable structured targeted surveillance of this pathogen on the ground.

Acknowledgments

The authors thank Danièle Giblot-Ducray and Kelly Hill for the critical feedback and support.

Funding

A. T. Gonzalez, R. Kimber, and B. Schwessinger were supported by an Australian Government grant from the Department of Agriculture, Fisheries and Forestry entitled "Automated air sampling for remote surveillance and high throughput processing of environmental samples for eDNA analyses". São Paulo Research Foundation (FAPESP) Grant 2019/13191-5. T. R. Boufleur was supported by FAPESP Grants 2022/11900-1 and 2021/01606-6.

Literature Cited

Baskarathevan, J., Taylor, R. K., Ho, W., McDougal, R. L., Shivas, R. G., and Alexander, B. J. R. 2016. Real-Time PCR Assays for the detection of *Puccinia psidii*. Plant Disease. 100:617–624.

Beenken, L. 2017. *Austropuccinia*: a new genus name for the myrtle rust *Puccinia psidii* placed within the redefined family Sphaerophragmiaceae (Pucciniales). Phytotaxa. 297:53.

Bini, A. P. 2016. Estudo molecular do desenvolvimento de *Puccinia psidii* Winter *in vitro*

e no processo de infecção em *Eucalyptus grandis*. Available at:

http://www.teses.usp.br/teses/disponiveis/11/11137/tde-10112016-161846/ [Accessed

September 19, 2022].

Boufleur, T. R., Morales, J. V. P., Martins, T. V., Gonçalves, M. P., Júnior, N. S. M., and

Amorim, L. 2023. A diagnostic guide for myrtle rust. Plant Health Progress. 24:242–251.

Carnegie, A. J., Lidbetter, J. R., Walker, J., Horwood, M. A., Tesoriero, L., Glen, M., et

al. 2010. *Uredo rangelii*, a taxon in the guava rust complex, newly recorded on

Myrtaceae in Australia. Australasian Plant Pathol. 39:463–466.

Coelho, M. A., Bakkeren, G., Sun, S., Hood, M. E., and Giraud, T. 2017. Fungal sex: the

Basidiomycota. Microbiol Spectr. 5.

Cuomo, C. A., Bakkeren, G., Khalil, H. B., Panwar, V., Joly, D., Linning, R., et al. 2017.

Comparative analysis highlights variable genome content of wheat rusts and divergence

of the mating loci. G3 Genes|Genomes|Genetics. 7:361–376.

Department of Agriculture, Fisheries and Forestry. 2023. Myrtle rust (exotic strains).

Biosecurity and Trade. Available at: https://www.agriculture.gov.au/biosecurity-

trade/pests-diseases-weeds/plant/myrtle-

rust#:~:text=Myrtle%20rust%20(exotic%20strains)%20is,damaging%20to%20our%20eu

calyptus%20trees [Accessed December 19, 2023].

Edgar, R. C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res. 32:1792–1797.

Edwards, R. J., Dong, C., Park, R. F., and Tobias, P. A. 2022. A phased chromosome-

level genome and full mitochondrial sequence for the dikaryotic myrtle rust pathogen,

*Austropuccinia   psidii.   :2022.04.22.489119   Available   at:

https://www.biorxiv.org/content/10.1101/2022.04.22.489119v1 [Accessed August 22,

2022].

Ferrarezi, J. A., McTaggart, A. R., Tobias, P. A., Hayashibara, C. A. A., Degnan, R. M.,

Shuey, L. S., et al. 2022. *Austropuccinia psidii* uses tetrapolar mating and produces

meiotic spores in older infections on *Eucalyptus grandis*. Fungal Genetics and Biology.

160:103692.

Graça, R. N., Ross-Davis, A. L., Klopfenstein, N. B., Kim, M.-S., Peever, T. L., Cannon,

P. G., et al. 2013. Rust disease of eucalypts, caused by *Puccinia psidii*, did not originate

via host jump from guava in Brazil. Mol Ecol. 22:6033–6047.

Holden, S., Bakkeren, G., Hubensky, J., Bamrah, R., Abbasi, M., Qutob, D., et al. 2023.

Uncovering the history of recombination and population structure in western Canadian

stripe rust populations through mating type alleles. BMC Biology. 21:233.

Huelsenbeck, J. P., and Ronquist, F. 2001. MRBAYES: Bayesian inference of

phylogenetic trees. Bioinformatics. 17:754–755.

Hussain, K. K., Malavia, D., Johson, E., Littlechild, J. A., Winlove, P., Vollmer, F., et al.

2020. Biosensors and diagnostics for fungal detection. Journal of Fungi. 6:349.

Hyde, K. D., Al-Hatmi, A. M. S., Andersen, B., Boekhout, T., Buzina, W., Dawson, T. L.,

et al. 2018. The world's ten most feared fungi. Fungal Diversity. 93:161–194.

IPPC. 2018. DP 26: *Austropuccinia psidii* .

Leite, T. F., Moon, D. H., Lima, A. C. M., Labate, C. A., and Tanaka, F. A. O. 2013. A

simple protocol for whole leaf preparation to investigate the interaction between

Puccinia psidii and *Eucalyptus grandis*. Australasian Plant Pathol. 42:79–84.

Luo, Z; McTaggart, A; Schwessinger, B. 2024. Genome biology and evolution of matyng

type loci in four cereal rust fungi. Available at:

https://www.biorxiv.org/content/10.1101/2023.03.02.530769v3.

Makinson, R. O., Pegg, G. S., and Carnegie, A. J. 2020. Myrtle Rust in Australia – a

National Action Plan.

McTaggart, A. R., Duong, T. A., Le, V. Q., Shuey, L. S., Smidt, W., Naidoo, S., et al.

2018. Chromium sequencing: the doors open for genomics of obligate plant pathogens.

BioTechniques. 65:253–257.

Morales, J. V. P., Boufleur, T. R., Gonçalves, M. P., Parisi, M. C. M., Loehrer, M.,

Schaffrath, U., et al. 2023. Differential aggressiveness of *Austropuccinia psidii* isolates

from guava and rose apple upon cross-inoculation. Plant Pathology. Available at:

https://onlinelibrary.wiley.com/doi/abs/10.1111/ppa.13850 [Accessed January 12, 2024].

Ranwez, V., Douzery, E. J. P., Cambon, C., Chantret, N., and Delsuc, F. 2018. MACSE

v2: Toolkit for the Alignment of Coding Sequences Accounting for Frameshifts and Stop

Codons. Mol Biol Evol. 35:2582–2584.

Rognes, T., Flouri, T., Nichols, B., Quince, C., and Mahé, F. 2016. VSEARCH: a

versatile open source tool for metagenomics. PeerJ. 4:e2584.

Roux, J., Granados, G. M., Shuey, L., Barnes, I., Wingfield, M. J., and McTaggart, A. R.

2016. A unique genotype of the rust pathogen, *Puccinia psidii*, on Myrtaceae in South

Africa. Australasian Plant Pathol. 45:645–652.

Sandhu, K. S., Karaoglu, H., Zhang, P., and Park, R. F. 2016. Simple sequence repeat

markers support the presence of a single genotype of *Puccinia psidii* in Australia. Plant

Pathol. 65:1084–1094.

Santos, A. P., Gomes, R. L., Furtado, E. L., Passos, J. R. S. 2020. Quantifying losses in

productivity by the rust in eucalypt plantations in Brazil. Fores. Ecol. Manag. 468:

118170.

Simpson, J. A., Thomas, K., and Grgurinovic, C. A. 2006. Uredinales species

pathogenic on species of Myrtaceae. Australas. Plant Pathol. 35:549–562.

Stewart, J. E., Ross-Davis, A. L., Gra□a, R. N., Alfenas, A. C., Peever, T. L., Hanna, J.

W., et al. 2017. Genetic diversity of the myrtle rust pathogen (*Austropuccinia psidii*) in

the Americas and Hawaii: Global implications for invasive threat assessments ed. M.

Cleary. For. Path. 48:e12378.

Wilson, A. M., Godlonton, T., van der Nest, M. A., Wilken, P. M., Wingfield, M. J., and

Wingfield, B. D. 2015. Unisexual reproduction in *Huntiella moniliformis*. Fungal Genetics

and Biology. 80:1–9.

Winter, G. R. 1884. Rabenhorstii fungi europaei et extraeuropaei exsiccati cura Dr. G.

Winter, CenturiaXXXI et XXXII. Hedwigia. 23:164–172.

Tables

Table 1. List of primers used in the present study.

Primers	Sequence (5' - 3')	Region	Changes
HDFor2DG (forward)[a]	ATACAGTTYAGGTTWTRGCG	*bW-HD*	C/T, A/T, A/G
HDMR1 (reverse)	GAAAGGAAATATTGCCACT	*bW-HD*	-
HDMF2 DG (forward)	YGACCGCCTTCCTTTGAG	*bE-HD*	C/T,
HDRev2DG (reverse)[a]	GTGTCSAAGCWACCAAAATC	*bE-HD*	C/G, A/T

409 [a] Primers used to amplify the full *HD* locus.

418 **Table 2.** Description of isolates and populations and of *A. psidii* and non-target rusts used in this

419 study

Organism	Code	Source	Country	Amplification[a]			Alleles			
				Full locus	*bW-HD1*	*bE-HD2*	*HD1*		*HD2*	
Isolates										
Austropuccinia psidii	Au3[b]	*Agonis flexuosa*	AUS	+	+	+	Au3 HapA	Au3 HapB	Au3 HapA	Au3 HapB
A. psidii	MF-1 [b]	*Eucalyptus grandis*	BR	+	+	+	MF-1 HapA	MF-1 HapB	MF-1 HapA	MF-1 HapB
A. psidii	APG1 (GM1)	*Psidium guajava*	BR	-	-	+	NA	NA	APG1 HapA	NA
A. psidii	LFNJM1 [b]	*Syzygium jambos*	BR	+	+	+	LFNJM1 HapA	LFNJM1 HapB	LFNJM1 HapA	LFNJM HapB
A. psidii	LFNJM3	*S. jambos*	BR	+	+	-	LFNJM3 HapA	NA	NA	NA
A. psidii	LFNJM4	*S. jambos*	BR	+	+	+	LFNJM4 HapA	LFNJM4 HapB	LFNJM4 HapA	LFNJM HapB
A. psidii	LFNJRM1	*S. samarangense*	BR	-	+	+	LFNJRM1 HapA	NA	LFNJRM1 HapA	LFNJR HapB

A. psidii	Apsidii AM	*S. jambos*	SA	NA	NA	NA	SA HapA	SA HapB	SA HapA	SA Hap
Populations (urediniospores)										
A. psidii	CA	*Plinia edulis*	BR	-	+	+	CA Hap1	CA Hap2	CA Hap1	CA Ha
A. psidii	CG	*Eugenia dysenterica*	BR	-	+	+	LFNSDP1 HapA	LFNSDP1 HapB	LFNSDP1 HapA	LFNSD HapB
Populations (infected leaves)										
A. psidii	LFNEP1	*Eugenia stipitata*	BR	-	+	+	LFNEP1 HapA	NA	LFNEP1 HapA	LFNEP HapB
A. psidii	3.1	*Pimenta dioica*	BR	-	+	+	3.1 HapA	NA	3.1 HapA	3.1 Hap
A. psidii	SYD	*Melaleuca quinquenervia*	AUS	-	+	+	SYD HapA	SYD HapB	SYD HapA	SYD HapB
Non-target rust species										
Miyagia psudosphaeria	-	-	-	NS	NP	NA	NA	NA	NA	NA
Puccinia triticina	-	-	-	NS	NS	NA	NA	NA	NA	NA
P. graminis f. sp. *avenae*	-	-	-	NS	NS	NA	NA	NA	NA	NA
P. striiformis f. sp. *tritici*	-	-	-	NS	NS	NA	NA	NA	NA	NA
Thekopsora minima	-	-	-	-	NS	NA	NA	NA	NA	NA

[a] Conventional polymerase chain reaction (PCR) with the three sets of primers developed in the

present study. Full *HD* locus, HDFor2DG/HDRev2DG amplifies a fragment of 3000 kb; *bW-HD,*

HDFor2DG/HDMR1 amplifies a fragment of 1500 kb; *bE-HD,* HDMF2/HDRev2DG amplifies a

fragment of 1400 kb; +, positive; -, negative.

[b] *A. psidii* and non-target rust isolates used to confirm the specificity of the designed primers.

"NS" is defined as NS specific amplification in at least one of the technical replicates, "NA" as

not available, "-" as no amplification, "+" as amplification of the expected size amplicon product.

Figure captions

Figure 1: Bayesian inference genealogical trees reconstructed from *HD* (homeodomain) genes

of *Austropuccinia psidii*. Genealogical trees reconstructed from the alignment of two *bW-HD1*

(A, C) or two *bE-HD2* **(B, D)** alleles per dikaryotic genome assembly of reference isolates **(A-B),**

and with consensus sequences for each allele *de novo* reconstructed from ONT long-read

amplicons of *bW-HD1* and *bE-HD2* (C-D). Lineage identification is indicated by the combination

of different colors in the bottom. Reference sequences are indicated in bold (C-D). Bayesian

posterior probabilities are shown on the nodes.