Mating-compatibility genes employed as diagnostic markers to identify novel incursions

of the myrtle rust pathogen *Austropuccinia psidii*

Jinghang Feng[1], Austin Bird[1], Zhenyan Luo[1], Rita Tam[1], Luc Shepherd[1,2], Lydia

Murphy[1], Lavi Singh[1], Abigail Graetz[1], Mareike Moeller[1], Lilian Amorim[3], Nelson Sidnei

Massola Júnior[3], Asad Prodhan[2], Louise Shuey[4], Douglas Beattie[5], Alejandro Trujillo

Gonzalez[6], Peri A. Tobias[7], Amanda Padovan[1], Rohan Kimber[8], Alistair McTaggart[9],

Monica Kehoe[2], Benjamin Schwessinger[1,†,*], Thaís R. Boufleur[1,3,*]

[1] Research School of Biology, The Australian National University, Canberra, ACT 2601,

Australia.

[2] DPIRD Diagnostics and Laboratory Services, Department of Primary Industries and Regional

Development; 3 Baron-Hay Court, South Perth, WA 6151, Australia.

[3] Department of Plant Pathology and Nematology, Luiz de Queiroz College of Agriculture,

University of São Paulo, Piracicaba, SP 13418-900, Brazil.

[4]Queensland Department of Agriculture and Fisheries, Ecosciences Precinct, 41 Boggo Rd,

Dutton Park QLD 4102, Australia.

[5] Department of Health, Australian Government.

[6] Centre for Conservation Ecology and Genomics, Institute for Applied Ecology, University of

Canberra, Bruce, Australian Capital Territory, Australia.

[7] School of Life and Environmental Sciences, Faculty of Science, The University of Sydney,

Camperdown, New South Wales 2006, Australia.

[8] Crop Sciences, South Australian Research and Development Institute (SARDI), Waite

Research Precinct, Urrbrae, SA 5064, Australia

[9] Centre for Horticultural Science, Queensland Alliance for Agriculture and Food Innovation, The

University of Queensland, Ecosciences Precinct, Dutton Park, Queensland, Australia

J. Feng, A. Bird, and Z. Luo contributed equally to this manuscript.

*B. Schwessinger and T. R. Boufleur share the last authorship.

[†]Corresponding author: B. Schwessinger; E-mail: benjamin.schwessinger@anu.edu.au

32   **ABSTRACT**

Austropuccinia psidii is the causal agent of rust fungal disease in over 480 species

within the family Myrtaceae and poses a serious threat to global ecosystems. Lineages

of *A. psidii* are structured by host in its native range, and some have success on new-

encounter hosts. For example, the pandemic biotype has spread beyond South

America, and more potential lineages represent substantial additional risk to biodiversity

and industries in the non-native range. Efforts to manage the threat of exotic *A. psidii*

incursions, including the differentiation of lineages, relies on variable microsatellite

markers. Testing these markers is time-consuming, and complex, particularly on a large

scale. We designed a novel diagnostic targeting the fungal mating-type *HD*

(homeodomain) transcription factor locus to address these limitations. The *HD* locus

(*bW1/2*-HD1 and *bE1/2*-HD2) is highly polymorphic, facilitating clear biological

predictions about its inheritance from founding populations, making it a suitable target

for diagnostics. For example, host-delimited lineages contain the same four *HD* alleles.

Conversely, if an isolate does not share all four alleles with the isolate belonging to the

pandemic biotype it is derived from another lineage, or there has been potential

recombination among lineages. Our lineage diagnostics relies on PCR amplification of

the *HD* locus followed by amplicon sequencing using Oxford Nanopore Technologies

(ONT) and comparative analysis. We validated our lineage-specific assay on four

isolates with existing genomes, uncharacterized isolates from Brazil, and directly from

infected leaf material from Australia and Brazil. We reconstructed *HD* alleles from

amplicons, confirmed their sequence identity relative to their reference and

reconstructed genealogies to infer their relationships. Genealogies using *HD* alleles

confirmed the variations at the *HD* loci among lineages/isolates. Our study establishes a

robust diagnostic tool, for differentiating pandemic and non-pandemic lineages based on

clear biological predictions. This tool holds promise for detecting new pathogen

incursions and can be refined for broader applications, including air-sample detection

and mixed-isolate infections.

Keywords: *Austropuccinia psidii*, Myrtaceae, Diagnostics, Oxford Nanopore

Technologies, mating type, Homeodomain genes.

Background

Austropuccinia psidii, the causal agent of myrtle rust in over 480 host species

within the Myrtaceae family (Soewarto et al. 2019; Carnegie and Giblin 2021), is among

the world's top ten priority fungal species for biosecurity (Hyde et al. 2018). Its high

virulence and rapid adaptability to new environments present a major threat to

biodiversity and industries (Chock 2020), especially in regions like Australia and New

Zealand, where species of the Myrtaceae family are prevalent (Hyde et al. 2018). For

eucalyptus, for example, losses in volume due rust severity can vary from 23% to 35%

(Santos et al. 2020).

Initially described in Brazil (Winter 1884), *A. psidii* remained limited to the

Americas for many decades before spreading to all continents, except Europe and

Antarctica (Simpson et al. 2006). In Australia, where the pathogen was first detected in

2010 (Carnegie et al. 2010), only the pandemic biotype (group of organisms with

identical genetic constitution) of the pathogen has been reported thus far, while isolates

belonging to exotic lineages of *A. psidii* are considered one of the most serious threats

to Australia's natural environment and commercial native forests (DAFF 2023; Makinson

et al. 2020). Disease symptoms and spore morphology are highly similar across isolates

of *A. psidii* belonging to different lineages, even in cases with strong host associations

(Morales et al. 2023; Boufleur et al. 2023; Ferrarezi et al. 2022). The disease is

predominantly caused by the clonal stage of the pathogen and started by

urediniospores. It is characterized by the initial appearance of small chlorotic spots that

develop into bright orange pustules that sporulate as the disease progresses generating

re-invective urediniospores under most conditions (Glen et al., 2007).

Early detection and diagnosis are crucial for tracking, and potentially limiting rust

fungi incursions (Hussain et al. 2020). Microsatellite markers have been used to

differentiate lineages of *A. psidii* (Stewart et al. 2017; Graça et al. 2013), but their

application can be time-consuming and complex, particularly on a large scale. The

current internationally approved assay to diagnose *A. psidii* is a species-specific qPCR

(Quantitative Polymerase Chain Reaction) (IPPC 2018; Baskarathevan et al. 2016),

however, the choice of gene lacks the variability needed to differentiate among

pathogen lineages (Beenken 2017; Boufleur et al. 2023; Bini 2016). Therefore, there is a

need for the identification of novel target regions that precisely identify *A. psidii* to

develop diagnostic tests that enable faster and more precise identification, including the

specific pathogen lineages.

Mating in fungi is controlled through genes expressed at mating-type (*MAT*) loci

(Wilson et al. 2015). In rust fungi, these are two unlinked loci, one contains pheromone

precursors and receptors (*P/R*) and the other contains homeodomain (*HD*) transcription

factors that closely linked via a short DNA sequence. The *HD* locus encodes *bW-HD1*

and *bE-HD2* genes which are highly multiallelic in rust fungi (Luo et al. 2024) and many

Basidiomycota (Coelho et al. 2017). These transcription factors form heterodimeric

complexes between alleles and regulate cellular development during mating and the

fungal life cycle (Coelho et al. 2017; Cuomo et al. 2017; Holden et al. 2023; Wilson et al.

2015). The analysis of *A. psidii* genomes confirmed physically unlinked, heterozygous

P/R and *HD* loci, being mate compatibility in this pathogen governed by two multiallelic

HD genes (*bW-HD1* and *bE-HD2*) and a biallelic *P/R* gene (Ferrarezi et al. 2022).

The aim of this study was to develop a highly sensitive assay for the detection

and identification of *A. psidii* lineages distinct from the pandemic biotype. This assay can

be used for monitoring existing incursions/outbreaks, and to help prevent and limit

further incursions of exotic lineages. Here we introduce novel primers designed to target

the mating-type *HD* locus of *A. psidii*. These primers are designed to be used in

combination with long-read sequencing such as those facilitated by Oxford Nanopore

Technologies (ONT).

Material and Methods

HD locus identification and primer design

The *HD* locus of three *A. psidii* lineages was identified on complete dikyarotic

genome assemblies, including Brazilian isolates belonging to two different lineages MF-

1 (from *Eucalyptus grandis*) (PRJNA215767, GCA_000469055.2) and LFNJM1

(unpublished data) from *Syzygium jambos* (Boufleur et al. 2023), and APG1 from

Psidium guajava 2/19/2024 12:36:00 PM (unpublished data), along with the Au3 isolate

that belongs to the pandemic biotype (Au3_v2) (PRJNA810573, GCA_023105745.1,

GCA_023105775.1) (Edwards et al. 2022), and the South African isolate Apsidii_AM,

that belongs to the South African Biotype (PRJNA480390, GCA_003724095.1)

(McTaggart et al. 2018). The *HD* loci containing regions were identified with BLASTx

(v.2.15.0) in combination with annotated *bW-HD1* and *bE-HD2 A. psidii* genes, as

described by Ferrarezi et al., (2022). As expected for the *HD* locus in dikaryotic genome

assemblies, two alleles of each *bW-HD1* and *bE-HD2* gene were retrieved. The alleles

of each gene (*bW-HD1* and *bE-HD2*) were aligned separately with MACSE (v.2.07)

(Ranwez et al. 2018) and two Bayesian inference genealogy trees were generated with

Mr. Bayes v.3.2.6 (Huelsenbeck and Ronquist 2001).

Primers were designed manually based on a multiple sequence alignment of

contigs containing the *HD* alleles of the pandemic Au3_v2 isolate, the South African

isolate Apsidii_AM, and the Brazilian isolate MF-1. Two pairs of degenerate primers

were designed to amplify *bW-HD1* and *bE-HD2* individually, and the combination of the

most forward and the most reverse primers was used to amplify the full-length *HD* locus

(Table 1). Regions that were fully or nearly fully conserved having a maximum of two

single nucleotide differences between all *HD* loci alleles were selected to allow

amplification of all alleles at the same time. The expected amplicon size was ~1600 bp

and ~1400 bp for each *bW-HD* and *bE-HD,* and ~3000 bp for the complete *HD* locus.

DNA extraction and HD locus amplification

The designed primers were tested with diverse samples, including four positive

controls of *A. psidii*, ten field samples and three non-target rust species (Table 2).

Genomic DNA (gDNA) was extracted directly from *A. psidii* urediniospores, from

infected leaf material or from urediniospores of non-target rust species (Table 2) with the

DNeasy Plant mini kit (Qiagen) according to the manufacturer's instructions. The

integrity and quality of the DNA was measured with a Nanodrop spectrophotometer

(Thermo Fisher Scientific) and checked by agarose gel (0.8%) electrophoresis stained

with SYBR Safe (Thermo Fisher Scientific). The DNA concentration was determinate

using the Qubit 4 (Thermo Fisher Scientific), and all samples were adjusted to 25 ng/μL

for further studies.

In the first round of PCR tests, the aim was to evaluate the amplification of (non-)

target sequences by the designed primers followed by sequence analysis using *A. psidii*

and three non-target rust isolates (Table 2, Figure 1A-B). PCR was performed on a

Mastecycler nexus X2 thermal cycler (Eppendorf). The reaction mixture, with a final

volume of 25 μL, included 5 μL of 5X reaction buffer, 0.5 μL of dNTPs [10 mM], 1.25 μL

of each primer [10 μM], 0.25 μL of Q5 High-Fidelity DNA Polymerase (NEB), 14.75 uL of

Nuclease Free Water (NFW) and 2 μL (up to 50 ng) of template DNA. The PCR

amplification had an initial denaturation step at 98 °C for 30 s, followed by 35 cycles of

denaturation at 98 °C for 30 s, annealing at 58 °C for 30 s and extension at 72 °C for 30

s, with a final extension step at a 72 °C for 2 min. Specificity tests were run in duplicate,

and PCR products were visualised on 2% agarose gel stained with SYBR safe (Thermo

Fisher Scientific).

In the second round of PCR tests, the performance of the primers was assessed

against single-spore isolates and field samples (urediniospores and infected leaves)

collected in Brazil and Australia (Table 2). Identical reaction conditions, as described

above, were used to amplify the *bW-HD* and *bE-HD* loci individually or the full-length *HD*

locus (Table 2).

ONT sequencing

For Oxford Nanopore sequencing, libraries were generated following

manufacturer instructions for V14 Ligation Sequencing of amplicons (Native Barcoding

Kit V14 96 - SQK-NBD114.96) with modifications as follows. An initial bead clean was

performed using 1.2 x of 2 % Sera-Mag beads to purify the PCR product and 200 fmol of

clean DNA carried through to End-prep reaction. The End-prep reaction was incubated

at 20 °C and then 65 °C for 15 min to maximize yield. One µL of DNA was barcode to 1

µL of the end-prepped PCR amplicons using 5 µL Blunt/TA ligase master mix (NEB) in

total reaction volume of 10 µL for 20 min at 20 °C. The reaction was stopped by adding

1 µL of EDTA to each ligation reaction. The individually barcoded PCR amplicons were

pooled, bead cleaned with 0.6 x volume 2% Sera-Mag beads and two washes of 70%

ethanol. The pool of barcoded PCR amplicons was eluted in 21 µL of nuclease-free

water. Library preparation was completed according to the manufacturers protocol.

Twenty fmol of the barcoded library were loaded on a MinION R10.4.1 Flowcell (FLO-

MIN114) and sequencing was ran using a MinION. Basecalling was performed with

Guppy v. 6.4.2 Super High Accuracy mode. All long-read amplicon datasets were

deposited to Zenodo (https://doi.org/10.5281/zenodo.10656657).

De novo reconstruction of HD loci and genealogies

A two-step filtering process was implemented on the base-called sequences. In

the initial step, sequencing reads were filtered based on their Phred quality scores, with

reads having a mean quality score below 15 being removed. This ensured that the

remaining reads had an average per base accuracy of ≥ 97%. The second filtering step

involved selecting sequencing reads with lengths around the expected amplicon length

for each *HD* gene region: 1500-1800 bp for *bW-HD1* and 1300-1500 bp for *bE-HD2*. The

VSEARCH clustering algorithm (Rognes et al. 2016) was applied for quality control of

sequencing reads of each sample, using global similarity during clustering. It is

important to note that the VSEARCH clustering algorithm could not recognize

sequences of the same gene in the opposite direction. Hence, we obtained two clusters

for each *bW-HD1* and *bE-HD2* alleles in the sample along with two consensus

sequences - one forward and the other reverse. The analysis code is available on

Github   (https://github.com/TheRainInSpain/Lineage-Specific-Marker.git).   And   the

dataset is available on (https://doi.org/10.5281/zenodo.10656657). Geneious software

(v.2023.2.1) was used to visualize the forward read consensus sequences.

The reconstructed amplicon and reference sequences were aligned (Table 2)

with MUSCLE (v.5.1) (Edgar 2004), and simple genealogical trees were reconstructed

using the Geneious tree builder (v.2023.2.1).

Results

Specificity of the diagnostic assay achieved through HD loci amplification and ONT

sequencing

During the initial phase of primer testing, *HD* PCRs were performed on positive

controls of *A. psidii* (Au3, MF-1, APG1 and LFNJM1) and negative controls with non-

target rust species (*Miyagia psudosphaeria, Puccinia striiformis* f. sp. *tritici, P. graminis*,

P. triticina, Thekopsora minima). In addition, we focused on the individual PCR

amplicons (*bW-HD* and *bE-HD*) because the amplification of the full-length locus was

not robust enough across samples and technical replicates. The sizes of the amplicons

were as expected in the positive control samples, being of ~1600 bp for *bW-HD*, ~1400

bp for *bE-HD*, and ~3000 kb for the full HD locus (Table 2). In addition, we observed

bands of variable sizes in some of our technical repeats of non-target species. All

samples were sequenced with our ONT amplicon sequencing workflow because our

assay does not rely exclusively on the PCR amplification product but requires that the

amplified sequences match the *A. psidii HD* sequences in a genealogical framework.

None of the non-target species amplicon sequences matched the full-length *A.*

psidii HD sequences. For each *A. psidii* isolate, four *HD* alleles (2x *bW-HD1* and 2x *bE-*

HD2) were reconstructed based on ONT sequencing results, as expected for dikaryotic

organisms. Isolates were considered identical if their four *HD* alleles were identical or

carried minor non-functional variation (for example synonymous variation, variation

outside the variable domain or within introns). The applicability of the lineage diagnostic

test was confirmed by comparing the *de novo* reconstructed consensus sequences

derived from ONT amplicon sequencing with the *HD* amplicon sequences derived from

the reference genomes (APG1, MF-1, LFNJM1, Au3 and Apsidii_AM). All *de novo*

reconstructed ONT amplicon sequences clearly grouped with their respective *HD* alleles

obtained from reference genomes (Table 2, Figure 1A-B). Moreover, the reference

trees for *bW-HD1* and *bE-HD2* revealed that the Brazilian isolates and the South African

isolate carry clearly distinct alleles when compared against the pandemic isolate (Figure

1A-B), corroborating variations previously described using microsatellite markers

(Stewart et al. 2017; Roux et al. 2016; Graça et al. 2013).

HD-locus of _A. psidii_ primers are capable to detect isolates that belong to different

lineages from field samples

The designed primers successfully amplified the individual _HD_ loci of DNA

extracted from different sources of field samples, including urediniospores and infected

leaf material. As observed previously, the full _HD_ locus amplification was not possible

for the LFNJRM1 isolate and the field samples (Table 2). The reconstructed amplicon

and reference sequences were aligned and revealed a pairwise identity of 67.9% to

100% for _bW-HD1_ and 76.9% and to 100% for _bE-HD2_ alleles across the analyzed

sample isolates. Our results revealed that an isolate collected from field samples in

Australia in 2022 (SYD) belong to the same lineage as the pandemic lineage because

all four _HD_ alleles clearly grouped with those derived from the pandemic reference

isolate, while isolates collected from field samples from Brazil, belong to different

lineages having at least two different _HD_ alleles (Figure 1C-D).

Discussion

Traditionally, _A. psidii_ lineages have been identified using microsatellite markers

(Stewart et al. 2017; Graça et al. 2013; Roux et al. 2016; Sandhu et al. 2016). In this

study, a diagnostic assay targeting the _HD_ locus of _A. psidii_, coupled with ONT

sequencing was developed. This method successfully amplified diverse copies of the

HD locus from *A. psidii* isolates collected in Australia and Brazil and is predicted to work

for the South African isolate.

The genealogical tree of *de novo* reconstructed amplicons confirmed the

biological expectation of lineage specific variation among isolates originating from single

spores, field samples from different hosts, and distinct geographic locations. While the

isolate collected from field samples in Australia belong to the same lineage as the

pandemic lineage, in Brazil, the center of origin of the disease, there was a strong

association of *HD* allele status with their original host species, as previously observed

(Morales et al. 2023; Graça et al. 2013; Stewart et al. 2017). Overall, the high allelic

diversity at the *HD* locus in our samples is in line with the previously identified diversity

observed at this locus in rust fungi including *A. psidii* and *Puccinia* spp. (Ferrarezi et al.

2022; Holden et al. 2023; Luo et al., 2024).

While amplifying the entire *HD* locus in a single PCR reaction proved challenging

with the field samples, successful amplification of individual *HD* amplicons was achieved

for all samples. This allows for the isolate identification when analyzing the

reconstructed *HD* alleles from each amplicon (*bW-HD1* and *bE-HD2*). In the future, our

primers and/or amplification conditions can be further improved to amplify the entire

locus of natural *A. psidii* populations within its center of diversity, as it is more user

friendly to have one big region to amplify and ID, rather than two. In addition, we

observed off target PCR amplification in some of our technical replicates of non-target

species, yet none of these sequences match the *A. psidii HD* sequence. Improved

primers or nested PCR reactions could improve PCR specificity increasing sensitivity

and throughput when applied under routine testing conditions.

To date, the non-pandemic isolates shared at most one out of four *HD* alleles

with the pandemic isolate (Au_3). This finding shows non-pandemic isolates can be

differentiated from the pandemic lineage based on their *HD* alleles. This protocol serves

as a valuable tool in detecting new incursions of the pathogen in regions where a single

lineage is present as for example in Australia where only a single incursion of the

pandemic biotype has been reported to date. Further, comparing *HD* allele identity with

known reference sequences could link novel incursions with related populations in

source regions. This could help in identifying risks in import pathways of this exotic

pathogen and improve risk mitigation strategies. In addition, the assay could potentially

detect recombination between populations if purified single pustule isolates were

analyzed. In the future, we anticipate that the diagnostic test can be refined to detect

urediniospores of *A. psidii* from complex samples derived from air-sampling or mixed

infections to enable structured surveillance of this pathogen to detect changes or

incursions which will greatly assist the processing of samples in a surveillance setting.

Literature Cited

Baskarathevan, J., Taylor, R. K., Ho, W., McDougal, R. L., Shivas, R. G., and
Alexander, B. J. R. 2016. Real-Time PCR Assays for the detection of *Puccinia psidii*.
Plant Disease. 100:617-624.
Beenken, L. 2017. *Austropuccinia*: a new genus name for the myrtle rust *Puccinia psidii*
placed within the redefined family Sphaerophragmiaceae (Pucciniales). Phytotaxa.
297:53.
Bini, A. P. 2016. Estudo molecular do desenvolvimento de *Puccinia psidii* Winter *in vitro*
e   no   processo   de   infecção   em   *Eucalyptus   grandis*.   Available   at:
http://www.teses.usp.br/teses/disponiveis/11/11137/tde-10112016-161846/   [Accessed
September 19, 2022].

Boufleur, T. R., Morales, J. V. P., Martins, T. V., Gonçalves, M. P., Júnior, N. S. M., and

Amorim, L. 2023. A diagnostic guide for myrtle rust. Plant Health Progress. 24:242-251.

Carnegie, A. J., Lidbetter, J. R., Walker, J., Horwood, M. A., Tesoriero, L., Glen, M., et

al. 2010. *Uredo rangelii*, a taxon in the guava rust complex, newly recorded on

Myrtaceae in Australia. Australasian Plant Pathol. 39:463-466.

Coelho, M. A., Bakkeren, G., Sun, S., Hood, M. E., and Giraud, T. 2017. Fungal sex: the

Basidiomycota. Microbiol Spectr. 5.

Cuomo, C. A., Bakkeren, G., Khalil, H. B., Panwar, V., Joly, D., Linning, R., et al. 2017.

Comparative analysis highlights variable genome content of wheat rusts and divergence

of the mating loci. G3 Genes|Genomes|Genetics. 7:361-376.

Department of Agriculture, Fisheries and Forestry. 2023. Myrtle rust (exotic strains).

Biosecurity and Trade. Available at: https://www.agriculture.gov.au/biosecurity-

trade/pests-diseases-weeds/plant/myrtle-

rust#:~:text=Myrtle%20rust%20(exotic%20strains)%20is,damaging%20to%20our%20eu

calyptus%20trees [Accessed December 19, 2023].

Edgar, R. C. 2004. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. Nucleic Acids Res. 32:1792-1797.

Edwards, R. J., Dong, C., Park, R. F., and Tobias, P. A. 2022. A phased chromosome-

level genome and full mitochondrial sequence for the dikaryotic myrtle rust pathogen,

Austropuccinia psidii. :2022.04.22.489119 Available at:

https://www.biorxiv.org/content/10.1101/2022.04.22.489119v1 [Accessed August 22,

2022].

Ferrarezi, J. A., McTaggart, A. R., Tobias, P. A., Hayashibara, C. A. A., Degnan, R. M.,

Shuey, L. S., et al. 2022. *Austropuccinia psidii* uses tetrapolar mating and produces

meiotic spores in older infections on Eucalyptus grandis. Fungal Genetics and Biology.

160:103692.

Graça, R. N., Ross-Davis, A. L., Klopfenstein, N. B., Kim, M.-S., Peever, T. L., Cannon,

P. G., et al. 2013. Rust disease of eucalypts, caused by *Puccinia psidii*, did not originate

via host jump from guava in Brazil. Mol Ecol. 22:6033-6047.

Holden, S., Bakkeren, G., Hubensky, J., Bamrah, R., Abbasi, M., Qutob, D., et al. 2023.

Uncovering the history of recombination and population structure in western Canadian

stripe rust populations through mating type alleles. BMC Biology. 21:233.

Huelsenbeck, J. P., and Ronquist, F. 2001. MRBAYES: Bayesian inference of
phylogenetic trees. Bioinformatics. 17:754-755.

Hussain, K. K., Malavia, D., Johson, E., Littlechild, J. A., Winlove, P., Vollmer, F., et al.
2020. Biosensors and diagnostics for fungal detection. Journal of Fungi. 6:349.

Hyde, K. D., Al-Hatmi, A. M. S., Andersen, B., Boekhout, T., Buzina, W., Dawson, T. L.,
et al. 2018. The world's ten most feared fungi. Fungal Diversity. 93:161-194.

IPPC. 2018. DP 26: *Austropuccinia psidii* .

Leite, T. F., Moon, D. H., Lima, A. C. M., Labate, C. A., and Tanaka, F. A. O. 2013. A
simple protocol for whole leaf preparation to investigate the interaction between
Puccinia psidii and *Eucalyptus grandis*. Australasian Plant Pathol. 42:79-84.

Luo, Z; McTaggart, A; Schwessinger, B. 2024. Genome biology and evolution of matyng
type loci in four cereal rust fungi. Available at:
https://www.biorxiv.org/content/10.1101/2023.03.02.530769v3.

Makinson, R. O., Pegg, G. S., and Carnegie, A. J. 2020. Myrtle Rust in Australia - a
National Action Plan.

McTaggart, A. R., Duong, T. A., Le, V. Q., Shuey, L. S., Smidt, W., Naidoo, S., et al.
2018. Chromium sequencing: the doors open for genomics of obligate plant pathogens.
BioTechniques. 65:253-257.

Morales, J. V. P., Boufleur, T. R., Gonçalves, M. P., Parisi, M. C. M., Loehrer, M.,
Schaffrath, U., et al. 2023. Differential aggressiveness of *Austropuccinia psidii* isolates
from guava and rose apple upon cross-inoculation. Plant Pathology. Available at:
https://onlinelibrary.wiley.com/doi/abs/10.1111/ppa.13850 [Accessed January 12, 2024].

Ranwez, V., Douzery, E. J. P., Cambon, C., Chantret, N., and Delsuc, F. 2018. MACSE
v2: Toolkit for the Alignment of Coding Sequences Accounting for Frameshifts and Stop
Codons. Mol Biol Evol. 35:2582-2584.

Rognes, T., Flouri, T., Nichols, B., Quince, C., and Mahé, F. 2016. VSEARCH: a
versatile open source tool for metagenomics. PeerJ. 4:e2584.

Roux, J., Granados, G. M., Shuey, L., Barnes, I., Wingfield, M. J., and McTaggart, A. R.
2016. A unique genotype of the rust pathogen, *Puccinia psidii*, on Myrtaceae in South
Africa. Australasian Plant Pathol. 45:645-652.

Sandhu, K. S., Karaoglu, H., Zhang, P., and Park, R. F. 2016. Simple sequence repeat

markers support the presence of a single genotype of *Puccinia psidii* in Australia. Plant

Pathol. 65:1084-1094.

Santos, A. P., Gomes, R. L., Furtado, E. L., Passos, J. R. S. 2020. Quantifying losses in

productivity by the rust in eucalypt plantations in Brazil. Fores. Ecol. Manag. 468:

118170.

Simpson, J. A., Thomas, K., and Grgurinovic, C. A. 2006. Uredinales species

pathogenic on species of Myrtaceae. Australas. Plant Pathol. 35:549-562.

Stewart, J. E., Ross-Davis, A. L., Graça, R. N., Alfenas, A. C., Peever, T. L., Hanna, J.

W., et al. 2017. Genetic diversity of the myrtle rust pathogen (*Austropuccinia psidii*) in

the Americas and Hawaii: Global implications for invasive threat assessments ed. M.

Cleary. For. Path. 48:e12378.

Wilson, A. M., Godlonton, T., van der Nest, M. A., Wilken, P. M., Wingfield, M. J., and

Wingfield, B. D. 2015. Unisexual reproduction in *Huntiella moniliformis*. Fungal Genetics

and Biology. 80:1-9.

Winter, G. R. 1884. Rabenhorstii fungi europaei et extraeuropaei exsiccati cura Dr. G.

Winter, CenturiaXXXI et XXXII. Hedwigia. 23:164-172.

Acknowledgments

393 The authors thank Danièle Giblot-Ducray and Kelly Hill for the critical feedback and

394 support.

Funding

A. T. Gonzalez, R. Kimber, and B. Schwessinger were supported by an Australian

Government grant from the Department of Agriculture, Fisheries and Forestry entitled

"Automated air sampling for remote surveillance and high throughput processing of

environmental samples for eDNA analyses". São Paulo Research Foundation

(FAPESP) Grant 2019/13191-5. T. R. Boufleur was supported by FAPESP Grants

2022/11900-1 and 2021/01606-6.

405
406   **Table 1.** List of primers used in the present study.

Primers	Sequence (5' - 3')	Region	Changes
HDFor2DG (forward)[a]	ATACAGTTYAGGTTWTRGCG	*bW-HD*	C/T, A/T, A/G
HDMR1 (reverse)	GAAAGGAAATATTGCCACT	*bW-HD*	-
HDMF2 DG (forward)	YGACCGCCTTCCTTTGAG	*bE-HD*	C/T,
HDRev2DG (reverse)[a]	GTGTCSAAGCWACCAAAATC	*bE-HD*	C/G, A/T

407   [a] Primers used to amplify the full *HD* locus.

408   **Table 2.** Description of isolates and populations and of *A. psidii* and non-target rusts used in this study

Organism	Code	Source	Country	Amplification[a]			Alleles			
				Full locus	*bW-HD1*	*bE-HD2*	*HD1*		*HD2*	
Isolates										
Austropuccinia psidii	Au3[b]	*Agonis flexuosa*	AUS	+	+	+	Au3_HapA	Au3_HapB	Au3_HapA	Au3_HapB
A. psidii	MF-1[b]	*Eucalyptus grandis*	BR	+	+	+	MF-1_HapA	MF-1_HapB	MF-1_HapA	MF-1_HapB
A. psidii	APG1 (GM1)	*Psidium guajava*	BR	-	-	+	NA	NA	APG1_HapA	NA
A. psidii	LFNJM1[b]	*Syzygium jambos*	BR	+	+	+	LFNJM1_HapA	LFNJM1_HapB	LFNJM1_HapA	LFNJM1_HapB
A. psidii	LFNJM3	*S. jambos*	BR	+	+	-	LFNJM3_HapA	NA	NA	NA
A. psidii	LFNJM4	*S. jambos*	BR	+	+	+	LFNJM4_HapA	LFNJM4_HapB	LFNJM4_HapA	LFNJM4_HapB
A. psidii	LFNJRM1	*S. samarangense*	BR	-	+	+	LFNJRM1_HapA	NA	LFNJRM1_HapA	LFNJRM1_HapB
A. psidii	Apsidii_AM	*S. jambos*	SA	NA	NA	NA	SA_HapA	SA_HapB	SA_HapA	SA_HapB
Populations (urediniospores)										
A. psidii	CA	*Plinia edulis*	BR	-	+	+	CA_Hap1	CA_Hap2	CA_Hap1	CA_Hap2
A. psidii	CG	*Eugenia dysentericus*	BR	-	+	+	LFNSDP1_HapA	LFNSDP1_HapB	LFNSDP1_HapA	LFNSDP1_HapB
Populations (infected leaves)										
A. psidii	LFNEP1	*Eugenia stipitata*	BR	-	+	+	LFNEP1_HapA	NA	LFNEP1_HapA	LFNEP1_HapB
A. psidii	3.1	*Pimenta dioica*	BR	-	+	+	3.1_HapA	NA	3.1_HapA	3.1_HapB
A. psidii	SYD	*Malaleuca quinquenervia*	AUS	-	+	+	SYD_HapA	SYD_HapB	SYD_HapA	SYD_HapB
Non-target rust species										
Miyagia psudosphaeria	-	-	-	NS	NP	NA	NA	NA	NA	NA
Puccinia triticina	-	-	-	NS	NS	NA	NA	NA	NA	NA
P. graminis f. sp. *avenae*	-	-	-	NS	NS	NA	NA	NA	NA	NA
P. striiformis f. sp. *tritici*	-	-	-	NS	NS	NA	NA	NA	NA	NA
Thekopsora minima	-	-	-	-	NS	NA	NA	NA	NA	NA

[a] Conventional polymerase chain reaction (PCR) with the three sets of primers developed in the present study. Full *HD*

locus, HDFor2DG/HDRev2DG amplifies a fragment of 3000 kb; *bW-HD,* HDFor2DG/HDMR1 amplifies a fragment of 1500

kb; *bE-HD,* HDMF2/HDRev2DG amplifies a fragment of 1400 kb; +, positive; -, negative.

[b] *A. psidii* and non-target rust isolates used to confirm the specificity of the designed primers.

"NS" is defined as NS specific amplification in at least one of the technical replicates, "NA" as not available, "-" as not

amplification,   "+"   as   amplification   of   the   expected   size   amplicon   product.

Figure 1: Bayesian inference genealogical trees reconstructed from *HD* (homeodomain) genes

of *Austropuccinia psidii*. Genealogical trees reconstructed from the alignment of two *bW-HD1*

(A, C) or two *bE-HD2* **(B, D)** alleles per dikaryotic genome assembly of reference isolates **(A-B)**,

and with consensus sequences for each allele *de novo* reconstructed from ONT long-read

amplicons of *bW-HD1* and *bE-HD2* (C-D). Lineage identification is indicated by the combination

of different colors in the bottom. Reference sequences are indicated in bold (C-D). Bayesian

posterior probabilities are shown on the nodes.