A circadian transcriptional sub-network and *EARLY FLOWERING 3* control timing of senescence and grain nutrition in bread wheat

Christopher R. Buckley[1], Joshua M. Boyte[1], Robert L. Albiston[1], Jessica Hyles[2], Jesse Beasley[1], Alexander AT Johnson[1], Ben Trevaskis[2], Alexandre Fournier-Level[1], Michael J. Haydon[1] *

[1]School of BioSciences, University of Melbourne, Parkville, VIC 3010, Australia
[2]CSIRO Agriculture and Food, GPO Box 1700, Canberra, ACT 2601, Australia
* Corresponding author: m.haydon@unimelb.edu.au

Abstract

Circadian clocks control daily and seasonal timing of physiology and development. Because of their influence on photoperiodic flowering, variants in circadian clock genes have been selected for phenology during domestication of cereal crops. To explore the potential impact of this genetic variation on circadian-regulated traits, we investigated the relationship of the circadian clock and leaf senescence in hexaploid bread wheat. Phenotyping of a collection of elite wheat cultivars identified significant variation in circadian rhythms which was associated with timing of senescence and nutrient mobilisation efficiency. RNA sequencing revealed substantial reorganisation of the circadian-regulated transcriptome during senescence and a transcriptional sub-network representing a link between the circadian oscillator and regulators of leaf senescence. We used genotypes of multiple circadian clock genes to assign cultivars to 'chronotypes', which could be used to predict circadian-regulated phenotypes. This identified a deletion variant of *EARLY FLOWERING 3-D1 (ELF3-D1)* attributed to a phenology locus, *Earliness per se (Eps-D1),* and we used near-isogenic lines (NILs) to show that it affects timing of senescence and grain protein content (GPC). Thus, there are potential consequences of circadian clock genes selected for phenology on other valuable crop traits.

Keywords

Introduction

Leaf senescence is a controlled process involving the degradation of chlorophyll, downregulation of photosynthesis and the transport of remobilised nutrients to nascent tissues[1]. Efficient mobilisation of nutrients from vegetative to reproductive tissues during leaf senescence is a key determinant of plant fitness. The relative timing of senescence affects a trade-off between resource accumulation and usage[2] and can be adjusted to balance these processes for adaptation to different environments[3–5]. In cereal crops, the timing of senescence has a major impact on yield traits. In bread wheat (*Triticum aestivum*), high grain protein content (GPC) is a valuable trait which can be achieved with earlier senescence, but typically comes at a cost of lower yield[6]. However, these relationships can be environment-dependent[7–9]. Increasing nutrient content without yield loss across multiple environments is a priority target for crop breeding.

The circadian clock is a biological timing mechanism, comprised of a highly influential gene regulatory network that functions to synchronise development and physiology with daily and seasonal environmental cycles. Plant circadian clocks heavily influence metabolism, transport and signalling underlying growth and development[10]. In numerous cereal crops, the role of the circadian clock in photoperiodic flowering has led to widespread selection of circadian clock gene variants[11]. However, the impact of these variants on other circadian-controlled traits has not been explored.

Since the circadian clock influences timing of senescence in *Arabidopsis thaliana*[12–14], we sought to explore the role of the circadian clock in wheat in controlling leaf senescence and determining grain nutrient content. We compiled a panel of 25 wheat cultivars and found these traits covaried with phenotypic and genotypic variation in the circadian clock. We compared the circadian transcriptomes of mature and senescent flag leaves and used genotypic information for circadian clock genes to define chronotypes within the wheat panel. These suggested a role for the clock gene *EARLY FLOWERING 3-D1* (*ELF3-D1*) and using near isogenic lines (NILs) of a widespread deletion variant in *ELF3-D1,* we found a functional contribution to timing of senescence and GPC.

Results

Circadian period and timing of senescence covary in Australian wheat cultivars

Since the circadian clock influences leaf senescence in Arabidopsis [13–15] and the pace of circadian rhythms shortens with age [16,17], we investigated the relationship between phenotypic variation of circadian rhythms in wheat and the timing of senescence. Using delayed leaf fluorescence (DF) (Fig. 1A), we first measured circadian rhythms in the second leaf of three wheat cultivars 14 days after sowing (DAS; seedling), 33 DAS (5-leaf stage) and 44 DAS (end of tillering). Relative amplitude error (RAE; a proxy for rhythm robustness) did not significantly differ by leaf age (Fig. 1B), but circadian period was significantly shorter in leaves of older plants (Fig. 1C). This demonstrates that changes occur in the circadian clock of wheat plants during ageing.

To assess the extent of phenotypic variation of circadian rhythms in wheat, we chose 15 elite cultivars grown across diverse environments of Australia and measured circadian rhythms of DF in glasshouse-grown plants in two years. Significant differences in mean circadian period were detected between cultivars in each year (Fig. 1D, Supplementary Fig. 1A). In year 1, mean circadian period differed by over four hours over the panel (Fig. 1D,E). Circadian period was mostly consistent within cultivars over both years, except for Viking and Scout, which might suggest strong gene-by-environment interactions in these cultivars (Supplementary Fig. 1B). Nevertheless, there is robust variation in circadian rhythms within elite wheat germplasm which could influence circadian-regulated agronomic traits.

To measure the timing of senescence in the 15 wheat cultivars, we monitored the decline of chlorophyll content in flag leaves after anthesis. There were significant differences in the timing of senescence between cultivars (Fig. 1F), with a maximum mean difference of 10.3 days. We also measured total ion content of mature flag leaves and harvested grain to calculate nutrient mobilisation efficiency. We detected substantial variation in mobilisation efficiency between cultivars (Fig. 1G;) and observed a strongly negative correlation with timing of senescence (Supplementary Fig. 2). Timing of senescence and nutrient mobilisation efficiency were both significantly correlated with circadian period in year one (Fig. 1H,I), such that cultivars with shorter period tend to senesce earlier and exhibit more efficient leaf-to-grain nutrient mobilisation. The relationship of these traits to circadian period was weaker in year two due to a strong year-effect for Scout and Viking. Without these cultivars, a stronger correlation was detected (Supplementary Fig. 1C,D). We also measured total GPC, an important trait which

Figure 1: Circadian period of wheat cultivars is associated with senescence and grain nutrition. A, Overview of delayed fluorescence (DF) workflow for measuring wheat circadian rhythms. **B,** Relative amplitude error (RAE) and **C**, circadian period of DF rhythms in 2nd leaf of three cultivars at 14, 33 and 44 days after sowing (DAS). Letters indicate significant differences as determined by one-way ANOVA followed by Tukey's HSD; $P < 0.05$. Error bars represent ±sd, n=15. **D,** Circadian period of DF in 15 Australian wheat cultivars in year one. Letters indicate significant differences as determined by Welch's t-test, $P < 0.05$. Error bars represent ±sd, n=8. **E,** Circadian rhythms of DF in the Supreme and Viking. Error bars represent ±sd, n=8. **F,** Time to senescence, (days after anthesis; DAA) in 15 wheat cultivars. Letters indicate significant differences as determined by one-way ANOVA followed by Tukey's HSD; $P < 0.05$. Error bars represent ±sd, n=6. **G,** Total nutrient remobilisation efficiency (ratio of leaf-to-seed nutrient content, see methods) in 15 wheat cultivars. **H**, Mean circadian period (year one) versus mean time to senescence and **I,** total remobilisation efficiency in 15 wheat cultivars. Significance assessed by Pearson's product-moment correlation.

depends on nutrient mobilisation (Supplementary Fig. 3A). GPC was significantly negatively correlated with circadian period in year two, but not year one (Supplementary Fig. 3B,C). Overall, our data suggest strong relationships between circadian period, leaf ageing and senescence, nutrient mobilisation efficiency and GPC in elite wheat cultivars.

Reshaping of the circadian transcriptome during leaf senescence

To explore transcriptional changes in the circadian network during leaf senescence we compared the transcriptomes of mature and senescent flag leaves in *cv.* Mace over 48 h in continuous light (Fig. 2A). Principal component analysis (PCA) separated each individual transcriptome by developmental stage (Supplementary Fig. 4A) and time of day (Supplementary Fig. 4B-E). We used BIO_CYCLE[18] and MetaCycle[19] to detect rhythmic transcripts and estimate their period, phase and amplitude. A total of 11,705 expressed transcripts were called rhythmic in mature flag leaves (15.7% of 74550 expressed) by both algorithms and 16,906 transcripts were rhythmic in senescent flag leaves (22.8% of 74272 expressed) (Fig. 2B). Since fewer rhythmic genes were detected in mature flag leaves compared to senescent flag leaves, the changes in the circadian transcriptome during senescence are unlikely to be due to a general decay in gene expression, we performed gene ontology (GO) enrichment analysis of transcripts that are rhythmic only in senescent flag leaves. Significantly enriched terms included "protein transport", "peptide metabolic process" and "nitrogen compound transport" (Fig. 2C), suggesting that the circadian network is connected to the reprogramming of metabolism and transport during senescence.

To identify potential regulators of the circadian transcriptome during leaf senescence, we compared periodicity (BIO_CYCLE's measure of rhythmicity, from 0 to 1) across families of transcription factors (TFs)[20]. *PSEUDO RESPONSE REGULATORs* (*PRRs*), which are core circadian oscillator genes[21], were the most rhythmic family in both tissue types, followed by Sigma70-like TFs, which regulate rhythmic chloroplast gene expression in Arabidopsis (Fig. 2D)[22]. Mean periodicity of the *WRKY* and *CCAAT-HAP2* families was significantly higher in senescent tissue than mature tissue (Fig. 2D). To summarise the overall changes in TF gene expression, we calculated metagene expression values of entire TF gene families (Fig. 2E) and observed a clear increase of *WRKY* metagene rhythmicity during senescence. By contrast, a similar change in rhythmicity was not observed for other TF families with roles in senescence (e.g. *NAC, AP2* and *MYB*[23]). However, we did detect a phase advance of metagenes for the *MYB, PRR* and *Sigma70* families (Fig. 2E). *WRKY* genes are known as regulators of leaf senescence[24] but our data suggests the influence of the circadian clock on their regulation might have been underestimated.

Among the 8,593 rhythmic transcripts in both tissues, the mean circadian period was ~0.5 h shorter in senescent flag leaves compared to mature flag leaves (Fig. 3A; Supplementary Fig. 5B). This is consistent with the expectation for shorter period in older leaves (Fig. 1C). In addition,

Figure 2: Circadian transcriptional network increases its output at leaf senescence. A,
Sampling design of comparative circadian senescence transcriptome. 'Sen' and 'Mat' refer to
the sowing dates of plants that were harvested for senescent and mature flag leaves,
respectively. **B,** Number of transcripts determined to be rhythmic by BIO_CYCLE[18] and
MetaCycle[19]. **C,** Gene ontology term enrichment analysis of transcripts that are uniquely
rhythmic in senescent flag leaves. **D,** Comparison of mean periodicity of transcription factor
families determined by BIO_CYCLE (0 = arrhythmic). Asterisks indicate significant differences
as determined by Welch's t-test, * = $P < 0.05$, *** = $P < 0.001$. **E,** Metagene expression (average
across entire gene family) of selected transcription factor families.a circadian phase advance of
~0.5 h (corrected for period) was also detected in senescent leaves (Supplementary Fig. 5A,B).
Amplitude differences were small between the two developmental stages, although mean
amplitude was higher in mature tissue (Supplementary Fig. 5C).

we noted that there was substantial variation in the period differences for individual transcripts
outside the global trend (Fig 3A). To identify patterns in differential period, we performed GO term
enrichment of transcripts with shorter (< -0.5 h), longer (> 0.5 h) or unchanged (< ±0.5 h) circadian
period in senescent leaves. Rhythmic genes with shorter period were significantly enriched for
GO categories related to primary metabolism (Fig. 3B,C). Among transcripts with longer period in

senescent tissue, terms related to photosynthesis were overrepresented, perhaps reflecting the
breakdown of normal rhythmic regulation of photosynthesis (Fig. 3B,C; Supplementary Fig. 3).
The two most strongly enriched GO terms within transcripts with unchanged period were "rhythmic
process" and "circadian rhythm" (Fig. 3B,C). This suggests that the global shortening of period in
the senescent transcriptome could be driven by amplitude or phase changes in the circadian
oscillator.

To identify other transcriptional changes between mature and senescent leaves, we used
LimoRhyde[25] to identify differentially expressed (DE) and differentially rhythmic (DR) genes.
Among DE genes in senescent leaves, 6,310 upregulated genes were enriched for GO terms for
transport and catabolism, and 8,785 downregulated genes were enriched for GO terms related to
biosynthesis (Supplementary Fig. 6). There were 1,743 DR genes, enriched for GO terms related
to photosynthesis, light signalling and circadian rhythms (Fig. 3D). When we compared DR genes
to the remainder of the rhythmic transcriptome, we observed an advanced mean phase in
senescent tissue, rather than change in period or amplitude (Fig. 3E).

A circadian-regulated sub-network in senescent leaves
We hypothesised that the phase-advanced DR genes, which are enriched for the "circadian
rhythm" GO term, represent a sub-network driving changes in the transcriptome. We identified
106 TFs among the DR genes, which were characterised by large phase advances
(Supplementary Fig. 7A). Several families of TFs were significantly over-represented, including
C2C2-Dof, TAZ, FHA, PRR and Sigma70 (Supplementary Fig. 7B). To analyse the interactions
between DR TFs, we modelled the putative interactions between members of this gene set using
STRING[26]. We identified 308 putative functional and physical interactions, which is 7x more than
expected (Hypergeo. test, $P < 1\times10^{-16}$). The network includes genes related to hormone signalling,
light signalling and photoperiodic flowering, and is centred around core clock genes; *PRRs*,
REVEILLEs (*RVEs*), *LUX ARRHYTHMO* (*LUX*) and *LATE ELONGATED HYPOCTYL* (*LHY*) (Fig.
3F). Evening elements (EEs), which are *cis*-regulatory motifs bound by clock TFs[27], were highly
significantly enriched among all DR gene promoters (Supplementary Table 1). These results
suggest that there is a circadian-controlled transcriptional subnetwork with advanced phase in
senescent tissue.

Figure 3: A circadian-regulated transcriptional subnetwork in senescent leaves. A,
Distribution of circadian period values of all rhythmic transcripts in mature and senescent flag
leaves. Asterisks indicate a significant difference between the mean circadian period values,
Welch's *t*-test, *P* < 0.001. **B,** GO term enrichment analysis of transcripts categorised according to
the change in circadian period between developmental stages; short period (Δperiod < -0.5 h),
long period (Δperiod > 0.5 h), unchanged period (Δperiod < ±0.5 h). O (observed) and E
(expected) number of transcripts. **C,** Proportions of transcripts in each Δperiod category for a
selection of GO terms. **D,** GO term enrichment analysis of differentially rhythmic (DR) genes. **E,**
Distribution of changes in circadian rhythm parameters for DR genes. Asterisks indicate
significant differences between DR genes and all other rhythmic genes as determined by Welch's
t-test, * = *P* < 0.05, *** = *P* < 0.001, ns = not significant. **F,** Putative protein-protein interaction
network of DR transcription factors. Combined score reflects the probability of interaction
determined by STRING[26]. Related processes are manually assigned after literature and orthology
database searches.

We performed a systematic analysis of circadian oscillator gene expression and identified 59 out
of 79 putative circadian oscillator genes (Supplementary Table 2) that were rhythmic in both

tissues (Supplementary Fig. 8). Mean period difference of this group was 0.52 h, which was similar to the global trend, and suggested that maintenance of period is not uniform across the oscillator (Fig. 3B). Strikingly, the mean phase advance of oscillator genes was 1.37 h, which is significantly larger than the global average (Welch's t-test, t (59.5) = 4.21, P < 0.001). To categorise the oscillator genes by rhythmic expression patterns, we performed PCA of changes in period, phase and amplitude (Fig. 4A). The three PCs were grouped into six transcript clusters of 1-20 genes by k-means (Fig. 4C). Clusters 5 and 6 are mainly comprised of genes expressed around dawn, including *RVEs*, *NIGHT LIGHT–INDUCIBLE AND CLOCK-REGULATED* (*LNKs*) and *LHY* genes (Fig. 4C) and have shorter period in senescing tissue, consistent with the global trend (Fig. 4B,D). Clusters 2, 3 and 4 have advanced phase in senescing tissue (Fig. 4B), similar to the identified subnetwork (Fig. 3F). These are mostly afternoon- or evening-phased genes and include the majority of *PRR* and *LUX* genes (Fig. 4C). Amplitude was increased in cluster 2 genes, and period was unchanged in clusters 2 and 4 (Fig. 4B,D). Thus, there appear to be distinct changes in oscillator genes during senescence according to their phase of expression and this suggests a restructuring of the circadian oscillator during leaf senescence.

Identification of chronotypes within elite wheat cultivars

Since the circadian clock appears to strongly influence the wheat transcriptome during leaf senescence, we sought to maximise the circadian gene variation within the wheat panel for further phenotyping. We explored SNP data for the OzWheat diversity panel comprised of 283 Australian cultivars. We found four novel non-synonymous SNPs in conserved regions of clock genes from distinct transcript clusters (Fig. 3C); *TaPRR59-B1* (Cluster 2), *TaLUX-B1* (Cluster 3), *TaTOC1-B1* (Cluster 4) and *TaPRR73-A1* (Cluster 6). We used these SNPs and three major polymorphic clock gene loci in wheat (*TaPpd-B1*, *TaPpd-D1* and *TaELF3-D1*[28–30]), to categorise cultivars into six chronotypes (Fig. 5A,B; Supplementary Table 3) and compiled a circadian diversity panel of 25 cultivars, representing all six chronotypes (Supplementary Table 4), which were phenotyped for circadian period and timing of senescence (Fig. 5E,F; Supplementary Fig. 9). There were significant differences between chronotypes for both circadian period and time to senescence (Fig. 5C,D). In turn, grain filling was also affected, as thousand grain weight (TGW) differed significantly between chronotypes (Supplementary Fig. 10A,B). However, there were no significant differences between chronotypes for GPC (Supplementary Fig. 10C,D). Although the circadian period values do not correlate with time to senescence and TGW for all chronotypes, chronotype 5 exhibited short period, early senescence and low TGW, while chronotype 6 showed both longer period and later senescence (Fig. 5C,D; Supplementary Fig. 10).

Figure 4: Reshaping of the circadian oscillator during leaf senescence. A, PCA of changes in circadian period, phase and amplitude between mature and senescent leaves for putative circadian oscillator genes. PCs generated were used to group transcripts into 6 clusters (represented by different colours). **B,** Mean change in circadian period, phase and amplitude for each transcript cluster. **C,** Names of genes within each transcript cluster. **D,** Transcript circadian rhythms for a representative gene from each cluster. Error bars represent ±sd.

ELF3-D1 **deletion impacts timing of senescence and GPC**

Chronotypes 2 and 5 both have a deletion in *TaELF3-D1* and relatively early senescence. To test the potential for this circadian polymorphism on timing of leaf senescence, we used four pairs of near-isogenic lines (NILs) that differ by variation at *TaELF3-D1* (Supplementary Table 3,4). Across the four pairs of NILs circadian period was not significantly different (Fig. 5G,H) and neither was TGW (Fig 5I,J). However, early senescence was detected for the *TaELF3-D1* deletion in three out of four NIL pairs (Fig. 5K,L). Similarly, GPC was reduced by the *TaELF3-D1* deletion in

267
Figure 5: Wheat clock gene variation affects time to senescence and GPC. A, Cultivars
from the OzWheat diversity panel were grouped into chronotypes by *k*-modes clustering based
on their genotype at seven clock gene markers. Multiple correspondence analysis was used to
visualise clusters in space. **B,** Mode allelic variation of each cluster (see Supplementary Table 3
for explanation of alleles). **C,** Circadian period (n=10) and **D**, time to senescence (n=5) of 25
wheat cultivars in a circadian diversity panel. Error bars represent ±sd. **E,** Circadian period and
F, time to senescence for the 25 cultivars by chronotype defined in **A**. Letters indicate significant
differences as determined by one-way ANOVA followed by Tukey's HSD; *P* < 0.05. **G,H,**
Circadian period, **I,J,** thousand grain weight (TGW), **K,L,** time to senescence and **M,N,** grain
protein content (GPC) of near isogenic lines (NILs) for variation at *TaELF3-D1*. Individual NIL
pairs are shown in **G,I,K,M.** Error bars represent ±sd, n=5 . Values for all four NILs with
functional (*WT*) and *ELF3-D1del (del)* are combined in **H,J,L,N.** Asterisks indicate significant
differences between *WT* and *del* alleles as determined by Welch's *t*-test, * = *P* < 0.05, *** = *P* <
0.001, ns = not significant.

the same three NIL pairs (Fig. 5M,N). The exception is NIL pair c, which carries a different
genotype for *TaPpd-D1* (Supplementary Table 5), a putative downstream target of *ELF3*. Thus,
the absence of an effect in this NIL pair could be explained by an epistatic interaction.
285
286   **Discussion**
287
We have used a diverse collection of elite wheat cultivars and comparative transcriptomics to
explore the influence of the circadian clock on leaf senescence in wheat and assess its impact on
grain quality. Our data reveal significant phenotypic variation in circadian rhythms within wheat
cultivars and identify strong correlations with timing of leaf senescence, nutrient mobilisation
efficiency and GPC. Comparison of the circadian transcriptome in mature and senescent leaves
identified a circadian-controlled subnetwork, centred around several core clock genes and
connected to possible transcriptional regulators of senescence. Using genotypic information for
circadian clock genes in these wheat cultivars we were able to define chronotypes which we show
could potentially be used to predict circadian phenotypes and yield traits. Phenotyping of NILs for
an extant deletion variant in *TaELF3-D1* revealed a functional contribution of this core circadian
clock gene to timing of senescence and GPC.
299
ELF3 is a core component of the circadian oscillator as part of the evening complex (EC), a
transcriptional repressor complex also comprised of ELF4 and LUX in Arabidopsis[31]. Null mutants
in *ELF3* in Arabidopsis and double null mutants in tetraploid wheat have arrhythmic circadian
phenotypes[30,32]. Hypomorphic alleles of *elf3* in Arabidopsis have short period phenotypes[33,34].
Chronotypes 2 and 5 have a *TaELF3-D1* deletion, and chronotype 5 had a relatively short period
(Fig 5). Since we did not detect significant differences between the *TaELF3-D1* NIL pairs, it seems
unlikely that this variant alone causes the short period phenotype. Notwithstanding that the
TaELF3-D1 deletion might be recessive to functional A and B homoeologues for circadian period,
there are several genotypic differences in clock genes between these two chronotypes, so the
short period of chronotype 5 could be due to genetic interactions between different clock alleles.
For example, chronotypes 2 and 5 have different alleles at *TaPpd-D1* and *TaPpd-B1*, which are
putative regulatory targets of the EC[35].
312
We found a significant effect of the *TaELF3-D1* deletion on timing of leaf senescence and GPC
for three out of four NIL pairs (Fig 5). This suggests that the widespread *TaELF3-D1* deletion,
attributed to *Earliness per se (Eps-D1)* locus[30,36], has a potential negative effect on grain quality,

independent of any effect on circadian period. The exception was NIL pair c, which has a promoter
deletion within *TaPpd-D1,* in contrast to the functional alleles present in the other three
(Supplementary Table 5). Thus, the effect of *TaELF3-D1* on senescence might depend on the
genotype at *TaPpd-1.* Notably, *TaPpd-B1* is among the *PRR* genes identified within the
transcriptional subnetwork (Fig. 3) which we predict to make a regulatory contribution to leaf
senescence.

Increased GPC is commonly associated with reduced yield in cereal crops[6] and this relationship
has been attributed to timing of senescence[2]. TGW, which is an important yield component, was
not significantly different between *TaELF3-D1* NILs (Fig 5), but accurate estimates will require
multi-year field trials. These will also be required to better assess the robustness of the
relationship between circadian period and timing of senescence, since we detected substantial
gene-by-environment effects for some cultivars (Supplementary Fig. 1).

The overall trend in the rhythmic transcriptome was of shorter circadian period in senescent tissue
(Fig. 3) which is consistent with expectations of accelerating rhythms with age[16,17] (Fig 1).
However, there was substantial variation across the transcriptome. For example, photosynthesis
genes had a longer period, which could be explained by general decay in expression and an
uncoupling from the core oscillator. Of particular note, genes that were relatively unchanged for
period were enriched for circadian genes and instead are characterised by an advanced phase
(Fig 3). Indeed, three of the circadian transcript clusters we defined had advanced phase in
senescent tissue, and include numerous *PRR* genes, EC genes and *GIGANTEA* (Fig 4). We
propose that the phase adjustment of these oscillator genes could be driving the period change
of the wider transcriptome. Interestingly, *ELF3* genes and *TaPpd-D1* are among the few oscillator
genes that have a substantially shorter period in senescent leaves and might represent a
regulatory link between the adjusted oscillator and senescence-regulated genes.

Our findings have potentially important implications for breeders and growers because they
suggest that extant variation in circadian clock genes in crops, which were selected for their
effects on phenology, can have a significant impact on other traits such as grain quality. The
extent of these effects is likely to depend on genetic interactions between other circadian loci and
so identifying chronotypes within wheat germplasm could be a useful tool to predict phenotypic
effects. The TFs within the subnetwork are candidates that could be targeted to fine-tune
circadian-regulated leaf senescence without affecting the core circadian clock.

Materials and methods

Plant materials

Phenotyping experiments in year one and two used a panel of 15 elite Australian wheat cultivars (Calingiri, Cobra, Corack, Gladius, Hydra, Mace, Magenta, Scout, Sunguard, Suntop, Supreme, Viking, Wyalkatchem, Yitpi and Zen) which were obtained from Australian Grains Genebank (AGG). In the third and fourth years, a 25-cultivar circadian diversity panel was developed by including cultivars from the OzWheat diversity panel developed by the Commonwealth Scientific and Industrial Research Organisation (CSIRO). These additional cultivars were Axe, Bayonet, Buckley, Cascades, EGA Wedgetail, Insignia, Lillimur, Takari, Wilgoyne and Yarralinka.

Four pairs of NILs with variation at *TaELF3-D1* were derived from a large NIL population with phenology gene variation developed by CSIRO[37,38]. The NILs were generated by crossing donors of phenology genes with the cultivar Sunstate, which carries the *TaELF3-D1del* allele. This was followed by five rounds of recurrent backcrossing to obtain average 97% genetic identity between NILs.

Growth conditions

For glasshouse experiments, seeds were sown in July-August (Parkville, Victoria). Growth room and growth cabinet experiments were set to 12 h light (200-280 µmol m^{-2} s^{-1}, 20ºC), 12 h dark (15ºC). In all phenotyping experiments, seeds were chilled at 4°C for 48 h and sown in a mix of sterilised potting soil with 4.5 g/L Osmocote 6-9 month slow-release fertiliser. Full details of growth conditions can be found in Supplementary Note 1.

Delayed leaf fluorescence (DF) imaging

Unless specified otherwise, DF was imaged in the second leaves of plants at the 5-leaf stage. In all experiments, 2.5 cm sections were dissected from excised leaves (cut 3 cm from the leaf tip) and placed on 12 cm^2 0.5% agar (type M; Sigma-Aldrich) Petri plates, arranged according to a randomised controlled block design. Leaf sections were imaged using a Retiga LUMO charge-coupled device (CCD) camera (Teledyne Photometrics) fitted with a 25 mm f/0.95 lens housed within light-tight dark boxes. Light (>45 µmol m^{-2}s^{-1}) was provided by two LB3 red/blue LED panels (Photek). The lights and camera were controlled by beanshell (bsh) scripts in µManager (version 2.0[39]). DF was measured in 120 h of continuous light (LL) conditions by 1 min camera exposure in the dark, following 59 min of light. Camera properties were binning = 4x4, gain = 1, readout

rate = 0.650195 MHz 16 bit. DF was measured in leaf sections from image stacks as integrated

density using ImageJ (NIH). Timeseries data were uploaded to BioDare2[40] for period analysis of

background-subtracted data using the fast Fourier transform non-linear least squares (FFT-NLLS)

method[41]. Data were detrended according to the amplitude and baseline detrending method and

analysis was performed using a 24 - 96 h window. Estimates of period that fell outside of the

range of 24 h ± 5 h or had relative amplitude error (RAE) > 0.6 were excluded. Time series

intensity data presented are Z-score normalised.

Quantifying time to senescence

Leaf chlorophyll content was recorded using a handheld SPAD meter (Konica Minolta).

Measurements were taken three times/week from heading (full emergence of ear). In year two,

five SPAD values per plant were measured from the midpoint of the flag leaf of the main tiller and

the median value was recorded. In year four, ten SPAD values per plant were measured across

the whole flag leaf of the main tiller and the mean value was recorded. Time to senescence was

defined as the number of days after anthesis (DAA) to reach 50% of the maximum chlorophyll

content. Plants with time to senescence values three standard deviations above or below the

overall mean were excluded.

Nutrient profiling of grain and leaf samples

Whole flag leaves were harvested at anthesis from six plants of each genotype and mature grains

were harvested from another six plants. Flag leaf harvest was performed between 9 am and 12

pm (approximately 2-5 h after dawn). For grain samples, all ripened grain heads from each plant

were harvested. Thousand grain weight (TGW) was recorded by counting and weighing 50 seeds

from each sample. Leaf and grain samples were dried at 65°C for 7 days. Threshed grain samples

were sieved with a 2 mm slotted sieve and then milled with a Tube Mill 'Control' (Ika). Each sample

was ground for 90 seconds at 25,000 rpm. Dried leaf samples were milled using a TissueLyser

(Qiagen), with each sample milled for 2 min at 21.5 Hz with a single 18 mm stainless steel ball

bearing. Powdered leaf and grain samples were digested in HNO_3 and ion concentration was

determined by inductively coupled plasma optical emission spectroscopy (ICP-OES). Total

remobilisation efficiency was calculated as the mean normalised ratio of seed-to-leaf

concentration across eight elements (P, S, K, Ca, Mg, Fe, Zn, Mn). Total protein of grain samples

was extracted following the protocol of Dupont *et al.*[42] and quantified by bicinchoninic acid (BCA)

assay.

Defining chronotypes

Clock gene markers were identified by searching for SNPs in the OzWheat diversity panel[43]. Genotyping of the entire OzWheat panel has been performed with allele-specific markers for agronomically important genes for phenology (e.g. *Ppd-1* and *ELF3*) and around 26K SNPs were identified by transcriptome sequencing (crown tissue from seedlings) of the panel. The OzWheat polymorphism database was filtered against a set of putative wheat clock genes (Supplementary Table 1), and novel non-synonymous SNPs in clock genes were retained. Kompetitive allele specific PCR (KASP) genotyping of clock gene markers (*TaPpd-B1, TaPpd-D1, TaELF3-D1*) and novel SNP markers (*TaPRR73-A1, TaPRR59-B1, TaTOC1-B1, TaLUX-B1*) was completed for the circadian diversity panel. Homoeologue-specific primers for each clock gene marker were designed using PolyMarker[44] (Supplementary Table 6).

Genotype data for the seven clock gene markers from cultivars in the OzWheat panel plus additional cultivars in this study were used as input for *k*-modes categorical clustering, using only cultivars with complete genotypic data for the seven genes. To choose the value for *k*, the total within-cluster simple-matching distance was calculated and plotted for a range of possible values of *k* (2–10). *k*-modes clustering was performed using the R/klaR package[45]. Chronotypes were visualised in space by multiple correspondence analysis (MCA), performed using the R/FactoMineR package[46]. To estimate the number of dimensions to retain for MCA, *k*-fold cross-validation was performed using the R/missMDA package[47], and the number of dimensions (3) that produced the smallest mean square error of prediction (MSEP) was used for MCA.

RNA-sequencing

Wheat seeds of the Australian cultivar Mace were sown in a single growth room in two staggered batches, 28 days apart. Leaves were sampled concurrently when 'mature' plants were at ear emergence (Zadok's stage 51-59) and 'senescent' plants had evident chlorophyll degradation in flag leaves. At dawn (*Zeitgeber* Time 0; ZT0), the growth room was set to continuous light and temperature (20ºC) and whole flag leaves from the primary tiller of four plants were collected every 2 h from ZT45 to ZT91 (48 h). Samples were immediately flash frozen in liquid nitrogen and stored at -80°C.

Frozen leaf tissue was ground to a fine powder in liquid nitrogen-cooled mortar and pestles. Lysate buffer was added to the tissue (350 μL per 100 mg of tissue) and the mixture was vortexed immediately. RNA was extracted using ISOLATE II RNA Plant Kit (Meridian Bioscience). RNA

from two individual leaf samples was equally pooled to provide two replicates for sequencing.

Library preparation and RNA-sequencing was performed by a commercial provider (Azenta).

Library preparations were constructed using NEBNext Ultra RNA Library Prep Kit for Illumina

(New England Biolabs; NEB). Libraries with different indices were multiplexed and sequenced on

an Illumina HiSeq platform (Illumina, San Diego, CA). Sequencing was carried out using a 2x150-

bp paired-end (PE) configuration. An average of 24 million 150-bp PE reads were generated for

each library.

Reads were filtered for quality and mapped to the *T. aestivum* cv. Mace (PGSBv2.1) genome

sequence and annotation using STAR with default parameters[48]. Mapped reads were quantified

and normalised using the cuffquant and cuffnorm modules of Cufflinks[49]. Normalised counts

(fragments per kilobase transcript per million reads; FPKM) outputted from cuffnorm were used

for downstream analyses. Only transcripts that were expressed (i.e. FPKM > 0) at least once in

each 24 h period (ZT45-ZT67 and ZT69-ZT91) were retained.

Parameterisation of transcript circadian rhythms

The packages BIO_CYCLE[18] and r/MetaCycle[19] were used to estimate rhythmicity and

parameterise circadian rhythms of transcripts. BIO_CYCLE was run using default parameters,

with minimum and maximum expected period values of 20 and 28 h. MetaCycle incorporates

different methods for rhythmicity detection to generate a combined prediction of rhythmicity.

JTK_CYCLE and Lomb-Scargle (LS) were selected for integration by MetaCycle. Default settings

were maintained for all other parameters. Transcripts with $q < 0.05$ were retained from both

BIO_CYCLE and MetaCycle, and the intersection of rhythmic genes from these datasets was

used for downstream analyses. All circadian rhythm parameter estimates (period, phase,

amplitude, periodicity) presented are derived from BIO_CYCLE. Phase values are adjusted to

period (phase = phase(24/period)). The mature-to-senescent change in each of these parameters

at the gene level (Δperiod, Δphase and Δamplitude) are used for several analyses. ΔPeriod and

Δphase are calculated as the difference between the mature and senescent value of these

parameters for a given gene (e.g. Mature period - senescent period). ΔAmplitude is the log ratio

of mature and senescent amplitude for a given gene (i.e. log(Mature amplitude: senescent

amplitude)).

Gene ontology (GO) term and TF family enrichment analysis

GO terms for the *T. aestivum* cv. Mace (PGSBv2.1) annotation were sourced from Ensembl Plants[50]. r/topGO was used to perform GO enrichment analysis of various gene sets using the 'classic' algorithm[51]. Fisher's exact test was selected to compare the enrichment of GO terms in each gene set to a gene universe of 87,493 unique wheat genes with GO term annotations. For enrichment analyses of TF families, Chinese Spring wheat genes annotated as TFs were sourced from https://opendata.earlham.ac.uk/wheat/under_license/toronto/Ramirez-Gonzalez_etal_2018-06025-Transcriptome-Landscape/data/data_tables/[20]. Mace orthologues of each gene were assigned through orthology data available from Ensembl Plants[50]. For enrichment analyses, 2 x 2 contingency tables were built for each TF family and analysed with Fisher's exact test (from r/stats). Enrichment analysis of TF families that are only rhythmic in senescent tissue was performed against all rhythmic TFs (rhythmic in senescent, mature or both). Enrichment analysis of differentially rhythmic TFs was performed against all rhythmic TFs (rhythmic in both tissues).

Cis-regulatory element enrichment

Promoter sequences (1000 bp upstream of the start codon) were obtained from all wheat genes using the 'retrieve sequences' tool from RSAT (Regulatory Sequence Analysis Tools[52]). Promoter sequences were subsetted into groups of interest (e.g. short period genes), and these groups were used as input for *cis*-regulatory element enrichment analysis. For this, promoter sequences from the target set of genes were used as input for RSAT peak-motifs oligo-analysis[53], with all genes that were rhythmic in either mature or senescent flag leaves used as the control set. Peak-motifs settings were kept as default, and significantly enriched oligos with E-value < 0.05 were retained. For each significantly enriched oligo, motif matching was performed with STAMP (Similarity, Tree-building, and Alignment of Motif Profiles[54]) against the PLACE database of plant motifs[55], using default settings. The top match for each oligo (with smallest *p*-value) was retained.

Differential expression and differential rhythmicity

LimoRhyde (linear models for rhythmicity, design) was used to test for statistically significant changes in rhythmicity between mature and senescent tissue (differential rhythmicity; DR)[25]. LimoRhyde first decomposes *Zeitgeber* Time into sine and cosine with a period of 24 h. Then, transcripts that exhibit DR are identified using a linear model that tests for significant interactions between developmental stage and the time components. LimoRhyde was also used to identify differentially expressed (DE) transcripts. For this, a linear model tests for interactions between developmental stage without interaction of the time components[25]. Code used to perform

LimoRhyde's functions was developed from a vignette from the LimoRhyde GitHub repository

(https://github.com/hugheylab/limorhyde). *p*-values were adjusted by the Benjamini-Hochberg

procedure with a false discovery rate FDR = 5%[56]. DR and DE genes were assigned using a *q*-

value threshold of 0.05. DE genes have log2(fold change) > 1.2 or < -1.2. To allow comparison of

circadian rhythm parameters (i.e. period, phase, amplitude), the DR gene set was filtered against

the set of genes that are rhythmic in both tissues.

STRING protein-protein network analysis

Protein-protein interactions of DR wheat TFs were analysed using STRING[26]. STRING uses an

early genome assembly and annotation for wheat (CSS[57]). CSS gene IDs for the DR gene list

were acquired via BLAST searches within STRING, though not all Mace gene annotations had

an equivalent locus in the CSS assembly. DR TFs with STRING entries were used to build a

protein-protein network in STRING. The network was built using both physical and functional

protein interactions, and the minimum weighted interaction score was set to 0.4.

Clustering of oscillator transcript responses

Mace orthologues of putative wheat circadian oscillator genes from Chinese Spring were

identified using orthology data from Ensembl Plants[50] (Supplementary Table 2). The list of genes

was filtered to include only transcripts that are rhythmic in both mature and senescent tissue.

Principal component analysis (PCA) was performed for the 59 clock genes in this list using three

variables: mature-to-senescent changes in circadian period, phase, and amplitude (Δperiod,

Δphase and Δamplitude). The three PCs from this analysis were used as input for *k*-means

clustering. To determine an optimal value of *k*, the ratio of within-cluster sum-of-squares (WCSS)

to total sum-of-squares (TSS) was plotted against a range of potential *k* values (1-8). PCA and *k*-

means were performed using r/stats.

Acknowledgements

Research was funded by the Research Continuity Scheme from the Faculty of Science, University

of Melbourne and the JN Peters Bequest to MJH, and an Alfred Nicholas Fellowship, Melbourne

Research Scholarship and Faculty of Science Postgraduate Writing-Up Award to CRB. This

research was not funded by the Australian Research Council.

References

1. Gan, S. & Amasino, R. M. Making Sense of Senescence (Molecular Genetic Regulation and
Manipulation of Leaf Senescence). *Plant Physiol.* **113**, 313–319 (1997).
2. Distelfeld, A., Avni, R. & Fischer, A. M. Senescence, nutrient remobilization, and yield in
wheat and barley. *J. Exp. Bot.* **65**, 3783–3798 (2014).
3. Chen, L. *et al.* Leaf senescence exhibits stronger climatic responses during warm than
during cold autumns. *Nat. Clim. Change* **10**, 777–780 (2020).
4. Estiarte, M. & Peñuelas, J. Alteration of the phenology of leaf senescence and fall in winter
deciduous species by climate change: effects on nutrient proficiency. *Glob. Change Biol.* **21**,
1005–1017 (2015).
5. Bucher, S. F. & Römermann, C. The timing of leaf senescence relates to flowering
phenology and functional traits in 17 herbaceous species along elevational gradients. *J.*
Ecol. **109**, 1537–1548 (2021).
6. Simmonds, N. W. The relation between yield and protein in cereal grain. *J. Sci. Food Agric.*
67, 309–315 (1995).
7. Christopher, J. T., Christopher, M. J., Borrell, A. K., Fletcher, S. & Chenu, K. Stay-green
traits to improve wheat adaptation in well-watered and water-limited environments. *J. Exp.*
Bot. **67**, 5159–5172 (2016).
8. Christopher, M. *et al.* QTL for stay-green traits in wheat in well-watered and water-limited
environments. *Field Crops Res.* **217**, 32–44 (2018).
9. Alhabbar, Z. *et al.* Associations of NAM-A1 alleles with the onset of senescence and
nitrogen use efficiency under Western Australian conditions. *Euphytica* **214**, 180 (2018).
10. Sanchez, S. E. & Kay, S. A. The Plant Circadian Clock: From a Simple Timekeeper to a
Complex Developmental Manager. *Cold Spring Harb. Perspect. Biol.* **8**, a027748 (2016).
11. Bendix, C., Marshall, C. M. & Harmon, F. G. Circadian Clock Genes Universally Control Key
Agricultural Traits. *Mol. Plant* **8**, 1135–1152 (2015).
12. Zhang, Y. *et al.* Circadian Evening Complex Represses Jasmonate-Induced Leaf
Senescence in Arabidopsis. *Mol. Plant* **11**, 326–337 (2018).
13. Sakuraba, Y. *et al.* Phytochrome-interacting transcription factors PIF4 and PIF5 induce leaf
senescence in Arabidopsis. *Nat. Commun.* **5**, 4636 (2014).
14. Kim, H. *et al.* Circadian control of ORE1 by PRR9 positively regulates leaf senescence in
Arabidopsis. *Proc. Natl. Acad. Sci.* **115**, 8448–8453 (2018).
15. Zhang, Y. *et al.* Circadian Evening Complex Represses Jasmonate-Induced Leaf
Senescence in Arabidopsis. *Mol. Plant* **11**, 326–337 (2018).
16. Kim, H., Kim, Y., Yeom, M., Lim, J. & Nam, H. G. Age-associated circadian period changes
in Arabidopsis leaves. *J. Exp. Bot.* **67**, 2665–2673 (2016).
17. Rees, H. *et al.* A high-throughput delayed fluorescence method reveals underlying
differences in the control of circadian rhythms in Triticum aestivum and Brassica napus.
Plant Methods **15**, 51 (2019).
18. Agostinelli, F., Ceglia, N., Shahbaba, B., Sassone-Corsi, P. & Baldi, P. What time is it?
Deep learning approaches for circadian rhythms. *Bioinformatics* **32**, i8–i17 (2016).
19. Wu, G., Anafi, R. C., Hughes, M. E., Kornacker, K. & Hogenesch, J. B. MetaCycle: an
integrated R package to evaluate periodicity in large scale data. *Bioinformatics* **32**, 3351–
3353 (2016).

20. Ramírez-González, R. H. *et al.* The transcriptional landscape of polyploid wheat. *Science*
361, eaar6089 (2018).
21. Nakamichi, N., Kita, M., Ito, S., Yamashino, T. & Mizuno, T. PSEUDO-RESPONSE
REGULATORS, PRR9, PRR7 and PRR5, Together Play Essential Roles Close to the
Circadian Clock of Arabidopsis thaliana. *Plant Cell Physiol.* **46**, 686–698 (2005).
22. Noordally, Z. B. *et al.* Circadian Control of Chloroplast Transcription by a Nuclear-Encoded
Timing Signal. *Science* **339**, 1316–1319 (2013).
23. Cao, J., Liu, H., Tan, S. & Li, Z. Transcription Factors-Regulated Leaf Senescence: Current
Knowledge, Challenges and Approaches. *Int. J. Mol. Sci.* **24**, 9245 (2023).
24. Guo, Y. & Gan, S. Leaf senescence: signals, execution, and regulation. *Curr. Top. Dev.*
Biol. **71**, 83–112 (2005).
25. Singer, J. M. & Hughey, J. J. LimoRhyde: A Flexible Approach for Differential Analysis of
Rhythmic Transcriptome Data. *J. Biol. Rhythms* **34**, 5–18 (2019).
26. Szklarczyk, D. *et al.* The STRING database in 2023: protein-protein association networks
and functional enrichment analyses for any sequenced genome of interest. *Nucleic Acids*
Res. **51**, D638–D646 (2023).
27. Harmer, S. L. *et al.* Orchestrated Transcription of Key Pathways in Arabidopsis by the
Circadian Clock. *Science* **290**, 2110–2113 (2000).
28. Cane, K. *et al.* Ppd-B1 and Ppd-D1 and their effects in southern Australian wheat. *Crop*
Pasture Sci. **64**, 100–114 (2013).
29. Beales, J., Turner, A., Griffiths, S., Snape, J. W. & Laurie, D. A. A Pseudo-Response
Regulator is misexpressed in the photoperiod insensitive Ppd-D1a mutant of wheat (Triticum
aestivum L.). *Theor. Appl. Genet.* **115**, 721–733 (2007).
30. Wittern, L. *et al.* Wheat EARLY FLOWERING 3 affects heading date without disrupting
circadian oscillations. *Plant Physiol.* **191**, 1383–1403 (2023).
31. Nusinow, D. A. *et al.* The ELF4-ELF3-LUX Complex Links the Circadian Clock to Diurnal
Control of Hypocotyl Growth. *Nature* **475**, 398–402 (2011).
32. Hicks, K. A. *et al.* Conditional Circadian Dysfunction of the Arabidopsis early-flowering 3
Mutant. *Science* **274**, 790–792 (1996).
33. Kolmos, E. *et al.* A Reduced-Function Allele Reveals That EARLY FLOWERING3
Repressive Action on the Circadian Clock Is Modulated by Phytochrome Signals in
Arabidopsis. *Plant Cell* **23**, 3230–3246 (2011).
34. Anwer, M. U. *et al.* Natural variation reveals that intracellular distribution of ELF3 protein is
associated with function in the circadian clock. *eLife* **3**, e02206 (2014).
35. Alvarez, M. A. *et al.* EARLY FLOWERING 3 interactions with PHYTOCHROME B and
PHOTOPERIOD1 are critical for the photoperiodic regulation of wheat heading time. *PLOS*
Genet. **19**, e1010655 (2023).
36. Zikhali, M., Wingen, L. U. & Griffiths, S. Delimitation of the Earliness per se D1 (Eps-D1)
flowering gene to a subtelomeric chromosomal deletion in bread wheat (Triticum aestivum).
J. Exp. Bot. **67**, 287–299 (2016).
37. Boden, S. A. *et al.* Ppd-1 is a key regulator of inflorescence architecture and paired spikelet
development in wheat. *Nat. Plants* **1**, 1–6 (2015).

38. Steinfort, U., Trevaskis, B., Fukai, S., Bell, K. L. & Dreccer, M. F. Vernalisation and
photoperiod sensitivity in wheat: Impact on canopy development and yield components.
Field Crops Res. **201**, 108–121 (2017).
39. Edelstein, A. D. *et al.* Advanced methods of microscope control using µManager software.
J. Biol. Methods **1**, e10 (2014).
40. Zielinski, T., Moore, A. M., Troup, E., Halliday, K. J. & Millar, A. J. Strengths and Limitations
of Period Estimation Methods for Circadian Data. *PLOS ONE* **9**, e96462 (2014).
41. Plautz, J. D. *et al.* Quantitative analysis of Drosophila period gene transcription in living
animals. *J. Biol. Rhythms* **12**, 204–217 (1997).
42. Dupont, F. M., Vensel, W. H., Tanaka, C. K., Hurkman, W. J. & Altenbach, S. B.
Deciphering the complexities of the wheat flour proteome using quantitative two-
dimensional electrophoresis, three proteases and tandem mass spectrometry. *Proteome*
Sci. **9**, 10 (2011).
43. Hyles, J. E. Genome To Phenome: Diversity For Adaptation And Phenology In Wheat
(Triticum aestivum L.). (The University of Sydney, Sydney, Australia, 2021).
44. Ramirez-Gonzalez, R. H., Uauy, C. & Caccamo, M. PolyMarker: A fast polyploid primer
design pipeline. *Bioinformatics* **31**, 2038–2039 (2015).
45. Weihs, C., Ligges, U., Luebke, K. & Raabe, N. klaR Analyzing German Business Cycles. in
Data Analysis and Decision Support (eds. Baier, D., Decker, R. & Schmidt-Thieme, L.) 335–
343 (Springer, Berlin, Heidelberg, 2005). doi:10.1007/3-540-28397-8_36.
46. Lê, S., Josse, J. & Husson, F. FactoMineR: An R Package for Multivariate Analysis. *J. Stat.*
Softw. **25**, 1–18 (2008).
47. Josse, J. & Husson, F. missMDA: A Package for Handling Missing Values in Multivariate
Data Analysis. *J. Stat. Softw.* **70**, 1–31 (2016).
48. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).
49. Trapnell, C. *et al.* Transcript assembly and quantification by RNA-Seq reveals unannotated
transcripts and isoform switching during cell differentiation. *Nat. Biotechnol.* **28**, 511–515
(2010).
50. Cunningham, F. *et al.* Ensembl 2022. *Nucleic Acids Res.* **50**, D988–D995 (2022).
51. Alexa, A., Rahnenführer, J. & Lengauer, T. Improved scoring of functional groups from gene
expression data by decorrelating GO graph structure. *Bioinformatics* **22**, 1600–1607 (2006).
52. Nguyen, N. T. T. *et al.* RSAT 2018: regulatory sequence analysis tools 20th anniversary.
Nucleic Acids Res. **46**, W209–W214 (2018).
53. Thomas-Chollier, M. *et al.* RSAT peak-motifs: motif analysis in full-size ChIP-seq datasets.
Nucleic Acids Res. **40**, e31 (2012).
54. Mahony, S. & Benos, P. V. STAMP: a web tool for exploring DNA-binding motif similarities.
Nucleic Acids Res. **35**, W253–W258 (2007).
55. Higo, K., Ugawa, Y., Iwamoto, M. & Higo, H. PLACE: A database of plant cis -acting
regulatory DNA elements. *Nucleic Acids Res.* **26**, 358–359 (1998).
56. Benjamini, Y. & Hochberg, Y. Controlling the False Discovery Rate: A Practical and
Powerful Approach to Multiple Testing. *J. R. Stat. Soc. Ser. B Methodol.* **57**, 289–300
(1995).

57. International Wheat Genome Sequencing Consortium (IWGSC). A chromosome-based draft
sequence of the hexaploid bread wheat (Triticum aestivum) genome. *Science* **345**, 1251788
(2014).

685
686