Functional avidity of anti-B7H3 CAR-T constructs predicts antigen density thresholds for triggering effector function

Marta Barisa[1],***, Elisa Zappa [2,6], Henrike Muller[1,6], Rivani Shah[1], Juliane Buhl[2,3] Benjamin Draper[1], Courtney Himsworth[1], Chantelle Bowers[1], Sophie Munnings-Tomes[1], Marilena Nicolaidou[1], Sonia Morlando[1], Katie Birley[1], Clara Leboreiro-Babe[1], Alice Vitali[1], Laura Privitera[1], Kyle O'Sullivan[1], Ailsa Greppi[1], Magdalena Buschhaus[4], Mario Barrera Román[2,3], Sam de Blank[2,3], Femke van den Ham[2], Brenna R. van 't Veld[2], Gabrielle Ferry[1], Laura K. Donovan[1], Louis Chesler[5], Jan Molenaar[2], Jarno Drost[2,3], Anne Rios[2,3], Kerry Chester[4], Judith Wienke[2],** and John Anderson[1,7],*

Affiliations

[1] Great Ormond Street Institute of Child Health, University College London, UK

[2] Princess Máxima Center for Pediatric Oncology, Utrecht, The Netherlands

[3] Oncode Institute, Utrecht, The Netherlands

[4] Cancer Institute, University College London, UK

[5] The Institute of Cancer Research, UK

[6] Equal contribution

[7] Lead contact

*Corresponding authors

Correspondence to:

j.anderson@ucl.ac.uk, j.wienke-4@prinsesmaximacentrum.nl, m.barisa@ucl.ac.uk

Abstract

Chimeric Antigen receptor T cell (CAR-T) treatments for solid cancers have been compromised by limited expansion and survival in the tumor microenvironment following interaction with antigen-expressing target cells. Using B7H3 as a model antigen with broad clinical applicability, we evaluated the relationship between the antibody/antigen affinity of three clinical candidate binders and the three following functional characteristics: functional avidity, prolonged cytotoxicity in tumoroid re-stimulation assays, and *in vivo* anti-tumoral responses. BEHAV3D video-microscopy assessed distinct CAR-T cell behaviors at single cell resolution. T cell exhaustion did not dictate effector function. Rather, we demonstrated a threshold avidity of CAR-T / tumor cell interaction, characterized by longer cumulative CD8[+] CAR-T / tumor target interaction times, and required for adequate CAR-T cell expansion to result in sustained tumor control upon re-challenge. These results provide new insights into design of CAR-T cells for antigen-dim cell targeting, and avoidance of antigen-dim tumor relapse.

Manuscript word count: 6,975

Introduction

Genetic modification of a cancer patients' T cells with Chimeric antigen receptors (CAR-T cells) for adoptive transfer immunotherapy is a rapidly expanding clinical and scientific field. Clinical studies targeting hematological malignancy using adoptively-transferred CAR-T cell therapy have resulted in FDA approvals of CAR-T products for the treatment of leukaemia, lymphoma and myeloma [1]. In contrast, far fewer durable clinical responses attributable to CAR-T cells are reported in solid cancers despite considerable efforts[2]. A major reason for the relative failure of CAR-T cell adoptive transfer in the solid cancer setting is suboptimal T cell functionality, evidenced by a lack of T cell proliferation and persistence. Reasons for this failure can likely be distilled down to inhibitory effects of the tumor microenvironment and inherent failure of CARs to provide optimal signals to the T cell upon target engagement.

CARs comprise an antigen-sensing ectodomain, which is typically a single chain variable fragment (scFv) of an antibody, and endodomains that are an amalgamation of ITAM-containing T cell receptor (TCR) signalling domains, most typically the ζ-chain of CD3, and co-stimulatory domains. CAR-T cells thereby commandeer the MHC-unrestricted antigen specificity of a monoclonal antibody and combine it with T cell signalling in a single molecule. The triggering of T cell activity is dependent on the formation of an effective immune synapse with an antigen-positive target cell. Once a productive synapse is formed, signalling occurs through engagement of the CD3ζ and co-stimulatory moieties with proximal signalling molecules, leading to T cell effector functions and a pro-inflammatory response.

For example, structural features of the CAR that influence persistence of effector function include hinge and transmembrane components, and choice and orientation of co-stimulatory endodomains. Moreover, certain scFv binders which signal in the absence of antigen contribute to excessive signalling and subsequent T cell dysfunction[3,4]. Studies in CD19-expressing leukaemia have shown that scFv affinity can affect the strength and duration of the immune synapse with consequences for subsequent disease control[5]. The quality and quantity

of immune synapse formation are influenced by binder affinity (both on- and off-rate), target antigen density and CAR expression density. Functional avidity (strength of association between a CAR-T cell and its cellular target), is similarly governed by both affinity of interaction and expression levels of CAR and antigen in the interacting cells. A detailed understanding is lacking of how best to select an optimal CAR scFv for solid tumor-targeting CAR-T cells, and how scFv affinity and avidity impact on sustained functionality in solid tumors.

To address these challenges, we chose a broadly expressed solid tumor cancer antigen that is targeted in a number of current CAR-T clinical studies in solid tumors. B7H3 is expressed on solid and hematological cancer in the adult and pediatric settings[6,7]. A member of the immunoglobulin superfamily and closely related to PD-L1, B7H3 is enriched on high-grade tumors and expressed at low levels on healthy tissue[8–13]. There are several B7H3 binders that have been or are being progressed to evaluation in clinical trials including MGA271, 376.96 and TE9 [7,9,14]. To specifically evaluate the impact of the scFv binding properties on CAR-T function we cloned these three different binders into identical viral backbones. All constructs had a 2nd generation 28ζ endodomain and CD8 hinge and transmembrane format, and hence we set out to determine generalizable relationships between binder properties with CAR-T function rather than evaluate the binders in the context of their optimized formats for clinical studies.

Contrary to our expectations based on preceding literature[5,15–17], we found that CAR-T incorporating either of two scFv's that formed high avidity interaction with B7H3-expressing cells mediated superior anti-tumor functionality compared to a low avidity scFv. Both high and low avidity interactions enabled acute cytotoxicity and cytokine production, but only high avidity interactions led to sustained CAR-T proliferation and consequent activity in the re-challenge setting. These differences in scFv-driven functional potential were most evident against B7H3^{med} or B7H3^{dim} targets, the consequence of which was emergence of antigen-dim escape variants with the low avidity binder. We postulate that, in order to mediate tumour

control, B7H3 CAR-T cells need to interact with their target cell with a supra-threshold avidity,

sufficient to drive initial activation and proliferation.

Results

MGA271, 376.96 and TE9 scFv's mediate divergent CAR-T responses against in vivo models

of solid tumours

MGA271[18], 376.98[19] and TE9[20] scFv's were cloned into a CD8-28ζ 2nd generation CAR format

containing a CD8α hinge-transmembrane, and expressed in SFG γ-retroviral vector with an

EF1α promoter as described previously[20]. A compact epitope tag, RQR8[21], was co-expressed

from the construct to monitor CAR-T transduction efficiency (Figure 1A). Comparable

transgene co-expression between the binders was validated by staining the CAR directly with

B7H3-His protein (Supplementary Figure 1). Against LAN-1 and Kelly neuroblastoma targets,

we observed enhanced CAR-T cytokine production for 28ζ compared to 4-1BBζ endodomain

for all the binders, consistent with previous literature including our own data[7,20,22], and which

was concurrent with higher checkpoint receptor expression (Supplementary Figure 2). CD28

co-stimulation was, therefore, chosen over 4-1BB.

We first evaluated TE9 *in vivo* performance in an orthotopic, B7H3-positive[23] Med8A model of

medulloblastoma in which NOD.Cg-Prkdc scid Il2rg tm1Wjl /SzJ (NSG) mice received

intraventricular CAR-T cells 48 hours after orthotopic cerebellar tumor engraftment. Used in

these non-stress conditions, TE9-28ζ fully cleared the tumors within a week of CAR-T infusion

(Figure 1B). Next, we evaluated the binders MGA271 and 376.96 in the same model but in

increased stress conditions of longer tumor engraftment and lower CAR-T cell dose. Here the

376.96-28ζ treatment was curative (Figure 1C) whilst MGA271-28ζ did not significantly

improve survival compared with non-transduced (NTD) T cells. This was unexpected, as both

CAR-T products were equivalently cytotoxic against Med8A in a single challenge effector:

target (E:T) ratio of 1:10 co-culture (Figure 1D). In a repeat experiment of the stress condition

Med8A model, TE9-28ζ CAR-T efficacy matched that of 376.96-28ζ CAR-T, with both binders

effecting more cures than MGA271-28ζ CAR-T (Supplementary Figure 3).

We have previously shown that MGA271, 376.96 and TE9 28ζ scFv CAR-T are equivalent in

short-term single challenge cytotoxicity assays against LAN-1 neuroblastoma cells, and that

TE9-28ζ CAR-T significantly enhance survival in a difficult-to-treat subcutaneous LAN1 NSG

model[20]. We, therefore, evaluated MGA271 and 376.96-28ζ CAR-T in the LAN-1 *in vivo* model

(Figure 1E). Intriguingly, in contrast with the Med8A model where differences between binders

were apparent, 376.96 and MGA271 binders performed similarly against LAN-1 tumors,

extending survival similarly to what was to described for TE9-28ζ[20]. Taken together, different

in vivo models establish that B7H3 CAR-T functionality can be affected by scFv selection, but

in a manner that is dependent on the choice of tumor target.

MGA271 CAR-T have a lower avidity than 376.96 and TE9 CAR-T for B7H3-expressing tumor

cells

Despite achieving similar high transduction efficiencies with all three binder CARs

(Supplementary Figure 1), we noted that MGA271 CAR-T staining with B7H3-his protein was

dimmer than that of matched 376.96 and TE9 CAR-T (Figure 2A, Supplementary Figure 4A).

We hypothesized that the strength of the different scFvs' binding to target, and its impact on

strength of immune synapse might correlate with the differences observed between binders

in the Med8A and LAN-1 *in vivo* studies. To test this, we evaluated the strength of cell-to-cell

interaction for each of these CAR-T cells with Med8A, LAN-1, SupT1 wild type (SupT1-WT),

and SupT1 transduced with 4-Ig human B7H3 (SupT1-B7H3hi), representing a range of

antigen expression (Figure 2B). To avoid differences in transduction efficiency biasing avidity

measurement, CAR-T of all scFv's were diluted with donor-matched NTD T cells to

standardize expression of the RQR8 marker gene (Supplementary Figure 4B).

Avidity measurements were carried out using a Lumicks z-Movi Cell Avidity Analyzer, which

measures the force (in piconewtons, pN) required to detach a population of effector cells from

their targets (Figure 2C). None of the CAR-T bound to B7H3[neg] SupT1-WT cells, whilst all three

bound B7H3[pos] cells, albeit with different avidities (Figure 2D,E). TE9 and 376.96 scFv CAR-

T consistently bound antigen-positive target cells with higher avidity than MGA271 or NTD

cells.

To investigate the relationship between antibody-to-antigen binding properties, avidity and

antigen target density, we measured the antibodies' affinities, as well as target B7H3

molecules per cell in the respective target cell lines. Target cells with high B7H3 per cell overall

formed higher avidity synapses with CAR-T, although the positive correlation between binding

strength and antigen density was more pronounced for TE9 and 376.96 than MGA271 (Figure

2F). Binder affinity of binders in IgG1 format for 4-Ig B7H3 was measured using Surface

Plasmon Resonance. TE9 had the highest affinity for B7H3 (0.44nM), similar to 376.96

(0.69mM) and higher than MGA271 (1.61nM) (Figure 2G, Supplementary Figure 5). It was of

interest that TE9 had both the highest on-rate and the highest off-rate, whilst 376.9 had

relatively slow on- and off-rate (Figure 2H). Since affinity is the ratio of off- and on-rates,

dissociation constant (K_D) values of TE9 and 376.9 were similar. Binder affinity for B7H3

correlated with the intensity of staining of CAR-T cells by recombinant B7H3 protein (Figure

2I).

Cytotoxic capacity of low-avidity MGA271 CAR-T deteriorates upon rechallenges with B7H3[dim]

but not B7H3[bright] tumors

To examine the impact of CAR-T binding avidity on effector function, the different constructs

were assessed against solid tumor target cells that express a range of B7H3 densities. We

selected five different B7H3[pos] target types which were either cell lines or primary neurosphere

cultures (defined hereafter as 'tumoroids'), derived from pediatric solid tumors with a broad

range of B7H3 antigen densities as determined using QuantiBright beads (Figure 3A).

CAR-T cytotoxicity was first evaluated at a range of E:T ratios in single-challenge short-term

assays. All binders displayed equivalent effective cytotoxicity against B7H3[hi] neuroblastoma

cell lines LAN-1 and Kelly (Figure 3B), but relative failure of the lower affinity MGA271 binder

was seen when targeting the lower B7H3 antigen density neuroblastoma tumoroids at low E:T

ratios (≤1:10) (Figure 3C).

Capacity of CAR-T constructs to maintain killing on repeat encounter with target cells is

essential for their clinical utility. Therefore, CAR-T were re-challenged over a four-week co-

culture as illustrated in Figure 3D. Allowing the CAR-T 6 days to effect cytotoxicity, 691-T

tumoroids were killed by both low-avidity CAR MGA271 and representative high-avidity CAR

376.96 following the first challenge (Figure 3E). However, MGA271 killing dropped drastically

after re-challenge and by the third challenge no MGA271-mediated 691-T killing was

observed, while high avidity 376.96 CAR-T maintained tumor control, similar to TE9 CAR-T in

an independent experiment (Supplementary Figure 6). Using 691-B tumoroids which have

lower antigen density, the high avidity binders 376.96 and TE9 completely eliminated targets

after a single challenge whilst MGA271 showed no cytotoxicity. All CAR constructs failed

against 691-B upon re-challenge (Figure 3E, Supplementary Figure 6). A similar trend was

observed for B7H3^{med} 103-T malignant rhabdoid tumoroids. All CARs were cytotoxic against

103-T, but the inferiority of MGA271 was the most pronounced upon challenge during the third

and final week (Figure 3F,G).

Potential explanations for CAR-T failure in repeat stimulation assays is T cell exhaustion or

differentiation to terminal effector cells. While we identified upregulation of

exhaustion/activation receptors (including PD-1, TIM-3 and LAG-3) on all scFv CAR-T cells

following challenge with neuroblastoma targets (Supplementary Figure 2 for Kelly and LAN-1

cell lines; Supplementary Figure 7 for 691 tumoroids), these were not consistently different

between the different binders, and, therefore, did not appear to account for functional

differences between the CAR products. There were similarly no consistent differences

between binders in memory markers CD45RA and CCR7, nor activation markers CD25 and

CD69 (Supplementary Figure 7). Collating data from the tumoroid co-cultures at 1:10 E:T ratio,

the differences in cytotoxicity between binders at low antigen density was pronounced at both

initial challenge and re-challenges, and is represented graphically in Figure 3H. At the lowest

antigen density (~1,000 molecules per cell for 691-B tumoroids), only the high avidity binders,

TE9 and 376.96, were capable of initiating killing at first challenge but failed to sustain

cytotoxicity at re-challenge. At ~5,000 molecules per cell or more, all the binders were effective

at initial challenge but only the two high avidity binders sustained killing over multiple rounds

of stimulation (Figure 3H).

The data prompted us to hypothesize that at high antigen density, binders perform similarly

because CAR-T signalling for all of them exceeds a threshold for proliferation and cytokine

secretion to initiate tumour control. At low antigen density, high avidity interaction is required

to exceed this threshold. Therefore, we next evaluated CAR-T effector functions other than

cytotoxicity, including IL-2 and IFN-γ production, and CAR-T cell proliferation.

High avidity interaction with target cells drives CAR-T proliferation and thereby sustained

effector function

To compare activation profiles, we first measured CAR-T cytokine secretion in response to

antigen-bright LAN-1 and Kelly neuroblastoma cell lines. Unexpectedly, despite having seen

high and equivalent killing of the targets by all three CARs in short-term killing assays (Figure

3B), the lower avidity MGA271 produced significantly lower levels of IFN-γ and IL-2 than TE9

and 376.96 CAR-T (Figure 4A,B). MGA271 CAR-T cell numbers declined throughout the

assay – in a manner that was indistinguishable from NTD T cells, whilst TE9 and 376.96 CAR-

T proliferated at each challenge with targets (Figure 4C).

The pattern of CAR-T persistence and proliferation in the 691-T or -B neuroblastoma tumoroid

re-challenge assays (Figure 4D and E) mirrored the differential cytokine responses of the

different binders. MGA271, but not TE9 and 376.96, numbers declined steadily over time –

even in conditions where the MGA271 CAR-T had initially mediated high level killing at first

challenge such as was seen against 691-T in Figure 3C and E.

To evaluate whether B7H3 CAR-T responses to brain tumor targets are similar, we identified

three different medulloblastoma models with a range of antigen densities (Figure 5A). All were

B7H3^{med} or B7H3^{med-hi}. CAR-T cytotoxicity upon single challenge with these targets was

equivalently high across all scFv's at a range of E:T ratios (Supplementary Figure 8). As was

the case with the neuroblastoma targets, however, there were differences in antigen induced

CAR-T proliferation between the binders. Low avidity MGA271 proliferated significantly less

than TE9 or 376.96 CAR-T, at a level that was not significantly greater than that of NTD T cells

(Figure 5B). Proliferation for all CARs was proportionate with antigen expression, and the

required B7H3 threshold for driving an increase in T cell numbers was higher for MGA271

than TE9 and 376.96 (Figure 5C). Linking CAR-T proliferation to avidity measurements from

Figure 2 revealed that for both Med8A and LAN-1 targets, avidity appeared predictive of

subsequent T cell proliferation (Figure 5D).

High avidity, determined by ScFv affinity and B7H3 antigen density, drives CAR-T activation

To identify the threshold of B7H3 antigen density for binder-dependent CAR-T functionality

within the context of the same target cell, we generated a cell line that expresses a range of

B7H3 densities by transducing SupT1 cells with 4-Ig B7H3 (Figure 6A). The SupT1 cells were

also transduced with a transgene encoding eGFP and luciferase to enable cytotoxicity

evaluation by luminescence. Resulting SupT1-B7H3^{range} contained cells with range of B7H3

antigen densities from negative to ultra-high (Figure 6B). CAR-T were challenged with SupT1-

B7H3^{range} cells at a 1:1 E:T ratio daily for a week to mimic continued exposure to tumor (Figure

6C).

While all CAR-T cells exhibited some cytotoxicity against SupT1-B7H3^{range} cells, MGA271

killing of targets was consistently lower than that of TE9 and 376.96 CAR-T (Figure 6D). Of

note, in contrast to previous targets assayed, MGA271 CAR-T kept up sustained, if low-level,

killing even following 7 challenges with targets. None of the CAR-T killed control SupT1-WT

cells (Supplementary Figure 9A). Substantial 376.96 and TE9 proliferation was seen as target

challenge progressed, whilst more modest MGA271 proliferation was observed (Figure 6E).

The antigen dependency of the proliferation was demonstrated by the lack of response in the

presence of SupT1-WT targets (Supplementary Figure 9B) or unstimulated conditions (Figure

6F). TE9 and 376.96 CAR-T produced significantly more IL-2 and IFN-γ than MGA271 after

the initial challenge with SupT1-B7H3^{range} targets (Figure 6G).

CAR-T / tumor co-cultures were harvested after the 5th challenge, and residual SupT1-

B7H3^{range} expression of B7H3 was measured. Figure 6H shows concatenated B7H3

expression on GFP-positive SupT1 cells across the CAR groups. While TE9 and 376.96 CAR-

T cultures contained few or no residual B7H3-positive targets, MGA271 CAR-exposed SupT1

cells maintained a range of B7H3 expression. Compared to the NTD T cell condition, MGA271

eliminated only the antigen-high SupT1 targets (Figure 6H,I).

We then aimed to determine the relationship between-T effector function (cytokine secretion

and proliferation) and CAR avidity. CAR-T fold-proliferation over 5 challenges with targets

correlated with IL-2 production at the time of the initial target challenge, though 376.96 CARs

were more consistent initial IL-2 producers than TE9 CARs (Figure 6J). The avidity of CAR-T

interaction with SupT1-B7H3^{hi} targets shown in Figure 2 correlated with both IL-2 production

and proliferation in response to SupT1-B7H3^{range} cells (Figure 6K,L). A similarly strong but

inverse correlation was observed between avidity of interaction and the antigen expression on

residual SupT1-B7H3^{range} cells following serial re-challenge (Figure 6M). Taken together, the

data indicate that for each cell target engaged by a standard CD28ζ CAR-T cell, there is a

threshold of avidity required for sufficient cytokine and proliferation response to allow for

expansion and effective cytotoxicity on target re-challenge. We next examined the impact of

scFv on B7H3 CAR-T functionality at the single cell level.

High 'on-rate' is associated with high CD8^{+} CAR-T / tumoroid contact duration and rapid

cytotoxicity against B7H3^{dim} targets

BEHAV3D, a multispectral, 3D image-based platform, is designed to live-track the efficacy

and mode of action of cellular immunotherapy at the single cell level [24]. BEHAV3D was used

to track anti-B7H3 CAR-T functionality over three challenges with 691-B and 691-T tumor

targets. Cultures were imaged during the first (day 0) and third (day 7) challenge with targets,

and the series of images then processed for CAR-T behavior using a bioinformatic pipeline,

developed by Anne Rios and colleagues[24] (the assay setup is illustrated in Figure 7A and B

and Supplementary Figure 10). CAR-T cell behavior was tracked across six different

parameters, shown in Figure 7B. Video compilations of day 0 and day 7 CAR-T / tumoroid co-

cultures can be viewed in Supplementary Videos 1-24.

CAR-T/tumoroid interactions leading to tumoroid death was evident for all conditions

(representative co-cultures images shown in Figure 7C). In analysis of behaviors against

B7H3^{dim} 691-B tumoroids, striking and unexpected differences were seen between the

binders, and between CD4$^+$ and CD8$^+$ T cells. Plotting the cumulative CD8$^+$ CAR-T/tumoroid

contact since initiation of the day 0 co-culture showed that TE9 and 376.96 CD8$^+$ CAR-T spent

significantly longer in contact with B7H3^{dim} 691-B tumoroids than MGA271 CD8$^+$ CAR-T did

(Figure 7D). A summation of total CD8$^+$ CAR-T / 691-B contact time during the entire day-0

analysis confirmed this difference between binders (Figure 7F). This cumulative contact score

correlated with the cytotoxicity of respective binder-CAR-T constructs against these B7H3^{dim}

691-B targets in an overnight 1:10 E:T killing assay (Figure 7D, E). In contrast, all CD8$^+$ CAR-

T showed similar contact time and cytotoxicity with higher antigen density 691-T tumoroids

(Figure 7D-F). Analyzing cytotoxicity by detection of organoid death during the imaging (tumor

cell conversion from 'yellow' to 'red' in Figure 7C) confirmed the observation that TE9 CAR-T

were the most rapid at killing B7H3^{dim} 691-B tumoroids followed by 376.96 and MGA271 CAR-

T cells. In contrast, all CAR-T cells were equally rapid at killing 691-T tumoroids of higher

antigen density (Figure 7G).

There were no differences between binders in terms of the number of new CAR-T contacts

during the 800 minute data collection, using either tumoroid model (Supplementary Figure

11A). Hence the finding of highest total T cell/tumoroid contact time of the TE9 CD8 CAR-T

cells (Figure 7F) implied TE9 was the binder conferring the longest duration of individual

contacts. Consistent with this observation, TE9 CD8+ CAR-T cells also showed the lowest

displacement distance (Figure 7H) and speed of displacement (Figure 7I). whilst MGA271 had

the shortest and 376.96 were intermediate (Figure 7F,H,I). This ranking of tumor interaction

duration of TE9 > 376.96 > MGA271 when targeting low antigen density targets correlated

with the cytotoxicity, avidity, and binder association rate constant (Supplementary Figure 11B).

These data suggest that high avidity binder CD8+ CAR-T cells stay longer in contact with low

antigen tumoroids to facilitate more efficient killing.

Compared to CD8+ CAR-T cells, CD4+ CAR-T cells showed greater displacement and reduced

tumoroid contact, and this was observed in both models of higher and lower antigen density

respectively (Figures 7F,H,I). Moreover, in CD4+ CAR-T, the influence of the respective

binders on tumoroid contact and displacement was variable, consistent with CD4+ CAR-T cells

having a lesser role in tumoroid contact and direct cytotoxicity, as described previously using

BEHAV3D analysis[24] (Figure 7F,H,I). While more mobile overall, CD4+ CAR-T cells did not

engage in CAR-T / CAR-T interaction more than CD8+ CAR-T cells (Supplementary Figure

11C).

Distinct CAR-T cell characteristics revealed by dimensionality reduction of behavioral

parameters

To extract behavioral patterns and dynamics, the day 0 BEHAV3D tumor challenge imaging

data was clustered using UMAP (Supplementary Figure 12A), revealing six CAR-T behavior

types: 'tumoroid engagers' (cluster 1), 'multi-taskers' (clusters 2, 3), 'CAR-T engagers'

(clusters 4,5) and 'dying T cells' (cluster 6) (Figure 7J). Fitting with the individual cell analysis,

the 'tumoroid engager' cluster 1 was relatively enriched in CD8$^+$ cells compared to CD4$^+$ and

with higher antigen density 691-T compared with 691-B targets (Figure 7K). Moreover, TE9

and 376.96 binder CD8$^+$ T cells consistently exhibited more cluster 1-type tumor engager

behavior than MGA271 CAR-T against both tumor targets, and further exhibited more 'multi-

tasker' behavior in the 691-B tumoroid co-culture. These differences between binders were

less apparent in the CD4$^+$ CAR-T context.

CAR-T behavior was then compared between the two analyzed timepoints (day 0 and day 7).

Day 7 CAR-T behavior was reduced using UMAP into 7 clusters across the two imaging

sessions (Supplementary Figure 12B), which grouped into 'organoid interaction' (clusters A,

B, C, D), 'CAR-T interaction (clusters E, F) and cell death (cluster G) (Supplementary Figure

12C). Strikingly, no consistent differences between the CAR-T binders or even CD4$^+$ *vs* CD8$^+$

T cells remained evident at day 7 (Supplementary Figure 12D). This narrowing of CAR-T

phenotype was evident when comparing the trajectory of tumoroid death over the day 0 and

day 7 imaging sessions, pooling data from all three binders and target types (Figure 7L). A

broad range of tumor killing trajectory was seen at day 0 whilst at day 7 all CARs mediated

rapid tumor clearance. This apparent ability of low avidity MGA271 CAR-T to control low

antigen density tumor at day 7 is most likely due to artificially resetting the effector to target

ratio at each tumor challenge with re-sorted viable CAR-T cells.

The dimensionality reduction BEHAV3D analysis reflects the broad conclusion that high

avidity of CD8$^+$ T CAR-T cells equips them for longer interactions with low antigen density

targets, which facilitates sustained cytotoxic activity. The data supports our hypothesis that

the differences in CAR-T functionality seen *in vivo* and *in vitro* assays are caused not by

variation in binder-driven functional exhaustion, but rather by the differential ability of the CAR-

T constructs to proliferate and engage in serial tumor cytotoxicity.

Discussion

CAR-T cell technology represents a highly promising cancer therapy as exemplified by FDA approvals for several agents in the hemato-oncology field. In solid cancers the pace of development has been slower. The solid cancer environment is immunologically hostile and hypoxic, which limits of capacity of infiltrating CAR-T Cells to engage in the sustained proliferation and cytotoxicity, which is a requirement for shrinking a solid tumor in which the ratio of tumor to CAR-T cells is stacked in favour of the former. CAR-T relative failure is likely compounded by T cell exhaustion, a phenomenon whereby T cells become transcriptionally programmed to hypo-functionality, as a result of repeated antigen stimulation. The capacity to engage in efficient serial killing through repeated successful interactions with tumor cells without inducing T cell exhaustion may be the defining feature of a successful solid cancer CAR-T construct.

The CAR-T to tumor interaction encompasses a CAR immune synapse, wherein the initial contact between T cell and tumor is stabilised by the CAR antibody to target antigen binding interaction, which develops into a cluster of engaged CAR constructs[25,26]. Although the biophysical properties of the CAR-T to antigen synapse are not as well characterized as native TCR immune synapses, it is likely that the strength of this interaction as well as the number of target antigens will determine whether the synapse becomes functional; *i.e.* triggers death of the target cell through release of cytolytic granules concomitant with T cell signalling to promote proliferation and cytokine secretion[25,26].

Successful formation of a synapse leads to the exclusion of phosphatases (*e.g.* CD45) and the recruitment of src family kinases (*e.g.* lck) resulting in phosphorylation of CD3ζ chain ITAMs within the CAR molecule, and subsequent signalling. An efficient and high-quality CAR-T immune synapse might be one that is strong enough to induce death of the target and short-lived enough to induce a "goldilocks" quantity and quality of T cell signalling that promotes activation without exhaustion. Immune synapse quality is likely affected by the nature of the

initiating antibody-to-antigen interaction. Recent progress in CAR-T / target interaction

tracking at the single cell level, on platforms such as BEHAV3D[24], has enabled a nascent

understanding of the kinetics of immune synapse formation and how synaptic formation

dynamics differ between CD4 and CD8 cells. The relationships between the synapse, the

binding properties of the CAR scFv's and the effect of synaptic signalling on T cell activation

and exhaustion remain poorly understood.

Functional avidity is the strength of interaction between a CAR-T cell and its tumor target,

which we have measured as the force required to disrupt the interaction. We hypothesized

that avidity would be influenced by three factors: the CAR surface expression on CAR-T cells,

the target antigen density on the tumor cells, and the affinity of the antibody to antigen

interaction at the single molecular interaction scale. We reasoned that this interplay of factors

would be such that none of the parameters alone would be predictive of CAR-T response, but

the interaction of the three factors to define functional avidity would be the closest predictor of

subsequent T cell function. We evaluated the relationship of antibody to antigen biding

properties as both affinity (Biacore surface plasmon resonance) and functional avidity

(Lumicks) measurements against a range of antigen densities. Two binders of similar affinity

(K_D) had quite differing rate constants of association and dissociation with similar avidity (TE9

and 376.96), whilst a third of lower affinity also demonstrated low avidity (MGA271).

Several previous studies have investigated avidity of CAR-T cells using the Lumicks platform.

For example, the group of Maher *et al* studied a range of anti-CD19 binders in the context of

leukemia therapy using conventional 2[nd] generation CARs co-expressing a 4-1BB chimeric

costimulatory receptor[27]. Interestingly, intermediate avidity CAR-T were identified to have the

greatest *in vivo* capacity to control tumor growth.

Using repeat stimulation assays against suspension cells, monolayers and three dimensional

tumoroid models to recapitulate the repeat antigenic engagement of the tumor environment,

we show that low avidity interaction on first tumor encounter results in a weak cytokine and

proliferative response. Weak initial response translates into ultimate failure of expansion and

tumor control in stress conditions. In contrast, higher avidity initial interaction in the low antigen

density setting results in effective proliferation and serial killing. In the setting of higher antigen

density, all CAR-T constructs have sufficiently high avidity initial interaction to induce

expansion and serial killing although low avidity MGA271 still ultimately has reduced

functionality at the end of serial challenges. Previous studies have identified fast off-rate as

linked to more efficient and effective CAR-T control of leukemia[5,15–17] and it is therefore an

interesting contrast in our tumoroid studies to note that the quickest construct for tumor control

of low antigen density had the longest organoid contact time as well as the fastest on- and off-

rates. This apparent discordance might be explained by the multiple cell contacts that single

$CD8^+$ CAR-T generate with the multi-cellular tumoroid structures as previously described using

BEHAV3D analysis[24]. Here, a more complex and dynamic interaction of an individual $CD8^+$

CAR-T cell with multiple tumor cell targets within a 3D structure might explain a more complex

relationship between avidity and success in tumor control.

Of note, in the current models which extend to seven repeat tumor challenges we have not

seen evidence of differences in T cell exhaustion between the three binders. This may be

because the repeat antigenic challenges were not sustained long enough and it is also

tempting to speculate that a higher affinity antibody / higher avidity CAR-T cell might have

induced exhaustion in the same repeat stimulation experiments. All binders evaluated showed

only modest increases in TIM-3$^+$PD-1$^+$ double-positive cells and little evidence of exhaustion

induced by signalling in the absence of antigen (tonic signalling). It will be important in future

studies to evaluate whether very high avidity binders and/or tonic signaling binders show

relative failure associated with increased exhaustion marker expression.

One notable observation was that the TE9 binder, which had similar affinity and avidity to the

376.96 binder, had both the highest on-rate and fastest on-rate. To determine if this impacted

on the CAR-T / tumor interaction time, we performed BEHAV3D video microscopy analysis.

Interestingly, we found that at higher antigen densities there were no difference in dwell time

between the binders. In contrast, at low antigen density, the CD8$^+$ 376.96 and TE9 CAR-T

cells had longer dwell times than MGA271 CAR-T, which was consistent with both their greater

cytotoxicity and the higher displacement score of the MGA271 CAR-T. The fast on- and off-

rates (in single molecule Biacore affinity studies) of TE9 appears to translate into longer

tumoroid contact, and leads us to speculate that avidity measurements may prove to be of

greater predictive value for CAR-T synapse formation characteristics. A further surprising

result was the short tumour dwell-time of TE9 CD4$^+$ cells, which appeared to be of less

importance in direct cytotoxicity. Further studies are required to confirm these observations

with other binders and targets, as well as mechanistic studies to determine the significance of

differences in avidity between CD4$^+$ and CD8$^+$ CAR-T cells.

B7H3 is an emerging attractive target for cancer immunotherapy due to its bright expression

on the surface of a high proportion of cancer cells, its association with poor prognosis, its

potential function as an immune regulator, and its relative absence on healthy tissues [28,29].

Interestingly, its cancer specificity seems to extend beyond cancer cells in the tumor

environment since several lines of evidence point to its expression in a variety of tumor

infiltrating cells, such as vascular endothelial cells. Hence, the clinical application may extend

beyond cancers in which B7H3 has a role as a driver oncogene. Indeed, its vascular

endothelial expression might mitigate tumor heterogeneity being a cause of antigen escape

variants which has been a significant consideration diminishing enthusiasm for clinical

progression with several alternate antigens. All three of the binders evaluated in this study are

of clinical relevance and are being evaluated in a range of clinical trials.

There are three important implications of these avidity findings for clinical activity of these

respective binders. The first is threshold of antigen density for activation. B7H3 is expressed

at low level on some healthy tissues. A lower avidity binder might have optimal activation

threshold to distinguish healthy from tumor tissue to avoid on-target off-tumor toxicity.

Secondly, the avidity threshold for activation will be affected by the respective signalling

characteristics of the CAR. For example, work using the MGA271 binder has shown that

substitution of the CD8 hinge and transmembrane regions (H/TM) as evaluated in this paper,

for CD28 H/TM, within the context of 41BB-CD3ζ signaling, results is significantly improved

function against low antigen density targets[6]. In current clinical trial NCT05474378 (Mackall

and colleagues), the configuration is with CD8 hinge/transmembrane with 41BB costimulation.

A second recruiting clinical study (SJ-19-0014; St Jude's, USA; DeRenzo *et al*) using MGA271

with CD28ζ is deploying an additional 41BBL co-stimulation *in-trans* module[22], whilst trial

NCT04185038[14] (Seattle, USA; Vitanza and team) makes use of MGA271 with CD28 H/TM in

a 2nd gen 41BB-CD3ζ configuration. Hence, none of the current MGA271 trials are using the

configuration in the current paper. Thirdly, higher expression of CAR through use of higher

MOI of virus could increase avidity of a given binder against targets. Artificially, we ensured

equivalent expression levels to allow fair comparison of avidity effects in relation to target

antigen density. Higher expression of CAR might increase avidity and functionality. What

seems clear is that all CARs require fine-tuning of binders to identify optimal activation

thresholds. Avidity measurements may prove an additional valuable parameter for screening

constructs to optimal signalling and activation.

Figures and Figure Legends

Figure 1. MGA271, 376.96 and TE9 scFv's mediate divergent CAR-T responses against _in_

vivo models of medulloblastoma and neuroblastoma. (**A**) Schematic of the retroviral

constructs. (**B**) Orthotopic "non-stress" condition Med8A tumor model pilot in NSG mice; N=3

for NTD (non-transduced T cells), N=4 for TE9-CAR (tumor luminescence comparison

between groups by Two-Way ANOVA). (**C**) Orthotopic "stress condition" Med8A tumor model

in NSG mice; N=4 mice for all conditions (Kaplan-Meyer comparison by Log-rank Mantel-Cox

test). (**D**) Overnight CAR-T cytotoxicity assay run in-parallel to the _in vivo_ study from panel (**C**)

(E:T ratio 1:10, N=1). (**E**) Subcutaneous LAN-1 tumor model in NSG mice. N=5 mice for all

conditions (Kaplan-Meyer curve comparison by Log-rank Mantel-Cox test).

Figure 2. 376.96 and TE9 but not MGA271 CAR-T form high avidity interactions with tumor

cells in a B7H3 density dependent manner. (**A**) Cryopreserved, thawed CAR-T cell surface

481 transgene expression was detected using flow cytometry with clone QBEND10 mAb to detect

482 RQR8 marker gene and histidine-x6 tagged B7H3 protein (B7H3-his) + anti-his mAb to detect

483 the CAR (mean ± SEM, N=3 donors, Two-Way ANOVA). (**B**) Target cells were stained with

484 anti-B7H3-PE mAb and analyzed by FACS. Shown on the staggered histogram are respective

485 geometric mean fluorescence intensities (gMFI) of B7H3-PE. (**C**) Principle of Lumicks z-Movi

486 avidity measurement platform. (**D**) CAR-T detachment plotted against force in pN applied to

487 the T cell / target interaction. The line indicates the mean T cell attachment of 4 independent

488 donors across 3 experimental replicates whilst shaded area represents SEM. (**E**) Avidity as

489 represented by attachment at 990-1,000 pN applied force: N=4 donors each with 3

490 experimental replicates, One-Way ANOVA. (**F**) Bound CAR-T (%) at 1,000 pN plotted against

491 QuantiBright defined B7H3 molecules per cell. The lines are linear regressions of the means

492 of 4 individual donors (Two-Way ANOVA). (**G**) Binder affinity for B7H3 (4-Ig) was measured

493 using whole antibodies in IgG1 format using Biacore Surface Plasmon Resonance (SPR). (**H**)

494 The association ('on-rate') and dissociation ('off-rate') rate constants for the different binders

495 were plotted. (**I**) Binder affinity was plotted against the gMFI of B7H3-his protein staining of

496 three independent donor CAR-T cell products.

Figure 3. TE9 and 376.96, but not MGA271 CAR-T, respond to B7H3 ultra-dim tumor targets

in serial re-challenge cytotoxicity assays. (**A**) Number of B7H3 molecules/cell (QuantiBright

flow cytometry assay) for range of targets. (**B**) CAR-T cell cytotoxicity against LAN-1 and Kelly

neuroblastoma cell lines (4h ^{51}Cr release assay). 'Specific cytotoxicity' denotes cytotoxicity

relative to non-transduced (NTD) T cell controls (mean ± SEM, N=3 donors, Two-Way

ANOVA). (**C**) CAR-T cell cytotoxicity against GFP/Luc-engineered tumoroids by overnight

luminescence-based assay. 'Specific cytotoxicity' denotes cytotoxicity relative to NTD T cell

controls (mean ± SEM, N=2 independent donors across 3 technical replicates, Two-Way

ANOVA). (**D-F**) Re-challenge assay timecourse (**D**), against neuroblastoma (**E**) and malignant

rhabdoid tumor (MRT) organoids (**F**); gray arrows indicate re-challenge with tumor. In (**E**), data

is mean ± SEM, N=2 independent T cell donors across 3 experimental replicates and 'specific

cytotoxicity' denotes cytotoxicity relative to NTD T cell controls; in (**F**) 'specific cytotoxicity'

denotes cytotoxicity relative to tumor alone controls plated in parallel, data is mean ± SEM,

N=1 independent T cell donor (same as one of the donors used to generate data for panel **E**)

across 3 experimental replicates. (**G**) Representative microscopy images of 103-T MRT

tumoroid being killed by CAR-T over time. (**H**) 1:10 E:T ratio cytotoxicity for the first and third

week challenge from the assays showed in panels (**E** and **F**) plotted against antigen density

(Quantibright); line = mean cytotoxicity, shaded area = SEM.

Figure 4. High avidity interaction with target cells drives CAR-T proliferation and resulting

sustained effector function. (**A**,**B**) CAR-T cells were stimulated with irradiated LAN-1 or Kelly

neuroblastoma cell lines at a 1:1 E:T ratio every 7 days ; IFN-γ (**A**) and IL-2 (**B**) were measured

in supernatants collected 24h after every stimulation (mean ± SEM, N=3 independent donors,

Two-Way ANOVA). (**C**) T cell numbers in the same co-cultures were evaluated at each time

point using Precision Count Beads and flow cytometry (mean ± SEM, N=3 independent

donors, Two-Way ANOVA). (**D**) Relative T cell proportions measured using flow cytometry in

691-B and 691-T tumor co-cultures over three weeks of weekly tumor challenges at an E:T

ratio of 1:10. Gated on singlet PBMC of harvested cultures, representative dot plots from one

donor CAR-T are shown.

Figure 5. High avidity interactions drive CAR-T proliferation and cytotoxicity in brain tumor

models. (**A**) B7H3 molecules per cell were quantified using a QuantiBright flow cytometry

assay; brain tumor targets in blue, neuroblastoma and rhabdoid tumor targets in gray. (**B**) T

cell proliferation in response to live brain tumor challenge at a 1:1 E:T ratio was quantified

after 6 days of co-culture (mean ± SEM, 3 independent T cell donors across two experimental

replicates, Kurskal-Wallis test). (**C**) T cell fold-expansion against the number of B7H3

molecules per target cell; dots represent mean proliferation of the different donors and

replicates; lines are simple linear regression curves. (**D**) Mean fold-proliferation of CAR-T cells

against Med8a and LAN-1 cells was plotted against the mean avidity at 1,000 pN for the two

cell lines. The dots represent mean proliferation of the different donors and replicates; the

lines are simple linear regression curves.

Figure 6. High avidity drives elimination of antigen-dim targets. (**A**) Constructs for generation

of B7H3 and GFP/luciferase sublines of SupT1 cells. (**B**) Expression of B7H3 in the unsorted

'SupT1-B7H3^{range}' and flow-sorted 'SupT1-B7H3^{hi}' cells. (**C**) Rechallenge assay experimental

schematic. For all subsequent data in the panels: N=4 independent donors across 4

experimental replicates. (**D**) Cytotoxicity was measured by luciferase signal as relative to

tumor luminescence in the presence of donor-matched non-transduced (NTD) T cells (mean

± SEM, Two-Way ANOVA). (**E**) T cell numbers in the same assay following either a range of

challenges with tumor targets or (**F**) target-free (mean ± SEM, Two-Way ANOVA). (**G**)

Cytokine measurements in co-culture supernatants following a single challenge wit SupT1-

B7H3^{range} targets at a 1:1 E:T ratio (mean ± SEM, Two-Way ANOVA). (**H**) B7H3 expression

on targets was measured using flow cytometry following the fifth challenge. Shown are

concatenated histograms that were generated by combining flow data adjusted for 10,000

SupT1 cells per sample. (**I**) The same SupT1-B7H3^{range} positivity for antigen as analysed in

each sample separately (mean represents geometric mean fluorescence intensity (gMFI) ±

SEM, Two-Way ANOVA). (**J**) Respective CAR fold-proliferation in response to 5 challenges

with SupT1-B7H3^{range} targets plotted against IL-2 production following the initial challenge. (**K**)

IL-2 production after first challenge plotted against the mean avidity for SupT1-B7H3^{hi} targets

at 1,000 pN. (**L**) Proliferation after 5 challenges with targets was plotted against the mean

synapse avidity that the respective CARs achieved when combined with SupT1-B7H3^{hi} targets

at 1,000 pN. (**M**) SupT1-B7H3^{range} cell B7H3 gMFI after 5 challenges with targets was plotted

against the mean synapse avidity that the respective CARs achieved when combined with

SupT1-B7H3^{hi} targets at 1,000 pN. In (**J**-**M**) each dot represents an individual donor

experimental replicate, the lines are simple linear regressions.

Figure 7. Higher CAR binder B7H3 avidity associated with increased CD8$^+$ CAR-T / tumoroid

contact duration and more rapid cytotoxicity against B7H3$^{ultra-dim}$ tumoroids. (**A**,**B**) experimental

overview. (**C**) Illustrative stills of tumoroid and T cell clusters at the start (day 0 = t0), and end

(day 7 = t1) 800 minute imaging sessions. (**D**) CD8$^+$ CAR-T cell cumulative contact time with

691-B and 691-T organoids tracked for the first 2h of the first tumor challenge on day 0. The

line represents the mean cumulative tumoroid contact score of all the individual CAR-T cells

that were successfully tracked for the full 120 min period (the number of cells tracked in each

of the conditions was a mean 31.33±10.8; curves were compared using simple linear

regressions). (**E**) CAR-T cells were incubated with tumoroids overnight at a 1:10 E:T ratio.

'Specific cytotoxicity' denotes tumor luminescence relative to matched non-transduced T cell

(NTD) controls (mean ± SEM, N=3 independent experimental replicates, Kruskal-Wallis test).

(**F-I**) Respective parameters were tracked for entire duration (800 minutes) of the first tumor

challenge at day 0. (**F**) CD4$^+$ and CD8$^+$ CAR-T cell cumulative contact time with 691-B and

691-T organoids (mean ± SEM, N = 31.5 ± 10.6 cells per T cell condition, Two-Way ANOVA).

(**G**) Cumulative tumoroid death (conversion from viability dye yellow to red) (a total of 398

incremental time-points were recorded to track tumoroid apoptosis over the assay; curve

comparison was carried out using a Friedman test for P value and rank sum differences

(RSD)). (**H**) CD4$^+$ and CD8$^+$ CAR-T displacement (mean ± SEM, N = 31.5 ± 10.6 cells per T

cell condition, Two-Way ANOVA). (**I**) Mean speed of CD4$^+$ and CD8$^+$ CAR-T displacement

(mean ± SEM, N = 31.5 ± 10.6 cells per T cell condition, Two-Way ANOVA). (**J**) Trends in

CAR-T behavior reduced into 6 clusters using UMAP analysis of the T cell behavior across all

the conditions observed in the initial tumor challenge at day 0 and represented as a heatmap.

(**K**) The CAR-T behaviors broken down by T cell, binder and target type. (**L**) Pooled data from

all CAR constructs against both targets comparing day 0 and day 7 challenge rate of organoid

death.

Acknowledgements

We thank Debarati Shome and Reza Nadafi of Lumicks for their support with our z-Movi

analysis of B7H3-CAR-T synapse avidities, and Lumicks more broadly for supporting us with

hardware and technical advice during the experiments. We're grateful to the Flow Cytometry

Core Facility at UCL GOS Institute of Child Health and to the Imaging Center at the Princess

Máxima Center for their support in running our studies. We thank the Princess Máxima Center

Imaging and Tumoroid facilities for support.

Funding

We are grateful for the following grant support. Anderson and Chesler: Stand up to Cancer/

CRUK Paediatric Cancer New Discoveries Challenge; Anderson, Chesler and Donovan: CwC

INSTINCT-MB award, Fight Kids Cancer, and LPT programme awards; Anderson: Research

into Childhood Cancer, NIHR GOSH BRC, GOSHCC 552864, AMR/LifeArc awards. Drost and

Buhl: Oncode Institute, Children Cancer-free Foundation (KiKa), Deutsche

Forschungsgemeinschaft (German Research Foundation, DFG). Molenaar: Villa Joep, Veni.

Author Contributions

MB, JW, and JA wrote the manuscript; MB, HM, EZ, MBR, and SbD analyzed data; GF

generated reagents; EZ, HM, RS, BD, CH, CB, SMT, MN, SM, KB, CLB, AV, LP, KoS, AG,

and MB performed experiments; JA, JW, JB, KC, AR, JD, JM, LC co-supervised the research.

Conflicts of Interest

MB, JA, KC, KB, MB hold patents in CAR-T technology development, including a pending

patent for the TE9 anti-B7H3 binder. JA holds founders shares in Autolus.

References

1. Research, C. for B. E. and. Approved Cellular and Gene Therapy Products. *FDA* (2023).

2. Albelda, S. M. CAR T cell therapy for patients with solid tumours: key lessons to learn and unlearn. *Nat Rev Clin Oncol* **21**, 47–66 (2024).

3. Weber, E. W. *et al.* Transient "rest" restores functionality in exhausted CAR-T cells via epigenetic remodeling. *Science* **372**, eaba1786 (2021).

4. Weber, E. W. *et al.* Pharmacologic control of CAR-T cell function using dasatinib. *Blood Adv* **3**, 711–717 (2019).

5. Ghorashian, S. *et al.* Enhanced CAR T cell expansion and prolonged persistence in pediatric patients with ALL treated with a low-affinity CD19 CAR. *Nature Medicine* **25**, 1408–1414 (2019).

6. Majzner, R. G. *et al.* Tuning the Antigen Density Requirement for CAR T-cell Activity. *Cancer Discov* **10**, 702–723 (2020).

7. Du, H. *et al.* Antitumor Responses in the Absence of Toxicity in Solid Tumors by Targeting B7-H3 via Chimeric Antigen Receptor T Cells. *Cancer Cell* **35**, 221-237.e8 (2019).

8. Theruvath, J. *et al.* Locoregionally administered B7-H3-targeted CAR T cells for treatment of atypical teratoid/rhabdoid tumors. *Nat Med* **26**, 712–719 (2020).

9. Majzner, R. G. *et al.* CAR T Cells Targeting B7-H3, a Pan-Cancer Antigen, Demonstrate Potent Preclinical Activity Against Pediatric Solid Tumors and Brain Tumors. *Clinical Cancer Research* **25**, 2560–2574 (2019).

10. Maachani, U. B. *et al.* B7-H3 as a Prognostic Biomarker and Therapeutic Target in Pediatric central nervous system Tumors. *Transl Oncol* **13**, 365–371 (2020).

11. Modak, S., Kramer, K., Gultekin, S. H., Guo, H. F. & Cheung, N. K. Monoclonal antibody 8H9 targets a novel cell surface antigen expressed by a wide spectrum of human solid tumors. *Cancer Res* **61**, 4048–4054 (2001).

12.   Chapoval, A. I. *et al.* B7-H3: A costimulatory molecule for T cell activation and IFN-γ

production. *Nature Immunology* **2**, 269–274 (2001).

13.   Wang, L., Kang, F.-B. & Shan, B.-E. B7-H3-mediated tumor immunology: Friend or

foe? *International Journal of Cancer* **134**, 2764–2771 (2014).

14.   Vitanza, N. A. *et al.* Intraventricular B7-H3 CAR T Cells for Diffuse Intrinsic Pontine

Glioma: Preliminary First-in-Human Bioactivity and Safety. *Cancer Discov* **13**, 114–131

(2023).

15.   Chmielewski, M., Hombach, A., Heuser, C., Adams, G. P. & Abken, H. T Cell

Activation by Antibody-Like Immunoreceptors: Increase in Affinity of the Single-Chain

Fragment Domain above Threshold Does Not Increase T Cell Activation against Antigen-

Positive Target Cells but Decreases Selectivity1. *The Journal of Immunology* **173**, 7647–

7653 (2004).

16.   Mause, E. R. V. *et al.* Combinatorial T cell engineering eliminates on-target off-tumor

toxicity of CD229 CAR T cells while maintaining anti-tumor activity. 2021.12.06.471279

Preprint at https://doi.org/10.1101/2021.12.06.471279 (2022).

17.   Olson, M. L. *et al.* Low-affinity CAR T cells exhibit reduced trogocytosis, preventing

rapid antigen loss, and increasing CAR T cell expansion. *Leukemia* **36**, 1943–1946

(2022).

18.   Loo, D. *et al.* Development of an Fc-enhanced anti-B7-H3 monoclonal antibody with

potent antitumor activity. *Clin Cancer Res* **18**, 3834–3845 (2012).

19.   Imai, K., Wilson, B. S., Bigotti, A., Natali, P. G. & Ferrone, S. A 94,000-dalton

glycoprotein expressed by human melanoma and carcinoma cells. *J Natl Cancer Inst* **68**,

761–769 (1982).

20.   Birley, K. *et al.* A novel anti-B7-H3 chimeric antigen receptor from a single-chain

antibody library for immunotherapy of solid cancers. *Molecular Therapy - Oncolytics* **26**,

429–443 (2022).

21.   Philip, B. *et al.* A highly compact epitope-based marker/suicide gene for easier and

safer T-cell therapy. *Blood* **124**, 1277–1287 (2014).

22.   Nguyen, P. *et al.* Route of 41BB/41BBL Costimulation Determines Effector Function

of B7-H3-CAR.CD28ζ T Cells. *Mol Ther Oncolytics* **18**, 202–214 (2020).

23.   Labanieh, L. *et al.* Enhanced safety and efficacy of protease-regulated CAR-T cell

receptors. *Cell* **185**, 1745-1763.e22 (2022).

24.   Dekkers, J. F. *et al.* Uncovering the mode of action of engineered T cells in patient

cancer organoids. *Nat Biotechnol* 1–10 (2022) doi:10.1038/s41587-022-01397-w.

25.   Sajman, J. *et al.* Nanoscale CAR Organization at the Immune Synapse Correlates

with CAR-T Effector Functions. *Cells* **12**, 2261 (2023).

26.   Xiong, Y., Libby, K. A. & Su, X. The physical landscape of CAR-T synapse.

Biophysical Journal (2023) doi:10.1016/j.bpj.2023.09.004.

27.   Halim, L. *et al.* Engineering of an Avidity-Optimized CD19-Specific Parallel Chimeric

Antigen Receptor That Delivers Dual CD28 and 4-1BB Co-Stimulation. *Frontiers in*

Immunology **13**, (2022).

28.   Kontos, F. *et al.* B7-H3: An Attractive Target for Antibody-based Immunotherapy.

Clinical Cancer Research **27**, 1227–1235 (2021).

29.   Zhou, W.-T. & Jin, W.-L. B7-H3/CD276: An Emerging Cancer Immunotherapy.

Frontiers in Immunology **12**, (2021).

30.   Schutgens, F. *et al.* Tubuloids derived from human adult kidney and urine for

personalized disease modeling. *Nat Biotechnol* **37**, 303–313 (2019).

31.   Calandrini, C. *et al.* An organoid biobank for childhood kidney cancers that captures

disease and tissue heterogeneity. *Nat Commun* **11**, 1310 (2020).

32.   Custers, L. *et al.* Somatic mutations and single-cell transcriptomes reveal the root of

malignant rhabdoid tumours. *Nat Commun* **12**, 1407 (2021).

33.   Koo, B.-K. *et al.* Controlled gene expression in primary Lgr5 organoid cultures. *Nat*

Methods **9**, 81–83 (2011).

34.   Johnson, L. S., Huang, L., Moore, P. A., Loo, D. T. & Chen, F. Z. Antibodies reactive

with B7-H3 and uses thereof. (2014).

35.   Dotti, G., Ferrone, S., Du, H., Wang, X. & Ferrone, C. Methods and compositions for

chimeric antigen receptor targeting cancer cells. (2019).

36.   Sabbatino, F. *et al.* Novel Tumor Antigen-Specific Monoclonal Antibody-Based

Immunotherapy to Eradicate Both Differentiated Cancer Cells and Cancer-Initiating Cells

in Solid Tumors. *Seminars in Oncology* **41**, 685–699 (2014).

37.   The z-Movi Cell Avidity Analyzer workflow. *LUMICKS*

https://lumicks.com/knowledge/the-z-movi-workflow/.

38.   Jason-Moller, L., Murphy, M. & Bruno, J. Overview of Biacore systems and their

applications. *Curr Protoc Protein Sci* **Chapter 19**, Unit 19.13 (2006).

Methods

Cells lines and culture conditions

All cell lines were cultured in humidified 37C^o in 5% CO_2 conditions. Cell lines were thawed

and passaged for at least two weeks prior to use in functional assays. Culture mycoplasma

tests were performed monthly using MycoAlert Detection Kit from Lonza. The following table

summarizes the characteristics and culture conditions of the cell lines that were used over the

course of this study.

Cell Line ID	Type of Cancer	Source	RRID	Culture Media
LAN-1 (LA-N-1)	Neuroblastoma	ECACC	CVCL_1827	RPMI1640 (Gibco) + 10% FCS (Gibco)
Med8A (MED-MEB-8A) wild type	Medulloblastoma	Gift from Michael Taylor, The Hospital for Sick Children, Canada	CVCL_M137	DMEM (Gibco) + 10% FCS + 1x Glutamax (Gibco)
Med8A GFP/Luc	Medulloblastoma	Derived from Med8A wild type by RD114 γ-retroviral transduction with GFP/Luc	None	DMEM (Gibco) + 10% FCS + 1x Glutamax (Gibco)
Kelly	Neuroblastoma	Gift from Andrew Stoker, UCL, UK	CVCL_2092	RPMI1640 + 10% FCS
SupT1 wild type	Childhood T lymphoblastic lymphoma	ECACC	CVCL_1714	RPMI1640 + 10% FCS
SupT1-B7H3^{hi}	Childhood T lymphoblastic lymphoma	Derived from SupT1 wild type by RD114 γ-retroviral transduction with 4-Ig B7H3 (Sinobiological), followed by flow cytometric sorting	None	RPMI1640 + 10% FCS
SupT1-B7H3^{range} GFP/Luc	Childhood T lymphoblastic lymphoma	Derived from SupT1 wild type by RD114 γ-retroviral transduction with GFP/Luc followed by flow cytometric sorting, then 4-Ig B7H3 (Sinobiological) without sorting	None	RPMI1640 + 10% FCS

691-T (AMC691-T) wild type	Neuroblastoma	Gift from Jan Molenaar (Utrecht)	None	'Neurosphere medium': DMEM-GlutaMAX supplemented with 20% Ham's F-12 nutrient mixture, 2% B-27 supplement minus vitamin A, 1% N-2 Supplement, 100 U/mL penicillin, 100 mg/mL streptomycin, 300 ng/mL insulin-like growth factor (IGF-1), 40 ng/mL fibroblast growth factor-basic (FGF-2), 20 ng/mL epidermal growth factor (EGF), 10 ng/mL platelet-derived growth factor-AA (PDGF-AA) and 10 ng/mL platelet-derived growth factor-BB (PDGF-BB)
691-T (AMC691-T) GFP/Luc	Neuroblastoma	Gift from Jan Molenaar (Utrecht)	None	'Neurosphere medium'
691-B (AMC691-B)	Neuroblastoma	Gift from Jan Molenaar (Utrecht)	None	'Neurosphere medium'
691-B (AMC691-B) GFP/Luc	Neuroblastoma	Gift from Jan Molenaar (Utrecht)	None	'Neurosphere medium'
103-T wild type	Malignant rhabdoid tumor	Gift from Jarno Drost (Utrecht)	None	'MRT medium'[30,31]: MRT tumoroids were cultured in droplets of growth factor-reduced basement membrane extract (BME) Type 2 (R&D Systems) in kidney tumoroid medium (KOM) (Advanced Dulbecco's Modified Eagle Medium (DMEM)/F12 (Gibco) containing 1X GlutaMAX (Thermo Fisher Scientific), 10 mM HEPES (Thermo Fisher Scientific) and 1X Penicillin-Streptomycin (P/S) (Merck Millipore), supplemented with 10% R-spondin–conditioned medium, 1.5% B27 supplement (Gibco), 50 ng/mL epidermal growth factor (EGF) (PeproTech), 50 ng/mL fibroblast growth factor (FGF-)2 (PeproTech), 1.25 mM N-acetylcysteine (Sigma), 10 M Rho-associated coiled–coil containing protein kinase (ROCK) inhibitor Y-27632 (Abmole) and 5 uM A83-01 (Tocris Bioscience))
103-T Luc	Malignant rhabdoid tumor	Gift from Jarno Drost (Utrecht); derived from 103-T wild type by lentiviral transduction with pLKO.1-UbC-luciferase blast and pLV-H2B-mNeon-puro cassettes[32,33]	None	'MRT medium'
D341 wild type	Medulloblastoma	Gift from Michael Taylor, The Hospital for Sick Children, Canada	CVCL_ 0018	DMEM (Gibco) + 10% FCS + 1x Glutamax (Gibco)

D341 GFP/Luc	Medulloblastoma	Derived from D341 wild type by RD114 γ-retroviral transduction with GFP/Luc	None	DMEM (Gibco) + 10% FCS + 1x Glutamax (Gibco)
D425 wild type	Medulloblastoma	Gift from Michael Taylor, The Hospital for Sick Children, Canada	CVCL_1275	DMEM (Gibco) + 10% FCS + 1x Glutamax (Gibco)
D425 GFP/Luc	Medulloblastoma	Derived from D425 wild type by RD114 γ-retroviral transduction with GFP/Luc	None	DMEM (Gibco) + 10% FCS + 1x Glutamax (Gibco)
293-T (HEK293T)	Transformed cell line	ATCC	CVCL_0063	IMDM + 10% FCS

CAR sequences

All CARs were expressed using an SFG backbone in a 2nd generation RD114 γ-retroviral

system. All plasmids were sequenced to verify absence of unexpected sequence mutations

prior to manufacture of virus. A diagram of construct structure is shown in Figure 1A, with all

anti-B7H3 CARs expressed in the following format: RQR8-T2A-scFv-CD8αLinker-CD8αTM-

CD28endodomain-CD3ζchain. The sequences for the scFv's were obtained from public

sources and are summarized in the below table. To insert these into the CAR backbone, the

scFv sequences were ordered as geneblocks and then restriction cloned into a CD8α-CD28-

CD3ζ format.

scFv ID	Full CAR Amino Acid Sequence	Source of scFv Sequence
MGA271	EVQLVESGGGLVQPGGSLRLSCAASGFTFSSFGMHWVRQAPGKGLEWVAYISSDSSAIYYADTVKGRFTISRDNAKNSLYLQMNSLRDEDTAVYYCGRGRENIYYGSRLDYWGQGTTVTVSSGGGGSGGGGSGGGGSDIQLTQSPSFLSASVGDRVTITCKASQNVDTNVAWYQQKPGKAPKALIYSASYRYSGVPSRFSGSGSGTDFTLTISSLQPEDFATYYCQQYNNYPFTFGQGTKLEIKAAADPTTTPAPRPPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLDFACDIYIWAPLAGTCGVLLLSLVITLYRTRSKRSRLLHSDYMNMTPRRPGPTRKHYQPYAPPRDFAAYRSRVKFSRSADAPAYQQGQNQLYNELNLGRREEYDVLDKRRGRDPEMGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRGKGHDGLYQGLSTATKDTYDALHMQALPPR*	Macrogenics Inc Patent 'US9441049B2'[34]; development described by Loo and colleagues[18]

_376.96_	EVQLVESGGGLVKPGGSLKLSCEASRFTFSSYAMSWVRQTPEKRLEWV AAISGGGRYTYYPDSMKGRFTISRDNAKNFLYLQMSSLRSEDTAMYYCA RHYDGYLDYWGQGTTLTVSSGGGGSGGGGSGGGGSDIVMTQSHKFMS TSIGARVSITCKASQDVRTAVAWYQQKPGQSPKLLIYSASYRYTGVPDRF TGSGSGTDFTFTISSVQAEDLAVYYCQQHYGTPPWTFGGGTKLEIKAAAD PTTTPAPRPPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLDFACDIYIWA PLAGTCGVLLLSLVITLYRTRSKRSRLLHSDYMNMTPRRPGPTRKHYQPY APPRDFAAYRSRVKFSRSADAPAYQQGQNQLYNELNLGRREEYDVLDK RRGRDPEMGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRRGK GHDGLYQGLSTATKDTYDALHMQALPPR*	University of North Carolina at Chapel Hill, General Hospital Corp Patent 'US10233226B2'[35]; development described by Imai[19] and Sabbatino[36]
_TE9_	QVQLQQSGAALVKPGTSVKLSCKASGYTFTSYWMHWVKQRPGQGLEWI GMIHPKSGSVDYNEKFTNKATLTGDKSSGTAYMQLSSLTSEDSAVYYCA RGGYGSPFDYWGQGTTVTVSSGGGGSGGGGSGGGGSENVLTQSPAIM SASPGEKVTMTCSASSSVSYMYWYQQKPGSSPRLLIYRTSNLASGVPAR FSGSGSGTSYSLTISSMEAEDAATYYCQQWSSNPPTFGGGTKLEIKRAA ADPTTTPAPRPPTPAPTIASQPLSLRPEACRPAAGGAVHTRGLDFACDIYI WAPLAGTCGVLLLSLVITLYRTRSKRSRLLHSDYMNMTPRRPGPTRKHY QPYAPPRDFAAYRSRVKFSRSADAPAYQQGQNQLYNELNLGRREEYDV LDKRRGRDPEMGGKPRRKNPQEGLYNELQKDKMAEAYSEIGMKGERRR GKGHDGLYQGLSTATKDTYDALHMQALPPR*	Developed in-house at UCL. Development described by Birley _et al_ [20]

CAR-T cell manufacture

Leukapheresis cones were acquired from NHS Blood and Transplant. PBMCs were separated

through Ficoll centrifugation using Lymphoprep (STEMCELL Technologies). PBMCs were

washed and residual red cells lysed with ACK Lysis buffer (Thermo Fisher Scientific). NK cells

were depleted using magnetic CD56 depletion beads (Miltenyi Biotec) and LD depletion

columns (Miltenyi Biotec). PBMCs were suspended in RPMI1640 containing 10% FCS and 1x

L-glutamine at a concentration of 1×10^6 cells/mL, and then activated with 0.5 mg/mL of anti-

CD3 (clone: OKT-3; Miltenyi Biotec) and anti-CD28 antibodies (clone: CD28.2; Miltenyi

Biotec). Forty-eight hours before transduction and on the day of transduction, 100 IU/mL

recombinant human IL-2 (Proleukin, Novartis) was added. T cells were transduced with the

above-mentioned γ-retroviral SFG construct following plating on retronectin (Takara Bio)-

coated plates and spinoculation. Transduction efficiency was measured three days after

transduction using flow cytometry by staining for RQR8 (clone: QBNED10; R&D BioSystems).

T cell populations were not corrected for transduction efficiency in functional assays.

Transduction efficiencies of >50% were routinely achieved. Expanding CAR-T cell IL-2 media

was replenished every two days. Expansions were harvested at day 10-12 following culture

initiation and cryopreserved in 10% DMSO complete media. When initiating *in vitro* or *in vivo*

functional assays, CAR-T cells were thawed in pre-warmed IL-2 (100 IU/mL) complete media

(RPMI1640, 10% FCS), rested overnight and then counted. CAR-T E:T ratios were based on

the number of viable T cells after overnight rest. Cryopreserved CAR-T recovery after

overnight rest (relative to the number of CAR-T that were frozen down) was routinely 50-60%,

with a ~90% viability.

Animal studies

Animal protocols were approved by local institutional research committees and in accordance

with UK Home Office guidelines. Mixed male and female NSG mice aged between 6 and 8

weeks were supplied by Charles River. All experiments were carried out under UK home office

licenses project license number 15981/01, and personal license numbers I13398879,

P2E645DD9, I83870811, I27052867, I8C20AD0D and I754094BC.

For the LAN-1 *in vivo* study, NSG mice were injected with 1 x 10^6 LAN-1-BFP/Luc in Matrigel

(Corning) subcutaneously into the flank. CAR-T cells (1 x 10^7) were injected intravenously into

the tail vain at day 10 following tumor engraftment (treatment schedule illustrated in Figure

1E). Tumor size was monitored twice a week with digital callipers. Mice were given 200 uL

luciferin (Thermo Fisher Scientific) into the scruff and imaged using a PhotonIMAGERTM

optical imaging system (Biospace Lab) weekly. Animal sacrifice was initiated upon reaching

either a tumor diameter of >15 mm in any direction (as measured by callipers) or the

appearance of tumor burden toxicity that exceeds our animal welfare standards (>15% weight

loss, change in activity, ataxia, dehydration, loss of normal gait, grimacing).

For the Med8A *in vivo* study, two types of model were run: the 'high-stress' model and the

'low-stress' model. In both models 5 x 10^4 Med8A cells re-suspended in PBS were injected

into the cerebellum of an NSG mouse. In the 'low-stress model' (illustrated in Figure 1B), two

days after tumor engraftment the mice were administered intraventricularly with 2.5 x 10^6 CAR-

T cells. In the 'high-stress model' (illustrated in Figure 1C) the mice were treated six days after

tumor engraftment, and with 0.5×10^6 CAR-T cells per animal. Mice were given 200 uL luciferin

(Thermo Fisher Scientific) into the scruff and imaged using a PhotonIMAGERTM optical

imaging system (Biospace Lab) weekly. Animal sacrifice was initiated upon reaching either a

tumor luminescence of 10E10 photons/second/cm^2 (via IVIS), or the appearance of tumor

burden toxicity that exceeds our animal welfare standards (>15% weight loss, doming of the

cranium, change in activity, ataxia, dehydration, loss of normal gait, grimacing).

CAR-T in vitro functional assays

A number of different functional CAR-T assays were performed. While high (>60%)

transduction efficiency of CAR-T was routinely achieved, transduced cell numbers were

normalised for functional assays to ensure that each binder condition receives the same

number of viable CAR-T cells. All data was collected in experimental duplicates or triplicates,

as indicated in figure legends.

CAR-T responses against LAN-1 and Kelly target cells were assessed as follows. For the

overnight (~18h) co-culture cytokine assays, CAR-T cells were co-cultured with LAN-1, Kelly

cells or no antigen stimulus in 48-well plates at an E:T ratio of 1:1. After 18 h, supernatant was

removed for ELISA and cells incubated with monensin (BioLegend). Checkpoint receptors PD-

1, TIM-3 and LAG-3 were detected by flow cytometry. Overnight cytotoxicity was tested using

a ^{51}Cr release assay. Target cells were incubated with ^{51}Cr for 1 h then washed and plated in

96-well plates. CAR-T cells or untransduced cells were plated at a range of E:T ratios. The

plates were incubated for 4 h at 37°C and then the supernatant was removed and incubated

with scintillation fluid (PerkinElmer) overnight at room temperature. ^{51}Cr released into the

supernatant was measured using a 1450 MicroBeta TriLux (PerkinElmer). For the re-

challenge assays, CAR-T cells were co-cultured with irradiated LAN-1, Kelly, or no target cells

in 24-well plates at an E:T ratio of 2:1. Cell medium was replenished every 2–3 days. CAR-T

cells were challenged with irradiated target cells every 6 days, cultured for a further 24 h, and

analyzed. Cells were pelleted and supernatant was removed every week for ELISA. CAR-T

cell proliferation was measured weekly by flow cytometry using Precision Count Beads

(BioLegend). The levels of cytokines IL-2 and IFN-γ were quantified using ELISA MAX Deluxe

Set Human IL-2 and ELISA MAX Deluxe Set Human IFN-γ (BioLegend).

SupT1-B7H3^{range} re-challenge assays were performed as follows (illustrated in Figure 6C). For

the overnight (~18h) co-culture cytokine assays, CAR-T cells were co-cultured with GFP/Luc-

SupT1-B7H3^{range} cells or GFP/Luc-SupT1-WT cells in 96-well plates at an E:T ratio of 1:1.

After 18 h, supernatant was removed for analysis with ELISA MAX Deluxe Set Human IL-2

and ELISA MAX Deluxe Set Human IFN-γ (BioLegend). Cytotoxicity at the various indicated

timepoints was tested using a luciferase-based assay. Briefly, 15ug/mL of D-luciferin (Perkin

Elmer) was added to each well to document the remaining number of living tumor cells. The

plate was then incubated for 5 min at 37°C in 5% CO_2 and run on a SpectraMax, Molecular

Devices analyzer, with 'no effector' tumor wells as controls for T cell-mediated tumor death.

CAR-T cells or untransduced cells were plated at a 1:1 E:T ratio, with daily re-challenges.

Cytotoxicity readings were done in this manner 18h after each new tumor re-stimulation. CAR-

T cell proliferation was measured weekly by flow cytometry using Precision Count Beads

(BioLegend).

Brain tumor challenge assays with GFP/Luc-Med8A, -D425 and -D341 cells were done by

plating CAR-T cells with targets at the indicated E:T ratios in 96-well plates. Cytotoxicity at

96h following tumor challenge was tested using a luciferase-based assay. Briefly, 15ug/mL of

D-luciferin (Perkin Elmer) was added to each well to document the remaining number of living

tumor cells. The plate was then incubated for 5 min at 37°C in 5% CO_2 and run on a

SpectraMax, Molecular Devices analyzer, with 'no effector' tumor wells as controls for T cell-

mediated tumor death. CAR-T cell proliferation was measured weekly by flow cytometry using

Precision Count Beads (BioLegend).

Malignant rhabdoid tumoroids were established, characterised and cultured as previously

described[30,31]. 103-T tumoroids were transduced with a construct encoding for luciferase were

cultured in KOM in BME droplets for 7 days in advance. 4 days before the co-culture the

tumoroids were disrupted mechanically into a single cell suspension by mechanical

dissociation and seeded in BME droplets. On the day of CAR-T challenge, MRT tumoroids

were collected, washed with cold medium to remove BME, and a fraction was dissociated into

a single cell suspension. Single tumor cells were counted to determine the number of cells

present in the MRT tumoroid suspension. MRT tumoroids were then seeded out with an

equivalent of 7500 single tumor cells per well in 50 uL of co-culture medium containing 10%

FBS and a 1:1 ratio of KOM and RPMI with 1X GlutaMAX and 1X P/S. To determine tumor

specific killing, a released luciferase assay was performed using the luciferase assay system

(Promega). Medium was removed and cells were washed with 100 uL of PBS. 20 uL of 1X

passive lysis buffer (Promega) were added and the plate was incubated at room temperature

for 15 min while shaking. In a LUMITRAC white polystyrene 96-well plate (Greiner) 100 uL of

the luciferase reagent were added to 20 uL of the lysed cells and the plate was measured at

the FluoSTAR Omega Microplate reader immediately. Luciferase signal was determined on

day 1 after co-culture as well as before re-challenge with new tumoroids.

To measure CAR-T cytotoxicity against (AMC) 691-B and 691-T neuroblastoma tumoroids,

GFP-Luc tumoroids were dissociated into single cells by re-suspending in 1 mL of Accutase

solution (Sigma-Aldrich) through a p10 tip on top of a p1000 tip to combine both enzymatic

and mechanic dissociation methods. Tumoroid cells in suspension were counted and plated

two days before the beginning of the experiment to ensure spheroid formation in 96-well white

flat bottom plates(Costar). On the day of initiation co-culture, CAR-T cells were counted and

plated with GFP/Luc tumoroids at the stated E:T ratios in half 'neurosphere medium'

(expanded on in the Cell Lines section of Methods and Materials) and half T cell medium

(RPMI- GlutaMAX 1640 + 25 mM Hepes (ThermoFisher) supplemented with 10% FCS

(Gibco), 1% penicillin/streptomycin (ThermoFisher) and 1% L- glutamine (ThermoFisher)

supplemented with 100 IU/mL IL-2 (Miltenyi Biotec)). For re-challenge, tumoroids were plated

2 days before each challenge, with the CAR-T culture added on top. Tumoroid re-challenge a

set E:T ratio was done every 2-3 days. Cytotoxicity was measured by adding 15ug/mL of D-

luciferin (Perkin Elmer) to each well to document the remaining number of living tumor cells,

with 'no effector' tumor wells as controls for T cell-mediated tumor death. Cells were then

incubated for 5' at 37°C in 5% CO_2 and readout performed in the FLUOstar Omega microplate

reader.

Flow cytometry staining and analysis

The following cell cytometers were used to collect flow cytometry data: BD ® LSR II Flow

Cytometer (BD Biosciences), FACSymphony™ A5 High-Parameter Cell Analyzer (BD

Biosciences) and CytoFLEX S V4-B2-Y4-R3 Flow Cytometer (Beckman Coulter). FlowJo

10.9.0 software for Mac (Becton Dickinson) was used to analyse flow cytometry data. All

staining was done in FACS buffer, PBS, 2% FCS with 2mM EDTA. Samples were either

analysed immediately following staining or fixed with Biolegend Fixation Buffer and analysed

within two weeks. The following table lists the flow cytometry staining reagents that were used:

Target	Product ID	Catalogue Number
Viability Dye	Fixable Blue Dead Cell Stain — LIVE/Dead™ (Thermo)	L34961
Flow count beads	Precision Count Beads (Biolegend)	424902
B7H3-his	B7H3-his protein to detect CAR at the cell surface (R&D Biosystems)	1949-B3-050/CF
His	Anti-his mAb clone J095G46-APC (BioLegend)	362605
Ag Density Quantification	BD Biosciences™ Quantibrite™ Phycoerythrin (PE) Beads; used to quantify B7H3 antigen density on target cell surface	10626384
Fixation buffer	Fixation buffer (Biolegend)	420801
Cell membrane dye	eBioscience Cell Proliferation Dye eFluor 450 (blue) (Thermo Fisher); used to label cells for BEHAV3D analysis	65-0842-85

Cell membrane dye	Calcein AM (green) (Thermo Fisher); used to label cells for BEHAV3D analysis	C1430
Cell membrane dye	CellTrace™ Far Red Cell Proliferation Kit, for flow cytometry; used to label cells for Lumicks z-Movi analysis	C34564
RQR8 marker gene	Clone QBEND10-APC (BioTechne)	FAB7227A
CD3	Clone OKT-3-Brilliant Violet 711 (Biolegend)	317328
CD4	Clone VIT4-VioBlue (Miltenyi Biotec),	130-113-219
B7H3	Clone MIH42-PE (BioLegend)	351003
PD-1	Clone EH12.1-Brillant Violet 605 (Beckton Dickinson)	563245
CD25	Clone 2A3-Briljant Violet 711 (Beckton Dickinson)	563159
CD4	Clone A161A1-FITC (Biolegend)	357406
CD8	Clone SK1-PerCP (Biolegend)	344706
TIM-3	Clone F38-2E2-PE (Biolegend)	345006
CCR7	Clone G043H7-PE/Dazzle594 (Biolegend)	353235
CD69	Clone FN50-PE-Cy7 (Beckton Dickinson)	561928
CD137	Clone 4B4-1-APC (Beckton Dickinson)	550890
CD3	Clone HIT3a-Alexa Fluor 700 (BioLegend)	300324
CD45RA	Clone HI100-APC/Cy7 (Beckton Dickinson)	750258

Lumicks CAR-T synapse avidity measurements

Proprietary z-Movi Lumicks protocols were followed under the supervision of Lumicks staff

present for the assays. Briefly, SupT1-WT, SupT1-B7H3^{hi}, LAN-1 and Med8A tumor cells were

seeded in a z-Movi microfluidic chip (Lumicks, Amsterdam, Netherlands) coated with poly-L-

lysine and cultured for 16 hours. The next day, flow sorted, transduction efficiency-normalized

and CellTrace Red (Biolegend)-labelled CAR-T cells were serially flowed in the chips and

incubated with the target cells for 10 minutes prior to initializing a 3-minute linear force ramp.

During the force ramp, the z-Movi device (Lumicks) captured a time series of images using a

bright field microscope integrated into the platform. Detached cells were levitated towards the

acoustic nodes, allowing the tracking of cells based on their XY positions. Changes in the Z-

position resulted in a change in the diffraction pattern, which allowed the distinction between

cells adhered to the substrate and cells suspended to the acoustic nodes. This information

was used to correlate cell detachment events with a specific rupture force. Cell detachment

was acquired using z-Movi Tracking and post experiment image analysis was done using Cell

Tracking offline analysis. Data were presented as 'mean % CAR-T attached cells' at the

maximum mean force applied, 1000 piconewtons, pN. This was repeated in experimental

triplicate for four biological T cell donors. A greater depth of z-Movi workflow is reported

online[37]. The software used to acquire the data was z-Movi Software (v1.0).

Binder affinity measurements

Biacore surface plasmon resonance (SPR) for anti-B7H3 antibodies was done as previously

described[38]. Briefly, MGA271, 376.96 and TE9 binders were expressed in human IgG1 format

(Evitria), and then evaluated by Antibody Analytics using Biacore SPR for binding kinetics to

4-Ig human B7H3. Anti-B7H3 antibodies were first adhered to plates and several

concentrations of Human B7-H3 analyte were passed over the surface to assess the binding

interactions. The data is shown in Figure 2G. Clone 9G8 mAb was used as an isotype control.

BEHAV3D CAR-T / tumoroid imaging

Tumoroids (691-B and -T) were grown in 'Neurosphere medium' (as described in the 'Cell

Lines' section), cultured for 3 weeks prior to the experiments and passaged once or twice a

week according to tumoroid size and confluence. The experimental timeline is illustrated in

Figure 7A. CAR T cells were thawed one day prior to the start of the experiment in RPMI-

GlutaMAX 1640 + 25 mM Hepes (ThermoFisher) supplemented with 10% FCS (Gibco), 1%

penicillin/streptomycin (ThermoFisher) and 1% L- glutamine (ThermoFisher), rested overnight

at a density of 1×10^7 cells/mL in 100 IU/mL IL-2 (Miltenyi Biotec)-supplemented medium. CAR

T cells for the initial challenge day 0 imaging experiment were stained with the following

antibody mix in FACS buffer (PBS 1x + 2% FCS + 2mM EDTA): anti-RQR8 mAb clone

QBEND10-APC (R&D Systems), anti-CD4 mAb clone VIT4-VioBlue (Miltenyi Biotec) and

mouse serum (Invitrogen). $RQR8^+CD4^+$ and $RQR8^+CD4^-$ CAR-T cells were FACS-sorted with

a Sony SH800S cell sorter into FCS, and rested overnight in a density of 1×10^7 cells/mL in

medium supplemented with 100 IU/mL IL-2 (Miltenyi Biotec).

For co-culture experiments, tumoroids were dissociated into single cells by pipetting 1 mL of

Accutase solution (Sigma-Aldrich, counted and plated two days before the beginning of the

experiment to ensure spheroid formation in either 12-well flat-bottom plates (for rec-hallenges)

(Greiner) or in a glass-bottom 96-well SensoPlate (Greiner) for imaging. The correct cell

number was determined considering organoid growth rates. Wild type organoids were used.

Sorted and rested CAR T cells were stained separately with eBioscience Cell Proliferation Dye

eFluor 450 (1:4000, ThermoFisher) or Calcein AM (1:3000, ThermoFisher) in PBS for 15

minutes at 37°C and mixed in a 1:1 ratio immediately before plating them with the organoids

in a 1:5 E:T ratio. The co-culture medium (100µL organoids + 100µL T cells) was then

supplemented with 2.5% basement membrane extract (BME, Cultrex), Nucred Dead 647

(three drops per mL, ThermoFisher), and TO-PRO-3 (1:5000, ThermoFisher). The

combination of Nucred Dead 647 and TO-PRO-3 has previously been described by Dekkers

et al[24]. The plate was placed in a SP8 confocal microscope (Leica) containing an incubation

chamber (37°C, 5% CO2) and imaged for thirteen hours in time series with two-minute

intervals (as shown in Supplementary Figure 10). Day 7 BEHAV3D imaging was the third

challenge with 691 tumoroids. On Day 6, cells from re-challenge plates were stained with the

following antibody mix in FACS buffer (PBS1x + 2% FCS + 2mM EDTA): anti-RQR8 mAb

clone QBEND10-APC (R&D Systems), anti-CD4 mAb clone VIT4-VioBlue (Miltenyi Biotec),

anti B7H3 mAb clone MIH42-PE (BioLegend) and mouse serum (Invitrogen). $B7H3^-$

$CD34^+CD4^+$ and $B7H3^-$ $CD34^+CD4^-$ CAR-T cells were FACS-sorted with the Sony SH800S

cell sorter into FCS, rested overnight at a density of 1×10^7 cells/mL in medium supplemented

with 100 IU/mL IL-2 (Miltenyi Biotec). On day 7, rested CAR-T cells were stained separately

and plated with WT organoids in a 1:5 E:T ratio, as described above. The plate was placed in

a SP8 confocal microscope (Leica) containing an incubation chamber (37°C, 5% CO2) and

imaged for thirteen hours in time series with a two-minute interval.

Image processing was done according to the protocol previously described by Dekkers *et al*[24]

using the Imaris (Oxford Instruments) version 10.0 software for 3D visualization, cell

segmentation and extraction of statistics. The Channel Arithmetics Xtension was used to

create additional channels for the specific identification of $CD8^+$ and $CD4^+$ T cells (live and

dead), organoids (live and dead), and to exclude debris. The Surface and ImarisTrack

modules were used to specifically detect and track T cells and/or organoids. For tracked T

cells, time-lapse statistics consisting of the coordinates of each cell, speed, square

displacement, distance to either organoids or the other T cell subset, and dead cell dye

channel intensity were exported into a metadata file for subsequent processing and analysis.

Data presentation

Data was graphed and analysed statistically using Graphpad Prism 10 for macOS software

(Dotmatics). Statistical tests were chosen as suitable to particular data sets, and statistical test

recommendations by Graphpad Prism 10 software were followed, based on data type,

replicates and degree of pairing. The number of biological and experimental replicates used

in each analysis is listed in respective figure legends. Data processing for the BEHAV3D

platform was done using 'R' programming language code that was written by and published

from the Rios group[24].