Title page

Universal protocol to separate and compare Extracellular Vesicles from human plasma and skeletal muscle biopsy

Authors: Valentina Mangolini[1,2], Annalisa Radeghieri[1,3], Simone Piva[4,5], Stefano Cattaneo[6], Marco Brucale[7,3], Francesco Valle[7,3], Arianna Balestri[8,3], Costanza Montis[8,3], Nicola Latronico[4,5,9], Paolo Bergese[1,3,10], Lucia Paolini[5,3]*.

Affiliations:

1. Dept. Molecular and Translational Medicine (DMMT), University of Brescia, Brescia, Italy.

2. IRCCS Fondazione Don Carlo Gnocchi ONLUS, Milano, Italy.

3. Center for Colloid and Surface Science, (CSGI), Sesto Fiorentino (FI), Italy.

4. ASST Spedali Civili di Brescia, Dept. of Emergency, Spedali Civili UniversityHospital, Brescia, Italy.

5. Dept. of Medical and Surgical Specialties, Radiological Sciences and Public Health (DSMC), University of Brescia, Brescia, Italy.

6. ASST Spedali Civili di Brescia, ASST Spedali Civili, Dept. of Bone and Joint Surgery, Brescia, Italy.

7. Consiglio Nazionale delle Ricerche, Istituto per lo Studio dei Materiali Nanostrutturati, Bologna, Italy.

8.Dept. of Chemistry "Ugo Schiff", University of Firenze, Firenze, Italy.

9. "Alessandra BONO" University Research Center on LOng Term Outcome (LOTO) in Survivors of Critical Illness, University of Brescia, Italy

10. IRIB - Institute for Research and Biomedical Innovation of CNR, Palermo Italy.

*Corresponding author: Lucia Paolini

e-mail address: lucia.paolini@unibs.it

Postal address: Department of Medical and Surgical Specialties, Radiological Sciences and Public Health, University of Brescia. Viale Europa, 11, 25123, Brescia, Italy.

Ethical Approval and Consent to Participate

Consent for Publication

All authors have reviewed and endorsed this version of the manuscript for publication.

Availability of Data and Materials

The CONAN and DLS datasets supporting the conclusions of this article are available in the Zenodo repository, https://doi.org/10.5281/zenodo.10579305

Conflict of Interest

The authors declare no conflicts of interest.

Funding

This work was supported by the Italian Ministry of Health (Ministero della Salute) under the Ricerca Finalizzata (RF) "Theory enhancing" grant: ID project RF-2021-12375279, BOW project, Horizon 2020 – Future and emerging technologies (H2020 –FETOPEN), ID: No. 952183 and "Alessandra Bono" Foundation;

ABSTRACT:

The majority of studies on extracellular vesicles (EVs) focused on samples isolated from liquid matrices, such as cell culture media and blood, due to their accessibility. However, recent research highlights the emerging roles of EVs derived from solid tissues, including the brain, muscles and tumors. Investigating EVs from the extracellular matrix of solid tissues offers insights into their microenvironment and potential biological influences on surrounding cells. This study presents a universal method for comparing EV-enriched samples from solid (human skeletal muscle biopsy) and liquid (human plasma) matrices, addressing technical challenges and minimizing biases in separation techniques. By employing optimized protocols and advanced analytical techniques, the study reveals differences in biomolecular composition, nanomechanical properties, particle yield, size distribution, and colloidal stability between human skeletal muscle and plasma EVs. Understanding these distinctions may contribute to the development of novel diagnostic assays for muscular pathologies and shed light on the roles of EVs in diverse tissue environments.

KEYWORDS

Manuscript text

Introduction

The majority of studies on EVs conducted thus far have mainly focused on EVs isolated from liquid matrices such as cell culture media, blood, and urine. This is ascribable to the fact that these samples are readily available and can be processed using established separation techniques[1,2]. As a result, our understanding of EVs has been limited to those found in these particular matrices.

In the latest years, the important roles of EVs in solid tissues[3,4], including human and non-human brains[5–10], cardiac[11] and skeletal muscles[12,13,14], liver[14,15], adipose tissue[16,17] and several tumors[18–21], are emerging.

The study of solid tissue EVs is expected to offer a glimpse into the EVs in the microenvironment where they are released. Moreover, it provides the potential to comprehend the mechanisms through which EVs, released in the extracellular matrix of a tissue, exert a biological influence on their cellular targets, which may be located in various organs. By investigating solid tissue EVs, we can gain insights into the complex interplay between EVs and their surrounding environment, shedding light on the underlying processes and effects they have on cells.

This increases the possibility to find specific signatures of pathological conditions within organs. By analyzing the distinct profiles of tissue-EV associated with various diseases, researchers can uncover biomarkers that serve as diagnostic indicators. This knowledge could aid in early disease detection but also provides a basis for developing novel liquid biopsy assays[20,22] using blood[23] or other biological fluids[24].

However, this perspective is at its infancy due to technical challenges[4] and the lack of rigorous approaches that allow comparing EVs from solid and liquid tissues, minimizing bias due to different separation techniques and highlighting differences due to the EV microenvironments.

Common techniques applied to separate EVs from liquid matrices imply several combinations of differential (ultra)centrifugation, density gradient, size exclusion chromatography, fluid-flow based separation, affinity-based separation[1] and microfluidics techniques[25]. Recently developed EV separation methods from solid tissues mainly involve tissue homogenization or tissue digestion with different enzymes followed by differential (ultra)centrifugation, filtration[3] and density gradients[4]. However, it is more and more evident among EV community that "the product is the process"[26,27], indeed the application of different separation methods inevitably leads to the recovery of different EV subpopulations, in terms of physical properties[28,29] (i.e. number density, size distribution,

purity from exogenous contaminants), biomolecular content[30–32] (i.e. proteins, metabolites, nucleic acids, biomolecular corona) and therefore biological activity[29,33.] In turn separation procedures affect EV functionality both *in vitro*[34] and *in vivo*[35] and this might be due also to the presence of EV co-isolates[26].

Therefore, there is the need of universal EV separation protocols that can be equally applied to liquid and solid matrices in order to compare EVs from these two different sources without the biases that inevitably different separation methods introduce.

This would allow the exploration of how the biomolecular and biophysical properties of EVs may vary between those present in the tissue environment and those circulating, which remains poorly investigated.

To this aim, we have chosen to analyze EVs from human skeletal muscle (SkM) and human plasma derived from blood. These samples have a great clinical relevance: SkM is not only the largest organ in the human body, playing a central role in whole-body energy metabolism, but also acts as a secretory organ, producing and releasing hundreds of products, including myokines and EVs (SkM-EVs) that can have autocrine, paracrine, and endocrine effects[12,13,36,37]. Plasma is the most studied source of EVs from bio-fluids[1,2].

Previous studies that compared EVs from plasma and tissues (particularly skeletal muscle) relied on EVs collected from tissues cultured *ex vivo*[13,38]. However, this approach introduces new variables related to the culture conditions, meaning that EVs isolated *ex vivo* may not accurately represent those directly obtained from the tissue's extracellular matrix. Another attempt used immunoaffinity-based separation methods[12], hence narrowing the analysis on subpopulations presenting specific surface markers. In addition, these studies have been conducted on animal models, especially rodents, but the translation of those data to human EVs have never been demonstrated.

To minimize modifications induced by differences in the separation process we standardized the procedure to the greatest practicable extent, optimizing the protocol to be applied universally both

to liquid and solid starting materials. This combined method allowed to obtain reproducible and comparable set of EVs from the two different sources. Indeed, SkM-nanoparticles were released from the extracellular matrix in a liquid phase, using mechanical and enzymatic dissociation. EVs from tissue and plasma were, then, enriched using a consistent protocol involving serial ultra-centrifugations and discontinuous sucrose gradient allowing to enrich EVs according to their density and size, general properties conserved in different subtypes of EVs.

We, then, characterized the preparations using orthogonal collective and single-particle analysis technique, dissecting sample characteristics at multi-scale levels: from the molecular scale (immunoassays to define their biochemical composition), to the meso-scale (determining samples colloidal properties in terms of purity from exogenous proteins, nano-mechanical features, concentration, size distribution and stability).

This allowed to define "on a common reference" specific biochemical and biophysical properties of EVs secreted by SkM cells and released in the interstitial space[5,7,12,19] in respect to EVs present in plasma, the intercellular matrix of blood. These characteristics have never been described and compared before for EVs from human SkM tissue and plasma.

Studies in this direction might be useful in understanding open questions in EV biology, for example if EVs circulating in the blood stream present different properties than EVs present in the extracellular matrix of different tissue. It has been demonstrated that EVs deriving from healthy or pathological tissues can be detected in circulation[38–41], which raises important considerations about how they enter the bloodstream and whether they need to acquire specific characteristics or undergo changes in their properties after entering circulation.

All this information could help in elucidating the EV roles in various physiological and pathological processes and to develop novel, non-invasive diagnostic and monitoring assays for many diseases including muscular pathologies, hereditary diseases such as metabolic myopathies, dystrophic

myopathies or acquired conditions as Intensive-care acquired weakness in critically ill patients with sepsis[42,43].

MATERIALS AND METHODS

All data were acquired and described following the MIRABEL[44] , MISEV 2018[2] and MISEV2023[1] international guidelines. We have submitted all relevant data of EV samples to the EV-TRACK[45] database (EV-TRACK ID: EV240003) and MIBlood-EV[46] (see Fig.1SI)

Subject Recruitment and Sample Collection

In this study, 5 plasma samples and 5 skeletal muscle (SkM) tissue biopsies were obtained from subjects hospitalized in the Intensive Care Unit (n=3) and Orthopedic ward (n=2) of the Spedali Civili University Hospital in Brescia, Italy. The Ethics Committee of Brescia approved the study in adherence with the Declaration of Helsinki and Good Clinical Practice (Protocol number 34 68). All traceable identifiers were removed before analysis to protect patient confidentiality; all samples were analyzed anonymously. Written informed consent was obtained from the patient before the collection of samples. The subjects were all male, with an age range of 53-75 years (average age 64,6 y/o). Both blood and SkM tissue biopsies were collected at the same time and processed within 30 minutes from the withdrawal.

Plasma was obtained as previously described from peripheral blood samples in EDTA[47]. Blood was collected through a peripheral venous catheter in one of the main venous trunks of the forearm and SkM tissue was collected from quadriceps femoris. See Supplementary Materials for more details.

Enrichment of EVs from human plasma

Plasma EVs were isolated through serial ultra-centrifugation and discontinuous sucrose gradient (DSG). All steps were performed at 4°C. Briefly, 1 ml plasma was centrifuged at 300 x g for 10 min. Supernatant (1 ml) was transferred to a new tube and centrifuged at 2,000 x g for 20 min.

Supernatant (1 ml) was transferred to a new tube and centrifuged at 16,500 x g for 20 min and finally supernatant was transferred to an appropriate tube and centrifuged at 118,000 x g for 2h and 30 min. The 16,500 x g centrifugation step allows sedimentation of Plasma 16k sample, while the 118,000 x g ultracentrifugation step allow sedimentation of Plasma 118k sample.

DSG was carried out as follows: Plasma 16k and Plasma 118k pellets, obtained as described above, were further processed adapting the protocols developed in previous studies[30,47]. Briefly, Plasma 16k or Plasma 118k samples were re☐suspended in 1,000 µl buffer A (10 mM Tris☐HCl 250 mM sucrose, pH 7.4), loaded on top of a discontinuous sucrose gradient 15% (600 µl), 20, 25, 30, 40, 60, 65% (400 µl), 70% (800 µl) sucrose in 10 mM Tris☐HCl, pH 7.4) and centrifuged at 230,000 x g for 16 h. For Western blot (WB) analysis of fractions 1-12 (250 µl) were precipitated by incubation with 10% trichloracetic acid for 4 h as described in Paolini et al.[48]. For all the other analyses, fractions from 6 to 9 (750 µl each) were ultracentrifuged and resupended in 100 µl HPLC water and aliquoted for further analyses. See Supplementary Materials for more details.

Enrichment of EVs from human SkM tissue

EVs were isolated from human SkM tissue after mechanical and enzymatic dissociation, as previously described[19] with some modifications. Frozen tissue was thawed 2 min at r.t., washed one with cold PBS 1x and weighted (5 samples, mean 0.89 gr). Tissue was transferred in a 100 mm dish with cold RPMI (medium/tissue ratio: 1ml RPMI/ 0.2 gr tissue) and gently cut into small pieces (2x2x2mm) with a scalpel. Tissue pieces were transferred in a 6-well plate with 0.2 gr of SkM tissue each well. RPMI medium was added to each well to reach 2 ml of total medium. SkM tissue was inoculated with DNase (40U/ml final concentration) and collagenase D (2 mg/ml final concentration) at 37°C for 30 minutes in gently rotation (78 rpm). Medium derived after the enzymatic dissociation was filtered in a 0,70 µm pore strain in order to eliminate the largest debris. The filtrate was used for EV-enrichment through serial ultra-centrifugation and DSG as described for plasma.

Tissue filtrate was centrifuged at 300 x g for 10 min. Supernatant was transferred to a new tube and centrifuged at 2,000 x g for 20 min. Again, supernatant was transferred to a new tube and centrifuged at 16,500 x g for 20 min in order to obtain Tissue 16K preparation. Finally, supernatant was ultracentrifuged at 118,000 x g for 2 h and 30 min to get Tissue 118K preparation. All centrifugation steps were performed at 4°C.

DSG: Tissue 16k and Tissue 118k pellet obtained were treated as described for plasma preparations. For WB analysis of gradient fractions see section above.

SkM tissue homogenate was obtained from SkM tissue before (THB) and after the mechanical and enzymatic dissociation (TH). See Supplementary Materials for more details.

SDS-PAGE and WB analysis

SDS–PAGE and WB were performed by standard procedures[49] Sucrose density gradient fractions from 1 to 12 (20 μl of each fraction) were loaded on acrylamide–bisacrylamide, 12.5% gel. Total plasma or tissue homogenate (30 μg, protein content determined by Bradford assay) were loaded in the same gel as positive control of antibody reactivity in Western blot. Samples were electrophoresed and analyzed by WB with the antibodies described in the text. See Supplemental Material for more details.

COlorimetric NANoplasmonic (CONAN) assay

The EV-enriched preparations (DSG fractions 6-9) were checked for purity from protein contaminants using the COlorimetric NANoplasmonic (CONAN) assay[50,51]. EVs were resuspended in 100 μl of HPLC water. Two microlitres of EV solution at serial dilutions in HPLC water (1:1, 2μl of starting sample – 1:10, 2 μl of sample diluted in HPLC H_2O – 1:100, 2 μl of sample diluted in HPLC water) were resuspended in 23 μl of water, mixed with 50 μl of gold nanoparticles 6 nM and 25 μl of PBS. The result of the assay was collected on Ensight Multi Mode Reader (Perkin Elmer). Measurements were performed for each sample in triplicate.

Bicinchoninic acid (BCA) and Bradford assay

Protein concentrations of samples (fractions 6-9) were determined with a Pierce™ BCA Protein

Assay Kit (ThermoFisher, Rockford, USA), and Bradford assay (Biorad) following the

manufacturer's instructions.

Atomic Force Microscopy (AFM) analysis

Quantitative AFM morphometry was performed as described elsewhere[52,53]. Five μL aliquots from

samples were deposited on glass coverslips previously functionalized with 0.01 mg/mL poly-L-

lysine and left to adsorb for 60' at 4°C. Aliquots were progressively diluted up to 200x in

successive depositions in order to maximise the surface density of isolated particles. Imaging was

performed in ultrapure water at room temperature on a Bruker Multimode8 AFM equipped with a

Nanoscope V controller, a sealed fluid cell and a type JV piezoelectric scanner using Bruker

ScanAsystFluid+ probes. Background subtraction was performed using Gwyddion[54] 2.61.

Quantitative morphometry was performed with custom Python scripts to recover the surface contact

angle (CA) and equivalent solution diameter (D) of several hundred individual objects for each

sample. The relative particle concentrations of samples were estimated as described in

Supplementary section.

Nanoparticle Tracking Analysis (NTA)

Pellets from 6 to 9 sucrose fractions obtained from both plasma and human SkM tissue were pooled

together, resuspended in 100 μl of sterile HPLC water and for the characterization by Nanoparticles

Tracking Analysis (NTA) using a NanoSight NS300 system (Malvern, Panalytical LTD, Malvern,

UK) to evaluate the concentration and size distribution of EVs. See Supplemental Material for more

details.

Dynamic light scattering (DLS) analysis and zeta potential measurement

DLS at $\theta = 90°$ and zeta potential measurements were performed using a Brookhaven Instrument 90 Plus (Brookhaven Instruments Corporation, Holtsville, NY). For DLS, each measurement was an average of four replicates of 1 minute each. The autocorrelation functions were analysed by Laplace inversion using the CONTIN algorithm, allowing an estimate of the particles size distribution. Zeta potentials were obtained from the electrophoretic mobility u, according to the Helmholtz-Smoluchowski equation: $\zeta = (\eta/\varepsilon) \times u$ where η is the viscosity of the medium and ε is the dielectric permittivity of the dispersing medium. The zeta potential values are given as the average of ten measurements, performed in water.

Statistical analysis.

Significant differences among datasets were determined with Student's t-test (Graph Pad). P values of less than 0.05 were considered statistically significant with $**p < 0.01$ $***p < 0.001$ and $****p < 0.0001$. Values were shown as mean ± standard error of the mean (SEM) of at least 3 experiments.

RESULTS AND DISCUSSION

Universal protocol for the enrichment of nanoparticles from human plasma and SkM tissue biopsies

Peripheral blood and SkM tissue biopsies were collected from 5 subjects. Blood was processed as described in Methods section to obtain plasma. SkM tissue was washed twice in cold solution NaCl 0.9% and sample tube was snap-frozen in liquid nitrogen. Both plasma and SkM tissue were stored at -80°C. Then, Plasma and SkM tissue biopsies were processed with a newly optimized combined protocol (Fig.1).

Figure 1. Rationale of the paper sketched. Enrichment and characterization of EVs from human plasma and SkM tissue. After isolation, samples were characterized for their biomolecular and biophysical properties. Icons made by Freepik and ITM2101 from www.flaticon.com.

SkM biopsy (mean weight 0.89 gr) was thawed and washed with cold PBS. The biopsy was then incubated in RPMI medium and processed through gentle mechanical and enzymatic dissociation with DNase and Collagenase for 30 min at 37°C. This process allowed to release nanoparticles embedded in the interstitial space surrounding tissue cells and to diffuse them in the media during the procedure[19]. The media containing SkM-EVs was filtered through a 70 μm pore cell strainer in order to eliminate the largest tissue debris. At this stage the same protocol for the EV isolation from

plasma and SkM was used. The SkM filtrate and 1ml of plasma were centrifuged at 300g for 10 min, the supernatant (SN), collected without disturbing the pellet, was centrifuged at 2,000g for 30 min. Then, the subsequent SN was centrifuged at 16,500g for 20 min. The subsequent pellet is referred in the text as the Plasma or Tissue 16k subpopulations. Finally, the SN obtained after the 16,500g centrifugation was centrifuged at 118,000g for 2h 30 min to obtain pellets referred in the text as Plasma or Tissue 118k subpopulations.

It is recognized that pellets derived from both 16k and 118k Plasma and Tissue are composed by a complex mixture of different biogenic nanoparticles[55]: extracellular particles (EP- i.e. extracellular vesicles), non-vesicular extracellular particles (NVEP- i.e. lipoproteins, exomeres[56]), cellular and tissue small debris. In order to refine the separation of EPs, all four samples were loaded, separately, on top of a discontinuous sucrose gradient (DSG) and centrifuged at 230,000g for 16h. This allows the separation of different nanoparticles both for their size and density[30,47,57,58]. Twelve fractions of equal volume (400 µl) were collected, processed as described in Methods section and analyzed as further described.

Selection of Plasma and Tissue-EV positive fractions by biochemical characterization

Western Blot analysis was conducted for the biochemical characterization of the twelve fractions deriving from the processing of the Plasma 16k, Plasma 118k, Tissue 16k and Tissue 118k pellets by DSG. This allowed to determine the gradient fractions enriched in EVs and fractions enriched in NEVPs markers. Different proteins were selected according to MISEV 2018 guidelines[2] and recent studies[59]: EV markers (tetraspannins CD63, CD81; protein involved in the vesicular trafficking such as Flotillin1; single-pass transmembrane protein such as ADAM10; component of the ESCRT I complex such as TSG101); lipoproteins markers (ApoB marker of Low-density and Very-Low density Lipoproteins -LDL and VLDL; APOA1 marker of High Density Lipoprotein-HDL); SkM cell and SkM-EVs[12,13] markers (Caveolin-3 and Beta-enolase); markers of co-isolates/contaminants[1,26] (hemoglobin A for blood; Fatty Acid binding protein (FABP) for adipose

tissue[60]; GM130 and Calnexin for intracellular compartments: the Golgi network and the endoplasmic reticulum respectively).

As shown in Fig. 2, fractions 4 and 5 (red box, corresponding to a density of 1.09-1.11 g/cm^3) resulted enriched in HDL marker ApoA1 both in Plasma 16k (Fig. 2A) and in Plasma 118k (Figure 2B) samples, while ApoB resulted enriched in fraction 4 and 5 for Plasma 118k preparation, and slightly present in fraction 6 and 7 in Plasma 16k preparation. Differences in lipoprotein distribution in gradient fractions could be due to donor-to-donor variations. On the other hand, both Plasma 16k and Plasma 118k fractions from 6 to 9 (corresponding to a density of 1.13-1.29 g/cm^3) resulted enriched in EV markers, but depleted of blood contaminants such as lipoproteins and hemoglobin A (Supplementary Figure 2-6). Fraction from Plasma samples showed weak signals from markers of the intracellular compartments GM130 and Calnexin, and only in fraction 6 (Fig. 2A, B).

Figure 2. Biochemical characterization of Plasma samples after sucrose gradient fractionation.

Equal volumes (20 μL) of each fraction both for Plasma 16k (A) and 118k (B) were loaded, 30 μg

of Plasma total protein (PL) were loaded as positive control. Samples were electrophoresed on

SDS-PAGE gel (12% (Acrylamide/Bis-acrylamide) and analyzed for the antibodies described in the figures. Un-cropped WB are available in the Supplementary Information (Fig.2-5SI). Each fraction density is indicated in g/cm^3. Red box: fractions discarded for further analyses; Green box: fractions collected and further processed as a single sample.

Similarly, fractions 6 to 9 of Tissue 16k (Fig. 3A) and Tissue 118k (Fig. 3B) samples, after DSG processing, resulted enriched in EV markers such as ADAM10, Flotillin-1, TSG101 and CD81. Fractions analyzed still retain the signal of surface proteins such as TSG101[61], transmembrane proteins such as CD81 and ADAM10[62]. These data indicate that the mechanical and enzymatic treatment of the SkM tissue did not alter the EV surface protein profile and we observed an enrichment of these proteins in the EV preparations as expected[63].

In addition, fractions 6-9 resulted enriched in SkM cell marker proteins such as Caveolin-3 and beta-enolase demonstrating the tissue-specific origin of EVs in the preparations[12]. Signal intensity of Caveolin-3 in Plasma (Fig. 2A and B, PL) is lower than the signal intensity in Tissue homogenate (Fig. 3A and B, TH) and a similar trend is reflected in gradient fractions from Plasma (Fig.2A,B) and Tissue samples respectively, indicating that, in this experimental conditions, Caveolin-3 could be considered a potential specific SkM-EV marker[12,14,37].

Figure 3. Biochemical characterization of Tissue samples after sucrose gradient fractionation.
Equal volume (20 µL) of each fraction both for Tissue 16k (A) and 118k (B), 30 µg of Tissue
Homogenate (TH) were loaded as positive controls. WB for the antibodies described in the figures.
Un-cropped WB are available in the Supplementary Information (Fig 6-8SI). Each fraction density
is indicated in g/cm^3. The green box indicates fractions that, for further analyses, have been
collected and further processed as a single sample.

Furthermore, in fraction 6-9 signal of lipoprotein markers and adipose tissue markers were not present (Supplementary Information- SI-, Fig. 6SI ,7SI), indicating a negligible blood and adipose tissue contaminants in the SkM tissue biopsies. Tissue samples retain some amount of cellular marker Calnexin[7,22] in fraction 6 to 9, even if in opposite tendency to the signal of EV markers. Quantification of the chemiluminescent signal of Calnexin in SkM tissue homogenate before (THB) and after (TH) enzymatic and mechanical treatment did not change significantly (Fig. 8SI), indicating that the processing did not alter tissue protein composition or induced significant tissue disruption. It is to note that, in our experimental conditions, the chemiluminescent signal of GM130 was not detectable neither in homogenates obtained from the SkM tissue before (data not shown) and after the enzymatic treatment, nor in the gradient fractions, indicating that probably this marker is unsuitable for determining the presence of intracellular co-isolates in human muscle samples, unlike SkM tissue from other species[12] (Fig. 6SI, 7SI).

According to the results described above, we decided to focus our attention to fractions from 6 to 9 (Fig. 2 and 3, green box) for all four preparations: these fractions are enriched in EV markers and contain nanoparticle with the typical density of EVs[30,47,57,58]. We discarded fractions 4 and 5 (Fig. 2, 3 red box) that still contain EV markers, but have lower density and, especially in plasma samples, are also enriched in lipoprotein markers. For further analyses fractions 6-9 have been collected and analyzed as a single sample and will be further referred as Plasma 16k fractions 6-9 post gradient (PL16kG), Plasma 118k fractions 6-9 post gradient (PL118kG), Tissue 16k fractions 6-9 post gradient (TS16kG), Tissue 118k fractions 6-9 post gradient (TS118kG).

Pure Tissue samples from exogenous protein contaminants have a higher protein yield than Plasma Samples

Preparations PL16kG, PL118kG, TS16kG, TS118kG were checked for their purity from exogenous protein contaminants (EPC) with the COlorimetric NANoplasmonic (CONAN) assay as previously described[50,51,64]. Serial dilutions (1:1, 1:10 and 1:100 in HPLC water) of the samples were analyzed: PL16kG and PL118kG (Fig. 4A) resulted with an AI% lower than 20% at the minimal dilution of the sample (1:1) and remained below the 20% (dotted line) even at higher dilutions, but with increasing AI%. This represents the trend of samples pure from any detectable EPC (EPC content ≤0.05 μg/μl)[47,51,65,66] where the starting concentration of EVs is optimal for the interaction with gold nanoparticles (AuNPs) and further dilution results in an increase of the AI%. TS16kG and TS118kG (Fig.4B) presented an AI% higher than 20% at the dilution 1:1, while it decreased under the 20% with the serial dilution 1:10 and 1:100, meaning that the samples are pure from EPC, but a further dilution is needed to fall the AI below the 20% dotted line. These data indicate that our experimental procedures allowed to obtain EVs from SkM tissue pure from EPC, and indicate that both Tissue-EV preparations are more concentrated than Plasma-EV preparations, since TS16kG and TS118kG reached an AI% below 20% after a dilution 1:10 of the starting material. The increase of AI% in further dilutions indicates that samples reached the optimal EV concentration for AuNPs interaction at dilution 1:10, while at dilution 1:100 we are shifting from the optimal range in terms of EV concentration to interact with AuNPs resulting in an increase of the AI%.

Figure 4. Purity assessment and protein concentration measurements. Fractions 6 to 9 after the

sucrose density gradient were collected and analyzed. Samples derived from Plasma (PL16kG,

PL118kG) or Tissue (TS16kG, TS118kG). Samples were checked for purity from exogenous

protein contaminants (EPC) with the CONAN assay. Aggregation Index ratios (AI%) of serial dilution (1:10, 1:100) of a representative sample are showed in the graph. The dotted line represents the CONAN assay threshold for EPC detection (AI% <20% EPC content ≤0.05 µg/µl). The AI% is inversely proportional to the preparation purity and samples with AI% below the 20% are considered pure. AI% values of a representative sample with assay internal controls (NormREF, black dots; IntREF, green dots) (A); Protein concentration with BCA (B) and Bradford assay (C); Protein concentration for each group was normalized by tissue mass (per 1 gr). Values were shown as mean ± SEM. Significant differences among datasets were determined with Student's t-test. P values: ****p < 0.0001. TS16kG, dark blue points; PL16kG, dark red squares; TS118kG, light triangle up; PL118kG, light red triangle down.

After the purity check with the CONAN assay we performed Bicinchoninic acid (BCA) and Bradford assays to determine the protein concentrations in the samples (very likely associated to EVs). In order to compare the protein yield of samples from SkM tissue and plasma, the concentration was normalized to one gram of starting material[7,19,38,67] (1 ml of Plasma and 1 gr of Tissue). BCA and Bradford results were comparable and in accordance to CONAN results. Tissue samples presented a significant higher protein yield than Plasma samples both with 16kG and 118kG preparations measured with BCA (Fig. 4C). It is to note that for the Bradford assay, these highly purified with low-yield samples challenged the assay sensitivity. Consequently, the protein concentration exhibited wide Standard Error of the Mean (SEM), especially in TS16kG and TS118kG, and in the case of the PL16kG sample, it was undetectable (Fig. 4D). However, despite these challenges, the trend of higher protein yield in Tissue samples compared to Plasma samples persisted, as indicated by BCA measurements.

Tissue and Plasma EV-enriched fractions display heterogeneous nanomechanical properties.

The morphology and nanomechanical characteristics of nanoparticles present in PL16kG, PL118kG, TS16kG and TS118kG preparations were evaluated using a high-throughput AFM imaging approach described previously[52,68]. Preliminary qualitative inspection of the resulting Atomic Force Microscopy (AFM) micrographs revealed that all samples contained abundant globular particles with a morphology broadly compatible with that typically displayed by EVs (Fig. 5A,B,C,D).

Figure 5. Nano-mechanical features of samples from Plasma and Tissue**.** Representative AFM

images of PL16kG (A), PL118kG (B), TS16kG (C), and TS118kG (D). Contact angle VS diameter

plot of several hundred individual particles from at least five different depositions for each sample as calculated via AFM morphometry of PL16kG (E), PL118kG (F), TS16kG (G) and TS118kG (H). Blue box represents area populated by structures with the typical EV-like characteristics: contact angle ranging from 75° to 135°, diameter from 30 to 300 nm.

We recently demonstrated that quantitative AFM morphometry can discriminate different subpopulations of nanoparticles in the same preparation on the basis of their nanomechanical properties. In particular, the stiffness of individual putative vesicles adhered to the substrate can be estimated via their equivalent spherical cap contact angle (CA). In samples derived from plasma, this technique was shown to discriminate EVs from NVEPs such as lipoproteins[53]. Briefly, since CAs were shown to be heavily influenced by the nanomechanical properties of their parent particle (*e.g.* the stiffness of intact EVs), collecting the morphometric measurements of individual objects from the same sample on a CA versus Diameter (D) plot results in the clustering of different subtypes of EVs and NVPs in specific zones of the plot[53].

Accordingly, the CA and D values of several hundred individual particles from PL16kG, PL118kG, TS16kG and TS118kG preparations were collected to evaluate the particle heterogeneity of each sample (Fig.5). All samples exhibited the typical log-normal distribution of sizes, containing abundant globular particles with D values below the range typically associated with intact EVs (<30 nm) and highly dispersed CA values surpassing those previously associated with lipoproteins (> 135°); although it is impossible to determine the biological identity of these particles via AFM, their CA and D ranges might be compatible with protein aggregates and/or other NVEPs. Given their higher stiffness compared to liposomes, they could be protein-enriched NEVPs, similar to exomeres[56].

Their overall similarity notwithstanding, the main differences between the four samples are found in the relative abundance, size and stiffness of particles with diameters above 50 nm. As shown in Fig.

5E, PL16kG was found to contain a large fraction of particles within typical CA and D ranges previously associated with intact EVs from a variety of sources[52,53,68] (blue box, CA=80°-130° and D>30nm). On the other hands, the regions with lower CA (<80 nm), ascribed to other biogenic nanoparticles such as lipoproteins, was practically devoid of particles, confirming results from WB. The corresponding PL118kG preparation (Fig. 5F) also contained a rich population of EV-compatible particles and a marginal presence of particles with lower CA values. We then applied the same analytical procedure to TS16kG (Fig. 5G) and TS118kG (Fig. 5H) samples. Despite being very similar to the Plasma preparations with abundant particles within typical CA and D ranges previously associated with intact EVs (blue box), both TS16kG and TS118kG samples showed a higher abundance of softer particles (CA below 80°, below the blue box) in the fraction of sizes above 50 nm. This means that the EV subpopulations in the samples enriched from SkM tissue present a more heterogeneous CA than EVs enriched from Plasma.

In addition, as outlined in the methods section, particle surface density calculations from AFM images allow estimating relative particle concentrations across different samples. With this method, the concentration of EVs in samples enriched from tissue (TS16kG and TS118kG) was found to be on average 10^2 times higher than the corresponding preparations from PL16kG and PL118kG as suggested by CONAN, BCA and Bradford assays (SI, Table 1).

Tissue samples have higher particles concentration, broader size distribution and higher colloidal stability than Plasma samples.

The analytical techniques described earlier provide an estimation of whether one sample is more concentrated than another in terms of nanoparticles and proteins. To confirm these data, we compared the particle concentration in the samples by Nanoparticle Tracking Analysis (NTA). Indeed, NTA results confirmed the data from CONAN, protein quantification and AFM, indicating a significant higher particle yield in TS16kG samples (mean concentration $6.3*10^{10}$ particles/gr of tissue) than PL16kG samples (mean concentration $2.3*10^{10}$ particles/gr of plasma) and TS118kG

(mean concentration 5.5×10^{10} particles/gr of tissue) samples respect to PL118kG samples (mean concentration 1.7×10^{10} particles/gr of plasma). It is to note that, to facilitate the comparison of particle yield between Plasma and Tissue samples, data have been normalized to one gram of starting material (Fig. 6A). Despite the differences in nature of the two, in our experimental conditions, all the orthogonal techniques implied confirmed that the EV yield from SkM is significantly higher than Plasma for both subpopulations analyzed.

Figure 6. Concentration and size distribution. Particle concentration for each sample were

measured by NTA. Particle concentration was normalized for starting material mass (1gr) (A).

Particle Size distribution was measured by NTA. Histograms represent the mean of the size distribution divided by range (30-200 nm; 201-400 nm; 401-600 nm) of the 16k and 118k samples from plasma and tissue B). Particle size was analyzed by NTA and modal distribution (nm) (C). Values in nm were shown as mean ± SEM. Significant differences among datasets were determined with Student's t-test. P values: **p< 0.01, ***p < 0.001 and ****p < 0.0001.

Samples size distribution was determined with orthogonal techniques: NTA, Dynamic Light Scattering (DLS) and AFM. Each method has its advantages and limits and the combined evaluation of data obtained with different approaches allows to reveals different aspect and characteristics.

The size of particles was analyzed using NTA and mean size distribution profiles of preparations are shown in Fig. 6B. According to these measurements, both TS16kG and TS118kG samples have a broader size distribution than PL16kG and PL118kG respectively, with presence of EVs in the range from 30-200 nm, from 201 to 400 nm and also from 401 to 600 nm, while the majority of EVs from Plasma samples are in the range from 30-200 and 201-400 nm (Figure 6B). The mode size was found to be significantly different from TS16kG and PL16kG samples, with TS16kG EVs diameter larger than PL16kG EVs (166 ± 4 nm and 131 ± 3 nm respectively, Fig. 6C) while no significant differences were found in the mode size of TS118kG and PL118kG preparations (148 ± 4 nm and PL118kG 143 ± 4 nm respectively, Fig. 6C). NTA results data indicated that in TS16kG preparation the mode size of EVs is significantly higher than TS118kG sample.

The Laplace inversion analysis using the CONTIN algorithm was applied to autocorrelation functions obtained through DLS for four preparations (PL16kG, PL118kG, TS16kG, and TS118kG, Fig. 7A,D,G,L) to qualitatively confirm the presence of submicrometer particles and possible aggregates that may be present in EV samples[69]. The DLS profiles show a polydisperse size distribution, likely arising from a combination of the scattering from single and aggregated EVs, as well as from the scattering from objects of similar refractive index (e.g., proteins). The scattered

intensity-weighted size distribution, depicted in Fig. 7B,E,H,M, provides a reliable analysis without assuming the composition of different populations. The results indicated that PL16kG and TS16kG samples are characterized by the presence of objects ranging from hundreds of nanometers to micrometers in diameter. In contrast, PL118kG and TS118kG preparations exhibit bi-modal size distributions with a small subpopulation (around 100 nm) and larger aggregates (diameter > 400 nm). It should be noted that DLS technique inherently tends to overestimate the presence of larger objects compared to smaller ones[69]. Therefore, a number-weighted distribution is also presented (Fig. 7C,F,I,N). In this scale, the majority of the scattering objects are around 100 nm diameter. This observation is consistent with the presence of small objects such as EVs, as corroborated by results from NTA and AFM.

Figure 7. Size distribution with DLS. Representative autocorrelation function of the scattered

intensity, with the fitting curve according to the Laplace Inversion via CONTIN algorithm

(A,D,G,L); corresponding scattered intensity-weighted (B,E,H,M) and number-weighted (C,F,I,N) size distributions of scattering objects of samples described in the text.

Size distributions obtained via AFM morphometry can be inferred from the abundance of individual particles along the horizontal (D) axis of plots in Fig. 5E-5H. While a direct quantitative comparison of sizes distributions obtained from AFM and NTA is often questionable, the general trends observed by both techniques coincide: both AFM and NTA attribute to TS16kG EVs a higher diameter mode than PL16kG EVs (Fig. 5G and 5E, respectively), while those of TS118kG and PL118kG did not differ significantly (Fig. 5H and 5F, respectively). AFM also attributes to TS16kG EVs (Fig. 5G, blue box) a more heterogeneous size distribution than PL16kG EVs (Fig. 5E, blue box), with more particles populating the area of diameter between 100 and 300 nm in TS16kG sample, while PL118kG and TS118kG EVs (Fig. 5F and 5H, respectively) seem to be more homogenous.

Finally, we analyzed the colloidal stability of the four preparations measuring the zeta-potential. All measurements were done in milliQ H_2O. TS16kG presented a mean zeta potential of -34.2 mV, PL16kG -13.7mV, TS118kG -24.2 and PL118G -10.6mV (Fig. 8). Notably, differences in colloidal stability between Plasma and Tissue-EVs enriched samples emerged. Both PL16kG and PL118kG exhibited a significantly higher zeta-potential compared to TS16kG and TS118kG, suggesting lower stability. This disparity implies that plasma EVs may be more susceptible to interactions with other components circulating in blood (i.e. other biogenic entities such as lipoproteins[70]) than EVs in the extracellular matrix of SkM tissue. It will be interesting to evaluate if this property is specific of blood EVs at the moment of the secretion or if EVs acquired it during the stay in a such a complex biofluid, and if these differences affect the biological functionality of EVs. For example, plasma EVs, with their lower stability, might be more prone to aggregation or faster clearance from circulation, whereas more stable tissue EVs may persist longer in the interstitial space. Tissue EVs, on the other hand, maybe more stable in their local environment due to less interaction with foreign

biogenic components. Further analyses should be conducted to assess how would these influence the development of EV-based diagnostics, particularly in distinguishing between tissue and blood-derived EVs, as stability could influences EV bioavailability and detectability in biofluids[71].

Figure 8. Zeta-potential measurement. Preparations were analyzed for their colloidal stability measuring z-potential with DLS. Values in mV were shown as mean ± standard error of the mean (SEM). Significant differences among datasets were determined with Student's t-test. P values: **p< 0.01.

CONCLUSIONS

We presented a new universal method to separate EV from liquid and solid matrices, that allowed to compared, for the first time, properties of EVs from human plasma and skeletal muscle tissue. The combination of differential (ultra)centrifugation steps and a top-down sucrose density gradient allowed the separation of EVs from their starting materials and to select gradient fractions that were positive for the presence of typical EV markers and cellular origin markers from SkM tissue. In addition, we demonstrated, with orthogonal techniques, that the selected gradient fractions are depleted from co-isolated NVEPs such as lipoproteins, and contaminants such as cellular debris and exogenous soluble proteins.

Furthermore, the characterization of the preparations showed differences in biomolecular and biophysical properties of EVs-enriched samples from human plasma and SkM. The observed differences regarding the nanomechanical properties, particle yield, size and colloidal stability are not biased by the separation method and thus we can confidently say that reflect specific characteristics of EVs-enriched samples from liquid and solid matrices. This poses new issues in understanding the determinants of these differences: the cellular origin, the molecular composition, the complex interaction between EVs and the biological environment. The last point highlights the importance of considering, in this scenario, the role of the biomolecular corona[64,72] the extravesicular cargo of molecules adsorbed to EVs, influenced by the surrounding matrix and that can change their physicochemical properties[73]. In addition, it will be interesting to compare the biophysical properties of EVs from different human tissues isolated with the same protocol to see if they are SkM tissue-specific or typical of all tissue derived EVs.

From a broader standpoint, our optimized method could be widely adopted in the EV field since it can be easily translated to other tissues. In this way, researchers may directly assess the traits that we highlighted to other solid and liquid matrices such as tumor tissues, urines and determine novel pivotal factors in the discovery of biomarkers[74]. Infact, the characterization of EV biophysical properties could serve, in the next future, as powerful diagnostic tools[73,74] for various pathological conditions, including, but not only related to SkM pathologies, complementing traditional biomolecular markers. Moreover, our results on stored tissue can be of interest for all researchers that can have access to existing biobanks thus increasing the number of samples that can be processed and used as source of EVs, and greatly increasing the possibility for developing EV-based diagnostic tools.

Acknowledgements

We thank Dr. Federica Bono from the Dept. of Molecular and Translational Medicine of the University of Brescia for the Ultra-Turrax T25 basic (IKA-Werke) homogenizer.

References

[1] Welsh, J. A. *et al.* Minimal information for studies of extracellular vesicles (MISEV2023): From basic to advanced approaches. *J Extracell Vesicles.* **13**, e12404 (2024).

[2] Théry, C. *et al.* Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *J Extracell Vesicles.* **7**, 1535750 (2018).

[3] Li, S.-R. *et al.* Tissue-derived extracellular vesicles in cancers and non-cancer diseases: Present and future. *J of Extracellular Vesicles.* **10**, e12175 (2021).

[4] Zhi, Z., *et al.* Research advances and challenges in tissue-derived extracellular vesicles. *Frontiers in Molecular Biosciences.* **9**, (2022).

[5] Perez-Gonzalez, R. *et al.* The Exosome Secretory Pathway Transports Amyloid Precursor Protein Carboxyl-terminal Fragments from the Cell into the Brain Extracellular Space. *J of Biological Chemistry.* **287**, 43108–43115 (2012).

[6] Vella, L. J. *et al.* A rigorous method to enrich for exosomes from brain tissue. *J of Extracellular Vesicles.* **6**, 1348885 (2017).

[7] Huang, Y. *et al.* Influence of species and processing parameters on recovery and content of brain tissue-derived extracellular vesicles. *J of Extracellular Vesicles.* **9**, 1785746 (2020).

[8] Brenna, S. *et al.* Characterization of brain-derived extracellular vesicles reveals changes in cellular origin after stroke and enrichment of the prion protein with a potential role in cellular uptake. *J Extracell Vesicles.* **9**, 1809065 (2020).

[9] Zhang, Z. *et al.* Comprehensive characterization of human brain-derived extracellular vesicles using multiple isolation methods: Implications for diagnostic and therapeutic applications. *J of Extracellular Vesicles.* **12**, 12358 (2023).

[10] Gomes, P. A. *et al.* A novel isolation method for spontaneously released extracellular vesicles from brain tissue and its implications for stress-driven brain pathology. *Cell Communication and Signaling.* **21**, 35 (2023).

[11] Liang, W. *et al.* Protocol to separate small and large extracellular vesicles from mouse and human cardiac tissues. *STAR Protocols.* **5**, 102914 (2024).

[12] Watanabe, S. *et al.* Skeletal muscle releases extracellular vesicles with distinct protein and microRNA signatures that function in the muscle microenvironment. *PNAS Nexus.* **1**, pgac173 (2022).

[13] Ismaeel, A. *et al.* Extracellular vesicle distribution and localization in skeletal muscle at rest and following disuse atrophy. *Skeletal Muscle.* **13**, 6 (2023).

[14] Matejovič, A. *et al.* Comparison of separation methods for tissue-derived extracellular vesicles in the liver, heart, and skeletal muscle. *FEBS Open Bio.* **11**, 482–493 (2021).

[15] Olofsson Bagge, R. *et al.* Three-dimensional reconstruction of interstitial extracellular vesicles in human liver as determined by electron tomography. *J Extracell Vesicles.* **12**, e12380 (2023).

[16] Jeurissen, S. *et al.* The isolation of morphologically intact and biologically active extracellular vesicles from the secretome of cancer-associated adipose tissue. *Cell Adh Migr.* **11**, 196–204 (2017).

[17] Crewe, C. *et al.* An Endothelial-to-Adipocyte Extracellular Vesicle Axis Governed by Metabolic State. *Cell.* **175**, 695-708.e13 (2018).

[18] Hurwitz, S. N. *et al.* Extraction of Extracellular Vesicles from Whole Tissue. *J Vis Exp.* **144**, 10.3791/59143 (2019).

[19] Crescitelli, R. *et al.* Isolation and characterization of extracellular vesicle subpopulations from tissues. *Nat Protoc.* **16**, 1548–1580 (2021).

[20] Jang, S. C. *et al.* Mitochondrial protein enriched extracellular vesicles discovered in human melanoma tissues can be detected in patient plasma. *J Extracell Vesicles.* **8**, 1635420 (2019). https://doi.org/10.1080/20013078.2019.1635420.

[21] Tassinari, S. *et al.* Profile of matrix-entrapped extracellular vesicles of microenvironmental and infiltrating cell origin in decellularized colorectal cancer and adjacent mucosa. *J of Extracellular Bio.* **3**, e144 (2024).

[22] Crescitelli, R. *et al.* Extracellular vesicle DNA from human melanoma tissues contains cancer-specific mutations. *Frontiers in Cell and Developmental Biology.* **10**, (2022).

[23] Cheng, L. *et al.* Small RNA fingerprinting of Alzheimer's disease frontal cortex extracellular vesicles and their comparison with peripheral extracellular vesicles, *J Extracell Vesicles.* **9**, 1766822 (2020).

[24] Dong, L. *et al.* Tumour tissue-derived small extracellular vesicles reflect molecular subtypes of bladder cancer. *J of Extracellular Vesicles.* **13**, e12402 (2024).

[25] Abreu, C. M. *et al.* Microfluidic platforms for extracellular vesicle isolation, analysis and therapy in cancer. *Lab Chip.* **22**, 1093–1125 (2022).

[26] Paolini, L. *et al.* Large⬚scale production of extracellular vesicles: Report on the "massivEVs" ISEV workshop. *J of Extracellular Bio.* **1**, e63; 10.1002/jex2.63 (2022).

[27] Buschmann, D. *et al.* Separation, characterization, and standardization of extracellular vesicles for drug delivery applications. *Advanced Drug Delivery Reviews.* **174**, 348-368 (2021).

[28] Jeon, H. *et al.* Effects of different separation methods on the physical and functional properties of extracellular vesicles. *PLOS ONE.* **15**, e0235793 (2020).

[29] Takov, K. *et al.* Comparison of small extracellular vesicles isolated from plasma by ultracentrifugation or size-exclusion chromatography: yield, purity and functional potential, *J Extracell Vesicles.* **8**, 1560809 (2019).

[30] Radeghieri, A. *et al.* Active antithrombin glycoforms are selectively physiosorbed on plasma extracellular vesicles. *J of Extracellular Bio.* **1**, e57 (2022).

[31] Tzaridis, T. *et al.* Extracellular Vesicle Separation Techniques Impact Results from Human Blood Samples: Considerations for Diagnostic Applications. *International Journal of Molecular Sciences.* **22**, (2021).

[32] Williams, Palviainen, C.M. *et al.* Metabolomics Applied to the Study of Extracellular Vesicles. *Metabolites.* **9**, 276 (2019).

[33] Paolini, L. *et al.* Residual matrix from different separation techniques impacts exosome biological activity. *Sci Rep.* **6**, 23550 (2016).

[34] Mol, E. A. *et al.* Higher functionality of extracellular vesicles isolated using size-exclusion chromatography compared to ultracentrifugation. *Nanomedicine: Nanotechnology, Biology and Medicine.* **13**, 2061–2065 (2017).

[35] Wolf, M. *et al.* A functional corona around extracellular vesicles enhances angiogenesis, skin regeneration and immunomodulation. *J of Extracellular Vesicles.* **11**, e12207 (2022).

[36] Rome, S. Muscle and Adipose Tissue Communicate with Extracellular Vesicles. *Int J Mol Sci.* **23**, 7052 (2022).

[37] Guescini, M. *et al*. Muscle Releases Alpha-Sarcoglycan Positive Extracellular Vesicles Carrying miRNAs in the Bloodstream. PLOS ONE. **10**, e0125094 (2015).

[38] Estrada, A.L. *et al*. Extracellular vesicle secretion is tissue-dependent ex vivo and skeletal muscle myofiber extracellular vesicles reach the circulation in vivo. Am J Physiol Cell Physiol. **322**, 246-259 (2022).

[39] Li, Y. *et al*. EV-origin: Enumerating the tissue-cellular origin of circulating extracellular vesicles using exLR profile. *Computational and Structural Biotechnology Journal*. **18**, 2851–2859 (2020).

[40] Su, Y., Chen, M., Xu, W., Gu, P. & Fan, X. Advances in Extracellular-Vesicles-Based Diagnostic and Therapeutic Approaches for Ocular Diseases. *ACS Nano*. **18**, 22793–22828 (2024).

[41] Mustapic, M. *et al*. Plasma Extracellular Vesicles Enriched for Neuronal Origin: A Potential Window into Brain Pathologic Processes. *Front. Neurosci*. **11**, (2017).

[42] Latronico, N., Rasulo, F. A., Eikermann, M. & Piva, S. Critical Illness Weakness, Polyneuropathy and Myopathy: Diagnosis, treatment, and long-term outcomes. *Crit Care*. **27**, 439 (2023).

[43] Lad, H. *et al*. Intensive Care Unit-Acquired Weakness: Not just Another Muscle Atrophying Condition. *Int J Mol Sci*. **21**, 7840 (2020).

[44] Faria, M. *et al*. Minimum information reporting in bio-nano experimental literature. *Nat Nanotechnol*. **13**, 777–785 (2018).

[45] Van Deun, J. *et al*. EV-TRACK: transparent reporting and centralizing knowledge in extracellular vesicle research. *Nat Methods*. **14**, 228–232 (2017).

[46] Lucien, F. *et al*. MIBlood-EV: Minimal information to enhance the quality and reproducibility of blood extracellular vesicle research. *Journal of Extracellular Vesicles*. **12**, 12385 (2023).

[47] Grossi, I. *et al*. MicroRNA-34a-5p expression in the plasma and in its extracellular vesicle fractions in subjects with Parkinson's disease: An exploratory study. *Int J Mol Med*. **47**, 533–546 (2021).

[48] Paolini, L. *et al*. The epsilon hinge-ear region regulates membrane localization of the AP-4 complex. *Traffic* **12**, 1604–1619 (2011).

[49] Salvi, A. *et al*. Analysis of a nanoparticle□enriched fraction of plasma reveals miRNA candidates for Down syndrome pathogenesis. *Int J Mol Med*. **43**, 2303–2318 (2019).

[50] Maiolo, D. *et al*. Colorimetric nanoplasmonic assay to determine purity and titrate extracellular vesicles. *Anal Chem*. **87**, 4168–4176 (2015).

[51] Zendrini, A. *et al*. Augmented COlorimetric NANoplasmonic (CONAN) Method for Grading Purity and Determine Concentration of EV Microliter Volume Solutions. *Frontiers in Bioengineering and Biotechnology*. **7**, (2020).

[52] Ridolfi, A. *et al*. AFM-Based High-Throughput Nanomechanical Screening of Single Extracellular Vesicles. *Anal. Chem*. **92**, 10274-10282 (2020).

[53] Ridolfi, A. *et al*. Particle profiling of EV-lipoprotein mixtures by AFM nanomechanical imaging. *Journal of Extracellular Vesicles* **12**, 12349 (2023).

[54] Nečas, D. *et al*. Gwyddion: an open-source software for SPM data analysis. *Open Physics* **10**, 181–188 (2012).

[55] Busatto, S. et al. The nanostructured secretome, Biomater. Sci. **8**, 39-63.

[56] Zhang, H. *et al*. Identification of distinct nanoparticles and subsets of extracellular vesicles by asymmetric flow field-flow fractionation. Nat Cell Biol. **20**, 332-343 (2018).

[57] Di Noto, G. *et al*. Enriched Serum Microvesicles Are Generated in Malignant Plasma Cell Dyscrasia. *PLOS ONE*. **8**, e70811 (2013).

[58] Di Noto, G. *et al*. Immunoglobulin free light chains and GAGs mediate Multiple Myeloma Extracellular Vesicles uptake and secondary NfkB nuclear translocation. *Frontiers in Immunology*. **5**, (2014).

[59] Jeppesen, D.K., Zhang, Q., Franklin, J.L. & Coffey, R.J. Extracellular vesicles and nanoparticles: emerging complexities. *Trends in Cell Biology*. **33**, 667-681 (2023).

[60] Yamamoto, T. *et al.* Classification of FABP isoforms and tissues based on quantitative evaluation of transcript levels of these isoforms in various rat tissues. *Biotechnol Lett* **31**, 1695–1701 (2009).

[61] Paolini, L. *et al*. Exosomes Secreted by HeLa Cells Shuttle on Their Surface the Plasma Membrane-Associated Sialidase NEU3. *Biochemistry*. **56**, 6401–6408 (2017).

[62] Kowal J. *et al*. Proteomic comparison defines novel markers to characterize heterogeneous populations of extracellular vesicle subtypes. *Proceedings of the National Academy of Sciences*. **113**, E968–E977 (2016).

[63] Lötvall, J. *et al*. Minimal experimental requirements for definition of extracellular vesicles and their functions: a position statement from the International Society for Extracellular Vesicles. *J of Extracellular Vesicles*. **3**, 26913 (2014).

[64] Musicò, A. *et al*. Surface functionalization of extracellular vesicle nanoparticles with antibodies: a first study on the protein corona "variable". *Nanoscale Advances* **5**, 4703–4717 (2023).

[65] Paolini, L. *et al*. Fourier-transform Infrared (FT-IR) spectroscopy fingerprints subpopulations of extracellular vesicles of different sizes and cellular origin. *J of Extracellular Vesicles*. **9**, 1741174 (2020).

[66] Mangolini, V. *et al*. Biochemical Characterization of Human Salivary Extracellular Vesicles as a Valuable Source of Biomarkers. *Biology*. **12**, 227 (2023).

[67] Huang, Y. *et al*. Brain Tissue-Derived Extracellular Vesicles in Alzheimer's Disease Display Altered Key Protein Levels Including Cell Type-Specific Markers. *J Alzheimers Dis*. **90**, 1057–1072 (2022).

[68] Borup, A. *et al*. Comparison of separation methods for immunomodulatory extracellular vesicles from helminths. *Journal of Extracellular Biology*. **1**, e41 (2022).

[69] Montis, C. *et al*. Size distribution of extracellular vesicles by optical correlation techniques. *Colloids and Surfaces B: Biointerfaces*. **158**, 331-338 (2017).

[70] Busatto, S. *et al*. Brain metastases-derived extracellular vesicles induce binding and aggregation of low-density lipoprotein. *Journal of Nanobiotechnology*. **18**, 162 (2020). https://doi.org/10.1186/s12951-020-00722-2.

[71] Rogers N.M.K. *et al*. Comparative electrokinetic properties of extracellular vesicles produced by yeast and bacteria. *Colloidal and Surfaces B: Biointerfaces*. **225**, 113249 (2023).

[72] Musicò, A. *et al*. Extracellular vesicles of different cellular origin feature distinct biomolecular corona dynamics. *Nanoscale Horiz*. **10**, 104–112 (2024).

[73] Radeghieri, A. *et al*. The biomolecular corona of extracellular nanoparticles holds new promises for advancing clinical molecular diagnostics. *Expert Rev Mol Diagn*. **23**, 471–474 (2023).

[74] Paolini, L. *et al*. Biophysical properties of extracellular vesicles in diagnostics, Biomark. *Med*. **12**, 383-391 (2018)

Authors' Contributions

CRediT taxonomy. VM Data curation, Formal analysis, Investigation, Methodology, Writing - review & editing; AR Conceptualization, Data curation, Formal analysis, Investigation, Funding acquisition, Writing - review & editing; SP Conceptualization, Data curation, Formal analysis, Funding acquisition, Writing - review & editing; SC Investigation, Methodology, Writing - review

& editing; NL Conceptualization; Data curation; Formal analysis; Funding acquisition, Writing - review & editing; MB Investigation, Methodology, Data curation, Writing - review & editing; FV Investigation, Methodology, Data curation; AB Investigation, Methodology, Writing - review & editing; CM Investigation, Methodology, Writing - review & editing; PB Conceptualization; Data curation; Formal analysis; Funding acquisition, Writing - review & editing; LP Conceptualization, Data curation, Formal analysis, Funding acquisition, Investigation; Methodology, Writing - original draft

Supplementary Materials

The datasets supporting the conclusions of this article are included within the article and its additional file named "Supplementary Materials".

Figure captions

Figure 1. Rationale of the paper sketched. Enrichment and characterization of EVs from human plasma and SkM tissue. After isolation, samples were characterized for their biochemical and biophysically properties. Icons made by Freepik and ITM2101 from www.flaticon.com.

Figure 2. Biochemical characterization of Plasma samples after sucrose gradient fractionation. Equal volume (20 μL) of each fraction both for Plasma 16k (A) and 118k (B) were loaded, 30 μg of Plasma total protein (PL) were loaded as positive control. Samples were electrophoresed on SDS-PAGE gel (12% (Acrylamide/Bis-acrylamide) and analyzed for the antibodies described in the figures. Un-cropped WB are available in the Supplementary Information (Fig.2-5SI). Each fraction density is indicated in g/cm^3. Red box: fractions discarded for further analyses; Green box: fractions collected and further processed as a single sample.

Figure 3. Biochemical characterization of Tissue samples after sucrose gradient fractionation. Equal volume (20 μL) of each fraction both for Tissue 16k (A) and 118k (B), 30 μg of Tissue Homogenate (TH) were loaded as positive controls. WB for the antibodies described in the figures. Un-cropped WB are available in the Supplementary Information (Fig 6-8SI). Each fraction density

is indicated in g/cm^3. The green box indicates fractions that, for further analyses, have been collected and further processed as a single sample.

Figure 4. Purity assessment and protein concentration measurements.

Fractions 6 to 9 after the sucrose density gradient were collected and analyzed. Samples derived from Plasma (PL16kG, PL118kG) or Tissue (TS16kG, TS118kG). Samples were checked for purity from exogenous protein contaminants (EPC) with the CONAN assay. Aggregation Index ratios (AI%) of serial dilution (1:10, 1:100) of a representative sample are showed in the graph. The dotted line represents the CONAN assay threshold for EPC detection (AI% <20% EPC content ≤0.05 μg/μl). The AI% is inversely proportional to the preparation purity and samples with AI% below the 20% are considered pure. AI% values of a representative sample with assay internal controls (NormREF, black dots; IntREF, green dots) (A); Protein concentration with BCA (B) and Bradford assay (C); Protein concentration for each group was normalized by tissue mass (per 1 gr). Values were shown as mean ± SEM. Significant differences among datasets were determined with Student's t-test. P values: ****p < 0.0001. TS16kG, dark blue points; PL16kG, dark red squares; TS118kG, light triangle up; PL118kG, light red triangle down.

Figure 5. Nano-mechanical features of samples from Plasma and Tissue**.** Representative AFM images of PL16kG (A), PL118kG (B), TS16kG (C), and TS118kG (D). Contact angle VS diameter plot of several hundred individual particles from at least five different depositions for each sample as calculated via AFM morphometry of PL16kG (E), PL118kG (F), TS16kG (G) and TS118kG (H). Blue box represents area populated by structures with the typical EV-like characteristics: contact angle ranging from 75° to 135°, diameter from 30 to 300 nm.

Figure 6. Concentration and size distribution. Particle concentration for each sample were measured by NTA. Particle concentration was normalized for starting material mass (1gr) (A). Particle Size distribution was measured by NTA. Histograms represent the mean of the size distribution divided by range (0-200 nm; 200-400 nm; 400-600 nm) of the 16k and 118k samples from plasma and tissueB). Particle size was analyzed by NTA and modal distribution (nm) (C).

Values in nm were shown as mean ± SEM. Significant differences among datasets were determined with Student's t-test. P values: **p< 0.01, ***p < 0.001 and ****p < 0.0001.

Figure 7. Size distribution with DLS. Representative autocorrelation function of the scattered intensity, with the fitting curve according to the Laplace Inversion via CONTIN algorithm (A,D,G,L); corresponding scattered intensity-weighted (B,E,H,M) and number-weighted (C,F,I,N) size distributions of scattering objects of samples described in the text.

Figure 8. Zeta-potential measurement. Preparations were analyzed for their colloidal stability measuring z-potential with DLS. Values in mV were shown as mean ± standard error of the mean (SEM). Significant differences among datasets were determined with Student's t-test. P values: **p< 0.01.

A) Tissue 16k

	1	2	3	4	5	6	7	8	9	10	11	12	TH (30μg)
20kDa													Caveolin 3
50kDa													Beta enolase
75kDa													ADAM10
50kDa													Flotillin-1
50kDa													TSG101
25kDa													CD81
75kDa													Calnexin
g/cm³	1.05	1.06	1.07	1.09	1.11	1.13	1.18	1.24	1.29	1.30	1.34	1.34	

B) Tissue 118k

	1	2	3	4	5	6	7	8	9	10	11	12	TH (30μg)
20kDa													Caveolin 3
50kDa													Beta enolase
75kDa													ADAM10
50kDa													Flotillin-1
50kDa													TSG101
25kDa													CD81
75kDa													Calnexin
g/cm³	1.05	1.06	1.07	1.09	1.11	1.13	1.18	1.24	1.29	1.30	1.34	1.34	

A)

B)

C)

BCA

D)

Bradford