Biallelic pathogenic variants in *DNAH3* cause male infertility in humans and mice

Xiang Wang[1, 2, 9], Gan Shen[1, 9], Yihong Yang[3, 9], Chuan Jiang [1, 9], Tiechao Ruan[4, 9], Xue Yang[1], Liangchai Zhuo[1], Yingteng Zhang[1], Yangdi Ou[5], Xinya Zhao[6], Shunhua Long[7, 8], Xiangrong Tang[7, 8], Tingting Lin[7, 8, *], Ying Shen[1, 2, *]

Author Affiliation

[1] Department of Obstetrics/Gynecology, Key Laboratory of Obstetric, Gynecologic and Pediatric Diseases and Birth Defects of Ministry of Education, West China Second University Hospital, Sichuan University, Chengdu 610041, China.

[2] NHC Key Laboratory of Chronobiology, Sichuan University, Chengdu 610041, China.

[3] Reproduction Medical Center of West China Second University Hospital, Key Laboratory of Obstetric, Gynecologic and Pediatric Diseases and Birth Defects of Ministry of Education, Sichuan University, Chengdu 610041, China.

[4] Department of Pediatrics, West China Second University Hospital, Sichuan University, Chengdu 610041, China.

[5] West China School of Medicine, Sichuan University, Chengdu 610041, China.

[6] West China School of Basic Medicine and Forensic Medicine, Sichuan University, Chengdu 610041, China.

[7] Chongqing Key Laboratory of Human Embryo Engineering, Center for Reproductive Medicine, Women and Children's Hospital of Chongqing Medical University, Chongqing 400010, China.

[8] Chongqing Clinical Research Center for Reproductive Medicine, Chongqing Health Center for Women and Children, Chongqing 400010, China.

[9] These authors contributed equally to this work.

[*] Correspondence to Tingting Lin (yuting9263@163.com) and Ying Shen (shenying01@scu.edu.cn)

Abstract

Axonemal protein complexes, including the outer and inner dynein arms (ODA/IDA), are highly ordered structures of the sperm flagella that drive sperm motility. Deficiencies in several axonemal proteins have been associated with male infertility, which is characterized by asthenozoospermia or asthenoteratozoospermia. Dynein axonemal heavy chain 3 (DNAH3) resides in the IDA and is highly expressed in the testis. However, the relationship between DNAH3 and male infertility is still unclear. Herein, we identified biallelic variants of *DNAH3* in four unrelated Han Chinese infertile men with asthenoteratozoospermia through whole-exome sequencing (WES). These variants contributed to deficient DNAH3 expression in the patients' sperm flagella. Importantly, the patients represented the anomalous sperm flagellar morphology, and the flagellar ultrastructure was severely disrupted. Intriguingly, *Dnah3* knockout (KO) male mice were also infertile, especially showing the severe reduction in sperm movement with the abnormal IDA and mitochondrion structure. Mechanically, nonfunctional DNAH3 expression resulted in decreased expression of IDA-associated proteins in the spermatozoa flagella of patients and KO mice, including DNAH1, DNAH6, and DNALI1, the deletion of which has been involved in disruption of sperm motility. Moreover, the infertility of patients with *DNAH3* variants and *Dnah3* KO mice could be rescued by intracytoplasmic sperm injection (ICSI) treatment. Our findings indicated that *DNAH3* is a novel pathogenic gene for asthenoteratozoospermia and may further contribute to the diagnosis, genetic counseling, and prognosis of male infertility.

Key words: DNAH3, Male infertility, IDA-associated proteins, asthenoteratozoospermia, WES.

Introduction

Infertility is a global public health and social problem that affects approximately one in six couples worldwide. (1) Male infertility, which accounts for half of infertile cases, is a multifactorial disease with common phenotypes, including oligo/azoospermia (poor sperm count or absence of spermatozoa); teratozoospermia (aberrant sperm morphology); asthenozoospermia (weakened sperm motility); and a combination of these phenotypes, such as asthenoteratozoospermia, oligoasthenozoospermia, oligoteratozoospermia and oligoasthenoteratozoospermia. (2, 3)

Asthenoteratozoospermia is one of the most common phenotypes of male infertility, and genetic factors have been established as the predominant cause of asthenoteratozoospermia. Multiple morphological abnormalities of the flagella (MMAF), a subtype of asthenoteratozoospermia, characterized by a mosaic of abnormalities of the flagellar morphology, including absent, short, coiled, bent and/or irregular flagella, is almost always caused by genetic defects. (4, 5) To date, more than 40 genes have been identified as pathogenic genes of MMAF, but these genes can only explain approximately 60% of MMAF-affected cases. (6-9) Therefore, the genetic basis of the remaining cases is still unknown.

The motility of a sperm is driven by its rhythmically beating flagella, and at the center of the flagella lies a conserved axonemal structure, containing the "9 + 2" microtubular arrangement: a ring of nine microtubule doublets (MTDs) surrounding a central pair (CP) of singlet microtubules. Each MTD consists of an A tubule and a B tubule, and the outer (ODA)

and inner (IDA) dynein arms are anchored along the A tubule. (10) The ODA and IDA are

ATPase-based protein complexes that drive the movement between the A tubule and the

neighboring B tubule of the next doublet, producing the original force for sperm motility.

(11, 12) Structural and functional abnormalities of the ODA and IDA have been

demonstrated to cause male infertility associated with asthenozoospermia and/or

asthenoteratozoospermia. (4, 13, 14)

The dynein axonemal heavy chain (DNAH) family comprises a series of proteins

(DNAH1–3, DNAH5–12, and DNAH17) that are precisely assembled with other axonemal

dynein motor proteins in the ODAs and IDAs of sperm flagella and motile cilia. (15-17) In

humans, DNAH1, DNAH2, DNAH6, DNAH7, DNAH8, DNAH10, DNAH12 and DNAH17

are highly expressed in the testis, and deficiency of these proteins has been demonstrated to

cause MMAF-associated asthenoteratozoospermia. (18-25) DNAH3 is an evolutionarily

conserved IDA-associated protein and is highly expressed in testes of humans and mice.

Deficient DNAH3 has been shown to impair sperm motility in *Drosophila* and cattle. (26, 27)

In humans, *DNAH3* has been identified as a novel breast cancer candidate gene. (28)

However, the role of DNAH3 in male reproduction in humans and mice remains largely

unknown.

In the present study, we identified four biallelic variations in *DNAH3* in four unrelated

Han Chinese patients with asthenoteratozoospermia using whole-exome sequencing (WES).

The spermatozoa of the patients showed extremely reduced sperm motility and a high

proportion of sperm tail defects characterized by the MMAF phenotype. We further

generated *Dnah3* knockout (KO) mice, and the male KO mice expectedly showed

aberrations in sperm movement, flagellar IDA, and mitochondrion. Moreover, the absence of

DNAH3 led to decreased expression of other IDA-associated proteins, including DNAH1,

DNAH6 and DNALI1. Importantly, good outcomes of intracytoplasmic sperm injection

(ICSI) treatment were observed in *DNAH3*-deficient patients and *Dnah3* KO mice. This

study revealed *DNAH3* as a novel pathogenic gene of asthenoteratozoospermia, and the

findings provide valuable suggestions for the clinical diagnosis and treatment of male

infertility.

Results

Identification of biallelic pathogenic variants of *DNAH3* in four unrelated infertile men

In the present study, we conducted whole-exome sequencing (WES) to identify potential

candidate genes associated with primary asthenoteratozoospermia. Interestingly, after

filtering, four biallelic variations in *DNAH3* (NG_052617.1, NM_017539.2, NP_060009.1)

were identified in four unrelated patients: a compound heterozygous mutation of c.3590C>T

(p.P1197L) and c.3590C>G (p.P1197R) in Patient 1, a homozygous missense mutation of

c.4837G>T (p.A1613S) in Patient 2, a compound heterozygous mutation of c.5587del

(p.L1863*) and c.10355C>T (p.S3452L) in Patient 3 and a compound heterozygous

mutation of c.2314C>T (p.R772W) and c.4045G>A (p.D1349N) in Patient 4 (**Figure 1A**).

Importantly, routine semen analysis revealed that all patients showed extremely reduced

sperm motility and a high proportion of sperm tail defects (**Table 1**). These variants either

were not recorded or had an extremely low frequency in multiple public population

databases and were predicted to be potentially deleterious by SIFT, PolyPhen-2 and

Mutation Tester (**Table 2**). Next, Sanger sequencing confirmed these variants in the probands, and their fertile parents carried the heterozygous variants (**Figure 1A**). Moreover, the variant sites are localized in several domains of the DNAH3 protein and are highly conserved across species (**Figure 1B**).

Strikingly, immunofluorescence staining revealed that DNAH3 was exclusively resided in the tail and concentrated in the midpiece of control sperm. However, the fluorescence signal of DNAH3 was hardly detected in the patients' spermatozoa (**Figure 1C**). Additionally, subsequent western blotting analysis yielded consistent results with immunofluorescence staining, indicating that these variants led to disrupted expression of DNAH3 (**Figure 1D**). These results suggested that biallelic variants in *DNAH3* disrupted DNAH3 expression and might be responsible for the infertility of the four patients.

Asthenoteratozoospermia phenotype is observed in patients with *DNAH3* variants

We next investigated the aberrant sperm morphology of the patients using Papanicolaou staining and SEM analysis. Notably, the tails of sperm from the patients exhibited a typical phenotype associated with MMAF, including coiled, short, bent, irregular, and/or absent flagella (**Figure 2A and B**). In addition, a fraction of defects in the sperm head were also present in the patients' sperm (**Figure 2A and B**).

TEM was employed to determine the ultrastructure of the sperm from the patients. Compared to the integrated and well-organized "9 + 2" axonemal arrangement of the sperm flagella from the normal control, spermatozoa from the patients showed absent or disordered CPs, MTDs, and outer dense fibers (ODFs) in different regions of the flagella (**Figure 3A**).

Interestingly, the IDAs of sperm flagella of the patients were hardly captured compared to

the control (**Figure 3A**). Additionally, in the midpiece of sperm flagella of the patients,

dissolved mitochondrial material was also observed evidently under TEM (**Figure 3A**). We

next conducted immunofluorescence staining to label the mitochondria of patients' sperm

with TOM20, a subunit of the mitochondrial import receptor. Remarkably, in contrast to the

robust TOM20 signals observed in the normal control, the TOM20 signals in the sperm from

the patients were considerably diminished, indicating a disrupted mitochondrial function

(**Figure 3B**). Together, these data suggested that *DNAH3* may function in sperm flagellar

development, and loss-of-function variants were associated with MMAF in humans.

DNAH3 is exclusively expressed in the sperm flagella of humans and mice

To further understand the function of DNAH3 in male reproduction, we explored the

expression pattern of DNAH3 in humans and mice. qPCR results revealed that *Dnah3* was

predominantly expressed in the mouse testis (**Figure S1A**). Moreover, when observing the

expression of *Dnah3* in testes from mice at different postnatal days, we found that *Dnah3*

expression was significantly elevated beginning on postnatal Day 22, peaked at postnatal

Day 30, and maintained a stable expression level thereafter (**Figure S1B**). In addition, germ

cells at different stages were isolated from the testes of humans and mice and were stained

with anti-DNAH3 antibody. The results showed that DNAH3 was expressed in the

cytoplasm of spermatocytes and spermatogonia and then obviously in the flagellum of early

and late spermatids (**Figure S2A and B**). These expression data suggest that DNAH3 may

play an important role in sperm flagellar development during spermatogenesis in humans

and mice.

Deletion of *Dnah3* causes male infertility in mice

Considering the absent expression of DNAH3 in the patient sperm, we generated *Dnah3* KO

mice using CRISPR–Cas9 technology to further confirm the essential role of DNAH3 in

spermatogenesis (**Figure S3A**). PCR, qPCR, and immunofluorescence staining were used to

confirm that *Dnah3* was null in KO mice (**Figure S3B-E**). The *Dnah3* KO mice survived

without any evident abnormalities in development and behavior. H&E staining further

revealed that there were no histological differences in the lung, brain, eye, or oviduct

between wild-type (WT) and *Dnah3* KO mice (**Figure S4A**). In addition, no obvious

abnormalities in ciliary development were observed in these organs in KO mice compared to

WT mice (**Figure S4B**). The *Dnah3* KO female mice were fertile with normal oocyte

development (**Figure S5A**). However, the *Dnah3* KO male mice were completely infertile

(**Figure 4A**). We next examined the testis and epididymis of *Dnah3* KO male mice to

elucidate the etiology of infertility. There was no detectable difference in the testis/body

weight ratio of *Dnah3* KO mice when compared to WT mice (**Figure S5B**).

Moreover, subsequent computer-assisted sperm analysis (CASA) also showed that sperm

isolated from the cauda epididymis were slightly decreased, and nearly all sperm were

completely immobile (**Table 3, Movie S1 and Movie S2**). Papanicolaou staining and SEM

analysis revealed morphological defects in partial spermatozoa from *Dnah3* KO mice,

including coiled, bent, and irregular flagella, as well as aberrant heads and acephalic

spermatozoa (**Figure S6A and B**).

TEM was further utilized to evaluate the sperm flagellar ultrastructure of *Dnah3* KO mice.

There were no obvious abnormalities of "9 + 2" microtube arrangement in most sperm from

the *Dnah3* KO mice when compared to WT mice (**Figure 4B**). However, in contrast to the

clear display of an IDA and an ODA on the A-tube of each microtubule doublet in the sperm

flagella of WT mice, the sperm flagella of *Dnah3* KO mice exhibited an absence of almost

all the IDAs (**Figure 4B**). In addition, the disrupted mitochondria of spermatozoa from

Dnah3 KO mice were also observed under TEM, as manifested by the dilated

intermembrane spaces and dissolved mitochondrial material (**Figure 4B and C**). We next

performed immunofluorescence staining to label SLC25A4, which is responsible for the

exchange of ATP and ADP across the mitochondrial inner membrane. Strikingly, compared

to the bright fluorescence signals in the midpiece of WT sperm, the signals *Dnah3* KO were

significantly diminished (**Figure 4D**), indicating impaired mitochondrial function.

Collectively, DNAH3 is essential for spermatogenesis, and its deficiency seriously damages

the sperm motility and IDAs in both humans and mice.

DNAH3 deficiency impairs IDAs related to the reduction of IDA-associated proteins

Considering the disrupted IDAs revealed by TEM analysis in both our patients and *Dnah3*

KO mice, we speculated whether the defective IDAs were attributed to the decreased

expression of the key IDA-associated proteins. The immunofluorescence data showed that

DNAH1/DNAH6 and DNALI1, corresponding to the heavy and light intermediate chains of

the IDAs, (16) respectively, were almost invisible along the sperm flagella of the patients

when compared to control (**Figure 5A-C**). Consistent results were obtained in our

subsequent western blotting analysis of sperm lysates from the patients (**Figure 5D-F**),

indicating that DNAH3 may manipulate the assembly of IDA through regulating the

expression of IDA-associated proteins. In contrast, DNAH8/DNAH17 and DNAI1,

corresponding to the heavy and intermediate chains of ODAs, (25) were readily detectable

in the patients' sperm flagella and were comparable to the control (**Figure S7A-C**),

suggesting that DNAH3 may not regulate the expression of ODA-associated proteins. We

also performed immunofluorescence staining and western blotting analysis of DNAH1,

DNAH6, DNALI1, DNAH8, DNAH17 and DNAI1 on sperm from *Dnah3* KO mice, and the

results observed were consistent with those of the patients (**Figure 6A-F, Figure S8A-C**).

These findings suggested that other IDA-associated proteins might be downstream effectors

of DNAH3, which needs more future research.

ICSI treatment of humans with *DNAH3* variants and *Dnah3* KO mice

ICSI treatment has been reported to be effective in asthenoteratozoospermia-associated

infertility. (29, 30) ICSI cycles were attempted for the patients after written informed

consent was obtained. The female partners all had normal basal hormone levels and

underwent a long gonadotrophin-releasing hormone agonist protocol (**Table 4**). The wife of

Patient 1 underwent one ICSI attempt. A total of 21 metaphase II (MII) oocytes were

retrieved and microinjected, of which 17 oocytes were successfully fertilized (17/21,

80.95%) and cleaved (17/17, 100%). Thirteen Day 3 (D3) embryos were formed, six of

which developed into blastocysts (8/13, 61.54%) after standard embryo culture. Two

blastocysts were transferred, one of which was implanted. She eventually achieved clinical

pregnancy, and the pregnancy is ongoing (**Table 4**). The partner of Patient 2 underwent two ICSI attempts. In her first ICSI attempt, six MII oocytes were retrieved, of which three were fertilized (3/6, 50%) and cleaved (3/3, 100%). After standard embryo culture, two D3 embryos were formed and transferred. However, this ICSI failed because no embryos were implanted. In her second ICSI attempt, all five MII oocytes were fertilized and cleaved (5/5, 100%). Five D3 embryos were obtained, of which two were transferred, but no embryos were implanted. The remaining three D3 embryos were cultured continuously, and two available blastocysts were formed and kept to be transferred in the future (**Table 4**). The partner of Patient 3 underwent one ICSI attempt. Of the 20 MII oocytes retrieved, 19 oocytes were fertilized (19/20, 95.0%) and cleaved (19/19, 100%). Fifteen D3 embryos were obtained, and 10 developed into available blastocysts (10/15, 66.7%). One blastocyst was transferred and implanted. She achieved clinical pregnancy, and the pregnancy is ongoing (**Table 4**). The wife of Patient 4 underwent four failed ICSI attempts. In her first two ICSI attempts, 13 and 12 MII oocytes were retrieved, of which five (5/12, 41.67%) and six (6/13, 46.15%), respectively, were fertilized and cleaved. Two available D3 embryos were obtained and transferred in both ICSI attempts, but no embryos were implanted. In her third ICSI attempt, of the eight MII oocytes retrieved, four (4/8, 50%) were fertilized and cleaved (4/4, 100%). However, no available D3 embryos were acquired. In her last ICSI attempt, seven MII oocytes were retrieved, of which three were fertilized (3/7, 42.68%) and two were cleaved (2/3, 66.7%), but no available D3 embryos were formed (**Table 4**). The vivid embryonic development of Patient 1 and Patient 3 after ICSI treatment was shown in **Figure 7A**.

We also carried out ICSI treatment on *Dnah3* KO male mice. Strikingly, favorable

outcomes of ICSI were obtained in *Dnah3* KO male mice. After injection of spermatozoa

from *Dnah3* KO male mice, pronuclei were observed in most embryos in both the KO and

WT groups, indicating a normal fertilization rate (**Figure 7B**). There was no difference in

the percentage of 2-cell and blastocyst-stage embryos between the KO and WT groups

(**Figure 7B**). Collectively, we observed successful ICSI outcomes in two out of four

DNAH3-deficient patients and *Dnah3* KO male mice and therefore suggested ICSI as an

optional treatment for infertile men harboring biallelic pathogenic variants in *DNAH3*, and

the additional female risk factors for infertility should not be excluded in the failed patients.

Discussion

In the present study, we identified pathogenic variants in *DNAH3* in unrelated infertile men

with asthenoteratozoospermia. These variations resulted in the almost absence of DNAH3

and sharply decreased the expression of other IDA-associated proteins, including DNAH1,

DNAH6 and DNALI1. Combined with similar findings in *Dnah3* KO mice, we

demonstrated that DNAH3 is fundamental for male fertility. Moreover, we suggest that ICSI

might be a favorable treatment for male infertility caused by DNAH3 deficiency. Our

findings identify a function for DNAH3 in male reproduction in humans and mice and may

provide a new view on the clinical practice of male infertility.

Primary ciliary dyskinesia (PCD, MIM: 244400) is a genetic disorder affecting at least

one in 7554 individuals. (31) The most common symptoms of PCD are recurrent infections

in airways due to malfunction of the motile cilia that are responsible for mucus clearance.

(32) It has been suggested that male infertility associated with sperm defects is highly

prevalent (up to 75%) among individuals with PCD. (33) Axonemal defects caused by

variants within *DNAH* family members, including *DNAH5*, *DNAH6*, *DNAH7*, *DNAH9* and

DNAH11, are causative factors for PCD. (34-36) Moreover, deficiency in these

PCD-causing *DNAH*s has also been associated with male infertility. (9, 14, 20, 21, 37-39)

Additionally, other *DNAH*s, such as *DNAH1*, *DNAH2*, *DNAH8*, *DNAH10*, *DNAH12* and

DNAH17, are suggested to be pathogenic genes of isolated male infertility. (18, 19, 22, 24,

25) These phenotype–genotype correlations may be attributed to the fact that ciliary and

flagellar axonemes have cell type-specific or cell type-enriched DNAHs. (40) DNAH3

resides in the IDA and is expressed in testis and ciliary tissues, including the lung, brain, eye,

and oviduct. However, despite its presence in these tissues, the relationship between

deficient DNAH3 and disease is unclear to date. Intriguingly, in our study, we observed male

infertility associated with abolished sperm movement and IDA structure in both patients

carrying biallelic *DNAH3* mutations and *Dnah3* KO mice. However, none of the patients

reported suffering from any of the principal PCD symptoms, and the *Dnah3* KO mice

showed normal ciliary development in the lung, brain, eye, and oviduct. These findings

suggested that DNAH3 may play a more important role in sperm flagellar development than

in other motile cilia functions. Given that DNAH3 is expressed in ciliary tissues, its role in

these tissues remains intriguing and could be elucidated through sequencing of larger

cohorts of individuals with PCD.

In our study, both *Dnah3* KO male mice and *DNAH3*-mutant men are infertile and

showed the immotile sperm with disrupted flagellar IDAs and mitochondria. However, there

are still some differences between their phenotypes. The *Dnah3* KO mice displayed

moderate sperm morphological defects, and the "9 + 2" microtube arrangement showed no

obvious abnormalities. While our patient exhibited a high proportion of sperm tail defects,

including coiled, short, bent, and irregular flagella, along with seriously defective "9 + 2"

microtube arrangement. Importantly, previous studies have suggested that the phenotypes

resulting from depletions of *Dnah*s in mice do not always resemble deficiencies in the

human orthologs. Specifically, *Dnah1*$^{-/-}$ mice was infertile due to decreased sperm motility,

without any observable structural defects of the axoneme. (41) In contrast, spermatozoa

from *DNAH1*-deficient men exhibited significant axonemal disorganization and MMAF

phenotype. (18) Combined with our findings, it is conceivable that DNAH3 is crucial for

human and mouse male reproduction, but may play distinct roles during flagellar axonemal

assembly.

ICSI has been an efficient treatment for male infertility. (42, 43) However, the

outcomes of ICSI for male infertility caused by variants in different *DNAH* genes are

variable. It has been demonstrated that infertile males with variants in *DNAH1*, *DNAH2*,

DNAH7, and *DNAH8* have a favorable prognosis, (22, 44-48) while patients with variants in

DNAH17 have poor outcomes after ICSI treatment. (25, 49) Meanwhile, the ICSI outcomes

in male infertility caused by *DNAH6* variants may depend on the specific mutation or be

controversial. (20, 50, 51) The patients with *DNAH3* mutations in our study experienced

different clinical outcomes of ICSI treatment. The partners of Patient 1 and Patient 3

achieved clinical pregnancy. The wives of Patient 2 and Patient 4 obtained favorable

fertilization and cleavage rates but experienced no clinical pregnancy due to the

nonimplantation of the transferred embryos. Remarkably, despite the diverse variants within

DNAH3 observed in the four patients, all variants led to a complete absence of DNAH3

expression. Therefore, these different ICSI outcomes might be attributed to additional

unexplained factors from the female partners. Considering the successful ICSI outcomes

observed in *Dnah3* KO mice, we suggested ICSI as an optimized treatment for infertile men

carrying variants in *DNAH3*. More cases are needed to precisely estimate the prevalence of

DNAH3 mutations and determine a prognosis for ICSI treatments.

In conclusion, our study revealed an unexplored role of DNAH3 in male reproduction

in humans and mice, suggesting *DNAH3* as a novel causative gene for human

asthenoteratozoospermia. Moreover, ICSI is as an optimized treatment for infertile men with

DNAH3 variants. This study expands our knowledge of the relationship between DNAH

proteins and disease, facilitating genetic counseling and clinical treatment of male infertility

in the future.

Methods

Human subjects

Four unrelated Han Chinese infertile men and their family members were recruited from

West China Second University Hospital of Sichuan University and Women and Children's

Hospital of Chongqing Medical University. All patients exhibited a normal karyotype (46

XY) without deletion of the azoospermia factor (AZF) region in the Y-chromosome. All of

the participants were provided informed consent, and the study was approved by the ethics

committee of West China Second University Hospital and The First Affiliated Hospital of

Chongqing Medical University.

Genetic analysis

Peripheral blood samples were obtained from the subjects to extract genomic DNA using a

DNA purification kit (TIANGEN, DP304). For WES, 1 μg of genomic DNA was utilized for

exon capture using the Agilent SureSelect Human All Exon V6 Kit and sequenced on the

Illumina HiSeq X system (150-bp read length). Mutations were annotated and filtered using

the ExAC browser, dbSNP, 1000 Genomes Project and HGMD. Then, PolyPhen-2, SIFT,

MutationTaster, and CADD were utilized for functional prediction. The candidate

pathogenic variants were verified by Sanger sequencing on DNA from the patients' families.

The primer pairs used for PCR amplification are listed in **Table S1**.

Electron microscopy

For scanning electron microscopy (SEM), sperm samples were fixed in glutaraldehyde

(2.5%, w/v) and dehydrated using an ethanol gradient (30, 50, 75, 85, 95, and 100% ethanol).

The samples were dried using a CO_2 critical-point dryer (Eiko HCP-2, Hitachi) and

observed under SEM (S-3400, Hitachi).

For transmission electron microscopy (TEM), sperm samples were fixed in

glutaraldehyde (3%, w/v) and osmium tetroxide (1%, w/v) and dehydrated with an ethanol

gradient. The samples were embedded in Epon 812. Ultrathin sections were stained with

uranyl acetate and lead citrate and analyzed under TEM (Tecnai G2 F20).

STA-PUT velocity sedimentation

Single testicular cells from obstructive azoospermia and 8-week-old C57BL male mice were obtained using the STA-PUT velocity sedimentation method as described previously. In brief, total spermatogenic cells were harvested by digesting seminiferous tubules with collagenase (Invitrogen, 17100017), trypsin (Sigma, T4799) and DNase (Promega, M6101) for 15 min each at 37 °C. Cells were diluted in bovine serum albumin (BSA, 3%, w/v) and filtered through an 80 mm mesh to remove fragments. Then, the cells were resuspended in BSA (3%, w/v) and loaded in an STA-PUT velocity sedimentation cell separator (ProScience) to obtain germ cells at different stages.

RNA isolation and quantitative PCR (qPCR)

Total RNA of mouse tissues was extracted using TRIzol reagent (Invitrogen,15596026,) and reverse-transcribed using the 1st Strand cDNA Synthesis Kit (Yeasen, HB210629) according to the manufacturer's instructions. qPCR was carried out on an iCycler RT–PCR Detection System (Bio-Rad Laboratories) using SYBR Green qPCR Master Mix (Bimake, B21202). Primer sequences are listed in **Table S1**.

Immunofluorescence staining

Sperm samples were fixed in paraformaldehyde (4%, w/v), permeabilized with Triton X-100 (0.3% v/v) and blocked with BSA (3%, w/v) at room temperature. Samples were incubated with primary antibodies, including DNAH1 (Cusabio, CSB-PA878961LA01HU, 1:100), DNAH3 (Cusabio, CSB-PA823461LA01HU, 1:100), DNAH6 (Proteintech, 18080-1-AP,

RRID: AB_2878493, 1:50), DNAH8 (Atlas, HPA028447, RRID: AB_10599600, 1:200),

DNAH17 (Proteintech, 24488-1-AP, RRID: AB_2879568, 1:50), DNAI1 (Proteintech,

12756-1-AP, RRID: AB_10643244, 1:50), DNALI1 (Proteintech, 17601-1-AP, RRID:

AB_2095372, 1:50), TOM20 (Proteintech, 11802-1-AP, RRID: AB_2207530, 1:50),

SLC25A4 (Signalway, 32484, RRID: AB_2941094, 1:100), lectin PNA (Invitrogen,

L-32460, 1:50) and alpha tubulin (Abcam, ab7291, RRID: AB_2241126, 1:500), overnight

at 4 °C. The next day, the samples were washed and incubated with the secondary antibody

Alexa Fluor 488 (Invitrogen, A11008, RRID: AB_143165, 1:1000) or Alexa Fluor 594

(Invitrogen, A11005, RRID: AB_141372, 1:1000), and the nuclei were labeled with

4′,6-diamidino-2-phenylindole (DAPI, Sigma–Aldrich, D9542). Image capture was

performed by a laser scanning confocal microscope (Olympus, FV3000).

For staining of mouse tissues, samples were first fixed in paraformaldehyde (4%, w/v)

and dehydrated with an ethanol gradient. Then, the samples were embedded in paraffin and

sliced into 5-μm sections. After deparaffinization and rehydration, sections were processed

with 3% hydrogen peroxide and incubated in sodium citrate for antigen repair. Subsequently,

sections were blocked with goat serum and incubated with primary antibodies against

DNAH3 (Cusabio, CSB-PA823461LA01HU, 1:100) or ac-Tubulin (Abcam, ab24610, RRID:

AB_448182, 1:500) at 4 °C overnight. The next day, the sections were incubated with the

secondary antibody Alexa Fluor 488, followed by labeling the nuclei with DAPI. Image

capture was performed using a fluorescence microscope (Zeiss, Ax10).

Western Blotting

Sperm samples were lysed in RIPA buffer (Beyotime, P0013B) to extract the total protein. For analysis of DNALI1, the protein samples were mixed with SDS loading buffer (P0015, Beyotime, China), boiled at 95 °C for 5 minutes, and separated by 12.5% SDS-PAGE. For analysis of DNAH1, DNAH3, DNAH6, the protein samples were mixed with NuPAGE™ LDS sample buffer (Invitrogen, NP0007), denatured at 70°C for 10 minutes, and separated by 3–8% NuPAGE™ Tris-Acetate gels (EA0375BOX, Invitrogen). Then the resolved proteins were transferred to 0.45 µm PVDF membranes (Merck Millipore, IPVH00010). The membranes were blocked, incubated with primary antibodies, including DNALI1 (Proteintech, 17601-1-AP, RRID: AB_2095372, 1:150), DNAH3 (Cusabio, CSB-PA823461LA01HU, 1:200), DNAH6 (Proteintech, 18080-1-AP, RRID: AB_2878493, 1:150) and alpha tubulin (Proteintech, 11224-1-AP, RRID: AB_2210206, 1:1000) at 4 °C overnight. The following day, membranes were washed and incubated with HRP-conjugated secondary antibody (Proteintech, SA00001-2, RRID: AB_2722564, 1:5000). Protein bands were visualized using enhanced chemiluminescence reagents (Millipore, WBKLS0500).

Histology hematoxylin-eosin (H&E) staining

Tissue samples from mice were fixed with 4% paraformaldehyde (w/v) overnight. Following dehydration by ethanol, the samples were embedded in paraffin and sliced into 5-µm sections. The sections were stained with hematoxylin and eosin and observed under a microscope (Zeiss, Axio Imager 2).

Generation of the *Dnah3* KO mouse model

Animal experiments in this study were approved by the Experimental Animal Management

and Ethics Committee of West China Second University Hospital, Sichuan University, and

complied with the Animal Care and Use Committee of Sichuan University. A *Dnah3*

knockout mouse model was generated by the CRISPR–Cas9 system. Briefly, Cas9 and

signal-guide RNAs (5'-GTATCAAGTGGATGTAAACC-3') were transcribed using T7

RNA polymerase in vitro and comicroinjected into the cytoplasm of single-cell C57BL/6J

mouse embryos to generate frameshift mutations by nonhomologous recombination through

introduction of a 1 bp insertion in exon 13. Then, the embryos were cultured and transferred

into the oviducts of pseudopregnant female mice at 0.5 days post-coitum. A mutation of

Dnah3 in the founder mouse and their offspring was confirmed using PCR and Sanger

sequencing. The primers used for the generation of animal models are listed in **Table S1.**

Intracytoplasmic sperm injection (ICSI)

ICSI was carried out using standard techniques. In brief, one-month-old female KM mice

were injected with 5 IU of equine chorionic gonadotropin (eCG) (ProSpec, HOR-272) to

induce superovulation. Metaphase II-arrested (MII) oocytes were acquired through another

injection of 5 IU human chorionic gonadotropin after 48 hours. MII oocytes were incubated

with Chatot-Ziomek-Bavister medium (Easycheck, M2750) at 37.5 °C and 5% CO_2 until use.

Mouse cauda epididymal spermatozoa were incubated in human tubal fluid (HTF) medium

(Easycheck, M1150) and then frozen and thawed repeatedly to remove sperm tails. For ICSI,

a single sperm head was microinjected into an MII oocyte by using a NIKON inverted

microscope and a Piezo (PrimeTech, Osaka, Japan) in Whitten's-HEPES medium containing

0.01% polyvinyl alcohol (Gibco,12360-038) and cytochalasin B (3.5 g/ml; Sigma–Aldrich,

C-6762). The successfully injected oocytes were transferred into G1-Plus medium (Vitrolife,

10132) and incubated at 37.5 °C and 5% CO_2. The animal experiments were approved by

the Experimental Animal Management and Ethics Committee of West China Second

University Hospital, Sichuan University.

Statistical analysis

Prism (version 8.4.0, GraphPad, Boston, MA, USA) and SPSS (version 18.0, IBM

Corporation, Armonk, NY, USA) were used to perform statistical analyses. All data are

presented as the means ± SEMs. Data from two groups were compared using an unpaired,

parametric, two-sided Student's t test, and a p value less than 0.05 was considered

statistically significant.

Ethics approval

This study was approved by Ethical Review Board of West China Second University

Hospital, Sichuan University. Informed consent was obtained from each participate in this

study before taking part.

Data availability

The published article includes all datasets generated or analyzed during this study. The

exome-sequencing data supporting the current study have not been deposited in a public

repository because of privacy issues but are available from the corresponding author on

request.

Competing interests

470 The authors declare that they have no conflict of interest.

Author contributions

Xiang Wang: Data curation; formal analysis; investigation; methodology; writing – original

draft. **Gan Shen**: Data curation; formal analysis; **Yihong Yang**: Resources; investigation;

methodology. **Chuan Jiang**: Data curation; formal analysis. **Tiechao Ruan**: Formal analysis;

methodology. **Xue Yang**: Validation; investigation. **Liangchai Zhuo**: Investigation;

Yingteng Zhang: Investigation, methodology. **Yangdi Ou**: Investigation. **Xinya zhao**:

Investigation. **Shunhua Long**: Methodology. **Xiangrong Tang**: Investigation. **Tingting Lin**:

Conceptualization; funding acquisition; project administration. **Ying Shen**:

Conceptualization; supervision; project administration; writing – review and editing.

Acknowledgements

481 The authors thank the patient and his family members for their voluntary participation. We

482 are grateful to Guiping Yuan from Analytical and Testing Center of Sichuan University and

483 Yan Liang from Research Core Facility of West China Hospital, Sichuan University for their

484 help with TEM images and preparing histology slides. This work was supported by

485 National Natural Science Foundation of China (82301807).

Reference

1.   Cox CM, Thoma ME, Tchangalova N, Mburu G, Bornstein MJ, Johnson CL, Kiarie J. Infertility
prevalence and the methods of estimation from 1990 to 2021: a systematic review and meta-analysis. Hum
Reprod Open. 2022;2022(4):hoac051.
2.   Eisenberg ML, Esteves SC, Lamb DJ, Hotaling JM, Giwercman A, Hwang K, Cheng YS. Male infertility.

Nat Rev Dis Primers. 2023;9(1):49.

3.   Agarwal A, Mulgund A, Hamada A, Chyatte MR. A unique view on male infertility around the globe.
Reprod Biol Endocrinol. 2015;13:37.

4.   Toure A, Martinez G, Kherraf ZE, Cazin C, Beurois J, Arnoult C, et al. The genetic architecture of
morphological abnormalities of the sperm tail. Hum Genet. 2021;140(1):21-42.

5.   Wang WL, Tu CF, Tan YQ. Insight on multiple morphological abnormalities of sperm flagella in male
infertility: what is new? Asian J Androl. 2020;22(3):236-45.

6.   Wang J, Wang W, Shen L, Zheng A, Meng Q, Li H, Yang S. Clinical detection, diagnosis and treatment of
morphological abnormalities of sperm flagella: A review of literature. Front Genet. 2022;13:1034951.

7.   Lu S, Gu Y, Wu Y, Yang S, Li C, Meng L, et al. Bi-allelic variants in human WDR63 cause male
infertility via abnormal inner dynein arms assembly. Cell Discov. 2021;7(1):110.

8.   Houston BJ, Riera-Escamilla A, Wyrwoll MJ, Salas-Huetos A, Xavier MJ, Nagirnaja L, et al. A
systematic review of the validated monogenic causes of human male infertility: 2020 update and a discussion
of emerging gene-disease relationships. Hum Reprod Update. 2021;28(1):15-29.

9.   Jiao SY, Yang YH, Chen SR. Molecular genetics of infertility: loss-of-function mutations in humans and
corresponding knockout/mutated mice. Hum Reprod Update. 2021;27(1):154-89.

10.   Leung MR, Zeng J, Wang X, Roelofs MC, Huang W, Zenezini Chiozzi R, et al. Structural specializations
of the sperm tail. Cell. 2023;186(13):2880-96 e17.

11.   Burgess SA, Walker ML, Sakakibara H, Knight PJ, Oiwa K. Dynein structure and power stroke. Nature.
2003;421(6924):715-8.

12.   Linck RW, Chemes H, Albertini DF. The axoneme: the propulsive engine of spermatozoa and cilia and
associated ciliopathies leading to infertility. J Assist Reprod Genet. 2016;33(2):141-56.

13.   Gunes S, Sengupta P, Henkel R, Alguraigari A, Sinigaglia MM, Kayal M, et al. Microtubular Dysfunction
and Male Infertility. World J Mens Health. 2020;38(1):9-23.

14.   Sironen A, Shoemark A, Patel M, Loebinger MR, Mitchison HM. Sperm defects in primary ciliary
dyskinesia and related causes of male infertility. Cell Mol Life Sci. 2020;77(11):2029-48.

15.   King SM. Axonemal Dynein Arms. Cold Spring Harb Perspect Biol. 2016;8(11).

16.   Walton T, Gui M, Velkova S, Fassad MR, Hirst RA, Haarman E, et al. Axonemal structures reveal
mechanoregulatory and disease mechanisms. Nature. 2023;618(7965):625-33.

17.   Aprea I, Raidt J, Hoben IM, Loges NT, Nothe-Menchen T, Pennekamp P, et al. Defects in the cytoplasmic
assembly of axonemal dynein arms cause morphological abnormalities and dysmotility in sperm cells leading
to male infertility. PLoS Genet. 2021;17(2):e1009306.

18.   Ben Khelifa M, Coutton C, Zouari R, Karaouzene T, Rendu J, Bidart M, et al. Mutations in DNAH1,
which encodes an inner arm heavy chain dynein, lead to male infertility from multiple morphological
abnormalities of the sperm flagella. Am J Hum Genet. 2014;94(1):95-104.

19.   Hwang JY, Nawaz S, Choi J, Wang H, Hussain S, Nawaz M, et al. Genetic Defects in DNAH2 Underlie
Male Infertility With Multiple Morphological Abnormalities of the Sperm Flagella in Humans and Mice. Front
Cell Dev Biol. 2021;9:662903.

20.   Tu C, Nie H, Meng L, Yuan S, He W, Luo A, et al. Identification of DNAH6 mutations in infertile men
with multiple morphological abnormalities of the sperm flagella. Sci Rep. 2019;9(1):15864.

21.   Gao Y, Liu L, Shen Q, Fu F, Xu C, Geng H, et al. Loss of function mutation in DNAH7 induces male
infertility associated with abnormalities of the sperm flagella and mitochondria in human. Clin Genet.
2022;102(2):130-5.

22.   Liu C, Miyata H, Gao Y, Sha Y, Tang S, Xu Z, et al. Bi-allelic DNAH8 Variants Lead to Multiple

Morphological Abnormalities of the Sperm Flagella and Primary Male Infertility. Am J Hum Genet. 2020;107(2):330-41.

23. Tu C, Cong J, Zhang Q, He X, Zheng R, Yang X, et al. Bi-allelic mutations of DNAH10 cause primary male infertility with asthenoteratozoospermia in humans and mice. Am J Hum Genet. 2021;108(8):1466-77.

24. Li Y, Wang Y, Wen Y, Zhang T, Wang X, Jiang C, et al. Whole-exome sequencing of a cohort of infertile men reveals novel causative genes in teratozoospermia that are chiefly related to sperm head defects. Hum Reprod. 2021;37(1):152-77.

25. Whitfield M, Thomas L, Bequignon E, Schmitt A, Stouvenel L, Montantin G, et al. Mutations in DNAH17, Encoding a Sperm-Specific Axonemal Outer Dynein Arm Heavy Chain, Cause Isolated Male Infertility Due to Asthenozoospermia. Am J Hum Genet. 2019;105(1):198-212.

26. Karak S, Jacobs JS, Kittelmann M, Spalthoff C, Katana R, Sivan-Loukianova E, et al. Diverse Roles of Axonemal Dyneins in Drosophila Auditory Neuron Function and Mechanical Amplification in Hearing. Sci Rep. 2015;5:17085.

27. Modiba MC, Nephawe KA, Mdladla KH, Lu W, Mtileni B. Candidate Genes in Bull Semen Production Traits: An Information Approach Review. Vet Sci. 2022;9(4).

28. Hamdi Y, Boujemaa M, Ben Rekaya M, Ben Hamda C, Mighri N, El Benna H, et al. Family specific genetic predisposition to breast cancer: results from Tunisian whole exome sequenced breast cancer cases. J Transl Med. 2018;16(1):158.

29. Tsai CC, Huang FJ, Wang LJ, Lin YJ, Kung FT, Hsieh CH, Lan KC. Clinical outcomes and development of children born after intracytoplasmic sperm injection (ICSI) using extracted testicular sperm or ejaculated extreme severe oligo-astheno-teratozoospermia sperm: a comparative study. Fertil Steril. 2011;96(3):567-71.

30. Colpi GM, Francavilla S, Haidl G, Link K, Behre HM, Goulis DG, et al. European Academy of Andrology guideline Management of oligo-astheno-teratozoospermia. Andrology. 2018;6(4):513-24.

31. Hannah WB, Seifert BA, Truty R, Zariwala MA, Ameel K, Zhao Y, et al. The global prevalence and ethnic heterogeneity of primary ciliary dyskinesia gene variants: a genetic database analysis. Lancet Respir Med. 2022;10(5):459-68.

32. O'Connor MG, Mosquera R, Metjian H, Marmor M, Olivier KN, Shapiro AJ. Primary Ciliary Dyskinesia. CHEST Pulmonary. 2023;1(1):100004.

33. Vanaken GJ, Bassinet L, Boon M, Mani R, Honore I, Papon JF, et al. Infertility in an adult cohort with primary ciliary dyskinesia: phenotype-gene association. Eur Respir J. 2017;50(5).

34. Hornef N, Olbrich H, Horvath J, Zariwala MA, Fliegauf M, Loges NT, et al. DNAH5 mutations are a common cause of primary ciliary dyskinesia with outer dynein arm defects. Am J Respir Crit Care Med. 2006;174(2):120-6.

35. Guan Y, Yang H, Yao X, Xu H, Liu H, Tang X, et al. Clinical and Genetic Spectrum of Children With Primary Ciliary Dyskinesia in China. Chest. 2021;159(5):1768-81.

36. Peng B, Gao YH, Xie JQ, He XW, Wang CC, Xu JF, Zhang GJ. Clinical and genetic spectrum of primary ciliary dyskinesia in Chinese patients: a systematic review. Orphanet J Rare Dis. 2022;17(1):283.

37. Fliegauf M, Olbrich H, Horvath J, Wildhaber JH, Zariwala MA, Kennedy M, et al. Mislocalization of DNAH5 and DNAH9 in respiratory cells from patients with primary ciliary dyskinesia. Am J Respir Crit Care Med. 2005;171(12):1343-9.

38. Fassad MR, Shoemark A, Legendre M, Hirst RA, Koll F, le Borgne P, et al. Mutations in Outer Dynein Arm Heavy Chain DNAH9 Cause Motile Cilia Defects and Situs Inversus. Am J Hum Genet. 2018;103(6):984-94.

39. Zuccarello D, Ferlin A, Cazzadore C, Pepe A, Garolla A, Moretti A, et al. Mutations in dynein genes in

patients affected by isolated non-syndromic asthenozoospermia. Hum Reprod. 2008;23(8):1957-62.

40.   Wallmeier J, Nielsen KG, Kuehni CE, Lucas JS, Leigh MW, Zariwala MA, Omran H. Motile ciliopathies.

Nat Rev Dis Primers. 2020;6(1):77.

41.   Neesen J, Kirschner R, Ochs M, Schmiedl A, Habermann B, Mueller C, et al. Disruption of an inner arm

dynein heavy chain gene results in asthenozoospermia and reduced ciliary beat frequency. Hum Mol Genet.

2001;10(11):1117-28.

42.   Pan MM, Hockenberry MS, Kirby EW, Lipshultz LI. Male Infertility Diagnosis and Treatment in the Era

of In Vitro Fertilization and Intracytoplasmic Sperm Injection. Med Clin North Am. 2018;102(2):337-47.

43.   Esteves SC, Roque M, Bedoschi G, Haahr T, Humaidan P. Intracytoplasmic sperm injection for male

infertility and consequences for offspring. Nat Rev Urol. 2018;15(9):535-62.

44.   Long S, Fu L, Ma J, Yu H, Tang X, Han W, et al. O-253   Novel biallelic variants in DNAH1 cause

multiple morphological abnormalities of sperm flagella with favorable outcomes of fertility after ICSI in Han

Chinese males. Human Reproduction. 2023;38(Supplement_1).

45.   Wambergue C, Zouari R, Fourati Ben Mustapha S, Martinez G, Devillard F, Hennebicq S, et al. Patients

with multiple morphological abnormalities of the sperm flagella due to DNAH1 mutations have a good

prognosis following intracytoplasmic sperm injection. Hum Reprod. 2016;31(6):1164-72.

46.   Li Y, Sha Y, Wang X, Ding L, Liu W, Ji Z, et al. DNAH2 is a novel candidate gene associated with

multiple morphological abnormalities of the sperm flagella. Clin Genet. 2019;95(5):590-600.

47.   Gao Y, Tian S, Sha Y, Zha X, Cheng H, Wang A, et al. Novel bi-allelic variants in DNAH2 cause severe

asthenoteratozoospermia with multiple morphological abnormalities of the flagella. Reprod Biomed Online.

2021;42(5):963-72.

48.   Wei X, Sha Y, Wei Z, Zhu X, He F, Zhang X, et al. Bi-allelic mutations in DNAH7 cause

asthenozoospermia by impairing the integrality of axoneme structure. Acta Biochim Biophys Sin (Shanghai).

2021;53(10):1300-9.

49.   Zheng R, Sun Y, Jiang C, Chen D, Yang Y, Shen Y. A novel mutation in DNAH17 is present in a patient

with multiple morphological abnormalities of the flagella. Reprod Biomed Online. 2021;43(3):532-41.

50.   Huang F, Zeng J, Liu D, Zhang J, Liang B, Gao J, et al. A novel frameshift mutation in DNAH6

associated with male infertility and asthenoteratozoospermia. Front Endocrinol (Lausanne). 2023;14:1122004.

51.   Li L, Sha YW, Xu X, Mei LB, Qiu PP, Ji ZY, et al. DNAH6 is a novel candidate gene associated with

sperm head anomaly. Andrologia. 2018.

Figure Legends

Figure 1. Identification of biallelic pathogenic variants in *DNAH3* from four unrelated infertile families. (**A**) Pedigrees of four families affected by *DNAH3* variants (M1–M7). Black arrows indicate the probands in these families. (**B**) Location of the variants and conservation of affected amino acids in DNAH3. Black arrows indicate the position of the variants. (**C**) Immunofluorescence staining of DNAH3 in sperm from the patients and normal control. Red, DNAH3; green, α-Tubulin; blue, DAPI; scale bars, 5 μm. (**D**) Western blotting analysis of DNAH3 expressed in spermatozoa from the patients and normal control.

Figure 2. **Defects in sperm morphology of the patients harboring *DNAH3* variants**. (**A**, **B**) Abnormal sperm morphology was observed through Papanicolaou staining (A), and SEM analysis (B) compared to normal control. Scale bars, 5 μm.

Figure 3. Ultrastructural and mitochondrial defects in sperm from infertile men with *DNAH3* variants. (**A**) TEM analysis of sperm obtained from a normal control and patients harboring *DNAH3* variants. Cross-sections of the midpiece, principal piece and endpiece of sperm from normal control showed the typical "9 + 2" microtubule structure, and an IDA and an ODA were displayed on the A-tube of each microtubule doublet. Cross-sections of the midpiece, principal piece and endpiece of sperm from the patients displayed absent or disordered CPs, MTDs and ODFs, as well as an evident missing of the IDAs in different pieces of the flagella. M, mitochondria sheath; ODF, outer dense fiber; MTD, microtubule doublets; CP, central pair; IDA, inner dynein arms; ODA, outer dynein arms. Scale bars, 200

nm. (**B**) Immunofluorescence staining of TOM20 in sperm from the patients and normal

control. Red, TOM20; green, α-Tubulin; blue, DAPI; scale bars, 5 μm.

Figure 4. *Dnah3* KO male mice are infertile. (**A**) Fertility of *Dnah3* KO mice. The KO

male mice were infertile (n = five biologically independent WT mice or KO mice; Student's

t test; *, P < 0.05; NS, no significance; error bars, s.e.m.). (**B**) TEM analysis of the

cross-sections of spermatozoa from *Dnah3* KO mice revealed an obvious absence of IDAs

in different pieces of the flagella compared to WT mice. M, mitochondrion sheath; ODF,

outer dense fiber; MTD, microtubule doublet; CP, central pair; IDA, inner dynein arm; ODA,

outer dynein arm. Scale bars, 200 nm. (**C**) Disrupted mitochondria were observed in

spermatozoa tail from *Dnah3* KO mice by TEM analysis. The yellow arrows indicate the

normal mitochondria. The red arrowheads indicate the dilated intermembrane spaces and

dissolved mitochondrial material. M, mitochondrion sheath. Scale bars, 200 nm. (**D**)

Immunofluorescence staining of SLC25A4 indicated impaired mitochondrial formation in

Dnah3 KO mice compared to WT mice. Red, SLC25A4; green, α-Tubulin; blue, DAPI;

scale bars, 5 μm.

Figure 5. **Immunofluorescence staining and western blotting analysis of IDA-associated**

proteins in spermatozoa obtained from normal control and patients with *DNAH3*

variants. (**A** – **C**) Immunofluorescence staining of DNAH1 (A), DNAH6 (B) and DNALI1

(C) in spermatozoa from patients and normal controls. Red, DNAH1 in (A), DNAH6 in (B),

DNALI1 in (C); green, α-Tubulin; blue, DAPI; scale bars, 5 μm. (**D** – **F**) Western blotting

analysis of DNAH1(D), DNAH6 (E), DNALI1 (F) in sperm lysates from the patients and normal control.

Figure 6. Immunofluorescence staining and western blotting analysis of IDA-associated proteins in spermatozoa from WT and *Dnah3* KO mice. (**A** – **C**) Immunofluorescence staining of DNAH1 (A), DNAH6 (B) and DNALI1 (C) in spermatozoa from *Dnah3* KO and WT mice. Red, DNAH1 in (A), DNAH6 in (B), DNALI1 in (C); green, α-Tubulin; blue, DAPI; scale bars, 5 μm. (**D** – **F**) Western blotting analysis of DNAH1(D), DNAH6 (E) and DNALI1 (F) in spermatozoa lysates from *Dnah3* KO and WT mice.

Figure 7. ICSI outcomes of *DNAH3*-deficient patients and *Dnah3* KO mice. (**A**) The embryonic development of Patient 1 and Patient 3 after ICSI treatment. MII, metaphase II; PN, pronucleus; scale bars, 40 μm. (**B**) There was no difference in the fertilization rate or 2-cell and blastocyst embryo formation rates between the *Dnah3* KO and WT groups (n = three biologically independent WT mice or KO mice; Student's *t* test; NS, no significance; error bars, s.e.m.).

672 **Tables**

673 **Table 1. Semen analysis of the patients in the present study.**

		Patient 1	Patient 2	Patient 3	Patient 4	Reference
Semen parameters	Semen volume (ml)	4.3	3.3	0.8	4.6	≥1.5
	Semen concentration (10^6/ml)	2.0	0.5	11.0	21.0	≥15.0
	Motility (%)	6.0	3.0	0	0	≥40.0
	Progressive motility (%)	0	2.3	0	0	≥30.0
Sperm morphology	Normal (%)	1.3	1.1	1.8	1.2	≥4.0
	Head defect (%)	14.2	18.3	10.3	15.7	-
	Tail defect (%)	91.3	87.5	96.5	97.5	-

674 -, not applicable.

675   **Table 2. Variants analysis of the patients in the present study.**

		Patient 1		Patient 2		Patient 3		Patient 4
Variant	cDNA mutation*	c.3590C>G	c.3590C>T	c.4837G>T	c.5587del	c.10355C>T	c.2314C>T	c.4045G>A
	Protein alteration	p.P1197R	p.P1197L	p.A1613S	p.L1863*	p.S3452L	p.R772W	p.D1349N
	Mutation type	Missense	Missense	Missense	Nonsense	Missense	Missense	Missense
Allele frequency	ExAC Browser	0	0	0.00037135	0	0.0001	0.0005	0.0002
	GnomAD	0	0	0.00026658	0	0.0001	0.0006	0.0002
	1000 Genomes Project	0	0	0.00099840	0	0.000199681	0.00139776	0.000798722
Function prediction	SIFT	Deleterious	Deleterious	Deleterious	/	Tolerated	Deleterious	Deleterious
	Polyphen-2	Probably damaging	Probably damaging	Probably damaging	/	Probably damaging	Probably damaging	Probably damaging
	Mutation Tester	Disease causing	Disease causing	/	/	Disease causing	Disease causing	Disease causing

676   *, NM_017539.2; /, not applicable.

678   **Table 3. Semen analysis using CASA in the *Dnah3* KO mice.**

	WT	KO	P[b] value
Semen parameters			
Sperm concentration (10^6 /ml) [a]	112.32 ± 18.26	105.17 ± 11.15	0.059
Motility (%) [b]	71.56 ± 3.97	4.37 ± 1.15	<0.01
Progressive motility (%) [b]	60.36 ± 4.32	4.37 ± 1.15	<0.01
Sperm locomotion parameters			
Curvilinear velocity (VCL) (μm/s) [b]	67.54 ± 6.79	9.07 ± 1.22	<0.01
Straight-line velocity (VSL) (μm/s) [b]	28.91 ± 4.86	2.68 ± 0.52	<0.01
Average path velocity (VAP) (μm/s) [b]	39.02 ± 5.31	3.85 ± 0.82	<0.01
Amplitude of lateral head displacement (ALH) (μm) [b]	0.71 ± 0.03	0.13 ± 0.04	<0.01
Linearity (LIN) [b]	0.43 ± 0.07	0.30 ± 0.02	0.037
Wobble (WOB, = VAP/VCL) [b]	0.58 ± 0.06	0.42 ± 0.05	0.024
Straightness (STR, = VSL/VAP)	0.74 ± 0.22	0.70 ± 0.13	0.80
Beat-cross frequency (BCF) (Hz) [b]	4.86 ± 0.12	0.73 ± 0.08	<0.01

679   [a] Epididymides and vas deferens.

680   [b] A significant difference, two-sided student's t-test. n = 3 biologically independent WT mice or KO

681   mice.

Table 4. Outcomes of ICSI treatment in the patients with *DNAH3* mutations.

	Patient	Patient 1	Patient 2		Patient 3	Patient 4			
Subjects	Female age (y)	24	30		30	36			
	Length of primary infertility history (y)	6	1		3	8			
Basal hormones	FSH (IU/L)	7.1	8.5		3.49	4.3			
	LH (IU/L)	4.44	2.5		4.2	2.6			
	E2 (pg/mL)	83	50		43.49	68			
	Prog (ng/mL)	0.2	0.6		0.3	0.3			
		Cycle 1	Cycle 1	Cycle 2	Cycle 1	Cycle 1	Cycle 2	Cycle 3	Cycle 4
ICSI Cycles	E2 level on the trigger day (pg/ml)	3366	1519.6	1582.4	>5000	4440	5000	3152	2206
	No. of follicles ≥ 14 mm on the trigger day	15	6	5	20	13	12	14	11
	No. of follicles ≥ 18 mm on the trigger day	10	3	4	8	11	9	7	5
	No. of oocytes retrieved	24	6	5	25	16	20	21	14
ICSI progress	No. of MII oocytes	21	6	5	20	12	13	8	7
	Fertilization rate (%)	80.95 17/21)	50 (3/6)	100 (5/5)	95 (19/20)	41.67 (5/12)	46.15 (6/13)	50 (4/8)	42.86 (3/7)
	Cleavage Rate (%)	100 (17/17)	100 (2/2)	100 (5/5)	100 (19/19)	100 (5/5)	100 (6/6)	100 (4/4)	66.7 (2/3)
	Available D3 embryos	13	2	5	15	2	2	0	0
	Blastocyst formation rate (%)	61.5 (8/13)	0	66.7 (2/3)	66.7 (10/15)	-	-	-	-
Clinical outcomes	No. of embryos transferred	2 blastocysts	0	2 D3	1 blastocyst	2 D3	2 D3	-	-
	Implantation rate (%)	50 (1/2)	0	0	100 (1/1)	0	0	-	-
	Clinical pregnancy	Yes	No	No	Yes	-	-	-	-
	No. of live birth	ongoing	-	-	ongoing	-	-	-	-

684 -, not applicable.

Supporting information

Figure S1. The expression of DNAH3 in mouse testis. (**A**) qPCR analysis revealed that *Dnah3* was highly expressed in the mouse testis. (**B**) qPCR analysis showed that *Dnah3* expression was significantly elevated beginning on postnatal Day 12, peaked at postnatal Day 30, and maintained a stable expression level thereafter.

Figure S2. DNAH3 is expressed during spermatogenesis in mice and humans. (**A**) Immunofluorescence staining of DNAH3 in isolated mouse germ cells. Pink, PNA; green, DNAH3; blue, DAPI; scale bars, 5 μm. (**B**) Immunofluorescence staining of DNAH3 in isolated human germ cells. Pink, PNA; green, DNAH3; blue, DAPI; scale bars, 5 μm.

Figure S3. Generation of *Dnah3* KO mice. (**A**) Schematic illustration of the strategy for the generation of *Dnah3* KO mice. (**B, C**) PCR sequencing (B) and qPCR (C) were used to confirm the genotype and KO efficiency (n = three biologically independent WT mice or KO mice; Student's *t* test; *, P<0.05; error bars, s.e.m.). (**D**) Immunofluorescence staining of DNAH3 in testis of *Dnah3* KO mice and WT mice. Green, DNAH3; blue, DAPI; scale bars, 75 μm. (**E**) Immunofluorescence staining of DNAH3 in spermatozoa isolated from the cauda epididymis of *Dnah3* KO mice and WT mice. Red, DNAH3; green, α-Tubulin; blue, DAPI; scale bars, 5 μm.

Figure S4. Ciliary development of *Dnah3* KO mice. (**A**) H&E staining of lung, brain, eye, and oviduct from *Dnah3* KO mice and WT mice. Scale bars, 100 μm. (**B**) Analysis of ciliary

development in the lung, brain, eye, and oviduct from *Dnah3* KO mice and WT mice by using immunofluorescence staining. Green, Ac-Tubulin; blue, DAPI; scale bars, 20 μm.

Figure S5. Fertility of *Dnah3* KO mice. (**A**) H&E staining of ovary tissue sections from 8-week-old *Dnah3* KO female mice and WT female mice. Scale bars, 75 μm (n = three biologically independent WT mice or KO mice). (**B**) Sizes of the testis and epididymis of the 8-week-old *Dnah3* KO and WT mice (n = three biologically independent WT mice or KO mice; Student's *t* test; NS, no significance; error bars, s.e.m.).

Figure S6. Morphology of sperm isolated from *Dnah3* KO mice. (A, B) Papanicolaou staining (A) and SEM analysis (B) revealed morphological defects in partial spermatozoa from *Dnah3* KO mice compared to WT mice. Scale bars in (A), 5 μm; scale bars in (B), 2.5 μm.

Figure S7. Immunofluorescence staining of ODA-associated proteins in spermatozoa obtained from variants within *DNAH3* patients. (**A** – **C**) The expression of DNAH8 (A), DNAH17 (B) and DNAI1 (C) in spermatozoa of the patients was comparable to that in normal controls. Red, DNAH8 in (A), DNAH17 in (B), DNAI1 in (C); green, α-Tubulin; blue, DAPI; scale bars, 5 μm.

Figure S8. Immunofluorescence staining of ODA-associated proteins in spermatozoa of *Dnah3* KO and WT mice. (**A** – **C**) The expression of DNAH8 (A), DNAH17 (B) and

DNAI1 (C) in spermatozoa from *Dnah3* KO mice was comparable to that in spermatozoa

from WT mice. Red, DNAH8 in (A), DNAH17 in (B), DNAI1 in (C); green, α-Tubulin; blue,

DAPI; scale bars, 5 μm.

Table S1. Primer pairs used in the present study.

Movie S1. CASA of sperm from WT mice. Sperm from the epididymis of WT mice were

collected, incubated, and recorded under a phase-contrast microscope. A normal quantity

and motility of sperm were observed in the WT mice (n = three biologically independent WT

mice).

Movie S2. CASA of sperm from *Dnah3* KO mice. Epididymal sperm of *Dnah3* KO mice

were collected, incubated in HTF medium at 37 °C for 10 minutes, and recorded under a

phase-contrast microscope. The movie showed a significantly reduced motility of sperm

from *Dnah3* KO (n = three biologically independent *Dnah3* mice).