# Genetic barcodes for ash (*Fraxinus*) species and generation of new wide hybrids

William J. Plumb[1,2,3], Laura J. Kelly[1,2], Joe Mullender[2], Robyn F. Powell[2], Miguel Nemesio-Gorriz[3], David Carey[4], Mary E. Mason[4], William Crowther[1], Jennifer Koch[4], Gerry C. Douglas[3], Richard J. A. Buggs*[1,2]

[1] School of Biological and Behavioural Sciences, Queen Mary University of London, London, UK.
[2] Royal Botanic Gardens, Kew, Richmond upon Thames, UK.
[3] Forestry Development Department, Teagasc, Ashtown, Dublin, Republic of Ireland
[4] United States Department of Agriculture, Forest Service, Northern Research Station, Delaware, OH, USA.

*author for correspondence: r.buggs@qmul.ac.uk

## Abstract

Native ash tree species in Europe and North America are being devastated by ash dieback and the emerald ash borer, respectively. As worldwide ash species differ in their level of susceptibility to these threats, hybrid breeding may allow resistance to be transferred among species. However, we do not know the extent to which distantly related ash species can be crossed, and many ash species are difficult to identify from morphology alone leading to some mislabelling in living collections. While sequences from the internal transcribed spacer (ITS) of nuclear ribosomal DNA can identify some *Fraxinus* species, this is often hindered by intragenomic variation in ITS sequences within the genus. Here, we develop a genetic barcode system for the identification of *Fraxinus* species based on three low-copy-number protein coding genes. We also conduct experimental crosses among ash species in different sections. Our barcodes are effective in identifying ash samples to sectional level and in some cases to species level, and can also identify hybrids. They highlight that *F. mandschurica, F. platypoda* and *F. chiisanensis* are frequently mistaken for one another in living collections. We succeeded in generating ten wide hybrid plants: two of *F. pennsylvanica* X *F. excelsior* and eight of section *Ornus* X *F. excelsior*. One hybrid from each of our crosses has survived natural ash dieback inoculation in Ireland. Our findings facilitate the deployment of global ash species diversity in response to alien pests and pathogens.

## 34 Societal impact statement

The world-wide diversity of ash trees includes genetic information encoding resistance to the ash
dieback fungus and the emerald ash borer beetle, which are currently devastating ash populations in
Europe and North America. In order to mobilise this genetic diversity to counter the devastation, we
need to be able to accurately identify ash species from around the world, and cross them with one
another. Here, we present a genetic barcoding system for ash species, and a series of hybridisation
experiments between European ash and other species. Two of the hybrids show early promise against
ash dieback.

## 42 Introduction

In the temperate regions of Europe, common ash (*Fraxinus excelsior*) is an ecologically and
economically important broadleaf tree species (Dobrowolska *et al.*, 2011; Mitchell *et al.*, 2014) but is
severely affected by the rapid spread of the non-native fungal species, *Hymenoscyphus fraxineus*,
causing symptoms commonly referred to as "ash dieback" (ADB) or "Chalara" (Pautasso *et al.*, 2013;
Gross *et al.*, 2014; Coker *et al.*, 2019). European ash populations are further threatened by an insect
species, the Emerald Ash Borer (EAB) (Orlova-Bienkowskaja & Bieńkowski, 2018) which has already
devastated ash populations in North America (Herms & McCullough, 2014). Worldwide species of ash
differ in their levels of resistance to ADB and EAB (Koch *et al.*, 2007; Kowalski *et al.*, 2015; Rigsby *et*
*al.*, 2016; Nielsen *et al.*, 2016; Plumb *et al.*, 2019; Kelly *et al.*, 2020). Thus some non-native *Fraxinus*
species could be suitable to replace susceptible populations (Jepson & Arakelyan, 2017; Marzano *et al.*,
2019; Lévesque *et al.*, 2023) or used as parents in hybrid breeding programmes to transfer resistance
among species (Koch *et al.*, 2007, 2012; Plumb *et al.*, 2019).

Several classifications of the genus *Fraxinus* have been published (e.g. Wenzig, 1883; Wesmael, 1892;
Lingelsheim, 1907; Miller, 1955; Nikolaev, 1981; Wallander, 2008, 2012) with differing species
divisions and sectional divisions. In this paper we follow the classification of Wallander (2013), where
the genus contains over 40 species, many of which are placed into six sections. Our only exception to
this is that we do not place *F. griffithii* into section *Ornus* due to its placement outside the *Ornus* clade
in the phylogeny of Kelly *et al.* (2020) and Hinsinger *et al.* (2013). *Fraxinus* species can be difficult to
tell apart as many of their most important morphological features are only visible at certain times of
year, or are age-dependent (Henry, 1914; Wallander, 2012). Some closely-related species are
morphologically similar and have been defined partly on the basis of their geographic distribution.
Some species show high intraspecies variability and phenotypic plasticity. It is hard to differentiate
between some *Fraxinus* species using DNA sequencing of the ITS region, as sequence similarity is high

between closely related species and intragenomic variation is present within some individuals

(Wallander, 2008). Ploidy level (and hence C-value) is a useful method to distinguish among certain

species but most species are diploid and others, such as *F. caroliniana*, include more than one ploidy

level (Whittemore *et al.*, 2018).

Hybridisation occurs between some species of *Fraxinus*. The sister species *F. excelsior* and *F.*

*angustifolia* naturally hybridise in Europe (Fernández-Manjarrés *et al.*, 2006; Heuertz *et al.*, 2006;

Thomasset *et al.*, 2011) and have also been crossed by breeders (Raquin *et al.*, 2002). Many of

the *Fraxinus* species in the section *Melioides* clade hybridise in areas of coexistence in North America

(Wallander, 2008). There is anecdotal evidence of natural hybridisation in regions of species overlap

between section *Melioides* and section *Ornus* (Wallander, 2012). Henry (Henry, 1914) documented

various artificial interspecific crosses of *Fraxinus* species, though the extent to which they were

successful is unclear. Hybrids of *F. nigra* X *F. mandshurica* (both within the section *Fraxinus*) were

produced by Davidson and two commercial clones, *Northern treasure* and *Northern gem*, were patented

(Davidson, 1999; Davidson & Ronald, 2001). Recently, Koch et al. have successfully produced new *F.*

*nigra* X *F. mandshurica* hybrids and second-generation backcrosses appear to be resistant to EAB

(Koch *et al.*, 2007, 2010). Within-section hybrids between *F. mandshurica* and *F. angustifolia*  subsp.

*syriaca* (syn. *F. sogdiana* used by authors of the study) have been produced and propagated using

embryo culture, with one clone showing photosynthetic heterosis (He *et al.*, 2021, 2023). Wider hybrids

between sections *Fraxinus* and *Melioides* have also been produced in China involving *F. mandshurica*

and *F. americana*, and *F. mandshurica* and *F. velutina*, using electrostatic treatments to overcome

fertility barriers; these hybrids showed heterosis in growth (Zeng *et al.*, 2015; He *et al.*, 2019).

Hybrid breeding programmes have been set up in response to other well-established pathogens of trees,

such as Dutch elm disease (Smalley & Guries, 1993; Santini *et al.*, 2007) and chestnut blight

(Anagnostakis, 2012), producing resistant trees. The process of interspecific hybridisation is usually not

as simple as identifying two trees of different species and cross-pollinating them. Prezygotic

incompatibilities include: flowering time differences between the species, failure of pollen germination

on the stigma of a different species, and reduction or halting of pollen tube growth.  Post-zygotically,

the hybrid endosperm or embryos may fail to develop. Interventions may allow some of these barriers

to be overcome, including: pollen storage, mentor pollen (Knox *et al.*, 1987) and embryo rescue

(Sharma *et al.*, 1996; Raquin *et al.*, 2002).

The growing number of research programmes that involve different species of ash and their hybrids

means that a method for their easy identification, usable on any tissue type at any time of year, is highly

desirable. Whole-genome sequences for 22 *Fraxinus* species have recently been published (Kelly *et al.*,

2020). This provided us with the opportunity to undertake genome-wide searches for nuclear loci

suitable for barcoding within the genus. Here, we describe development of a set of barcoding regions

using single-copy genes in *Fraxinus*. We show that these barcodes can identify specimens to sectional

level and often to species level. We also produce new inter-sectional *Fraxinus* hybrids, and verify their

parentage using our new barcodes.

## Materials and Methods

***Fraxinus* accessions**

We collected materials from *Fraxinus* accessions in arboreta and botanic gardens (Supplementary

Table 1) and from the United Stated Department for Agriculture Forest Service ash breeding program at

the Northern Research Station (in Delaware, Ohio). We focused on these, rather than herbarium

specimens, as they are potential sources for: experiments, future planting stocks and hybridisation

programmes. Some of these specimens were used in separate studies as sources of scion material to set

up assays for EAB susceptibility in Ohio (Kelly *et al.*, 2020) and trials for ADB susceptibility in

England (see: www.forestresearch.gov.uk/research/testing-a-range-of-ash-species-for-tolerance-to-ash-

dieback/). The living collections in the British Isles that contributed materials were: Royal Botanic

Gardens Kew (including Wakehurst), Cambridge University Botanic Garden, Westonbirt Arboretum,

Earth Trust, RHS Wisley, Ness Botanic Gardens, Royal Botanic Garden Edinburgh, Dawyck Botanic

Garden, The National Botanic Gardens Glasnevin (Ireland), The John F. Kennedy Arboretum (Ireland)

and a private collection belonging to Gerry Douglas (Dublin, Ireland). We did not have access to some

accepted taxa of *Fraxinus*, mainly due to poor representation of species native to Mexico and South

America in British and Irish collections. As these species do not grow well in European conditions, they

are unlikely to be part of the solution to ADB and EAB. DNA was extracted from a total of 150

specimens from living collections, using Qiagen DNeasy® Plant Mini Kit (QIAGEN Ltd., Manchester,

UK) or CTAB protocols (Kelly et al., 2020).

126
**Selection of candidate barcoding genes**

A genome assembly for *F. excelsior* was published in 2017 (Sollars *et al.*, 2017) based on a tree grown

from seed collected in a woodland in Gloucestershire, UK. For a further 28 of the samples we used,

Kelly et al (2020) had previously sequenced whole genomic DNA (see Column E in Supplementary

Table 1) and produced genome assemblies. The species identifications of these samples had been

checked by Dr Eva Wallander (an expert on the genus) based on morphology and ITS, and where these

differed from the labels in the living collections we took the identification by Wallander to be correct.

In the case of accessions 1973-6204 and chi-12 there was uncertainty about the identity and Kelly et al

(2020) initially  labelled them as *Fraxinus* sp. From the whole genome assemblies of these 29 *Fraxinus*

| 136 | samples, Kelly et al (2020) had identified groups of putatively orthologous protein coding gene |
| 137 | sequences in them and three outgroups (*Olea europaea*, *Erythranthe guttata* and *Solanum* |
| 138 | *lycopersicum*) using OMA (Altenhoff *et al.*, 2019). Full details of methods can be found in Kelly *et al.* |
| 139 | (2020), but briefly, the full-length sequences (i.e. including both exons and introns, where present) for |
| 140 | OMA groups that included sequences from all 29 *Fraxinus* samples and the three outgroups were |
| 141 | aligned using MUSCLE (Edgar, 2004) via GUIDANCE (Sela *et al.*, 2015) and unreliably aligned |
| 142 | positions removed. OMA groups with alignments shorter than 300 characters in length, or which |
| 143 | included sequences with <10% non-gap characters were excluded, leaving a set of 1385 OMA groups. |
| 144 | |
| 145 | The alignments for these groups were examined using a pre-release version of CONTEXT, a |
| 146 | phylogenomic dataset browser (v0.8. pre-release) (https://github.com/lonelyjoeparker/qmul-genome- |
| 147 | convergence-pipeline/blob/master/CONTEXT.md) to generate statistical information about each OMA |
| 148 | group including longest sequence length, longest un-gapped region, number of variant nucleotides and |
| 149 | amino acids, and the mean entropy values for nucleotides and amino acids. We selected candidates for |
| 150 | further analysis based on their: degree of variance (25% or above), greatest overall length, and length of |
| 151 | the longest un-gapped region (being longer than 800bp). For these candidate OMA groups, maximum |
| 152 | likelihood trees were generated using RAXML-ng version 0.8.0 (Kozlov *et al.*, 2019), using the |
| 153 | following parameter settings: --model GTR+G --tree pars{10},rand{10} --bs-trees 1000. We examined |
| 154 | the trees for congruence with the species-tree from Kelly *et al.* (2020). We checked for putative |
| 155 | paralogs by searching the sequence from *F. excelsior* against its reference genome (assembly version |
| 156 | BATG0.5) using BLAST and OMA groups were excluded if the sequence matched another region of |
| 157 | the genome with total coverage of greater than 75% of the query sequence length. The sequences for the |
| 158 | three outgroups were removed from the remaining candidate OMA groups and new alignments |
| 159 | generated using the MUSCLE tool from EMBL-EBI (https://www.ebi.ac.uk/Tools/msa/muscle/ |
| 160 | accessed October 2017). |
| 161 | |
| 162 | **Primer design and testing** |
| 163 | |
| 164 | We selected four variable OMA groups for primer design. These were OG8679 (a pentatricopeptide |
| 165 | repeat, according to the functional annotation of BATG0.5), OG9972 (a histidine phosphotransferase |
| 166 | protein), OG22833 (a protein of unknown function)  and OG8143 (a polyadenylate-binding protein 1). |
| 167 | Primers were designed using Primer3web v4.0.0 (http://bioinfo.ut.ee/primer3/ accessed September |
| 168 | 2017) and purchased from Eurofins MWG Operon (Ebersberg, Germany). The primers were provided |
| 169 | lyophilised, and Milli-Q® $H_2O$ was used to reconstitute them to a concentration of 100 pmoles/μl. We |
| 170 | tested these by performing PCR on our DNA extractions from 151 trees in living collections. |
| 171 | |

Each PCR reaction contained <100/ng genomic DNA, 1μl forward primer 10pmol/μl, 1μl reverse

primer 10pmol/μl, 25μl DreamTaq Green PCR Master Mix (2X) (DreamTaq DNA Polymerase), 2X

DreamTaq Green buffer, dNTPs, and four mM $MgCl_2$ (Thermo Fisher), 10μl 5X TBT-PAR (containing

trehalose, bovine serum albumin (BSA), and polysorbate-20 (Tween-20), and 2μl DMSO (Dimethyl

sulfoxide). Reactions were made up to 50μl with Milli-Q® $H_2O$. The conditions used were: an initial

denaturation stage at 94ºC for 6 min; followed by 38 cycles of denaturation at 94ºC for 1 min, annealing

at 52ºC for 1 min and extension at 72ºC for 2 min; with a final extension at 72ºC for 7 min. PCRs were

carried out on a GeneAmp® PCR System 9700 thermocycler (Applied Biosystems, Warrington, UK).

Agarose gel electrophoresis was used to determine the presence of correctly sized PCR products. Gels

were prepared by dissolving 1.5g molecular grade agarose (Bioline Reagents Ltd., London, UK) in

150ml 0.5x Tris-borate EDTA (TBE) (27 g of Tris base, 13.75 g of boric acid and 10 ml of 0.5 M

Ethylenediaminetetraacetic acid (EDTA) (pH 8.0), made up to 1 litre using Milli-Q® $H_2O$) using a

microwave to heat the solution. Once the agarose was fully dissolved it was left to cool to 55°C before

mixing 6μl 10mg/ml ethidium bromide (EtBr) into the solution. The solution was then poured into a

prepared gel tray and left to cool at room temperature until it had solidified. Once the gel had set, it was

placed into a Mini-Sub® Cell gel tank (Bio-Rad Laboratories Ltd., Hemel Hempstead, UK) and

submerged in 0.5x TBE buffer. Gels were run at 100V / 600mA for 30 min with the amplicon samples

and a 1 Kb Plus DNA Ladder (Thermo Fisher). The gels were visualised using a UV trans-illuminator

Gel Doc 2000 and Quantity One 1-D analysis v4.6.5 software (Bio-Rad).

For sequencing, the PCR products were purified using a NucleoSpin® Gel and PCR Clean-up kit

(Macherey-Nagel). Sequencing reactions were made up of 20-25/ng of the purified PCR product, 0.5ul

at 5pmol/μl of appropriate primer, 1ul 5x sequencing buffer, 0.25ul DMSO (Dimethyl sulfoxide) and

0.25ul BigDye® Direct Sanger Sequencing Kit Thermo Fisher Scientific UK Ltd (Loughborough).

Cycle sequencing was performed using the following parameters: 35 cycles of denaturation at 96ºC for

10 s, annealing at 52ºC for 5 s and extension at 60ºC for 4 min using a GeneAmp® PCR System 9700

thermocycler (Applied Biosystems, Warrington, UK). Samples were Sanger-sequenced using a 3730

Series Genetic Analyzer Thermo Fisher Scientific UK Ltd (Loughborough). The electropherograms

were trimmed and the forward and reverse sequence for each sample were assembled into contigs using

Geneious® ver8.1.9 software (Biomatters, Ltd., Auckland, New Zealand).

We aligned sequence data for the four barcode regions from all successfully sequenced samples using

MUSCLE (Edgar, 2004) within JALVIEW (Procter *et al.*, 2021). We trimmed the alignment to the ends

of the outermost PCR primers. Each sample was given a label that began with an abbreviation of its

taxonomic section (DIP = *Dipetala*, FRA = *Fraxinus*, MEL = *Melioides*, ORN = *Ornus*, PAU =

*Pauciflorae*, SCI = *Sciadanthus*, INC = *Incertae sedis*), followed by its species ID, followed by its

sequencing tube number. For the samples that had been used for whole genome sequencing and barcode
primer design, the label contained the section and species allocated after identification by Eva
Wallander using morphology and ITS data. For the other samples, the label contained section and
species according to labels in living collections. We manually sorted the alignment by section and
species. When this was done it was apparent that in all barcodes there were SNV alleles specific to
sections and some species, but that some samples were anomalous. The anomalous samples were given
the prefix ANOM and were reassigned to different sections according to the alleles they contained, with
their new section abbreviation placed at the start of their label. Within sections, we then sought to
manually cluster species, where possible, according to species-specific SNV alleles. Where a species
included no samples that had been whole genome sequenced, we assumed that the majority of living
collection labels that matched the correct section were also correct at a species level, allowing us to
provisionally assign some of the anomalous samples to species. For each sample, we noted the SNV
loci in each barcode that allowed assignment to section and, where relevant, species. Once we had as far
as possible identified each sample (that had not been included in the whole genome sequencing panel)
using barcodes, we compared these identifications, where possible, to independent identifications of the
samples made by Eva Wallander using morphology and ITS, and also flow cytometry assessment of
genome size.
226
227 **New hybrid crosses**
228
In order to produce new interspecies crosses, we took scion material from a collection of mature *F.*
*excelsior* plus trees selected from Irish forests for good phenotypic traits. These were: five
predominantly female trees (Fex 32, 076, 8X and G 40) and five predominantly males (Fex 359, 42,
405-14 and G1). We took scion material from the John F. Kennedy Arboretum, Co. Wexford, Ireland,
from trees labelled as: *F. pennsylvanica* (JFK.02931), *F. paxiana* (JFK.02928), *F. floribunda*
(JFK.02916), *F. japonica* (a synonym for *F. chinensis* in (Wallander, 2012))(JFK.02920), *F. chinensis*
(JFK.02906), *F. chinensis* subsp. *rhynchophylla* (JFK.02908), *F. retusa henryana* (a synonym for *F.*
*floribunda* in (Wallander, 2012))(JFK2005.0132), *Fraxinus nigra* Marsh. × *Fraxinus mandshurica*
(Canadian Hybrid No 8921 "Northern Treasure" JFK.2001.0159), *F. americana* (JFK.02899). From
Glasnevin we took scions from: *F texensis* (a synonym for *F. albicans* in (Wallander,
2012))(XX.011046 - A2?) and *F. mandschurica* (1934.011053.A2). We also took scion material from
trees labelled as *F. mandshurica* (T1-T3) in Dr Gerry Douglas' private garden.
241
Scions were cut to length 15-20 cm and ~0.5cm diameter, with at least two sets of lateral buds and one
terminal bud. These scions were cleft-grafted to rootstocks at Teagasc, Kinsealy, Ireland between 2011-
2015 as follows. Two year old trees from naturally occurring Irish *F. excelsior* populations were
sourced from a commercial nursery. During the period February-March, as the scion and rootstocks lay

dormant, the  rootstock trees were cut to 5-10 cm above soil level on the main stem, and cleft. The basal

end of the scion was cut to two opposing smooth-tapered cuts approximately 2-4 cm in length from the

lowest set of buds using a sharp grafting knife. The cut scions were inserted into the rootstock cleft,

with the cambium of the scions positioned into direct contact with the cambium of the rootstock. The

graft union was bound using elastic grafting bands. The entire scion and graft union was dipped into

molten paraffin wax (45°C).  Grafted plants were potted into 3L pots with commercial-grade peat

compost and maintained in the greenhouse. Sprouts that arose from below the graft union were

routinely removed as these can lead to graft rejection. Grafted trees were later moved to the Teagasc

Ashtown Research Centre, Ireland in 2018, following the Kinsealy site's closure.

Cross-pollination was carried out between March and April of 2015 as some of the grafted trees came

into flower. The grafted plants were at least two years old at the time of flowering and crossing. The

following non-native accessions were used in the crosses:  trees labelled as *F. mandshurica* in Dr Gerry

Douglas' private garden were used as male parents (no female flowers were available), *F. pennsylvanica*

(JFK.02931) used as a female parent, *F. chinensis* (JFK.02906) was used as a male and a female parent,

*F. retusa henryana* (JFK2005.0132) was used as a female parent,  *F. japonica* (JFK.02920) was used as

a male parent, and *Fraxinus nigra* Marsh. X *Fraxinus mandshurica* (Canadian Hybrid No 8921

"Northern Treasure" JFK.2001.0159) was used as a female parent. Each cross was with *F. excelsior* as

the other parent. In addition to the grafted *F. excelsior* plus trees 076, 8X, G40, 32, 42, 405-14, G1, and

359, we also used *F. excelsior* plus tree 98 which was growing outside nearby.

Pollen was collected from male flowers at anthesis by gently tapping inflorescences so that pollen fell

into glass petri dishes (60 x 17mm; Thermo Fisher). Dishes were labelled with the pollen donors'

accession number. Glass was used instead of plastic petri dishes for collection and storage, to avoid

static electricity making the pollen challenging to manipulate. Pollen was stored for future use by

wrapping parafilm around the petri dish's lid, leaving a small opening for drying. Wrapped dishes were

then placed in a glass desiccator containing silica gel and stored in a refrigerator at 4°C until required in

the same season.  In some cases when the non-native species were hermaphrodite, the male flowers

were emasculated to prevent self-fertilisation. Pollinations were  performed at least twice for each cross

using pollen collected on the same day or from refrigerator-stored pollen and within a few days of the

initial pollination event. Stigmas on selected female trees were checked daily over the period in which

the inflorescence was maturing and pollinated when some of them showed a glistening surface. Pollen

was applied using a tiny paintbrush that was lightly brushed against the stigmas to release a puff of

pollen. A tag was placed around each inflorescence of the pollinated tree, with an allocated number

corresponding to the parental details for the cross. In one case two different pollen donors *F. excelsior*

405-14 and *F. excelsior* G1 were used.  Pollen of *F. excelsior* 405-14 was used on the first day and

pollen of  *F. excelsior* G1 was used on a subsequent day for the same cross as 'mentor pollen' may

facilitate hybridisation in some cases. The pollination period was generally in the last two weeks of

March when stigmas were deemed to be most receptive. After the first pollination event the trees were

moved to a separate section of the greenhouse to minimise potential contamination by nearby ash trees

which may have been flowering. After that, the pollinated trees were grown in the glasshouse for the

summer period in which unpollinated pistils abscised naturally. As *F. excelsior* plus tree 98 was field

grown, the pollinated female inflorescences were enclosed and sealed in pollination bags and unbagged

inflorescences were used as open pollinated (OP) intra-species controls under the assumption that any

ambient pollen would be from *F. excelsior.*

The pollinated inflorescences were observed for the development of the samaras over several months

during which many immature/unfertilised samaras abscised naturally. During the last week of August

and the first two in September, an assessment was made of seed production from crosses. Full samaras

were excised, and surface sterilised in 7% (w/v) calcium hypochlorite $Ca(ClO)_2$ solution for 20 min,

followed by three rinses in sterile water and aseptic air drying. Thereafter the samaras were dissected

aseptically under a stereomicroscope, and the seeds were removed and distributed onto B5 medium for

germination (Gamborg et al. 1968) in small Petri plates (5 cm) containing 6-7 ml of medium with

sucrose in concentrations of 2%, 3%, 4%, 5% (w/v) and solidified with phytagel 3 g $L^{-1}$. Culture

conditions were at $22 \pm 1^{o}C$ with 16-hour photoperiod using Philips® cool white fluorescent tubes

giving a photon flux density of 58.6 µmol m-1s-1. Seed germination and viability were assessed after

approximately six weeks and after this period, ungerminated seeds were either transferred to fresh

media, or the embryos were excised from seeds for embryo rescue culturing. In some cases we did not

attempt whole seed germination but embryos were excised and cultured directly after seed collection

and surface sterilisation. Embryo rescue was performed under a stereomicroscope in a laminar flow

hood under aseptic conditions. An incision was made along the longitudinal axis of the seed and into

the endosperm and the outer seed coat was peeled off. The seed was prised open at the cotyledon end to

reveal the embryo attached within one half of the seed.  The embryo was gently prised off the

endosperm using the tip of a scalpel and placed directly into B5 medium. Whole seeds and isolated

embryos were transferred to fresh media every 6-8 weeks at least six times. We routinely used B5

medium with 3% sucrose in the latter culture periods. Germinated seedlings were transferred to glass

jars (volume, 150 ml) on WPM medium salts (Lloyd and McCown 1980) and vitamins of B5 medium

(Gamborg et al. 1968) supplemented with 3% w/v activated charcoal and then weaned to the

glasshouse.

The *Fraxinus* genetic barcodes were used to identify successful new hybridisations. DNA extraction,

PCR and sequencing was performed in the same way as it was for the living collections (see above).

Sanger sequence reads were then trimmed, cleaned and assembled into contigs. These were viewed

using Geneious prime version 8.1 and we looked for double peaks at SNP loci that differed between the

putative parental species sequenced in the test panel. We carried out an initial screening using the

barcode region OG9927 and hybrids found with this locus were also sequenced with the barcoding

regions OG22833 and OG8143. The new hybrids were grown in an area of high ADB disease pressure,

and records were kept of death of trees due to ADB.

**Ploidy investigations**

Where polyploidy has been reported for certain *Fraxinus* taxa, measurement of genome sizes using

flow cytometry can aid in the identification of species and their hybrids. In such cases, nuclear DNA

contents were estimated following the one-step flow cytometry procedure (Dolezel *et al.*, 2007).

Approximately 1 cm$^2$ of the sample leaf material was incubated for 30 seconds on ice in 1 ml of

'general-purpose buffer' (GPB) (Loureiro *et al.*, 2007) supplemented with 3% PVP-40, after which leaf

material from the calibration standard *Petroselinum crispum* (Mill.) Fuss or *Pisum sativum* was added

(Obermayer *et al.*, 2002). This material was chopped together rapidly using a fresh razor blade. Another

1 ml of the isolation buffer was then added to the chopped material, and the homogenate was filtered

through a 30µm nylon mesh (Celltrics 30µM mesh, Sysmex, Goritz, Germany). Following filtering, 100

µl propidium iodide (1 mg/mL) was added, and the sample was incubated on ice for 10 minutes before

analysis. The relative fluorescence of 5000 particles was then recorded using a Partec Cyflow SL3 flow

cytometer (Partec GmbH, Münster, Germany) fitted with a 100 mW green solid-state laser (532 nm,

Cobolt Samba, Solna, Sweden) and the output histograms were analysed with the FlowMax software

v.2.4 (Partec GmbH, Münster, Germany).

# Results

**Testing of barcodes**

Each barcode region (Table 1) amplified successfully and yielded sequence data for the majority of our 150 DNA samples from 45 *Fraxinus* species or subspecies (see Supplementary Table 1). Usable sequences were generated for 147 samples with OG9927, 140 samples with OG22833, 134 samples with OG8143 and 149 for OG8679.  If DNA quality was poor, the second primer sets for OG9927 and OG8679 tended to produce lower-quality reads. Trimmed alignments of all clean sequences generated can be found in Supplementary Files 1-4, ordered by section and species. These are also available at https://doi.org/10.5281/zenodo.10066755. Examination of these alignments showed that all of the barcodes contained SNV alleles that were specific to taxonomic sections and present in all samples within those sections. There were also many SNV alleles that were specific to species, though these were not always present in all samples from a species.

Supplementary Table 1 gives a list of all tree accessions used, and details of the extent to which they can be identified to section and species levels using each DNA barcode. Barcodes OG9972, OG22833 and OG8143 gave conflicting sectional results for only three samples. In contrast, barcode OG8679 gave many double peaks in chromatograms and gave sequence results that sometimes conflicted with the other barcodes for nine samples. In the sequence alignment for OG8679, several SNP loci had three or even four alleles present among the samples. To give one example, site 546 was:  G in the majority of samples,  T in all four samples of *F. paxiana*, one of eight individuals of *F. floribunda*, one of six individuals of *F. excelsior,* C in two of three individuals of *F. lanuginosa*, A in all individuals later identified as *F. chiisanensis* (except one which was R (A/G)) and one of eight individuals of *F. quadrangulata* and in two of four individuals of *F. platypoda*, R in one of seven *F. nigra* and one of five *F. latifolia*. Those individuals with an A at position 546 all had an A at position 592, a G at position 657 and a T at position 674. Those that had an R at position 546 had an A or R at position 592, a K at position 657 and a T or Y at position 674. Such a pattern is best explained by the presence of two paralogs, with some samples showing amplification of both, and some only amplifying one copy, potentially due to loss or mutation of the primer site at the other copy. There were no cases in which OG8679 was able to give us sectional or species assignments for a sample that were unavailable from one of the other barcodes. Therefore we decided to discard OG8679 as a barcode and focus only on OG9972, OG22833 and OG8143.

Using OG9972, OG22833 and OG8143, 148 samples could be assigned to section (taking a majority

vote in the three cases of sectional conflict) and 82 could be assigned to species. Those 66 samples that

could not be identified to species level were all in the sections *Melioides*, *Fraxinus*, *Dipetalae* or *Ornus*.

These sections are known to have similar species that sometimes hybridise. No alleles were found to be

specific to *F. excelsior* (in section *Fraxinus*), the most widespread species in Europe, or *F.*

*pennsylvanica* and *F. americana* (in section *Melioides*), the most widespread species in North America,

probably due to widespread hybridisation with other species found within their ranges.

Of the 148 samples assigned to section or species level by the three barcodes, 70 were also identified by

Eva Wallander using morphology and ITS. Of these, only three were assigned to a different section by

the two methods of identification. One of these three cases was also one of the three cases where the

three barcodes were in conflict, and in this case one of the three barcodes agreed with Eva Wallander's

identification: this is Kew accession 1928-8802 *F. angustifolia* subsp. *oxycarpa* (in section *Fraxinus*)

which was assigned to the section Fraxinus by OG9927 but to Dipetala by OG22833 and OG8147, and

identified as *F. angustifolia* by Eva Wallander. It is possible that we made an error in labelling the tubes

used for OG22833 and OG8147. The other two disagreements between the barcode identifications and

those of Eva Wallander were for RHS Wisley W862006 *F. greggii* (*Pauciflorae*) and Kew 1981-8204

*F. latifolia* (*Melioides*) which were both assigned to the section *Fraxinus* by OG9927 and OG8147 but

identified as *F. cuspidata* (*Incertae sedis*) and *F. latifolia* (*Melioides*) by Eva Wallander. Again, errors

may have been responsible for this conflict. In the 67 out of 70 cases where the barcodes and Eva

Wallander's identifications were to the same section, 34 were identified to species level by the both

methods. In 30 cases these agreed. The cases of disagreement were for *F. pennsylvanica*/*F.*

*caroliniana*/*F. velutina* which are all closely related within section *Melioides*, or *F. chinensis*/*F.*

*baroniana*/*F. floribunda* in section *Ornus*.

All three barcodes were each able to identify most samples to sectional level. However, section

*Fraxinus* was hard to identify using OG22833 and OG8143. For section *Fraxinus,* OG9972 performed

best, with three clear section-specific alleles (Table 2). Overall, OG9972 identified 48 samples to

species level, OG22833 identified 63 samples to species level, and OG8143 identified 46 samples to

species level. Within section *Dipetala*, OG9972 performed best at identifying species and OG8143

could not distinguish any species. Within section *Fraxinus*, OG9972 and OG22833 both performed

better than OG8143, but OG8143 was particularly useful for distinguishing *F. mandschurica*. In section

*Melioides*, all barcodes were unable to distinguish species, with the exception of *F. velutina*, which

could be identified by OG22833. In section *Ornus*, the barcodes differed widely in their patterns of

diversity: OG9972 was good for identifying *F. floribunda* and *F lanuginosa*; OG22833 was good for

distinguishing *F. chinensis* and OG8143 was good for distinguishing *F. chinensis and F. ornus*. Within

the small section *Pauciflorae*, all barcodes could distinguish *F. greggii* but only OG8143 could

distinguish *F. gooddingii*. All barcodes performed well at distinguishing the species not assigned to sections (*Incertae sedis*) but OG8143 was alone in being unable to identify *F. griffithii* to species level.

Of our 148 sample identifications with barcodes, 26 conflicted with the labels in the living collections that they were from. Twenty of these conflicts involved assignments of different sections, and six were assignments of different species within sections. Of the 20 sectional conflicts, seven identifications were also available from Eva Wallander, of which five agreed with the barcodes. Of the two cases where the barcodes differed from both the label and Wallander, one was the aforementioned case of Kew accession 1928-8802 *F. angustifolia* subsp. *oxycarpa* where OG9927 differed from the other two barcodes in its assignment, in agreement with Wallander and the label. The other case was Kew 1981-8204 *F. latifolia* (*Melioides*) where Wallander agreed with the label but the barcodes suggested section *Fraxinus*.

There were two cases where the tree label said *F. platypoda* (*Incertae sedis*) but the barcodes indicated *F. mandschurica* (Fraxinus). These were JFK.02932 and USDA FS plat-3, and for the latter, the barcode identification was confirmed by Wallander. The barcodes assigned Kew 1900-44501 *F. quadrangulata* (*Dipetala*) and Kew 2011-1622 *F. quadrangulata* (*Dipetala*) to section *Fraxinus*. A sample labelled *F. greggii* (Cambridge 19860253*A) was identified as *F. cuspidata* (*Incertae sedis*) by the barcodes. A sample labelled *F. chinensis* subsp. *chinensis* (*Ornus*)(Westonbirt 56.0455) was identified as *F. pennsylvanica* (*Melioides*) by both the barcodes and Wallander. A sample labelled *F. platypoda* (*Incertae sedis*)(USDA FS plat-1) was identified as *F. pennsylvanica* (*Melioides*) by both the barcodes and Wallander (it is now relabelled in the USDA FS collection as pe-103). Three samples labelled as *F. nigra* (*Fraxinus*)(Wakehurst 1996-5167 and 1996-5169 in replicate) were identified as section *Melioides* by the barcodes, and 1996-5169 was identified as *F. pennsylvanica* (*Melioides*) by Wallander. One sample labelled *F. americana* (Melioides)(JFK.02899) was identified as *F. chinensis* (Ornus) by the barcodes; morphological evaluation of this accession is needed. Seven samples labelled as *F. mandschurica* (GD-T3, GD-T1 and Kew 1989-3691, 1989-8285, 1992-375, 1992-376 and 1992-377) in section *Fraxinus* were identified by the barcodes as being not in the section *Fraxinus*. Closer examination of their genome size and bud morphology (see details below) allowed them to be identified as *F. chiisanensis* (*Incertae sedis*).

Of the six cases where the barcodes assigned a different species within the same section as the living collection label, three of them were also examined by Wallander who in two cases made the same identification as the label (USDA FS PE_48 FRAX10 *F. pennsylvanica* which the barcodes assigned to *F. velutina* and USDA FS Bar-2 FRAX28 *F. baroniana* which the barcodes assigned to *F. chinensis*), suggesting the possibility of a DNA tube labelling error. In one case Wallander made the same identification as the barcodes (Kew 1973-6204 FRAX29 *F. bungeana* which the barcodes assigned to

*F. ornus*), suggesting that the barcoding was accurate. The other three cases involved: JFK.02906 *F.*

*chinensis* being assigned to *F. ornus*; Kew 2004-1231 *F. paxiana* and USDA FS BUN-5 *F. bungeana*

being assigned to *F. chinensis*. Details of the SNP loci used for each assignment can be found in

Supplementary Table 1 and seen in Supplementary Files 1-3.

*Fraxinus excelsior* (USDA FS ex-12) had several positions in OG9927 that were diagnostic for *F.*

*cuspidata* or were heterozygous for a base found in section *Fraxinus* and a base found in at least one

individual of *F. cuspidata*, suggesting it could be a hybrid between these two species. However in

OG22833 and OG8143 there was no evidence of this, suggesting possible contamination in the OG9927

PCR reaction.

**Genome size measurements**

Genome sizes were measured for several *Fraxinus* accessions. These results are shown in

Supplementary Table 1, column Y, with inferred ploidy level from these genome sizes in column Z.

The majority of samples had 1C genome sizes between 700Mb and 1000Mb and were inferred to be

diploid. Some, but not all, accessions identified by barcodes as *F. chinensis* were hexaploid: Westonbirt

5.0456, USDA FS chi-16, JFK.02920, JFK.02908 and 2GD were hexaploid but JFK.02899 and USDA

FS bar-2 (FRAX 28) were diploid. Kew's 1992-377 and the samples in Gerry Douglas' garden, all

labelled as *F. mandschurica* but with anomalous barcode sequences, were hexaploid (unlike JFK 02932

identified by barcodes as *F. mandschurica* which had a diploid C-value). The other polyploids found

were all samples for which the barcodes had failed to sequence: *F. biltmoreana* Westonbirt 5.0463

(FRAX17) was hexaploid, *F. profunda* Wakehurst 1993-1493 appeared to be octoploid, *F. lanuginosa*

RBGE 20071296-G (FRAX22) was tetraploid, and *F. uhdei* USDA FS uhdei-ponto (FRAX34) was

hexaploid; it is likely that the barcodes did not work well in these samples due to divergent

homeologous copies of the barcode sequences among their sub-genomes.

*Fraxinus chiisanensis*

Closer analysis of the morphology of the samples labelled as *F. mandschurica* at Kew and in Gerry

Douglas's garden but with anomalous barcodes and hexaploidy showed them to have naked buds which

is characteristic of *F. chiisanensis* (Wallander, 2012), a rare endemic of Korea (Kim *et al.*, 2022). The

seeds for the trees in Gerry Douglas' garden were sourced from South Korea. A chromosome squash on

a seedling from one of the Kew trees, which was hermaphroditic (1992-377), was also consistent with

hexaploidy. A previous unpublished study also suggested that *F. chiisanensis* is hexaploid (mentioned

in Siljak-Yakovlev *et al.*, 2014). Interestingly, the pattern of barcode polymorphism within these

putative *F. chiisanensis* samples showed some unique alleles (OG9972: 1563, 1091, 1187, 764;

OG8143: 390, 709) but also patterns of polymorphism involving alleles common in different sections

(OG9972: 123, 153, 381, 405, 1048, 1487, 1698). In OG8143, *F. chiisanensis* shared alleles with

*Melioides* at some loci (OG8143: 262-267, 801) but shared alleles with everything but *Melioides* at

other loci (OG8143: 200, 237, 736-738). In OG22833, *F. chiisanensis* shared alleles with everything

but section *Ornus* at 66-67, everything but *Melioides* at 144, everything but *Melioides* and *Dipetala* at

366, everything but *Ornus* and *Sciadanthus* at 372 and everything but *Pauciflorae* at 272. Thus, within

its six genome copies, *F. chiisanensis* harbours a complex mix of variation found scattered across other

sections of the genus.

**Hybrid crosses**

Due to the barcoding and flow cytometry results reported above, several of the trees that were grafted

for hybridisation were re-identified, including some that were used as parents in our crosses. *Fraxinus*

*chinensis* (JFK.02906) was identified as diploid *F. ornus*. *Fraxinus japonica* (JFK.02920) (which, as

noted above, is a synonym for *F.  chinensis* (Wallander, 2012)) was identified as the hexaploid *F.*

*chinensis.* The trees labelled  *F. mandshurica* in Dr Gerry Douglas' private garden were identified as

hexaploid *F. chiisanensis*. *Fraxinus retusa henryana* (a synonym for *F. floribunda* in (Wallander,

2012))(JFK2005.0132), was confirmed as a diploid in section *Ornus* but could not be assigned to a

species; we cannot be sure if it is *F. floribunda* or *F. ornus*. *Fraxinus pennsylvanica* (JFK.02931) was

confirmed as being in section *Melioides* but could not be assigned to a species by the barcodes; we

therefore have no reason to believe that the label is incorrect. *Fraxinus nigra* Marsh. $\times$ *F. mandshurica*

(Canadian Hybrid No 8921 JFK.2001.0159) was not examined using barcodes, but we are confident

that its parentage is reliably known (Davidson & Ronald, 2001).

Of the 35 crosses and controls that were performed (Supplementary Table 2), 756 seeds were produced,

of which 363 were propagated, producing 131 seedlings of which 85 survived to be viable plants. We

carried out an initial screening using the barcode region OG9927 on 84 seedlings that were alive at the

time of the screening. This identified 16 putative hybrids showing heterozygosity at sites that varied

among parental species. We then also sequenced these individuals with the barcoding regions OG22833

and OG8143. Region OG2833 was found to produce clear sequence reads of a comparable length to the

average sequence length for the pure *Fraxinus* species. However, for the barcode region OG8143 we

found only the forward primer was able to produce useable sequence reads for all of the putative

hybrids, the useable region after sequence clean-up was found to be ~650bp, this is ~100bp short of the

length of the full region.

None of the crosses that used non-native species as pollen donors produced hybrid offspring. In all of

these cases, the offspring were sired by *F. excelsior* pollen. This includes all crosses involving *F.*

*chiisanensis*, and flow cytometry confirmed that the progeny of the *F. excelsior* mother tree used in this

cross were all diploid. There were 15 crosses done with non-native species as the maternal parent. Four

of these produced no progeny. Seven of them had progeny that were confirmed as hybrid. We identified

14 intersectional hybrids; this included two plants of *F. pennsylvanica* X *F. excelsior*, six of *F. ornus* X

*F. excelsior* and five section *Ornus* X *F. excelsior.*

Both of the *F. pennsylvanica* X *F. excelsior* hybrids displayed vigorous growth and had no ash dieback

symptoms in 2021, despite growing in an area with prevalent spore inoculum. One of these was still

alive and healthy in July 2023. Most of the hybrids produced between *F. ornus* or section *Ornus* and *F.*

*excelsior* died of ash dieback symptoms. Four were dead by the end of summer 2020. In 2021, one had

severe symptoms, three had mild symptoms, and three were healthy. One was still alive and healthy in

July 2023.

Sequence data for these two surviving hybrids for OG22833 and OG9927 are included in

Supplementary Files 1 and 2. For OG9927, in the *F. pennsylvanica* X *F. excelsior* hybrid, positions

416, 454, 603, 648, 743, 839, 1148, 1193 and 1482 were heterozygous. These sites are all diagnostic for

section *Fraxinus* or section *Melioides* (see Table 2) confirming that this individual is a hybrid. In the

section *Ornus* X *F. excelsior* hybrid, positions 1148, 1465, 1482, 1538 and 1660 are heterozygous,

confirming its hybridity between section *Ornus* and section *Fraxinus*. For OG22833, in the *F.*

*pennsylvanica* X *F. excelsior* hybrid, positions 144, 180 and 366 were heterozygous, which are

diagnostic for *Melioides* (Table 2) and position 272. In the section *Ornus* X *F. excelsior* hybrid,

positions 66 and 67, which are diagnostic for section *Ornus* (Table 2), were heterozygous, as were

positions 116, 266 and 332 which are fixed in one of the sections but polymorphic in the other.

# Discussion

We have developed three low-copy nuclear gene barcodes that can delineate taxonomic sections and in

some cases species within the genus *Fraxinus*, allowing identification of ash samples and verification of

hybrids. The cases where the barcodes did not perform well were certain polyploid species, and within

large sections that contain many closely related species with a shared geographic distribution. Sample

identifications using the barcodes sometimes differed from living collection labels but seldom differed

from identifications by Eva Wallander who is a taxonomic expert on the genus.

Use of these barcodes on living collections shows that some ash species are commonly mislabelled. The

rare Korean endemic *F. chiisanensis* is commonly labelled as *F. mandschurica*, and genuine samples of

*F. mandschurica* are sometimes labelled as *F. platypoda*. These, and other mis-labellings of Asiatic

species, may reflect lower acquaintance of European curators with Asiatic species than with European

and American species.  Several cases of mislabelling in living collections involve a local species being

given an exotic label: in some cases this may be because exotic scion material had been grafted to a

local rootstock, but shoots from the rootstock have become dominant and the exotic material has not

survived. Other cases of mislabelling may be *ad hoc* errors within collections. Our findings highlight

the need for verification of living collection labels before undertaking experiments or initiating

breeding programmes.

We generated and verified wide hybrids of ash, among sections *Fraxinus*, *Melioides* and *Ornus*: these

were from *F. pennsylvanica* X *F. excelsior* and section *Ornus* X *F. excelsior* crosses. Many hybrids

have been reported or made previously within the genus *Fraxinus* (see Introduction) but ours is the first

study we are aware of to have confirmed the production of intersectional hybrid plants between sections

*Ornus* and *Fraxinus*. Unlike the wide hybridisations previously conducted between sections *Fraxinus*

and *Melioides* (*F. mandshurica* X *F. americana*, and *F. mandshurica* X *F. velutina*)(Zeng *et al.*, 2015;

He *et al.*, 2019), we did not use electrostatic treatments to overcome fertility barriers.

One of our major objectives when setting up crosses was to hybridise *F. excelsior* with east Asian ash

species possessing high resistance to both EAB and ADB. We have not achieved this, partly due to

initial misidentification of parental trees, some of which were labelled as Asiatic species, but turned out

not to be. We attempted many crosses with *F. chiisanensis* pollen, but these were unsuccessful. Our *F.*

*chiisanensis* trees did not generate any female flowers: we might have had more success with female

flowers as exclusion of ambient pollen would then have been less of an issue. We cannot exclude the

possibility that there are intrinsic incompatibilities between *F. excelsior* and *F. chiisanensis*. While *F.*

*excelsior* is a diploid, *F. chiisanensis* is hexaploid and our barcodes suggest it could have a complex

hybrid origin.

Whilst the majority of our hybrids died from ash dieback, one individual from each of these crosses

remained alive and healthy in summer 2023. Although *F. pennsylvanica* is highly susceptible to EAB

(Koch *et al.*, 2015; Orlova-Bienkowskaja & Bieńkowski, 2016), it does appear to be less susceptible to

ADB than *F. excelsior* (Nielsen *et al.*, 2016; Drenkhan *et al.*, 2017; Plumb *et al.*, 2019). Thus, crosses

between *F. pennsylvanica* and *F. excelsior* may have promise for ADB resistance breeding, especially

as *F. pennsylvanica* is a good timber species. Section *Ornus* contains species with low susceptibility to

ADB and EAB (Nielsen *et al.*, 2016; Drenkhan *et al.*, 2017; Plumb *et al.*, 2019; Kelly *et al.*, 2020), so

hybrids from this section may also be of interest.

The barcodes and hybrids presented here provide methodological and biological resources for the use of

world-wide ash germplasm for resistance to pests and pathogens and adaptation to climate change. The

success of our wide hybridisations suggests that hybridisations may be possible throughout the genus

*Fraxinus*, allowing effective mobilisation of genetic diversity throughout the genus in hybrid breeding

programmes. Planting of non-native species, or hybrid breeding programmes between *F. excelsior* and
non-native species, may be part of the solution to the threats of ADB and EAB, especially for
commercial timber production and rapid carbon sequestration.
597

# 598 Acknowledgements

We thank Eva Wallander for her expert identifications of numerous samples used in this study. We
thank Jaume Pellicer for advice and assistance with flow cytometry. We thank Timothy Baxter, Samuel
Brockington, Peter Brownless, Dan Crowley, Dawn Edwards, Simon Honey, Ross Irvine, Richard
Jinks, Penny Jones, Tony Kirkham, Hugh McAllister, Iain Parkinson and Sara Redstone for help with
obtaining *Fraxinus* materials from UK collections. We thank Colin Kelleher, National Botanic Gardens
of Ireland, Glasnevin for provision of scion wood of exotic species and John McNamara for technical
assistance in grafting. R.J.A.B. and L.J.K. acknowledge funding from the Erica Waltraud Albrecht
Endowment Fund and the Living with Environmental Change Tree Health and Plant Biosecurity
Initiative – Phase 2 (grant no. BB/L012162/1), funded jointly by BBSRC, Defra, ESRC, Forestry
Commission, NERC and the Scottish Government. W.J.P. was supported by a Walsh Fellowship from
Teagasc, and funding from Defra.
610

# 611 Author contributions

R.J.A.B., W.J.P., L.J.K, J.K. and G.C.D. planned and designed the research. W.J.P., J.M., D.C., and
W.C. extracted DNA. R.F.P. conducted flow cytometry. G.C.D. conducted hybridisations. W.J.P.,
L.J.K, M.N.G., M.E.M. and J.K. collated samples. R.J.A.B., W.J.P., L.J.K, J.K. and G.C.D. wrote the
manuscript.

# Tables

**Table 1. Primers designed and tested for *Fraxinus* species barcoding.**

Each primer's location is given based on *F. excelsior* BATG0.5 genome. OG numbers refer to the

Orthology group number assigned to genes in Kelly et al 2020.

| *F. excelsior* position and gene name | Primer name | Primer sequence (5'-3') | Tm* | Expected amplicon length |
|---|---|---|---|---|
| Contig238 91392-93122 histidine phosphotransferase protein OG9927 | 9927(1)fwd | TGGAAGGCAGAGTTTCTAGGG | 59.8°C | ~800bp |
| | 9927(1)rev | GAGCAGAGTGMTGCATTGAA | 56.3°C | |
| | 9927(2)fwd | GCTTGGGCRTTGTCAAAGAT | 56.3°C | ~800bp |
| | 9927(2)rev | TTAGCATRGTATGCCCCAGT | 56.3°C | |
| Contig5804 11231-10691 Unknown function OG22833 | 22833fwd | TTGACATGGGATTYGAGATGT | 54.9°C | ~500bp |
| | 22833rev | ACCATAAAGCTGCACAAACCA | 55.9°C | |
| Contig803 88622-87665 polyadenylate-binding protein 1 OG8143 | 8143fwd | CACCTCGAACCTTAYCTGCT | 58.3°C | ~800bp |
| | 8143rev | TCTCGYAGTCCTTCRATGGT | 57.3°C | |
| Contig382 73390-74770 Pentatricopeptide repeat OG8679 | 8679(1)fwd | CAAAAACCAAAGCSGAAATC | 53.2°C | ~800bp |
| | 8679(1)rev | TTCCCATCTTYCTCATGCTC | 56.3°C | |
| | 8679(2)fwd | ACRCTTTTGACCATGTATGTTG | 55.6°C | ~800bp |
| | 8679(2)rev | GCCTGTCTGAAAACCCAAGT | 57.3°C | |

620 *Tm, melting temperature.

**Table 2. Diagnostic SNV allele locations for sections and species of the genus *Fraxinus* using three barcodes.** Locations refer to positions in the alignments in Supplementary Files 1-3, where the first position in the first base in the forward primer. Locations in brackets mean that there is an allele unique to the species but not fixed within the species, so its presence is diagnostic but its absence is not. The plus sign is used for loci that must be in a particular combination to identify a species or section. The sections are according to Wallander (2013) classification with the exception of *F. griffithii* which we place as *Incertae Sedis* due to its placement in the species-tree inferred in Kelly *et al.* (2020).

| Section | Species | OG9972 SNPs | OG22833 SNPs | OG8143 SNPs |
|---|---|---|---|---|
| **Dipetala** | | 100, 342, 1093, 1405-1407 | 147, 182, 419 | 109, 122, (227), 303, 484, 522 |
| | *F. anomala* | 1238 | | |
| | *F. quadrangulata* | (268), (498), (1057) | (443), (465) | (529) |
| **Fraxinus** | | 618+743, 1482, (347), (1538) | 66-67 + 362-364 + 366-373, (352) | 304, (360), (648) |
| | *F. angustifolia* | (805), (897), (1256) | (369), (412) | |
| | *F. mandschurica* | (452), (574), (749) | 324 | (129), (239) (262), 315, (767) |
| | *F. nigra* | 1663 | (275), 340, 425, 446 | |
| **Melioides** | | 603, 648, 839, 955 1048, 1149, 1193 | 144, 180, 366 | 200, 237, 757 |
| | *F. caroliniana* | 676 | | |
| | *F. velutina* | | 24, 110 | |
| **Ornus** | | 381 | 66, 67, (255) | (156), (174), (205), (249), (321), (374), (676) |
| | *F. chinensis* | | | (156) |
| | *F. floribunda* | (207), (361), (1605) | (194) | |
| | *F. ornus* | | | (427), (750) |
| | *F. lanuginosa & F. sieboldiana* | | | 169 |
| **Pauciflorae** | | 293, 316, 567, 729, 1071, 1163, 1470, 1701, 1716 | 275, 364, 426 | 109, 526, 535 |
| | *F. greggii* | 1686 | 160, 190, 223 | 725 |
| **Sciadanthus** | | | | |
| | *F. xanthoxyloides* | 698, 1577 | 338, 381, 509 | 782 |
| **Incertae sedis** | | (1619) | | |
| | *F. chiisanensis* | (153), (241), (463), (603), (624), (764), (1091), (1185), (1187), (1563) | 66-67 + 144 + 366 + 372 + 272 | (390), 709 |
| | *F. cuspidata* | 313, 319, 323, (537), 1152, 1335, 1412 | 45, 182, 223, 359, 415 | 86, (173-226), 263, (340), 421, 495, 559, 621 |
| | *F. griffithii* | 150, 408, 490, 894, 1140-1154, 1185, 1468, 1496, 1564 | 43, 76, 219, 362, 469, 494 | |
| | *F. platypoda* | 1438, 1469 | 38 | 182, 409 |

# References

**Altenhoff AM, Levy J, Zarowiecki M, Tomiczek B, Warwick Vesztrocy A, Dalquen DA, Müller S, Telford MJ, Glover NM, Dylus D,** *et al.* **2019**. OMA standalone: orthology inference among public and custom genomes and transcriptomes. *Genome research* **29**: 1152–1163.

**Anagnostakis SL. 2012**. Chestnut Breeding in the United States for Disease and Insect Resistance. *Plant disease* **96**: 1392–1403.

**Coker TLR, Rozsypálek J, Edwards A, Harwood TP, Butfoy L, Buggs RJA. 2019**. Estimating mortality rates of European ash (Fraxinus excelsior) under the ash dieback (Hymenoscyphus fraxineus) epidemic. *Plants, People, Planet* **1**: 48–58.

**Davidson CG. 1999**. 'Northern treasure' and 'Northern gem' hybrid ash. *HortScience* **34**: 151–152.

**Davidson CG, Ronald WG. 2001**. Hybrid ash tree named 'Northern Treasure'. *US Patent App. 09/233,440*.

**Dobrowolska D, Hein S, Oosterbaan A, Wagner S, Clark J, Skovsgaard JP. 2011**. A review of European ash (Fraxinus excelsior L.): implications for silviculture. *Forestry* **84**: 133–148.

**Dolezel J, Greilhuber J, Suda J. 2007**. Estimation of nuclear DNA content in plants using flow cytometry. *Nature protocols* **2**: 2233–2244.

**Drenkhan R, Agan A, Palm K, Rosenvald R, Jürisoo L, Maaten T, Padari A, Drenkhan T. 2017**. Overview of ash and ash dieback in Estonia. In: Vasaitis R, Enderle R, eds. Dieback of European Ash (Fraxinus spp.): Consequences and Guidelines for Sustainable Management. Swedish University of Agricultural Sciences, 115–124.

**Edgar RC. 2004**. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic acids research* **32**: 1792–1797.

**Fernández-Manjarrés JF, Gerard PR, Dufour J, Raquin C, Frascaria-Lacoste N. 2006**. Differential patterns of morphological and molecular hybridization between Fraxinus excelsior L. and Fraxinus angustifolia Vahl (Oleaceae) in eastern and western France. *Molecular ecology* **15**: 3245–3257.

**Gross A, Holdenrieder O, Pautasso M, Queloz V, Sieber TN. 2014**. Hymenoscyphus pseudoalbidus, the causal agent of European ash dieback. *Molecular plant pathology* **15**: 5–21.

**Henry A. 1914**. The artificial production of vigorous trees. *Department of Agriculture and Technical Instruction for Ireland Journal* **XV**: 34–52.

**Herms DA, McCullough DG. 2014**. Emerald ash borer invasion of North America: history, biology, ecology, impacts, and management. *Annual review of entomology* **59**: 13–30.

**Heuertz M, Carnevale S, Fineschi S, Sebastiani F, Hausman JF, Paule L, Vendramin GG. 2006**. Chloroplast DNA phylogeography of European ashes, Fraxinus sp. (Oleaceae): roles of hybridization and life history traits. *Molecular ecology* **15**: 2131–2140.

**He L, Xu Y, Zeng F, Tian H, Xiao Y, Liu H, Yu L, Zhan Y. 2021**. Establishment of a

microppropagation supporting technology for the Fraxinus mandshurica × Fraxinus sogdiana. *In*
*Vitro Cellular & Developmental Biology - Plant* **57**: 307–318.

**He L, Yan J, Lu H, Su W, Gao S, Wang Y, Zhan Y, Zeng F**. **2023**. Phenotypic Differences of
Leaves and Transcriptome Analysis of Fraxinus mandshurica × Fraxinus sogdiana F1 Variety.
*Forests, Trees and Livelihoods* **14**: 1554.

**He L, Zeng F, He Z, Li Y, Li S, Zhan Y**. **2019**. Interspecific hybridizations of Fraxinus L. (F.
mandshurica × F. americana and F. mandshurica × F. velutina) and heterosis analysis and
selection of F1 progenies. *Canadian journal of forest research. Journal canadien de la*
*recherche forestiere* **49**: 1265–1276.

**Hinsinger DD, Basak J, Gaudeul M, Cruaud C, Bertolino P, Frascaria-Lacoste N,**
**Bousquet J**. **2013**. The phylogeny and biogeographic history of ashes (fraxinus, oleaceae)
highlight the roles of migration and vicariance in the diversification of temperate trees. *PloS*
*one* **8**: e80431.

**Jepson PR, Arakelyan I**. **2017**. Developing publicly acceptable tree health policy: public
perceptions of tree-breeding solutions to ash dieback among interested publics in the UK.
*Forest Policy and Economics* **80**: 167–177.

**Kelly LJ, Plumb WJ, Carey DW, Mason ME, Cooper ED, Crowther W, Whittemore AT,**
**Rossiter SJ, Koch JL, Buggs RJA**. **2020**. Convergent molecular evolution among ash
species resistant to the emerald ash borer. *Nature ecology & evolution* **4**: 1116–1128.

**Kim C, Kim D-K, Sun H, Kim J-H**. **2022**. Phylogenetic relationship, biogeography, and
conservation genetics of endangered Fraxinus chiisanensis (Oleaceae), endemic to South
Korea. *Plant diversity* **44**: 170–180.

**Knox RB, Gaget M, Dumas C**. **1987**. Mentor Pollen Techniques. In: Bourne GH, Jeon KW,
Friedlander M, eds. International Review of Cytology. Academic Press, 315–332.

**Koch JL, Carey DW, Knight KS, Poland T, Herms DA, Mason ME**. **2012**. Breeding
strategies for the development of emerald ash borer-resistant North American ash. In:
Proceedings of the 4th International Workshop on the Genetics of Host--Parasite Interactions
in Forestry: Disease and Insect Resistance in Forest Trees. USDA Forest Service, Pacific
Southwest Research Station Albany, California, 235–239.

**Koch J I., Carey DW, Mason ME**. **2007**. Development of novel ash hybrids to introgress
resistance to emerald ash borer into north American ash species. In: Mastro V, Lance D,
Reardon R, Parra G, eds. Emerald ash borer research and development meeting. U.S.
Department of Agriculture, Forest Service, Forest Health Technology Enterprise Team, 36–37.

**Koch JL, Carey DW, Mason ME, Nurul Islam-Faridi M**. **2010**. Overcoming obstacles to
interspecies hybridization of ash. In: Michler CH, Ginzel MD, eds. General Technical Report -
Proceedings. Proceedings of symposium on ash in North America; 2010 March 9-11; West
Lafayette, IN. Newtown Square, PA: U.S. Department of Agriculture, Forest Service, 41–44.

**Koch JL, Carey DW, Mason ME, Poland TM, Knight KS**. **2015**. Intraspecific variation in
Fraxinus pennsylvanica responses to emerald ash borer (Agrilus planipennis). *New Forests*
**46**: 995–1011.

**Kowalski T, Bilański P, Holdenrieder O**. **2015**. Virulence of Hymenoscyphus albidus and H.
fraxineus on Fraxinus excelsior and F. pennsylvanica. *PloS one* **10**: e0141592.

**Kozlov AM, Darriba D, Flouri T, Morel B, Stamatakis A**. **2019**. RAxML-NG: a fast, scalable

and user-friendly tool for maximum likelihood phylogenetic inference. *Bioinformatics* **35**:
4453–4455.

**Lévesque M, Bustamante Eduardo JI, Queloz V**. **2023**. Potential alternative tree species to
Fraxinus excelsior in European forests. *Frontiers in Forests and Global Change* **6**: 1048971.

**Lingelsheim A**. **1907**. Vorarbeiten zu einer Monographie der Gattung Fraxinus. *Botanische*
*Jahrbucher fur Systematik, Pflanzengeschichte und Pflanzengeographie* **40**: 185–223.

**Loureiro J, Rodriguez E, Dolezel J, Santos C**. **2007**. Two new nuclear isolation buffers for
plant DNA flow cytometry: a test with 37 species. *Annals of botany* **100**: 875–888.

**Marzano M, Woodcock P, Quine CP**. **2019**. Dealing with dieback: forest manager attitudes
towards developing resistant ash trees in the United Kingdom. *Forestry* **92**: 554–567.

**Miller GN**. **1955**. The genus Fraxinus, the Ashes, in North America, North of Mexico. *Cornell*
*University Agricultural Experimental Station Memoir* **355**: 1–64.

**Mitchell RJ, Beaton JK, Bellamy PE, Broome A, Chetcuti J, Eaton S, Ellis CJ, Gimona A,**
**Harmer R, Hester AJ,** *et al.* **2014**. Ash dieback in the UK: A review of the ecological and
conservation implications and potential management options. *Biological conservation* **175**:
95–109.

**Nielsen LR, McKinney LV, Hietala AM, Kjær ED**. **2016**. The susceptibility of Asian,
European and North American Fraxinus species to the ash dieback pathogen
Hymenoscyphus fraxineus reflects their phylogenetic history. *European journal of forest*
*research*: 1–15.

**Nikolaev EV**. **1981**. The genus Fraxinus (Oleaceae) in the flora of the USSR. *Botanicheskii*
*zhurnal* **66**: 1419–1432.

**Obermayer R, Leitch IJ, Hanson L, Bennett MD**. **2002**. Nuclear DNA C□values in 30
Species Double the Familial Representation in Pteridophytes. *Annals of botany* **90**: 209–217.

**Orlova-Bienkowskaja MJ, Bieńkowski AO**. **2016**. The life cycle of the emerald ash
borerAgrilus planipennisin European Russia and comparisons with its life cycles in Asia and
North America. *Agricultural and forest entomology* **18**: 182–188.

**Orlova-Bienkowskaja MJ, Bieńkowski AO**. **2018**. Modeling long-distance dispersal of
emerald ash borer in European Russia and prognosis of spread of this pest to neighboring
countries within next 5 years. *Ecology and evolution* **8**: 9295–9304.

**Pautasso M, Aas G, Queloz V, Holdenrieder O**. **2013**. European ash (Fraxinus excelsior)
dieback – A conservation biology challenge. *Biological conservation* **158**: 37–49.

**Plumb WJ, Coker TLR, Stocks JJ, Woodcock P, Quine CP, Nemesio□Gorriz M, Douglas**
**GC, Kelly LJ, Buggs RJA**. **2019**. The viability of a breeding programme for ash in the British
Isles in the face of ash dieback. *Plants People Planet* **58**: 227.

**Procter JB, Carstairs GM, Soares B, Mourão K, Ofoegbu TC, Barton D, Lui L, Menard A,**
**Sherstnev N, Roldan-Martinez D,** *et al.* **2021**. Alignment of Biological Sequences with
Jalview. *Methods in molecular biology* **2231**: 203–224.

**Raquin C, Jung-Muller B, Dufour J, Frascaria-Lacoste N**. **2002**. Rapid seedling obtaining
from European ash species Fraxinus excelsior (L.) and Fraxinus angustifolia (Vahl.). *Annals of*
*forest science* **59**: 219–224.

**Rigsby CM, Herms DA, Bonello P, Cipollini D**. **2016**. Higher activities of defense-associated
enzymes may contribute to greater resistance of Manchurian ash to Emerald Ash Borer than a
closely related and susceptible congener. *Journal of chemical ecology* **42**: 782–792.

**Santini A, Fagnani A, Ferrini F, Ghelardini L, Mittempergher L**. **2007**. 'Fiorente' and 'Arno'
Elm Trees. *HortScience: a publication of the American Society for Horticultural Science* **42**:
712–714.

**Sela I, Ashkenazy H, Katoh K, Pupko T**. **2015**. GUIDANCE2: accurate detection of
unreliable alignment regions accounting for the uncertainty of multiple parameters. *Nucleic*
*acids research* **43**: W7–14.

**Sharma DR, Kaur R, Kumar K**. **1996**. Embryo rescue in plants—a review. *Euphytica/*
*Netherlands journal of plant breeding* **89**: 325–337.

**Siljak-Yakovlev S, Temunović M, Robin O, Raquin C, Frascaria-Lacoste N**. **2014**.
Molecular–cytogenetic studies of ribosomal RNA genes and heterochromatin in three
European Fraxinus species. *Tree genetics & genomes* **10**: 231–239.

**Smalley EB, Guries RP**. **1993**. Breeding elms for resistance to Dutch elm disease. *Annual*
*review of phytopathology* **31**: 325–354.

**Sollars ESA, Harper AL, Kelly LJ, Sambles CM, Ramirez-Gonzalez RH, Swarbreck D,**
**Kaithakottil G, Cooper ED, Uauy C, Havlickova L, *et al.* 2017**. Genome sequence and
genetic diversity of European ash trees. *Nature* **541**: 212–216.

**Thomasset M, Fernandez-Manjarrés JF, Douglas GC, Frascaria-Lacoste N, Raquin C,**
**Hodkinson TR**. **2011**. Molecular and Morphological Characterization of Reciprocal F1 Hybrid
Ash (Fraxinus excelsior × Fraxinus angustifolia, Oleaceae) and Parental Species Reveals
Asymmetric Character Inheritance. *International journal of plant sciences* **172**: 423–433.

**Wallander E**. **2008**. Systematics of Fraxinus (Oleaceae) and evolution of dioecy. *Plant*
*systematics and evolution = Entwicklungsgeschichte und Systematik der Pflanzen* **273**: 25–
49.

**Wallander E**. **2012**. Systematics and floral evolution in Fraxinus (Oleaceae). *Belgische*
*Dendrologie Belge* **2013**: 39–58.

**Wenzig T**. **1883**. Die Gattung Fraxinus Tourn. neu bearbeitet. *Botanische Jahrbucher fur*
*Systematik, Pflanzengeschichte und Pflanzengeographie* **4**: 165–188.

**Wesmael A**. **1892**. Monographie des espèces du genre Fraxinus. *Bulletin de la Société*
*Royale de Botanique de Belgique / Bulletin van de Koninklijke Belgische Botanische*
*Vereniging* **31**: 69–117.

**Whittemore AT, Campbell JJN, Xia Z-L, Carlson CH, Atha D, Olsen RT**. **2018**. Ploidy
Variation in Fraxinus L. (Oleaceae) of Eastern North America: Genome Size Diversity and
Taxonomy in a Suddenly Endangered Genus. *International journal of plant sciences* **179**:
377–389.

**Zeng F-S, Li L-L, Liang N-S, Wang X, Li X, Zhan Y-G**. **2015**. Salt tolerance and alterations
in cytosine methylation in the interspecific hybrids of Fraxinus velutina and Fraxinus
mandshurica. *Euphytica/ Netherlands journal of plant breeding* **205**: 721–737.