Assessment of the Histone Mark-based Epigenomic Landscape in Human Myometrium at
Term Pregnancy

San-Pin Wu[1,*], Elvis Quiroz[1,*], Tianyuan Wang[2], Skylar Montague Redecke[1], Xin Xu[3], Lin
Lin[1,4], Matthew L. Anderson[5], Francesco J. DeMayo[1,#]

[1]Reproductive and Developmental Biology Laboratory, [2]Biostatistics & Computational Biology
Branch, and [3]Epigenomic and DNA Sequencing Core Laboratory, National Institute of
Environmental Health Sciences, National Institutes of Health, Research Triangle Park, North
Carolina. [4]School of Nursing, University of California San Francisco, San Francisco, California.
[5]Department of Obstetrics & Gynecology, University of South Florida Morsani College of
Medicine and Moffitt Cancer Center, Tampa, Florida.

*These authors contributed equally.

#Correspondence

ORCID:

San-Pin Wu: 0000-0001-9626-784X

Abstract

The myometrium plays a critical role during pregnancy as it is responsible for both the structural

integrity of the uterus and force generation at term. Emerging studies in mice indicate a dynamic

change of the myometrial epigenome and transcriptome during pregnancy to ready the

contractile machinery for parturition. However, the regulatory systems underlying myometrial

gene expression patterns throughout gestation remain largely unknown. Here we investigated

human term pregnant nonlabor myometrial biopsies for transcriptome, enhancer histone mark

cistrome, and chromatin conformation pattern mapping. More than thirty-thousand putative

enhancers with H3K27ac and H3K4me1 double positive marks were identified in the

myometrium. Enriched transcription factor binding motifs include known myometrial regulators

AP-1, STAT, NFkB, and PGR among others. Putative myometrial super enhancers are mostly

colocalized with progesterone receptor occupying sites and preferentially associated with highly

expressing genes, suggesting a conserved role of PGR in regulating the myometrial

transcriptome between species. In human myometrial specimens, inferred PGR activities are

positively correlated with phospholipase C like 2 (*PLCL2*) mRNA levels, supporting that PGR

may act through this genomic region to promote *PLCL2* expression. PGR overexpression

facilitated *PLCL2* gene expression in myometrial cells. Using CRISPR activation, we assessed

the functionality of a PGR putative enhancer 35-kilobases upstream of the contractile-restrictive

gene *PLCL2*. In summary, results of this study serve as a resource to study gene regulatory

mechanisms in the human myometrium at the term pregnancy stage for further advancing

women's health research.

Introduction

The myometrium is the muscular component of the uterus. Throughout gestation, the

myometrium maintains contractile quiescence until labor, where uterine contractility facilitates

parturition. Pre-term birth, often characterized by aberrant myometrial activity before term,

results in the termination of pregnancy prior to 37 weeks gestation. Complications associated

with pre-term birth account for 35% of neonatal death and contribute to lifelong physical and

socioeconomic multi-morbidities to surviving neonates and mothers (Chawanpaiboon, Vogel et

al. 2019). It is therefore critical to understand the molecular mechanisms underlying the

maintenance and shift of myometrial quiescence throughout gestation.

The myometrium undergoes extensive structural and functional remodeling in preparation for

parturition through genomic regulatory mechanisms which influence gene expression

throughout pregnancy. Major transcription factor families have been identified to contribute

throughout the remodeling process. Studies on the role of steroid hormone receptors in

myometrial remodeling suggest that the withdrawal of functional progesterone signaling at term

results from a stoichiometric shift favoring the PGR-A isoform over PGR-B. This shift is

associated with increased activation of estrogen receptor alpha (ESR1) expression at term

(Mesiano, Chan et al. 2002) (Merlino, Welsh et al. 2007) (Nadeem, Shynlova et al. 2016).

Activator protein 1 (AP-1) complex subunits have been observed to act as PGR coregulators

(Dai, Litman et al. 2003) and have dynamic expression patterns throughout gestation in both

humans and rodent models. For example, FOS:JUN heterodimers are implicated to be critical

for the initiation of labor through transcriptional regulation of gap junction proteins such as Gja1

(Gap junction alpha 1) (Mitchell and Lye 2001) (Mitchell and Lye 2005) (Balducci, Risek et al.

1993).

Contributing to the dynamic nature of the myometrial transcriptome during term is the

epigenome and its reprogramming. Studies investigating epigenetic markers related to gene

activation in the mouse myometrium have revealed that the promoters for contractility-driving

genes, such as Gja1, are epigenetically activated well before the onset of labor (Shchuka, Abatti

et al. 2020). Considering this, the myometrial epigenome's role in parturition disorders, such as

preterm birth, has thus been under investigation. For example, it is known that altered DNA

methylation is linked with preterm birth (Barcelona de Mendoza, Wright et al. 2017), including at

the promoters of genes associated with myometrial contraction (Erickson, Myatt et al. 2023) and

fetal membrane rupture (Wang, Ogawa et al. 2008). However, differential myometrial DNA

methylation at CpG islands in the promoters of contractility-driving genes is not thought to be a

major contributor to preterm birth (Mitsuya, Singh et al. 2014). Given that DNA methylation

mediated gene regulation often occurs outside of CpG islands (Irizarry, Ladd-Acosta et al. 2009),

there is still work to be done at this interface. Regardless, whether through epigenetic or transcriptomic

gene regulatory programs, the molecular mechanisms underlying the regulation of genes critical for

myometrial reprogramming and quiescent maintenance are still poorly understood.

In this study, we aim to improve our understanding of the cellular processes involved in shaping

the myometrium's dynamic state of quiescence at pregnancy through the epigenetic and

transcriptomic profiling of the human myometrium. Because of the relevance of genomic

regulatory mechanisms in coordinating myometrial activity, we identified candidate cis-acting

regulatory regions using chromatin immunoprecipitation sequencing (ChIP-Seq) assays of

surrogate histone enhancer markers H3K27ac (Creyghton, Cheng et al. 2010) and H3K4me1

(Heintzman, Stuart et al. 2007) alongside chromatin conformation capture assays in term

pregnant myometrial tissues. Using this data, we investigated cis-acting regulatory regions for

the gene phospholipase C like 2, which encodes for the protein PLCL2 and has been implicated

in the modulation of calcium signaling (Uji, Matsuda et al. 2002) and maintenance of myometrial

quiescence (Peavey, Wu et al. 2021). Putative enhancers upstream of the PLCL2 transcriptional

start site were subjected to functional assessment using CRISPR activation-based assays.

Here, we identified a genomic region 35-kilobases upstream of the *PLCL2* transcriptional start

site involved in *PLCL2* transcriptional regulation. Furthermore, we aimed to characterize this

genomic region through the identification of candidate *PLCL2* transcriptional regulators co-

localizing with this cis-acting element using integrative cistrome and transcriptome analysis and

have demonstrated PGR to be a direct regulator for *PLCL2* expression. These findings build

upon our understanding of myometrial remodeling throughout gestation and will be pertinent for

the development of medical interventions aiming to address pre-term birth.

Results

<u>Epigenomic Landscape in Term Pregnant Myometrial Specimens</u>

To better understand the regulatory network shaping the myometrial transcriptome before labor,

we analyzed transcriptome and putative enhancers in individual human myometrial specimens.

Using RNA-seq, we identified actively expressed RNAs, while ChIP-seq for H3K27ac and

H3K4me1 was used to map putative enhancers. Active genes were associated with nearby

putative enhancers based on their genomic proximity. Additionally, chromatin looping patterns

were mapped using Hi-C to further link active genes and putative enhancers within the same

chromatin loops. The ChIP-seq assay identified an average of 44238 H3K27ac and 74325

H3K4me1 positive genomic regions in three term-pregnant, nonlabor myometrial biopsies

(Table 1). An RNA-Seq assay revealed an average of 12157 active genes in these specimens

that manifested expression levels of fragments per kilobase of transcript per million mapped

reads (FPKM) greater or equal to 1 (Table 1). A High-throughput Chromosome Conformation

Capture (Hi-C) assay further found a total of 27162 chromatin loops from two of the three

myometrial specimens (Table 1, subject 1 and 3). We failed to identify chromatin loops in the

second subject's biopsy due to limited sample availability. Together, these results delineate a

map of H3K27ac and H3K4me1 positive signals in the human term pregnant myometrial tissue

before the onset of labor, which we use as a resource to investigate the molecular mechanisms

that prepare the myometrium for subsequent parturition.

When comparing the present study to previous findings (Dotts, Reiman et al. 2023), 8563

genomic regions carry common H3K27ac-positive histone marks. However, significant

variations on the number and location of H3K27ac-positive intervals are present across the six

samples among these two studies (Figure S1). Depending on the number of mapped intervals,

the 8563 common regions constitute between 20.7% to 71.5% of total H3K27ac-positive

intervals across the six myometrial specimens (Figure S1). Notably, the fraction of intervals

commonly identified in the present study's three specimens ranges between 56.4% and 70.0%

of each of their total H3K27ac-positive regions, while the Dotts dataset has a wider range

between 28.0% and 74.0%. With a less stringent criterium, most the H3K27ac-positive intervals

are found in at least two samples, ranging between 75.4% and 98.3% (Figure S1). These data

together highlight the degree of variation on mapping the epigenome among specimens and

datasets.

<u>Putative Enhancers for Gene Regulation in the Human Myometrium</u>

Since H3K27ac and H3K4me1 marked genome regions are associated with enhancers for gene

regulation (Creyghton, Cheng et al. 2010) (Heintzman, Stuart et al. 2007), we define the regions

that have overlapping H3K27ac and H3K4me1 marks as putative myometrial enhancers at the

term pregnant nonlabor stage (Dataset S1). Among the three specimens, 13114 putative

enhancers are commonly present (Figure 1A). A significant variance on the location of

myometrial putative enhancers is also seen among biopsies as evident by the observation that

these common putative enhancers make up less than half of the total putative enhancers in

each individual specimen (Figure 1A). More than one-third of these 13114 common putative

enhancers are located within a 100-kilobase vicinity of actively high-expressing genes (FPKM

>= 15), while a much smaller fraction of them (13.3%) are associated with actively low-

expressing genes (Figure 1B). Motif enrichment analysis on these common putative enhancers

further reveals an overrepresentation of binding motifs for myometrial transcription factors AP-1,

STAT5, and NFκB, steroid hormone receptors PGR and NR3C1, and smooth muscle

transcription regulators SRF and ELK1 (Figure 1C and Dataset S2). These findings collectively

suggest that the putative myometrial enhancers bring together smooth muscle and hormonal

control programs for the regulation of myometrial gene expression at term pregnancy.

Super enhancers house hormone-dependent gene regulatory programs for female reproductive

tract homeostasis (Shin, Willi et al. 2016) (Hewitt, Grimm et al. 2020). Across the three human

subjects, 540 putative super enhancers are commonly identified in the term pregnant nonlabor

myometrial specimens (Figure 2A and Dataset S3). More than 40% of the 540 putative super

enhancers are located within a 100-kilobase distance to high-expressing genes (FPKM >= 15),

while only 7.3% of putative myometrial super enhancers are found near low-expressing genes

(5 > FPKM >=1) (Figure 2B). Compared with the regular putative enhancers, the putative

myometrial super enhancers are found more frequently near active genes that are expressed at

relatively higher levels (Figure 1B and Figure 2B). Notably, 76% of the putative super-enhancers

co-localize with known PGR-occupied regions in human myometrial tissue, compared to 20%

co-localization observed in regular enhancers (Figure S2). Further examining the myometrial

active genes that are associated with putative super enhancers revealed an enrichment of gene

functions in cytoskeleton organization, extracellular-receptor interaction, transcription regulation,

mRNA stability, and estrogen signaling (Figure 2C and Dataset S4). Taken together, these

results establish the association among the putative myometrial enhancers, the potential

regulatory program within these super enhancers, and the cellular functions of the genes they

may control.

Cis-acting elements for the control of the contractile gene *PLCL2*

We previously demonstrated the positive correlation of PLCL2 and PGR expression in a mouse

model and PLCL2's function on negatively modulating oxytocin-induced myometrial cell

contraction (Peavy et al., 2021). However, the mechanism that underlies PGR's regulation of

PLCL2 remains unclear. Taking advantage of the mapped myometrial cis-acting elements, we

aimed to identify the cis-acting elements that may contribute to the PLCL2 transcriptional

regulation with a special interest on the PGR-related enhancers.

The Hi-C assay identified a chromatin loop with one end at the *PLCL2* transcription start site

(TSS) and the other end at approximately 90-kilobases upstream of the TSS (Figure 3A). Within

this chromatin loop, ChIP-Seq results revealed seven genomic regions that are marked with

H3K27ac in all three myometrial specimens (Figure 3A). Six of the seven regions are also co-

localized with previously published genome occupancy of transcription regulators curated by the

ReMap Atlas (Hammal, de Langen et al. 2022) (Figure 3A). For the purpose of functional

screening, we focus on H3K27ac signals instead of using H3K27ac/H3K4me1 double positive

criterium to cast a wider net. Eight guide RNAs (gRNAs) were designed to target these seven

candidate regions for CRISPR activation (CRISPRa) based screening of *PLCL2* gene regulation

(Figure 3A). The two previously reported gRNAs (PLCL2-7 and PLCL2-8) were able to elevate

the endogenous *PLCL2* mRNA levels significantly higher than the non-targeting gRNA in

immortalized human myometrial cells (hTERT-HM) (Figure 3B) (Peavey, Wu et al. 2021).

gRNAs PLCLC2-1, PLCL2-3, PLCL2-5, and PLCL2-6 also significantly increased *PLCL2*

expression above the levels of the non-targeting group (Figure 3B). Moreover, results from the

Perturb-Seq assay found that the PLCL2 gene ranks high in the differentially expressed gene

list of the PLCL2-5 gRNA expressing cells compared with the non-targeting gRNA expressing

cells (Dataset S4), in line with the qRT-PCR assay finding in which the gRNA PLCL2-5 induced

PLCL2 expression to the greatest extent among the upstream non-coding genome targeting

gRNA (Figure 3B). Moreover, gRNA PLCL2-5 was capable of mediating the increase of the

PLCL2 transcript abundance in another uterine mesenchymal cell type T-HESC (Figure 3C).

Collectively, these findings support the PLCL2-5 targeted genomic region as a cis-acting

element for regulation of the *PLCL2* gene.

<u>PGR as an upstream regulator for the *PLCL2* gene in the human myometrium</u>

After determining its cis-acting element role, the PLCL2-5 targeted genomic region was further

examined to identify upstream regulators that control myometrial *PLCL2* expression. This was

achieved by first identifying transcription factor occupancy in this genome region in any

tissues/cells that are documented in public databases (ReMAP Atlas) (Hammal, de Langen et

al. 2022), followed by filtering for factors that have known mRNA and/or protein expression in

hTERT-HM cells and tissues using publicly available transcriptomic and histological data

including the Human Protein Atlas (Human Protein Atlas proteinatlas.org) (Uhlen, Fagerberg et

al. 2015) and published RNA-Seq datasets (Stanfield, Amini et al. 2019) (Wu, Anderson et al.

2020). The resulting candidate upstream regulators for the myometrial *PLCL2* gene are

documented in Table 2. Since the *PLCL2* mRNA abundance is much lower in the hTERT-HM cells than in the myometrial tissues, these candidate upstream regulators were further grouped into candidate activators and repressors based on an assumption that a candidate activator would be expressed higher in human myometrial tissue, and a candidate repressor being expressed higher instead in the hTERT-HM cells, holding the *PLCL2* mRNA levels down. Candidate activators for the *PLCL2* gene include known myometrial regulators such as PGR, estrogen receptor alpha (ESR1), and AP-1 (Table 2). Interestingly, the mediator complex protein subunit MED1, cohesion complex members RAD21 and SMC3, and many epigenomic modifiers are among the list of the candidate repressors (Table 2). These findings not only provide candidates for subsequent functional assessment, but also highlight potential pathways for future investigations on the regulation of the contractile gene *PLCL2* expression in the myometrium.

We previously demonstrated the regulation of mouse *Plcl2* gene by the myometrial PGR (Wu, Wang et al. 2022). To further test whether such a regulatory mechanism is also present in the human, the CRISPRa technology was employed to increase the expression of PGR mRNA and proteins in hTERT-HM cells (Figure 4A and Figure 4B). Given that the PGR640 gRNA led to higher PGR protein production (Figure 4B), this gRNA was utilized to stimulate myometrial *PGR* expression for subsequent experiments. Indeed, RT-qPCR results showed that PGR overexpression increased *PLCL2* mRNA abundance in hTERT-HM cells (Figure 4C), demonstrating a consistent finding with the mouse model.

To further explore this regulatory relationship in the human myometrial tissue, we examined the degree of correlation between the inferred PGR activity and the *PLCL2* transcript abundance in myometrial biopsies. Gene expression profiles of myometrial specimens from 13 proliferative-phase, 6 secretory-phase, 14 post-menopausal, and 10 Provera-treated human subjects were determined by the RNA-Seq assay (Dataset S6) to take advantage of the wide spectrum of

progesterone signaling activities in these samples. Inferred myometrial PGR activities,

represented as T-scores (Dataset S6), were derived from this normalized gene expression

matrix of 43 human myometrial specimens with the previously published mouse myometrial *Pgr*

gene signature (Wu, Wang et al. 2022) (Li, Bushel et al. 2021). A Pearson correlation analysis

found a positive correlation (r = 0.47) between the T-scores and normalized *PLCL2* mRNA

levels in these 43 specimens (Figure 4D). Together, these in vitro and in vivo findings

collectively support that PGR is a *PLCL2* activator, likely acting through the 35-kb upstream cis-

acting element.

Discussion

Temporally and spatially regulated gene expression is the foundation of tissue phenotype

manifestation. Shchuka and colleagues utilize the mouse model to show that the myometrial

epigenome is readily programed for parturition-associated gene induction days ahead of the

term pregnancy (Shchuka, Abatti et al. 2020), laying the ground for subsequent laboring event.

We chose to study the human myometrium at the term pregnant nonlabor stage in order to

examine the association between the epigenome landscape and the gene expression pattern

not only for preparing the myometrium for parturition, but also for maintaining myometrial

quiescence. We also performed a mullti-omic study on a myometrial specimen from each

individual human subject to better align all the datasets together, reflecting specimen and

subject individuality. While the variation among specimens is substantial, we were able to

identify myometrial cis-acting elements commonly shared among three individuals. As a proof of

principle, we subsequently used this information to functionally screen for genomic regions that

may mediate the regulation of the contractile-suppressing gene *PLCL2*. The H3K27ac marked

35-kilobase upstream region as a result of the screening then serves as a bridge to identify

PGR as one of the upstream regulators for modulating *PLCL2* gene expression in human

myometrium. These findings showcase the value of these datasets as a resource for future

investigations on the mechanisms of myometrial gene regulation.

Variations across cistromic datasets for the human myometrium are significant (Figure 1 and

Figure S1) (Dotts, Reiman et al. 2023). Contributing factors likely include but are not limited to

subject-to-subject differences and batch effects from sample preparations. Taking the H3K27ac

marked genomic regions as an example, regions commonly identified in the specimens of this

study's three subjects account for 55.9%, 67.7%, and 57.9% of total H3K27ac-positive intervals

in each individual sample, despite our best attempt to reduce the batch effect. Heterogeneity on

cell type compositions, likely due to the sampling location, could be a major contributing factor

to the variation. It is also speculated that the genomic regions with subject-to-subject variances

could be transiently present and thus the signal could not be captured across samples. In

addition, the underlying health conditions, medications, and environmental challenges among

others may also affect the epigenomic profile. Variations could also arise from differences in the

detection threshold. We identified an average of 44238 H3K27ac marked regions per specimen,

while Dotts and colleagues found nearly 19000 per sample (Dotts, Reiman et al. 2023). Reagent

and sample handling could confound with biological variances of specimens leading to this

difference. Future investigations using standardized sample preparation protocols at the single

cell resolution may help to reduce this variation.

Results from the CRISPRa-based Perturb-Seq assay demonstrate the capability of this

technology to screen for cis-acting elements that are in topological vicinity of the gene of

interest. This assay is particularly useful on genes that are expressed at low levels in the

screening platform, i.e., the cells. The dCas9VPR activator can generate satisfactory signal-to-

noise ratios to be detected by single cell RNAseq. The advantage of assaying with this system

is that each individual cell serves as a container to generate data points instead of relying on

multiple wells on cell culture plates. Multiplexing with multiple gRNAs can be achieved by using

the gRNA sequence as a unique barcode of the cell, enabling the simultaneous collection of

numerous data points for each individual gRNA. In the present study, 56.7% of tested cells

carried one species of detectable gRNAs, permitting the study of the resulting effect under a

single gRNA. Moreover, another 27.0% of assayed cells have more than one species of

detectable gRNAs, which open the possibility to study interactions between different cis-acting

elements. However, we failed to use this approach for the purpose of identifying target genes of

a putative enhancer of interest (Dataset S4). Challenges such as difficulties on defining the

criteria of the associated genes and unsatisfactory signal-to-noise ratios on genes already at

high baseline levels should be considered in future experimental designs. Taken together,

results of the present study support the use of the CRISPRa-based Pertub-Seq assay under a

defined scope.

PLCL2 and its paralog PLCL1 are catalytically inactive members of the phospholipase C family

functioning as negative regulators of calcium signaling (Kanematsu, Takeuchi et al. 2005)

(Takenaka, Fukami et al. 2003) (Uji, Matsuda et al. 2002). Both of them are expressed in human

myometrial tissues and are able to attenuate oxytocin-induced muscle cell contraction (Peavey,

Wu et al. 2021). *PLCL1* mRNA abundance is higher in the myometrium at term pregnancy than

the non-gravid stage, while *PLCL2* transcript levels remain comparable between the two stages

(Wu, Anderson et al. 2020). Data from cultured human endometrial stromal cells and the

genetically engineered mouse model reveal progesterone signaling as the upstream regulator of

PLCL1 and *PLCL2*, respectively (Muter, Brighton et al. 2016) (Peavey, Wu et al. 2021). The

present study further demonstrates the PGR-dependent regulation of the *PLCL2* gene in the

human myometrial cells, suggesting that this pathway is conserved between human and mouse.

Importantly, the identification of the cis-acting element 35-kilobase upstream of the *PLCL2* gene

opens an avenue to investigate the impact of interactions between PGR and other myometrial

transcription regulators on the mechanism of action that controls the myometrial contractile

machinery. The future findings in this space could provide insight on progesterone signaling

modification and identify progesterone signaling modifiers for the development of novel therapy

that targets parturition disorders.

Materials and Methods

Collection of myometrial specimens

Permission to collect human tissue specimens was prospectively obtained from individuals

undergoing hysterectomy or cesarean section for benign clinical indications (H-33461). Gravid

myometrial tissue was obtained from the margin of the hysterotomy in women undergoing term

cesarean sections (>38 weeks estimated gestational age) without evidence of labor. Non-gravid

myometrial tissue was collected from pre-menopausal women undergoing hysterectomy for

benign conditions. Specimens from gravid women receiving treatment for pre-eclampsia,

eclampsia, pregnancy-related hypertension, or pre-term labor were excluded.

Human Cell Lines

Human immortalized myometrial cells (hTERT-HM) (Condon, Yin et al. 2002) cells and human

immortalized endometrial cells (T-hESC) (ATCC, CRL-4003) were cultured in DMEM/F-12

(Invitrogen, Grand Island, NY, USA) supplemented with 10% Fetal Bovine Serum (FBS, Gibco)

and antibiotics (10 000 IU/mL penicillin, 10 000 IU/ mL streptomycin; Life Technologies, Grand

Island, NY). Cell culture media was filtered using the 0.22um Rapid-Flow™ Sterile Disposable

Filter Units (Nalgene). Cells were incubated at 5% CO_2 and 37°C.

RNA Isolation and RT-qPCR

RNA was isolated from cells using TRIzol Reagent (Invitrogen) and the RNeasy mini RNA

isolation kit (Qiagen, Hilden, Germany). cDNA was prepared using Moloney Murine Leukemia

Virus reverse transcriptase (Thermo Fisher Scientific) with Random Hexamers (Invitrogen,

Waltham, MA, USA) according to manufacturer protocol. For quantitative analysis of mRNA,

SsoAdvanced Universal SYBR Green Supermix (BioRad, Hercules, CA, USA) was used

according to manufacturer instructions. Each reaction was performed in technical duplicates

using the standard curve-based method (Larionov, Krause and Miller 2005). Relative expression

of genes of interest were normalized to the 18S rRNA. Briefly, reaction samples were prepared

to a volume of 20ul with 5uM of both forward and reverse primers, cDNA, and a final 1x

concentration of the SYBR Green Supermix. The reaction was heated to 95°C for 30 seconds,

followed by 39 cycles of denaturation at 95°C for 15 seconds and annealing and elongation at

60°C for 30 seconds. Temperature cycles were performed on the CFX Connect^{TM} Real-Time

PCR Detection System (Bio-Rad).

Western Blot Assay

Protein was isolated from cells using the RIPA Lysis and Extraction Buffer (Thermo Scientific)

with the following specifications: approximately 300,000 cells were pelleted and lysed with 100ul

of complete Pierce RIPA buffer. Protein was quantified using the BCA Protein Assay Kit

(Thermo Scientific, Waltham, MA, USA) as instructed by the manufacturer, and 40ug of protein

were loaded per lane on a Mini-PROTEAN TGX Pre-cast Protein gel (Bio-Rad, #4568094) with

the Precision Plus Protein Dual Color Standards Ladder (Bio-Rad #1610374). Protein lysates

from MCF7 cells were used as positive controls, while protein lysates from HEK293T cells were

used as negative controls. Gels were transferred to nitrocellulose membrane using the Turbo-

Blot transfer system (BioRad) according to the manufacturer's instructions. The membrane was

blocked with 5% milk (Santa Cruz Biotech, Santa Cruz, CA, USA) in TBST (20 mM Tris, pH 7.4

(Lonza, Morrisville, NC, USA), 140 mM NaCl (Lonza), 1% TWEEN-20 (Sigma). PGR protein

was detected using Monoclonal Mouse Anti-Human Progesterone receptor clone PgR 1294

diluted 1:1000 in milk overnight at 4C. Bands were detected using Donkey Anti-Mouse 926-

32212 (Lot # C60524-15) diluted 1:20,000 in milk for 45 minutes.

Blots were imaged using Odyssey Fc Imager (LI-COR Biosciences) using 800nm channel for 10

minutes, and 700nm channel for 30 seconds (to image ladder). The control protein (B-actin) was

detected similarly as above using the following antibodies: Actin (I-19)-R sc-11616-R

(Lot#DO406) Rabbit polyclonal IgG Santa Cruz (1:1000 dilution) and Goat anti-rabbit 926-32211

(Lot#DOO304-15) (1:20,000 dilution).

RNA-Seq Library Preparation

RNA-Seq libraries were prepared according to the Illumina TruSeq Stranded mRNA protocol

Document # 1000000040498 v00. Libraries were sequenced using the Illumina NextSeq 500

and NovaSeq 6000 systems. The summary table is available in Dataset S7.

ChIP-Seq Library Preparation

The ChIP-Seq assay was performed by the Active Motif service laboratory using snap-freeze

human myometrial specimens. The H3K27Ac ChIP reactions were conducted with 10 µg of

tissue chromatin and 4 µg of H3K27Ac antibody (Active Motif, cat # 39133). The H3K4me1 ChIP

reactions were carried out by using 10 ug of tissue chromatin and 4 ul of H3K4me1 antibody

(Active Motif, cat # 39297). Libraries were prepared by a custom Illumina library with the

standard Illumina PE adaptors (Bentley, Balasubramanian et al. 2008). Libraries were

sequenced using the Illumina NextSeq 500 and NovaSeq 6000 systems.

Hi-C Library Preparation

Snap-freeze human myometrial specimens were shipped to the Arima Genomics and the Active

Motif service laboratories for preparation of the Hi-C library using the Arima-HiC Kit (Arima

Genomics A510008), protocol version A160132 v00. Libraries were sequenced in the NIEHS on

the Illumina NovaSeq 6000 platform.

CRISPR Activation (CRISPRa) assay

Acquisition of guide RNA expression vectors for CRISPRa assay

All gRNA expression vectors were synthesized by and acquired from VectorBuilder. gRNA

expressing vectors for PGR targeting (PGR640A and P6SM), non-targeting control, and the

empty backbone were purchased from VectorBuilder under the catalog numbers VB191117-

1498rhp (PGR640), VB210824-1268wmh (P6SM), VB191117-1500trj, and VB190918-1522gwq,

respectively.

Acquisition of dCas9-VPR-mCherry for CRISPRa assay

IGI-P0492 pHR-dCas9-NLS-VPR-mCherry was a gift from Jacob Corn (Addgene plasmid #

102245 ; http://n2t.net/addgene:102245 ; RRID:Addgene_102245).

Production of Lentiviruses

All lentiviruses were packaged in HEK293T/17 cells (ATCC # CRL-11268) according to

published method in Current Protocols in Neuroscience (Chen, Haam et al. 2019). Briefly, 293T

cells were transiently transfected with pMD2G, psPAX2 and a transfer vector containing the

desired gene using Lipofectamine 2000. Supernatant was collected 48 hours post transfection

and concentrated by centrifugation at 50,000 g for 2 hours. Pellets were resuspended in PBS

and used for infection. To determine titer, HEK293T/17 cells were infected with lentiviral

samples. Five days post infection, Qiagen DNeasy Blood & Tissue Kits was used to isolate

chromosomal DNA from infected cells. All titers were determined by performing droplet digital

PCR (ddPCR) to measure the number of lentiviral particles that integrated into the host genome.

CRISPRa viral transduction

hTERT-HM cells were infected with lentiviral gRNA expression vectors with a multiplicity of

infection of 4 (MOI = 4) and with dCas9-VPR-mCherry at an MOI = 4. gRNA expression vector

and dCas9-VPR-mCherry expression vectors, following lentiviral transduction, confer GFP and

mCherry fluorescent markers, respectively.

Fluorescence Activated Cell Sorting for CRISPRa assay

Cells were examined using a BD FACSAriaII cell sorter (Becton Dickinson Biosciences, San

Jose, CA) equipped with FACSDiVa software. Initially, a "scatter" gate was set on a forward

scatter (FSC-A) versus side scatter (SSC-A) dot plot to isolate the principal population of cells

free of debris. Subsequently, cells were consecutively gated on a side scatter height (SSC-H)

versus width (SSC-W), then a forward scatter height (FSC-H) versus width (FSC-W) dot plot to

isolate single cells. GFP expressing cells were excited off a 488 nm laser and detected using a

525/50 nm filter. mCherry expressing cells were excited off a 561 nm laser and detected using

a 610/20 nm filter. Double positive cells were collected based off non-transfected controls and

used for further cultural/biochemical analysis.

Perturb-Seq library preparation

The cells in suspension were counted and examined for viability with trypan blue staining using a TC-20 cell counter (Bio-Rad). Approximately 16,500 live cells at 1×10^6 cells/ml concentration with 65% or above viability were loaded into the Single Cell Chip to generate single cell emulsion in Chromium Controller with Chromium Single Cell 3' Library & Gel Bead Kit v3.1 (10x Genomics, Cat. 1000268). Reverse transcription of mRNA and cDNA amplification were carried out following the manufacture's instruction (10x Genomics, Cat. 1000268, Cat. 1000262 with 10x Genomics protocol CG000316). The amplified cDNA was separated into CRISPR sgRNA derived cDNA and transcriptome derived cDNA. The CRISPR sgRNA derived cDNA was used to make NGS sequencing libraries. The transcriptome derived cDNA was further fragmented to construct NGS libraries. Both libraries were then sequenced together with the molar ratio of 1 to 4 by the NIEHS Epigenomics and DNA Sequencing Core Laboratory with the parameters recommended in the manufacture's instruction.

Human enhancer perturb-seq data analysis

The raw sequencing FASTQ files generated from both the transcriptome and CRISPR screening libraries were processed together by Cell Ranger software (version 4.0.0, 10X Genomics). The "cellranger count" pipeline used STAR for aligning the reads to the human reference, GRCh38 "refdata-gex-GRCh38-2020-A" (10X Genomics), and associated gene expression profile with guide RNA (gRNA) identity by unique barcode in each cell. Seurat software (version 3.6.3) was utilized to perform clustering analysis on the combined dataset (Satija, Farrell et al. 2015). We applied the SCTransform package to normalize gene expression counts across cells (Hafemeister and Satija 2019). The cells were clustered based on number of

unique gRNA type detected. The cell populations containing more than one type of gRNA were

excluded from further analyses. We applied two approaches to determine the association

between gRNAs and target genes:

A. For each cell cluster having gRNA with known target genes, such as PLCL1 and PLCL2, we

compared the expression of target genes in these clusters to the cluster with scramble control

gRNA. We defined individual gRNA activation as a fold change greater than 1.5.

B. For cell clusters having gRNAs with unknown target genes, we identified differentially

expressed genes by performing pair-wise comparisons between each cluster and the cluster

with scramble control gRNA. This analysis was conducted using the "FindAllMarkers" function in

the Seurat package. Target genes were defined as those with a fold change greater than 1.5

and an adjusted p-value less than 0.05.

Additional perturb-seq data analysis is provided in Dataset S9, with a summary table available

in Dataset S7

ChIP-Seq

The raw ChIP-seq reads (51-bp, single-end) were filtered with average quality scores greater

than 20. Adaptor sequences were trimmed from reads using cutadapt (v1.12). Then the reads

were aligned to the human reference genome (hg38) using Bowtie (v1.1.2) (Langmead, Trapnell

et al. 2009), requiring unique mapping and allowing up to 2 mismatches per read (-m 1 -v 2).

Duplicated reads with identical sequences were removed using Picard tools. To visualize the

read coverage, BigWig files were generated from the bedgraph files of each sample using

bedGraphToBigWig. These bigWig files were displayed as custom tracks on the UCSC genome

browser.

The normalization of sequencing depth across all ChIP-seq datasets were achieved by down

sampling to 20 million uniquely mapped reads per sample. The initial peaks of each sample

were identified using MACS2 with a cutoff of adjusted p-value < 0.0001 (Zhang, Liu et al. 2008).

The gene associated with each peak was predicted by searching the transcription start site

(TSS) of nearby genes within a 100Kb range using HOMER (Heinz, Benner et al. 2010). The

summary table is available in Dataset 7.

Identification of union peaks between H3K27ac and H3K4me1 peaks

The initial union peaks between H3K27ac and H3K4me1 in each sample were identified using

the "merge" function of BedTools. The common union peaks from multiple samples were used

for downstream analysis.

Identification of union peaks for term pregnant myometrial specimens

Additional H3K27ac ChIP-seq and related input data were obtained from GEO (GSE202027;

patient 1, 4, and 7). The same ChIP-seq analysis pipeline described above was applied to those

samples. To obtain a comprehensive set of peaks across all H3K27ac ChIP-seq datasets, the

peak intervals from each dataset were merged using BedTools.

Identification of super enhancers

H3K27ac-positive enhancers were defined as regions of H3K27ac ChIP-seq peaks in each

sample. The enhancers within 12.5Kb were merged by using bedtools merge function with

parameter "-d 12500". The combined enhancer regions were called super enhancers if they

were larger than 15Kb. The common super enhancers from multiple samples were used for

downstream analysis (Whyte, Orlando et al. 2013).

Colocalization of super enhancers and PGR genome occupancy was compared by calling

peaks from previously published PGR ChIP-seq data (GSM4081683 and GSM4081684). The

percentages of enhancers and super enhancers that manifest PGR occupancy were calculated

by overlapping the genomic regions in each category with PGR occupancy regions.

RNA-Seq analysis of Term Myometrial Specimens

The raw paired-end reads (51-bp, paired-end) were initially processed by applying a filter with

an average quality score of 20. Adaptor sequences were subsequently removed from reads

using cutadapt version 1.12. The processed reads were then aligned to the human reference

genome (hg38) using the STAR aligner version 2.5.2b. The gene expression levels in each

sample were determined by counting the total number of paired-end reads mapped to each

gene using DESeq2 R package version 1.12.4 (Love, Huber and Anders 2014). The gene

expression count matrix was normalized based on ratio of total mapped read pairs in each

sample to 56.5 million. Only the genes with average normalized count across all samples larger

than one were used for further analysis. The Bioconductor package edgeR was applied to the

gene expression count matrix to detect differentially expressed genes between groups of

interest (Robinson, McCarthy and Smyth 2010). The false discovery rate (FDR) of differentially

expressed genes was estimated using the Benjamini and Hochberg method.

RNA-Seq analysis of Provera Treated Human Myometrial Specimens

The raw paired-end reads (76-bp, paired-end) were processed by applying a filter with an

average quality score of 20. Adaptor sequences were removed from reads using cutadapt

(v1.12). The processed reads were then aligned to the human reference genome (hg38) using

the STAR aligner (v2.5.2b). Fragments Per Kilobase Million (FPKM) was calculated for each

gene in each sample by using Cufflinks version 2.0.2.

Hi-C

The raw reads (51-bp, paired-end) generated from HiC library were mapped to human reference

genome (hg38; Genome Reference Consortium Human GRCh38 from December 2013) using

HiCUP version 0.7.1 (Wingett, Ewels et al. 2015). The uniquely mapped di-tag passing quality

filtering with distance larger than 10Kb were used for downstream analysis. Chromatin loops

were identified using "hiccups" function of Juicer version 1.8.9 with default parameters (Durand,

Shamim et al. 2016). The A/B compartments were predicted at 100Kb resolution using

"eigenvector" function of Juicer. The Hi-C quality control data are provided in Dataset S8, and

the summary table is available in Dataset 7.

DNA-binding factor motif enrichment analysis

Enriched motifs were identified by HOMER (Hypergeometric Optimization of Motif EnRichment)

v4.11 with default background sequences matching the input sequences (Heinz, Benner et al.

2010).

Inferred myometrial PGR activities and the correlation analysis

The inferred PGR activities were represented by the T-score, which was derived by inputting the

mouse myometrial *Pgr* gene signature, based on the differentially expressed genes between

control and myometrial *Pgr* knockout groups at mid-pregnancy (Wu, Wang et al., 2022), into the

SEMIPs application (Li, Bushel et al., 2021). The T-scores were computed using this signature

alongside the normalized gene expression counts (FPKM) from 43 human myometrial biopsy

specimens.

Data Availability

All the Nest Generation Sequencing Data were deposited to the National Center for

Biotechnology Information Gene Expression Omnibus under the accession number GSE244735

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE244735).

Author Contribution

SPW, LL, and FJD conceived the study. MLA provided crucial reagent. SPW, EQ, SMR, and XX

performed experiments. SPW, TW, EQ, and SMR conducted data analyses. SPW, TW, EQ, XX,

MLA, and SMR wrote the initial draft.

Acknowledgement

We thank the following National Institute of Environmental Health Sciences (NIEHS) Core

facilities for exceptional technical supports: Epigenomic and DNA Sequencing Core, Viral Vector

Core, Flowcytometry Core, and Integrative Bioinformatics Supportive Group. We thank Ms.

Olivia Emery for her technical assistance. This study is partly supported by the Intramural

Research Program of the National Institute of Environmental Health Sciences Z1AES103311

560 **Table**

561 **Table 1.** Epigenome and transcriptome profile of the individual myometrium biopsy. Active

562 genes are defined as FPKM ≥ 1. N.D., not determined.

Subject	1	2	3
Number of H3K27ac intervals	47,223	39,465	46,026
Number of H3K4me1 intervals	72,091	74,511	76,374
Number of active genes	12,809	11,761	11,902
Number of chromatin loops	10,321	N.D.	16,841

565   **Table 2.** Candidate upstream regulators that mediate PLCL2-5 enhancer's regulatory effect on

566   PLCL2 expression.

Candidate Activators	Description
PGR	Steroid Hormone Receptor
ESR1	
FOS, JUN	AP-1 Transcription Factor Subunit, bZIP Transcription Factor
JUN	
JUNB	
JUND	
MAF	bZIP Transcription Factor
BHLHE40	bHLH Transcription Factor, Involved in CLOCK gene regulation
TCF21	bHLH Transcription Factor
MAX	bHLHZ Transcription Factor
FOXO1	Forkhead Box Transcription Factor
MRTFB	Myocardin Family, SRF Trancriptional Co-Activator
STAT3	Transcriptional Regulator
ERG	ETS Transcriptional Regulator

Candidate Repressors	Description
ARID1A	SWI/SNF Family of Epigenetic Modifiers
BRD2	Epigenetic Modifier, Chromatin Reader
BRD4	
KMT2A	Component of MLL Epigenetic Modification complex
SMARCB1	Component of BAF Epigenetic Modification Complex
ZMYM3	Component of Epigenetic Modification Complexes
CREB1	bZIP Transcription Factor, Interacts with Epigenetic Modifiers
CREBBP	CREB Binding Protein, Epigenetic Modifier
HDAC2	Histone Deacetylase, Epigenetic Modifier
HDAC3	
CLOCK	bHLH Transcription Factor Family, Rhythmic Epigenetic Modifier
MED1	Mediator Complex Subunit
RELB	NF-KB Transcription Factor Subunit
CHD4	Component of NuRD Epigenetic Modification Complex
RAD21	Cohesin Complex Member
SMC3	
REST	KLF Sliencing Transcription Factor
TCF4	bHLH Transcription Factor
ZNF687	Zinc Finger Protein, Transcriptional Regulator

Legend

Figure 1. Putative enhancers in term pregnant human myometrial tissues. (A) Distinct and

common putative enhancers in term pregnant myometrial biopsies from three subjects.

Genomic regions with H3K27ac and H3K4me1 double positive histone marks are defined as

putative active enhancers. (B) Association of commonly shared putative enhancers with active

genes. The association between an interval and an active gene is defined by locating within 100

kb vicinity of each other. (C) Over-Represented transcription factor binding motifs in putative

enhancers. A subset of enriched motifs that are relevant to myometrial homeostasis in the

13090 H3K4me1/H3K27ac-positive putative enhancer regions is displayed.

Figure 2. Putative super enhancers in term pregnant human myometrial tissues. (A) Number of

super enhancers mapped in tissues of each individual human subject. (B) Association of

commonly shared super enhancers with active genes in human myometrium. (C) Selected

enriched gene ontology terms in the 346 active genes that are associated with super enhancers.

Figure 3. Identification of enhancers for the *PLCL2* gene. (A) UCSC Genome Browser track

view of the human PLCL2 locus marked with gRNA targeting locations. (B-C) Relative PLCL2

mRNA levels measured by qRT-PCR in hTERT-HM cells (B) or T-HESC cells (C) that express

denoted gRNAs with the CRISPR activator (N=3 with technical duplicates). ****, $p < 0.0001$; ***,

$p < 0.001$; **, $p < 0.01$; *, $p < 0.05$ by unpaired t-test.

Figure 4. **PGR regulation of *PLCL2* expression**. (A) *PGR* mRNA abundance in hTERT-HM

cells with the PGR-promoter targeting (PGR640) or non-targeting gRNAs under the CRISPR

activation system. (N=3 with technical qPCR duplicates). (B) Protein abundance of in hTERT-

HM cells with the PGR-promoter targeting (PGR640 and P6SM) or non-targeting gRNAs under

the CRISPR activation system. Protein extracts from unmanipulated MCF7 and HEK293T cells

serve as positive and negative control for the PGR presence. (C) *PLCL2* mRNA abundance in

in hTERT-HM cells with the PGR-promoter targeting (PGR640) or non-targeting gRNAs under

the CRISPR activation system. (N=3 with technical qPCR duplicates). (D) Pearson correlation

between *PLCL2* mRNA levels and inferred PGR activities (T-Scores) in 43 human myometrial

specimens. **, $p < 0.01$; *, $p < 0.05$ by by Mann-Whitney test (A) and unpaired t-test (C).

Figure S1. Subject variations on H3K27ac-positive histone marks.

Figure S2. **PGR occupancy in myometrial enhancers**. (A) Percentages of enhancers and

super enhancers that manifest PGR occupancy. PGR genome occupancy data was previously

published in NCBI GEO accession numbers GSM4081683 and GSM4081684.

Dataset S1. H3K27ac and H3k4me1 double positive enhancers in term pregnant not in labor

human myometrial specimens.

Dataset S2. Enrichment of known transcription factor binding motifs in putative myometrial

enhancers.

Dataset S3. Super enhancers in term pregnant not in labor human myometrial specimens.

Dataset S4. Active genes associated with super enhancers in the term nonlabor myometrium.

Dataset S5. CRISPRa dependent gene expression patterns in hTERT-HM cells and gRNA

information. DEG, differentially expressed genes between denoted gRNAs.

Dataset S6. Normalized gene expression counts across the 43 human myometrial specimens

and the PGR T-Scores of individual specimens.

Dataset S7. Summary table of sequencing datasets.

Dataset S8. Summary table of Hi-C quality control metrics.

Dataset S9. Cell counts per gRNA and protospacer call frequencies per cell for Perturb-seq
analysis.

References

Balducci, J., B. Risek, N. B. Gilula, A. Hand, J. F. Egan and A. M. Vintzileos (1993). "Gap junction formation
in human myometrium: a key to preterm labor?" Am J Obstet Gynecol **168**(5): 1609-1615.
Barcelona de Mendoza, V., M. L. Wright, C. Agaba, L. Prescott, A. Desir, C. A. Crusto, Y. V. Sun and J. Y.
Taylor (2017). "A Systematic Review of DNA Methylation and Preterm Birth in African American
Women." Biol Res Nurs **19**(3): 308-317.
Bentley, D. R., S. Balasubramanian, H. P. Swerdlow, G. P. Smith, J. Milton, C. G. Brown, K. P. Hall, D. J.
Evers, C. L. Barnes, H. R. Bignell, J. M. Boutell, J. Bryant, R. J. Carter, R. Keira Cheetham, A. J. Cox, D. J.
Ellis, M. R. Flatbush, N. A. Gormley, S. J. Humphray, L. J. Irving, M. S. Karbelashvili, S. M. Kirk, H. Li, X. Liu,
K. S. Maisinger, L. J. Murray, B. Obradovic, T. Ost, M. L. Parkinson, M. R. Pratt, I. M. Rasolonjatovo, M. T.
Reed, R. Rigatti, C. Rodighiero, M. T. Ross, A. Sabot, S. V. Sankar, A. Scally, G. P. Schroth, M. E. Smith, V.
P. Smith, A. Spiridou, P. E. Torrance, S. S. Tzonev, E. H. Vermaas, K. Walter, X. Wu, L. Zhang, M. D. Alam,
C. Anastasi, I. C. Aniebo, D. M. Bailey, I. R. Bancarz, S. Banerjee, S. G. Barbour, P. A. Baybayan, V. A.
Benoit, K. F. Benson, C. Bevis, P. J. Black, A. Boodhun, J. S. Brennan, J. A. Bridgham, R. C. Brown, A. A.
Brown, D. H. Buermann, A. A. Bundu, J. C. Burrows, N. P. Carter, N. Castillo, E. C. M. Chiara, S. Chang, R.
Neil Cooley, N. R. Crake, O. O. Dada, K. D. Diakoumakos, B. Dominguez-Fernandez, D. J. Earnshaw, U. C.
Egbujor, D. W. Elmore, S. S. Etchin, M. R. Ewan, M. Fedurco, L. J. Fraser, K. V. Fuentes Fajardo, W. Scott
Furey, D. George, K. J. Gietzen, C. P. Goddard, G. S. Golda, P. A. Granieri, D. E. Green, D. L. Gustafson, N.
F. Hansen, K. Harnish, C. D. Haudenschild, N. I. Heyer, M. M. Hims, J. T. Ho, A. M. Horgan, K. Hoschler, S.
Hurwitz, D. V. Ivanov, M. Q. Johnson, T. James, T. A. Huw Jones, G. D. Kang, T. H. Kerelska, A. D. Kersey, I.
Khrebtukova, A. P. Kindwall, Z. Kingsbury, P. I. Kokko-Gonzales, A. Kumar, M. A. Laurent, C. T. Lawley, S.
E. Lee, X. Lee, A. K. Liao, J. A. Loch, M. Lok, S. Luo, R. M. Mammen, J. W. Martin, P. G. McCauley, P.
McNitt, P. Mehta, K. W. Moon, J. W. Mullens, T. Newington, Z. Ning, B. Ling Ng, S. M. Novo, M. J. O'Neill,
M. A. Osborne, A. Osnowski, O. Ostadan, L. L. Paraschos, L. Pickering, A. C. Pike, A. C. Pike, D. Chris
Pinkard, D. P. Pliskin, J. Podhasky, V. J. Quijano, C. Raczy, V. H. Rae, S. R. Rawlings, A. Chiva Rodriguez, P.
M. Roe, J. Rogers, M. C. Rogert Bacigalupo, N. Romanov, A. Romieu, R. K. Roth, N. J. Rourke, S. T.
Ruediger, E. Rusman, R. M. Sanches-Kuiper, M. R. Schenker, J. M. Seoane, R. J. Shaw, M. K. Shiver, S. W.
Short, N. L. Sizto, J. P. Sluis, M. A. Smith, J. Ernest Sohna Sohna, E. J. Spence, K. Stevens, N. Sutton, L.
Szajkowski, C. L. Tregidgo, G. Turcatti, S. Vandevondele, Y. Verhovsky, S. M. Virk, S. Wakelin, G. C.
Walcott, J. Wang, G. J. Worsley, J. Yan, L. Yau, M. Zuerlein, J. Rogers, J. C. Mullikin, M. E. Hurles, N. J.
McCooke, J. S. West, F. L. Oaks, P. L. Lundberg, D. Klenerman, R. Durbin and A. J. Smith (2008). "Accurate
whole human genome sequencing using reversible terminator chemistry." Nature **456**(7218): 53-59.
Chawanpaiboon, S., J. P. Vogel, A. B. Moller, P. Lumbiganon, M. Petzold, D. Hogan, S. Landoulsi, N.
Jampathong, K. Kongwattanakul, M. Laopaiboon, C. Lewis, S. Rattanakanokchai, D. N. Teng, J.
Thinkhamrop, K. Watananirun, J. Zhang, W. Zhou and A. M. Gulmezoglu (2019). "Global, regional, and
national estimates of levels of preterm birth in 2014: a systematic review and modelling analysis."
Lancet Glob Health **7**(1): e37-e46.
Chen, S. H., J. Haam, M. Walker, E. Scappini, J. Naughton and N. P. Martin (2019). "Production of Viral
Constructs for Neuroanatomy, Calcium Imaging, and Optogenetics." Curr Protoc Neurosci **87**(1): e66.
Condon, J., S. Yin, B. Mayhew, R. A. Word, W. E. Wright, J. W. Shay and W. E. Rainey (2002). "Telomerase
immortalization of human myometrial cells." Biol Reprod **67**(2): 506-514.
Creyghton, M. P., A. W. Cheng, G. G. Welstead, T. Kooistra, B. W. Carey, E. J. Steine, J. Hanna, M. A.
Lodato, G. M. Frampton, P. A. Sharp, L. A. Boyer, R. A. Young and R. Jaenisch (2010). "Histone H3K27ac
separates active from poised enhancers and predicts developmental state." Proc Natl Acad Sci U S A
107(50): 21931-21936.

Dai, D., E. S. Litman, E. Schonteich and K. K. Leslie (2003). "Progesterone regulation of activating protein-
1 transcriptional activity: a possible mechanism of progesterone inhibition of endometrial cancer cell
growth." J Steroid Biochem Mol Biol **87**(2-3): 123-131.
Dotts, A. J., D. Reiman, P. Yin, S. Kujawa, W. A. Grobman, Y. Dai and S. E. Bulun (2023). "In Vivo Genome-
Wide PGR Binding in Pregnant Human Myometrium Identifies Potential Regulators of Labor." Reprod Sci
30(2): 544-559.
Durand, N. C., M. S. Shamim, I. Machol, S. S. Rao, M. H. Huntley, E. S. Lander and E. L. Aiden (2016).
"Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C Experiments." Cell Syst **3**(1): 95-
98.
Erickson, E. N., L. Myatt, J. S. Danoff, K. M. Krol and J. J. Connelly (2023). "Oxytocin receptor DNA
methylation is associated with exogenous oxytocin needs during parturition and postpartum
hemorrhage." Commun Med (Lond) **3**(1): 11.
Hafemeister, C. and R. Satija (2019). "Normalization and variance stabilization of single-cell RNA-seq
data using regularized negative binomial regression." Genome Biol **20**(1): 296.
Hammal, F., P. de Langen, A. Bergon, F. Lopez and B. Ballester (2022). "ReMap 2022: a database of
Human, Mouse, Drosophila and Arabidopsis regulatory regions from an integrative analysis of DNA-
binding sequencing experiments." Nucleic Acids Res **50**(D1): D316-D325.
Heintzman, N. D., R. K. Stuart, G. Hon, Y. Fu, C. W. Ching, R. D. Hawkins, L. O. Barrera, S. Van Calcar, C.
Qu, K. A. Ching, W. Wang, Z. Weng, R. D. Green, G. E. Crawford and B. Ren (2007). "Distinct and
predictive chromatin signatures of transcriptional promoters and enhancers in the human genome." Nat
Genet **39**(3): 311-318.
Heinz, S., C. Benner, N. Spann, E. Bertolino, Y. C. Lin, P. Laslo, J. X. Cheng, C. Murre, H. Singh and C. K.
Glass (2010). "Simple combinations of lineage-determining transcription factors prime cis-regulatory
elements required for macrophage and B cell identities." Mol Cell **38**(4): 576-589.
Hewitt, S. C., S. A. Grimm, S. P. Wu, F. J. DeMayo and K. S. Korach (2020). "Estrogen receptor alpha
(ERalpha)-binding super-enhancers drive key mediators that control uterine estrogen responses in
mice." J Biol Chem **295**(25): 8387-8400.
Irizarry, R. A., C. Ladd-Acosta, B. Wen, Z. Wu, C. Montano, P. Onyango, H. Cui, K. Gabo, M. Rongione, M.
Webster, H. Ji, J. Potash, S. Sabunciyan and A. P. Feinberg (2009). "The human colon cancer methylome
shows similar hypo- and hypermethylation at conserved tissue-specific CpG island shores." Nat Genet
41(2): 178-186.
Kanematsu, T., H. Takeuchi, M. Terunuma and M. Hirata (2005). "PRIP, a novel Ins(1,4,5)P3 binding
protein, functional significance in Ca2+ signaling and extension to neuroscience and beyond." Mol Cells
20(3): 305-314.
Langmead, B., C. Trapnell, M. Pop and S. L. Salzberg (2009). "Ultrafast and memory-efficient alignment
of short DNA sequences to the human genome." Genome Biol **10**(3): R25.
Larionov, A., A. Krause and W. Miller (2005). "A standard curve based method for relative real time PCR
data processing." BMC Bioinformatics **6**: 62.
Li, J., P. R. Bushel, L. Lin, K. Day, T. Wang, F. J. DeMayo, S. P. Wu and J. L. Li (2021). "Structural Equation
Modeling of In silico Perturbations." Front Genet **12**: 727532.
Love, M. I., W. Huber and S. Anders (2014). "Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2." Genome Biol **15**(12): 550.
Merlino, A. A., T. N. Welsh, H. Tan, L. J. Yi, V. Cannon, B. M. Mercer and S. Mesiano (2007). "Nuclear
progesterone receptors in the human pregnancy myometrium: evidence that parturition involves
functional progesterone withdrawal mediated by increased expression of progesterone receptor-A." J
Clin Endocrinol Metab **92**(5): 1927-1933.

Mesiano, S., E. C. Chan, J. T. Fitter, K. Kwek, G. Yeo and R. Smith (2002). "Progesterone withdrawal and
estrogen activation in human parturition are coordinated by progesterone receptor A expression in the
myometrium." J Clin Endocrinol Metab **87**(6): 2924-2930.
Mitchell, J. A. and S. J. Lye (2001). "Regulation of connexin43 expression by c-fos and c-jun in myometrial
cells." Cell Communication & Adhesion **8**(4-6): 299-302.
Mitchell, J. A. and S. J. Lye (2005). "Differential activation of the connexin 43 promoter by dimers of
activator protein-1 transcription factors in myometrial cells." Endocrinology **146**(4): 2048–2054.
Mitsuya, K., N. Singh, S. R. Sooranna, M. R. Johnson and L. Myatt (2014). "Epigenetics of human
myometrium: DNA methylation of genes encoding contraction-associated proteins in term and preterm
labor." Biol Reprod **90**(5): 98.
Muter, J., P. J. Brighton, E. S. Lucas, L. Lacey, A. Shmygol, S. Quenby, A. M. Blanks and J. J. Brosens
(2016). "Progesterone-Dependent Induction of Phospholipase C-Related Catalytically Inactive Protein 1
(PRIP-1) in Decidualizing Human Endometrial Stromal Cells." Endocrinology **157**(7): 2883-2893.Nadeem,
L., O. Shynlova, E. Matysiak-Zablocki, S. Mesiano, X. Dong and S. Lye (2016). "Molecular evidence of
functional progesterone withdrawal in human myometrium." Nat Commun **7**: 11565.
Peavey, M. C., S. P. Wu, R. Li, J. Liu, O. M. Emery, T. Wang, L. Zhou, M. Wetendorf, C. Yallampalli, W. E.
Gibbons, J. P. Lydon and F. J. DeMayo (2021). "Progesterone receptor isoform B regulates the Oxtr-Plcl2-
Trpc3 pathway to suppress uterine contractility." Proc Natl Acad Sci U S A **118**(11).
Robinson, M. D., D. J. McCarthy and G. K. Smyth (2010). "edgeR: a Bioconductor package for differential
expression analysis of digital gene expression data." Bioinformatics **26**(1): 139-140.
Satija, R., J. A. Farrell, D. Gennert, A. F. Schier and A. Regev (2015). "Spatial reconstruction of single-cell
gene expression data." Nat Biotechnol **33**(5): 495-502.
Shchuka, V. M., L. E. Abatti, H. Hou, N. Khader, A. Dorogin, M. D. Wilson, O. Shynlova and J. A. Mitchell
(2020). "The pregnant myometrium is epigenetically activated at contractility-driving gene loci prior to
the onset of labor in mice." PLoS Biol **18**(7): e3000710.
Shin, H. Y., M. Willi, K. HyunYoo, X. Zeng, C. Wang, G. Metser and L. Hennighausen (2016). "Hierarchy
within the mammary STAT5-driven Wap super-enhancer." Nat Genet **48**(8): 904-911.
Stanfield, Z., P. Amini, J. Wang, L. Yi, H. Tan, M. R. Chance, M. Koyuturk and S. Mesiano (2019). "Interplay
of transcriptional signaling by progesterone, cyclic AMP, and inflammation in myometrial cells:
implications for the control of human parturition." Mol Hum Reprod **25**(7): 408-422.
Takenaka, K., K. Fukami, M. Otsuki, Y. Nakamura, Y. Kataoka, M. Wada, K. Tsuji, S. Nishikawa, N. Yoshida
and T. Takenawa (2003). "Role of phospholipase C-L2, a novel phospholipase C-like protein that lacks
lipase activity, in B-cell receptor signaling." Mol Cell Biol **23**(20): 7329-7338.
Uhlen, M., L. Fagerberg, B. M. Hallstrom, C. Lindskog, P. Oksvold, A. Mardinoglu, A. Sivertsson, C. Kampf,
E. Sjostedt, A. Asplund, I. Olsson, K. Edlund, E. Lundberg, S. Navani, C. A. Szigyarto, J. Odeberg, D.
Djureinovic, J. O. Takanen, S. Hober, T. Alm, P. H. Edqvist, H. Berling, H. Tegel, J. Mulder, J. Rockberg, P.
Nilsson, J. M. Schwenk, M. Hamsten, K. von Feilitzen, M. Forsberg, L. Persson, F. Johansson, M. Zwahlen,
G. von Heijne, J. Nielsen and F. Ponten (2015). "Proteomics. Tissue-based map of the human proteome."
Science **347**(6220): 1260419.
Uji, A., M. Matsuda, T. Kukita, K. Maeda, T. Kanematsu and M. Hirata (2002). "Molecules interacting with
PRIP-2, a novel Ins(1,4,5)P3 binding protein type 2: Comparison with PRIP-1." Life Sci **72**(4-5): 443-453.
Wang, H., M. Ogawa, J. R. Wood, M. S. Bartolomei, M. D. Sammel, J. P. Kusanovic, S. W. Walsh, R.
Romero and J. F. Strauss, 3rd (2008). "Genetic and epigenetic mechanisms combine to control MMP1
expression and its association with preterm premature rupture of membranes." Hum Mol Genet **17**(8):
1087-1096.Whyte, W. A., Orlando, D. A., Hnisz, D., Abraham, B. J., Lin, C. Y., Kagey, M. H., Rahl, P. B., Lee,
T. I., & Young, R. A. (2013). "Master Transcription Factors and Mediator Establish Super-Enhancers at
Key Cell Identity Genes." Cell 153(2):307-319.

Wingett, S., P. Ewels, M. Furlan-Magaril, T. Nagano, S. Schoenfelder, P. Fraser and S. Andrews (2015).
"HiCUP: pipeline for mapping and processing Hi-C data." <u>F1000Res</u> **4**: 1310.
Wu, S. P., M. L. Anderson, T. Wang, L. Zhou, O. M. Emery, X. Li and F. J. DeMayo (2020). "Dynamic
transcriptome, accessible genome, and PGR cistrome profiles in the human myometrium." <u>FASEB J</u>
34(2): 2252-2268.
Wu, S. P., T. Wang, Z. C. Yao, M. C. Peavey, X. Li, L. Zhou, I. V. Larina and F. J. DeMayo (2022).
"Myometrial progesterone receptor determines a transcription program for uterine remodeling and
contractions during pregnancy." <u>PNAS Nexus</u> **1**(4): pgac155.
Zhang, Y., T. Liu, C. A. Meyer, J. Eeckhoute, D. S. Johnson, B. E. Bernstein, C. Nusbaum, R. M. Myers, M.
Brown, W. Li and X. S. Liu (2008). "Model-based analysis of ChIP-Seq (MACS)." <u>Genome Biol</u> **9**(9): R137.

A

29681   27884   28163

13114

— Subject 1
— Subject 2
— Subject 3

B

Intervals not associated with active genes

33.2%

33.7%

13.3%   19.8%

Intervals associated with top third active genes (FPKM >= 15)

Intervals associated with bottom third active genes (5 > FPKM >= 1)

Intervals associated with middle third active genes (15 > FPKM >= 5)

C

Motif	Consensus Sequence
Known Myometrium Associated Transcription Factors	
AP-1	
STAT5	
NFkB	
Steroid Hormone Receptors	
PGR	
GRE	
Transcription Factors For Regulation of Smooth Muscle Genes	
CArG (SRF)	
ELK1	

Figure 1

A

Super Enhancers

1233   932   1106

B

Intervals not associated with active genes — 32.4%

Intervals associated with top third active genes (FPKM >= 15) — 40.3%

Intervals associated with bottom third active genes (5 > FPKM >= 1) — 7.3%

Intervals associated with middle third active genes (15 > FPKM >= 5) — 20.0%

C

Gene Ontology Term	P-value	Genes
Actin cytoskeleton organization	9.22E-05	PTPN1, SHC1, ARHGEF17, ACTN1, FHL3, ACTN4, SSH1, PDLIM1, INF2, TYROBP, PDLIM2, CAPZB, MYH9, FLNB
ECM-receptor interaction	0.024366	COL1A1, COL4A1, COL6A2, THBS1, HSPG2, CD44
DNA-binding transcription factor activity	1.18E-03	ZFHX3, NFIX, SMAD3, ZNF581, EPAS1, RFX2, FOS, NR2F2, HIC1, FOXP1, ELK3, MNT, DBP, SP1, NFIC, KLF9, ATOH8, CREB3L2, RARA, ERF, HOXD9, MEF2D
3'-UTR-mediated mRNA destabilization	0.037196	ZFP36, ZFP36L2, ZFP36L1
Estrogen signaling pathway	0.004193	SHC1, SP1, GREB1, CREB3L2, RARA, FOS, ADCY7, HSPA1B, GNAI2

Figure 2

Figure 3

Figure 4