Gene expression signatures of response to fluoxetine treatment: systematic review and meta-analyses

David G. Cooper, Ph.D.; J. Paige Cowden, Pharm.D.; Parker A. Stanley; Jack T. Karbowski; Victoria S. Gaertig; Caiden J. Lukan; Patrick M. Vo; Ariel D. Worthington; Caleb A. Class, Ph.D.*

Butler University, Department of Pharmaceutical Sciences

*Corresponding Author:
Caleb A. Class
4600 Sunset Ave.
Indianapolis, IN, USA, 46208
317-940-9831
cclass@butler.edu

Running Title: Gene expression signatures of fluoxetine response

ABSTRACT

Background: Selecting the best antidepressant for a patient with major depressive disorder (MDD) remains a challenge, and some have turned to genomic (and other 'omic) data to identify an optimal therapy. In this work, we synthesized gene expression data for fluoxetine treatment in both human patients and rodent models, to better understand biological pathways affected by treatment, as well as those that may distinguish clinical or behavioral response.

Methods: Following the PRISMA guidelines, we searched the Gene Expression Omnibus (GEO) for studies profiling humans or rodent models with treatment of the antidepressant fluoxetine, excluding those not done in the context of depression or anxiety, in an irrelevant tissue type, or with fewer than three samples per group. Included studies were systematically reanalyzed by differential expression analysis and Gene Set Enrichment Analysis (GSEA). Individual pathway and gene statistics were synthesized across studies by three p-value combination methods, and then corrected for multiple testing.

Results: Of the 74 data series that were screened, 20 were included: 18 in rodents, and two in tissue from human patients. Studies were highly heterogeneous in the comparisons of both treated vs. control samples and responders vs. non-responders, with 737 and 356 pathways, respectively, identified as significantly different between groups in at least one study. 19 pathways were identified as consistently different in responders vs. non-responders, including Toll Like Receptor (TLR) and other immune pathways. Signal transduction pathways were identified as consistently affected by fluoxetine treatment in depressed patients and rodent models.

Discussion: These meta-analyses confirm known pathways and provide new hints toward antidepressant resistance, but more work is needed. Most included studies involved rodent models, and both patient studies had small cohorts. Additional large-cohort studies applying additional 'omics technologies are necessary to understand the intricacies and heterogeneity of antidepressant response.

INTRODUCTION

Despite decades of advances in the treatment of depression, approximately half of patients do not respond to their first prescribed antidepressant(1,2). Nonresponsive patients may spend years cycling through the dozens of possible therapies before finding one that works; some will be treatment-resistant, which is associated with higher risk of suicide(3). Current approaches to treatment-resistant depression (TRD) include precision medicine--where pharmacogenetic tests are used to better match patients to

treatments--and the development of new drugs and non-pharmacological options to combat TRD directly(4–6). These advances have led to modest improvements in treatment response; for example, one clinical trial demonstrated a 5% improvement in both response and remission when the antidepressant was prescribed based on a genetic test involving 59 alleles and variants across 8 genes, versus treatment as usual(4). Meanwhile, many novel or repurposed drugs are under investigation or in trials as alternative or adjunct therapies for depression; while this will provide additional options for those suffering from TRD, selecting from the increasing number of approved antidepressants will remain challenging(5).

Selective serotonin reuptake inhibitors (SSRI's) are currently the most commonly prescribed antidepressant class, in large part due to relatively high tolerability and efficacy(7,8). As the first SSRI to win regulatory approval in the United States, and one of the best tolerated, fluoxetine remains one of the most prescribed antidepressants, and many research studies have been conducted to understand its therapeutic efficacy and mechanism of action(9,10). The advent and uptake of computational chemistry and "omics" methods have helped increase scientific knowledge in this and many other areas, providing leads for new drugs, better insights into their mechanisms of action, and potential signatures of response(5,11,12). However, these methods have not been the panaceas that some may have hoped, as heterogeneity between patients, the need to consider multiple levels of biology, and environmental and other factors can be difficult to capture(2). So while progress has been made understanding the mechanism of action of fluoxetine and other antidepressants, exact causes of differences in response between patients remain enigmatic.

The use of systematic reviews in mental disorders and other fields has dramatically increased in recent years due to the wealth of clinical data being generated across fields and the publication of the PRISMA guidelines(13–16). However, these guidelines have been applied less frequently in the use of gene expression data. Kontou *et al.* identified and meta-analyzed four studies containing gene expression data in patients with myocardial infarction and healthy controls(17). Pisanu *et al.* conducted a systematic review with qualitative synthesis of 123 gene expression studies in patients with psychiatric disorders characterized for response(18). Recently, Ibrahim *et al.* conducted a non-PRISMA systematic review and meta-analysis of gene expression signatures corresponding to fluoxetine treatment in rodents(19). These studies have focused on a specific tissue type in either humans or rodents, and translating between organisms remains difficult despite recent advances. Although over 15,000 *homo sapiens* genes have orthologues in *mus musculus*, the biological effects underlying drug treatment frequently are not shared; in cancer for example, fewer than 8% of drugs successfully translate from animal models to clinical trials

in human patients(20,21). Alternatively, meta-analyses considering biological pathways allow us to focus on the shared biological effects between studies, which likely are more consistent across organisms(22,23). In this systematic review and meta-analysis, we will apply this approach to summarize and potentially identify new biological pathways of antidepressant treatment and response.

The objectives of this systematic review are to assess and synthesize the evidence for gene expression modification by fluoxetine treatment, and whether gene expression levels distinguish clinical or behavioral response to treatment. In this paper, we use "Response Signatures" to refer to gene expression changes that differ between those with a good vs. poor response to fluoxetine, while "Treatment Signatures" signifies gene expression changes that differ with fluoxetine treatment vs. control. To our knowledge, this is the first meta-analysis of gene expression data to investigate behavioral or clinical response to an antidepressant, as well as the first to integrate data across multiple organisms and tissue types. Results of the meta-analyses may be applied to improving the prediction and assessment of fluoxetine response shortly after treatment, and they may provide hints for combination therapies or new drug development for improved response. Additionally, protocols developed for this study can be applied, improved, and expanded for future systematic studies involving other drugs and 'omics data types.

METHODS

The Gene Expression Omnibus (GEO) database was identified as the main source for this systematic review, due to its primary focus on gene expression data and the relative consistency of data deposits and formatting(24). PubMed was identified as a contingent database in case the GEO search resulted in fewer than five studies that passed screening. As one of the most prescribed and studied antidepressants, fluoxetine was selected as the focus of this systematic review; we were more inclusive regarding organisms and tissue type, allowing us to identify both heterogeneity and potential consistencies. The GEO search was conducted on 4 April 2023, using the following keywords: *(fluoxetine) OR (selective serotonin reuptake inhibitor) OR (ssri)*. Results were filtered within GEO using *Entry type = Series* (to return full data sets rather than individual samples), *Organism = homo sapiens, mus musculus,* or *rattus norvegicus*, and *Study type = Expression profiling by array* or *expression profiling by high throughput sequencing*. The resulting data series were manually filtered by authors DC and CC based on the exclusion criteria: studies were excluded if they were not primarily focused on depression or anxiety, were not conducted in a relevant tissue type or genetic background (cancer cell lines, for example), did not involve treatment with fluoxetine, or if there were fewer than three samples per group. Decisions were made

based on review of the GEO series abstract and sample metadata. If the decision was unclear, any associated publication and supplementary materials were consulted to gain additional information. For the synthesis, studies would be grouped based on the comparison: good vs. poor response, fluoxetine treatment vs. control, or fluoxetine treatment vs. control (within depressed patients and rodent models, and separately within unstressed rodents). Studies would be synthesized both across and within organism.

Data extraction, transformation, synthesis, and assessment were conducted using R version 4.2.3(25). Each individual analysis was completed by one author and then checked by DC or CC. All analysis code and results are available for download (see *Code Availability*). Gene expression data were directly downloaded using the R library GEOquery, and then checked for completeness(26). Data sets that were not retrieved properly using GEOquery were instead downloaded directly from the GEO website and then imported into R for analysis. Basic quality control was assessed by visualizing and comparing gene expression distributions, principal components analysis, and between-sample correlations. Clear outlier samples were removed prior to any subsequent analyses. Differential expression analysis was conducted using the R libraries *DESeq2* for RNA sequencing data, or *limma* for microarray data(27,28). All differential expression analyses were conducted as *group* vs. *group*, with the main variables being treatment status or response (as appropriate). Other variables sought were tissue type, time point, and participant/rodent ID. For each comparison, the effect measures used in subsequent steps were the log2 fold change (log2FC), nominal p-value, and t statistic. When multiple tissue types were present in a study, separate group vs. group analyses were conducted within each tissue. Risk of bias was assessed by evaluating the provided characteristics of the study populations.

Differential expression results for each comparison were summarized to the pathway level using the *fgsea* implementation of Gene Set Enrichment Analysis (GSEA), with gene sets defined by the Consensus Pathway Database (CPDB)(29–31). The CPDB integrates gene set and pathway information from 31 sources, including metabolic pathways, signaling pathways, and gene regulation; it includes orthologous gene sets for human and mouse in the Reactome and KEGG pathway databases(32,33), allowing for the synthesis of human and rodent models at the pathway level (for rat data sets, gene ID's were converted to mouse ID's by homology prior to gene set analysis). A ranked list of differential expression t-statistics from each comparison was input to *fgsea*, and relevant effect measures for each pathway were the normalized enrichment score (NES), providing the magnitude of gene expression enrichment in one group or the other, and nominal p-value.

Results across comparisons were synthesized using the *MetaDE* R package(34). P-value combination methods were selected due to substantial heterogeneity between studies (organism, tissue type, and gene expression platform). Within organisms, p-values were synthesized at both the gene and pathway level, while across-organism syntheses were only conducted at the pathway level. Fisher's method was used to identify genes or pathways that were differentially expressed in *any study*, while Wilkinson's method (*Max-P*) was used to identify genes or pathways that were consistently differentially expressed *across studies*(35–37). The resulting meta-analysis p-values were corrected for false discovery using the method of Benjamini and Hochberg to result in q-values(38). As an *ad hoc* intermediate method, we also selected genes or pathways with nominal $p<0.05$ in greater than half of synthesized comparisons (Frequency of 50%, or *Freq50*). A depiction of these methods is provided in **Figure 2A**. Synthesis results were displayed using Venn diagrams to show how many genes/pathways were identified by each meta-analysis method, and bar graphs showing pathways identified by Max-P with $q < 0.05$.

Heterogeneity among study results was investigated using subgroup analysis (meta-analysis within specific organisms or tissue types), as well as heatmaps of the NES of pathways across studies. Additionally, consistency among the direction of results was estimated by simple vote-counting across comparisons: a pathway or gene was assigned +1 for each comparison where it was significantly enriched in responders with nominal $p<0.05$, or -1 if enriched in non-responders. These were summed across all comparisons, and numbers further from zero indicate greater consistency in results that are statistically significant within comparisons.

Sensitivity analyses were performed to assess the meta-analyses. Of 9 response vs. non-response comparisons, 8 involved mice while the other included human patients, so we looked at inclusion/exclusion of the patient data set. We further explored the effect of tissue type by removing the data set profiling blood in mice, resulting in 7 comparisons in brain tissue only. Of the 13 treated vs. untreated experiments in unstressed mouse models, 12 involved microarray profiling in male mice, while the other used RNA-Seq in female mice; we conducted sensitivity analysis to investigate the effects of inclusion/exclusion of this last study in meta-analysis. A sensitivity analysis compared meta-analysis results of treatment signatures when DS19 (containing 27 comparisons from various brain regions) was removed. Finally, the consistency of direction of effect across studies, as evidenced by vote-counting, was used to estimate certainty of the meta-analyses. For genes or pathways identified as differentially expressed in all or most studies, we generally expect them to agree in the direction of the effect (i.e.

overexpressed in treatment or control). This is not assured, as differing and even opposite effects have been reported in some cases(39,40).

RESULTS

The initial keyword search on resulted in 1958 entries, which was filtered to 84 data sets (known in GEO as "series") using automatic filtering for Entry Type, Study Type, and Organism (**Figure 1**). Many of the automatically excluded entries were individual samples: in almost all cases, these will also be included in one of the returned series, so they can be considered as duplicates, and any individual samples that are not part of a relevant data series are not of interest for this meta-analysis study. 10 "SuperSeries" were removed as duplicates, as they contained one or more individual series returned by the search. 74 data series were manually assessed for the exclusion criteria (**Supplementary Table 1**), resulting in 20 selected for inclusion(41–56) (**Table 1, Figure 1**). Of the 20 included data sets, two involved tissue from patients diagnosed with Major Depressive Disorder (MDD): one profiled gene expression in whole blood from adolescent females before and after eight weeks of continuous fluoxetine treatment(53), while the other profiled lymphoblastoid cell lines (LCL's) developed from 10 patients with known antidepressant response based on change in score by Hamilton Depression Rating Scale (HDRS). These LCL's were treated with fluoxetine or control *in vitro* for three weeks(43). Of the 10 patients in this study, six were classified as responders, while the other four were classified as non-responders.

The remaining 18 studies profiled gene expression with fluoxetine treatment in rodent models (11 in mice, seven in rats). Stress models were applied in nine of these studies to achieve anxious and/or depressive behavior: five employed chronic mild stress(44,49,56–58), while the others used either chronic restraint stress(50), single housing(45), injection with corticosterone(42), or selective breeding for high anxiety(51). Five studies included a classification of response in the treated animals, which was assessed using a behavioral method such as the open-field test for measurement of anxiety(45) or the forced swim test (FST) for despair(42). One study did not induce stress but did measure behavioral response: anxiety was measured by the open field test after three weeks of fluoxetine or vehicle treatment, and response was defined based on the ratio of scores between the fluoxetine and vehicle groups(41). One mouse study profiled whole blood, but the majority collected samples from one or more brain regions or neuronal cell types for gene expression profiling: the most common was the hippocampus or a hippocampal region (particularly dentate gyrus, included in five studies(42,44,49,52,55)). One profiled fluoxetine effects across

27 brain regions in rats, identifying region-specific differential expression signatures in both bulk and single-cell RNA-Seq data (we did not include single-cell data in this work) (**Supplementary Table 2**)(55).

As is common with rodent models, males were utilized in 17 of the 18 studies, demonstrating a substantial risk of bias. One study profiled gene expression with fluoxetine and imipramine treatment in female mice, which had been shown to exhibit behavioral despair by FST and tail suspension test (TST) after reduction in *Brd1* expression (this was not observed in male mice with reduced *Brd1* expression)(54). Conversely, the two human studies were biased toward female inclusion, with one including adolescent females only(53), and the other including eight females and two males in the microarray analysis(43). For the seven of the 20 data sets synthesized for Response Signatures, all six mouse data sets employed males, while the patient cohort was biased toward females.

Response Signatures

Meta-analysis by Fisher's method, the Max-P method, and the Freq50 method (**Figure 2A**) were applied on nine comparisons from seven data series to synthesize gene expression signatures associated with clinical or behavioral response to fluoxetine (**Figure 2B**). Systematic re-analysis of individual comparisons resulted in widely varying numbers of differentially expressed genes and pathways after false discovery correction for multiple testing (**Figure 2C**). DS6a had the greatest number of significant genes at 85, while four comparisons showed no statistically significant differential expression among genes. Conversely, statistically significant pathway enrichment was observed for all studies except DS4. Nominal p-values were synthesized across all comparisons for each Reactome and KEGG pathway and then corrected for false discovery to result in q-values. After false discovery correction, meta-analysis identified 356 pathways (over 30% of the pathways analyzed) enriched in at least one comparison with q<0.05 (Fisher's method), 19 pathways enriched across comparisons (Max-P), and 28 pathways enriched in at least 5 comparisons (Freq50) (**Figure 2D**). Pathways involving metabolism of proteins or RNA, transcription, or the immune system were most likely to be enriched (**Figure 2E**).

We then considered direction of enrichment through simple vote-counting across comparisons: a pathway was assigned +1 if significantly enriched in responders, or -1 if enriched in non-responders. The distribution of vote sums for each pathway across comparisons is provided in **Figure 2F**. Pathways that were not identified as enriched by either meta-analysis method had distributions centered around zero, with 92% of pathways ranging from -1 to +1, indicating the expected low agreement. Those identified as significant by meta-analysis had a wider range, indicating more consistent enrichment in responders or

non-responders. However, the greatest vote sum magnitude was -5 (of a possible 9 if all comparisons were significant in the same direction), indicating moderate certainty of results.

The Normalized Enrichment Scores (NES) by GSEA for pathways enriched by Fisher's method are presented in **Figure 2G**. Strong heterogeneity is observed across comparisons, but some patterns can be detected: the two studies profiling expression in mouse blood or human lymphoid-derived cells cluster together, while the seven from brain tissue comprise the other cluster. A group of immune pathways show strong enrichment in resistant blood samples, but weak, opposite enrichment across brain samples from responding mice. Other clusters of pathways show heterogeneous patterns within the set of comparisons profiling brain tissue, and none are enriched in the same direction across all comparisons, as evidenced by the distribution of vote sums.

Next, we focused on pathways that were identified as consistently enriched across comparisons with the Max-P method. Meta-analysis statistics for pathways with $q<0.05$ are presented in **Figure 3A,** and a network diagram showing similarity between these gene sets is presented in **Figure 3B** (a heatmap showing overlap between gene sets is provided in **Supplementary Figure 1**). Signal Transduction (a top-level Reactome pathway consisting of over 2000 genes) was the most consistently enriched pathway with $q<0.001$ by Max-P and a vote sum of +3, indicating somewhat consistent enrichment in good responders. 10 immune pathways were identified as consistently enriched, including five toll-like receptor (TLR) pathways and two MyD88 cascade pathways; these seven pathways are highly similar, sharing over 95% of genes. Additionally, C-type lectin receptors (CLRs), Downstream signaling of B Cell Receptor (BCR), and the top-level Immune System pathway were identified by meta-analysis. These were slightly enriched in good responders by total vote, except for the CLRs pathway with a vote sum of -1 (significantly enriched in non-responders in patient-derived LCL's). Additionally, the NF-kappa B signaling pathway from KEGG was enriched in good responders with a vote sum of +3.

Pathways related to metabolism of proteins or RNA were most consistently enriched in resistant samples, including the substantially overlapping pathways related to 40S and 60S ribosomal subunits (both with vote sums of -3), L13a-mediated translational silencing of Ceruloplasmin expression (-3), and Nonsense Mediated Decay independent of the Exon Junction Complex (-4). These were enriched in non-responders in both comparisons involving blood-derived tissue, as well as some (but not all) of the comparisons in brain tissue. Additionally, Extension of Telomeres ($q = 0.02$, 48 genes, with negligible overlap with the other enriched pathways shown in **Figure 3B**) was enriched by Max-P but with a vote sum of -1.

Meta-analysis was conducted for differential expression at both the gene and pathway level for the eight comparisons from six mouse data sets. More pathways were identified as consistently differentially expressed by Max-P (27 with only mouse studies vs. 19 including the patient cohort), although many of the uniquely identified pathways also involved TLR cascades (**Supplementary Figure 2**). Individual gene meta-analysis proved less consistent, with only 40 genes identified as differentially expressed in *any* comparison by Fisher's method, four differentially expressed in over half of the comparisons, and none identified as consistent by the Max-P method (**Figure 3C**). A meta-volcano plot of these results is presented in **Figure 3D**. Fosl2 is both highly enriched by Fisher's method ($q<0.001$) and consistently overexpressed in good responders (vote sum of +5). Notch4 (+4), Golm1 (-4), and Adamts (-4) are the other genes that are most consistently differentially expressed across comparisons in mice.

Because of the clear clustering by tissue, we conducted two sensitivity analyses successively removing individual comparisons from the meta-analysis. The previously discussed meta-analysis in mouse studies demonstrated the effects of removing the single patient data set; we then conducted a second analysis synthesizing the seven mouse comparisons in brain tissues only. Overlap between pathways identified by Fisher's and Max-P methods can be seen in **Supplementary Figure 2**. As individual data sets were removed, pathways identified by Fisher's method decreased, as this method identifies whether a pathway is enriched in *any* of the studies. By Max-P, only two pathways were identified in the full set but not in the subsets, while ten additional pathways were identified when the patient comparison was removed (as previously discussed). Meta-analysis of the seven brain comparisons did not identify any additional pathways by Max-P: thus, we can conclude that the Max-P meta-analysis of responders vs. non-responders was only slightly sensitive to the inclusion of gene expression profiling in blood samples, as all pathways identified as consistently enriched in responders or non-responders in brain were also enriched in blood (although not necessarily in the same direction).

Finally, we compared results from two studies that profiled gene expression in known or inferred responders and non-responders that had not previously received fluoxetine: one in patient-derived LCL's and the other in mouse cortex samples (mouse response was inferred based on overall response measured for multiple mice from each strain)(41). 13 pathways were identified as enriched by GSEA with $q<0.05$ in both comparisons, and all were enriched in good responders (**Figure 3E**). These included pathways related to metabolism and mitochondrial translation, in addition to gene signatures of Alzheimer's disease and Parkinson's disease.

Treatment Signatures

The same meta-analyses were applied on 55 comparisons of treated vs. untreated/control samples from 20 data sets, to synthesize gene expression changes due to treatment (**Figure 4A**). First, each data set was systematically reanalyzed, again providing widely varying results (**Figure 4B**): after false discovery correction, the number of differentially expressed genes ranged from zero (in 19 comparisons) to 8663, and the number of significantly enriched pathways ranged from zero (in 7 comparisons) to 971. Fisher's meta-analysis identified 737 pathways as enriched in at least one comparison; of these, 31 were enriched with nominal p<0.05 in more than half of the comparisons, and 10 were identified as consistently enriched by the Max-P method (**Figure 4C**).

The pathways identified as enriched in any of the 55 comparisons (65% of all pathways analyzed) are represented by all pathway categories, but most frequently components of Neuronal System or the KEGG database (**Figure 4D**). Heterogeneous enrichment patterns are again observed across studies (**Figure 4E**). Tissue type again played a role, as three of the four comparisons from blood or LCL's clustered tightly together, and most comparisons from hippocampus and cortex clustered together. Comparisons profiled in the same study frequently clustered together as well, evidenced by the data set ID's.

Overall, pathways identified only by Fisher's method have a wider distribution of vote sums (but still highly concentrated between +6 and -6, of a possible 55 for perfect agreement) than those that are not significantly enriched, while those identified by Max-P are skewed toward the left indicating downregulation by treatment (**Figure 4F**). The 10 pathways enriched by Max-P (**Figure 4G**) mainly come from the Signal Transduction category of Reactome or the KEGG Database, in addition to N-glycan antennae elongation in the medial/trans-Golgi (q=0.001, vote sum of -3). G alpha (i) signaling events are identified as most enriched by q-value and somewhat consistently decreased by treatment (q=0.001, -11). GABAergic synapse is most consistently decreased by treatment with a vote sum of -24 (q=0.006). There was not substantial overlap between genes within enriched pathways (**Figure 4H** and **Supplementary Figure 3**).

Subset meta-analysis was conducted for 13 comparisons from 10 data sets profiling fluoxetine treatment of stressed mice, rats, or human MDD patients. 731 pathways were identified as enriched in at least one comparison, and 101 were identified as consistently enriched by Max-P (**Figure 5A**). Greater enrichment is observed in immune pathways in the subset analysis as compared with the full analysis (**Figure 5B**). Generally low agreement between studies is observed, with vote sums ranging from -2 to +2 for 98.7% of

pathways not identified as enriched by meta-analysis (**Figure 5C**). 21% of pathways identified by both Fisher's and the Max-P method had vote sums outside this range.

Of the pathways identified by Max-P, we highlighted those with absolute vote sums of 3 or greater in **Figure 5D**, with gene set similarity presented in **Figure 5E** (all 101 pathways are presented in **Supplementary Figure 4**). Most significantly enriched are two NTRK signaling pathways, both with q<0.001 and vote sums of +4 and +6 respectively (of 13 possible), indicating moderate certainty. Other signal transduction pathways show a decrease with treatment, including Intracellular signaling by second messengers (q<0.001, vote sum of -3) and PIP3 activates AKT signaling (q=0.001, -5). The top-level Neuronal System gene set is significantly enriched with treatment (q=0.005, +6); conversely, the component Long-term Potentiation pathway is decreased (q=0.003, -4). A network graph showing overlap between all pathways identified by Max-P indicates groups of connected pathways from the KEGG Database and Reactome Signal Transduction, in addition to smaller groups of other pathways (**Figure 5E**).

A summary of all within-species meta-analysis results (bottom level of **Figure 4A**) is presented in **Supplementary Figure 5**. Profiling treatment effects in stressed mice identified the greatest number of differentially expressed genes, with 11,541 identified by Fisher's method and 1729 identified as consistently differentially expressed by Max-P (**Figure 5F**). Genes identified by both Fisher's method and Max-P had a wider distribution of Vote Sums, with 39.8% with absolute value of 3 or greater; 95.7% of those not identified by either method ranged from -1 to +1 (**Figure 5G**). Seven genes were significantly affected by treatment in the same direction across all seven comparisons: Bdnf, Zfp703, Raogap1, Rcan1, Dock10, St8sia4, and Lrrn3 (**Figure 5H**). Of these, Brain-derived neurotrophic factor (Bdnf) was identified as most significantly affected by Max-P (q<<0.001).

Overall, low agreement is observed between the meta-analysis of stressed rodents or MDD patients and the meta-analysis of unstressed rodents. No pathways overlap by Max-P, although 425 pathways overlapped by Fisher's method (**Supplementary Figure 6**). We conducted a sensitivity analysis removing DS19 (profiling across 27 brain regions), as this was over half of the comparisons included in the synthesis, some of which were reported as minimally affected by fluoxetine(55). We expected our Max-P results to be sensitive to comparisons in less-affected brain regions--conversely, we saw that fewer pathways were consistently enriched upon removal of this study (**Supplementary Figure 7**). Considering the gene level, 149 genes were consistently differentially expressed in the original meta-analysis, vs. none when DS19 was removed. Thus, we can conclude that meta-analysis of treatment effects was sensitive to DS19. As

seen in **Figure 4E**, comparisons across the 27 tissue types in DS19 showed largely consistent patterns, likely contributing to the observed result.

Finally, we conducted separate analyses and meta-analyses within responder and non-responder groups across studies that provided response information, to identify gene expression changes that may indicate response to treatment. The Max-P method identified 22 pathways that were consistently changed across both groups, 86 pathways that were specific to responders, and 29 that were specific to non-responders (**Figure 6A**). Of the pathways that consistently changed with treatment in responders, we focused on the 17 that were consistently *unchanged* in non-responders (**Figure 6B**, q > 0.05 in non-responders by Max-P and Fisher's method). This included eight pathways from the KEGG database, five signal transduction pathways, two immune pathways, and two others. However, vote sums ranged from -3 to +3, indicating weak certainty of evidence. Conversely, only the estrogen signaling pathway was consistently changed in non-responders but consistently unchanged in responders based on the same criteria.

DISCUSSION

Across nine independent comparisons from six gene expression data sets, 19 pathways were identified as consistently enriched by the Max-P meta-analysis method. This was surprising, as these studies profiled different tissue types and employed varying stress models in mice--in addition to one which did not induce stress, and another using samples from depressed patients. This also demonstrated the ability of pathway methods vs. individual gene meta-analysis in identifying consistency: among the eight comparisons in mice, zero genes were identified with consistent differences between responders and non-responders by Max-P, and only four were identified as differentially expressed in greater than half of comparisons. Less surprisingly, over three hundred pathways were identified as enriched in at least one comparison by Fisher's method, demonstrating the expected heterogeneity in behavioral response markers between studies. Some of this heterogeneity can be explained by tissue type, as comparisons from similar tissues clustered nearest each other (**Figure 2G**). Thus, we also conducted separate meta-analysis excluding studies profiling expression in blood, but saw only a slight effect on the meta-analysis result by Max-P.

Immune pathways were well represented among the 19 pathways that were consistently different between responders and non-responders, including seven Toll like receptor pathways, and pathways involving the C-type lectin receptor and B-cell receptor. Immune and inflammatory processes have been frequently implicated in the development of depression and antidepressant response(59–65). Additionally, one of the included studies specifically focused on involvement of inflammation via the TLR

pathway in the pathogenesis of depression and highlighted the potential contribution of acupuncture in relieving this inflammation(50). However, the direction of the change across the meta-analyzed studies was not consistent across all studies: while immune pathways were strongly upregulated in resistant samples in the two studies profiling gene expression in blood (one derived from human patients), these pathways were weakly upregulated in responding samples in studies profiling brain tissues. A recent systematic review has noted that inflammatory markers are only weakly correlated between peripheral and cerebrospinal fluid(66).

Protein and RNA metabolism pathways were among the most consistently enriched in samples resistant to fluoxetine treatment across organisms, in both studies profiling blood and most profiling brain tissue types. Specifically, two pathways involving the 40S and 60S ribosomal subunits were identified: a few studies have previously identified ribosomal proteins and pathways as implicated in response to antidepressants(67,68). These signatures may also be connected to immune response; although they share very few genes with the identified immune pathways, there is some evidence that ribosomes regulate gene expression involved in immune response(69).

Tissue heterogeneity again played a role when considering gene expression changes due to fluoxetine treatment, with blood derived tissues showing distinct overall effects from those in the brain (**Figure 4E**). However, considering the 55 widely varying comparisons of fluoxetine treatment vs. control, ten pathways were identified as consistently differentially expressed with the Max-P method (**Figure 4G**). Of these, the GABAergic synapse was overwhelmingly downregulated by fluoxetine. GABAergic neurons have been implicated in depression and antidepressant response, and positive modulators of the GABA$_A$ receptor have been approved by the FDA for postpartum depression(70–73). Importantly, meta-analysis limited to depressed patients and stressed mouse models did not identify this pathway as consistently affected, highlighting the difficulty of translating these observations to clinical application.

When considering only stressed models, many more pathways (101) are identified as consistently affected by fluoxetine by the Max-P method (**Figure 5A**). Further separation into treatment effects of responders and non-responders demonstrated a greater number of pathways consistently affected by treatment in responders than non-responders (**Figure 6A**). The 17 pathways that were consistently affected by treatment in responders but *never* affected by treatment in non-responders (**Figure 6B**) may provide hints in this area, although the agreement between studies as evidenced by the vote sums is not substantial. Treatment effects were again observed in signal transduction and immune pathways, while pathways related to cancer and addiction were also slightly affected. Antidepressants have recently been

demonstrated to inhibit liver and lung cancer through the mTOR pathway, although previous evidence has also indicated positive associations between antidepressant use and cancer in some cases(74–76).

While the PRISMA guidelines are more commonly applied for systematic review of clinical studies, we felt that they provided a strong framework for conducting these meta-analyses. We intentionally used broad inclusion criteria for organism and tissue types because we wanted to see if it was possible to observe consistent changes even across very heterogeneous studies; the use of Fisher's meta-analysis method, subgroup analyses, and sensitivity analyses then let us specifically explore the expected heterogeneity. Reporting bias is challenging to assess in gene expression analysis, where thousands of statistics are generated for each study. Yousefi *et al*. have assessed risk of reporting bias when classification algorithms are applied to gene expression data(77), but the authors are not aware of methods to detect reporting bias of the gene expression data itself; this will be a valuable tool as systematic reviews of gene expression studies become more prevalent.

This study did include multiple limitations that should be addressed to better understand antidepressant effectiveness. Only two studies provided gene expression data for responders and non-responders prior to treatment, so it is not possible to derive strong inferences for predictive markers of fluoxetine response; even large-scale patient studies have struggled to identify predictive markers to this point(18,78). Only two patient cohorts were included in our meta-analyses; other studies have been conducted in patients, but they tend to focus on a small number of biomarkers or otherwise do not submit full expression data to GEO(18). Most data sets profiled in our meta-analyses contained sample sizes of five or fewer per group, although some samples were pooled across multiple rodents. Additionally, this systematic review was not prospectively registered, and the reviewers did not work independently (note that all exclusion decisions are documented in **Supplementary Table 1**).

The included studies showed substantial bias based on sex, with 17 of 18 rodent studies profiling males, while the two patient studies were biased toward female inclusion. Results of Max-P meta-analysis provide some evidence for conserved response and treatment signatures, but it is not possible to identify sex-dependent biological signatures with these data. Previous work has shown both consistency and divergence between males and females regarding clinical response, molecular signatures, and adverse effects from antidepressants, and this remains an important area of study(39,79–82).

Considering the meta-analysis methods, reliance on the Max-P method means that we are particularly sensitive to a single aberrant study resulting in high p-values due to data quality or an unconsidered factor,

which may not have been apparent during screening. For this reason, all code and results are provided in our registered repository for further inspection and analysis, including individual data set analyses, as well as other meta-analyses by Fisher's method (which is more sensitive to studies that may have extremely low p-values) and the Freq-50 method (which may be the most robust but does not test a specific hypothesis).

Our meta-analyses have emphasized some known pathways in antidepressant response and unearthed a few new routes of potential investigation, but a true understanding of antidepressant response will require additional large-cohort studies focusing on transcriptomic data. However, recent studies of antidepressant response in over 100 patients have resulted in few, if any, individual biomarkers of response, indicating that there is not a single genetic signature of response, but a variety of 'omic and environmental factors that may contribute(18,78). Emerging technologies and approaches have begun to allow us to understand genomics and transcriptomics at a deeper level, including single-cell profiling, alternative splicing, and various epigenetic factors(55,83–85). And as technology improves our ability to quantify protein and metabolite levels, it will be valuable to incorporate these data, as this will bring us even closer to the true biology underpinning these complex disorders.

Acknowledgements

The authors gratefully acknowledge C. Andrew Class, M.D. (Ascension St. Vincent) and Marcos Oliveira, Ph.D. (Butler University) for helpful comments and conversations. This work was supported by the American Association of Colleges of Pharmacy (AACP New Investigator Award to CAC), the Butler University Holcomb Awards Committee, and departmental funds. Funders were not directly involved in conducting the project.

Competing Interests

None declared.

Code Availability

All analyses completed in this work are included at https://github.com/DavidGCooper/FLX-MetaDE with DOI 10.5281/zenodo.10668845(86).

REFERENCES

1. Rush AJ. STAR*D: What Have We Learned? AJP. 2007 Feb;164(2):201–4.

2. Akil H, Gordon J, Hen R, Javitch J, Mayberg H, McEwen B, et al. Treatment resistant depression: A multi-scale, systems biology approach. Neuroscience & Biobehavioral Reviews. 2018 Jan 1;84:272–88.

3. Bergfeld IO, Mantione M, Figee M, Schuurman PR, Lok A, Denys D. Treatment-resistant depression and suicidality. Journal of Affective Disorders. 2018 Aug 1;235:362–7.

4. Greden JF, Parikh SV, Rothschild AJ, Thase ME, Dunlop BW, DeBattista C, et al. Impact of pharmacogenomics on clinical outcomes in major depressive disorder in the GUIDED trial: A large, patient- and rater-blinded, randomized, controlled study. Journal of Psychiatric Research. 2019 Apr 1;111:59–67.

5. Borbély É, Simon M, Fuchs E, Wiborg O, Czéh B, Helyes Z. Novel drug developmental strategies for treatment-resistant depression. British Journal of Pharmacology. 2022;179(6):1146–86.

6. Farah WH, Alsawas M, Mainou M, Alahdab F, Farah MH, Ahmed AT, et al. Non-pharmacological treatment of depression: a systematic review and evidence map. BMJ Evidence-Based Medicine. 2016 Dec 1;21(6):214–21.

7. Cipriani A, Furukawa TA, Salanti G, Chaimani A, Atkinson LZ, Ogawa Y, et al. Comparative efficacy and acceptability of 21 antidepressant drugs for the acute treatment of adults with major depressive disorder: a systematic review and network meta-analysis. The Lancet. 2018 Apr 7;391(10128):1357–66.

8. Liu Z, Zhang Y, Franzin L, Cormier JN, Chan W, Xu H, et al. Trends and variations in breast and colorectal cancer incidence from 1995 to 2011: A comparative study between Texas Cancer Registry and National Cancer Institute's Surveillance, Epidemiology and End Results data. International Journal of Oncology. 2015 Apr;46(4):1819–26.

9. Kryst J, Majcher-Maślanka I, Chocyk A. Effects of chronic fluoxetine treatment on anxiety- and depressive-like behaviors in adolescent rodents - systematic review and meta-analysis. Pharmacol Rep. 2022 Oct;74(5):920–46.

10. Benfield P, Heel RC, Lewis SP. Fluoxetine. Drugs. 1986 Dec 1;32(6):481–508.

11. Casarotto PC, Girych M, Fred SM, Kovaleva V, Moliner R, Enkavi G, et al. Antidepressant drugs act by directly binding to TRKB neurotrophin receptors. Cell. 2021 Mar 4;184(5):1299-1313.e19.

12. Mariani N, Cattane N, Pariante C, Cattaneo A. Gene expression studies in Depression development and treatment: an overview of the underlying molecular mechanisms and biological processes to identify biomarkers. Transl Psychiatry. 2021 Jun 8;11(1):1–23.

13. Hoffmann F, Allers K, Rombey T, Helbach J, Hoffmann A, Mathes T, et al. Nearly 80 systematic reviews were published each day: Observational study on trends in epidemiology and reporting over the years 2000-2019. Journal of Clinical Epidemiology. 2021 Oct 1;138:1–11.

14. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ. 2021 Mar 29;372:n71.

15. Noetel M, Sanders T, Gallardo-Gómez D, Taylor P, Cruz B del P, Hoek D van den, et al. Effect of exercise for depression: systematic review and network meta-analysis of randomised controlled trials. BMJ. 2024 Feb 14;384:e075847.

16. Meehan AJ, Lewis SJ, Fazel S, Fusar-Poli P, Steyerberg EW, Stahl D, et al. Clinical prediction models in psychiatry: a systematic review of two decades of progress and challenges. Mol Psychiatry. 2022 Jun;27(6):2700–8.

17. Kontou P, Pavlopoulou A, Braliou G, Bogiatzi S, Dimou N, Bangalore S, et al. Identification of gene expression profiles in myocardial infarction: a systematic review and meta-analysis. BMC Medical Genomics. 2018 Nov 27;11(1):109.

18. Pisanu C, Severino G, De Toma I, Dierssen M, Fusar-Poli P, Gennarelli M, et al. Transcriptional biomarkers of response to pharmacological treatments in severe mental disorders: A systematic review. European Neuropsychopharmacology. 2022 Feb 1;55:112–57.

19. Ibrahim EC, Gorgievski V, Ortiz-Teba P, Belzeaux R, Turecki G, Sibille E, et al. Transcriptomic Studies of Antidepressant Action in Rodent Models of Depression: A First Meta-Analysis. International Journal of Molecular Sciences. 2022 Jan;23(21):13543.

20. Breschi A, Gingeras TR, Guigó R. Comparative transcriptomics in human and mouse. Nat Rev Genet. 2017 Jul;18(7):425–40.

21. Mak IW, Evaniew N, Ghert M. Lost in translation: animal models and clinical trials in cancer treatment. Am J Transl Res. 2014 Jan 15;6(2):114–8.

22. Shen K, Tseng GC. Meta-analysis for pathway enrichment analysis when combining multiple genomic studies. Bioinformatics. 2010 May 15;26(10):1316–23.

23. Manoli T, Gretz N, Gröne HJ, Kenzelmann M, Eils R, Brors B. Group testing for pathway analysis improves comparability of different microarray datasets. Bioinformatics. 2006 Oct 15;22(20):2500–6.

24. Barrett T, Wilhite SE, Ledoux P, Evangelista C, Kim IF, Tomashevsky M, et al. NCBI GEO: archive for functional genomics data sets—update. Nucleic Acids Research. 2013 Jan 1;41(D1):D991–5.

25. R Core Team. R: A Language and Environment for Statistical Computing [Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2023. Available from: https://www.R-project.org/

26. Davis S, Meltzer PS. GEOquery: a bridge between the Gene Expression Omnibus (GEO) and BioConductor. Bioinformatics. 2007 Jul 15;23(14):1846–7.

27. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology [Internet]. 2014 [cited 2016 Feb 19];15(12). Available from: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4302049/

28. Ritchie ME, Phipson B, Wu D, Hu Y, Law CW, Shi W, et al. limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Research. 2015;43(7):e47.

29. Subramanian A, Tamayo P, Mootha VK, Mukherjee S, Ebert BL, Gillette MA, et al. Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. Proceedings of the National Academy of Sciences. 2005 Oct;102(43):15545–50.

30. Sergushichev A. An algorithm for fast preranked gene set enrichment analysis using cumulative statistic calculation. bioRxiv [Internet]. 2016; Available from: http://biorxiv.org/content/early/2016/06/20/060012

31. Herwig R, Hardt C, Lienhard M, Kamburov A. Analyzing and interpreting genome data at the network level with ConsensusPathDB. Nat Protoc. 2016 Oct;11(10):1889–907.

32. Fabregat A, Jupe S, Matthews L, Sidiropoulos K, Gillespie M, Garapati P, et al. The Reactome Pathway Knowledgebase. Nucleic Acids Research. 2018 Jan;46(Database issue):D649–55.

33. Kanehisa M, Furumichi M, Tanabe M, Sato Y, Morishima K. KEGG: new perspectives on genomes, pathways, diseases and drugs. Nucleic Acids Research. 2017 Jan;45(D1):D353–61.

34. Wang X, Kang DD, Shen K, Song C, Lu S, Chang LC, et al. An R package suite for microarray meta-analysis in quality control, differentially expressed gene analysis and pathway enrichment detection. Bioinformatics. 2012 Oct 1;28(19):2534–6.

35. Toro-Domínguez D, Villatoro-García JA, Martorell-Marugán J, Román-Montoya Y, Alarcón-Riquelme ME, Carmona-Sáez P. A survey of gene expression meta-analysis: methods and applications. Briefings in Bioinformatics. 2021 Mar 1;22(2):1694–705.

36. Wilkinson B. A statistical consideration in psychological research. Psychological Bulletin. 1951;48(2):156–8.

37. Mosteller F, Fisher RA. Questions and answers. American Statistical Association bulletin. 1948;2:30–1.

38. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. Journal of the Royal Statistical Society Series B (Methodological). 1995;57(1):289–300.

39. Seney ML, Huo Z, Cahill K, French L, Puralewski R, Zhang J, et al. Opposite Molecular Signatures of Depression in Men and Women. Biological Psychiatry. 2018 Jul 1;84(1):18–27.

40. Bylund DB, Reed AL. Childhood and adolescent depression: why do children and adults respond differently to antidepressant drugs? Neurochem Int. 2007 Oct;51(5):246–53.

41. Benton CS, Miller BH, Skwerer S, Suzuki O, Schultz LE, Cameron MD, et al. Evaluating genetic markers and neurobiochemical analytes for fluoxetine response using a panel of mouse inbred strains. Psychopharmacology. 2012 May 1;221(2):297–315.

42. Samuels BA, Leonardo ED, Dranovsky A, Williams A, Wong E, Nesbitt AMI, et al. Global state measures of the dentate gyrus gene expression system predict antidepressant-sensitive behaviors. PLoS One. 2014;9(1):e85136.

43. Breitfeld J, Scholl C, Steffens M, Brandenburg K, Probst-Schendzielorz K, Efimkina O, et al. Proliferation rates and gene expression profiles in human lymphoblastoid cell lines from patients with depression characterized in response to antidepressant drug therapy. Transl Psychiatry. 2016 Nov 15;6(11):e950.

44. Hervé M, Bergon A, Le Guisquet AM, Leman S, Consoloni JL, Fernandez-Nunez N, et al. Translational Identification of Transcriptional Signatures of Major Depression and Antidepressant Response. Front Mol Neurosci. 2017 Aug 8;10:248.

45. Sargin D, Chottekalapanda RU, Perit KE, Yao V, Chu D, Sparks DW, et al. Mapping the physiological and molecular markers of stress and SSRI antidepressant treatment in S100a10 corticostriatal neurons. Mol Psychiatry. 2020 May;25(5):1112–29.

46. Schmidt EF, Warner-Schmidt JL, Otopalik BG, Pickett SB, Greengard P, Heintz N. Identification of the Cortical Neurons that Mediate Antidepressant Responses. Cell. 2012 May 25;149(5):1152–63.

47. Sarkar A, Chachra P, Kennedy P, Pena CJ, Desouza LA, Nestler EJ, et al. Hippocampal HDAC4 Contributes to Postnatal Fluoxetine-Evoked Depression-Like Behavior. Neuropsychopharmacol. 2014 Aug;39(9):2221–32.

48. Korostynski M, Piechota M, Dzbek J, Mlynarski W, Szklarczyk K, Ziolkowska B, et al. Novel drug-regulated transcriptional networks in brain reveal pharmacological properties of psychotropic drugs. BMC Genomics. 2013 Sep 8;14(1):606.

49. Patrício P, Mateus-Pinheiro A, Irmler M, Alves ND, Machado-Santos AR, Morais M, et al. Differential and Converging Molecular Mechanisms of Antidepressants' Action in the Hippocampal Dentate Gyrus. Neuropsychopharmacol. 2015 Jan;40(2):338–49.

50. Wang Y, Jiang H, Meng H, Lu J, Li J, Zhang X, et al. Genome-wide transcriptome analysis of hippocampus in rats indicated that TLR/NLR signaling pathway was involved in the pathogenisis of depressive disorder induced by chronic restraint stress. Brain Res Bull. 2017 Sep;134:195–204.

51. Malik VA, Zajicek F, Mittmann LA, Klaus J, Unterseer S, Rajkumar S, et al. GDF15 promotes simultaneous astrocyte remodeling and tight junction strengthening at the blood-brain barrier. J Neurosci Res. 2020 Jul;98(7):1433–56.

52. Hagihara H, Ohira K, Miyakawa T. Transcriptomic evidence for immaturity induced by antidepressant fluoxetine in the hippocampus and prefrontal cortex. Neuropsychopharmacology Reports. 2019;39(2):78–89.

53. Torres T, Boloc D, Rodríguez N, Blázquez A, Plana MT, Varela E, et al. Response to fluoxetine in children and adolescents: a weighted gene co-expression network analysis of peripheral blood. Am J Transl Res. 2020 May 15;12(5):2028–40.

54. Rajkumar AP, Qvist P, Donskov JG, Lazarus R, Pallesen J, Nava N, et al. Reduced Brd1 expression leads to reversible depression-like behaviors and gene-expression changes in female mice. Transl Psychiatry. 2020 Jul 17;10(1):1–14.

55. Rayan NA, Kumar V, Aow J, Rastegar N, Lim MGL, O'Toole N, et al. Integrative multi-omics landscape of fluoxetine action across 27 brain regions reveals global increase in energy metabolism and region-specific chromatin remodelling. Mol Psychiatry. 2022 Nov;27(11):4510–25.

56. Demin KA, Krotova NA, Ilyin NP, Galstyan DS, Kolesnikova TO, Strekalova T, et al. Evolutionarily conserved gene expression patterns for affective disorders revealed using cross-species brain transcriptomic analyses in humans, rats and zebrafish. Sci Rep. 2022 Dec 2;12:20836.

57. Willner P. The chronic mild stress (CMS) model of depression: History, evaluation and usage. Neurobiol Stress. 2016 Aug 24;6:78–93.

58. Wang QS, Yan K, Li KD, Gao LN, Wang X, Liu H, et al. Targeting hippocampal phospholipid and tryptophan metabolism for antidepressant-like effects of albiflorin. Phytomedicine. 2021 Nov 1;92:153735.

59. Beurel E, Toups M, Nemeroff CB. The Bidirectional Relationship of Depression and Inflammation: Double Trouble. Neuron. 2020 Jul 22;107(2):234–56.

60. Maes M. Major depression and activation of the inflammatory response system. Adv Exp Med Biol. 1999;461:25–46.

61. Miller AH, Maletic V, Raison CL. Inflammation and Its Discontents: The Role of Cytokines in the Pathophysiology of Major Depression. Biol Psychiatry. 2009 May 1;65(9):732–41.

62. Mamdani F, Berlim MT, Beaulieu MM, Labbe A, Merette C, Turecki G. Gene expression biomarkers of response to citalopram treatment in major depressive disorder. Transl Psychiatry. 2011 Jun;1(6):e13.

63. Guilloux JP, Bassi S, Ding Y, Walsh C, Turecki G, Tseng G, et al. Testing the Predictive Value of Peripheral Gene Expression for Nonremission Following Citalopram Treatment for Major Depression. Neuropsychopharmacology. 2015 Feb;40(3):701–10.

64. Raison CL, Rutherford RE, Woolwine BJ, Shuo C, Schettler P, Drake DF, et al. A Randomized Controlled Trial of the Tumor Necrosis Factor-alpha Antagonist Infliximab in Treatment Resistant Depression: Role of Baseline Inflammatory Biomarkers. JAMA Psychiatry. 2013 Jan;70(1):31–41.

65. Fuh SC, Fiori LM, Turecki G, Nagy C, Li Y. Multi-omic modeling of antidepressant response implicates dynamic immune and inflammatory changes in individuals who respond to treatment. PLoS One. 2023 May 15;18(5):e0285123.

66. Gigase FAJ, Smith E, Collins B, Moore K, Snijders GJLJ, Katz D, et al. The association between inflammatory markers in blood and cerebrospinal fluid: a systematic review and meta-analysis. Mol Psychiatry. 2023 Apr;28(4):1502–15.

67. Powell TR, Murphy T, de Jong S, Lee SH, Tansey KE, Hodgson K, et al. The genome-wide expression effects of escitalopram and its relationship to neurogenesis, hippocampal volume, and antidepressant response. American Journal of Medical Genetics Part B: Neuropsychiatric Genetics. 2017;174(4):427–34.

68. Marchetti L, Lauria M, Caberlotto L, Musazzi L, Popoli M, Mathé AA, et al. Gene expression signature of antidepressant treatment response/non-response in Flinders Sensitive Line rats subjected to maternal separation. European Neuropsychopharmacology. 2020 Feb 1;31:69–85.

69. Zhou X, Liao WJ, Liao JM, Liao P, Lu H. Ribosomal proteins: functions beyond the ribosome. Journal of Molecular Cell Biology. 2015 Apr 1;7(2):92–104.

70. Luscher B, Maguire JL, Rudolph U, Sibille E. GABAA receptors as targets for treating affective and cognitive symptoms of depression. Trends Pharmacol Sci. 2023 Sep;44(9):586–600.

71. Funayama Y, Li H, Ishimori E, Kawatake-Kuno A, Inaba H, Yamagata H, et al. Antidepressant Response and Stress Resilience Are Promoted by CART Peptides in GABAergic Neurons of the Anterior Cingulate Cortex. Biological Psychiatry Global Open Science. 2023 Jan 1;3(1):87–98.

72. Duman RS, Sanacora G, Krystal JH. Altered Connectivity in Depression: GABA and Glutamate Neurotransmitter Deficits and Reversal by Novel Treatments. Neuron. 2019 Apr 3;102(1):75–90.

73. Mullard A. FDA approves first oral drug for postpartum depression, but rejects it for major depressive disorder. Nature Reviews Drug Discovery. 2023 Aug 11;22(10):774–774.

74. Wang C, Vegna S, Jin H, Benedict B, Lieftink C, Ramirez C, et al. Inducing and exploiting vulnerabilities for the treatment of liver cancer. Nature. 2019 Oct;574(7777):268–72.

75. Shao S, Zhuang X, Zhang L, Qiao T. Antidepressants Fluoxetine Mediates Endoplasmic Reticulum Stress and Autophagy of Non–Small Cell Lung Cancer Cells Through the ATF4-AKT-mTOR Signaling Pathway. Frontiers in Pharmacology [Internet]. 2022 [cited 2024 Feb 16];13. Available from: https://www.frontiersin.org/journals/pharmacology/articles/10.3389/fphar.2022.904701

76. Steingart AB, Cotterchio M. Do antidepressants cause, promote, or inhibit cancers? J Clin Epidemiol. 1995 Nov;48(11):1407–12.

77. Yousefi MR, Hua J, Sima C, Dougherty ER. Reporting bias when using real data sets to analyze classification performance. Bioinformatics. 2010 Jan 1;26(1):68–76.

78. Nøhr AK, Lindow M, Forsingdal A, Demharter S, Nielsen T, Buller R, et al. A large-scale genome-wide gene expression analysis in peripheral blood identifies very few differentially expressed genes related to antidepressant treatment and response in patients with major depressive disorder. Neuropsychopharmacol. 2021 Jun;46(7):1324–32.

79. Kornstein SG, Schatzberg AF, Thase ME, Yonkers KA, McCullough JP, Keitner GI, et al. Gender Differences in Treatment Response to Sertraline Versus Imipramine in Chronic Depression. AJP. 2000 Sep;157(9):1445–52.

80. Babaaeyan H, Sakhaie N, Sadegzadeh F, Saadati H, Niapour A. Cardiac and hepatic side effects of fluoxetine in male and female adolescent rats. Fundam Clin Pharmacol. 2023 Nov 8;

81. Sramek JJ, Cutler NR. The impact of gender on antidepressants. Curr Top Behav Neurosci. 2011;8:231–49.

82. Mitic M, Brkic Z, Lukic I, Adzic M. Convergence of glycogen synthase kinase 3β and GR signaling in response to fluoxetine treatment in chronically stressed female and male rats. Behav Brain Res. 2017 Aug 30;333:295–303.

83. Belzeaux R, Lin R, Ju C, Chay MA, Fiori LM, Lutz PE, et al. Transcriptomic and epigenomic biomarkers of antidepressant response. Journal of Affective Disorders. 2018 Jun 1;233:36–44.

84. François BL, Zhang L, Mahajan GJ, Stockmeier CA, Friedman E, Albert PR. A Novel Alternative Splicing Mechanism That Enhances Human 5-HT1A Receptor RNA Stability Is Altered in Major Depression. J Neurosci. 2018 Sep 19;38(38):8200–10.

85. Piechota M, Golda S, Ficek J, Jantas D, Przewlocki R, Korostynski M. Regulation of alternative gene transcription in the striatum in response to antidepressant drugs. Neuropharmacology. 2015 Dec 1;99:328–36.

86. Cooper DG, Class CA. DavidGCooper/FLX-MetaDE: FLX-MetaDE [Internet]. Zenodo; 2024 [cited 2024 Feb 16]. Available from: https://zenodo.org/records/10668846

FIGURE CAPTIONS

Figure 1. PRISMA flow diagram of systematic search and screening for gene expression studies assessing fluoxetine treatment.

Figure 2. (A) Demonstration of meta-analysis methods used in this study to synthesize pathway analyses and differential expression analyses. Heatmap on left shows GSEA nominal p-values for four example pathways in five comparisons. Checkboxes indicate which meta-analysis method(s) would identify that pathway as significantly enriched. **(B)** Individual comparisons used in meta-analysis for gene expression signatures of behavioral or clinical response. **(C)** Number of genes differentially expressed (left) and pathways enriched (right) in each of the nine comparisons of samples from responders vs. non-responders (q < 0.05). "Up" indicates greater expression in responders. **(D)** Pathways identified as significantly enriched (q < 0.05) across the nine comparisons by the three meta-analysis methods. **(E)** Percentage of pathways significantly enriched in responders or non-responders from each of the Reactome categories ("Top Level") or KEGG Database. x-axis indicates percentage of pathways from a certain category identified as enriched. **(F)** Density plot showing distribution of Vote Sums across pathways, colored by meta-analysis result. Positive scores indicate enrichment in responders, negative in non-responders **(G)** Heatmap of Normalized Enrichment Score (NES) across comparisons for pathways identified as enriched in any comparison (Fisher's q < 0.05). Positive NES indicates enrichment in responders.

Figure 3. (A) q-value by Max-P and Vote Sums for pathways identified as enriched across the nine comparisons of responders vs. non-responders (q < 0.05 by Max-P). **(B)** Network diagram showing similarity between pathways, with labels provided in *3A*. Pathways sharing more genes are plotted closer together. **(C)** Meta-analysis results for genewise differential expression in eight mouse comparisons. No genes were identified as consistently differentially expressed by Max-P. **(D)** Volcano plot of mouse differential expression meta-analysis results. **(E)** 13 pathways enriched in responders vs. non-responders in two comparisons of samples not treated with fluoxetine (q < 0.05 in each).

Figure 4. (A) Individual comparisons used in meta-analysis for gene expression signatures of fluoxetine treatment effects. **(B)** Number of genes differentially expressed (left) and pathways enriched (right) in each of the nine comparisons of samples from fluoxetine treated vs. control samples (q < 0.05). "Up" indicates increased expression with fluoxetine. **(C)** Pathways identified as significantly enriched (q < 0.05) across the 55 comparisons by the three meta-analysis methods. **(D)** Percentage of pathways significantly affected by fluoxetine treatment from each of the Reactome categories ("Top Level") or KEGG Database. **(E)** Heatmap of Normalized Enrichment Score (NES) across comparisons for pathways identified as enriched in any comparison (Fisher's q < 0.05). Positive NES indicates increased expression with fluoxetine treatment. **(F)** Density plot showing distribution of Vote Sums across pathways, colored by meta-analysis result. **(G)** q-value by Max-P, and Vote Sums for pathways identified as significant across the 55 comparisons of treated vs. untreated (q < 0.05 by Max-P). **(H)** Network diagram showing similarity between pathways, with labels provided in *4G*. Pathways sharing more genes are plotted closer together.

Figure 5. (A) Pathways identified as significantly enriched (q < 0.05) across the 13 comparisons of treatment effects in MDD patients or stressed rodent models. **(B)** Percentage of pathways significantly affected by fluoxetine treatment in MDD patients or stressed rodent models from each of the Reactome categories ("Top Level") or KEGG Database. **(C)** Density plot showing distribution of Vote Sums across pathways, colored by meta-analysis result. Positive scores indicate increased expression with fluoxetine. **(D)** q-value by Max-P and Vote Sums for pathways identified as significant across the 13 comparisons in MDD patients or stressed rodents (q < 0.05 by Max-P), subset to include only those with absolute Vote Sum ≥ 3. **(E)** Network diagram showing similarity between pathways, with labels provided in *5D*. **(F)** Genes identified as differentially expressed (q < 0.05) across the seven comparisons of treatment effects in stressed mice. **(G)** Density plot showing distribution of Vote Sums across genes, colored by meta-analysis result. **(H)** Volcano plot of meta-analysis results for differential expression of fluoxetine vs. control in stressed mice. Genes with absolute vote sum of 7 are labeled.

Figure 6. (A) Comparison of pathway enrichment by treatment for responders and non-responders, by Max-P. Dotted lines indicate q = 0.05, and counts of pathways within each quadrant are provided. **(B)** q-value by Max-P and Vote Sums for pathways identified as significant across the eight comparisons of treatment vs. control in responders (q < 0.05 by Max-P), but consistently *not* significant across eight comparisons of treatment vs. control for non-responders (q > 0.05 by both Max-P and Fisher's method).

Supplementary Figure 1. Overlap of genes included in pathways identified as significant across the nine comparisons of responders vs. non-responders (q < 0.05 by Max-P). (*Top*) All included genes. (*Bottom*) Subset of genes appearing in at least two of the pathways.

Supplementary Figure 2. Overlap of pathways identified for responders vs. non-responders for all nine comparisons, with two sensitivity analyses (meta-analysis with human comparison removed, and meta-analysis with blood comparisons removed). Vote sums are indicated in parentheses.

Supplementary Figure 3. Overlap of genes included in pathways identified as significant across all comparisons of fluoxetine treated vs. untreated samples (q < 0.05 by Max-P).

Supplementary Figure 4. q-value by Max-P and Vote Sums for all pathways identified as significant across the 13 comparisons in MDD patients or stressed rodents (q < 0.05 by Max-P).

Supplementary Figure 5. Number of genes ("DE") or pathways ("GSEA") identified as statistically significant in each meta-analysis, with q < 0.05. Same data are presented in linear (*top*) and log10 scale (*bottom*).

Supplementary Figure 6. Number of pathways identified as statistically significant for the full meta-analysis of treated vs. untreated samples, as well as subset analyses of stressed rodents or depressed patients, and unstressed rodents. Results of Fisher's meta-analysis (*top*) and Max-P (*bottom*).

Supplementary Figure 7. Sensitivity analysis removing DS19 (profiling of 27 brain regions by Rayan *et al.***(55)**). Gene (*left*) and pathway-level (*right*) analyses are presented. Dashed lines indicate q=0.05, diagonal line indicates y=x. Number in each quadrant indicates number of genes or pathways in that quadrant. Results of Fisher's meta-analysis (*top*) and Max-P (*bottom*).

Figure 1

Table 1. 20 data sets included in synthesis. Letters in parentheses in "Comparisons" column are used as reference in subsequent figures.

ID	Comparisons	GEO Series Accession Number	Citation	Platform	Organism	Tissue(s)	Samples	Stress Method / Diagnosis	Fluoxetine Treatment	Drug Response Determination
DS1	(a-d) Cortex (a) Response (b) Treatment (c) Non-responders (d) Responders	GSE28644	Benton *et al.*[41]	Microarray	*Mus musculus*	Cortex	60 (30/group)	N/A	3-weeks 18 mg/kg/day	Open-field test
DS2	(a, c, e, g) Dorsal Dentate Gyrus (b, d, f, h) Ventral Dentate Gyrus (a-b) Reponse (c-d) Treatment (e-f) Non-responders (g-h) Responders	GSE43261	Samuels *et al.*[42]	Microarray	*Mus musculus*	Dorsal dentate gyrus, Ventral dentate gyrus	38 (4 or 8/group)	Corticosterone	>1-week 160 mg/L in drinking water	Novelty suppressed feeding, Forced swim test
DS3	(a-d) Lymphoblastoid Cell Lines (a) Response (b) Treatment (c) Non-responders (d) Responders	GSE83386	Breitfeld *et al.*[43]	Microarray	*Homo sapiens*	Lymphoblastoid cell lines	20 (10/group)	Major Depressive Disorder	3-weeks 0.5 µg/mL	Hamilton Depression Rating Scale
DS4	(a, c, e, g, i) Anterior Cingulate Cortex (b, d, f, h, j) Dentate Gyrus (a-b) Reponse (c-d) Stressed Treatment (e-f) Unstressed Treatment (g-h) Non-responders (i-j) Responders	GSE84183	Hervé *et al.*[44]	Microarray	*Mus musculus*	Anterior cingulate cortex, Dentate gyrus	64 (8/group)	7-weeks unpredictable chronic mild stress	5-weeks 120 mg/L in drinking water	Coat state measurement
DS5	(a-e) Blood (a) Reponse (b) Stressed Treatment (c) Unstressed Treatment (d) Non-responders (e) Responders	GSE84184	Hervé *et al.*[44]	Microarray	*Mus musculus*	Whole blood	32 (8/group)	7-weeks unpredictable chronic mild stress	5-weeks 120 mg/L in drinking water	Coat state measurement
DS6	(a, c, e, g) S100a10 Whole Cell (b, d, f, h) S100a10 TRAP (a-b) Reponse (c-d) Treatment (e-f) Non-responders (g-h) Responders	GSE202172	Sargin *et al.*[45]	RNA-Seq	*Mus musculus*	S100a10 cortical cells	32 (4/group)	7-weeks single-housing	3-weeks 167 mg/L in drinking water	Homecage time spent in shelter zone
DS7	(-) S100a10 Cells (-) Treatment	GSE35761	Schmidt *et al.*[46]	Microarray	*Mus musculus*	S100a10 cortical cells	6 (3/group)	N/A	15-18 days 167 mg/L in drinking water	None
DS8	(-) Glt25d2 Cells (-) Treatment	GSE35763	Schmidt *et al.*[46]	Microarray	*Mus musculus*	Glt25d2 cortical cells	6 (3/group)	N/A	15-18 days 167 mg/L in drinking water	None
DS9	(-) Hippocampus (-) Treatment	GSE42940	Sarkar *et al.*[47]	Microarray	*Rattus norvegicus*	Hippocampus	8 (4/group)	N/A	3-weeks 10 mg/kg/day	None
DS10	(a-d) Striatum (a) Treatment 1 hour (b) Treatment 2 hours (c) Treatment 4 hours (d) Treatment 8 hours	GSE48951	Korostynski *et al.*[48]	Microarray	*Mus musculus*	Striatum	60 (12/group)	N/A	1-dose 20 mg/kg	None
DS11	(-) Dentate Gyrus (-) Treatment	GSE56028	Patrício *et al.*[49]	Microarray	*Rattus norvegicus*	Dentate gyrus	21 (3/group)	6-weeks unpredictable chronic mild stress	2-weeks 10 mg/kg/day	None
DS12	(-) Brain (-) Treatment	GSE86392	Wang *et al.*[50]	RNA-Seq	*Rattus norvegicus*	Hippocampus, Frontal cortex, Pituitary gland	12 (3/group: 1/tissue)	4-weeks of chronic restraint stress	4-weeks 10 mg/kg/day	None
DS13	(-) Glioma (-) Treatment	GSE89873	Malik *et al.*[51]	Microarray	*Rattus norvegicus*	C6 glioma cells (model of astrocytes)	28 (4 or 6/group)	N/A	2-hours 25 µM	None
DS14	(-) Hippocampus (-) Treatment	GSE109445	N/A	RNA-Seq	*Rattus norvegicus*	Hippocampus	12 (3/group)	5-weeks chronic unpredictable stress	1-week 10 mg/kg/day	None
DS15	(-) Prefrontal Cortex (-) Treatment	GSE118668	Hagihara *et al.*[52]	Microarray	*Mus musculus*	Prefrontal cortex	16 (8/group)	N/A	3-weeks 15 mg/kg/day	None
DS16	(-) Dentate Gyrus (-) Treatment	GSE118669	Hagihara *et al.*[52]	Microarray	*Mus musculus*	Dentate gyrus	16 (8/group)	N/A	3-weeks 15 mg/kg/day	None
DS17	(-) Blood (-) Treatment	GSE128387	Torres *et al.*[53]	Microarray	*Homo sapiens*	Whole blood	32 (15-17/group)	Major Depressive Disorder	8-weeks, dosage not specified	None
DS18	(-) Amygdala (-) Treatment	GSE150431	Rajkumar *et al.*[54]	RNA-Seq	*Mus musculus*	Amygdala	48 (8/group)	N/A	2 days, 5 mg/kg/day	None
DS19	(a-aa) 27 Brain Regions *See Supplementary Table 2* (a-aa) Treatment	GSE194289	Rayan *et al.*[55]	RNA-Seq	*Rattus norvegicus*	27 brain regions	212 (~4/group)	N/A	6-weeks 18 mg/kg/day	None
DS20	(-) Hippocampus (-) Treatment	GSE205325	Demin *et al.*[56]	RNA-Seq	*Rattus norvegicus*	Hippocampus	21 (3/group)	12-weeks chronic unpredictable stress	4-weeks 5 mg/kg/day	None

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Supplementary Figure 1

Zoomed in on genes that appear in at least 2 gene sets

Signal Transduction

Immune System

Nonsense Mediated Decay (NMD) independent of the Exon Junction Complex (EJC)

Formation of a pool of free 40S subunits

GTP hydrolysis and joining of the 60S ribosomal subunit

L13a-mediated translational silencing of Ceruloplasmin expression

NF-kappa B signaling pathway

Cell death signalling via NRAGE, NRIF and NADE

Toll Like Receptor 9 (TLR9) Cascade

Toll Like Receptor 7/8 (TLR7/8) Cascade

MyD88 dependent cascade initiated on endosome

TRAF6 mediated induction of NFkB and MAP kinases upon TLR7/8 or 9 activation

Toll Like Receptor 5 (TLR5) Cascade

MyD88 cascade initiated on plasma membrane

Toll Like Receptor 10 (TLR10) Cascade

Extension of Telomeres

Phospholipase C-mediated cascade; FGFR4

C-type lectin receptors (CLRs)

Downstream signaling events of B Cell Receptor (BCR)

Top_Level

- Cell Cycle
- Immune System
- KEGG Database
- Metabolism of proteins
- Metabolism of RNA
- Signal Transduction

0.8

0.6

0.4

0.2

Top_Level

Supplementary Figure 2

Fisher

Max-P

TLR2 Cascade (2), TLR3 Cascade (3), TLR4 Cascade (4), TLR1:TLR2 Cascade (2), TLR6:TLR2 Cascade (2) TRIF-mediated TLR4 signaling (5) Cell-extracellular matrix interactions (1) MyD88-independent TLR4 cascade (4)

Nonsense Mediated Decay independent of the Exon Junction Complex (-4) Phospholipase C-mediated cascade; FGFR4 (0)

Clathrin-mediated endocytosis (1) Removal of the Flap Intermediate (-1)

Supplementary Figure 3

Supplementary Figure 4

Linear x-axis

Log-scale x-axis

Gene Level
(Rat Only)

Pathway Level
(Mouse and Rat)