Sniffing can be initiated by dopamine's actions on ventral striatum neurons

Natalie L. Johnson[1], Anamaria Cotelo-Larrea[1], Lucas A. Stetzik[1], Umit M. Akkaya[2], Zihao

Zhang[1], Marie A. Gadziola[3], Adrienn G. Varga[4], Minghong Ma[5], and Daniel W. Wesson[1,*]

[1]Department of Pharmacology and Therapeutics, Center for Smell and Taste, Center for

Addiction Research and Education; University of Florida College of Medicine, Gainesville,

FL 32610, USA

[2]Department of Computer Engineering, Gebze Technical University, 41400 Kocaeli, Turkey

[3]Department of Psychology, University of Toronto Scarborough, Toronto, ON, Canada

[4]Department of Neuroscience, Breathing Research and Therapeutics Center; University of

Florida College of Medicine, Gainesville, FL 32610, USA

[5]Department of Neuroscience, Perelman School of Medicine, University of Pennsylvania,

Philadelphia, PA 19104, USA

[*]Corresponding author: Email: danielwesson@ufl.edu

Abstract

Sniffing is a motivated behavior displayed by nearly all terrestrial vertebrates. While sniffing is associated with acquiring and processing odors, sniffing is also intertwined with affective and motivated states. The neuromodulatory systems which influence the display of sniffing are unclear. Here, we report that dopamine release into the ventral striatum is coupled with bouts of sniffing and that stimulation of dopaminergic terminals in these regions drives increases in respiratory rate to initiate sniffing whereas inhibition of these terminals reduces respiratory rate. Both the firing of individual neurons and the activity of post-synaptic D1 and D2 receptor-expressing neurons in the ventral striatum are also coupled with sniffing and local antagonism of D1 and D2 receptors squelches sniffing. Together, these results support a model whereby sniffing can be initiated by dopamine's actions upon ventral striatum neurons. The nature of sniffing being integral to both olfaction and motivated behaviors implicates this circuit in a wide array of functions.

Introduction

Sniffing is a volitional behavior displayed by nearly all terrestrial vertebrates and is defined by the rapid inhalation and exhalation of air through the nose. The commonplace function ascribed to sniffing is that it aids in perceiving odors[1,2]. This is supported by studies reporting that rodents increase their respiratory frequency from ~2-4Hz at rest, up into an around 6-12Hz range when sampling an odor[3–6]. In addition to supporting olfaction, sniffing is also intertwined with affective and motivated states wherein sniffing is considered an appetitive behavior[7–9]. For instance, rodents, dogs, and even semi-aquatic vertebrates increase their sniffing frequency while foraging for food and in anticipation of reinforcers in instrumental tasks[5,6,9,10]. The frequency of sniffing also appears to influence emotional states[11,12]. Furthermore, sniffing serves as a "master clock" for essential physiological rhythms including whisking, wherein the rhythm of sniffing is coupled with whisking and also other head movements[13,14]. Therefore, it is clear that sniffing holds a ubiquitous position in nature.

The automatic coordination of respiration, and the tight control of the inspiratory / post-inspiratory / expiratory rhythm, are afforded by brainstem structures which elegantly time breaths, control their depth, and send these instructive signals toward the breathing plant by means of the spinal and cranial nerves[15,16]. While likely many of the respiratory brainstem structures that participate in the control of basal breathing are integral for the orchestration of sniffing, the exact mechanisms and neural basis for sniffing still remain unknown. Although sniffing is associated with increased bursting in the inspiratory rhythm generator, the pre-Bötzinger complex[17], optogenetic stimulation of this region has yet to reproduce the rapid breathing rates that characterize sniffing[18–20], unlike a

63 subset of neurons in the pontine respiratory modulator, parabrachial nucleus[21]. Additionally,

64 subsets of neurons in the expiratory control center, the retrotrapezoid nucleus, increase their

65 activity shortly before the onset of sniffing[22]. These neurons provide input to the pre-Bötzinger

66 complex[23], as well as the facial motor nucleus, another region recognized to play a direct role in

67 controlling sniffing[24]. While these findings together begin to uncover the circuitry for sniffing,

68 more work needs to be done to pinpoint all the components, and specifically the neuromodulatory

69 circuitry, which invigorates an animal to go from basal breathing which serves the purpose of gas

70 exchange, to engage in the voluntary act of sniffing.

72 We reasoned the mesolimbic dopamine (DA) system as a prime candidate for the invigoration of

73 sniffing. Electrical stimulation of the ventral tegmental area (VTA) elicits vigorous sniffing[25].

74 Neurons synthesizing DA in the VTA project into the basal forebrain, including into the ventral

75 striatum. The ventral striatum, considered a limbic-motor interface[26], is well-known to integrate

76 cortical and limbic circuitry to adaptively guide behavior[27,28]. The major subcomponents of the

77 ventral striatum, including the nucleus accumbens (NAc) and the tubular striatum (TuS; also

78 known as the olfactory tubercle) are recipient of dense DAergic inputs from the VTA. Neurons in

79 both the NAc and TuS are largely spiny projection neurons expressing D1- or D2-like receptors

80 which mediate actions of DA[29–31].

82 Through studies of DA's role in the ventral striatum, we have come to learn that DA's actions in

83 the ventral striatum influence a wide-range of behaviors and states, such as aversion behaviors,

84 and also cellular plasticity needed for learning[32–37]. DA in the ventral striatum is also a major

85 mediator of motivated states and reinforcement with manipulation of DA, and D1 and D2 neural

activity in the ventral striatum guiding those functions[36,38–40]. Since sniffing is a motivated

behavior, it is reasonable to believe the ventral striatum's role in motivational control is in-part a

regulatory system which influences the occurrence of sniffing. Indeed, DAergic manipulations in

the ventral striatum also influence exploratory behaviors in open-field and other gross motor

behavior assays and recently it was reported that DAergic inputs regulate whisking[41]. Mesolimbic

DA thus seems a quintessential neural system for supporting sniffing. Here, we investigated the

role of mesolimbic DA in the display of sniffing behavior to find that sniffing can be initiated by

DA's actions upon ventral striatum neurons.

Results

Sniffing frequency is associated with mesolimbic DA release

We initially sought to understand the relationship between sniffing and DA release in the ventral

striatum. To address this, we first performed a tracing study to identify ventral striatum subregions

wherein DA inputs terminate for subsequent monitoring of DA during behavior. It is well known

in both mice and rats that the ventral striatum receives dense input from mesolimbic DAergic

neurons[28,42,43] and that DA dynamics may differ between ventral striatum subregions[44,45]. In

contrast to the innervation of the dorsal striatum and NAc, quantitative analyses of DA input to

the TuS in mice is especially lacking. To address this, we injected AAV1-hSyn-FLEX-mGFP-2A-

Synaptophysin-mRuby into the VTA of *DAT^IRES-Cre* mice (**Fig 1A & Supplementary Fig 1**).

This approach allowed for visualization of fluorescent puncta, which indicate DAergic presynaptic

terminals, throughout the TuS and NAc (**Fig 1B**). The injections resulted in GFP expression

throughout the VTA (**Supplementary Fig 1**). We quantified fluorescent puncta in the NAc Core

(NAcC), NAc Shell (NAcSh), medial TuS (mTuS), and lateral TuS given different roles for these

subregions in the regulation of motivated behaviors (**Fig 1C**)[46–49]. As expected, we observed robust

fluorescence, indicating synaptophysin+ DAT VTA terminals, in the TuS and NAc, most notably

in the anterior aspect of the mTuS and NAc (**Fig 1C**) with 113% and 78% more innervation in the

anterior versus posterior halves of the mTuS and NAcSh, respectively (the anterior or posterior-

most 200μm were used for quantification of these regions). Following this we focused analyses on

the anterior aspect of both structures. While the mTuS was rich with synaptophysin+ DAT VTA

terminals, this contrasted to the lateral TuS which was largely void of innervation (**Fig 1C & 1D**,

rmANOVA $F(1.52, 7.6)=31.74$ $p=0.0003$, Tukey's multiple comparisons ant mTuS vs. ant LatTuS

$p=0.014$). Comparable amounts of synaptophysin+ DAT VTA terminals were found in the anterior

NAcSh and anterior NAcC (**Fig 1C & 1D**, Tukey's multiple comparisons antNAcSh vs. antNAcC,

$p=0.073$). The density of input across the anteromedial aspects of the TuS, NAcSh, and NAcC

were all similar (**Fig 1C & 1D;** Tukey's multiple comparisons anterior mTuS vs. anterior NAcSh

$p=0.217$; anterior mTuS vs. anterior NAcC $p=0.878$). Together this work adds to literature

indicating that the anterior NAcSh, NAcC, and mTuS are the major recipients of VTA DAergic

input in the ventral striatum and provide quantitative support for focusing on the medial aspects of

these regions in mice.

Having established this, we next sought to investigate the temporal relationship between

mesolimbic DA release and sniffing behavior. We took advantage of whole-body

plethysmography to assess respiratory cycles in awake mice and identify bouts of sniffing. In this

method, breaths are detected by a pressure transducer which accesses the inside of the chamber

(**Fig 2A**). The technical needs of our experiments require a patch cable attached to the mouse in

the chamber (*e.g.,* **Fig 2C**) which may impair fidelity of the respiratory signal since the cable

would need to exit the plethysmography chamber which is normally air-tight. Further, we wished

to identify temporal relationships between sniffing and other signals (DA, photostimulation, *etc*)

and this mandates that first we establish a temporal relationship between breaths detected with

whole-body plethysmography and breathing of mice which occurs only through their noses since

they are obligate nasal air breathers. Therefore, we simultaneously monitored respiration via the

plethysmograph as well as by means of a chronically-implanted cannula into the dorsal nasal recess

of awake mice (**Fig 2A**)[6]. For the intranasal measures, a flexible tube was connected to the nasal

cannula which passed through the top of the plethysmograph via a snug 3D-printed fitting which

allowed for mice to explore and freely-rotate within the plethysmograph. The end of this tube

terminated into a pressure transducer for amplification of intranasal pressure changes. Using this

simultaneous approach, we confirmed in three mice that whole-body plethysmography, even with

the patch cable freely passing through the chamber, reliably detected intranasal respiratory cycles,

and faithfully followed during high frequency sniffing bouts (**Fig 2Bi & 2Biii**). The only notable

difference we observed for the purpose of this study was a slight temporal lag with the peaks of

the whole-body plethysmograph signal trailing those of the intranasal signal (**Fig 2Bii**). This lag

is anticipated since the origin of the signal is from within the mouse (intranasal) versus from within

a chamber (plethysmograph). Across the three mice as they spontaneously transition in and out of

sniffing bouts, the mean lag ranged from 0.028 ± 0.007s to $.036 \pm 0.0004$s (using data with

respiratory rates ranging from ~2-12Hz, as in **Fig 3Biii**). While solely at resting respiration (2-

4Hz), the lag between the two signals increased and ranged from 0.053 ± 0.001s to 0.10 ± 0.001s

(across-animal range). Nevertheless, in events wherein respiratory cycles were detected in the

nose, in >99% of the cases, they were also detected by the plethysmograph (3,239 breaths across

three mice, a restricted range of data from one mouse shown in **Fig 2Bii**).

Having confirmed the ability to accurately detect both individual respiratory cycles and sniffing

bouts alike, we next monitored the relationship between DA levels in the ventral striatum as they

relate to sniffing. We injected mice into either the medial TuS, the NAcSh, or NAcC with an AAV

encoding $GRAB_{DA}$ (pAAV.hSyn-GRAB_DA1h)[50] and in the same surgery implanted into the

structure receiving the AAV, a 400µm optical fiber for later fiber photometry. Other than for the

purpose of showing example traces, for other analyses we subtracted the isosbestic 405nm signal

from the $GRAB_{DA}$ signal. Following several weeks for surgical recovery and viral expression, we

placed the mice in a plethysmograph for simultaneous monitoring of respiration and DA levels

(**Fig 2C**). Since mice rapidly transition in and out of sniffing bouts[6], we restricted our analyses to

more stable/pronounced bouts versus short bouts by only defining a sniff bout as the occurrence

of sniffing $\geq$6Hz for $\geq$1s in duration. Further, in some instances, mice continuously sniff for

numerous seconds and then reengage in another bout shortly thereafter. Since these may reflect

periods of more generalized arousal versus spontaneous exploratory behavior, we excluded sniff

bouts occurring in proximity to these long (>10s) sniffing bouts. We analyzed the relationship

between DA levels and sniffing in several manners, and in all analyses, we accounted for the lag

between when a nasal inhalation occurs and when the plethysmograph detects the inhalation by

subtracting the conservative 28ms lag (see **Fig 2Bii**) from the onset event of every peak-detected

respiratory cycle.

We analyzed DA levels across trials wherein mice in the plethysmograph spontaneously sniffed,

were delivered odors to evoke sniffing, or were delivered an auditory/somatosensory stimulus by

connecting a vibration motor to the plethysmograph which modestly resonated the chamber while

simultaneously providing a small auditory 'buzz' cue due to its motor (**Supplementary Fig 2**). All

mice were confirmed to have implants extending into the mTuS, NAcC, or the NAcSh (**Fig 2D**).

First, we looked at the relationship between spontaneous sniffing like that displayed by mice

during self-motivated states of exploration. As shown in the example traces from a mouse

implanted into the TuS (**Fig 2E**), whereas no detectable changes in DA levels were observed when

mice were simply breathing at baseline frequencies ('respiration at rest'; **Fig 2Ei &**

Supplementary Fig 3A), spontaneous sniffing was associated with elevations in the $GRAB_{DA}$

signal (**Fig 2Eii**). Across mice, $GRAB_{DA}$ levels during spontaneous sniffing were elevated in both

the TuS and the NAcSh, but in contrast were decreased in the NAcC (**Fig 2Fi, 2Gi, 2Hi**; TuS

paired $t(6)=11.85$ $p<0.0001$; NAcSh paired $t(6)=6.56$, $p=0.0006$; NAcC paired $t(6)=12.15$,

$p<0.0001$). Only in the TuS, was the average frequency of the spontaneous sniffing bout correlated

with the amplitude of the $GRAB_{DA}$ signal (**Fig 2Fii, 2Gii, & 2Hii**; TuS Pearson's $r(83)=0.29$,

$p=0.008$), but not in the NAcSh or NAcC (NAcSh Pearson's $r(101)=0.12$, $p=0.2493$; NAcC

Pearson's $r(102)=0.038$, $p=0.703$). Notably, neither the isosbestic 405nm signal acquired from

these same animals, nor the DA-insensitive control virus, $GRAB_{DA}$-mut, acquired in a separate

cohort of 9 mice, showed comparable changes during spontaneous sniffing (**Supplementary Fig**

4).

Next, we investigated the relationship between odor-evoked sniffing and DA levels (**Fig 3A**). Each

mouse was presented with pseudorandom trials of odors, including R(+)-limonene, thioglycolic

acid, and peanut oil, each five times across a behavior session. These odors were selected based

upon their known positive hedonic qualities perceived by mice, in order to elicit vigorous sniffing

[51–53]. As shown in the example in **Figure 3B** from a mouse implanted in the TuS, the first

presentation of thioglycolic acid was associated with robust increases in sniffing, which was

accompanied by an increase in the $GRAB_{DA}$ signal. Across all groups (TuS, NAcSh, NAcC),

sniffing frequency was elevated compared to baseline for the first presentation of odors (TuS

paired $t(6)=19.23$, $p<0.0001$; NAcSh paired $t(6)=13.27$, $p<0.0001$; NAcC paired $t(6)=14.35$,

$p<0.0001$), which habituated across subsequent presentations (TuS rmANOVA

$F(2.26,13.55)=14.64$, $p=0.0003$; NAcSh rmANOVA $F(2.44, 14.65)=32.65$, $p<0.0001$; NAcC

rmANOVA $F(1.59, 9.55)=23.5$, $p=0.0003$) (**Figs 3Ci, 3Di, & 3Ei**). Simultaneously during novel

odor-evoked sniffing bouts (trial #1 of odor), we observed elevations in the $GRAB_{DA}$ signal in

both the TuS and NAcSh (**Fig 3Cii & 3Dii**; TuS paired $t(6)=6.54$, $p=0.0006$; NAcSh paired

$t(6)=4.38$, $p=0.005$). The increases rose soon around sniff bout onset and decayed within a few

seconds. In contrast, the NAcC displayed reduction in the $GRAB_{DA}$ signal upon sniffing (**Fig 3Eii**;

paired $t(6)=8.35$, $p=0.0002$).

The elevation in the $GRAB_{DA}$ signal was not solely due to odor input since the signal became

attenuated across repeated odor deliveries in the TuS and NAcSh **(Fig 3Ciii & 3Diii**; trials #1-5;

TuS rmANOVA $F(2.08, 12.48)=7.942$, $p=0.006$; NAcSh rmANOVA: $F(1.90,11.39)=5.563$,

$p=0.022$), whereupon trial 5, the $GRAB_{DA}$ signal was observed near baseline levels. In the NAcC,

reductions in the $GRAB_{DA}$ signal was not relieved with repeated odor presentations (**Fig 3Eiii**;

rmANOVA $F(1.98, 11.90)=0.5760$, $p=0.58$). Also supporting that the change in amplitude of the

$GRAB_{DA}$ signal is related to sniffing, we found that the frequency of odor-evoked sniffing

correlated with the peak of the $GRAB_{DA}$ signal in both the TuS and NAcSh (TuS Pearson's

$r(34)=0.65$, $p<0.0001$; NAcSh Pearson's $r(34)=0.51$, $p=0.002$), but not in the NAcC (Pearson's

$r(34)=-0.14$, $p=0.414$) (**Figs 3Civ, 3Div, & 3Eiv**). Interestingly here, the odor-evoked sniffing in

the NAcSh was correlated with GRAB$_{DA}$ levels which was not the case with spontaneous sniffing

(**Fig 2G**), suggesting in the NAcSh the relationship may be dependent upon odor.

Finally, we assessed the impact of the multimodal buzz stimulus on GRAB$_{DA}$ levels. While not as

dramatic nor reliable as an odor (as in **Fig 3Ci, 3Di, & 3Ei**), the buzz stimulus evoked sniffing

greater than baseline in the majority of mice (**Fig 3Fi, 3Gi, & 3Hi**). Correspondingly, DA was

elevated upon initial buzz-evoked sniffing in the TuS and the NAcSh (**Figs 3Fii & Gii**; TuS paired

$t(6)=6.77$, $p=0.0005$; NAcSh paired $t(6)=3.17$, $p=0.019$). The buzz-evoked sniffing did not

strongly habituate across repeated trials, and likewise, GRAB$_{DA}$ levels also remained consistent

across trials (**Fig 3Fi & 3Fiii**, **Fig 3Gi & 3Giii**; TuS sniffing rmANOVA $F(2.57, 15.4)=1.351$,

$p=0.293$; TuS GRAB$_{DA}$ rmANOVA $F(1.90, 11.38)=0.49$, $p=0.616$; NAcSh sniffing rmANOVA

$F(2.43, 14.58)=2.97$, $p=0.0752$; NAcSh GRAB$_{DA}$ rmANOVA $F(1.32, 7.94)=0.27$, $p=0.6786$).

Interestingly, in the NAcSh, there was a biphasic increase in GRAB$_{DA}$ with the first phase

occurring upon buzz onset, and the second upon buzz offset (**Fig 3Gii**). Whereas in the NAcSh,

GRAB$_{DA}$ levels measured upon either phase were not correlated with the frequency of sniffing

(**Fig 3Giv**; during buzz Pearson's $r(34)=-0.17$, $p=0.316$; upon buzz offset Pearson's $r(34)=-0.074$,

$p=0.673$), the GRAB$_{DA}$ levels upon buzz strongly correlated with frequency of sniffing in the TuS

(**Fig 3Eiv**; Pearson's $r(34)=0.43$, $p=0.011$). Similar to that seen for both spontaneous sniffing and

odor-evoked sniffing, GRAB$_{DA}$ levels in the NAcC were reduced upon buzz-evoked sniffing (**Fig**

3Hii; paired $t(6)=3.23$, $p=0.018$) which similarly did not habituate across trials (**Fig 3Hii & 3Hiii**;

rmANOVA $F(2.19, 13.14)=0.44$, $p=0.668$). There was no correlation between buzz-evoked

sniffing frequency and GRAB$_{DA}$ levels in the NAcC (**Fig 3Hiv**; Pearson's $r(34)=-0.22$, $p=0.194$).

When comparing the peak GRAB$_{DA}$ responses across all regions and all conditions, we found

similar evoked responses in the TuS and NAcSh to novel odor (trial #1 of odor) presentations and

during spontaneous sniffing (**Fig 3I**; two-way rmANOVA, main effect of region $F(2,18)=58.45$,

$p<0.0001$; Tukey's multiple comparisons TuS vs. NAcSh odor $p=0.99$; TuS vs. NAcSh spont

$p=0.219$). As with spontaneous sniffing, neither odor- nor buzz-evoked sniffing resulted in

comparable changes in the separate cohort of 9 mice wherein we recorded GRAB$_{DA}$-mut signal

(**Supplementary Fig 4**). Taken together, these results indicate that sniffing often corresponds with

increases in DA release in two specific sub-regions of the ventral striatum, and that the level of

DA correlates to the frequency of sniffing.

DA release into the ventral striatum elicits sniffing

Given the relationship between DA release in the ventral striatum and the occurrence and

frequency of sniffing, we next sought to determine whether this release of DA is sufficient to elicit

sniffing. To assess this, we injected a Cre-dependent Channelrhodopsin2 (AAV5-EF1α-DIO-

hChR2(E123T/T159C)-EYFP) unilaterally into the VTA of *DAT^{IRES}-Cre* mice (**Supplementary**

Fig 5) and implanted a 300μm optical fiber terminating over the TuS, NAcSh, or NAcC to drive

terminal release of DA upon light stimulation (**Fig 4A & 4B**). Control mice were injected with

AAV solely expressing EYFP in the VTA (AAV5-EF1α-DIO-eYFP) and also received an optical

fiber terminating over the TuS, NAcSh, or NAcC. One second long, 25Hz 473nm light stimulation

was delivered through the optical fiber while mice were in the plethysmograph. This 25Hz stimulus

was selected based upon prior work using a similar stimulus paradigm to elicit VTA DA release

into the ventral striatum[39,40,54]. As shown in **Figure 4C**, in an example mouse, photostimulation of

DAergic terminals in the TuS evoked a brief bout of sniffing. In this example mouse, the average

latency to onset of a sniff bout was 240 ± 22.5ms (across trial range: 130-330ms; bout onset defined

as the moment of achieving $\geq$6Hz respiration). Further, the average amount of time spent sniffing

(time $\geq$6Hz) during photostimulation was 672 ± 59.6ms (across trial range: 310-1000ms). The light

evoked sniffing in this example mouse was also reliable across repeated trials (**Fig 4C**).

In all three ChR2-injected groups of mice, we observed a rapid increase in respiratory frequency

upon photostimulation (**Fig 4D**). This photostimulation-evoked display of sniffing was largely

maintained across repeated trials (**Fig 4D & 4E;** testing for stability in Hz across trials: mixed-

effects analysis $F(4.09, 85.87)=2.34$, $p=0.058$). While anticipated, EYFP control mice displayed a

slight increase in respiratory frequency during photostimulation, likely due to the arousing nature

of the visual stimulus leaking from the fiber implant. Alternatively, given that light stimuli can

evoke DA release in the ventral striatum[55] it is also possible that light-triggered DA release could

have contributed to the increase in respiratory frequency seen in EYFP controls. Nonetheless, the

frequency of the sniffing bout was lower in EYFP mice than in all ChR2-treated groups (**Fig 4D,**

right, Fig 4E & F). As shown in **Figure 4F**, the temporal dynamics of photostimulation-evoked

sniffing in ChR2-injected mice was remarkably similar. The average latency to onset of a sniff

bout was 279.4 ± 13.3ms SEM (inter-mouse range: 225-362.5ms) for TuS, 264.7 ± 14.83ms SEM

(inter-mouse range: 202.5-299ms) for NAcSh, and 263.3 ± 7.52ms SEM (inter-mouse range:

242.2-290ms) for NAcC (**Fig 4Gi**). The average latency to the onset of a sniff bout in EYFP

controls (339.4 ± 18.72ms SEM, inter-mouse range: 274.2-405ms) was significantly greater than

in the ChR2-treated groups (all groups with means <290ms as in **Fig 4Gi;** one-way ANOVA $F(3,$

$25)=5.11$, $p=0.007$, Tukey's multiple comparisons test TuS vs. EYFP $p=0.0273$, NAcSh vs. EYFP

$p=0.0141$, NAcC vs. EYFP $p=0.122$). Further, the average amount of time spent sniffing during

photostimulation was 534.1 ± 37.8ms (inter-mouse range 314.3-743ms) for TuS, 647.7 ± 53.7ms

(inter-mouse range 479-834ms) for NAcSh, and 520.1 ± 85.7ms (inter-mouse range 254.4-755ms)

for NAcC. All ChR2-treated mice spent more time sniffing than EYFP controls, and there were no

significant differences between ChR2-treated groups (**Fig 4Gii**) (one-way ANOVA

$F(3,28)=14.02$, $p<0.0001$; Tukey's multiple comparisons test TuS vs. EYFP $p=0.0001$, NAcSh vs.

EYFP $p<0.0001$, NAcC vs. EYFP $p=0.0017$). Further, the frequency of photostimulation-evoked

sniffing (measured as the peak Hz during photostimulation) did not differ between ChR2-treated

groups and was significantly greater than EYFP controls (**Fig 4Giii**; one-way ANOVA $F(3,$

$28)=14.17$, $p<0.0001$; Tukey's multiple comparisons test TuS vs. EYFP $p<0.0001$, NAcSh vs.

EYFP $p<0.0001$; NAcC vs. EYFP $p=0.0010$). Finally, we explored the reliability whereby a

DAergic terminal photostimulation would elicit a sniffing bout. The probability that

photostimulation elicited sniffing was robust, and significantly greater than in EYFP controls –

indeed with many animals in each ChR2-treatment group displaying a sniff bout to *every*

photostimulation trial and the vast majority to >80% of trials (**Fig 4Giv**; one-way ANOVA $F(3,$

$28)=11.01$, $p<0.0001$; Tukey's multiple comparisons TuS vs. EYFP $p=0.0002$, NAcSh vs. EYFP

$p=0.0002$, NAcC vs. EYFP $p=0.004$).

Since VTA neurons are known to co-release glutamate in addition to dopamine[35,56,57], and knowing

that ventral striatum neurons express glutamatergic receptors, we next sought to isolate the

influence of the above effect (**Fig 4**) on DA versus glutamate. We crossed *VGluT2 fl/fl* mice with

mice expressing Cre recombinase in all cells expressing tyrosine hydroxylase (*TH-Cre*), the rate-

limiting enzyme upstream for DA synthesis. This approach ensures that *VGluT2* is lacking from

tyrosine hydroxylase expressing neurons, thus eliminating the possibility of glutamate co-release

from TH+ neurons. As above, we injected *VGluT2fl/fl* x *TH-Cre* mice with the same ChR2-

expressing virus or associated control virus unilaterally into the VTA and implanted an optical

fiber terminating over the TuS to drive terminal excitation. Since photostimulation in all three

ventral striatum subregions of ChR2-expressing mice in the prior experiment drove sniffing, here

we excluded the NAc groups for simplicity. As shown in **Supplementary Figure 6**,

photostimulation of DAergic terminals in *VGluT2fl/fl* x *TH-Cre* mice elicited sniffing with

comparable temporal dynamics, frequency, and probability as in *DAT^{IRES}-Cre* mice (**see Fig 4**) and

at greater amounts in *VGluT2fl/fl* x *TH-Cre* mice than in EYFP controls.

We next wanted to determine whether DA release in the ventral striatum serves to 'initiate'

sniffing, or if it further can persistently drive sniffing over extended time courses. To test this, in

the same ChR2-treated *DAT^{IRES}-Cre* mice as in **Figures 4B-G,** we delivered repeated presentations

of the 25Hz photostimulation for either 120ms, 1s, 2s, or 4s. We predicted that if DA initiates

sniffing we would see transiently-evoked sniffing across all stimulation durations, whereas if DA

drives sniffing, we would see the duration of sniffing bouts track the duration of the

photostimulation. As shown in **Figure 4H**, we found that the dynamics of the sniffing bout were

equivalent regardless of photostimulation duration. Indeed, in all three ChR2-treated groups,

photostimulation of DAergic terminals elicited a statistically similar frequency of sniffing (TuS

mixed-effects analysis; $F(1.78, 6.53)=0.94$ $p=0.426$; NAcSh rmANOVA $F(1.63, 4.89)=0.924$,

$p=0.436$; NAcC rmANOVA $F(1.93, 5.79)=0.24$, $p=0.789$), and similar percentage of time sniffing

(TuS mixed-effects analysis $F(1.18, 4.32)=1.57$, $p=0.284$; NAcSh rmANOVA $F(1.50, 4.5)=0.77$

$p=0.479$; NAcC rmANOVA $F(1.43, 4.30)=1.81$, $p=0.404$). These results indicate that DA release

into the ventral striatum initiates sniffing, but that it does not persistently drive a sniff bout.

Stimulation of respiratory brain stem regions known to support breathing can generate sniffing patterns even in anesthetized mice (*e.g.* [15,16,58]). Is mesolimbic DA input to the ventral striatum capable of initiating sniffing even in anesthetized mice? To test this, following the above experiments, *DAT^IRES-Cre* mice injected with ChR2 were deeply anesthetized with urethane and photostimulation was delivered. The baseline respiratory pattern and rate in these mice were both stable and slow (<1.25Hz), indicating deep sedation. Breathing rate under anesthesia showed no change upon photostimulation of DAergic terminals in either the TuS or the NAcSh and NAcC (**Fig 5A - C**), suggesting that the VTA→ventral striatum DAergic circuit is a component of the voluntary, not automatic, respiration-control system.

Sniffing bouts can be displayed in response to a variety of stimuli an animal may perceive in its environment. While not well-studied, it is possible the temporal dynamics of sniffing bouts may differ based upon what elicited the sniffing. We wanted to know if the sniffing bouts triggered by DA release into the ventral striatum are similar, or distinct from, those which are elicited in response to a stimulus or those bouts spontaneously generated. To address this, we used a machine learning approach (**Fig 5D**). We converted 1s long sniffing traces (the 1s following bout onset) into 1sec long instantaneous sniffing frequency arrays (10ms bin size). Next, we used K-nearest neighbors with Diffusion Time Warping (DTW)[59–61] to calculate the difference in sniffing frequency dynamics between frequency arrays. With DTW, the smaller the value reported in each time bin corresponds to the dynamics between arrays being more similar at that moment. The model was trained on DTW arrays from odor-evoked, spontaneously-generated, and photostimulation-evoked data (*n*=63-70 arrays/group). Following, we fed the model DTW arrays

from each group to test how distinct the sniffing dynamics are between groups. In addition to using

the complete 'real' data set, we generated synthetic data using SMOTE (synthetic minority

oversampling technique[62]) to balance the sample numbers to 70 and separately, tested the model's

classification precision when providing it balanced samples of reduced numbers of 40 or 60 (**Fig**

5E). We found that photostimulation-evoked sniffing was highly distinct from all other types of

sniffing. In all runs, precision in classifying opto-evoked sniffing even exceeded the precision of

classifying odor-evoked sniffing, which is known to be highly stereotyped (**Figs 5Ei & 5Eii**). As

a comparison, spontaneously evoked sniffing was the most poorly classified, likely due to the

variability in these self-motivated bouts. While buzz-evoked sniffing was classified better than

spontaneous, the model struggled to classify it as well as opto-evoked sniffing. Both SMOTE and

the 60 sample runs yielded robust classification of photostimulation-evoked sniffing, with F1

scores (a single metric of model performance) of 0.9 in the 60-sample run, and 1.0 in the SMOTE

run. This result implies that DA release into the ventral striatum engages a distinct respiratory

pattern from that when an animal engages in self-motivated exploration or upon perception of an

environmental stimulus (*i.e.*, an odor).

As an additional experiment, we used an inhibitory optogenetic approach to suppress DAergic

terminals in either the NAcSh or the TuS to test whether this achieves reductions in sniffing. We

targeted these two ventral striatum subregions based upon their prominent elevations in DA upon

sniffing (**Figs 2 & 3**). We injected a Cre-dependent halorhodopsin (AAV5-EF1α-DIO-eNpHR 3.0-

EYFP) bilaterally into the VTA of *DAT^{IRES}-Cre* or *DAT^{IRES}-Cre x Ai9* mice (**Supplementary Fig**

7) and implanted bilateral 400µm diameter optical fibers terminating over the TuS or NAcSh (**Fig**

6A & 6B) to suppress terminal release of DA upon light stimulation. An example of a bilateral

cannula implant in the TuS is shown in **Figure 6C**. We sought to understand if suppressing

DAergic terminals could reduce sniffing both when mice spontaneously engage in sniffing, and

when mice sniff in response to a stimulus. To target spontaneous sniffing bouts, 5s long, 560nm

constant light stimulation was delivered through the optical fibers every 20s while mice were in

the plethysmograph. Since spontaneous bouts of sniffing can sometimes be short (~several

hundreds of milliseconds), we provided light every 20s without considering if the animal was

engaging in a sniff bout or not. We reasoned this would be a more objective stimulation paradigm

than if we were to trigger light off of every bout onset (*e.g.*, in a closed loop paradigm) which

could result in some 'learning' of the mouse to change its behavior.

As shown in the example in **Figure 6Di**, 560nm light stimulation suppressed sniffing frequency

during spontaneous events. Indeed, compared to control mice which received 5s long, 405nm

constant light stimulation (light stimulation unable to activate eNpHR[63]), photoinhibition of

DAergic terminals in the ventral striatum modestly suppressed respiratory frequency (**Fig 6Dii**; 2-

way rmANOVA, effect of light stimulation $F(1, 10)=11.08$, $p=0.0076$; Sidak's multiple

comparisons 560nm before light vs. during light $p=0.0037$). To test effects of DAergic inhibition

on stimulus-evoked sniffing, each mouse was presented with pseudorandom trials of odors,

including R(+)-limonene, thioglycolic acid, and peanut oil, each five times across a behavior

session. 560nm or 405nm light was delivered concurrently with odor and buzz delivery. For this

we used a longer light duration (7s, constant) to ensure that light was on upon the possible moment

the animal may perceive the stimulus and would typically engage in sniffing. Whereas inhibition

of DAergic terminals in the ventral striatum suppressed spontaneous sniffing (**Fig 4D**), there was

no effect of photoinhibition on the frequency of odor- or buzz-evoked sniffing in the TuS or NAcSh

(**Figs 6E & 6F**, two-way rmANOVA: NAcSh odor no effect of light $F(1, 4)=1.009$, $p=0.3719$;

NAcSh buzz no effect of light $F(1, 4)=1.117$, $p=0.3502$; TuS odor no effect of light $F(1, 5)=0.025$,

$p=0.8816$; TuS buzz no effect of light $F(1, 5)=4.696$, $p=0.083$). Taken together with the results of

the optical stimulation, these results cumulatively support that DA release in the ventral striatum

elicits sniffing.

Ventral striatum neural activity is associated with sniffing

VTA DA is released onto spiny projection neurons in the ventral striatum expressing either the D1

or D2 DA receptor. Is the activity of these neurons associated with sniffing? To test this, we

performed two different experiments. First, we took advantage of an existing data set[64] wherein

extracellular neural activity was recorded from the TuS and respiration simultaneously sampled

from an intranasal cannula in behaving c57bl/6j mice throughout their engagement in an olfactory

discrimination task. In this task, mice were head-fixed which allowed for the snout to remain stable

for precision in the odor delivery. A subset of animals contributed respiratory data, and of them,

three mice also each contributed multiple single-units ($n=20$ total). As expected from head-fixed

animals[65,66], we observed bouts of sniffing in response to odor delivery and also during the inter-

trial interval separating odors (variable 15-17sec window). Further, as expected in head-fixed

animals and also seen in mice in the plethysmograph (*e.g.,* **Fig 3**), odor delivery significantly

increased the median of the instantaneous respiratory frequency (Wilcoxon rank-sum test, $z=-19.8$,

rank sum=3.2×10^7, $p<0.0001$, **Fig 7C**).

As shown in **Figures 7A & B**, we found that the firing of ventral striatum neurons occurs in

relation to both bouts of sniffing and even proximal to individual breaths which consisted of a

short duration respiratory cycle (*viz.*, "sniffs"). Each of the neurons in this example displayed a

significant increase in firing in the 200ms following either a sniff or a sniff bout (**Fig 7B**; unit 1

$t(27)=5.44$, $p<0.0001$; unit 2 $t(27)=2.29$, $p=0.049$; unit 3 $t(27)=-3.46$, $p=0.001$). Looking across

the entire population of neurons, 19/20 (including the three from **Fig 7B**) were significantly

modulated relative to a short respiratory cycle ($p<0.05$, 2-tailed *t*-tests of firing rate 200ms before

vs. 200ms during sniffing across sniff occurrences). Among these, they were more likely to fire

within a short respiratory cycle during the inter-trial interval when the background respiratory rate

is slower (**Fig 7D**, mixed ANOVA, F(3, 76)=31.48, $p<0.0001$).

We then explored the relationship between the probability of a change in firing frequency relative

to either the first or last short duration breath in a bout of sniffing. To do this we identified

respiration cycles containing an 'initial spike' and three respiration cycles before and after that

initial spike. We observed that the initial spike occurs before the respiration cycle gets shorter (**Fig**

7Ei). This was especially the case during background breathing in the inter-trial interval when

breathing is slower, wherein cycles +1, 2, and 3 (relative to the initial spike) were significantly

higher in frequency than cycle 0 (ANOVA $F(6, 126)=6.37$, $p<0.0001$), while cycle 0 did not differ

from previous cycles. In the odor presentation period, however, although ANOVA analysis

showed there was a significant main effect of respiration cycles (ANOVA $F(6, 112)=4.43$,

$p=0.0005$), post hoc comparison showed that cycle 0 was not significantly different from the

adjacent cycles. Finally, we performed an additional analysis to determine if the last action

potential in a burst was "locked" to the cessation of a sniff bout. **Figure 7Eii** shows that there is

no change in respiration cycles aligned to the last spike, whether in the inter-trial interval or odor

presentation period. Thus, ventral striatum neural activity appears closely orchestrated with the

occurrence of sniffs.

As a second experiment to example the relationship between sniffing and recruitment of post-

synaptic ventral striatum neurons, we injected a Cre-dependent AAV expressing GCamP6f

(AAV.Syn.Flex.GCaMP6f.WPRE.SV40) into the TuS of D1-Cre and D2-Cre mice[67] which also

were subsequently implanted with a 400µm core optical fiber over the TuS (**Fig 8A & B**). An

extra-nasal flow sensor was positioned near the animal's nose to monitor sniffing[68]. This allowed

us to extend the results of the extracellular recordings (though they provided greatest temporal

resolution) by analyzing specifically D1 and D2 ventral striatum neuron activity across trials

wherein mice spontaneously entered into bouts of sniffing and also during odor-evoked sniffing.

Similar to that observed with DA release into the ventral striatum (**Fig 2**), the activity of both D1

and D2 neurons increased upon spontaneous sniffing bouts (**Fig 8C**). D1 and D2 neuron evoked

responses were comparable in their temporal dynamics, and both groups displayed similar evoked

frequency of sniffing (**Fig 8C**; unpaired $t(16)=1.084$, $p=0.294$). The activity of D2 neurons also

positively correlated with the frequency of spontaneous sniffing (**Fig 8Civ & 8Cv**) (D1 Pearson's

$r(37)=0.19$, $p=0.241$; D2 Pearson's $r(68)=0.48$, $p<0.0001$).

Novel odor delivery (*viz.*, the first presentation of a given odor) evoked sniffing in both D1- and

D2-Cre mice (paired $t(11)=11.58$, $p<0.0001$), which habituated across subsequent presentations of

the same odor (**Fig 8Di**) (rmANOVA $F(1.73, 19.07)=35.82$, $p<0.0001$). During novel odor-evoked

sniffing bouts, D1 and D2 neurons increased in their activity (**Fig 8Dii & 8Div**); D1-Cre: paired

$t(9)=7.03$, $p<0.0001$; D2-Cre: paired $t(9)=9.88$, $p<0.0001$). Reminiscent of changes in DA release

across repeated trials of odor-evoked sniffing (**Fig 3**), the activity of D1 and D2 neurons similarly

reduced across repeated presentations (**Fig 8Diii & 8Dv**; D1-Cre: rmANOVA $F(1.42,$

$12.77)=15.69$, $p=0.0008$; D2-Cre: rmANOVA $F(1.80, 16.21)=25.01$ $p<0.0001$). The evoked

neural activity and its subsequent adaptation across trials was similar in D1- and D2 mice (**Fig 8vi**;

two-way ANOVA, no effect of genotype $F(1, 18)=1.387$, $p=0.254$). These results indicate that the

activity of DA receptor-expressing ventral striatum neurons post-synaptic from midbrain DAergic

terminals is associated with sniffing.

DA receptor antagonism in the ventral striatum reduces both the occurrence and frequency of

sniffing

Ventral striatum neural activity can be influenced by a wealth of

neurotransmitters/neuromodulators arising from numerous bottom-up and top-down brain regions.

For instance, NAc activity is influenced not only by DA, but also by glutamatergic inputs from the

frontal cortex[69]. While the above results suggest that sniff-evoked changes in mesolimbic DA

influx to the ventral striatum engages postsynaptic neurons, those results still beg the question: is

DA binding to DA receptors on ventral striatum neurons needed for an animal to initiate sniffing?

We implanted c57bl/6j mice of both sexes with bilateral indwelling infusion cannulae into either

the NAcSh or TuS (**Fig 9A**). We focused on antagonism of DA receptors in the TuS and NAcSh

since it was within these regions wherein we found the strongest correlation between DA levels

and sniffing frequency (**Figs 3 & 4**). Since TuS and NAcSh neurons may express D1, D2, or D3

receptors[70–74], across days in a counter-balanced within subject's design, each mouse was

intracranially infused with SCH23390, raclopride, or PG01037 to selectively inhibit D1, D2, or

D3 receptors, respectively. The infusion occurred 10min prior to placing the mouse into the

plethysmograph for respiratory monitoring, wherein we measured both spontaneous and sensory-

evoked sniffing bouts (**Fig 9B**). We analyzed for effects of D1, D2, and D3 antagonism on both

the number of sniff bouts and their frequency, displayed spontaneously and stimulus-evoked. DA

receptor antagonism has some known sedative effects[75], and notably, DA receptor antagonism had

some effect on baseline breathing, with specifically D2 receptor antagonism in the NAcSh

influencing breath frequency (Wilcoxon signed-rank test W=-279, p =0.012; **Supplementary Fig**

3).

We found that D1 receptor binding contributed to spontaneous sniffing in that D1 receptor

antagonism in the TuS reduced the number of spontaneous sniffing bouts (**Fig 9Ci;** two-way

ANOVA, main effect of treatment $F_{(1, 24)}$=12.62, p=0.002; Sidak's multiple comparisons TuS

VEH vs. SCH p=0.023). D1 receptor binding also contributed to sensory-evoked sniffing in that

D1 receptor antagonism in the TuS reduced the number of buzz-evoked bouts (**Fig 9Ei**; two-way

ANOVA, main effect of treatment $F_{(1, 24)}$=4.30, p=0.049; Sidak's multiple comparisons TuS

VEH vs. SCH p=0.048).

Likewise, we found that D2 receptor binding contributed to spontaneous sniffing in that D2

receptor antagonism in both the TuS and NAc reduced the number and frequency of spontaneous

sniffing bouts (**Fig 9Cii**; two-way rmANOVA, main effect of treatment $F_{(1, 24)}$=48.64, p<0.0001;

Sidak's multiple comparisons TuS VEH vs. RAC p<0.0001, NAc VEH vs. RAC p=0.002). D2

receptor binding in the TuS and NAc also contributed to odor-evoked sniffing in that D2 receptor

antagonism reduced the number of sniffing bouts (**Fig 9Dii**; two-way rmANOVA, main effect of

treatment $F_{(1, 24)}$=31.26, p<0.0001; Sidak's multiple comparisons TuS VEH vs. RAC p<0.0001,

NAc VEH vs. RAC $p=0.017$). In contrast, we saw no effect of D3 receptor inhibition on

spontaneous or odor-evoked sniffing in either the TuS or NAcSh (**Fig 9Ciii, 9Diii, 9Eiii**).

Interestingly, D2 antagonism in both the TuS and NAc not only reduced the number of odor-

evoked bouts displayed by mice, but in both regions, the odor-evoked bouts which were displayed

were reduced in frequency (**Supplementary Fig 8Bi & 8Bii**; D1 two-way rmANOVA main effect

of treatment $F(1, 24)=7.75$, $p=0.010$; Sidak's multiple comparisons TuS VEH vs. SCH $p=0.228$,

NAcSh VEH vs. SCH $p=0.056$; D2 two-way rmANOVA, main effect of treatment $F(1, 24)=96.64$,

$p<0.0001$; Sidak's multiple comparisons TuS VEH vs. RAC $p<0.0001$, NAcSh VEH vs. RAC

$p<0.0001$). This was not observed during spontaneous sniffing bouts or buzz-evoked sniffing bouts

pointing to a unique system which supports the frequency of odor-evoked sniffing compared to

spontaneous sniffing which was influenced solely in terms of the total number of bouts. Altogether,

these results confirm that mesolimbic DA initiates sniffing by its actions upon D1 and D2 receptors

in the TuS and NAcSh.

Discussion

DA appears to partake in numerous functions[33,34,76,77]. For instance, DA can influence the

perceived salience of stimuli wherein DA might make certain smells stand out more[39], making

them more noticeable during sniffing. DA might enhance our sensitivity to sensory stimuli,

including odors, allowing them to be better detected at low levels[78]. DA is also involved in arousal

and attention mechanisms wherein it could heighten our alertness and focus on sensory inputs,

including odors and other cues[79,80]. Indeed, DA supports cognitive functions in humans during

non-olfactory functions such as during gazing[81]. Further, DA is involved in synaptic plasticity and

learning from experiences and adapting to new information. When sniffing odors, DA release, including upon ventral striatum neurons, could contribute to the brain's ability to form associations and learn information[30]. Likewise, when we encounter pleasant or novel odors, DA release might contribute to feelings of pleasure and satisfaction[53]. Indeed, DA is a key transmitter in the brain's reward system and is connected to motivation and exploration[82], wherein it may influence our motivation to sniff, encouraging us to explore and investigate different odor landscapes. Furthermore, several lines of evidence have documented the orchestration of sniffing during motivated behaviors and even upon anticipation of a reinforcer. These functions of DA stated, no prior work has mechanistically linked DA with the highly conserved and ethologically-relevant behavior of sniffing. Our results support a model wherein DA acts upon the ventral striatum, specifically upon neurons with D1 and D2 receptors in the TuS and NAcSh, to initiate sniffing and even increase the frequency of sniffing. This finding perhaps is analogous to the known role of DA in guiding attention in other sensory modalities, including in humans[83,84] We predict the promotion of sniffing by DA is an essential component linking the historic cooccurrence of sniffing with motivated states.

Stimulus active sampling behaviors have a tight linkage to both the acquisition and processing of environmental information, and these are modulated by ascending monoaminergic neurotransmission. For instance, stimulation of the locus coeruleus in rats increases the responsiveness of barrel cortex neurons to whisker deflection[85]. In zebra finches, auditory coding in the finch auditory cortex homologue is modulated by norepinephrine[86]. Likewise, visual processing in the lateral geniculate nucleus of cats is modulated by dopamine[87]. In terms of active sampling behaviors and their regulation by monoaminergic transmission, rhythmic licking in mice

can be suppressed by DA D1 receptor antagonism in the motor cortex[88]. The rhythmic movement

of the vibrissae ("whisking") is perhaps the most well-studied in terms of its modulation by

monoaminergic inputs. Serotonergic input from the dorsal raphe nucleus, whose neurons fire

rhythmically within the whisking frequency bands, generates whisking in rats and whisking can

be diminished by serotonin receptor antagonism in the dorsal raphe[89]. Also, recent evidence

indicates that mesolimbic DA input to the NAc in mice promotes orofacial behaviors including

whisking[41]. Our results add to these prior studies by establishing a necessary and sufficient role

for mesolimbic DA in the initiation of sniffing behavior in that not only does photostimulation of

DAergic terminals initiate sniffing bouts, but also, blockade of DAergic receptors and optical

inhibition of DAergic terminals suppresses the normal occurrence of sniffing and its frequency. In

line with these results, VTA→ventral striatum DA may serve as a pathway to facilitate adaptive

sniffing to promote olfactory search behavior, and moreover may shape odor processing given the

regulation of odor coding and perception by the respiratory rhythm[90–92]. Given the extensive

innervation of the olfactory bulb by neuromodulatory systems, including norepinephrine, and the

presence of DAergic interneurons in the olfactory bulb, one might speculate that odor processing

and perception are resultant from a delicate balance of neuromodulators[93], which not only

recognize a novel odor, but also consequentially instruct novel motor patterns. The time-course of

stimulation-evoked sniffing we report herein supports that indeed DAergic release into the ventral

striatum may not be the initial driver of novel-odor evoked sniffing, which can be as fast as

~100ms[94].

We found heterogeneity of DA release in ventral striatum subregions. Previous work from

Menegas and colleagues reported that DA is only released in the tail of the striatum upon novel

odor presentation (which likely would entail sniffing), with negligible changes in DA levels seen

in the ventral striatum[32]. It is interesting to note that in their study, ventral striatum recordings were

largely acquired from the NAcC, wherein we similarly did not see increases in DA levels.

Furthermore, the results of Menegas and colleagues even suggest subtle reductions in DA levels,

which agrees with our current findings from the NAcC. We expand upon this work by investigating

DA dynamics across all three ventral striatal regions which include the NAcC, NAcSh, and TuS

while animals sniff. Our findings are reminiscent of those monitoring DA release during other

behaviors which has uncovered heterogeneous levels of DA in the NAcC versus NacSh in manners

likely supported by differing VTA neural pathways innervating these subregions[43,44,95]. Here we

found that in comparison to the increases in DA upon sniffing in both the TuS and NAcSh, that

the NAcC experienced reductions in DA release (**Figs 2 & 3**). This implies that sniffing increases

tonic firing of VTA DAergic neurons, or suppresses firing of DAergic neurons, in the NAcSh and

TuS and NAcC, respectively. That optical excitation of the terminals in any of the regions equally

initiates sniffing suggests that this is overriding changes in VTA neuron firing which give rise to

DA dynamics in the ventral striatum and also hints that heterogeneity in subregion DA dynamics

are not a byproduct of interregional differences in DA clearance (*e.g.,* by different kinetics or

function of the dopamine transporter). It is tempting to speculate that different DA dynamics in

these subregions might stem from unique inputs from areas that may signal salience or arousal

which would synapse on VTA GABAergic neurons, including projections from the TuS and NAc

themselves[95]. Furthermore, different DA neuron populations in the VTA display heterogenous

responses to novelty vs. reward[32,80,96,97], which might stem from functional subtypes of DAergic

neurons. It will be important to investigate the differing VTA neuron populations which may

underly these subregion-specific differences in DA dynamics in the ventral striatum during

sniffing.

What 'type' of sniffing does mesolimbic DA initiate? The machine learning model trained on

different experimenter classified sniffing patterns (**Fig 5D**) points to mesolimbic DA-evoked

sniffing as its own type of sniffing, unique from that a mouse would self-motivate upon the desire

to explore (spontaneous) and that evoked by a sensory stimulus (like an odor or buzz). So, what

type is it? Sniffing is a dynamic behavior which can be displayed in unique manners based upon

context. For instance, dogs sniff at lower frequencies when searching the air for odors than when

searching on the ground wherein they elevate their sniffing rate[98]. Rats also display sniffing in a

distinct frequency band when transitioning from sampling odors to anticipating a reinforcer in an

instrumental task in manners which suggests sniffing reflects strategic 'modes' or states[5]. Our

results are in-line with these prior works and suggest that mesolimbic DA-evoked sniffing reflects

a unique sniffing mode. Several reports have established links between specific brain regions, and

neurons within those brain regions, in regulating adaptive respiration, although the role of those

regions does not appear to initiate sniffing as much as shape more specific respiratory parameters

(*e.g.* [99]). Our results wherein we varied photostimulation of DAergic terminals uncovered that the

duration of evoked-sniffing bouts did not increase with more prolonged optical stimulation (**Fig**

4H). Considering that VTA DAergic neurons fire in characteristic phasic manners[100,101], this

suggests that DA release into the ventral striatum is more so to initiate and possibly elevate sniffing

frequency than to maintain its tonic display.

There are a few caveats to this study we believe important to note. First, optogenetic stimulation

of DAergic terminals, while herein an effective means at robustly driving sniffing, is likely driving

DA release onto postsynaptic neurons in supraphysiological manners. While we selected a

stimulation paradigm established by others to be effective on TuS projecting VTA terminals, it is

nevertheless likely DA is released at higher than usual levels. Nevertheless, both our

pharmacological data (**Fig 9**) and our work wherein we removed *VGluT* from tyrosine hydroxylase

expressing neurons (**Supplementary Fig 6**) all agree that DA is contributing to the initiation of

sniffing. It is also possible that some behavioral effects from the photostimulation are due to

backpropagation of activity into the VTA itself. Again though, the pharmacological results indicate

DA input to the ventral striatum is important for sniffing. Second, while DA release into the ventral

striatum initiates sniffing, and while in several of our analyses, DA levels were correlated with

sniffing frequency, this was not always the case. For instance, for the buzz stimulus experiments,

large DA levels were found despite the mice only displaying modest frequencies of sniffing. This

corresponds with other studies which have reported DA release in response to a variety of salient

stimuli. Finally, while D2 receptors are enriched on spiny projection neurons in the TuS and NAc,

they also serve autoreceptor functions on VTA DAergic terminals, including those innervating

these regions. So, while we know local D1 antagonism suppressed sniffing, it is possible the effects

of D2 antagonism are due to contributions of antagonizing D2 receptors on both the VTA terminals

(which would enhance DA release) and on spiny projection neurons.

Antagonizing DAergic receptors versus inhibiting DAergic terminals resulted in different effects

upon respiration. Whereas pharmacological inhibition of TuS D1 receptors and NAc D1 and D2

receptors influenced specific occurrences of sniffing, including both spontaneous and stimulus-

evoked sniffing (**Fig 9**), optical inhibition of DAergic terminals only impacted spontaneous events

(**Fig 6**). We reason this may be due to a few possibilities. In our photoinhibition experiments, we

targeted the TuS and NAcSh separately. Given the transient and specific nature of the

photoinhibition paradigm, it is likely that the inhibition of DA release in the TuS could not override

the resultant effects of DA release in the NAcSh on sniffing, or *vice versa*, therefore resulting in

modest changes in respiration. Why then did we see robust effects when we pharmacologically

inhibited DA receptors in a single region (**Fig 9 & Supplementary Fig 8**)? We reason this may

be because a direct antagonist infusion would inhibit a great number of post-synaptic receptors,

and this coupled with potential influences of the antagonists on autoreceptor function and the

massive collateralization of spiny projection neurons upon each other would yield more dramatic

changes in the ability of the cells to act upon presynaptic DA.

A major question is how the receipt of DA upon spiny projection neurons might ultimately drive

changes in the respiratory rhythm. Both the TuS and NAc have outputs into downstream structures

important for homeostatic regulation/control and limbic function. Ventral striatum spiny

projection neurons also feedback upon the VTA in manners regulating DA release and this too

could support, even if just bisynaptically, modulation of respiratory central pattern generators

which are sensitive to dopamine[102,103]. Perhaps one clue may reside from our results wherein we

attempted to stimulate sniffing under anesthesia and found that DA release into the TuS or NAc is

incapable of initiating sniffing (**Fig 5A & B**). This suggests that the functional 'sniffing' output

from the ventral striatum is within a structure which can drive changes in breathing but not under

anesthesia. It is not uncommon for stimulation of some respiratory control centers to be incapable

of altering respiration under anesthesia[21,103]. For instance, the ventral striatum projects to

hypothalamic nuclei that are known to innervate respiratory control nuclei in the respiratory pons

and could support entrainment of breathing at sniffing frequencies. Interestingly, these

hypothalamic targets are also state dependent in that changes in breathing frequency do not occur

upon stimulation under anesthesia [21]. Thus, there seems to be somewhat of a paradox in that in

some respiratory structures, including the PreBötzinger complex, high frequency stimulation is

incapable of driving respiration at frequencies greater than 6Hz[104,105]. Identifying what

downstream circuits link the TuS and NAc to respiratory brain stem circuits to orchestrate sniffing

remains an important goal. Our results point to D1 and D2 ventral striatum neurons being part of

a feedforward/feedback loop between the basal ganglia and respiratory control regions with

DAergic VTA neurons being an intermediary circuit node. While the bio-sensor based fiber

photometry recordings we used to capture changes in DA and ventral striatum D1 and D2 neurons

suggest that these signals may at times lag the onset of a sniff bout, we did not correct for the lag

inherit in photometry recordings. Despite that, in several of the examples provided, DA levels

begin to rise immediately upon sniff bout onset (*e.g.*, **Fig 2Eii**). Further, the TuS single unit

recordings offered temporal dynamics which point to firing of at least some TuS neurons

immediately prior to the detectable increase in sniffing. This ultimately leads us to speculate that

a signal arrives at the ventral striatum (regarding a novel odor, or a change in environment or

internal state) which is ultimately routed through basal ganglia to initiate sniffing by means of

actions in the respiratory brainstem. At that point, DA is released by activation of VTA DAergic

neurons (perhaps as even modulated by ventral striatum input itself, e.g.,[106,107]), which then, based

on our results, would further drive ("invigorate") sniffing by its actions on those same ventral

striatum neurons. If future studies support this reciprocal DAergic loop model for sniffing,

naturally they would need to incorporate inputs/outputs to frontal cortices, hypothalamic

structures, motor cortex, and other regions integral for the plethora of behaviors and states which

might coincide with sniffing. While in many of our measures the frequency of sniffing positively

correlated with the peak magnitude of DA levels (*e.g.,* **Figs 2 & 3**), some instances of sniffing did

not co-occur with major rises in DA (*e.g.,* **Supplementary Fig 2 and also as shown in Figs 2Fii**

& 2Gii) and so it is likely multiple circuits are at play which link DA with sniffing.

In conclusion, we report evidence that mesolimbic DA input into the ventral striatum contributes

to the display of the widely conserved behavior of sniffing. The nature of sniffing being integral

to both olfaction and motivated behaviors implicates this circuit in a wide array of functions.

Especially in macrosmatic animals, like the mice studied herein, olfaction is crucial for appropriate

social behaviors, avoiding predation, and foraging. Based upon our evidence, we propose it is

likely this circuit provides a neuromodulatory means whereby motivated states signaled by DA

release tune the occurrence and frequency of sniffing in order for animals to thrive in the above

ethological scenarios.

<u>Methods</u>

Animals

Male and female mice were housed on a 12:12hr light-dark cycle with *ad libitum* access to food

and water. All behavioral testing took place during the light cycle. Mice with viral injections alone

were group housed (≤5 mice/cage) and any mouse with a chronic implant was single housed

following its surgical procedure. Food (Envigo Teklad Global, 18% rodent diet irradiated pellet,

Cat # 2018; Indianapolis, IN) and reverse osmosis water were available *ad libitum* except during

behavioral monitoring/recordings. Mice were housed in standard shoebox sized plastic cages

(Allentown Jag 75 micro-vent system; Allentown, PA; L: 29.2cm W: 18.5cm H:12.7cm) on a rack

made for individually vented cages. Corncob bedding and a Nestlet^{TM} (Ancare; Bellmore, NY)

were in each cage along with manzanita sticks (Bio-Serve). Mice were housed on a 600-1800hr

light cycle with lights on during the daytime. Temperature in the room averaged $70 \pm 2°F$, with

30-70% humidity and 10-15 room air changes per hour. The housing room was specific pathogen

free. Observation occurred from 1000-1700hrs. All animal care was conducted with in the

AAALAC animal research program of the University of Florida, in accordance with the guidelines

from the Guide for the Care and Use of Laboratory Animals and approved by the University of

Florida Institutional Animal Care and Use Committee.

Mouse lines included the following transgenic lines which all were maintained on a c57bl/6j

background (Jackson labs, strain #000664; RRID:IMSR_JAX:000664) and were bred in house

within a University of Florida vivarium. *drd1-Cre* and *drd-2-Cre* lines[67] were obtained from the

UC Davis Mutant Mouse Regional Resource Center (D1: EY262Gsat/Mmucd; D2: ER44Gsat).

DAT^{IRES}-Cre mice[108] (strain #006660; RRID: IMSR_JAX:006660), *TH-Cre* mice[109]; strain

#008601; RRID: IMSR_JAX:008601), *VGluT2 fl/fl mice*[110]; strain #012898; RRID:

IMSR_JAX:012898), and *Ai9* TdTomato Cre reporter mice[111] (strain # 007909,

IMSR_JAX:007909) were obtained from Jackson labs. A total of 178 mice (mixed sexes) were

used in this study for experimentation. Out of these, a total number of 51 were excluded from data

analyses due to experimental issues (*e.g.,* surgical complications, off-target brain implants, viral

injections which missed the target).

For all surgical procedures, under aseptic conditions, mice were anesthetized with Isoflurane (2-

4% in oxygen, IsoFlo®, Patterson Veterinary) and mounted in a stereotaxic frame where they were

maintained under Isoflurane. Body temperatures were maintained with a 38°C heating pad

underneath. A local anesthetic, bupivacaine hydrochloride (Marcaine™, 5mg/kg subcutaneously,

s.c., Patterson Veterinary), and the analgesic meloxicam (5mg/kg, s.c., Patterson Veterinary) were

administered before incision and exposure of the skull. Mice receiving chronic implants (*e.g.,*

cannulae, optic fibers) were given meloxicam (0.5mg/kg, s.c.) for 3 days following surgery.

For viral injections, craniotomies were made over the ventral tegmental area (VTA) (-

3.6AP/0.35ML/-4.75DV), tubular striatum (TuS) (1.4AP/1ML/-4.90DV), nucleus accumbens

shell (NAcSh) (1.4AP/1ML/-4.40DV), and/or nucleus accumbens core (NAcC),(1.4AP/1ML/-

3.90DV) and a pulled glass micropipette containing the AAV was lowered into the region of

interest. Viruses were all delivered via the pulled pipette attached to a Nanoject III (Drummond

Scientific) at a rate of 2nl/s unless otherwise noted. After waiting 10min, the pipette was slowly

withdrawn from the brain. Unless followed by an implant in the same surgery, craniotomies were

sealed with dental wax, and the incision closed with wound clips.

For intranasal 'sniff' cannula implantation, a ~1mm diameter hole was drilled through the right

nasal bone, 1.5mm anterior the frontal/nasal fissure and 0.5mm lateral following prior methods [6].

A stainless-steel guide cannula (22GA, #C313G, P1 Technologies, Roanoke, VA) cut to extend

0.5mm below the pedestal was lowered through the craniotomy and secured with dental cement.

A dummy cannula (#C313DC, P1 Technologies, Roanoke, VA) cut to extend 0.2mm past the guide

cannula was also inserted when not recording from the mouse.

To record respiration, the dummy cannula was removed and a pressure transducer was connected

(CPXL04GF, Honeywell) to the cannula via a flexible piece of polyethylene tubing. The mouse

was then placed in the plethysmograph and recorded for 15min. Plethysmograph-recorded

transients were detected using a flow transducer (Data Sciences International) and digitized at

300Hz (0.1-12Hz band pass; Tucker Davis Technologies, Alachua, FL), following a 500x gain

amplification (Cyngus Technology Inc, Southport, NC). Intranasal pressure transients were

amplified and digitized using the same settings.

Anatomical Tracing

For synaptophysin injections, a craniotomy was made over the VTA, and a micropipette containing

100nL of pAAV-hSyn-FLEx-mGFP-2A-Synaptophysin-mRuby (Addgene 71760-AAV1; titer:

$9.8x10^{12}$vg/mL) was injected unilaterally into the VTA of DAT^{IRES}-Cre mice.

3 weeks following injections, mice were overdosed with sodium pentobarbital (Fatal-Plus,

Patterson Veterinary) and transcardially perfused with 0.9% saline followed by 10% phosphate-

buffered formalin. Brains were removed and stored in a 10% formalin in 30% sucrose solution at

4°C prior to sectioning. Brains were coronally sectioned at 40μm and sections were stored in Tris-

buffered saline with 0.03% sodium azide.

Image acquisition and quantification

Brain regions of interest were identified using the mouse brain atlas[112]. For the TuS, 12 images

were acquired spanning 1.94mm - 0.14mm anterior to Bregma. For the nucleus accumbens, a total

of 8 images were acquired spanning 1.94mm – 0.74mm anterior Bregma.

Images were acquired with a Nikon Eclipse Ti2e fluorescent microscope. For brains with

synaptophysin injections (*n*=6 mice), images were acquired at 20x magnification across the

hemisphere ipsilateral to the injection site, and Z-stacked at every 4μm. For quantification, regions

of interest (ROIs) were drawn around areas of interest, namely the NAcC and NAcSh along with

the medial and lateral TuS. To determine the medial and lateral portions, the TuS was divided into

thirds and the most medial or lateral third was used to ensure distinctness between the two

regions[113]. Images were preprocessed to decrease average background fluorescence and enhance

contrast. A semiautomated counting algorithm created within NIS elements software (Nikon) was

used to detect and count fluorescent puncta, allowing for an unbiased estimation of puncta

numbers. Puncta were identified and quantified based on their size and fluorescence intensity via

bright-spot detection.

To reconstruct synaptophysin injection sites **(Supplementary Fig 1)**, the VTA was imaged from

-2.92 to -3.80mm posterior bregma, as identified with use of a brain atlas[112]. Next, an experimenter

(A. C.-L.) identified and marked GFP+ cell bodies in VTA schematics within Adobe Illustrator.

Fiber photometric recording in the plethysmograph

Virus injection and optical fiber implant

C57bl/6j mice were injected unilaterally in the NAcSh, NAcC, or TuS (n=7/group) with 100nL

pAAV.hSyn-GRAB_DA1h (Addgene 113050, titer: 1×10^{13}vg/mL)[50]. For controls, c57bl/6j mice

were injected unilaterally in NAcSh, NAcC, or TuS (n=2-4/group) with 100nL pAAV.hSyn-

GRAB_DA-mut (Addgene 140555, titer: 1×10^{13} vg/mL). Viral injections were performed as

previously described (see 'Animals'). In the same surgery, a 400µm core, 0.48NA optical fiber

secured in a 2.5mm outer diameter metal ferrule was chronically implanted to terminate in the

TuS, NAcSh, or NAcC and affixed to the skull with dental cement. Animals were recorded from

3-5 weeks following surgery.

Photometry

465nm (GRAB$_{DA}$/GRAB$_{DA}$-mut excitation wavelength, driven at 210Hz) and 405nm (UV

excitation wavelength, driven at 330Hz; control) light emitting diodes (TDT, RZ10x) were coupled

to a fluorescent minicube (Doric lenses, FMC6) via 200µm, 0.22NA FCM optic fiber patch cords.

Both excitation and emission light were directed through a single 400µm, 0.57NA patch cord

connected to the animal by the implanted 2.5mm metal ferrule. Emission light was directed

through the mini cube and coupled to femtowatt photoreceivers to monitor GFP and UV signals

(TDT, RZ10x). The emission signals were low-pass filtered (6th order), pulse demodulated, and

digitized (1kHz) using a Tucker Davis Technologies processor (TDT). All parameters for imaging

and acquisition of photometry data were consistent across all groups.

Respiratory data acquisition

To simultaneously monitor photometry signals and respiration, mice were gently tethered to the

patch cord and placed in a whole-body plethysmograph (Data Sciences International, St. Paul,

MN). Respiratory transients were detected using a flow transducer (Data Sciences International)

and digitized at 300Hz (0.1-12Hz band pass; Tucker Davis Technologies, Alachua, FL), following

a 500x gain amplification (Cyngus Technology Inc, Southport, NC). Within all plethysmography

recordings, we analyzed the instantaneous frequencies of the peak-detected respiratory signal to

identify periods indicative of sniffing[6]. Since mice rapidly transition in and out of sniffing bouts [6],

we restricted our analyses to more stable/pronounced bouts versus short bouts by only defining a

sniff bout as the occurrence of sniffing ≥6Hz for ≥1s in duration. Further, in some instances, mice

continuously sniff for numerous seconds and then reengage in another bout shortly thereafter.

Since these may reflect periods of more generalized arousal versus spontaneous exploratory

behavior, we excluded sniff bouts occurring in proximity to these long (>10s) sniffing bouts. In a

subset of mice, an observer *post-hoc* monitored video (10Hz, 640x480 resolution) which was

recorded simultaneously with respiratory signal from the plethysmograph to validate that focusing

on >6Hz changes in respiration was indicative of 'active sampling' behaviors from the mouse

(including snout directed investigation and head-turning) and found 1:1 correspondence

supporting use of respiratory signals from the plethysmograph as suitable for detecting active

investigation (of course in this set-up we could not see minor facial, motor or vibrissal

movements).

Behavioral monitoring

Behavioral sessions lasted approximately 30min during which mice were presented with both

olfactory and non-olfactory stimuli. To allow for examination of spontaneous sniffing, we included

5min periods where no stimuli were presented at the beginning (0-5min), middle (12-17min), and

end (28-33min) of a session. Spontaneous bouts for analysis were identified between 2-5min, 14-

17min, and 30-33min to ensure that they were far removed from any stimuli presentations.

Between 5-12min, 2 trials of 4 different stimuli were pseudorandomly presented. The final 3 trials

of pseudorandom stimuli presentations occurred between 17-28min. Each stimulus was presented

for 2s with a 60s inter-trial interval.

Stimulus delivery

Odors R(+)-limonene and thioglycolic acid (Sigma-Aldrich, >97% purity), and that of peanut oil

(La Tourangelle®, Woodland, CA), were presented to mice via a custom air-dilution odor delivery

device connected to the plethysmograph (see Results for rationale in selecting these specific

odors). During experimentation, 2mL of all odors were contained in 40mL glass headspace vials

at room temperature. Computer-controlled and user-initiated openings of valves allowed for odors

to pass through headspace vials at a flow rate of 50mL/min, following which they were mixed with

a carrier stream of clean filtered room air delivered by a pump (1L/min; Tetra Whisper, Melle,

Germany) before being introduced into the plethysmograph. All odors were handled within

independent tubing to prevent cross-contamination. Following each odor presentation, odor-

vaporized air was passively cleared from the plethysmograph through an exhaust outlet at the

chamber's ceiling since the filtered room air is continuously delivered. Serving as a non-olfactory

stimulus, a vibration motor (3V, 12000 RPM; BestTong), referred to as "buzz," was secured to the

outside of the plethysmograph. During trials of pseudorandom stimulus presentations, each

stimulus was presented for 2s with a 60s inter-trial interval.

Fiber photometric recording in head-fixed mice

Virus injection and optical fiber implant

Mice for this method were also used for data collection and their results published (albeit from

different stimulus conditions wherein analyses were performed on well-learned odor-associated

pairs) in a prior study[40]. As we previously described in that study, *D1-Cre* and *D2-Cre* mice were

injected in the TuS with 1μL AAV5-EF1α-DIO-GCaMP6f-WPRE (obtained from the Penn Vector

Core in the Gene Therapy Program of the University of Pennsylvania, 5.43×10^{13}GC/mL). Viral

injections were performed as described above (see 'Animals'). Two weeks later, a 400μm core,

0.48NA optical fiber secured in a 2.5mm metal ferrule was chronically implanted to terminate in

the TuS. A plastic head-bar was also affixed to the skull with dental cement for later head fixation.

Animals were given a week for surgical recovery before undergoing behavioral testing.

In vivo fiber photometry

465nm (GCaMP excitation wavelength, driven at 210Hz) and 405nm (UV excitation wavelength,

driven at 330Hz; control) light emitting diodes were coupled to a 5 port minicube (Doric lenses,

FCM5) using 400μm, 0.48NA FCM optic fiber patch cords. Both excitation and emission light

were directed through a patch cord connected to the animal by the implanted 2.5mm metal ferrule.

Emission light was directed through the 5-port mini cube and coupled to femtowatt photoreceivers

to monitor GFP and UV signals. The data were then digitized and pulse demodulated at 1kHz

using a Tucker Davis Technologies processor (TDT). All parameters for imaging and acquisition

of photometry data were consistent across D1- and D2-Cre mice.

Behavioral sessions

Animals were head-fixed and presented with sets of 3 novel odors, 4 times each. To enhance each

animal's contribution to a more rigorous data set, animals were tested a second time at least 24hrs

later with sets of 3 different novel odors. Odors were delivered for 2s duration with a $17 \pm 2s$ inter-

trial interval through an odor port positioned 1cm from the animal's snout.

Stimulus delivery

Odors were presented through an olfactometer and delivered through an odor port positioned 1cm

from the animal's snout. Odors included ethyl butyrate, 1,7-octadiene, isopentyl acetate, heptanal,

2-heptanone, R(+)-limonene, ethyl propionate, S(-)-limonene, methanol, methyl valerate, 2-

butanone, 1,4-cineole, butanal, propyl acetate, allylbenzene, allyl bromide, isobutyl propionate,

and 2-methylbutyraldehyde (Sigma Aldrich, all > 97% purity). Odors were diluted to 1Torr in

mineral oil and further diluted by mixing 100mL odor vaporized N_2 with 900mL medical grade

N_2 (Airgas). Stimuli were therefore presented at a total flow rate of 1L/min. Odor was continuously

flowing to the odor port but removed by a vacuum prior to reaching the animal. Odor sets used

varied across animals and across sessions.

Respiratory data acquisition

922 Respiration was recorded via an extranasal flow sensor (CPXL04GF, Honeywell) positioned near

923 the animal's nose as described[68]. A 1/8" inner diameter hose which terminated into the flow sensor

924 was placed ~7-10mm from the nose of the mouse during head fixation.

Optogenetic stimulation in the plethysmograph

Virus injection and fiber implant

DAT^IRES-Cre x Ai9 mice or *VGluT2 fl/fl x TH-Cre* mice were injected unilaterally in the VTA with

400nL AAV5-EF1α-DIO-hChR2(E123T/T159C)-EYFP (Addgene 35509, titer: 1.3×10^{13} vg/mL)

or AAV5-EF1α-DIO-EYFP (for controls, UNC vector Core, titer: 3.5×10^{12}vg/mL). While we used

DAT to drive Cre expression in DAergic neurons in other experiments in this study, here we used

the gene encoding tyrosine hydroxylase (*TH*) and specially *VGluT2 fl/fl x TH-Cre* mice since this

line was already available in-house. While this *TH*-based strategy may not had captured the

identical cell population as in the DAT-based strategy, and *VGluT2* may have been removed from

non-DA cells in the VTA[114], this mouse line allowed us to isolate the influence of DA on sniffing

from glutamate. Viral injections were performed as previously described. In the same surgery, a

300µm core 0.39 NA optical fiber in a ceramic ferrule was chronically implanted to terminate in

the TuS, NAcSh, or NAcC in *DAT^{IRES}-Cre x Ai9* mice or in the TuS for *VGluT2 fl/fl x TH-Cre* and

affixed to the skull with dental cement. Animals were tested no sooner than 4 weeks following

surgery.

To reconstruct ChR2.EYFP injection sites in the VTA of *DAT^{IRES}-Cre* mice (**Supplementary Fig**

3), the VTA was imaged from -2.92 to -3.80mm posterior bregma, as identified with use of a brain

atlas[112]. An independent rater used Adobe Illutrator to trace density of expression across images.

Light stimulation and behavioral monitoring

Optogenetic stimulation (25Hz, 15ms pulse width, 1s duration, 18mW[3]) was delivered by a 473nm

laser (CrystaLaser, DL473-100) and through a 300µm, 0.39NA fiber connected the laser to a fiber

optic rotary joint, with a 400µm, 0.47NA fiber connecting the rotary joint to the mouse (brass

mating sleeve). This paradigm was based on prior studies[39,40,54] and pilot work done in the lab.

Animals (*DAT^{IRES}-Cre*: TuS *n*=11, NAcSh *n*=6, NAcC *n*=6, EYFP controls *n*=9; *VGluT2 fl/fl x*

TH-Cre: TuS *n*=4, EYFP controls *n*=4) received one habituation day wherein they were gently

tethered to the path cable and placed in the plethysmograph for 20mins. The following day, animals

were again tethered, placed in the plethysmograph, and given a brief, 5min habituation period.

Following this, photostimulations (7-10/mouse) were delivered with a 2 ± 1min inter-trial interval.

A subset of *DAT^IRES^-Cre* x *Ai9* (TuS: *n*=5, NAcSh: *n*=4, NAcC: *n*=4, EYFP controls: *n*=6)

underwent testing with varying durations of photostimulation. Stimulations at 25Hz with a 15ms

pulse width were delivered for durations of 120ms, 1s, 2s, or 4s. After a habituation day, mice

received a single stimulation parameter per day over four consecutive days. Each mouse underwent

the 1s duration on day 1, and the following 3 stimulation paradigms were randomized across mice

over the remaining 3 testing days. Photostimulations (9-11/mouse) were delivered with a 2 ± 1min

inter-trial interval.

Concluding testing, the same mice were then deeply anesthetized with urethane (1g/kg i.p.; Sigma

Life Science), tethered, and placed in the plethysmograph and photostimulated as described above.

Mice were transcardially perfused following recordings and within 30mins of urethane

administration.

Optogenetic inhibition in the plethysmograph

Virus injection and fiber implant

DAT^IRES^-Cre x Ai9 mice or *DAT^IRES^-Cre* mice were injected bilaterally in the VTA with 300nL

AAV5-EF1α-DIO-eNpHR 3.0-EYFP (Addgene 26966, titer: 1.0×10^{13} GC/mL). Viral injections

were performed as previously described earlier in this study. In the same surgery, 400μm core,

0.39 NA optical fibers in ceramic ferrules were chronically implanted to bilaterally terminate in

the TuS or NAcSh and affixed to the skull with dental cement. To afford precision in the bilateral spacing, the ferrules were first affixed in a 3D-printed adapter to ensure perfect 2.0mm from epicenter-to-epicenter spacing (tip-to-tip). Animals were tested no sooner than 4 weeks following surgery. Representative images of the primary injection site location of eNpHR 3.0-EYFP in the VTA were confirmed and imaged by an independent experimenter (**Supplementary Fig 7**).

Light stimulation and behavioral monitoring

Optogenetic stimulation (7-9mW^3, 5-7s duration, constant, see Results for rationale) was delivered by 405nm or 560nm LEDs (ThorLabs: Lx405 and Lx560). A 400μm, 0.48NA fiber connected the LED to the mouse.

To examine whether optogenetic inhibition of DAergic terminals in the ventral striatum had any effect on odor- or buzz-evoked sniffing, animals then underwent a similar behavioral session as previously described (see Fiber photometric recording in the plethysmograph, Behavioral monitoring**).** Animals first received one habituation day wherein they were gently tethered to the path cable and placed in the plethysmograph for 20mins. Following this, in a counterbalanced, within-subjects design, animals were tested over a 2-week period wherein they received concurrent 405nm or 560nm light delivery (7s duration, constant), initiated at the onset of both odor and buzz delivery.

Separately, the same mice were again tested in a counterbalanced, within subject's design over 2 days. Behavioral sessions lasted ~10 minutes. Mice were gently tethered and placed in the

998   plethysmograph. Starting at minute 2, 5s (constant) of 405nm or 560nm light was delivered every

999   20s for 8 minutes, for a total of 24 light deliveries.

1000

Machine learning to classify respiration

1002   The study utilized a dataset comprising differently evoked sniffing data which consist of 261

1003   samples, each 1s long. The dataset consisted of time series data representing respiratory

1004   frequencies (10ms bins).

1005

Dynamic Time Warping (DTW) was employed as the distance metric for measuring the similarity

between the time series data (respiratory frequency). DTW is known for its ability to align and

compare time series data. DTW involves finding the optimal alignment between sets of data points

of two sequences by warping the time axes. This allows the comparison of sequences with different

lengths or different temporal structures. DTW creates a cost matrix, where each element represents

the distance or dissimilarity between corresponding points in each of the two time series. Dynamic

programming is used to efficiently find the optimal alignment by considering all possible

alignments and selecting the one with the lowest cumulative distance [59]

1014

The k-Nearest Neighbors (k-NN) algorithm was utilized for classification of unknown data points

based on the majority class of its k-nearest neighbors in the feature space. In the context of time

series data, DTW can be used as a distance metric in the k-NN algorithm instead of using

traditional Euclidean distance. For a given query time series, the algorithm identifies the k-NN

from the training set based on DTW distance[60]. For each query time series, the k-NN were

identified and a majority voting scheme was employed to assign the class label.

1021

Prior to applying the DTW k-NN algorithm, time series data were applied min-max normalization

to ensure a uniform scale across all features. Subsequently, the dataset was divided into training

and testing sets, with 80% of the samples allocated for training (208 samples) and 20% for testing

(53 samples). The performance of the DTW k-NN algorithm was evaluated in terms of its

accuracy, precision, recall, and F1-score. The max warping window was set at 7 and k at 3 which

identified the combination that maximized the performance metrics during experiments.

1028

The experiments were conducted using Google Colab (Google Brain Team, Mountain View,

CA, USA). The DTW k-NN algorithm[61] was implemented in Python, utilizing the code available

in the GitHub repository: https://github.com/markdregan/K-Nearest-Neighbors-with-Dynamic-

Time-Warping. The dataset consisted of 63 photostimulation-evoked, 65 odor-evoked, 63 buzz-

evoked, and 70 spontaneous respiratory/sniff bouts (N=261 total samples). SMOTE[62] was applied

to generate synthetic data samples to bootstrap all groups to a total of 70 samples.

1035

Single unit recordings of ventral striatum neuron activity in head-fixed mice

These mice were used for data collection and their results published in a prior study from our

group, with exception of any neural analyses aligned to intranasal respiration as done herein[64]. As

we described in that study, c57bl/6j mice were implanted with eight-channel tungsten electrode

arrays in the TuS and implanted with intranasal cannula for simultaneous respiratory monitoring.

A plastic head-bar was also affixed to the skull with dental cement for later head-fixation. Animals

were given a week for surgical recovery before undergoing behavioral testing wherein water-

motivated mice were shaped to lick for a fluid reward for one odor, not a different one. Each animal

had 2-3 recording sessions on different days. Single neurons were sorted in Spike2, and then

spiking and respiration were analyzed in MATLAB.

1046

1047   Units from the same animal but not from the same recording day were limited to different channels.

1048   The spontaneous period was within the intertrial interval excluding the five seconds from odor

1049   onset to avoid any swallowing or licking effect. The odor presentation period was defined as the

1050   time from odor onset to two seconds after odor onset. The respiration cycle was the interval

1051   between two positive peaks in respiration signals.

1052

1053   *Sniffing burst detection*

1054   A high-pass filter (5 Hz) was applied to the respiration signal. The root mean square (RMS) was

1055   calculated using a 1s time window to quantify respiratory activity. The RMS was then transferred

1056   into the z-score. A threshold was set to 1 for sniffing onset detection.

1057

Spike ratio and initial and terminal spike analyses

Respiration cycles were divided into four groups based on their duration, each containing 25% of

the respiratory cycles. The spike number was then counted in each group for each unit. To establish

control, 100 sets of randomly generated spikes were created, with the number of spikes in each set

matching the actual spike number of the unit. The spike ratio was calculated as the spike number

divided by the mean of the randomly generated spike number in each percentile group. The

percentiles of the spontaneous and odor presentation periods were calculated separately. The

initial spike was defined as the first spike without any spikes in the previous five respiration cycles.

The terminal spike was the spike that had at least five spikes in the previous five respiration cycles

and was not followed by any spikes in the five subsequent respiration cycles.

1068

Pharmacological manipulations

Intracranial cannulae implantation

Bilateral craniotomies were made to target the the TuS and NAcSh. Stainless steel bilateral guide

cannulae (26GA, 2.0mm center-to-center distance, 5.3mm long from the base of the pedestal, P1

Technologies, Roanoke, VA) were lowered either to target the TuS (*n*=13) or the NAcSh (*n*=13).

Cannulae were affixed to the skull via dental cement, and dummy cannulae with a 0.5mm

projection from the end of the guide cannulae were then inserted and dust caps were added.

1076

Selective antagonists and intracranial infusions

SCH23390 (Tocris Bioscience, Bristol, UK) was dissolved in 0.9% sterile saline to a concentration

of 0.5µg/µL. 0.1µL was infused to deliver 0.05µg/hemisphere. Raclopride (Tocris Bioscience,

Bristol, UK) was dissolved in 0.9% saline + 5% dimethyl sulfoxide to a concentration of

23.8µg/µL, and 0.21µL was infused to deliver 5µg/hemisphere. PG01037 (Tocris Bioscience,

Bristol, UK) was dissolved in 0.9% saline + 12% DMSO to a concentration of 20µg/µL. 0.15µL

was infused to achieve 3µg/hemisphere. Internal cannulae that were cut at lengths projecting

0.5mm past the guide cannulae and thus terminating in the TuS or NAc were connected to 25GA

polyurethane tubing (Instech Laboratories, Plymouth Meeting, PA) backfilled with dH2O. 1µL

Hamilton syringes (Reno, NV) were used to inject total volumes ranging from 0.1-0.21µL. Mice

were gently scruffed, dust caps and dummy cannulae were removed, and the internal cannulae was

inserted before mice were placed back in their home cages for infusions. All drugs were infused

at a rate of 0.1µL/min followed by a 2min rest period before the internal cannulae was removed,

dummy cannulae and dustcap replaced. For all groups, behavioral testing began 10min following

the start of drug infusion. All mice were tested across a 6 week period and recieved all selective

antagonists and their corresponding vehicles, with 7 days between testing sessions.

1093

Behavioral methods

To monitor spontaneous and sensory-evoked sniffing behavior, mice were placed in the

plethysmograph and underwent behavioral testing and stimulus delivery as previously described

to monitor odor-, buzz-, and spontaneously-evoked sniffing (see Fiber photometric recording in

the plethysmograph, Behavioral monitoring). Baseline breathing rates were identified as periods

far removed from stimulation windows (*i.e.,* 2-5min, 14-17min, 30-33min).

1100

Histology

Mice receiving synaptophysin injections were perfused 3 weeks following injection. Mice

undergoing behavior were perfused following the last day of behavioral monitoring. All mice were

overdosed with sodium pentobarbital (Fatal-Plus, Patterson Veterinary) and transcardially

perfused with 0.9% saline followed by 10% phosphate-buffered formalin. Brains were removed

and stored (at 4ºC) in a 10% formalin in 30% sucrose solution prior to sectioning. Brains were

coronally sectioned at 40µm and sections were stored in Tris-buffered saline with 0.03% sodium

azide. Microscopic examination of tissue occurred on a Ti2e inverted fluorescent microscope, and

only mice with on-target manipulations (cannulae locations, viral injections, viral injections +

optic fiber placements) contributed to data sets. Verification of brain regions and target sites was

aided by use of a brain atlas[112].

1112

Quantification and Statistical Analysis

Analysis of respiratory data.

1115 Inhalation peaks (sampled at 300Hz) were detected offline in Spike2 (Cambridge Electronic

1116 Design, Cambridge, UK). First, the respiratory data were filtered (0.1-12Hz band pass). Second,

1117 the maximum point of each respiratory cycle was identified, and instantaneous frequency was then

1118 calculated based off of the duration between one peak and its preceding peak. Data were down-

1119 sampled to 100Hz for subsequent analyses.

1120

Analysis of fiber photometry data.

Offline using a custom script in Spike 2, the GFP and UV signals were filtered (2^{nd} order, 25Hz

low pass) and smoothed with a sliding window (9 bins averaged). A first-degree polynomial fit

was used to find the relationship between the signal and control channels and generate a scaled

control channel. The fitted control channel was subtracted from the signal channel and z-scored

$\Delta F/F$ traces were calculated using a 3s pre-event baseline window ($GRAB_{DA}$ analysis) or 2s pre-

event baseline window (GCaMP6f for D1- and D2-Cre analysis).

1128

1129 Based on a series of pilot experiments, we found that on average, tethered animals in the

1130 plethysmograph typically do not perceive odors until 3s following odor control valve opening (as

1131 indicated by their reflexive/orienting sniffing responses). To account for this delay, the 3s after

1132 valve opening and odor onset serve as the pre-event baseline. Buzz-evoked $GRAB_{DA}$ signals were

1133 analyzed relative to the buzz onset, and spontaneous sniffing was analyzed relative to bout onset.

1134

Peak GRAB$_{DA}$ signals and average sniff frequency were calculated within the 4s following odor

analysis onset, 2s buzz duration, or spontaneous bout. Spontaneous sniffing bout durations varied

and were identified based on prolonged periods of sniffing at frequencies ≥6Hz for ≥1s. Given the

highly dynamic nature of sniffing and the rapid shift between respiratory frequencies in mice, we

identified individual spontaneous sniff bouts wherein the mouse was sniffing at frequencies <6Hz

for at least 2s prior to bout onset.

1141

1142 For D1- and D2-Cre GCaMP6f photometry, because animals were head-fixed and odor was

1143 delivered directly to the nose with high temporal precision, the 2s before odor delivery was used

1144 as the trial specific baseline for the z-scored ΔF/F values.

1145

1146 *Additional statistical methods*

1147 Data were analyzed in GraphPad Prism. All performed statistical tests were two-sided. Sample

1148 sizes are consistent with those reported in the field. All data are reported as mean ± SEM unless

1149 otherwise reported. Specific tests and *p* values used can be found in the Results section or the

1150 figure legends.

References

1. Wachowiak, M. All in a Sniff: Olfaction as a Model for Active Sensing. *Neuron* **71**, 962–973 (2011).

2. Uchida, N., Kepecs, A. & Mainen, Z. F. Seeing at a glance, smelling in a whiff: rapid forms of perceptual decision making. *Nat Rev Neurosci* **7**, 485–491 (2006).

3. Welker, W. I. Analysis of sniffing in the albino rat. *Behavior* **22**, 223–244 (1964).

4. Youngentob, S. L., Mozell, M. M., Sheehe, P. R. & Hornung, D. E. A quantitative analysis of sniffing strategies in rats performing odor discrimination tasks. *Physiol Behav* **41**, 59–69 (1987).

5. Kepecs, A., Uchida, N. & Mainen, Z. F. Rapid and precise control of sniffing during olfactory discrimination in rats. *J Neurophysiol* **98**, 205–213 (2007).

6. Wesson, D. W., Donahou, T. N., Johnson, M. O. & Wachowiak, M. Sniffing behavior of mice during performance in odor-guided tasks. *Chem Senses* **33**, 581–596 (2008).

7. Clarke, S. & Trowill, J. A. Sniffing and motivated behavior in the rat. *Physiol Behav* **6**, 49–52 (1971).

8. Isles, A. R., Baum, M. J., Ma, D., Keverne, E. B. & Allen, N. D. Urinary odour preferences in mice. *Nature* **409**, 783–784 (2001).

9. Ikemoto, S. & Panksepp, J. The relationship between self-stimulation and sniffing in rats: does a common brain system mediate these behaviors? *Behav Brain Res* **61**, 143–162 (1994).

10. Catania, K. C. Olfaction: Underwater 'sniffing' by semi-aquatic mammals. *Nature* **444**, 1024–1025 (2006).

11. Moberly, A. H. *et al.* Olfactory inputs modulate respiration-related rhythmic activity in the prefrontal cortex and freezing behavior. *Nat Commun* **9**, 1528 (2018).

12. Ashhad, S., Kam, K., Del Negro, C. A. & Feldman, J. L. Breathing Rhythm and Pattern and Their Influence on Emotion. *Annu Rev Neurosci* **45**, 223–247 (2022).

13. Moore, J. D. *et al.* Hierarchy of orofacial rhythms revealed through whisking and breathing. *Nature* **advance on**, (2013).

14. Kleinfeld, D., Deschênes, M. & Ulanovsky, N. Whisking, Sniffing, and the Hippocampal θ-Rhythm: A Tale of Two Oscillators. *PLoS Biology* vol. 14 Preprint at https://doi.org/10.1371/journal.pbio.1002385 (2016).

15. Ramirez, J.-M. & Baertsch, N. A. The Dynamic Basis of Respiratory Rhythm Generation: One Breath at a Time. *Annu Rev Neurosci* **41**, 475–499 (2018).

16. Del Negro, C. A., Funk, G. D. & Feldman, J. L. Breathing matters. *Nat Rev Neurosci* **19**, 351–367 (2018).

17. Moore, J. D. *et al.* Hierarchy of orofacial rhythms revealed through whisking and breathing. *Nature* **497**, 205–10 (2013).

18. Vann, N. C., Pham, F. D., Dorst, K. E. & Del Negro, C. A. Dbx1 Pre-Bötzinger Complex Interneurons Comprise the Core Inspiratory Oscillator for Breathing in Unanesthetized Adult Mice. *eNeuro* **5**, (2018).

19. de Sousa Abreu, R. P., Bondarenko, E. & Feldman, J. L. Phase- and state-dependent modulation of breathing pattern by preBötzinger complex somatostatin expressing neurons. *J Physiol* **600**, 143–165 (2022).

20. Oliveira, L. M., Baertsch, N. A., Moreira, T. S., Ramirez, J.-M. & Takakura, A. C. Unraveling the Mechanisms Underlying Irregularities in Inspiratory Rhythm Generation in a Mouse Model of Parkinson's Disease. *J Neurosci* **41**, 4732–4747 (2021).

21. Arthurs, J. W., Bowen, A. J., Palmiter, R. D. & Baertsch, N. A. Parabrachial tachykinin1-expressing neurons involved in state-dependent breathing control. *Nat Commun* **14**, 963 (2023).

22. Bhandare, A. *et al.* Analyzing the brainstem circuits for respiratory chemosensitivity in freely moving mice. *Elife* **11**, (2022).

23. Rosin, D. L., Chang, D. A. & Guyenet, P. G. Afferent and efferent connections of the rat retrotrapezoid nucleus. *J Comp Neurol* **499**, 64–89 (2006).

24. Bhandare, A. *et al.* Analyzing the brainstem circuits for respiratory chemosensitivity in freely
moving mice. *Elife* **11**, (2022).
25. Rossi, J. & Panksepp, J. Analysis of the relationships between self-stimulation sniffing and brain-
stimulation sniffing. *Physiol Behav* **51**, 805–813 (1992).
26. Mogenson, G., Jones, D. & Yim, C. From motivation to action: Functional interface between the
limbic system and the motor system. *Prog Neurobiol* **14**, 69–97 (1980).
27. Haber, S. N. *Neuroanatomy of Reward: A View from the Ventral Striatum. Neurobiology of
Sensation and Reward* (CRC Press, Boca Raton (FL), 2011).
28. Ikemoto, S. Dopamine reward circuitry: Two projection systems from the ventral midbrain to the
nucleus accumbens-olfactory tubercle complex. *Brain Res Rev* **56**, 27–78 (2007).
29. Mingote, S. *et al.* Functional Connectome Analysis of Dopamine Neuron Glutamatergic
Connections in Forebrain Regions. *Journal of Neuroscience* **35**, 16259–16271 (2015).
30. Wieland, S. *et al.* Phasic Dopamine Modifies Sensory-Driven Output of Striatal Neurons through
Synaptic Plasticity. *The Journal of Neuroscience* **35**, 9946–9956 (2015).
31. Zahm, D. S. & Heimer, L. The ventral striatopallidothalamic projection. III. Striatal cells of the
olfactory tubercle establish direct synaptic contact with ventral pallidal cells projecting to
mediodorsal thalamus. *Brain Res* **404**, 327–331 (1987).
32. Menegas, W., Babayan, B. M., Uchida, N. & Watabe-Uchida, M. Opposite initialization to novel
cues in dopamine signaling in ventral and posterior striatum in mice. *Elife* **6**, e21886 (2017).
33. Wise, R. A. Dopamine, learning and motivation. *Nat Rev Neurosci* **5**, 483–494 (2004).
34. Morales, M. & Margolis, E. B. Ventral tegmental area: cellular heterogeneity, connectivity and
behaviour. *Nat Rev Neurosci* **18**, 73–85 (2017).
35. Stuber, G. D., Hnasko, T. S., Britt, J. P., Edwards, R. H. & Bonci, A. Dopaminergic Terminals in
the Nucleus Accumbens But Not the Dorsal Striatum Corelease Glutamate. *Journal of
Neuroscience* (2010) doi:10.1523/jneurosci.1754-10.2010.
36. Lobo, M. K. & Nestler, E. J. The Striatal Balancing Act in Drug Addiction: Distinct Roles of
Direct and Indirect Pathway Medium Spiny Neurons. *Front Neuroanat* **5**, 41 (2011).
37. Phillips, P. E. M., Stuber, G. D., Heien, M. L. A. V, Wightman, R. M. & Carelli, R. M. Subsecond
dopamine release promotes cocaine seeking. *Nature* **422**, 614–618 (2003).
38. Murata, K. *et al.* Opposing roles of dopamine receptor D1- and D2-expressing neurons in the
anteromedial olfactory tubercle in acquisition of place preference in mice. *Front Behav Neurosci*
13, (2019).
39. Zhang, Z. *et al.* Activation of the dopaminergic pathway from VTA to the medial olfactory
tubercle generates odor-preference and reward. *Elife* **6**, e25423 (2017).
40. Gadziola, M. A. *et al.* A neural system for encoding the association of odors with rewarded
outcomes and promoted behavioral engagement. *Cell Rep* **32**, 107919 (2020).
41. Li, W.-R. *et al.* Neural mechanisms underlying uninstructed orofacial movements during reward-
based learning behaviors. *Current Biology* (2023) doi:10.1016/j.cub.2023.07.013.
42. Watabe-Uchida, M., Zhu, L., Ogawa, S. K., Vamanrao, A. & Uchida, N. Whole-Brain Mapping of
Direct Inputs to Midbrain Dopamine Neurons. *Neuron* **74**, 858–873 (2012).
43. Beier, K. T. *et al.* Circuit Architecture of VTA Dopamine Neurons Revealed by Systematic Input-
Output Mapping. *Cell* **162**, 622–634 (2015).
44. Badrinarayan, A. *et al.* Aversive Stimuli Differentially Modulate Real-Time Dopamine
Transmission Dynamics within the Nucleus Accumbens Core and Shell. *The Journal of
Neuroscience* **32**, 15779–15790 (2012).
45. Lammel, S., Ion, D. I., Roeper, J. & Malenka, R. C. Projection-Specific Modulation of Dopamine
Neuron Synapses by Aversive and Rewarding Stimuli. *Neuron* **70**, 855–862 (2011).
46. Floresco, S. B. The Nucleus Accumbens: An Interface Between Cognition, Emotion, and Action.
Annu Rev Psychol **66**, 25–52 (2015).
47. Wesson, D. W. The Tubular Striatum. *Journal of Neuroscience* **40**, 7379–7386 (2020).

48.   Ikemoto, S. Involvement of the Olfactory Tubercle in Cocaine Reward: Intracranial Self-
Administration Studies. *The Journal of neuroscience* **23**, 9305–9311 (2003).

49.   Murata, K., Kanno, M., Ieki, N., Mori, K. & Yamaguchi, M. Mapping of Learned Odor-Induced
Motivated Behaviors in the Mouse Olfactory Tubercle. *The Journal of Neuroscience* **35**, 10581–
10599 (2015).

50.   Sun, F. *et al.* A Genetically Encoded Fluorescent Sensor Enables Rapid and Specific Detection of
Dopamine in Flies, Fish, and Mice. *Cell* **174**, 481-496.e19 (2018).

51.   Mandairon, N., Poncelet, J., Bensafi, M. & Didier, A. Humans and Mice Express Similar
Olfactory Preferences. *PLoS One* **4**, e4209 (2009).

52.   Jagetia, S. *et al.* Inter- and intra-mouse variability in odor preferences revealed in an olfactory
multiple-choice test. *Behavioral Neuroscience* **132**, 88–98 (2018).

53.   Midroit, M. *et al.* Neural processing of the reward value of pleasant odorants. *Current Biology* **31**,
1592–1605 (2021).

54.   Oettl, L.-L. *et al.* Phasic dopamine reinforces distinct striatal stimulus encoding in the olfactory
tubercle driving dopaminergic reward prediction. *Nat Commun* **11**, 3460 (2020).

55.   Gonzalez, L. S. *et al.* Ventral striatum dopamine release encodes unique properties of visual
stimuli in mice. *Elife* **12**, (2023).

56.   Sulzer, D. *et al.* Dopamine Neurons Make Glutamatergic Synapses in vitro. *The Journal of*
Neuroscience **18**, 4588 LP – 4602 (1998).

57.   Hnasko, T. S. *et al.* Vesicular glutamate transport promotes dopamine storage and glutamate
corelease in vivo. *Neuron* **65**, 643–656 (2010).

58.   Liu, S. *et al.* Divergent brainstem opioidergic pathways that coordinate breathing with pain and
emotions. *Neuron* **110**, 857-873.e9 (2022).

59.   Sakoe, H. & Chiba, S. Dynamic programming algorithm optimization for spoken word
recognition. *IEEE Trans Acoust* **26**, 43–49 (1978).

60.   Bagnall, A., Lines, J., Bostrom, A., Large, J. & Keogh, E. The great time series classification bake
off: a review and experimental evaluation of recent algorithmic advances. *Data Min Knowl Discov*
31, 606–660 (2017).

61.   Regan, M. K-Nearest-Neighbors-with-Dynamic-Time-Warping. https://github.com/markdregan/K-
Nearest-Neighbors-with-Dynamic-Time-Warping (2018).

62.   Chawla, N. V, Bowyer, K. W., Hall, L. O. & Kegelmeyer, W. P. SMOTE: synthetic minority over-
sampling technique. *J. Artif. Int. Res.* **16**, 321–357 (2002).

63.   Zhang, F. *et al.* Multimodal fast optical interrogation of neural circuitry. *Nature* **446**, 633–639
(2007).

64.   Gadziola, M. A., Tylicki, K. A., Christian, D. L. & Wesson, D. W. The Olfactory Tubercle
Encodes Odor Valence in Behaving Mice. *The Journal of Neuroscience* **35**, 4515–4527 (2015).

65.   Wesson, D. W., Verhagen, J. V & Wachowiak, M. Why Sniff Fast? The Relationship Between
Sniff Frequency, Odor Discrimination, and Receptor Neuron Activation in the Rat. *J Neurophysiol*
101, 1089–1102 (2009).

66.   Verhagen, J. V, Wesson, D. W., Netoff, T. I., White, J. A. & Wachowiak, M. Sniffing controls an
adaptive filter of sensory input to the olfactory bulb. *Nat Neurosci* **10**, 631–639 (2007).

67.   Gong, S. *et al.* Targeting Cre Recombinase to Specific Neuron Populations with Bacterial
Artificial Chromosome Constructs. *The Journal of Neuroscience* **27**, 9817 LP – 9823 (2007).

68.   Jordan, R., Kollo, M. & Schaefer, A. T. Sniffing Fast: Paradoxical Effects on Odor Concentration
Discrimination at the Levels of Olfactory Bulb Output and Behavior. *eNeuro* **5**, ENEURO.0148-
18.2018 (2018).

69.   Scofield, M. D. *et al.* The Nucleus Accumbens: Mechanisms of Addiction across Drug Classes
Reflect the Importance of Glutamate Homeostasis. *Pharmacol Rev* **68**, 816–871 (2016).

70.   Le Moine, C. & Bloch, B. D1 and D2 dopamine receptor gene expression in the rat striatum:
Sensitive cRNA probes demonstrate prominent segregation of D1 and D2 mRNAS in distinct
neuronal populations of the dorsal and ventral striatum. *J Comp Neurol* **355**, 418–426 (1995).

71. Mansour, A. *et al.* A comparison of D1 receptor binding and mRNA in rat brain using receptor autoradiographic and in situ hybridization techniques. *Neuroscience* **45**, 359–371 (1991).

72. Bouthenet, M. L. *et al.* Localization of dopamine D3 receptor mRNA in the rat brain using in situ hybridization histochemistry: comparison with dopamine D2 receptor mRNA. *Brain Res* **564**, 203–219 (1991).

73. Cepeda, C. *et al.* Differential electrophysiological properties of dopamine D1 and D2 receptor-containing striatal medium-sized spiny neurons. *European Journal of Neuroscience* **27**, 671–682 (2008).

74. Zhang, Y.-F. *et al.* Ventral striatal islands of Calleja neurons control grooming in mice. *Nat Neurosci* **24**, 1699–1710 (2021).

75. Starr, B. S. & Starr, M. S. Differential effects of dopamine D1 and D2 agonists and antagonists on velocity of movement, rearing and grooming in the mouse. Implications for the roles of D1 and D2 receptors. *Neuropharmacology* **25**, 455–463 (1986).

76. Markowitz, J. E. *et al.* Spontaneous behaviour is structured by reinforcement without explicit reward. *Nature* **614**, 108–117 (2023).

77. Berke, J. D. What does dopamine mean? *Nature Neuroscience* vol. 21 787–793 Preprint at https://doi.org/10.1038/s41593-018-0152-y (2018).

78. Ennis, M. *et al.* Dopamine D2 receptor-mediated presynaptic inhibition of olfactory nerve terminals. *J Neurophysiol* **86**, 2986–2997 (2001).

79. Nieoullon, A. Dopamine and the regulation of cognition and attention. *Prog Neurobiol* **67**, 53–83 (2002).

80. Bromberg-Martin, E. S., Matsumoto, M. & Hikosaka, O. Dopamine in Motivational Control: Rewarding, Aversive, and Alerting. *Neuron* **68**, 815–834 (2010).

81. Westbrook, A. *et al.* Dopamine promotes cognitive effort by biasing the benefits versus costs of cognitive work. *Science (1979)* **367**, 1362–1366 (2020).

82. Legault, M. & Wise, R. A. Novelty-evoked elevations of nucleus accumbens dopamine: Dependence on impulse flow from the ventral subiculum and glutamatergic neurotransmission in the ventral tegmental area. *European Journal of Neuroscience* **13**, 819–828 (2001).

83. Madras, B. K., Miller, G. M. & Fischman, A. J. The dopamine transporter and attention-deficit/hyperactivity disorder. *Biol Psychiatry* **57**, 1397–409 (2005).

84. Dang, L. C., O'Neil, J. P. & Jagust, W. J. Dopamine supports coupling of attention-related networks. *J Neurosci* **32**, 9582–7 (2012).

85. Motaghi, S., Sheibani, V., Farazifard, R. & Joneidi, H. Electrical stimulation of locus coeruleus strengthens the surround inhibition in layer V barrel cortex in rat. *Neurosci Lett* **401**, 280–284 (2006).

86. Lee, V., Pawlisch, B. A., Macedo-Lima, M. & Remage-Healey, L. Norepinephrine enhances song responsiveness and encoding in the auditory forebrain of male zebra finches. *J Neurophysiol* **119**, 209–220 (2018).

87. Zhao, Y., Kerscher, N., Eysel, U. & Funke, K. D1 and D2 receptor-mediated dopaminergic modulation of visual responses in cat dorsal lateral geniculate nucleus. *J Physiol* **539**, 223–238 (2002).

88. Chen, K., Vincis, R. & Fontanini, A. Disruption of Cortical Dopaminergic Modulation Impairs Preparatory Activity and Delays Licking Initiation. *Cerebral Cortex* **29**, 1802–1815 (2019).

89. Hattox, A., Li, Y. & Keller, A. Serotonin Regulates Rhythmic Whisking. *Neuron* **39**, 343–352 (2003).

90. Shusterman, R., Smear, M. C., Koulakov, A. A. & Rinberg, D. Precise olfactory responses tile the sniff cycle. *Nat Neurosci* **14**, 1039–1044 (2011).

91. Cury, K. M. & Uchida, N. Robust Odor Coding via Inhalation-Coupled Transient Activity in the Mammalian Olfactory Bulb. *Neuron* **68**, 570–585 (2010).

92. Iwata, R., Kiyonari, H. & Imai, T. Mechanosensory-Based Phase Coding of Odor Identity in the Olfactory Bulb. *Neuron* **96**, 1139-1152.e7 (2017).

93. Linster, C. & Cleland, T. A. Neuromodulation of olfactory transformations. *Curr Opin Neurobiol* **40**, 170–177 (2016).

94. Wesson, D. W., Carey, R. M., Verhagen, J. V & Wachowiak, M. Rapid encoding and perception of novel odors in the rat. *PLoS Biol* **6**, e82 (2008).

95. Yang, H. *et al.* Nucleus Accumbens Subnuclei Regulate Motivated Behavior via Direct Inhibition and Disinhibition of VTA Dopamine Subpopulations. *Neuron* **97**, 434-449.e4 (2018).

96. Tapper, A. R. & Molas, S. Midbrain circuits of novelty processing. *Neurobiol Learn Mem* **176**, 107323 (2020).

97. Azcorra, M. *et al.* Unique functional responses differentially map onto genetic subtypes of dopamine neurons. *Nat Neurosci* **26**, 1762–1774 (2023).

98. Steen, J. B., Mohus, I., Kvesetberg, T. K. & Walloe, L. Olfaction in bird dogs during hunting. *Acta. Physiol Scand* **157**, 115–119 (1996).

99. Angelakos, C. C. *et al.* A cluster of neuropeptide S neurons regulates breathing and arousal. *Current Biology* **33**, 5439-5455.e7 (2023).

100. Schultz, W., Dayan, P. & Montague, P. R. A Neural Substrate of Prediction and Reward. *Science (1979)* **275**, 1593–1599 (1997).

101. Grace, A. & Bunney, B. The control of firing pattern in nigral dopamine neurons: burst firing. *The Journal of Neuroscience* **4**, 2877–2890 (1984).

102. Sugita, T. *et al.* Breathing is affected by dopamine D2-like receptors in the basolateral amygdala. *Respir Physiol Neurobiol* **209**, 23–27 (2015).

103. Jhang, J., Liu, S., O'Keefe, D. D. & Han, S. A top-down slow breathing circuit that alleviates negative affect. *bioRxiv* (2023).

104. de Sousa Abreu, R. P., Bondarenko, E. & Feldman, J. L. Phase- and state-dependent modulation of breathing pattern by preBötzinger complex somatostatin expressing neurons. *J Physiol* **600**, 143–165 (2022).

105. Vann, N. C., Pham, F. D., Dorst, K. E. & Del Negro, C. A. Dbx1 Pre-Bötzinger Complex Interneurons Comprise the Core Inspiratory Oscillator for Breathing in Unanesthetized Adult Mice. *eNeuro* **5**, ENEURO.0130-18.2018 (2018).

106. Zhang, Y.-F. *et al.* Ventral striatal islands of Calleja neurons bidirectionally mediate depression-like behaviors in mice. *Nat Commun* **14**, 6887 (2023).

107. Edwards, N. J. *et al.* Circuit specificity in the inhibitory architecture of the VTA regulates cocaine-induced behavior. *Nat Neurosci* **20**, 438 (2017).

108. Bäckman, C. M. *et al.* Characterization of a mouse strain expressing Cre recombinase from the 3′ untranslated region of the dopamine transporter locus. *genesis* **44**, 383–390 (2006).

109. Savitt, J. M., Jang, S. S., Mu, W., Dawson, V. L. & Dawson, T. M. Bcl-x Is Required for Proper Development of the Mouse Substantia Nigra. *The Journal of Neuroscience* **25**, 6721–6728 (2005).

110. Tong, Q. *et al.* Synaptic Glutamate Release by Ventromedial Hypothalamic Neurons Is Part of the Neurocircuitry that Prevents Hypoglycemia. *Cell Metab* **5**, 383–393 (2007).

111. Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nat Neurosci* **13**, 133–140 (2010).

112. Paxinos, G. & Franklin, K. *The Mouse Brain in Stereotaxic Coordinates*. (Academic Press, San Diego, 2000).

113. Martin-Lopez, E., Xu, C., Liberia, T., Meller, S. J. & Greer, C. A. Embryonic and postnatal development of mouse olfactory tubercle. *Molecular and Cellular Neuroscience* **11**, 82–96 (2019).

114. Lammel, S. *et al.* Diversity of Transgenic Mouse Models for Selective Targeting of Midbrain Dopamine Neurons. *Neuron* **85**, 429–438 (2015).

Acknowledgements:

We thank Jacqueline Sipio for help with histology and Ellyse Thomas for expert animal care assistance.

This work was supported by R01DC014443 and R01DC016519 to D.W.W., R01DA049545, R01DA049449 and R01NS117061 to D.W.W. and M.M., R01DC006213 to M.M., and R00HL159232 to A.G.V.. N.L.J. was supported by NIDCD T32015994 and F31DC020364.

Author Contributions:

Conceptualization: NLJ & DWW
Methodology: NLJ, ACL, LAS, AGV, UMA, ZZ, MAG
Investigation: NLJ, ACL, LAS, ZZ, UMA, MAG
Visualization: NLJ, DWW, ACL
Funding acquisition: NLJ, DWW, MM, AGV
Supervision: DWW
Writing – original draft: NLJ
Writing – review & editing: NLJ, DWW, AGV, MM, UMA, LAS, ACL, ZZ

Competing interests: Authors declare that they have no competing interests.

 Figures & Legends

1422

1423

Figure 1. Organization of midbrain dopamine input into the tubular striatum and nucleus
accumbens. A. Schematic of tracing paradigm. *DAT^{IRES} -Cre* mice were injected into their VTA
with AAV.FLEX.hSyn.mGFP.synaptophysin.mRuby allowing for mGFP expression in DAT+ cell
bodies and synaptophysin.mRuby+ puncta in the terminals of DAT+ cells. Approximately 3 weeks
later, coronal sections throughout the ventral striatum were collected for later fluorescence imaging
and quantification of synaptophysin.mRuby+ puncta. **B.** Example of synaptophysin.mRuby
expression throughout the ventral striatum, with key regions of interest outlined by dashed lines.
C. Quantification of synaptophysin+ puncta throughout the anterior-posterior spans of regions of
interest (*n*=6 mice). Data [puncta #'s/mm^2] are represented as mean ± SEM. Data smoothed for
visualization purposes only. **D.** Histograms representing data in (**C**) with regions of interest
segregated into their most anterior and posterior aspects (anterior most half vs. posterior most half).
Data are mean ± SEM, points = individual mice. *$p<0.5$. abbreviations: VTA (ventral tegmental
area), NAcSh (nucleus accumbens shell), NAcC (nucleus accumbens core), mTuS (medial aspect
of the tubular striatum), LatTuS (lateral aspect of the tubular striatum), PCX (piriform cortex). (the
anterior or posterior-most 200μm were used for quantification of these regions).

1439

Figure 2. Self-motivated spontaneous sniffing is associated with DA levels. A. Diagram of the
setup for simultaneous measurement of respiration by means of both whole-body plethysmography
and an intranasal pressure cannula. Three mice were implanted with a cannula in its dorsal nasal
recess and placed in the plethysmograph. Intranasal pressure changes were monitored by a pressure
sensor connected to the cannula by a flexible tube passed through a fitted plug at the top of the
plethysmograph and compared to whole-body changes. Image made with BioRender. **Bi**. Overlaid
representative respiratory traces acquired from the plethysmograph (black) and intranasal cannula
(blue) from an awake mouse. An example of high frequency sniffing is shown zoomed in and the
expanded inset shows fidelity between the whole-body pressure and intranasal pressure signal. **Bii.**
Example of the latency difference between respiratory peaks recorded by intranasal pressure and
plethysmograph pressure during a high frequency sniffing bout with detected respiratory events
(*n*=125 respiratory cycles). Overlaid respiratory traces (top) shows an intranasal pressure peaks
aligned to time 0, and a plethysmograph pressure peak detected <60ms later. **Biii**. Histograms
comparing respiratory frequencies measured from the plethysmograph and intranasal cannula. The
plethysmograph reliably detected sniffing at both low and high respiratory frequencies with >99%
accuracy (out of 3,239 respiratory events collected from 3 mice). **C.** Schematic of fiber photometry
and plethysmograph system used to simultaneously record GRAB$_{DA}$ signals and sniffing behavior
in unrestrained mice. Animals were injected with AAV.hSyn-GRAB_DA1h and implanted with
an optic fiber in the TuS, NAcSh, or NAcC. 3-5 weeks later, animals were placed in the
plethysmograph and 465nm and 405nm light were passed through dichroic mirrors before reaching
the animal via patch cable and implanted optical fiber. GRAB$_{DA}$ and UV emissions were amplified
by photoreceivers before being digitized along with simultaneous respiratory data. Image made
with BioRender. **D.** Optic fiber implant locations in the TuS, NAcSh, and NAcC (*n*=7 mice/group).
Within each region, ellipse color corresponds to same colored data in **Figures 2F-H and 3C-H**.
E. Example of raw 465nm GRAB$_{DA}$ (green) and 405nm UV (purple) signals recorded in the TuS
during a spontaneous sniff bout. Raw respiratory trace (resp) is shown below with interpolated
instantaneous frequency above (Inst Hz- black trace). Sniff duration is indicated by the grey shaded

box. **Fi.** Averaged z-scored ΔF/F traces show an increase in GRAB$_{DA}$ levels in TuS during spontaneous sniff bouts. Average bout duration indicated by overlaid box plot. **Fii**. Significant positive correlation between GRAB$_{DA}$ levels in the TuS (the peak z-score during a spontaneous sniff bout) and average sniff frequency during bout (r=0.29, p=0.008). **Gi**. Averaged z-scored ΔF/F traces similarly show an increase in GRAB$_{DA}$ levels in the NAcSh during spontaneous sniff bouts. Average bout duration indicated by overlaid box plot. **Gii**. There was no correlation between GRAB$_{DA}$ levels in the NAcSh during spontaneous bout and sniffing frequency (r=0.12, p=0.249). **Hi**. Conversely, averaged z-scored ΔF/F traces show a decrease in GRAB$_{DA}$ levels in the NAcC during spontaneous sniff bouts. Average bout duration indicated by overlaid box plot. **Hii**. There was no significant correlation between GRAB$_{DA}$ levels in the NAcC during spontaneous bout and sniffing frequency (r=0.038, p=0.703). Abbreviations: resp (respiratory trace), Inst Hz (instantaneous frequency). Data in (Fi), (Gi), and (Hi) smoothed for visualization purposes only.

Figure 3. Bidirectional dopamine dynamics upon sensory-driven sniffing. A. Schematic of fiber photometry and plethysmograph system used to simultaneously record GRAB$_{DA}$ signals in the TuS, NAcSh, and NAcC and sniffing behavior in unrestrained mice (n=7/group) as used in Fig 2. Additionally, an olfactometer was used to deliver odors to animals in the plethysmograph, and a vibration motor (buzz) was attached to the outside of the plethysmograph. Image made with BioRender. **B.** Example of raw 465nm GRAB$_{DA}$ (green) and 405nm UV (purple) signals recorded in the TuS during an odor-evoked bout. Raw respiratory trace (resp) is shown below with interpolated instantaneous frequency above (Inst Hz- black trace). Grey shaded box denotes odor analysis window, starting 3 seconds following odor onset to account for the delay of odor delivery to the plethysmograph. **C-E.** Odor-evoked sniffing behavior and GRAB$_{DA}$ signals from mice with injections and optic ferrule implants in the TuS (C), NACSh (D), and NAcC (E). Average instantaneous sniffing frequency of TuS (Ci), NAcSh (Di), and NAcC (Ei) implanted mice across repeated odor presentations. Mean shown in black with individual mouse data across trials shown in light grey. Averaged z-scored ΔF/F across all 5 odor presentations in the TuS (Cii), NAcSh (Dii), and NAcC (Eii). Averaged peak of the z-scored GRAB$_{DA}$ response across trials in the TuS (Ciii), NAcSh (Diii), and NAcC (Eiii). Mean shown in black with individual mouse data across trials color coded to ellipse colors shown in **Figure 2D**. Significant positive correlations between GRAB$_{DA}$ levels (peak z-score during odor-evoked activity) and average sniff frequency during bout in the TuS, (r=0.65, p<0.0001, Civ) and NAcSh (r=0.51, p=0.002, Div) but not in the NAcC (r=-0.14, p=0.414, Eiv). **F-H.** Buzz-evoked sniffing behavior and GRAB$_{DA}$ signals from same mice with injections and implants in the TuS (F), NAcSh (G), and NAcC (H). Average instantaneous sniffing frequency of TuS (Fi), NAcSh (Gi), and NAcC (Hi) implanted mice across repeated odor presentations. Mean shown in black with individual mouse data across trials shown in light grey. Averaged z-scored ΔF/F across all 5 odor presentations in the TuS (Fii), NAcSh (Gii), and NAcC (Hii). Horizontal line depicts 2s buzz duration. Averaged peak of the z-scored GRAB$_{DA}$ response across trials in the TuS (Fiii), NAcSh (Giii), and NAcC (Hiii). Mean shown in black with individual mouse data across trials color coded to ellipse colors shown in **Figure 2D**. Significant positive correlation between GRAB$_{DA}$ levels and sniffing frequency in the TuS, (r=0.43, p=0.011, Fiv), but not the NAcSh (during buzz: r=-0.17, p=0.316; upon buzz offset: r=-0.074, p=0.673, Giv) or NAcC (r=-0.22, p=0.194, Hiv). **I.** Comparison of the peak-evoked GRAB$_{DA}$ responses across the TuS, NAcSh, and NAcC during odor-, buzz-, and spontaneously-evoked sniffing. Data are represented as mean ± SEM. Points = individual mice. ****p<0.0001,

***p<0.001, **p<0.01. Data shown in (Cii), (Dii), (Eii), (Fii), (Gii), and (Hii) are smoothed for visualization purposes only.

Figure 4. Initiation of sniffing by dopamine release into the ventral striatum. A. Schematic of optogenetically-evoked sniffing paradigm. *DAT^{IRES}-Cre x Ai9* mice were injected with AAV-EF1α-DIO-hChR2(E123T/T159C)-EYFP in the VTA, and optic fibers terminating in the TuS, NAcSh or NAcC were implanted in the same surgery. Control mice were injected with AAV-Ef1α-DIO-EYFP. At least 4 weeks later, mice were tethered and placed in the plethsymograph to optogenetically stimulate DA terminal release while monitoring sniffing. Image (right) made with BioRender. **B.** Optic fiber implant locations in the TuS, NAcSh, and NAcC (*n*=6-11 mice/group). **C.** Example respiratory data from a TuS implanted mouse across 10 trials of light stimulation. Timing of the 1s long, 25Hz photostimulation depicted by grey shaded box and dotted blue horizontal line. Raw respiratory traces are shown on the left, and corresponding instantaneous frequency for each trial shown in 2-dimensional histograms on the right. Dotted white line in 2-dimenional histogram depicts stimulation onset. **D**. 2-dimensional histograms across all mice tested within each group, showing individual trial data within mice. Example data from (C) denoted by the white star. Separation between mice marked by dark blue line. Stimulation onset is denoted by the vertical dotted white line, and 1s, 25Hz stimulation shown by the dotted blue trace. **E.** Mean sniffing frequency ± SEM during photostimulation across 10 trials in each group. **F.** Averaged z-scored respiratory frequency ± SEM of photostimulation-evoked sniffing. Stimulation onset denoted by vertical dotted black line and 25Hz, 1s duration by horizontal blue line. Group color legend matches what is shown in (E). **G**. Parameters of optogenetically-evoked sniffing during 1s photostimulation. All data presented as mean ± SEM, points = individual mice. ****p<0.0001, ***p<0.001, **p<0.01. **Gi.** Mean latency to sniff across groups. Animals that failed to reach 6Hz on >40% of trials were excluded from the EYFP control group. **Gii**. Mean percentage of time spent sniffing across groups. **Giii.** Max sniffing frequency reached during photostimulation. **Giv**. Mean probability that photostimulation will evoke sniffing. **H.** Averaged z-scored respiratory frequency during different photostimulation durations (100ms, 1s, 2s, 4s) compared to responses of EYFP controls stimulated for 1s in the TuS (Hi), NAcSh (Hii), and NAcC (Hiii). Insets in each panel show the percentage of time across all groups, calculated in a 4s window, that mice spend sniffing upon photostimulation.

Figure 5. State-dependence of a unique dopamine-evoked sniffing pattern. A. Example respiratory behavior of a TuS implanted mouse under urethane anesthesia. Raw respiratory traces (left) and corresponding 2-dimensional histogram (right) across 10 trials of photostimulation. 1s long, 25Hz photostimulation duration depicted by grey shaded box and dotted blue horizontal line. Dotted white line in the 2-dimensional histogram = photostimulation onset **B.** 2-dimensional histograms across 5 representative mice per group showing individual trial data within mice. Starred data represents example data from (A). Separation between mice denoted by dark blue line. Stimulation onset is denoted by the vertical dotted white line, and 1s, 25Hz stimulation shown by the dotted blue trace. Color scale matches that of **Figures 4C and D**. Little to no change in coloring highlights the substantial lack of any modulation in breathing rate under anesthesia. **C.** Averaged z-scored respiratory frequency of photostimulation-evoked sniffing. Stimulation onset denoted by vertical dotted black line and duration by horizontal blue dotted line. Inset shows the evoked z-scored average across all groups during light stimulation. **D.** Overview of machine learning applied to respiratory traces starting with dynamic time warping (DTW) between a 1s

odor-evoked sniff bout and 1s opto-evoked sniff bout. Bin size=10ms. 1s-long respiratory traces (100Hz) were converted to instantaneous frequency traces (also 1s long) and subjected to DTW to assess distance in proximal time-points. Subsequently, a model was trained on experimenter-identified traces from either buzz-, odor-, spontaneous-, or photostimulation-evoked respiration that were also previously subjected to DTW (see Methods). **Ei.** F1 score (single metric of model performance) for sniff types based on number of samples used per class. SMOTE=70 samples. **Eii.** Confusion matrix from SMOTE analyzed data displaying the classifier accuracy in predicting optogenetic-, odor-, buzz-, or spontaneously-evoked sniffing bouts. Classifier was trained on 80% of samples and tested on remaining 20%.

Figure 6. Inhibition of DA release into the ventral striatum and sniffing. A. Schematic of optogenetically-inhibited sniffing paradigm. DAT^{IRES}-*Cre* and DAT^{IRES}-*Cre x Ai9* mice were injected with AAV-EF1α-DIO-eNpHR 3.0-EYFP bilaterally in the VTA, and bilateral optic fibers terminating the TuS or NAcSh were implanted in the same surgery. At least 4 weeks later, mice were tethered and placed in the plethysmograph to optogenetically inhibit DA terminal release while monitoring sniffing. 560nm light was used to activate eNpHR and 405nm light was used for controls. Image (right) made with BioRender. **B.** Bilateral optic fiber implant locations in the TuS and NAcSh (*n*=5-6/group). **C.** Representative bilateral implant in the TuS. **Di.** Example of respiration during photoinhibition periods. Raw respiratory trace (resp) is shown with interpolated instantaneous frequency above (Inst Hz). 560nm LED duration indicated by red shaded box. **Dii.** Normalized respiratory frequency during the 3s before light stimulation compared to respiratory frequency during 5s light stimulation (405nm stimulation – left, 560nm stimulation – right). Blue dots indicate TuS implanted mice, green dots indicate NAcSh implanted mice. **p<0.01. **E.** Average instantaneous sniffing frequency of TuS (Ei) and NAcSh (Eii) implanted mice across repeated odor presentations with concurrent delivery of 560nm or 405nm light. Mean during 560nm concurrent light delivery shown in red, with individual mouse data across trials shown in light red. Mean during 405nm concurrent light delivery shown in gray, with individual mouse data across trials shown in light grey. **F.** Average instantaneous sniffing frequency of TuS (Fi) and NacSh (Fii) implanted mice across repeated buzz presentations with concurrent delivery of 560nm (red) or 405nm (gray) light.

Fig. 7. Ventral striatum neuron activity is coupled with sniffing. Ai. Continuous recording of intranasal respiration (blue, inhale = upward deflection) along with single-unit activity from three units in the TuS. Grey shaded boxes indicate periods wherein non-novel odors were delivered. **Aii** is a magnified inset to help illustrate the temporal relationship between the three units in **Ai** and sniffing. **B.** Overdrawn waveforms and raster plots of spiking aligned to sniffing onset from the three units in **A** to illustrate that spiking often occurred following sniffing onset. **C.** Distribution of respiratory frequencies from the mice (*n*=3) during both spontaneous and odor-delivery periods (see Methods). **D.** Spike ratio histogram showing that more spiking occurs (*n*=20 units, across 3 mice) during higher frequency respiration than lower frequency respiration, and that this was greater when animals spontaneously transitioned into sniffing during the inter-trial interval than when sniffing during an odor – likely because the different basal respiratory rates in those states (as shown in **C**). **Ei** and **Eii.** Histograms of the firing frequency of the units in D, relative to either the first spike or final spike (Ei or Eii, respectively) indicating that the first

1603 spike is predictive that the subsequent breath frequencies will be faster than the breaths prior to
1604 the first spike. *** = (ANOVA $F(6, 112)=4.43$, $p=0.0005$; see Results).

1605

1606

Figure 8. Sniffing is associated with recruitment of D1 and D2 spiny projection neurons. A.
Schematic of the fiber photometry system for use in head-fixed mice. D1- or D2- cre mice were
injected with AAV.hSyn.Flex.GCaMP6f.WPRE.SV40, and implanted with an optical fiber
terminating in the TuS. Mice also received a headbar for later fixation. Odors were delivered
directly to the animal's nose, and respiration was recorded with an extranasal flow sensor. Image
made with BioRender. **B.** Optic fiber implant locations in the TuS across D1- and D2-cre mice.
C. GCaMP activity of D1 and D2 neurons in the TuS during spontaneously-evoked sniffing. **Ci.**
Averaged z-scored ΔF/F of D1 and D2 neurons aligned to spontaneous sniff bout onset (D1, $n=4$,
2 points/mouse. D2, $n=5$, 2points/mouse. Overlaid box plots represent average bout duration.
Cii. Average peak of GCaMP evoked z-scored activity during spontaneous sniffing. **Ciii.** Mean
sniffing instantaneous frequency during spontaneous bouts. **Civ.** There was no correlation
between GCaMP levels in D1 neurons (peak z-score during a bout) during spontaneous sniffing
and (average sniff frequency during the bout, $p>0.05$. **Cv.** There was a significant positive
correlation between GCaMP levels in D2 neurons during spontaneous bouts and sniffing
frequency ($p<0.0001$). **D.** GCaMP activity of D1 and D2 neurons in the TuS during odor-evoked
sniffing. **Di.** Average instantaneous sniffing frequency of both D1-cre ($n=1$, 2 sessions/mouse)
and D2-cre ($n=5$, 2 sessions/mouse) mice across repeated odor presentations, with individual
mouse data across trials shown in light grey. **Dii.** Averaged z-scored ΔF/F across 4 odor
presentations in D1-mice. Odor duration depicted by horizontal line. **Diii.** Averaged peak of the
z-scored GCaMP response in D1-cre mice across trials. **Div.** Averaged z-scored ΔF/F across 4
odor presentations in D2-mice. Odor duration depicted by horizontal line. **Dv**. Averaged peak of
the z-scored GCaMP response in D2-cre mice across trials. **Dvi.** Overlayed averages of peak z-
score GCaMP responses from D1- and D2-cre mice across trials. Data in (Ci),(Dii) and (Div)
smoothed for visualization purposes only.

1631

Figure 9. Dopamine binding to D1 and D2 ventral striatum receptors contributes to sniffing.
A. Bilateral cannulae implant locations in the TuS and NAcSh. Colored for visualization purposes
only. Position relative to bregma. B. Diagram of pharmacological manipulation and behavioral
paradigm. Animals were implanted with stainless steel, bilateral cannulae with tips terminating in
the TuS ($n=13$) or NAcSh ($n=13$). After surgical recovery, animals underwent behavioral testing.
In a counterbalanced, within-subjects design, animals were administered an intracranial infusion
of a selective D1, D2, or D3 antagonist or corresponding vehicle control 10min prior to being
placed in the plethysmograph. Vehicle = sterile saline + 0-12% DMSO. During a behavioral
session, animals were presented with both olfactory and non-olfactory (buzz) stimuli. Image made
with BioRender. **C.** Impact of selective antagonism of D1- (Ci), D2- (Cii), and D3-receptors (Ciii)
on spontaneous sniffing. D3 antagonism had no effect on spontaneous sniffing in the TuS or
NAcSh (Ciii, $p>0.05$). **D.** Impact of selective antagonism of D1- (Di), D2- (Dii), and D3-receptors
(Diii) on odor-evoked sniffing. D1 and D3 antagonism had no effect on odor-evoked sniffing in
the TuS or NAc (Di & Diii, $p>0.05$). **E.** Impact of selective antagonism of D1- (Ei), D2- (Eii), and
D3-receptors (Eiii) on buzz-evoked sniffing. D2 and D3 antagonism had no effect on buzz-evoked
sniffing in the TuS or NAc (Eii & Eiii, $p>0.05$). Data are mean (dark bolded line) ± SEM, lighter

lines = individual mice. Abbreviations: SCH (D1 antagonist SCH23390), RAC (D2 antagonist
raclopride), PG0 (D3 antagonist PG010370), VEH (vehicle). ****$p<0.0001$, **$p<0.01$, *$p<0.05$.
1650

Johnson et al., Figure 8

Johnson et al, Supp Figure 1

n=13 mice / region, 9 mins of "baseline" respiration / mouse

Johnson et al, Supp Figure 5

Johnson et al., Supp Figure 7