Title: Time-restricted feeding leads to sex- and organ-specific responses in the murine digestive system

Short title: Food entrainment in the gut and liver

Lalita Oparija-Rogenmozere[1,2,¶,#a]*, Madeleine R. Di Natale[1,2,¶], Billie Hunne[1], Ada Koo[1], Mitchell Ringuet[1], Therese E. Fazio Coles[1], Linda J. Fothergill[1], Rachel McQuade[3,4,&], John B. Furness[1,2,&]

[1]Department of Anatomy and Physiology, The University of Melbourne, Parkville, Victoria 3052, Australia

[2]The Florey Institute of Neuroscience and Mental Health, Parkville, Victoria 3052, Australia

[3]Gut Barrier and Disease Laboratory, Department of Anatomy & Physiology, The University of Melbourne, Parkville, VIC 3010, Australia

[4]Australian Institute for Musculoskeletal Science (AIMSS), Western Centre for Health Research and Education (WCHRE), Sunshine Hospital, St Albans, VIC 3021, Australia

[#a]Current Address: Department of Biomedicine, University of Basel, 4058 Basel, Switzerland

*Corresponding author. Mailing address: Dr. Lalita Oparija-Rogenmozere, Department of Biomedicine, University of Basel, 4058 Basel, Switzerland. E-mail: lalita.oparija-rogenmozere@unibas.ch

[¶]These authors contributed equally to this work.

[&]These authors also contributed equally to this work.

Abstract

Food intake is one of the main zeitgebers in the digestive system; however, little is known about organ- and sex-specific differences in food-driven regulation. We placed male and female C57Bl/6 mice on time-restricted feeding (TRF), limiting the food intake period to 8 hours. Food was added either at dark (ZT12) or light (ZT0) onset for 14 days. Afterwards, an additional 4-hour delay in the feeding period was introduced for half of the mice, and the TRF regime continued for another 14 days.

TRF from ZT12 to ZT20 led to the highest weight gain in females but the lowest in males while improving intestinal transepithelial resistance (TEER) in both sexes. However, it also led to the disappearance of food-anticipatory response in several hepatic genes. Delaying the start of TRF until ZT16 led to an increase in weight gain and a decrease in fasting plasma glucose levels in male mice, as well as to strong entrainment of metabolism-related hepatic and duodenal genes in both sexes.

The alignment of food intake with the early lights-on phase (ZT0-ZT8) caused only minor changes in physiological responses. However, it did lead to an overall downregulation of hepatic and an upregulation of duodenal and gastric genes, with additional loss of food-anticipatory gene expression in both sexes. Delaying the start of food intake until ZT4 was highly detrimental, causing an increase in fasting blood glucose levels, a decrease in TEER, and further disruptions in gene expression patterns in the stomach and liver. In contrast, the duodenum was able to restore its food-driven gene expression.

These results demonstrate that the adjustment to food intake time in mice is highly sex- and organ-specific. Our chosen TRF regimes were not able to

50 synchronize food-anticipatory responses between the liver and gut. Instead, we

51 observed that organs entrain to food intake at different rates.

Keywords

53 Food entrainment, time-restricted feeding, digestive system, gut, liver, sex-

54 specific entrainment, food intake time, peripheral zeitgebers, nutrient

55 metabolism

Introduction

The ability to measure time has played such an importance in our species that clocks are one of the oldest human inventions. We even have devoted an entire subset of science, called Chronometry, to studying timekeeping and developing new methods and standards. And yet, our oldest and arguably most important clock remains understudied.

Self-sustaining and cell-autonomous circadian clocks are essential parts of our biology [1], with early studies revealing diurnal oscillations in the body temperature [2-4], food intake [5], and specific organ functions, for example, renal excretion of water and salts [6, 7]. By now, it has been shown that the circadian clock drives rhythmic expression of approximately 43% of our protein-coding genes and more than a thousand noncoding RNAs [8], regulating major physiological functions across most tissues and organs (for a recent review on the molecular clock and its regulatory role see: [9]).

Simplistically, the mammalian molecular circadian clock is a transcriptional feedback loop, expressed in almost every cell and driving expression patterns of multiple clock-controlled genes with approximately 24-hour rhythmicity. The positive limb of this feedback loop consists of two master transcription factors: CLOCK (circadian locomotor output cycles kaput) and BMAL1 (brain and muscle ARNT-like protein), which form a heterodimer. This heterodimer then binds to the promoters of its target genes, including those comprising the negative limb of the feedback loop: *Per1*, *Per2*, *Per3* (further: PERs – mammalian analogs of the *Drosophila period* gene) and *Cry1*, *Cry2* (further: CRYs). PERs and CRYs dimerize, translocate back to the nucleus, and inhibit the activity of CLOCK-BMAL1 heterodimer, subsequently inhibiting their expression (for extensive reviews on core clock function and mechanism, see

[1, 10-14]). There are also multiple secondary loops modulating the output of

the core clock, for example, those involving orphan nuclear receptors REV-

ERBs and/or retinoic acid receptor-related orphan receptors (RORs). Both of

these receptor families have been suggested to provide a link between nutrient

metabolism and the circadian clock, especially regarding lipid homeostasis [15,

16]. In addition to the translational regulation, the circadian clock is also

regulated post-translationally, with multiple kinases, phosphatases, microRNAs,

and RNA-binding protein complexes modulating various aspects and the output

of the core clock [1].

Such modulation mechanisms allow for an alignment of circadian rhythms to

various cues. In humans, the rhythm can be influenced by age, race,

geographical location, and sex [17-20]. In addition, our daily schedules cause

recurring variations in environmental and nutritional stimuli, leading to further

rhythm adaptation and synchronization. Such stimuli, affecting the timing of any

biological rhythm within an organism, are called "zeitgebers" ("time-givers"), a

term coined in 1945 [21].

The daily light-dark cycle is the best-researched zeitgeber, entraining the

central circadian oscillator that is located in the suprachiasmatic nucleus (SCN)

of the hypothalamus. SCN then conveys the generated rhythms to the periphery

– directly via neural and hormonal signaling and indirectly via regulation of body

temperature. Concurrently, peripheral tissues and organs, for example, the

gastrointestinal (further: GI) tract and liver, also possess self-sustaining

circadian rhythms with organ-specific phase shifts [22]. These peripheral clocks

can entrain to such zeitgebers as food (content and intake time), exercise, and

drugs of abuse, potentially leading to their uncoupling from the central clock [23-

30]. Circadian misalignment arising from conflicting zeitgeber signals

contributes to increased risks of cardiovascular, metabolic, immune,

neurological, and even psychiatric disorders (for reviews on health impact, see

[31, 32]). In contrast, synchronization between central and peripheral rhythms,

for example, by aligning food intake to the light-dark cycle, can be beneficial for

general health and well-being (for reviews, see [33, 34]).

Unfortunately, the research on food entrainment of peripheral rhythms is still

fragmented. In the liver, food entrainment has been shown to modulate gene

expression and metabolic state to account for the anticipated food volume and

intake time, even if it requires uncoupling from the SCN-driven rhythm and

exacerbating inflammatory responses [23, 35-40]. Ironically, organs comprising

the GI tract, which is essential for food uptake, breakdown, and absorption,

have been somewhat neglected in food-driven regulation research. Still, several

studies have shown that food entrainment regulates GI hormone and short-

chain fatty acid secretion [41-43], gut microbiota, lipidome and inflammation [43-

45], gastric contractility [43], and expression of clock and clock-controlled genes

in small and large intestine [25, 46-49] (for a recent review, see [50]).

Synchronization of rhythms across several peripheral organs or tissues in

response to food intake time is also poorly studied, with existing reports

suggesting that entrainment rates might be organ-specific [23, 46, 51].

Yet another understudied area is sex-specificity in daily rhythms, mostly due to

an underrepresentation of female animal models or an overrepresentation of

one sex in human studies. There is, however, a physiological basis to expect

sex-specific responses, as SCN morphology and neuropeptide expression in

rodents differ between males and females [52, 53]. Both rodent and human

studies have also shown sex-specific differences in corticosteroid secretion,

activity levels, circadian period length, and the stability of the rhythms (for

reviews, see [54-56]). When it comes to interaction between sex and food entrainment, male mice seem to develop an earlier and higher-amplitude food-anticipatory behavior when compared to females [57]. In addition, differences in weight gain, clock gene expression, and ghrelin secretion have also been observed [58]. Still, sex-specific differences in food entrainment remain enigmatic.

In this study, we aimed to investigate food-driven, sex-specific regulation of weight gain, glucose handling, intestinal barrier function, and genes involved in nutrient metabolism in the murine digestive system. Adult male and female mice were placed on an 8 h time-restricted feeding (TRF) regime starting either at the onset of the dark (ZT12) or light (ZT0) phase, with or without an additional 4 h feeding delay after 14 days of entrainment. We observed that food intake time not only regulates weight gain, fasting glucose values, and intestinal transepithelial resistance (further: TEER) but also leads to organ- and sex-specific alterations of the core clock (*Clock, Bmal1*) and metabolism-related gene expression.

Materials and Methods

Ethical approval

All experimental procedures and handling involving mice were approved by the University of Melbourne Animal Ethics Committee (application #1914983) and performed in compliance with the Australian code for the care and use of animals for scientific purposes.

Experimental animal origin and housing

In total, 180 male and 180 female C57Bl/6 mice (RRID:IMSR_JAX:000664) from the Animal Resources Centre (Canning Vale, WA, Australia) were used in

the study. All mice arrived at the local animal facility aged 5-7 weeks and were

kept under a 12:12 h light-dark (further: LD) cycle, with standard chow

(Barastoc 102108, Ridley Corporation, Melbourne, Australia) and water

provided *ad libitum*. After three days of post-arrival adjustment to the local

facility, mice were transferred to either LEDDY cages (black cages with

individual lighting systems) (Cat# GM500) or to the Aria ventilated cabinet (Cat#

9BIOC44R4Y1, both from Tecniplast, Buguggiate, Italy), and the light onset of

12:12 h LD cycle was adjusted such that the sample collection from day-fed and

night-fed mice could be done in parallel. Mice were adjusted to the new 12:12 h

LD cycle and housing conditions for 1 week, based on observations that young

mice re-entrain within 5-10 days, especially regarding peripheral rhythms [59]

and their activity [60]. Chow and water were provided *ad libitum* during this time.

After this week, mice were weighed at the end of the dark phase (Zeitgeber time

(ZT) 24). Obtained values were recorded as pre-time-restricted feeding weights

(further: pre-TRF) and used as a baseline for later weight gain calculations.

Food entrainment of mice

After a 1-week adjustment to the new LD cycle, mice were subjected to TRF for

14 days (Fig 1, Pre-shift days). During this time, mice were fed standard chow

for 8 h starting either from the dark onset at ZT12 (Fig 1, Restricted night-fed,

grey bar) or the light onset at ZT0 (Fig 1, Restricted day-fed, yellow bar),

followed by a 16 h fasting period. To ensure that mice did not feed on chow

debris during fasting, TRF was done by using two cages: a chow-containing

cage ("food cage") and a cage that never contained any chow ("empty cage").

Mice were placed in the "food cage" during the feeding period and in the "empty

cage" during the fasting period. Water was always provided *ad libitum* for the

whole duration of the experiment. Mice were weighed weekly at the end of their

respective feeding period (Fig 1, blue arrows). Food intake was not measured,

as mice were group-housed.

Fig 1. Food entrainment schedule. Bars indicate the placement and length of

the feeding period. Arrows show weekly weight measurement timepoints. Mice

were sacrificed on the 3rd, 7^{th,} or 14th day after feeding shift. Letters indicate

sacrifice timepoints on each day: A = food anticipation, F = food intake, P =

postprandial. n = 5 per sacrifice timepoint per sex.

After 14 days of the initial entrainment, the 8-hour feeding period was delayed

by 4 h for half of the mice (Fig 1, Day 15). The new feeding groups were labeled

as 'Restricted shifted night-fed' (Fig 1, ZT16-ZT0, dark grey bar) and 'Restricted

shifted day-fed' (Fig 1, ZT4-ZT12, orange bar). The other half of the mice

continued with their initial feeding regimen. Weekly weight gain measurements

were continued as previously stated (Fig 1, blue arrows). On the 3rd, 7^{th,} and

14th day after the shift, mice were sacrificed for organ collection. Three

timepoints were chosen to assess food-driven regulation on each sacrifice day:

1) food anticipation: after 16 h of fasting, right before the anticipated transfer to

the "food cage" (Fig 1, "A"); 2) food intake: 4 h after the mice were placed in the

"food cage" (Fig 1, "F"); 3) postprandial period: 4 h after the mice were placed in

the "empty cage" (Fig 1, "P"). On each sacrifice day, n=5 female and n=5 male

mice were sacrificed at each timepoint.

Mice were anesthetized with isoflurane (Cat# FGISO0250, Pharmachem, Eagle

Farm, Australia), followed by a cervical dislocation. Upon sacrifice, their body

weight was measured, and the following organs and tissues were collected and

snap-frozen using liquid nitrogen: the right lobe of the liver, the dorsal half of the

stomach, and the duodenal mucosa. To collect duodenal mucosa, the first 2 cm

of the duodenum were opened and scraped with a surgical scalpel at a 45-

degree angle. The ileum (5 cm proximal from the cecum) was collected only

during the food intake timepoint (Fig 1, "F") and used to measure intestinal

permeability and TEER.

Blood glucose measurements

217 Blood glucose was measured at the food anticipation timepoint after 16 h

218 fasting (Fig 1, "A"). A drop of blood from the tail vein was applied on the Accu-

219 Chek® Performa Test strip (Cat# 06454038020), and glucose concentration

220 was measured with Accu-Chek® Performa (Cat# 05894964014) blood glucose

221 meter (both from Roche Diagnostics, Manheim, Germany).

Intestinal electrical resistance measurements

The ileum collected during the food intake timepoint (Fig 1, "F") was cut in half.

Both pieces were placed in Krebs-Henseleit buffer (11.1 mM glucose, 118 mM

NaCl, 4.8 mM KCl, 1.0 mM NaH_2PO_4, 1.2 mM $MgSO_4$, 25 mM $NaHCO_3$, 2.5 mM

$CaCl_2$, pH 7.4), opened along the mesenteric border and pinned (full-thickness)

onto Ussing chamber sliders (Cat# P2311, 0.3 cm^2 apertures, Physiological

Instruments, San Diego, USA). Sliders were inserted in the middle of two-part

Ussing chambers (EasyMount Diffusion Chambers, Physiologic Instruments,

San Diego, USA), and 5 ml Krebs-Henseleit buffer solution was added to the

serosal side of the tissue. The mucosal side of the tissue received 5 ml of

modified Krebs-Henseleit buffer, where glucose was substituted with 11.1 mM

mannitol. This was done to avoid apical uptake of glucose while still maintaining

an osmotic balance. Buffers in both chambers were kept at 37^{o}C and bubbled

with carbogen (5% CO_2, 95% O_2) to maintain an optimal pH level. A

multichannel voltage-current clamp (VCC MC6, Physiologic Instruments, San

Diego, USA) was applied to each chamber through a set of four electrodes (2

voltage sensing and 2 current passing electrodes) and agar bridges (3%

agarose/3 M KCl in the tip and backfilled with 3 M KCl), installed on opposite

sides of the tissue. The tissue was left to equilibrate for 20 min before clamping

the voltage to 0 V.

To calculate TEER ($\Omega \cdot cm^2$), 2-sec pulses of 2 mV were administered to tissue

every 60 sec for 1 h, and the measured net resistance was multiplied by the

surface area. Voltage and short circuit current (Isc) measurements were

recorded using a PowerLab amplifier and the software LabChart® 5

(RRID:SCR_017551, both ADInstruments, Sydney, Australia).

RNA extraction

Total RNA from approx. 20 mg of the right lobe of the liver, dorsal half of the

stomach, and scraped duodenal mucosa was extracted using ISOLATE II RNA

Mini Kit (Cat# BIO-52073, Meridian Bioscience, Cincinnati, USA) with the

following adjustments to the manufacturer's instructions. First, tissue pieces

were immediately snap-frozen in 2 ml screw-cap tubes containing Lysing Matrix

D (Cat# 6540-434, Lot# 99999, MP Biomedicals, Irvine, USA) and stored at -

80^{o}C until further processing. Second, the volume of the provided lysis buffer

was increased to 500 µl per sample, and 5 µl of β-mercaptoethanol was added.

Lysis buffer was applied while the samples were still frozen and immediately

followed by lysis at 6000 rpm for 2 x 30 sec in a Precellys 24 homogenizer

(RRID:SCR_022979, Cat# 03119-200-RD010, Bertin Technologies SAS,

Montigny-le-Bretonneux, France). The volume of RNAse-free Ethanol (Cat#

EA043, Chem-supply, Gillman, Australia), used for RNA binding, was also

increased to 500 µl per sample, and lysate-ethanol filtration through the

provided column was done in two consecutive steps. The centrifugation time for

all washing steps was increased to 1 min to ensure better removal of wash

buffers. Samples were eluted in 100 µl (liver) or 60 µl (stomach, duodenal

mucosa) of RNase-free water (Cat# 10977-015, Lot# 2186758, Invitrogen by

Life Technologies, Grand Island, USA). The elution step was repeated using the

initial eluate to increase the RNA yield. The quality and quantity of the extracted

RNA were assessed using a 2200 Tape Station (RRID:SCR_014994, Agilent

Technologies, Santa Clara, USA) and a Nanodrop ND-1000 UV

spectrophotometer (RRID:SCR_016517, NanoDrop Technologies, Wilmington,

USA), respectively. The isolation was randomized and done in small batches

(24 samples) to ensure fast processing and good RNA quality.

The synthesis of cDNA was performed using 100 ng RNA in a 20 µl total

reaction volume using an iScript Reverse Transcription Supermix kit (Cat#

1708841, Bio-Rad, Hercules, USA) without any changes from the

manufacturer's instructions. Samples were split across 96-well plates, with 6

negative controls included on each plate, these were randomly chosen RNA

samples where no cDNA synthesizing enzyme was added. Synthesis reaction

was carried out using PCRExpress (Cat# 630-003, Thermo Hybaid, Franklin,

USA) or Bio-Rad T100™ (RRID:SCR_021921, Bio-Rad, Hercules, USA)

thermal cyclers.

282 Determination of gene expression

After cDNA synthesis, 10 µl of each cDNA sample was delivered to the

Translational Research Facility within Monash Health Translational Precinct to

determine gene expression levels using the Fluidigm Digital Array Integrated

Fluidic Circuits [61] (further: IFCs). First, all cDNA samples underwent quality

control, where expression of either *Gapdh* (forward primer:

TGACCTCAACTACATGGTCTACA, reverse: CTTCCCATTCTCGGCCTTG) or

β-actin (forward primer: GGCTGTATTCCCCTCCATCG, reverse:

CCAGTTGGTAACAATGCCATGT) was tested as SYBR assays using

QuantStudio 6 Flex RealTime PCR reader (RRID:SCR_020239, Thermo Fisher

Scientific, Waltham, USA). Afterwards, samples underwent pre-amplification

according to the manufacturer's instructions [62]. In short, all 24 TaqMan

assays for genes of interest (Table 1) were first pooled and diluted in Tris-EDTA

buffer (pH 8.0) to a final concentration of 180 nM per assay. Then 3.75 µl of

gene assay mix was added to 1.25 µl of cDNA sample and pre-amplified for 14

cycles using Veriti™ 96-well Thermal Cycler (RRID:SCR_021097, Cat# 9902,

Thermo Fisher Scientific, Waltham, USA). Pre-amplified cDNA samples were

further diluted 1:5 with Tris-EDTA buffer (pH 8.0) and loaded on the 192.24

Dynamic array IFC (Cat#100-6266, multiple lots used, Fluidigm, San Francisco,

USA) together with gene assays, following Fluidigm® 192.24 Real-Time PCR

Workflow Quick Reference PN 100-6170 [62]. The qPCR reaction was

performed using the Biomark™ HD system (RRID:SCR_022658, Cat# BMKHD,

Fluidigm, San Francisco, USA).

Table 1. TaqMan Gene assays used to determine relative gene expression levels

Assigned group by function	Gene	Gene accession number	Assay ID	Tested in samples from:
Reference gene	*18s (Rn18s)*	NR_003278.3	Mm04277571_s1	Liver, stomach, duodenum
Core clock genes	*Clock*	NM_001289826.1	Mm00455950_m1	
	Bmal1 (Arntl)	NM_001243048.1	Mm00500226_m1	
Glucose sensing, transport and metabolism	*Tas1r2*	NM_031873.1	Mm00499716_m1	
	Sglt1 (Slc5a1)	NM_019810.4	Mm00451203_m1	
	Glut2 (Slc2a2)	NM_031197.2	Mm00446229_m1	
	Pepck (Pck1)	NM_011044.2	Mm01247058_m1	
	Gys2	NM_145572.2	Mm01267381_g1	Liver

Amino acid and peptide transport	Pept1 (Slc15a1)	NM_053079.2	Mm04209483_m1	Liver, stomach, duodenum
	B⁰at1 (Slc6a19)	NM_028878.3	Mm01352157_m1	
Lipid metabolism	Mttp	NM_001163457.1	Mm00435015_m1	
	Fasn	NM_007988.3	Mm00662319_m1	
	Cyp7a1	NM_007824.2	Mm00484152_m1	
	Pparα	NM_001113418.1	Mm00440939_m1	
Biosynthesis of NAD+	Srebf1	NM_011480.3	Mm00550338_m1	Liver
	Nampt	NM_021524.2	Mm00451938_m1	
Hormone signalling	Ffg21	NM_020013.4	Mm00840165_g1	
	Igf-1	NM_001111274.1	Mm00439560_m1	
	Tph1	NM_001136084.2	Mm01202614_m1	Stomach and duodenum
	Sst	NM_009215.1	Mm00436671_m1	
	Ghrl	NM_001286404.1	Mm00612524_m1	
	Gast	NM_010257.3	Mm00439059_g1	Stomach
	Lepr	NM_001122899.1	Mm00440181_m1	
	Pga5	NM_021453.4	Mm01208256_m1	
	Gip	NM_008119.2	Mm00433601_m1	Duodenum
	Gcg	AF276754.1	Mm00801714_m1	
	Cck	NM_031161.4	Mm00446170_m1	
Pro-/Anti-inflammatory response	TNFα	NM_013693.3	Mm99999068_m1	Liver, stomach, duodenum
	Il-6	NM_031168.1	Mm00446190_m1	
	Il-17a	NM_010552.3	Mm00439619_m1	
	Crp	NM_007768.4	Mm00432680_g1	Liver
	Gpx1	NM_008160.6	Mm00656767_g1	
	Gpx2	NM_030677.2	Mm00850074_g1	Stomach and duodenum

307   *18s* was used as a reference gene for the calculation of relative expression

308   ratios (R), using the following formula: $R = 2^{-(Ct(test\ RNA)-Ct(18S\ RNA))}$. All relative

309   expression ratios were further normalized against the average value shown by

310   night-fed males at the food anticipation timepoint (Fig 1, point "A") on the 3rd day

311   after the feeding shift.

Data analysis and statistics

Experimental data analysis was performed using GraphPad Prism v8.2 (RRID:SCR_002798, GraphPad Software, San Diego, CA, USA). To demonstrate weight gain, data from each mouse within the same feeding group and sex (n=90 for pre-TRF, week 1 and week 2; n = 30 for week 3, n = 15 for week 4) were pooled and shown as a mean group value and 95% Confidence Interval, CI. Significant differences in weekly weight gain between the night-fed group and other feeding groups were assessed by Two-way analysis of variance (ANOVA) with Šidák's multiple comparison test.

Measured TEER values were submitted to an outlier test, with all values outside of mean +/- 2*(standard deviation interval) excluded from further analysis. This was done to exclude tissue pieces that might have been damaged during the resistance measurements.

Weight at the time of sacrifice (Fig 2B), fasting blood glucose values, and TEER values were shown as separate data points per individual mouse, together with the mean group value and standard deviation (Further: SD). Two-way ANOVA was used together with Dunnett's multiple comparison test to determine differences between all experimental groups independently of the sacrifice day. However, only significant differences between the night-fed group and other TRF regimes on each sacrifice day and the differences between values recorded on various sacrifice days within the same feeding group are shown in the figure.

To show gene expression differences between male and female mice, normalized expression ratios of mice representing the same feeding group, sacrifice timepoint, and sex were pooled together and shown as a tile within a heatmap. Fold changes in gene expression were shown as Log2 values. To

further demonstrate the differences in gene expression between the food

anticipation, intake, and postprandial period (=timepoint-specific differences),

the nonparametric Kruskal-Wallis test was used together with Dunn's multiple

comparison test (food anticipation timepoint vs food intake or vs postprandial

period). The p-values were shown as: $*p \le 0.05$, $**p \le 0.01$, $***p \le 0.001$, $***p \le$

0.0001. Differences in gene expression introduced by the 4 h feeding shift were

assessed with the same statistical test, comparing values between shifted and

non-shifted groups at the same timepoint (e.g., night-fed postprandial vs shifted

night-fed postprandial). The p-values were shown as: $^{£}p \le 0.05$, $^{££}p \le 0.01$, $^{£££}p$

≤ 0.001, $^{££££}p \le 0.0001$.

Results

Food-driven regulation of weight gain, fasting plasma glucose levels, and intestinal permeability

TRF-induced changes in weight gain are sex-specific

First, all mice had their baseline (pre-TRF) weights recorded. One day later, 180 male and 180 female mice started time restricted-feeding (further: TRF) as either day-fed (further: DF) or night-fed (further: NF) and remained on this regime for 2 weeks. 45 females and 45 males from each TRF regime then underwent an additional 4 h delay shift in the mealtime, creating shifted night-fed (further: SNF) and shifted day-fed (further: SDF) groups. The remaining mice stayed on their initial TRF regime (Fig 1). The total entrainment length was 4 weeks. Weight gain was assessed weekly at the end of the respective feeding period (Fig 1, blue arrows). Afterwards, experimental weights were related to their pre-TRF values and expressed as a change in percentage (%) over time. Weight gain in the NF group was used as a control value in all statistical tests, allowing to determine the impact of lights-on feeding and delay shifts; moreover, this regimen is also the closest to the natural food intake pattern in mice [63].

We observed TRF regime- and sex-specific differences in weight gain. At the end of the 2^{nd} entrainment week, DF males showed a significant 2.9% increase in weight when compared to their NF counterparts. However, this difference disappeared after longer entrainment (Fig 2A, bottom panel, yellow vs grey). In contrast, DF female mice showed significantly delayed weight gain (-4.4%) on week 3 but then also caught up to the NF group (Fig 2A, top panel, yellow vs grey) after longer entrainment.

Fig 2. Physiological effects of time-restricted feeding. A: Post-feeding weights were normalized to individual pre-food entrainment (pre-TRF) values. Mean (95% CI) shown, n = 90 mice per sex at pre-TRF, week 1 and week 2; week 3 n = 30 mice per sex; week 4 n = 15 mice per sex. Two-way ANOVA with Šidák's multiple comparison test, showing differences between the night-fed group and other TRF regimes on each sacrifice week, $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$. **B**: Differences in recorded sacrifice weights. Values normalized to individual pre-TRF values. **C**: Fasting glucose values recorded at food anticipation timepoint. **D**: Transepithelial electrical resistance (TEER) in the ileum, recorded at food intake timepoint. **B to D:** Mean (SD) shown, n = 5 mice per sex per timepoint. Two-way ANOVA with Dunnett's multiple comparison test, showing differences between the night-fed group and other TRF regimes on each sacrifice day and differences between various sacrifice days (Day 3 as a control) within each feeding group, $*p<0.05$, $**p<0.01$, $***p<0.001$, $****p<0.0001$.

The additional delay of mealtime caused sex-specific effects as well. In females, we observed similar weight gain between shifted and non-shifted counterparts, but male mice seemed to be more responsive, demonstrating a significantly increased weight gain in both shifted groups one week after the mealtime delay (Fig 2A, in orange vs grey). Overall, it appears that food intake in the late lights-on phase (SNF) was the only condition leading to a similar weight gain pattern in both sexes.

Feeding during the lights-on phase leads to weight instability in female mice

The bodyweight was also recorded at each sacrifice timepoint: food anticipation (after 16 h fast, when mice expect to be fed, Fig 1, "A"), food intake (4 h after food was added, Fig 1, "F") and postprandial period (4 h after food was removed, Fig 1, "P") on all sacrifice days. The earliest sacrifice was performed on Day 3 after the delayed feeding shift (Day 3), followed by sacrifices on Day 7 and Day 14.

We then compared the sacrifice weights to pre-TRF values (Fig 2B, blue dashed line) and observed that DF and SDF mice showed a more severe weight loss at the food anticipation timepoint, with some even reaching or dipping below the pre-TRF weights. This effect was less pronounced in male mice, where it disappeared under longer entrainment, while females did not show any stabilization. On the contrary – by the end of the experiment, DF and SDF females showed significantly lower food anticipation weights when compared to their NF counterparts (Fig 2B, top panel, Food anticipation).

Weights at the food intake timepoint (Fig 2B, Food intake) tended to increase over time, especially in male mice. There were no differences in weights between various TRF regimes; individual food intake was not measured.

Weights during the postprandial period (Fig 2B, Postprandial) did not differ between TRF regimes, suggesting similar rates of weight loss after 4h fasting.

TRF leads to a sex-specific regulation of fasting blood glucose levels

Fasting blood glucose was measured at the food anticipation timepoint (Fig 1, "A"). We observed that female DF mice showed significantly lower fasting blood glucose levels on Day 3 (=18 days of entrainment for this TRF regime) when

compared to their NF counterparts (Fig 2C, grey vs yellow). Concurrently, SNF led to significantly lower glucose levels in male mice (Fig 2C, grey vs black). Both of these differences disappeared under longer entrainment and were not present in the opposite sex, suggesting that male and female glucose homeostasis entrains to SNF and DF at different rates. Interestingly, delaying the food intake until the late lights-on phase quickly restored the blood glucose levels in female mice to NF-like-values (Fig 2C, grey vs orange vs yellow), while males showed no difference between DF and SDF.

Prolonged food intake during the lights-on phase negatively impacts intestinal barrier function

Ileum from mice sacrificed during the food intake timepoint (Fig 1, "F") was used to determine TEER. All feeding groups, independently of sex, showed similar TEER values on sacrifice Day 3 (Fig 2D, Day 3). However, we did observe changes under prolonged TRF, especially in male mice. NF males showed an increased TEER over time, while DF and SDF led to a significant decrease by Day 14 (Fig 2D). Female mice showed higher variability, with DF causing significantly lower TEER only on Day 7 but SDF – on Day 14 (Fig 2D). Overall, it appeared that prolonged SDF is detrimental for ileal TEER in both sexes, but NF allows to maintain or even improve the barrier function.

TRF-driven gene expression in the digestive system

Expression of genes involved in the molecular clock, nutrient metabolism, and immune response was determined in the liver, stomach, and duodenal mucosa. Tissue samples were collected from all experimental animals on sacrifice days 3, 7, and 14 (post-shift), at food anticipation, food intake, and postprandial timepoints (Fig 1, "A", "F" and "P", respectively). Gene expression was first

normalized against housekeeping gene *18s* and then to the expression level in

NF male mice at food anticipation timepoint on sacrifice Day 3. Afterwards, the

data from each experimental group (same sex, sacrifice day, and timepoint)

was pooled and shown as a single tile within the figure heatmaps. Significant

changes in expression between food anticipation and one of the other

timepoints within the same TRF group were classified as "timepoint-specific"

and shown as *p. Concurrently, significant gene expression differences between

shifted and non-shifted groups at the same timepoint were classified as "shift-

specific" and shown as $^{\$}$p.

Food-driven regulation of *Bmal1* and *Clock* is organ-specific

To determine whether the chosen TRF regimes were able to entrain peripheral

core clock genes, we looked at the expression of *Bmal1* and *Clock* in the liver,

stomach, and duodenum. We observed that only *Bmal1*, but not *Clock*, showed

significant differences between food anticipation and feeding and/or

postprandial timepoints in both sexes and all tested organs (Fig 3, *p).

Fig 3. Changes in *Bmal1* and *Clock* expression in male and female mice

under TRF. Sample collection timepoints shown at the top: A = food

anticipation, F = food intake, P = postprandial period. Feeding conditions shown

on the left: NF = night-fed (ZT12 – ZT20), SNF = shifted night-fed (ZT16 –

ZT24), DF = day-fed (ZT0 – ZT8), SDF = shifted day-fed (ZT4 – ZT12). Relative

gene expression values (Log2) were normalized to NF male mice on sacrifice

day 3 at the food anticipation timepoint and pooled by sex, timepoint, and

sacrifice day. The mean value of each experimental group is shown as an

individual tile (n= 4-5 mice). Non-parametric Kruskal-Wallis test with Dunn's

multiple comparison test used to show timepoint-specific (food anticipation vs

intake or vs postprandial period) differences in gene expression on each

sacrifice day, *p<0.05, **p<0.01, ***p<0.001. The same test was used to assess

shift-specific differences (non-shifted vs shifted) at each timepoint, $^{\$}$p<0.05,

$^{\$\$}$p<0.01.

It has been shown that in *ad libitum* fed mice under a standard 12:12h light

cycle, *Bmal1* expression in the liver reaches its peak in the late dark phase

(between ZT20 and ZT0) and the nadir right before the dark onset (between

ZT8 and ZT12) [64]. We observed the peak expression of the hepatic *Bmal1* at

the food intake or postprandial timepoint under all tested TRF regimes,

indicating a response to the food entrainment and possible uncoupling from the

light-driven rhythm under DF and SDF conditions. Some timepoint-specific

responses were also present for the hepatic *Clock* gene; however, they were

not as persistent.

In contrast to the liver, *Bmal1* expression in the stomach and duodenum

responded poorly to daytime feeding, with similar expression levels at all

timepoints or with the disappearance of changes over time, respectively (Fig 3).

However, in NF and SNF male and female mice, we still observed gastric and

duodenal *Bmal1* upregulation at the postprandial timepoint. The *Clock* gene did

not exhibit recurrent, timepoint-specific changes in expression levels, paralleling

previous observations that this gene is not cyclic in the stomach [65].

The 4 h feeding delay had little effect on *Bmal1* or *Clock* expression at tested

timepoints, as significant differences between shifted and non-shifted groups

appeared rarely and somewhat randomly (Fig 3, differences shown as $^{\$}$p). A

more interesting consequence of TRF was the overall lower expression of both

clock genes in the liver under DF and SDF regimes (compared to their NF

counterparts), while in the stomach and duodenum, we observed the opposite

(Fig 3).

Sex-specific differences in gene expression patterns were also present. In some

cases, for example, for hepatic *Bmal1* under NF, hepatic *Clock,* and gastric and

duodenal *Bmal1* under NF and SNF, it appeared that in female mice, timepoint-

specific changes were more likely to be present early (on Day 3 post-shift) and

then weaken/disappear over time, while in males the response appeared later,

but persisted over longer entrainment.

Food intake during the lights-on phase downregulates hepatic gene expression

Similar to our observations with clock genes, also hepatic nutrient metabolism-

and inflammatory response-related genes showed an overall lower expression

in DF and SDF male and female mice (Fig 4). In addition, even though

timepoint-specific expression differences were present for multiple genes, not

all of them aligned their expression pattern to every TRF regime tested, leading

to an overall poor response to food entrainment.

Fig 4. Hepatic gene expression changes in male and female mice under

TRF. Sample collection timepoints shown at the top: A = food anticipation, F =

food intake, P = postprandial period. Feeding conditions shown on the left: NF =

night-fed (ZT12 – ZT20), SNF = shifted night-fed (ZT16 – ZT24), DF = day-fed

(ZT0 – ZT8), SDF = shifted day-fed (ZT4 – ZT12). Relative gene expression

values (Log2) were normalized to NF male mice on sacrifice day 3 at the food

anticipation timepoint and pooled by sex, timepoint, and sacrifice day. The

mean value of each experimental group is shown as an individual tile (n= 4-5

mice). Non-parametric Kruskal-Wallis test with Dunn's multiple comparison test

used to show timepoint-specific (food anticipation vs intake or vs postprandial

period) differences in gene expression on each sacrifice day, *p<0.05, **p<0.01,

***p<0.001. The same test was used to assess shift-specific differences (non-

shifted vs shifted) at each timepoint, $p<0.05, $$p<0.01.

Surprisingly, *Pparα* and *Srebf1*, the principal transcription factors regulating

hepatic lipid metabolism [66] that have been observed to be responsive TRF

[39, 67], showed discontinuous timepoint-specific changes in gene expression

(Fig 4, *p) with only SNF females (*Pparα*) exhibiting downregulation at

postprandial period on more than one sacrifice day. The 4h food intake delay

also led to only minor changes, mostly by upregulating gene expression in SNF

male mice over prolonged entrainment, when compared to their non-shifted

counterparts (Fig 4, $p).

However, not every gene associated with nutrient metabolism was so

unresponsive to TRF. Genes encoding fatty acid synthase (*Fasn*), glycogen

synthase (*Gys2*), and nicotinamide adenine dinucleotide (NAD^+) biosynthetic

enzyme (*Nampt*) showed TRF-induced expression changes, similar to

observations before [39]. In addition, responses were also seen from genes

encoding phosphoenolpyruvate carboxykinase (*Pepck*) and fibroblast growth

factor 21 (*Fgf21*) (Fig 4, *p). Although we observed some sex- and timepoint-

specific gene expression differences under DF and SDF regimes for all these

genes, the most stable and continuous response in both sexes was seen under

SNF. An exception here was *Fgf21*, showing strong downregulation during food

intake and/or postprandial period under all TRF regimes, especially under

longer entrainment. Since *Fgf21* is involved in the regulation of nutrient and

energy homeostasis and has a fasting-induced secretion (for a review on its

role, see [68]), it might have been entrained to the daily 16 h fasting period in

our mice, resulting in food-anticipatory upregulation.

Not much is known about the food-driven regulation of hepatic (anti-

)inflammatory markers. Unfortunately, we were not able to clarify it either, as

$TNF\alpha$ showed an inconsistent upregulation at the food intake timepoint under

SNF and SDF, and no timepoint-specific pattern was present in $Gpx1$

expression.

Sex-specific differences seemed to follow a pattern observed for clock genes,

with females showing early timepoint-specific differences that tended to

disappear over time, but males developing late and/or more stable responses

(Fig 4., e.g., $Fasn, Pepck, Gys2, Nampt)$. Male mice also appeared to be more

likely to up- or down-regulate their gene expression in response to mealtime

shift, especially after 4 h food intake delay in the dark phase (SNF vs NF, $^{\$}$p,

see Fig 4: $Ppar\alpha, Srebf1, Fasn, Cyp7\alpha1, Mttp, Gys2)$.

TRF fails to elicit gene responses in the stomach

As a part of the digestive tract, the stomach is directly exposed to the ingested

food and has been shown to synchronize its hormone secretion to mealtime (for

a review, see [69]). And yet, gastric gene expression in our study was largely

unresponsive to TRF (Fig 5), a phenomenon also observed in rats [70]. Only

one major response was present in our mice – food intake during the lights-on

phase caused an overall gene upregulation (Fig 5, DF, SDF), following the

pattern seen for the gastric clock genes (Fig 3).

Fig 5. Gene expression changes in the stomach of male and female mice

under TRF. Sample collection timepoints shown at the top: A = food

anticipation, F = food intake, P = postprandial period. Feeding conditions shown

on the left: NF = night-fed (ZT12 – ZT20), SNF = shifted night-fed (ZT16 –

ZT24), DF = day-fed (ZT0 – ZT8), SDF = shifted day-fed (ZT4 – ZT12). Relative

gene expression values (Log2) were normalized to NF male mice on sacrifice

day 3 at the food anticipation timepoint and pooled by sex, timepoint, and

sacrifice day. The mean value of each experimental group is shown as an

individual tile (n= 4-5 mice). Non-parametric Kruskal-Wallis test with Dunn's

multiple comparison test used to show timepoint-specific (food anticipation vs

intake or vs postprandial period) differences in gene expression on each

sacrifice day, *$p<0.05$, **$p<0.01$, ***$p<0.001$. The same test was used to assess

shift-specific differences (non-shifted vs shifted) at each timepoint, $^{\$}p<0.05$,

$^{\$\$}p<0.01$, $^{\$\$\$}p<0.001$.

Despite the lack of stable and continuous TRF-induced changes in expression,

multiple genes appeared to follow similar response patterns. For example,

genes encoding major GI hormones or their precursors, such as *Ghrl, Pga5,*

and *Lepr* (encodes leptin receptor), as well as genes encoding (anti-

)inflammatory markers *Gxp2, Il-6* and *TNFα* and transcription factor *Pparα,* all

showed downregulation at the postprandial period in DF male mice on Day 14

and a considerable overlap in other timepoint-specific changes (Fig 5, *p). A

similar expression pattern was seen also for *Tph1,* which encodes tryptophan

hydroxylase – a rate-limiting protein in serotonin synthesis.

Gene encoding gastrin (*Gast*), a hormone stimulating gastric acid release and

gastric mucosal growth [71], showed a chaotic timepoint-specific gene

expression pattern as well, with changes appearing and disappearing between

sacrifice days and TRF regimes. Some studies have suggested that *Pparα*

could regulate *Gast* expression and/or translation [72, 73]; however, we

observed no similarities between their gene responses.

An unexpected observation was the strong timepoint-specific gene expression

pattern in gene *Pepck*, which encodes a rate-limiting enzyme for

gluconeogenesis. Downregulation during food intake was seen in SNF males

and females, as well as in NF females and DF and SDF males after prolonged

entrainment (Fig 5). Although gluconeogenesis can occur in the small intestine

[74], especially after gastric bypass surgery [75], currently, there is no evidence

of glucose production in the stomach, making us wonder why such gene

regulation would be necessary. However, *Pepck* has been proposed to play a

role also in the regulation of the TCA cycle flux [76] and nutrient processing

[77]; therefore, its timepoint-specific regulation in the stomach might be

important for gluconeogenesis-independent aspects of nutrient metabolism.

Gene expression changes induced by 4 h mealtime delay were uncommon in

the stomach, appearing only on select timepoints and sacrifice days (Fig 5, [$]p).

The lack of stable timepoint- and shift-specific expression patterns was present

in both sexes, obscuring any further sex-specific difference detection.

Duodenal genes involved in nutrient metabolism respond to

food entrainment

The duodenum is responsible for digestion and absorption of nutrients; thus, it

would not be surprising if food intake and nutrient-related signals would regulate

its gene expression. Indeed, duodenal genes involved in glucose and amino

acid uptake and/or metabolism were responsive to TRF (Fig 6). Some

similarities to our observations in the stomach were also present: daytime

feeding led to an overall gene upregulation, and the genes encoding GI

hormones or their precursors (*Cck, Gip, Ccg, Tph1*) showed poor responses to

TRF (Figs 5-6).

Fig 6. Duodenal gene expression changes in male and female mice under

TRF. Sample collection timepoints shown at the top: A = food anticipation, F =

food intake, P = postprandial period. Feeding conditions shown on the left: NF =

night-fed (ZT12 – ZT20), SNF = shifted night-fed (ZT16 – ZT24), DF = day-fed

(ZT0 – ZT8), SDF = shifted day-fed (ZT4 – ZT12). Relative gene expression

values (Log2) were normalized to NF male mice on sacrifice day 3 at the food

anticipation timepoint and pooled by sex, timepoint, and sacrifice day. The

mean value of each experimental group is shown as an individual tile (n= 4-5

mice). Non-parametric Kruskal-Wallis test with Dunn's multiple comparison test

used to show timepoint-specific (food anticipation vs intake or vs postprandial

period) differences in gene expression on each sacrifice day, $*p<0.05$, $**p<0.01$,

$***p<0.001$. The same test was used to assess shift-specific differences (non-

shifted vs shifted) at each timepoint, $^{\$}p<0.05$, $^{\$\$}p<0.01$.

Genes encoding large amino acid (B^0at1), peptide (*Pept1*) and glucose (*Sglt1*)

transporters, a sweet taste receptor (*Tas1r2*), an anti-inflammatory marker

(*Gxp2*), a gluconeogenesis regulator (*Pepck*) and a transcription factor

peroxisome proliferator-activated receptor alpha (*Pparα*) all had similar

timepoint-specific expression patterns (Fig 6, *p). Most of these genes were

downregulated during food intake and/or postprandial timepoint under NF and

SNF in both sexes. Interestingly, while DF did not always lead to timepoint-

specific differences (especially in females), SDF was able to induce very similar

expression patterns as NF, suggesting successful alignment to food-intake time.

It is also interesting that apart from *Bmal1*, *Pepck* was the only other gene

responding to TRF in all tested organs and both sexes.

Sex-specific differences were present as well. Females appeared to preserve or

even increase the timepoint-specific gene expression changes during prolonged

NF, while males showed stronger responses under prolonged DF and SDF.

Shift-specific differences seemed to be more prevalent in male mice (Fig 6, $^\$$p),

especially in the SNF group, peaking on Day 7 post-shift. Some genes (*B^0at1,*

Cck, Gip, Tph1, and *Glut2)* showed upregulation at food anticipation timepoint

in SDF females when compared to their non-shifted counterparts; however, this

difference was only present on the earliest sacrifice day.

Overall, we observed that different organs entrain to food intake time at various

rates. None of the used TRF regimes were able to synchronize timepoint-

specific differences across the liver, stomach, and duodenum. In addition,

responses were also sex-specific, with male mice appearing more sensitive to 4

h delay shift in the mealtime and female mice often exhibiting earlier but less

stable changes in gene expression.

Discussion

This study describes how food intake time affects weight gain, fasting glucose levels, intestinal permeability, and the expression of genes involved in the molecular clock, nutrient metabolism, and inflammatory processes in the digestive system of male and female mice. We utilized a TRF model to limit the food consumption to 8 h in a 24 h period, with the initial experimental groups starting their feeding period either at ZT12 (NF) or ZT0 (DF). After 14 days of entrainment, food intake time was delay shifted by 4 h for half of the mice (Fig 1), creating two new experimental groups: SNF (ZT16 start) and SDF (ZT4 start).

TRF impact on weight gain, glucose levels, and intestinal permeability

Observed changes in physiological parameters (relative to the NF group) during the full food entrainment period have been summarized in Table 2.

Feeding group	Weight gain		Fasting blood glucose		TEER	
	Females	*Males*	*Females*	*Males*	*Females*	*Males*
Night-fed (NF)	highest	lowest	No change over time		-	Increases over time
Shifted night-fed (SNF)	-	early ↑, then -	-	early ↓, then -	-	-
Day-fed (DF)	early ↓, then -	-	-	-	-	↓
Shifted day-fed (SDF)	↓	↑	early ↑, then -	-	↓	↓↓

Table 2. Cumulative changes in physiological parameters relative to the NF group

Our data reveal the presence of sex-specific responses to TRF, especially regarding weight gain. When mice were fed during the lights-on phase (DF, SDF), weight gain slowed in females, but increased in males. Delay shifting the

mealtime by 4 h, independently of the light-dark phase, led to a further weight

gain increase in males (Fig 2A). Struggle for DF and SDF females to retain

weight was also highlighted by the 16 h fasting period, leading to the lowest

weights at the food anticipation timepoint (Fig 2B). Weight gain in DF and

daytime-snacking male mice has been observed before [40, 78, 79], aligning

with our observations; however, we are unaware of studies looking at weight

gain in female mice under TRF. It has been observed that, under light phase

feeding, the majority of physical activity still occurs during the dark phase [80-

82], therefore, further measurements of locomotion and, ideally, firing rates

within the central clock neurons would be needed to determine if DF and SDF

females stay more active during the dark phase when no food is available to

replenish the lost calories. TRF-induced sex-specific changes in food intake

could also explain the observed differences, measuring individual consumption

would be necessary to properly interpret the weight gain data reported here.

Fasting blood glucose levels also showed sex-specific adjustment to TRF. DF in

female mice and SNF in male mice led to a significant decrease in fasting blood

glucose levels, although these differences disappeared under longer

entrainment (Fig 2C). This data contradicts previous observations of DF

inducing higher fasting blood glucose [83], and suggests that glucose levels can

normalize over long-term food entrainment. However, result interpretation would

greatly benefit from measuring individual food intake and glucose levels at other

timepoints.

Not much is known about TRF-driven changes in intestinal barrier function and

epithelial resistance. Studies in this area often use disease models and have

revealed that dark-phase TRF preserves intestinal integrity in mouse colitis

model [84]. Our NF male mice also showed a significant increase in TEER over

time (Fig 2D), complementing these observations. In contrast, circadian

misalignment, caused by changes in the LD cycle and 'wrong-time eating', has

been shown to downregulate the expression of tight junction proteins,

increasing intestinal permeability [85, 86] and contributing to inflammatory

bowel disease (for a review, see [87]). We also observed a significant decrease

in TEER under prolonged SDF of both sexes and DF of males when food- and

light-driven regulation would be misaligned (Fig 2D). Further measurements of

expression of tight junction proteins and their upstream effectors (such as

AMPK [88]), changes in microbiota, and TEER during acute fasting-refeeding

(although fasting alone seems to have no effect on ileal permeability [89]) would

aid in explaining how TRF regulates intestinal barrier function.

723 Hepatic, gastric, and duodenal gene expression in response to
724 TRF

Overall, we observed three distinct gene expression responses to food intake

time, summarized in Table 3. For a gene to be classified as non-responsive, it

had to show only one significant timepoint- or shift-specific expression

difference for the whole duration of the experiment. Interestingly, although it

was possible for genes to simultaneously show both timepoint- and shift-specific

changes, they seemed to occur independently from each other.

Table 3. Summary of gene expression responses to food entrainment. Data pooled from all

sacrifice days and TRF regimes.

Organ	Non-responsive genes	Timepoint-specific		Shift-specific	
		Females	Males	Females	Males
Liver	*Tas1r2* *Pept1, B⁰at1* *Il-17a, Il-6, Crp, Igf-1*	*Clock, Pepck, Mttp (males only), Srebf1 (males only)*			
		Bmal1, Glut2, Gys2, Pparα, Fasn, Cyp7a1, Nampt, Fgf21, TNFα			*Bmal1, Glut2, Sglt1, Gys2, Pparα, Fasn,*

					Cyp7a1, *Gpx1*
Stomach	*Tas1r2, Sglt1, Glut2* *Pept1, B⁰at1* *Mttp, Fasn, Cyp7a1* *Il-17a*	*Bmal1*, *Clock*, *Pepck*, *Gast*, *TNFα*			
		Pparα, *Tph1*, *Lepr*, *Ghrl*		*Il-6, Gpx2*	
		Sst	*Il-6, Gpx2*	*Tph1*	*Pparα*, *Sst*, *Pga5*, *Ghrl*
Duodenum	*Cyp7a1, Mttp, Fasn* *Sst, Ghrl* *Il-6, Il-17a*	*Tas1r2*, *Pepck*			
		Bmal1, *Sglt1*, *Glut2*, *Pept1*, *B⁰at1*, *Pparα*, *Gpx2*		*Gcg*	
		Gcg	*Clock*, *Gip*, *Cck*, *Tph1*, *TNFα*	*Glut2*	*Bmal1*, *Sglt1*, *Pept1*, *B⁰at1*, *Pparα*, *Gip*, *Cck*, *Tph1*, *TNFα*

Colors mark different assigned groups by gene product function: light blue – core clock genes;

purple – glucose sensing, transport, and metabolism; orange – lipid metabolism; pink –

biosynthesis of NAD+; red – amino acid and peptide transport, green – hormone signaling; grey

– pro-/anti-inflammatory response.

Our data (Figs 3-6) led to three major conclusions: 1) digestive organs entrain

to TRF at different rates; 2) DF drives an overall gene downregulation in the

liver and an upregulation in the stomach and duodenum when compared to NF;

3) male and female mice can exhibit opposite timepoint-specific gene

expression patterns, with males being more likely to also develop shift-specific

responses.

Organ- and tissue-specific patterns in gene expression have been observed

before, with surprisingly little overlap in cyclic genes and their phases, even if

the tested genes are ubiquitously expressed and exposed to the same

conditions [8, 22, 90-92]. This is similar to our observations, where the same

gene could show timepoint-specific differences in one organ, but not in the

others (e.g., *Pparα*) and where genes belonging to the same signaling pathway

had no similarity in their responses (e.g., hepatic *Fgf21* and its upstream

regulator *Pparα* [93]). At the same time, the gene *Pepck*, encoding a rate-

limiting enzyme in gluconeogenesis, showed a similar timepoint-specific

expression pattern in all organs, especially under SNF (Figs 4-6), demonstrating

potential organ-to-organ communication. It appears that TRF is unable to

synchronize timepoint-specific gene expression across the whole digestive

system.

However, there was one major impact of TRF that spanned across two organs

and all tested genes in both sexes. We did observe an overall gene

upregulation in the stomach and duodenum and downregulation in the liver

under DF and SDF (Figs 4-6). In the literature, such responses are rarely

described, and their causes thus have not been discussed. A study by Deota et

al. did show that TRF-specific gene up-/down-regulation patterns are similar

between murine stomach and jejunum, but not liver and that specific gene

groups can be upregulated in one organ, but downregulated in another [92],

somewhat aligning with our observations. It has also been shown that fasting

can induce gene expression changes in an organ-specific manner [94]; thus,

further studies applying acute fasting and TRF in constant conditions would be

needed to differentiate between fasting and TRF as causes for large-scale gene

expression changes. In addition, transcriptomics and/or proteomics of multiple

organs could provide further information on TRF-driven peripheral gene

expression.

Although sex-specificity in circadian behavior has been reported before (for a

review, see [95]), sex-specificity in TRF responses (or any other peripheral

Zeitgeber, for that matter) has been understudied. Studies profiling metabolites

have observed considerable sex-specific differences under TRF and other diets,

especially in the liver [96, 97]. Also, the impact of fasting and refeeding on

weight gain, energy expenditure, and regulation of appetite and nutrient

metabolism has been shown to be sex-specific [98]. Our study revealed a

considerable overlap of timepoint-specific responses in males and females;

however, sex-specificity was strongly present in expression pattern changes

over time and shift-specific responses. Long-term TRF studies with back-and-

forth shifts would be beneficial to determine how fast males and females adjust

to changes in food intake time and how stable the developed responses are.

Such studies could also provide insights into the overall health impact of shifted

or restricted feeding, which could be particularly important for shift workers and

those often traveling across time zones.

Overall, our study demonstrates the complexity of TRF responses which are

sex-, organ- and gene-specific and depend on the length of entrainment. It

stresses the importance of including females in TRF and other Zeitgeber

studies. Further details and integrative exploration of TRF under constant

conditions and acute fasting-refeeding phases would be necessary to delineate

the food-, fasting- and light-driven changes. In addition, increased sampling

frequency would provide a more detailed timeline of the (sex-specific)

adjustment to food intake time to fully unravel the interplay between sex,

circadian rhythms, and food intake.

Acknowledgments

We thank Single Cell and High Throughput Genomics Platform at Monash Health Translation project, especially Dr. Sen Wang, for performing the IFC analysis. We also thank Prof. Frédéric Gachon from the Institute of Bioscience at University of Queensland for the constructive exchange regarding experimental approaches and methodology.

Competing interests

The authors of this study confirm there are no competing interests.

Author contributions

All experiments were carried out at the Department of Anatomy and Physiology, The University of Melbourne, and the Florey Institute of Neuroscience and Mental Health. L.O-R., M.R.D., R.M., and J.B.F. designed the study and necessary experiments. M.R.D., L.O-R., R.M., B.H., A.K., M.R., T.E.F.C., and L.J.F carried out the experiments. L.O-R., M.R.D., R.M, and J.B.F. analyzed and interpreted the data. L.O-R., M.R.D., and R.M. wrote and critically revised the manuscript. All authors approved the final version of the manuscript. All persons qualifying for authorship are listed as such.

Funding

This study was supported by the Swiss National Science Foundation Grant No. 187739 to L.O-R.

References

1. Albrecht U. Timing to Perfection: The Biology of Central and Peripheral Circadian Clocks. Neuron. 2012;74(2):246-60. doi: https://doi.org/10.1016/j.neuron.2012.04.006.

2. Kleitman N, Engelman T. Diurnal cycle in activity and body temperature of rabbits. Fed Proc. 1947;6(1):143.

3. Rodbard S. Body temperature-arterial pressure relationship as a basis for physiological interpretation of diurnal rhythm. Fed Proc. 1947;6(1):191.

4. Mellette H, Hutt B, Askovitz, SI, Horvath S. Diurnal variations in body temperatures. J Appl Physiol. 1951;3(11):665-75.

5. Goetzl F, Stone F. Diurnal variations in acuity of olfaction and food intake. Gastroenterology. 1947;9(4):444-53.

6. Sirota J, Baldwin D, Villarreal, H. Diurnal variations of renal function in man. J Clin Invest. 1950;29(2):187-92.

7. Mills J. Diurnal rhythm in urine flow. J Physiol. 1951;113(4):528-36.

8. Zhang R, Lahens NF, Ballance HI, Hughes ME, Hogenesch JB. A circadian gene expression atlas in mammals: implications for biology and medicine. Proc Natl Acad Sci U S A. 2014;111(45):16219-24. Epub 20141027. doi: 10.1073/pnas.1408886111. PubMed PMID: 25349387; PubMed Central PMCID: PMCPMC4234565.

9. Fagiani F, Di Marino D, Romagnoli A, Travelli C, Voltan D, Di Cesare Mannelli L, et al. Molecular regulations of circadian rhythm and implications for physiology and diseases. Signal Transduction and Targeted Therapy. 2022;7(1):41. doi: 10.1038/s41392-022-00899-y.

10. Takahashi J, Lowrey P. Genetics of the mammalian circadian system: Photic entrainment, circadian pacemaker mechanisms, and posttranslational regulation Annu Rev Genet. 2000;34:533-62.

11. Buhr E, Takahashi J. Molecular components of the mammalian circadian clock. Handb Exp Pharmacol. 2013;217:3-27.

12. Buijs F, Leon-Mercado L, Guzman-Ruiz M, Guerrero-Vargas N, Romo-Nava F, Buijs R. The circadian system: a regulatory feedback network of periphery and brain Physiology. 2016;31(3):170-81.

13. Hastings M, Maywood E, Brancaccio M. Generation of circadian rhythms in the suprachiasmatic nucleus Nat Rev Neurosci. 2018;19(8):453-69.

14. Cox K, Takahashi J. Circadian clock genes and the transcriptional
architecture of the clock mechanism J Mol Endocrinol. 2019;63(4):R93-
R102.

15. Duez H, Staels B. The nuclear receptors Rev-erbs and RORs integrate
circadian rhythms and metabolism. Diab Vasc Dis Res. 2008;5(2):82-8.

16. Solt L, Kojetin D, Burris T. The REV-ERBs and RORs: molecular links
between circadian rhythms and lipid homeostasis. Future Med Chem.
2011;3(5):623-38.

17. Mills J. Circadian rhythms during and after three months in solitude
underground. J Physiol. 1964;174:217-31.

18. Forni D, Pozzoli U, Cagliani R, Tresoldi C, Menozzi G, Riva S, et al.
Genetic adaptation of the human circadian clock to day-length latitudinal
variations and relevance for affective disorders Genome Biol.
2014;15(10):DOI: 10.1186/s13059-014-0499-7

19. Eastman C, Tomaka V, Crowley S. Circadian rhythms of European and
African-Americans after a large delay of sleep as in jet lag and night work.
Sci Rep. 2016;6(36716):DOI: 10.1038/srep36716

20. Crowley S, Eastman C. Free-running circadian period in adolescents and
adults J Sleep Res. 2018;27(5):DOI: 10.1111/jsr.12678

21. Aschoff J. Zeitgeber der tierischen Tagesperiodik. Naturwissenschaften.
1945;41:49-56.

22. Mure LS, Le HD, Benegiamo G, Chang MW, Rios L, Jillani N, et al. Diurnal
transcriptome atlas of a primate across major neural and peripheral tissues.
Science. 2018;359(6381). Epub 20180208. doi: 10.1126/science.aao0318.
PubMed PMID: 29439024; PubMed Central PMCID: PMCPMC5924732.

23. Damiola F, Le Minh N, Preitner N, Kornmann B, Fleury-Olela F, Schibler U.
Restricted feeding uncouples circadian oscillators in peripheral tissues from
the central pacemaker in the suprachiasmatic nucleus. Genes Dev.
2000;14(23):2950-61. doi: 10.1101/gad.183500. PubMed PMID: 11114885.

24. Welsh DK, Takahashi JS, Kay SA. Suprachiasmatic nucleus: cell autonomy
and network properties. Annu Rev Physiol. 2010;72:551-77. doi:
10.1146/annurev-physiol-021909-135919. PubMed PMID: 20148688;
PubMed Central PMCID: PMCPMC3758475.

25. Balakrishnan A, Stearns AT, Ashley SW, Tavakkolizadeh A, Rhoads DB.
Restricted feeding phase shifts clock gene and sodium glucose

cotransporter 1 (SGLT1) expression in rats. J Nutr. 2010;140(5):908-14.
Epub 20100303. doi: 10.3945/jn.109.116749. PubMed PMID: 20200113;
PubMed Central PMCID: PMCPMC2855260.

26. Kuroda H, Tahara Y, Saito K, Ohnishi N, Kubo Y, Seo Y, et al. Meal
frequency patterns determine the phase of mouse peripheral circadian
clocks. Scientific Reports. 2012;2(1):711. doi: 10.1038/srep00711.

27. Schibler U, Gotic I, Saini C, Gos P, Curie T, Emmenegger Y, et al. Clock-
Talk: Interactions between Central and Peripheral Circadian Oscillators in
Mammals. Cold Spring Harb Symp Quant Biol. 2015;80:223-32. Epub
20151218. doi: 10.1101/sqb.2015.80.027490. PubMed PMID: 26683231.

28. Lewis P, Korf HW, Kuffer L, Groß JV, Erren TC. Exercise time cues
(zeitgebers) for human circadian systems can foster health and improve
performance: a systematic review. BMJ Open Sport Exerc Med.
2018;4(1):e000443. Epub 20181205. doi: 10.1136/bmjsem-2018-000443.
PubMed PMID: 30687511; PubMed Central PMCID: PMCPMC6330200.

29. Gillman AG, Rebec GV, Pecoraro NC, Kosobud AEK. Circadian
entrainment by food and drugs of abuse. Behav Processes. 2019;165:23-8.
Epub 20190524. doi: 10.1016/j.beproc.2019.05.017. PubMed PMID:
31132444; PubMed Central PMCID: PMCPMC6703158.

30. Begemann K, Neumann A-M, Oster H. Regulation and function of extra-
SCN circadian oscillators in the brain. Acta Physiologica.
2020;229(1):e13446. doi: https://doi.org/10.1111/apha.13446.

31. Baron KG, Reid KJ. Circadian misalignment and health. Int Rev Psychiatry.
2014;26(2):139-54. doi: 10.3109/09540261.2014.911149. PubMed PMID:
24892891; PubMed Central PMCID: PMCPMC4677771.

32. Fishbein AB, Knutson KL, Zee PC. Circadian disruption and human health.
The Journal of Clinical Investigation. 2021;131(19). doi:
10.1172/JCI148286.

33. Asher G, Sassone-Corsi P. Time for food: the intimate interplay between
nutrition, metabolism, and the circadian clock. Cell. 2015;161(1):84-92. doi:
10.1016/j.cell.2015.03.015. PubMed PMID: 25815987.

34. Voigt RM, Forsyth CB, Keshavarzian A. Circadian rhythms: a regulator of
gastrointestinal health and dysfunction. Expert Rev Gastroenterol Hepatol.
2019;13(5):411-24. Epub 2019/03/25. doi:
10.1080/17474124.2019.1595588. PubMed PMID: 30874451.

35. López-Olmeda JF, Tartaglione EV, de la Iglesia HO, Sánchez-Vázquez FJ.
Feeding entrainment of food-anticipatory activity and per1 expression in the
brain and liver of zebrafish under different lighting and feeding conditions.
Chronobiol Int. 2010;27(7):1380-400. doi: 10.3109/07420528.2010.501926.
PubMed PMID: 20795882.

36. Gómez-Boronat M, De Pedro N, Alonso-Gómez Á L, Delgado MJ, Isorna E.
Nuclear Receptors (PPARs, REV-ERBs, RORs) and Clock Gene Rhythms
in Goldfish (Carassius auratus) Are Differently Regulated in Hypothalamus
and Liver. Front Physiol. 2022;13:903799. Epub 20220606. doi:
10.3389/fphys.2022.903799. PubMed PMID: 35733989; PubMed Central
PMCID: PMCPMC9207440.

37. Díaz-Muñoz M, Vázquez-Martínez O, Aguilar-Roblero R, Escobar C.
Anticipatory changes in liver metabolism and entrainment of insulin,
glucagon, and corticosterone in food-restricted rats. Am J Physiol Regul
Integr Comp Physiol. 2000;279(6):R2048-56. doi:
10.1152/ajpregu.2000.279.6.R2048. PubMed PMID: 11080068.

38. Oyama Y, Iwasaka H, Koga H, Shingu C, Matsumoto S, Noguchi T.
Uncoupling of peripheral and master clock gene rhythms by reversed
feeding leads to an exacerbated inflammatory response after polymicrobial
sepsis in mice. Shock. 2014;41(3):214-21. doi:
10.1097/shk.0000000000000094. PubMed PMID: 24300828.

39. Greenwell BJ, Trott AJ, Beytebiere JR, Pao S, Bosley A, Beach E, et al.
Rhythmic Food Intake Drives Rhythmic Gene Expression More Potently
than the Hepatic Circadian Clock in Mice. Cell Reports. 2019;27(3):649-
57.e5. doi: https://doi.org/10.1016/j.celrep.2019.03.064.

40. Jang H, Lee G, Kong J, Choi G, Park YJ, Kim JB. Feeding Period
Restriction Alters the Expression of Peripheral Circadian Rhythm Genes
without Changing Body Weight in Mice. PLOS ONE. 2012;7(11):e49993.
doi: 10.1371/journal.pone.0049993.

41. Dailey MJ, Stingl KC, Moran TH. Disassociation between preprandial gut
peptide release and food-anticipatory activity. Endocrinology.
2012;153(1):132-42. Epub 20111129. doi: 10.1210/en.2011-1464. PubMed
PMID: 22128024; PubMed Central PMCID: PMCPMC3249668.

42. Segers A, Desmet L, Sun S, Verbeke K, Tack J, Depoortere I. Night-time
feeding of Bmal1-/- mice restores SCFA rhythms and their effect on ghrelin.

J Endocrinol. 2020;245(1):155-64. doi: 10.1530/joe-20-0011. PubMed
PMID: 32045364.

43. Laermans J, Broers C, Beckers K, Vancleef L, Steensels S, Thijs T, et al.
Shifting the circadian rhythm of feeding in mice induces gastrointestinal,
metabolic and immune alterations which are influenced by ghrelin and the
core clock gene Bmal1. PLoS One. 2014;9(10):e110176. Epub 20141016.
doi: 10.1371/journal.pone.0110176. PubMed PMID: 25329803; PubMed
Central PMCID: PMCPMC4199674.

44. Rust BM, Picklo MJ, Yan L, Mehus AA, Zeng H. Time-Restricted Feeding
Modifies the Fecal Lipidome and the Gut Microbiota. Nutrients. 2023;15(7).
Epub 20230323. doi: 10.3390/nu15071562. PubMed PMID: 37049404;
PubMed Central PMCID: PMCPMC10096715.

45. Rangan P, Choi I, Wei M, Navarrete G, Guen E, Brandhorst S, et al.
Fasting-Mimicking Diet Modulates Microbiota and Promotes Intestinal
Regeneration to Reduce Inflammatory Bowel Disease Pathology. Cell Rep.
2019;26(10):2704-19.e6. doi: 10.1016/j.celrep.2019.02.019. PubMed PMID:
30840892; PubMed Central PMCID: PMCPMC6528490.

46. Polidarová L, Sládek M, Soták M, Pácha J, Sumová A. Hepatic, duodenal,
and colonic circadian clocks differ in their persistence under conditions of
constant light and in their entrainment by restricted feeding. Chronobiol Int.
2011;28(3):204-15. doi: 10.3109/07420528.2010.548615. PubMed PMID:
21452916.

47. Sládek M, Rybová M, Jindráková Z, Zemanová Z, Polidarová L, Mrnka L, et
al. Insight into the circadian clock within rat colonic epithelial cells.
Gastroenterology. 2007;133(4):1240-9. Epub 20070602. doi:
10.1053/j.gastro.2007.05.053. PubMed PMID: 17675004.

48. Hoogerwerf WA, Hellmich HL, Cornélissen G, Halberg F, Shahinian VB,
Bostwick J, et al. Clock gene expression in the murine gastrointestinal tract:
endogenous rhythmicity and effects of a feeding regimen.
Gastroenterology. 2007;133(4):1250-60. Epub 20070712. doi:
10.1053/j.gastro.2007.07.009. PubMed PMID: 17919497.

49. Pan X, Terada T, Okuda M, Inui K. The diurnal rhythm of the intestinal
transporters SGLT1 and PEPT1 is regulated by the feeding conditions in
rats. J Nutr. 2004;134(9):2211-5. doi: 10.1093/jn/134.9.2211. PubMed
PMID: 15333706.

50. Segers A, Depoortere I. Circadian clocks in the digestive system. Nature
Reviews Gastroenterology & Hepatology. 2021;18(4):239-51. doi:
10.1038/s41575-020-00401-5.

51. Manella G, Sabath E, Aviram R, Dandavate V, Ezagouri S, Golik M, et al.
The liver-clock coordinates rhythmicity of peripheral tissues in response to
feeding. Nature Metabolism. 2021;3(6):829-42. doi: 10.1038/s42255-021-
00395-7.

52. Carmona-Alcocer V, Brown LS, Anchan A, Rohr KE, Evans JA.
Developmental patterning of peptide transcription in the central circadian
clock in both sexes. Front Neurosci. 2023;17:1177458. Epub 20230519.
doi: 10.3389/fnins.2023.1177458. PubMed PMID: 37274219; PubMed
Central PMCID: PMCPMC10235759.

53. Bailey M, Silver R. Sex differences in circadian timing systems: Implications
for disease. Frontiers in Neuroendocrinology. 2014;35(1):111-39. doi:
https://doi.org/10.1016/j.yfrne.2013.11.003.

54. Yan L, Silver R. Neuroendocrine underpinnings of sex differences in
circadian timing systems. J Steroid Biochem Mol Biol. 2016;160:118-26.
Epub 20151022. doi: 10.1016/j.jsbmb.2015.10.007. PubMed PMID:
26472554; PubMed Central PMCID: PMCPMC4841755.

55. Silver R. Cells have sex chromosomes and circadian clocks: Implications
for organismal level functions. Physiology & Behavior. 2018;187:6-12. doi:
https://doi.org/10.1016/j.physbeh.2017.11.016.

56. Joye DAM, Evans JA. Sex differences in daily timekeeping and circadian
clock circuits. Seminars in Cell & Developmental Biology. 2022;126:45-55.
doi: https://doi.org/10.1016/j.semcdb.2021.04.026.

57. Aguayo A, Martin CS, Huddy TF, Ogawa-Okada M, Adkins JL, Steele AD.
Sex differences in circadian food anticipatory activity are not altered by
individual manipulations of sex hormones or sex chromosome copy number
in mice. PLOS ONE. 2018;13(1):e0191373. doi:
10.1371/journal.pone.0191373.

58. Li Z, Wang Y, Sun KK, Wang K, Sun ZS, Zhao M, et al. Sex-related
difference in food-anticipatory activity of mice. Horm Behav. 2015;70:38-46.
Epub 20150228. doi: 10.1016/j.yhbeh.2015.02.004. PubMed PMID:
25736535.

59. Sellix MT, Evans JA, Leise TL, Castanon-Cervantes O, Hill DD, DeLisser P,
et al. Aging differentially affects the re-entrainment response of central and
peripheral circadian oscillators. J Neurosci. 2012;32(46):16193-202. doi:
10.1523/jneurosci.3559-12.2012. PubMed PMID: 23152603; PubMed
Central PMCID: PMCPMC3507430.

60. Pilorz V, Cunningham PS, Jackson A, West AC, Wager TT, Loudon AS, et
al. A novel mechanism controlling resetting speed of the circadian clock to
environmental stimuli. Curr Biol. 2014;24(7):766-73. Epub 20140320. doi:
10.1016/j.cub.2014.02.027. PubMed PMID: 24656826; PubMed Central
PMCID: PMCPMC3988889.

61. Ramakrishnan R, Qin J, Jones RC, Weaver LS. Integrated Fluidic Circuits
(IFCs) for digital PCR. Methods Mol Biol. 2013;949:423-31. doi:
10.1007/978-1-62703-134-9_27. PubMed PMID: 23329458.

62. Fluidigm. Gene Expression with the 192.24 IFC Using Fast TaqMan Assays
(Biomark HD only): Fluidigm; 2015 [cited 2022]. Available from:
https://fluidigm.my.salesforce.com/sfc/p/#700000009DAw/a/4u0000019jzR/
7DaOMZh8GOtlvivpWQ5SIX5Xj9wPhQx9c1D4yMF0nVc.

63. Xie X, Kukino A, Calcagno HE, Berman AM, Garner JP, Butler MP. Natural
food intake patterns have little synchronizing effect on peripheral circadian
clocks. BMC Biology. 2020;18(1):160. doi: 10.1186/s12915-020-00872-7.

64. Smith JG, Koronowski KB, Mortimer T, Sato T, Greco CM, Petrus P, et al.
Liver and muscle circadian clocks cooperate to support glucose tolerance in
mice. Cell Rep. 2023;42(6):112588. Epub 20230601. doi:
10.1016/j.celrep.2023.112588. PubMed PMID: 37267101; PubMed Central
PMCID: PMCPMC10592114.

65. Hoogerwerf WA, Hellmich HL, Cornélissen G, Halberg F, Shahinian VB,
Bostwick J, et al. Clock Gene Expression in the Murine Gastrointestinal
Tract: Endogenous Rhythmicity and Effects of a Feeding Regimen.
Gastroenterology. 2007;133(4):1250-60. doi:
https://doi.org/10.1053/j.gastro.2007.07.009.

66. Knight BL, Hebbachi A, Hauton D, Brown AM, Wiggins D, Patel DD, et al. A
role for PPARalpha in the control of SREBP activity and lipid synthesis in
the liver. Biochem J. 2005;389(Pt 2):413-21. doi: 10.1042/bj20041896.
PubMed PMID: 15777286; PubMed Central PMCID: PMCPMC1175119.

67. Rivera-Zavala JB, Báez-Ruiz A, Díaz-Muñoz M. Changes in the 24⬚h
Rhythmicity of Liver PPARs and Peroxisomal Markers When Feeding Is
Restricted to Two Daytime Hours. PPAR Research. 2011;2011(1):261584.
doi: https://doi.org/10.1155/2011/261584.

68. Flippo KH, Potthoff MJ. Metabolic Messengers: FGF21. Nat Metab.
2021;3(3):309-17. Epub 20210318. doi: 10.1038/s42255-021-00354-2.
PubMed PMID: 33758421; PubMed Central PMCID: PMCPMC8620721.

69. Patton DF, Mistlberger RE. Circadian adaptations to meal timing:
neuroendocrine mechanisms. Frontiers in Neuroscience. 2013;7. doi:
10.3389/fnins.2013.00185.

70. Davidson AJ, Poole AS, Yamazaki S, Menaker M. Is the food-entrainable
circadian oscillator in the digestive system? Genes, Brain and Behavior.
2003;2(1):32-9. doi: https://doi.org/10.1034/j.1601-183X.2003.00005.x.

71. Walsh JH. Role of Gastrin as a Trophic Hormone. Digestion.
2009;47(Suppl. 1):11-6. doi: 10.1159/000200509.

72. Saha L. Role of peroxisome proliferator-activated receptors alpha and
gamma in gastric ulcer: An overview of experimental evidences. World J
Gastrointest   Pharmacol   Ther.   2015;6(4):120-6.   doi:
10.4292/wjgpt.v6.i4.120. PubMed PMID: 26558146; PubMed Central
PMCID: PMCPMC4635152.

73. Bakke I, Hammer TA, Sandvik AK, Waldum HL. PPAR alpha stimulates the
rat gastrin-producing cell. Mol Cell Endocrinol. 2002;195(1-2):89-97. doi:
10.1016/s0303-7207(02)00187-9. PubMed PMID: 12354675.

74. Troy S, Soty M, Ribeiro L, Laval L, Migrenne S, Fioramonti X, et al.
Intestinal Gluconeogenesis Is a Key Factor for Early Metabolic Changes
after Gastric Bypass but Not after Gastric Lap-Band in Mice. Cell
Metabolism. 2008;8(3):201-11. doi: 10.1016/j.cmet.2008.08.008.

75. Vily-Petit J, Barataud A, Zitoun C, Gautier-Stein A, Serino M, Mithieux G.
Intestinal gluconeogenesis shapes gut microbiota, fecal and urine
metabolome in mice with gastric bypass surgery. Scientific Reports.
2022;12(1):1415. doi: 10.1038/s41598-022-04902-y.

76. Montal ED, Dewi R, Bhalla K, Ou L, Hwang BJ, Ropell AE, et al. PEPCK
Coordinates the Regulation of Central Carbon Metabolism to Promote
Cancer Cell Growth. Mol Cell. 2015;60(4):571-83. Epub 20151017. doi:

10.1016/j.molcel.2015.09.025. PubMed PMID: 26481663; PubMed Central PMCID: PMCPMC4656111.

77. Potts A, Uchida A, Deja S, Berglund ED, Kucejova B, Duarte JA, et al. Cytosolic phosphoenolpyruvate carboxykinase as a cataplerotic pathway in the small intestine. Am J Physiol Gastrointest Liver Physiol. 2018;315(2):G249-g58. Epub 20180406. doi: 10.1152/ajpgi.00039.2018. PubMed PMID: 29631378; PubMed Central PMCID: PMCPMC6139646.

78. Acosta-Rodríguez VA, de Groot MHM, Rijo-Ferreira F, Green CB, Takahashi JS. Mice under Caloric Restriction Self-Impose a Temporal Restriction of Food Intake as Revealed by an Automated Feeder System. Cell Metabolism. 2017;26(1):267-77.e2. doi: https://doi.org/10.1016/j.cmet.2017.06.007.

79. Begemann K, Heyde I, Witt P, Inderhees J, Leinweber B, Koch CE, et al. Rest phase snacking increases energy resorption and weight gain in male mice. Molecular Metabolism. 2023;69:101691. doi: https://doi.org/10.1016/j.molmet.2023.101691.

80. Dattolo T, Coomans CP, van Diepen HC, Patton DF, Power S, Antle MC, et al. Neural activity in the suprachiasmatic circadian clock of nocturnal mice anticipating a daytime meal. Neuroscience. 2016;315:91-103. Epub 20151214. doi: 10.1016/j.neuroscience.2015.12.014. PubMed PMID: 26701294.

81. Acosta-Rodríguez VA, Rijo-Ferreira F, van Rosmalen L, Izumo M, Park N, Joseph C, et al. Misaligned feeding uncouples daily rhythms within brown adipose tissue and between peripheral clocks. Cell Reports. 2024;43(8). doi: 10.1016/j.celrep.2024.114523.

82. Abe T, Kazama R, Okauchi H, Oishi K. Food deprivation during active phase induces skeletal muscle atrophy via IGF-1 reduction in mice. Arch Biochem Biophys. 2019;677:108160. Epub 20191019. doi: 10.1016/j.abb.2019.108160. PubMed PMID: 31639326.

83. Mukherji A, Kobiita A, Chambon P. Shifting the feeding of mice to the rest phase creates metabolic alterations, which, on their own, shift the peripheral circadian clocks by 12 hours. Proceedings of the National Academy of Sciences. 2015;112(48):E6683-E90. doi: doi:10.1073/pnas.1519735112.

84. Zhang X, Zou Q, Zhao B, Zhang J, Zhao W, Li Y, et al. Effects of alternate-
day fasting, time-restricted fasting and intermittent energy restriction DSS-
induced on colitis and behavioral disorders. Redox Biol. 2020;32:101535-.
Epub 2020/04/10. doi: 10.1016/j.redox.2020.101535. PubMed PMID:
32305005.

85. Deaver JA, Eum SY, Toborek M. Circadian Disruption Changes Gut
Microbiome Taxa and Functional Gene Composition. Frontiers in
Microbiology. 2018;9. doi: 10.3389/fmicb.2018.00737.

86. Tran L, Jochum SB, Shaikh M, Wilber S, Zhang L, Hayden DM, et al.
Circadian misalignment by environmental light/dark shifting causes
circadian disruption in colon. PLOS ONE. 2021;16(6):e0251604. doi:
10.1371/journal.pone.0251604.

87. Post Z, Zilberstein NF, Keshavarzian A. The circadian rhythm as
therapeutic target in inflammatory bowel disease. Journal of the Canadian
Association of Gastroenterology. 2024. doi: 10.1093/jcag/gwae027.

88. Sun X, Yang Q, Rogers CJ, Du M, Zhu M-J. AMPK improves gut epithelial
differentiation and barrier function via regulating Cdx2 expression. Cell
Death & Differentiation. 2017;24(5):819-31. doi: 10.1038/cdd.2017.14.

89. Nakamura C, Ishizuka N, Yokoyama K, Yazaki Y, Tatsumi F, Ikumi N, et al.
Regulatory mechanisms of glucose absorption in the mouse proximal small
intestine during fasting and feeding. Sci Rep. 2023;13(1):10838. Epub
20230705. doi: 10.1038/s41598-023-38024-w. PubMed PMID: 37407613;
PubMed Central PMCID: PMCPMC10322893.

90. Storch KF, Lipan O, Leykin I, Viswanathan N, Davis FC, Wong WH, et al.
Extensive and divergent circadian gene expression in liver and heart.
Nature. 2002;417(6884):78-83. Epub 20020421. doi: 10.1038/nature744.
PubMed PMID: 11967526.

91. Doherty CJ, Kay SA. Circadian control of global gene expression patterns.
Annu Rev Genet. 2010;44:419-44. doi: 10.1146/annurev-genet-102209-
163432. PubMed PMID: 20809800.

92. Deota S, Lin T, Chaix A, Williams A, Le H, Calligaro H, et al. Diurnal
transcriptome landscape of a multi-tissue response to time-restricted
feeding in mammals. Cell Metab. 2023;35(1):150-65.e4. doi:
10.1016/j.cmet.2022.12.006. PubMed PMID: 36599299; PubMed Central
PMCID: PMCPMC10026518.

93. Badman MK, Pissios P, Kennedy AR, Koukos G, Flier JS, Maratos-Flier E. Hepatic Fibroblast Growth Factor 21 Is Regulated by PPARα and Is a Key Mediator of Hepatic Lipid Metabolism in Ketotic States. Cell Metabolism. 2007;5(6):426-37. doi: https://doi.org/10.1016/j.cmet.2007.05.002.

94. Kinouchi K, Magnan C, Ceglia N, Liu Y, Cervantes M, Pastore N, et al. Fasting Imparts a Switch to Alternative Daily Pathways in Liver and Muscle. Cell Reports. 2018;25(12):3299-314.e6. doi: https://doi.org/10.1016/j.celrep.2018.11.077.

95. Krizo JA, Mintz EM. Sex Differences in Behavioral Circadian Rhythms in Laboratory Rodents. Frontiers in Endocrinology. 2015;5. doi: 10.3389/fendo.2014.00234.

96. Johnson H, Yates T, Leedom G, Ramanathan C, Puppa M, van der Merwe M, et al. Multi-Tissue Time-Domain NMR Metabolomics Investigation of Time-Restricted Feeding in Male and Female Nile Grass Rats. Metabolites. 2022;12(7). Epub 20220716. doi: 10.3390/metabo12070657. PubMed PMID: 35888782; PubMed Central PMCID: PMCPMC9321200.

97. Wells A, Barrington WT, Dearth S, May A, Threadgill DW, Campagna SR, et al. Tissue Level Diet and Sex-by-Diet Interactions Reveal Unique Metabolite and Clustering Profiles Using Untargeted Liquid Chromatography–Mass Spectrometry on Adipose, Skeletal Muscle, and Liver Tissue in C57BL6/J Mice. Journal of Proteome Research. 2018;17(3):1077-90. doi: 10.1021/acs.jproteome.7b00750.

98. de Souza GO, Wasinski F, Donato J. Characterization of the metabolic differences between male and female C57BL/6 mice. Life Sciences. 2022;301:120636. doi: https://doi.org/10.1016/j.lfs.2022.120636.