Spatial analysis reveals combinative role for natural killer and CD8 T cells in antitumor immunity despite profound MHC class I loss in non-small cell lung cancer

Coordinated NK and CD8 T effector cells deliver type I anti-tumoral immunity

MHC I⁺ Tumor cell

IFNγ

IFNγ⁺ NK cell

IFNγ⁺ CD8 T cell

NK cell-driven lymphocyte response despite significantly diminished immunity in MHC-I⁻ tumors

CD8 T cells

CD4 T cells (Th1?)

MHC I⁻ Tumor cell

IFNγ⁺ NK cells

Authors

Remziye E. Wessel, Nardin Ageeb, Joseph M. Obeid, Ileana Mauldin, Kate A. Goundry, Mahdin Hossain, Ryan D. Gentzler, Nolan A. Wages, Craig L. Slingluff Jr, Timothy N.J. Bullock, Sepideh Dolatshahi†, and Michael G. Brown†

†Co-correspondence:
mgb4n@virginia.edu and sd8us@virginia.edu

In brief
Tumor-infiltrated natural killer (NK) and CD8 T cells display increased evidence of co-localization and coordinated effector activity in human MHC-I-bearing NSCLC tumors.

Spatial analysis reveals combinative role for natural killer and CD8 T cells in antitumor immunity despite profound MHC class I loss in non-small cell lung cancer

Running Title: NK and CD8 T cells antitumor immunity in NSCLC

Remziye E. Wessel,[1] Nardin Ageeb,[2] Joseph M. Obeid,[3] Ileana Mauldin,[4] Kate A. Goundry,[1] Mahdin Hossain,[5] Chad Lehman,[5] Ryan D. Gentzler,[6] Nolan A. Wages,[7] Craig L. Slingluff Jr,[4] Timothy N.J. Bullock,[5,8] Sepideh Dolatshahi,[1,5†] and Michael G. Brown [5,9-11†]

[1]Department of Biomedical Engineering, University of Virginia School of Medicine, Charlottesville, Virginia 22908

[2]Department of Biology, University of Virginia School of Medicine, Charlottesville, Virginia 22908

[3] Department of Thoracic Surgery, Temple University Hospital, Philadelphia, Pennsylvania 19140

[4]Department of Surgery, University of Virginia School of Medicine, Charlottesville, Virginia 22908

[5]Beirne B. Carter Center for Immunology Research, University of Virginia School of Medicine, Charlottesville, Virginia 22908

[6]Department of Medicine, Hematology and Oncology, University of Virginia School of Medicine, Charlottesville, Virginia 22908

[7]Department of Biostatistics, Virginia Commonwealth University, Richmond, Virginia 23298-0032

[8]Department of Pathology, University of Virginia School of Medicine, Charlottesville, Virginia 22908

[9]Department of Medicine, Nephrology, University of Virginia School of Medicine, Charlottesville,

Virginia 22908

[10]Center for Immunity, Inflammation and Regenerative Medicine, University of Virginia School of

Medicine, Charlottesville, Virginia 22908

[11]Department of Microbiology, Immunology and Cancer Biology, University of Virginia School of

Medicine, Charlottesville, Virginia 22908

[†]Shared senior authorship.

Keywords: MHC class I, CD8 T cells, natural killer cells, tumor-infiltrating lymphocytes, inter-

feron-gamma, multiplex immunofluorescence, patient survival, tissue micro-array, spatial analy-

sis, systems immunology, intratumor heterogeneity, cell co-localization

Availability of data and material – Quantitative IHC and mIF data are available in a public, open-

access repository (UVA Libra). Algorithms developed for this study (O-PLSDA and cell neighbor-

hood scoring) are available on GitHub (https://github.com/Dolatshahi-Lab). IHC and mIF images

will be made available by the authors upon reasonable request.

Competing interests – CLS - Research support to the University of Virginia from Celldex (fund-

ing, drug), Merck (funding, drug), Theraclion (device staff support); Funding to the University of

Virginia from Polynoma for PI role on the MAVIS Clinical Trial; Funding to the University of Virginia

for roles on Scientific Advisory Boards for Immatics and CureVac. CLS also receives licensing fee

payments through the UVA Licensing and Ventures Group for patents for peptides used in cancer

vaccines. RDG – Research support to the University of Virginia from Pfizer, Amgen, Chugai,

Merck, AstraZeneca, Janssen, Daiichi Sankyo, Alliance Foundation, Takeda, ECOG/ACRIN,

Jounce Therapeutics, Bristol Myers Squibb, SWOG, Helsinn, Dizal Pharmaceuticals, and Mirati.

RDG received payment for service on Scientific Advisory Boards including AstraZeneca, Takeda,

Gilead, Janssen, Mirati, Daiichi Sankyo, Sanofi, Oncocyte, Jazz Pharmaceuticals, Blueprint Med-

icines, and Merus.

Funding – This work was funded by a Cancer Team Science Award from the University of Vir-

ginia Comprehensive Cancer Center (TNJB & MGB), and a Collaborative Research Award from

the University of Virginia Beirne Carter Center for Immunology Research (TNJB, SD & MGB).

The UVA Comprehensive Cancer Center Support Grant (P30CA044579) from the NCI addition-

ally supported work performed by the Biorepository and Tissue Research Facility (BTRF) and

Molecular Immunologic and Translational Science (MITS) cores. REW received training support

from National Institute of General Medical Sciences of the National Institutes of Health (T32-

GM145443). JMO received training support from National Cancer Institute T32 CA163177 and

the University of Virginia Department of Surgery. MH received training support through the

University of Virginia College Science Scholar Summer Research stipend. CL received

training support through the University of Virginia School of Medicine Summer Medical

Research Internship Program.

Authors contributions – REW, SD and MGB wrote the manuscript with input from all authors.

REW, NA, JMO, IM, KAG, CL, RDG, CLS, TNJB, SD and MGB contributed to the conception,

design, and development of the project. REW, NA, JMO, IM, KAG, MH, and NW performed ex-

periments and/or analyzed data. All authors read and approved the final version of the manuscript.

Acknowledgements – This work has been presented at the 2023 American Association of Im-

munologists (AAI) Annual Meeting (Abstract 62.08). We thank the University of Virginia BTRF and

MITS cores at UVA for outstanding technical support. We thank Gabriel Hanson for helpful dis-

cussion.

Authors' information (optional) –

Michael G. Brown – ORCID iD 0000-0003-2872-7685

Timothy N.J. Bullock – ORCID iD 0000-0001-6141-3261

Sepideh Dolatshahi – ORCID iD 0000-0003-0226-0933

Ryan D. Gentzler – ORCID iD 0000-0002-8983-6677

Joseph Obeid – ORCID iD 0000-0002-6111-9289

Craig L. Slingluff – ORCID ID 0000-0002-6664-4373

Remziye E. Wessel – ORCID iD 0000-0002-8596-4700

List of abbreviations – Non-small cell lung cancer, NSCLC; squamous cell carcinoma, SCC;

MHC class I, MHC-I; natural killer cell, NK cell; multiplexed immunofluorescence, mIF; interferon

gamma, IFNγ; checkpoint inhibitor immunotherapy, CPI; orthogonalized partial least squares dis-

criminant analysis, OPLSDA; variable importance in projection, VIP.

ABSTRACT

Background. MHC class I (MHC-I) loss is frequent in NSCLC and renders tumor cells resistant to T cell lysis. NK cells kill MHC-I deficient tumor cells, and although previous studies indicate their presence in NSCLC tumor margins, they were also functionally impaired. The goal of this study was to evaluate whether NK cell and CD8 T cell infiltration and activation vary with MHC-I expression. **Methods**. We used single-stain immunohistochemistry (IHC) and Kaplan-Meier analysis to test the effect of NK cell and CD8 T cell infiltration on overall and disease-free survival. To delineate immune covariates of MHC-I-disparate lung cancers, we used multiplexed immunofluorescence (mIF) imaging followed by multivariate statistical modeling. To identify differences in infiltration and intercellular communication between IFNγ-activated and non-activated lymphocytes, we developed a computational pipeline to enumerate single cell neighborhoods from mIF images followed by multivariate discriminant analysis. **Results**. Spatial quantitation of tumor cell MHC-I expression revealed intra- and inter-tumoral heterogeneity, which was associated with the local lymphocyte landscape. IHC analysis revealed that high numbers of CD56$^+$ cells in patient tumors were associated with disease-free and overall survival (p<0.05). The overall survival association strengthened considerably with high counts of both CD56$^+$ and CD8$^+$ cells (p<1x10^{-4}). mIF imaging and multivariate discriminant analysis revealed an enrichment of both CD3$^+$CD8$^+$ T cells and CD3$^-$CD56$^+$ NK cells in MHC-I-bearing tumors (p<0.05). To infer associations of functional cell states and local cell-cell communication, we analyzed spatial single cell neighborhood profiles to delineate the cellular environments of IFNγ$^{+/-}$ NK cells and T cells. We discovered that both IFNγ$^+$ NK and CD8 T cells were more frequently associated with other IFNγ$^+$ lymphocytes in comparison to IFNγ$^-$ NK cells and CD8 T cells (p<1x10^{-30}). Moreover, IFNγ$^+$ lymphocytes were most often found clustered near MHC-I$^+$ tumor cells. **Conclusions**. We demonstrate that tumor-infiltrating NK cells and CD8 T cells jointly affect vital control of NSCLC tumor progression. Robust association of NK and CD8 T cells was most evident in MHC-I-bearing tumors, especially in the

presence of IFNγ. Frequent co-localization of IFNγ[+] NK cells with other IFNγ[+] lymphocytes in

near-neighbor analysis indicates lymphocyte activation is coordinately regulated in NSCLC.

What is already known on this topic – MHC-I loss occurs frequently in human NSCLC and

corresponds with a waning immune response in the tumor microenvironment (TME). NK cells

recognize "missing-self" targets and thus may be leveraged to target NSCLC tumors with sub-

stantial class I loss. While NK cell presence at tumor margins has been documented in the NSCLC

TME, they have been shown to lose function in this environment.

What this study adds – We developed spatial statistical analysis pipelines that leverage the local

heterogeneity of the tumor microenvironment at single cell resolution to rigorously test whether

NK cells and T cells together contribute antitumoral immunity in human lung cancer. We discov-

ered that a high density of tumor-infiltrating NK cells corresponded with disease-free survival, and

this association was considerably increased in patients with high coincident T cells, especially

those in central tumor. Intriguingly, both cell types were further found clustered together in MHC-

I-bearing tumors, especially when both were also expressing IFNγ, thus suggesting coordinated

lymphocyte activities may give rise to enhanced immune control of human lung cancer.

How this study might affect research, practice, or policy – This study provides a rationale for

developing novel immunotherapies that simultaneously increase NK and T cell anti-tumoral im-

munity. Robust associations linking NK cells with survival and IFNγ activity in NSCLC, including

in tumors with profound MHC-I loss, provides a basis for further research to evaluate the effect of

NK cells in MHC-I-deficient, CD8 T cell refractory NSCLC tumors.

BACKGROUND

Lung cancer is the leading cause of cancer-related death in the United States and globally, and

non-small cell lung cancer (NSCLC) accounts for approximately 85% of all lung cancers (1, 2).

Adenocarcinoma (AdenoCA) and squamous cell carcinoma (SCC) together account for the ma-

jority (~90%) of all NSCLC subtypes (3, 4). Both tumor types are infiltrated by diverse immune

cell types, including T cells and NK cells with cytokine and cytotoxic effector activity (5), highlight-

ing the opportunity to harness these cytolytic cells to mediate more efficacious control of human

lung cancer.

Tumor cell MHC class I (MHC-I) loss to evade recognition by cytotoxic T cells is common

in NSCLC, occurring in over 30% of human lung cancer cells and with over 90% having defects

of one or more MHC-I antigens and other molecules associated with antigen presentation (23).

This constitutes a logical evolutionary step given the propensity of NSCLC to have a large reper-

toire of antigenic mutations that can be presented to T cells on MHC-I (24). Despite the frequency

of MHC-I loss, its impact on lymphocyte infiltration and patient survival has not been resolved (25-

31).

The density of CD8 T cells infiltrating NSCLC tumors generally has been associated with

a favorable prognosis (6-8). However, assessment of patient risk based on this determination

alone is challenging due to factors such as the heterogeneity of intratumoral CD8 T cell infiltrates,

the inherent difficulty in reliably sampling and assessing tumor-infiltrating lymphocytes, a plausible

role for other immune cells, and because T cell exhaustion can contribute to poor immunological

control of lung cancers (9-11), thus implicating additional risk determinants. On the other hand,

NK cells have been shown to provide protection against hematological malignancies or metasta-

ses from primary tumors, yet their antitumor activity has been less evident in solid tumors (12,

13). Given their ability to recognize and lyse 'missing-self' targets, NK cells represent a presently

unharnessed, yet logical candidate to improve anti-tumor immunity in MHC-I deficient, CD8 T cell

refractory lung tumors (32).

Prior work has shown that NK cells are prevalent in β2-microglobulin (β2m-)deficient

NSCLC tumors (30), but intratumoral NK cells from NSCLC patients were found to have reduced

IFNγ production and cytolytic capability compared to circulating NK cells from the same patients

(33). Nonetheless, tumor-infiltrating NK cells activated by cytokines or stimulated via direct recog-

nition of tumor cell expressed ligands of NK activating receptors have been shown to enhance

tumor-specific T cells and control of tumor progression , possibly via cytokines to enhance recruit-

ment or activation of classical dendritic cells (cDC[OBJ][OBJ]. Understanding NK cell and CD8 T cell

interactions that support the anti-tumor immune response in relation to MHC-I expression in the

NSCLC tumor microenvironment (TME) is thus a priority.

Hypothesizing that NK cells and CD8 T cells might together drive anti-tumor immunity in

NSCLC in relation to the level of tumor cell MHC-I expression, we investigated whether tumor-

infiltrating NK cells and CD8 T cells were associated with patient survival, and if tumor cell MHC-

I expression corresponded with variations in lymphocyte presence, activation status, or near-

neighbor groupings in the NSCLC TME. Our findings suggest that novel strategies designed to

target and leverage both T cells and NK cells in combined immune therapy may provide enhanced

antitumor immunity and increased T cell activation, even in CD8 T cell refractory tumors.

MATERIALS AND METHODS

Immunohistochemistry. The AJCC 7[th] edition was used for clinical and pathological staging. An

experienced pathologist outlined tumor areas and tumor margins. To identify regions of interest

for further analysis, Hematoxylin-eosin staining was performed on sections from formalin-fixed,

paraffin embedded (FFPE) tissues. Central tumor (CT) and peripheral tumor (PT) 1 mm cores

were then extracted from the FFPE tissues based on the pathologist's annotation and used to

construct tissue microarray (TMA) samples from each of the 151 tumors as described (34).TMA

tissue sections were deparaffinized and rehydrated (34). Heat-induced antigen retrieval was per-

formed in citrate buffer for most antibodies (Abs), or in EDTA/high pH buffer for Abs to PH:10 as

described (34). NSCLC tissues were incubated with primary Abs to β2-microglobulin (mAb NAMB-

1, kindly provided by Soldano Ferrone), CD3 (mAb DAKO, clone 7.2.38), CD8 (clone C8/144B,

DAKO Carpinteria, CA), CD56 (mAb NCL-L-CD56-1B6, Leica), Granzyme B (Abcam ab4059),

and HLA class I heavy chains (HC) (HC10 (35), Ferrone lab). Enzyme-linked secondary antibod-

ies and 3,3'-diaminobenzidine (Dako) were used to develop the staining. Ab-stained slides were

counterstained with hematoxylin (Dako) and permanently mounted as described (34). Images

were acquired using a Nikon Digital Sight DS-Fi1 camera (Nikon Corporation, Tokyo, Japan), an

Eclipse 400 microscope (Nikon), or a NIS-Elements Viewer version 3.2 (Nikon). Staining control

tissues included tonsil, lymph node, and placenta. Tissues incubated with phosphate buffered

saline (PBS) instead of primary Ab and then stained with secondary Abs only served as negative

staining controls.

TMA Immunotyping. Tumor cell antigen expression (e.g., HLA class I) was assessed by a

pathologist and scored by their expression intensity (0 = no staining, 1 = mild staining, 2 = strong

staining) in each core. Scores from two cores per patient were summed to a total score that ranges

from zero to four in each tumor to account for tumor heterogeneity. Automated CD8 counting was

performed using a Leica SCN400 Scanner (Leica, Nussloch, Germany) and Digital Imaging Hub

(Leica) and validated by manual counting by a pathologist. Immune cell counts were reported as

counts per millimeter squared (mm^2).

mIF imaging and geospatial analysis. Five (5) µm-thick sections were cut from FFPE NSCLC

patient tissues from Cohort 2 (**table 2**). Human lymph node samples were used as a positive

control. mIF was performed according to the manufacturer's protocol using the OPAL Multiplex

Manual IHC kit and antigen retrieval (AR) buffers AR6 and AR9 (Akoya Biosciences, Marlborough,

Massachusetts, USA) and DIVA Decloaker AR buffer (Biocare Medical, Pacheco, California,

USA). Staining sequence, antibodies and AR buffers used are as follow: AR9, CD3 (1:100, Cell

Marque MRQ-39) Opal520; AR9, CD8 (1:500, clone C8/144B Agilent Technologies, Santa Clara,

California, USA) Opal540; AR6, CD56 (1:1000, CD56 (Cell Marque MRQ-42) Opal570; DIVA,

IFNγ (1:1000 NeoBiotechnologies 345 MSM4 P2 IFNG/466) Opal620; AR6, MHCI (1:15,000 HLA

class I heavy chains (HC10,(35) kindly provided by Soldano Ferrone) Opal650; AR6 Pancyt-

okeratin (1:200 MA513203, Invitrogen) Opal690, and AR6, spectral DAPI (Akoya Biosciences,

Marlborough, Massachusetts, USA) as described previously (36).

Stained slides were mounted using prolong diamond antifade (Life Technologies, Carls-

bad, California, USA) and scanned using the PerkinElmer Vectra 3.0 system and Vectra software

(Akoya Biosciences, Marlborough, Massachusetts, USA). Regions (3 mm^2) were identified using

Phenochart software and tissue images were acquired at 20x magnification with the Vectra 3.0

system. The images were spectrally unmixed using single stain positive control images in the

InForm software (Akoya Biosciences, Marlborough, Massachusetts, USA). A tissue classifier was

trained on panCK$^+$ tumor cells and panCK$^-$ stroma using HALO software (Indica Labs, Albuquer-

que, New Mexico, USA). Immune cells were enumerated using HALO software (Highplex FL

module). Lymphocyte density was quantified as density of cells expressing a given marker

(cells/mm^2) within stromal or tumor compartments of each region.

To increase statistical power while avoiding size-related representational biases in re-

sected patient tissues, a patient-agnostic, tissue regional analysis was developed to analyze

scanned images of lung adenocarcinoma and SCC patient tumor biopsies as independent entities

(**supplemental figure S1**). In total, 394 regions across 29 adenocarcinoma tumors and 108 re-

gions across 7 SCC tumors were collected. Regions with evident tumor cell CD56 expression or

fewer than 50 cells enumerated were excluded. The remaining 361 adenocarcinoma regions and

89 SCC regions were analyzed. To validate that over-representation of tumors with a greater

number of images did not bias our results, we iteratively sampled a random subset of images with

equal representation from all tumors and observed similar results.

Correlation analyses. Spearman correlation analyses of Cohort 1 and Cohort 2 were performed

in R statistical computing software (version 4.1.2) using the "corr" package. Visualizations were

generated using "ggplot2" and "corrplot" packages. Slides or regions with missing data were omit-

ted only from analyses requiring the use of those missing variables.

Survival analyses. Univariate and multivariate survival analysis was performed using R software

version 4.2.1. *Kaplan-Meier analysis*. To test the effect of NK cell and CD8 T cell abundances on

survival, tumors from patients in Cohort 1 were stratified at the median normalized cell counts

(cells/mm^2) averaged between CT and PT regions (CT NK cells + PT NK cells) / 2). The effect of

MHC-I expression on overall survival (OS, the time from resection to death by any cause) and

disease-free survival (DFS, the time from surgical resection to disease recurrence or death by

any cause) was determined by comparing patients with percent MHC-I expression above and

below the median value averaged between CT and PT. Significance was determined by a log-

rank test ($p < 0.05$). Tumors with missing data for any of these markers were excluded from those

respective analyses. *Cox proportional hazards modeling*. Proportional hazards assumptions were

tested using Schoenfeld residual analysis for univariate and multivariate analysis using a signifi-

cance level of 0.01. Multivariate Cox proportional hazards models assessed the relationship be-

tween survival endpoints (OS and DFS) and NK cell, CD8 T cell abundances, and MHC-I

expression, after adjusting for age and sex. Associations were considered significant for two-

sided *p* values ≤ 0.05.

Cellular neighborhood analysis. *Custom cell-cell neighborhood scoring algorithm*. Intercellular

geospatial colocalizations in NSCLC tumors from Cohort 2 were determined in Python using sin-

gle cell 2-dimensional coordinates obtained from HALO. The Euclidian distance between each

cell and every other cell on the slide was computed; nearest neighbors were defined as cells with

a center-to-center Euclidian distance of less than 30 µm from the center cell (**supplemental fig-**

ure S2). The nearest neighbors of each phenotype were enumerated to yield the neighborhood

profile for every individual cell. *Intercellular K function analyses.* Intercellular geospatial colocali-

zations in NSCLC tumors from Cohort 2 were further analyzed using the Spatstat package in R

version 4.2.1 (37). The K function was used to determine the number of nearest neighbors for a

given cell type as a function of radius up to 200 µm for each tumor region separately.

Multivariate discriminant analyses. Orthogonalized Partial Least Squares Discriminant Anal-

yses (OPLSDA) two-component (latent variable) models were generated in MATLAB (Release

R2022a, MathWorks, Natick, MA) using scripts developed in-house.

All data input to OPLSDA were log-transformed, centered, and scaled. First, OPLSDA

models were constructed on cell densities per region (cells/mm^2) to discriminate between tumor

regions from Cohort 2 with panCK+MHC-I+ cell counts above or below the median for AdenoCA

and SCC tumors separately. Cells localized to the tumor nest were normalized to tumor classified

area as determined by HALO software, and likewise for stromal cells. Second, OPLSDA models

were constructed to discriminate the cellular neighborhood profiles between activated and inacti-

vated NK cells (CD3$^-$ CD56$^+$ IFNγ±) and CD8 T cells (CD3$^+$ CD8$^+$ IFNγ±). Prediction accuracy

was determined using a random 5-fold cross validation framework. Significance was determined

empirically by comparing the constructed model's mean squared error against one thousand ran-

domly permuted null models (38, 39). Univariate comparisons between model features were con-

ducted using a non-parametric Wilcoxon rank sum test with a Benjamini-Hochberg false discovery

rate correction controlled at $p < 0.05$.

RESULTS

Patient characteristics. Specimens used to construct the TMA were collected from patients who

were treated at the University of Virginia between 2011 and 2014 and were analyzed by single-

stain immunohistochemistry (IHC) along with retrospective review of clinical records (IRB-HSR#:

18346) (Cohort 1, n = 151) (**table 1**). Whole tumor resections were collected prospectively from

patients treated at the University of Virginia between 2014 and 2018 and were analyzed by mul-

tiplexed immunofluorescence (mIF) (Cohort 2, n = 36). Two patients in Cohort 2 were treated with

Pembrolizumab, one was treated with Atezolizumab, and one was treated with Nivolumab. No

patients in Cohort 1 received immunotherapy.

Table 1. NSCLC patient characteristics.

Clinical feature *		Cohort 1 (n = 151) (Retrospective)			Cohort 2 (n = 36) (Prospective)			Total (Cohort 1 + Cohort 2)
		AdenoCA (n=93)	SCCa (n=58)	Total Cohort 1	AdenoCA (n=28)	SCCa (n=8)	Total Cohort 2	
Biologic sex	Female	49 (53%)	21 (36%)	70 (46%)	20 (71%)	5 (63%)	25 (69%)	95 (51%)
Age in years, mean +/- (range)		67 ± 9 (49-89)	66 ± 9 (38-84)	67 ± 9 (38-89)	66 ± 11 (47-85)	62 ± 12 (55-81)	65 ± 11 (47-85)	66 ± 9 (38-89)
AJCC v7 Stage at excision	Stage I	57 (61%)	36 (62%)	93 (62%)	7 (25%)	0 (0%)	7 (19%)	100 (53%)
	Stage II	20 (22%)	13 (22%)	33 (22%)	10 (36%)	2 (25%)	12 (33%)	45 (24%)
	Stage III	13 (14%)	8 (14%)	21 (14%)	11 (39%)	5 (63%)	16 (44%)	37 (20%)
	Stage IV	3 (3%)	1 (2%)	4 (3%)	0 (0%)	1 (13%)	1 (3%)	5 (3%)
Largest Tumor Diameter* in cm, Mean +/- SD (range)		2.7 ± 1.8 (0.6-12)	3.7 ± 2.4 (0.9-10)	3.1 ± 2.1 (0.6-12)	3.29 ± 1.58 (1.3-6.3)	3.03 ± 1.58 (1.2-5.5)	3.23 ± 1.56 (1.2-6.3)	3.12 ± 1.99 (0.6-12)
Prior chemotherapy		5 (6%)	1 (2%)	6 (4%)	6 (21%)	3 (38%)	9 (25%)	15 (8%)
Treated with Immunotherapy		0 (0%)	0 (0%)	0 (0%)	4 (14%)	0 (0%)	4 (11%)	4 (2%)
* number (%) unless otherwise specified								

Broad MHC class I expression variation in human NSCLC is associated with variation in intra-tumoral lymphocytes. To examine the MHC-I expression in NSCLC tumors and its relationship with occupation by CD56[+] NK cells and CD8 T cells, we performed IHC on Cohort 1 (**table 1**) staining for anti-MHC-I heavy chain (HC), β2-microglobulin (β2m), CD56, and CD8. Cell counts were enumerated in CT and PT cores (**Methods**). We observed wide-ranging MHC-I HC expression levels in both AdenoCA and SCC, with greater than 90% MHC-I HC loss in 12% of AdenoCA and 22% of SCC tumors (**figure 1A,B**). At the other extreme, 16 AdenoCAs (17%) and 10 SCCs (17%) had MHC-I expression in 100% of cells. We additionally observed heterogeneity in tumor cell MHC-I HC expression within the tumors, as CT and PT cores of the same tumor often displayed distinct MHC-I HC expression patterns (**figure 1A,B light bars vs dark bars**). HC10 reacts with unfolded HLA-A, -B and -C heavy chains (33), so it may not identify HLA Class I molecules folded with β2m; thus, we analyzed the correlation of β2m with HLA I HC expression. HLA HC and β2m staining significantly correlated in both CT and PT regions in adenocarcinoma and SCC (**figure 1C,E**), thus suggesting that loss of HC10 staining was primarily due to decreased surface expression of folded MHC-I dimers. Nonetheless, broad variability in MHC-I HC for each β2m classification suggested other regulators of MHC-I expression and/or antigen presentation may have affected MHC-I loss in Cohort 1. MHC-I loss thus is heterogeneously manifest across NSCLC patient tumors comprising both AdenoCA and SCC, and these may develop or progress differently in defined TME regions within individual patients.

Given its significant loss in NSCLC, we asked whether tumor cell MHC-I expression may be associated with NK cell and CD8 T cell infiltrates into the TME. In pairwise Spearman correlations of Cohort 1 TMA features, we found that the density of CD56[+] and CD8[+] cells significantly correlated with expression levels of both MHC-I and β2m in both types of NSCLC (**figure 1D,F**). Moreover, intra-tumoral CD56[+] cells correlated with CD8[+] cells, most profoundly in PT regions, suggesting both cell types may interact and play a role in NSCLC tumor control.

Concurrent tumor-infiltrating CD8$^+$ and CD56$^+$ immune cells are associated with patient survival in human NSCLC independent of MHC-I expression. Prior work has shown that the density of tumor-infiltrating CD8 T cells is generally associated with better patient survival (6, 7). Given the effect of tumor cell MHC-I expression on CD8 T cell and NK cell recognition, we asked if MHC-I expression was associated with patient outcome. We did not detect any significant association between and DFS or OS and MHC-I expressed above or below the median in CT, PT, or averaged across CT and PT (**figure 2A,B, supplemental figure S3**). We thus investigated whether densities of intra-tumoral CD8$^+$ or CD56$^+$ cells were associated with DFS or OS outcomes for Cohort 1 patients (**figure 2C-F**). Intriguingly, high intra-tumoral CD56$^+$ cell counts averaged across CT and PT regions were positively associated with DFS, but a similar association was not significant for CD8$^+$ cell counts (**figure 2C,E**), consistent with our prior work showing that CD8 density across tumor, or in the invasive margins (ie, PT regions) was not correlated with patient survival (10). Both CD8$^+$ and CD56$^+$ cells averaged between CT and PT independently corresponded with OS (**figure 2D,F**). Similar trends were observed when considering CT and PT separately to varying degrees of statistical significance (**supplemental figures S4-S5**). Further assessment revealed that simultaneous CD8$^+$ and CD56$^+$ counts above their medians were more strongly associated with prolonged OS (**figure 2H**). Their joint impact on DFS was less striking, but evident when considering density of CD8$^+$ and CD56$^+$ cells in CT only (**figure 2G, supplemental figure S6**). Observations from Kaplan-Meier analyses were consistent when adjusting for patient age and sex in a Cox proportional hazards model, and statistical power was insufficient to adjust for cancer stage and prior chemotherapy treatment (**supplemental table 1**). Collectively, these data demonstrate that combined presence of NK cells and CD8 T cells predicts better OS, pointing towards potential complementary antitumor activities.

mIF imaging reveals the extent of MHC class I expression variation in patient resected tumor tissues. To further explore the relationship between lymphocyte infiltration and activation and tumor cell MHC-I expression considering its vast heterogeneity in NSCLC and the increased prognostic power observed with simultaneous infiltration of NK cells and CD8 T cells, we implemented mIF imaging with combined spatial analysis of whole tumor sections.

Like the TMA analysis, we observed broadly variable MHC-I expression by mIF as shown in representative SCC and AdenoCA images (**figure 3A**). Because our prior work demonstrated that CD8 T cell infiltrations in NSCLC are highly heterogeneous across tumor regions (10), we developed a tumor regional approach to delineate spatial relationships between NK cells, T cells and tumor cell MHC-I expression in 3 mm^2 regions by tumor type (**supplemental figure S1, and see Methods**). Cohort-wide MHC-I expression per region revealed total MHC-I loss in 26% of regions and >90% loss in 47% of regions, with similar distributions observed in AdenoCA and SCC (**figure 3B**). Wide-ranging inter- and intra-patient MHC-I expression variability was observed with the mean expression per patient ranging from 0.1%-81% of panCK+ cells, and at least one region per tumor displaying >40% MHC-I loss (**figure 3C**). Tumor cell MHC-I loss was thus common among NSCLC patients and heterogeneous across regions of tumor resections.

Tumor cell MHC-I deficiency is associated with decreased NK and T cell presence and activity in the NSCLC TME. We next investigated how NK cell or CD8 T cell infiltration and activation state corresponded to tumor cell MHC-I variability by building OPLSDA models to compare the presence of T cells and NK cells in MHC-I-disparate tumor regions in AdenoCA (**figure 4A-D**) or SCC (**figure 4E-H**) tumors. Lymphocyte activation was inferred by IFNγ staining; though IFNγ is a secreted cytokine and as such its source cannot be definitively determined by mIF, we observed IFNγ most frequently in the presence of NK cells and T cells, indicative of activation (**supplemental figure S7, figure 5**). Three tumors displayed strong IFNγ staining in tumor nests

lacking substantial lymphocyte infiltration, possibly indicative of a non-lymphocyte source of IFNγ

in these tumors.

Our resulting OPLSDA models were able to discriminate between tumor regions with high

and low MHC-I expression based on lymphocyte infiltrates with 79% (AdenoCA) or 87% (SCC)

accuracy, revealing distinct immune signatures of MHC-I high or low tumors (**figure 4A,E**). Vari-

able importance in projection (VIP) scores from the AdenoCA model revealed that stromal CD3$^-$

CD56$^+$ NK cells and as well as both stroma- and tumor-associated CD56$^+$CD3$^+$ cells were en-

riched in tumor regions with tumor MHC-I expression below the median, whereas tumor-associ-

ated CD3$^+$CD8$^-$ (putative CD4+) and CD3$^+$CD8$^+$ T cells were enriched in regions with MHC-I ex-

pression above the median (VIP > 1) (**figure 4B**). Tumor regions with high MHC-I expression also

displayed a greater density of IFNγ$^+$ CD8 T cells and NK cells (**figure 4D**), and Spearman corre-

lation analysis revealed that tumoral IFNγ$^+$ CD8 T cell densities were positively correlated with

IFNγ$^+$ NK cells in the stroma and tumor nest (**figure 4C,G, blue, bolded text**).

VIP scores from the SCC model indicated that CD3$^+$CD8$^-$ and CD3$^+$CD8$^+$ T cells, CD3$^-$

CD56$^+$ NK cells, and CD56$^+$CD3$^+$ cells were among the top features contributing to the separation

between MHC-I replete and deficient tumors (VIP > 1) (**figure 4F,H**). Akin with the adenocarcino-

mas, the only lymphocyte population found enriched in MHC-I low SCC tumor regions were IFNγ$^-$

stromal NK cells (**figure 4H**), indicating that inactive NK cells are abundant in the stroma of tumors

with MHC-I loss for both AdenoCA and SCC subtypes. We observed a clear lack of immune

infiltration into tumor nests with MHC-I loss in SCC tumors (**supplemental figure S8**). Addition-

ally, we observed significant negative pairwise correlations between MHC-I$^-$ tumor cell counts

and IFNγ$^+$ NK cells, CD56$^+$CD3$^+$ cells, CD3$^+$CD8$^+$ T cells, and CD3$^+$CD8$^-$ T cells, whereas stromal

IFNγ$^-$ NK cells were the only immune population positively correlated with MHC-I$^-$ cell densities

(**figure 4G**). This uniquely positive correlation further distinguishes the presence of NK cells in

the stroma of SCC tumors with substantial MHC-I downregulation. Together, these data indicate

that functional, tumor-infiltrating NK cells and T cells marked by IFNγ expression were best asso-

ciated with and coinhabited tumor regions expressing high MHC-I (**figure 4I**).

Local neighborhood analysis of IFNγ$^+$ NK cells and CD8 T cells in the NSCLC TME reveals

tightly clustered lymphocytes marked by IFNγ staining near MHC-I± tumor cells. Given the

observed differences in lymphocyte presence in MHC-I-disparate NSCLC tumors, we questioned

whether IFNγ$^+$ NK cells and CD8 T cells differ from IFNγ$^-$ lymphocytes with respect to their prox-

imity to other immune cells and MHC-I expressing tumor cells. To uncover potential functional

differences in cell-cell communication between activated NK cells and CD8 T cells, we determined

the cellular neighborhood profile of these effector cell types using spatial, cell-centric analyses of

the mIF images (**supplemental figure S2**). The cellular neighborhood profile scores of both tu-

mor- and stroma-associated cells were input into an OPLSDA model to delineate colocalization

differences between IFNγ± NK cells (**figure 6A-C**) and IFNγ± CD8 T cells (**figure 6E-G**). Both

models accurately distinguished between IFNγ$^+$ and IFNγ$^-$ lymphocytes based on their neighbor-

hood profiles with cross-validation accuracy scores of 96% and 97%, respectively (**figure 6A,E**).

Comparing the two models, we observed that both IFNγ$^+$ NK cells and CD8 T cells associated

with all IFNγ$^+$ lymphocyte subsets more so than IFNγ$^-$ cells (**figure 6B,F**). In agreement with our

regional analyses (**figure 4**), we found that NK cells and CD8 T cells were more likely to be IFNγ$^+$

in the tumor as opposed to the surrounding stroma (**figure 6D,H**). Moreover, we observed that

IFNγ$^+$ NK cells and IFNγ$^+$ CD8 T cells in the tumor nests colocalized with MHC-I$^+$ tumor cells more

frequently than did IFNγ$^-$ lymphocytes, suggesting preferential localization of activated cells to

tumors with MHC-I expression (**figure 6D,H**). Strikingly, IFNγ$^+$ NK cells and CD8 T cells were

more frequently associated with MHC-I$^-$ tumor cells compared to their IFNγ$^-$ counterparts, indi-

cating infiltration of IFNγ-secreting lymphocytes into both MHC-I$^+$ and MHC-I$^-$ tumor nests (**figure**

6I). To validate these observations over larger radii (<200 μm), we quantified the number of r-

neighbors—i.e., the number of neighbors in radius r around a center cell (37). We observed IFNγ$^+$

CD8 T cells had fewer MHC-I$^-$ neighbors than MHC-I$^+$ neighbors, and their association with MHC-

I$^-$ tumor cells was less than what was expected by random chance (**figure 6J**). This may signify

deliberate enrichment of IFNγ$^+$ CD8 T cells in MHC-I-bearing tumors. In contrast, IFNγ$^+$ NK cells

associated similarly with MHC-I$^+$ and MHC-I$^-$ tumor cells, suggesting that NK cells may become

or remain activated in the MHC-I-deficient TME. Marked colocalization of IFNγ$^+$ NK cells with other

IFNγ$^+$ lymphocytes including NK cells, CD8$^-$ T cells, and CD8$^+$ T cells additionally suggests that

lymphocytes may be activated en masse in MHC-I-bearing tumor regions that can trigger NK cell

activation. IFNγ produced in these clusters of activated lymphocytes thus may be indicative of

type I immune responses in the TME.

DISCUSSION

While targeting immune checkpoints with CPI has had a substantive impact on the treatment of

NSCLC, a minority of patients display durable responses. Whether this is due to hindered immune

infiltration, functional exhaustion, tumor cell evasion of immune recognition by loss of MHC-I ex-

pression, or alternative mechanisms remains to be determined. Understanding the basis of this

selectivity will increase the appropriate targeting of CPI or combination therapies to patients most

likely to benefit and will ultimately lead to development of innovative approaches to expand the

proportion of patients who receive clinical benefit.

We hypothesized that coordinated activities of both CD8 T cells and NK cells may jointly

enhance targeted lysis of NSCLC tumor cells while preventing tumor cell MHC-I loss. We ana-

lyzed resected patient (non-CPI treated) tumor tissues by both traditional IHC and mIF to simul-

taneously probe NK cells and CD8 T cells *in situ*, their effector activity indicated by IFNγ staining,

and their co-localization in MHC-I disparate NSCLC tumors. To define differences in lymphocyte

activation and intercellular communication in tumors with and without MHC-I expression, we

leveraged the inherent heterogeneity of the TME by developing a suite of spatial analysis methods

spanning whole tumors, tumor sub-regions, and single-cell enumeration of NK cell and T cell

neighborhoods. This use of integrative systems consistently pointed towards coordinated anti-

tumor activity of NK cells and T cells, particularly in tumors with MHC-I expression.

Consistent with prior work (23, 25, 28, 30), both cohorts displayed stark inter-patient and

intra-tumoral heterogeneity in MHC-I expression. Although MHC-I loss can lead to the develop-

ment of CD8[+] T cell refractory tumors, we did not observe an association between tumor cell

MHC-I expression and NSCLC patient survival in this study. Others have speculated cohort-spe-

cific composition, clinicopathologic characteristics, genetic contribution and/or survival/events

may obfuscate any potential association (28-30). MHC-I loss thus may indirectly impact patient

survival by regulating lymphocyte occupancy and activation.

IHC analysis of Cohort 1 revealed that tumors with high counts of both CD8[+] and CD56[+]

cells were more strongly associated with OS compared to tumors with high CD56[+] or CD8[+] cells

only. Using mIF to concurrently enumerate NK cell and CD8 T cell infiltrates in Cohort 2, we

observed positively correlated tumor-infiltrating NK cells and CD8 T cells with greater densities in

tumor nests expressing high levels of MHC-I, indicating coordinated infiltration. Furthermore, spa-

tial analysis of local cellular neighborhoods revealed infiltrated NK and CD8 T cells are present in

clusters marked by IFNγ activity together with CD8[−] T cells. Though much less frequent, similar

immune clusters were observed in tumors with MHC-I loss, suggesting that an effector response

involving NK cells, CD8 T cells, and CD8[−] T cells can be active, and possibly leveraged, even in

a setting considered refractory to lymphocyte immune control.

Whether intra-tumoral IFNγ promotes tumor cell MHC-I expression, immune recruitment

and co-localization, or coordinate lymphocyte effector activity awaits further analysis, but these

findings suggest that CPI plus novel approaches designed to jointly elicit NK and T cell effector

activities may be key to developing next generation immunotherapies. Antigen-stimulated NK

cells have been shown to provide support to CD8 T cell antitumor immunity in a mouse model of

lung adenocarcinoma (14). Likewise, human induced pluripotent stem cell (iPSC-)derived NK

cells have been shown to enhance T cell recruitment and activation to tumors, coinciding with

improved T cell sensitivity to CPI and enhanced antitumor control (40). Cytokine activation can

additionally invigorate NK cells to curb or regress tumor outgrowth, even in MHC-I⁻ tumors (17).

The potential benefit of NK cells in NSCLC is bolstered by prior work that showed that human NK

cell cytolytic activity is associated with lower cancer risk (41), whereas NK cell deficiency is asso-

ciated with increased cancer risk (42-44). Our data contrast recent observations in soft tissue

sarcoma that CD56⁺ NK cell infiltrates indicate a poor prognosis and do not colocalize with CD8

T cells or MHC-I⁺ tumor cells, indicating that the role of NK cells suggested by our results may be

context specific (45). This apparent synergy between NK cells and cytotoxic T cells in the NSCLC

TME highlights a vital clinical value in developing CPI treatments which stimulate both lymphocyte

populations.

Our findings that NK cells had infiltrated into tumor nests and displayed evidence of IFNγ

activity in NSCLC begs the question, why might tumoral NK cells become ineffective drivers of

anti-tumor immunity? Emerging evidence suggests that NK cell activity may be hindered in the

lung cancer TME. Human NKp46⁺ NK cells were shown to infiltrate the invasive margin of lung

tissue stroma surrounding tumor nests in NSCLC patients with no apparent effect on clinical out-

comes (33). These intra-tumoral NK cells displayed inferior effector activities in comparison to

their counterpart NK cells in peripheral blood (33). Whether due to suppressive mechanisms as-

sociated with the lung environment itself (46) or more direct suppression via tumor cells remains

unresolved as cancer cells can downregulate or secrete NK receptor ligands to evade natural

killing, and the TME is rife with suppressive factors which can impede NK cell activity (47). Prior

work in mouse models has further shown NK cells also undergo disarming in MHC-I⁻ tumors

rendering them hyporesponsive and unable to mediate tumor cell clearance (17, 48). As our study

affirms the significant association with survival and notable infiltration into MHC-I$^+$ tumor nests of

NK cells in NSCLC, future studies are needed to uncover the root causes of NK cell disarming in

the lung cancer TME to fully realize their potential as anti-tumor effectors.

Though these multifaceted spatial analyses consistently revealed NK cell and CD8 T cell

co-occurrence within NSCLC tumors, there are several limitations to consider. The TMA data from

Cohort 1 were manually generated by a pathologist and consequently may be impacted by sub-

jective bias; however, this concern is mitigated as automated analysis yielded similar results. mIF

imaging was limited to six phenotypic markers plus nuclear staining, thus constraining analysis of

the full complexity of the tumor-associated immune milieu. Consequently, we relied on IFNγ ex-

pression as a marker of lymphocyte activity and were unable to test for expression of other cyto-

toxic markers and inhibitory ligands which may obscure true activity levels of cytotoxic T cells and

NK cells. Ongoing work with higher dimensional imaging (e.g., spatial transcriptomics) will provide

a more comprehensive view of immune cell diversity, occupancy, proliferation, and activation in

the TME. As IFNγ is a secreted cytokine, we were unable to definitively determine its source.

Nevertheless, analysis of our data without three select tumors that displayed strong IFNγ staining

in the absence of lymphocytes yielded similar results and thus non-lymphocyte IFNγ was not a

major confounder.

Another important limitation of our study is that the cohort was composed of only complete

resections and thus there was no opportunity assess tissue after chemotherapy or immunother-

apy treatment. For instance, it remains to be determined whether NK cells drive treatment-related

recruitment and/or activation of CD8 T cells in the TME, or vice versa. Likewise, whether stark

differences in lymphocyte populations and activity in tumors with loss of MHC-I expression results

from differences in lymphocyte infiltration, expansion, or retention in the TME remains unclear.

Other outstanding questions include whether patient tumors are infiltrated by circulating NK cells

with higher baseline cytotoxic activity, by lung resident NK cells with lower baseline cytotoxic

activity, or both. Whether these different types of NK cells could affect other tumor-infiltrating

immune cells, such as helper T cells, and whether there are specific interactions that either cause

or arise from the spatial associations observed in the data remains to be determined. Future

mechanistic studies in animal models will help to delineate the effect of MHC-I on the balance of

NK cells and their impact on the tumor-immune landscape.

Notwithstanding these challenges, we have shown that tumor cell MHC-I loss is common

in NSCLC patient tissues across two separate cohorts, an effect that negatively impacted lym-

phocyte occupancy and activity in the TME. While we did not find MHC-I to be predictive of patient

survival, CD8+ and CD56+ cells were independently associated with survival and had an in-

creased effect in tumors with simultaneously high CD8+ and CD56+ counts. This evidence of

lymphocyte activation and intercellular communication in the presence of MHC-I$^+$ NSCLC tumors

highlights both the vital importance and promise for developing novel immunotherapies that lev-

erage both NK cells and CD8 T cells to enhance immunologic control of NSCLC.

REFERENCES

1.   Molina JR, Yang P, Cassivi SD, Schild SE, Adjei AA. Non-small cell lung cancer:
epidemiology, risk factors, treatment, and survivorship. Mayo Clinic proceedings.
2008;83(5):584-94.
2.   Siegel RL, Miller KD, Fuchs HE, Jemal A. Cancer statistics, 2022. CA: a cancer
journal for clinicians. 2022;72(1):7-33.
3.   Chen Z, Fillmore CM, Hammerman PS, Kim CF, Wong KK. Non-small-cell lung
cancers: a heterogeneous set of diseases. Nat Rev Cancer. 2014;14(8):535-46.
4.   Travis WD, Brambilla E, Noguchi M, Nicholson AG, Geisinger K, Yatabe Y, et al.
International Association for the Study of Lung Cancer/American Thoracic
Society/European Respiratory Society: international multidisciplinary classification of lung
adenocarcinoma: executive summary. Proc Am Thorac Soc. 2011;8(5):381-5.
5.   Stankovic B, Bjorhovde HAK, Skarshaug R, Aamodt H, Frafjord A, Muller E, et al.
Immune Cell Composition in Human Non-small Cell Lung Cancer. Frontiers in
immunology. 2018;9:3101.
6.   Geng Y, Shao Y, He W, Hu W, Xu Y, Chen J, et al. Prognostic Role of Tumor-
Infiltrating Lymphocytes in Lung Cancer: a Meta-Analysis. Cellular physiology and
biochemistry : international journal of experimental cellular physiology, biochemistry, and
pharmacology. 2015;37(4):1560-71.

7.   Zeng DQ, Yu YF, Ou QY, Li XY, Zhong RZ, Xie CM, et al. Prognostic and predictive
value of tumor-infiltrating lymphocytes for clinical therapeutic research in patients with
non-small cell lung cancer. Oncotarget. 2016;7(12):13765-81.
8.   Schalper KA, Brown J, Carvajal-Hausdorf D, McLaughlin J, Velcheti V, Syrigos KN,
et al. Objective measurement and clinical significance of TILs in non-small cell lung
cancer. J Natl Cancer Inst. 2015;107(3).
9.   Esendagli G, Bruderek K, Goldmann T, Busche A, Branscheid D, Vollmer E, et al.
Malignant and non-malignant lung tissue areas are differentially populated by natural killer
cells and regulatory T cells in non-small cell lung cancer. Lung cancer. 2008;59(1):32-40.
10.   Obeid JM, Wages NA, Hu Y, Deacon DH, Slingluff CL, Jr. Heterogeneity of CD8(+)
tumor-infiltrating lymphocytes in non-small-cell lung cancer: impact on patient prognostic
assessments and comparison of quantification by different sampling strategies. Cancer
immunology, immunotherapy : CII. 2017;66(1):33-43.
11.   Federico L, McGrail DJ, Bentebibel SE, Haymaker C, Ravelli A, Forget MA, et al.
Distinct tumor-infiltrating lymphocyte landscapes are associated with clinical outcomes in
localized non-small-cell lung cancer. Annals of oncology : official journal of the European
Society for Medical Oncology. 2022;33(1):42-56.
12.   Guillerey C, Huntington ND, Smyth MJ. Targeting natural killer cells in cancer
immunotherapy. Nat Immunol. 2016;17(9):1025-36.
13.   Malmberg KJ, Carlsten M, Bjorklund A, Sohlberg E, Bryceson YT, Ljunggren HG.
Natural killer cell-mediated immunosurveillance of human cancer. Semin Immunol.
2017;31:20-9.
14.   Schmidt L, Eskiocak B, Kohn R, Dang C, Joshi NS, DuPage M, et al. Enhanced
adaptive immune responses in lung adenocarcinoma through natural killer cell
stimulation. Proc Natl Acad Sci U S A. 2019;116(35):17460-9.
15.   Liu RB, Engels B, Arina A, Schreiber K, Hyjek E, Schietinger A, et al. Densely
granulated murine NK cells eradicate large solid tumors. Cancer research.
2012;72(8):1964-74.
16.   Nicolai CJ, Wolf N, Chang IC, Kirn G, Marcus A, Ndubaku CO, et al. NK cells
mediate clearance of CD8(+) T cell-resistant tumors in response to STING agonists.
Science immunology. 2020;5(45).
17.   Ardolino M, Azimi CS, Iannello A, Trevino TN, Horan L, Zhang L, et al. Cytokine
therapy reverses NK cell anergy in MHC-deficient tumors. J Clin Invest.
2014;124(11):4781-94.
18.   Bottcher JP, Bonavita E, Chakravarty P, Blees H, Cabeza-Cabrerizo M,
Sammicheli S, et al. NK Cells Stimulate Recruitment of cDC1 into the Tumor
Microenvironment Promoting Cancer Immune Control. Cell. 2018;172(5):1022-37 e14.
19.   Kyrysyuk O, Wucherpfennig KW. Designing Cancer Immunotherapies That
Engage T Cells and NK Cells. Annu Rev Immunol. 2023;41:17-38.
20.   Topalian SL, Hodi FS, Brahmer JR, Gettinger SN, Smith DC, McDermott DF, et al.
Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. The New England
journal of medicine. 2012;366(26):2443-54.
21.   Brahmer JR, Tykodi SS, Chow LQ, Hwu WJ, Topalian SL, Hwu P, et al. Safety and
activity of anti-PD-L1 antibody in patients with advanced cancer. The New England
journal of medicine. 2012;366(26):2455-65.

22.   Santarpia M, Giovannetti E, Rolfo C, Karachaliou N, Gonzalez-Cao M, Altavilla G,
et al. Recent developments in the use of immunotherapy in non-small cell lung cancer.
Expert Rev Respir Med. 2016;10(7):781-98.
23.   Garrido F, Aptsiauri N, Doorduijn EM, Garcia Lora AM, van Hall T. The urgent need
to recover MHC class I in cancers for effective immunotherapy. Current opinion in
immunology. 2016;39:44-51.
24.   Thunnissen E, van der Oord K, den Bakker M. Prognostic and predictive
biomarkers in lung cancer. A review. Virchows Arch. 2014;464(3):347-58.
25.   Redondo M, Ruiz-Cabello F, Concha A, Cabrera T, Perez-Ayala M, Oliva MR, et
al. Altered HLA class I expression in non-small cell lung cancer is independent of c-myc
activation. Cancer research. 1991;51(9):2463-8.
26.   Korkolopoulou P, Kaklamanis L, Pezzella F, Harris AL, Gatter KC. Loss of antigen-
presenting molecules (MHC class I and TAP-1) in lung cancer. British journal of cancer.
1996;73(2):148-53.
27.   Delp K, Momburg F, Hilmes C, Huber C, Seliger B. Functional deficiencies of
components of the MHC class I antigen pathway in human tumors of epithelial origin.
Bone Marrow Transplant. 2000;25 Suppl 2:S88-95.
28.   Ramnath N, Tan D, Li Q, Hylander BL, Bogner P, Ryes L, et al. Is downregulation
of MHC class I antigen expression in human non-small cell lung cancer associated with
prolonged survival? Cancer immunology, immunotherapy : CII. 2006;55(8):891-9.
29.   Li X, Guo F, Liu Y, Chen HJ, Wen F, Zou B, et al. NLRC5 expression in tumors
and its role as a negative prognostic indicator in stage III non-small-cell lung cancer
patients. Oncol Lett. 2015;10(3):1533-40.
30.   Datar IJ, Hauc SC, Desai S, Gianino N, Henick B, Liu Y, et al. Spatial Analysis and
Clinical Significance of HLA Class-I and Class-II Subunit Expression in Non-Small Cell
Lung Cancer. Clinical cancer research : an official journal of the American Association for
Cancer Research. 2021;27(10):2837-47.
31.   Kikuchi E, Yamazaki K, Torigoe T, Cho Y, Miyamoto M, Oizumi S, et al. HLA class
I antigen expression is associated with a favorable prognosis in early stage non-small cell
lung cancer. Cancer Sci. 2007;98(9):1424-30.
32.   Vivier E, Raulet DH, Moretta A, Caligiuri MA, Zitvogel L, Lanier LL, et al. Innate or
adaptive immunity? The example of natural killer cells. Science. 2011;331(6013):44-9.
33.   Platonova S, Cherfils-Vicini J, Damotte D, Crozet L, Vieillard V, Validire P, et al.
Profound coordinated alterations of intratumoral NK cell phenotype and function in lung
carcinoma. Cancer research. 2011;71(16):5412-22.
34.   Erdag G, Schaefer JT, Smolkin ME, Deacon DH, Shea SM, Dengel LT, et al.
Immunotype and immunohistologic characteristics of tumor-infiltrating immune cells are
associated with clinical outcome in metastatic melanoma. Cancer research.
2012;72(5):1070-80.
35.   Stam NJ, Spits H, Ploegh HL. Monoclonal antibodies raised against denatured
HLA-B locus heavy chains permit biochemical characterization of certain HLA-C locus
products. J Immunol. 1986;137(7):2299-306.
36.   Mauldin IS, Mahmutovic A, Young SJ, Slingluff CL, Jr. Multiplex
Immunofluorescence Histology for Immune Cell Infiltrates in Melanoma-Associated
Tertiary Lymphoid Structures. Methods Mol Biol. 2021;2265:573-87.

37.   Baddeley A, Turner R. spatstat: An R Package for Analyzing Spatial Point
Patterns. . Journal of Statistical Software. 2005;12:1-42.
38.   Dolatshahi S, Butler AL, Siedner MJ, Ngonzi J, Edlow AG, Adong J, et al. Altered
Maternal Antibody Profiles in Women With Human Immunodeficiency Virus Drive
Changes in Transplacental Antibody Transfer. Clinical infectious diseases : an official
publication of the Infectious Diseases Society of America. 2022;75(8):1359-69.
39.   Grace PS, Dolatshahi S, Lu LL, Cain A, Palmieri F, Petrone L, et al. Antibody
Subclass and Glycosylation Shift Following Effective TB Treatment. Frontiers in
immunology. 2021;12:679973.
40.   Cichocki F, Bjordahl R, Gaidarova S, Mahmood S, Abujarour R, Wang H, et al.
iPSC-derived NK cells maintain high cytotoxicity and enhance in vivo tumor control in
concert with T cells and anti-PD-1 therapy. Science translational medicine. 2020;12(568).
41.   Imai K, Matsuyama S, Miyake S, Suga K, Nakachi K. Natural cytotoxic activity of
peripheral-blood lymphocytes and cancer incidence: an 11-year follow-up study of a
general population. Lancet. 2000;356(9244):1795-9.
42.   Orange JS. Natural killer cell deficiency. The Journal of allergy and clinical
immunology. 2013;132(3):515-25.
43.   Moon WY, Powis SJ. Does Natural Killer Cell Deficiency (NKD) Increase the Risk
of Cancer? NKD May Increase the Risk of Some Virus Induced Cancer. Frontiers in
immunology. 2019;10:1703.
44.   Mace EM, Orange JS. Emerging insights into human health and NK cell biology
from the study of NK cell deficiencies. Immunol Rev. 2019;287(1):202-25.
45.   Cruz SM, Sholevar CJ, Judge SJ, Darrow MA, Iranpur KR, Farley LE, et al.
Intratumoral NKp46(+) natural killer cells are spatially distanced from T and MHC-I(+)
cells with prognostic implications in soft tissue sarcoma. Frontiers in immunology.
2023;14:1230534.
46.   Marquardt N, Kekalainen E, Chen P, Kvedaraite E, Wilson JN, Ivarsson MA, et al.
Human lung natural killer cells are predominantly comprised of highly differentiated
hypofunctional CD69(-)CD56(dim) cells. The Journal of allergy and clinical immunology.
2017;139(4):1321-30 e4.
47.   Granzin M, Wagner J, Kohl U, Cerwenka A, Huppert V, Ullrich E. Shaping of
Natural Killer Cell Antitumor Activity by Ex Vivo Cultivation. Frontiers in immunology.
2017;8:458.
48.   Marcus A, Mao AJ, Lensink-Vasan M, Wang L, Vance RE, Raulet DH. Tumor-
Derived cGAMP Triggers a STING-Mediated Interferon Response in Non-tumor Cells to
Activate the NK Cell Response. Immunity. 2018;49(4):754-63 e4.
687

Figure 1. MHC class I expression in NSCLC exhibits intra- and inter-tumor heterogeneity and correlates with intra-tumoral lymphocytes. (A & B) MHC-I is variably expressed in tumors and across AdenoCA **(A)** and SCC **(B)** patients from Cohort 1. Bar plots represent the percent of tumoral MHC-I⁺ cells per patient in the central tumor (CT) and peripheral tumor (PT). OS (overall survival) and DFS (disease-free survival) are shown for each patient below the bar graphs (black, survived; gray, did not survive; white, data not available). **(C & E)** β2M expression (0, low; 1, moderate; 2, high) in PT (left) and CT (right) correlates positively with MHC-I expression in AdenoCA **(C)** and SCC **(E)** patients from Cohort 1. R, Spearman correlation coefficient. **(D & F)** The correlation heatmaps show Spearman correlations between all features from Cohort 1 in AdenoCA **(D)** and SCC **(F)** patients. Circle color is proportional to correlation strength, and circle radius is proportional to significance. Insignificant correlations (p > 0.05) are indicated by white shading, and identity correlations are indicated with light grey.

Figure 2. Probability of survival increases with increasing intra-tumoral CD8⁺ and CD56⁺ lympho-cytes. Kaplan-Meier analysis predicts the probability of DFS or OS in Cohort 1 patients with respect to intra-tumoral MHC-I expression level **(A & B)**, CD56⁺ cell counts **(C & D)**, CD8⁺ cell counts **(E & F)**, or coincident CD56⁺ and CD8⁺ cell counts **(G & H)** above and below the median averaged between CT and PT regions. Significance was determined by a log-rank test.

Figure 3. mIF imaging reveals extent of MHC class I expression variation in resected patient tumor tissues. (A) The representative images show an example of MHC-I loss in AdenoCA (left) and SCC (right). A representative image classified as MHC-I high or low (stratified at the median count of panCK+MHC-I+ tumor cells) is show for each tumor type. **(B & C)** The histograms show percentage of MHC-I+ tumor cells in each region in adenocarcinoma (top) and SCC (bottom) patients from Cohort 2. Percentages are defined as the number of MHC-I+ PanCK+ cells per total PanCK+ cells. **(C)** The violin plots show the percentage of MHC-I+ tumor cells per region for each patient where each dot represents a region and the white circles represent mean of all regions from a patient. Patients are sorted by average percentage of MHC-I+ cells across their regions. AdenoCA and SCC patients are both shown.

MAD17-205

Figure 4. Tumor infiltrated lymphocytes and IFNγ expression are associated with high tumor cell
MHC-I expression. An OPLSDA model was constructed to discriminate between tumor regions with high
or low MHC-I expression, defined as regions with MHC-I$^+$ tumor cell densities (cells/mm^2 classified tumor)
above or below the cohort median in **(A-D)** AdenoCA and **(E-H)** SCC. A 5-fold cross validation yielded
models with 79% (adenocarcinoma) and 87% (SCC) classification accuracy. Both models outperformed all
of 1000 randomly permuted models ($p < 0.001$). **(A & E)** The scatter plots show the X scores, where each
point represents one tumor region. LV1 & LV2, latent variable 1 and 2. **(B & F)** The bar plots show the
Variable Importance in Projection (VIP) scores, artificially oriented in the direction of loadings on LV1 and
colored according to their association with MHC-I high or MHC-I low regions. |VIP score| > 1 indicates a
variable with greater than average influence on the separation between groups. **(C & G)** The correlation
heatmaps show the Spearman correlations among immunologic features in tumor (T) and stroma (S) re-
gions. Color intensity indicates correlation coefficient, and the circle size is inversely proportional to the p
value. Insignificant correlations ($p > 0.05$) are indicated by white shading, and identity correlations are indi-
cated with light grey. **(D & H)** The scatter plots show univariate comparisons between model features de-
termined by a non-parametric Wilcoxon rank sum test followed by multiple comparison correction using the
Benjamini-Hochberg method (*, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.0001$). Only features with an adjusted $p <$
0.05 are shown. **(I)** Representative image shows variation in lymphocyte occupancy and IFNγ staining in
different regions of the same tumor that do not express (top) or express (bottom) MHC-I.

MAD17-473

MAD17-450

MAD17-205

739 **Figure 5. NK cells and CD8 T cells colocalize in areas with high IFNγ staining.** The mIF images show
740 representative instances of CD56+CD3− NK cells and CD3+CD8+ T cells clustered together with evidence
741 of IFNγ staining. Each row is drawn from a different patient tumor tissue. The box inset highlights instances
742 of NK cell and CD8 T cell colocalization.

MAD16-101

Figure 6. Activated IFNγ+ NK cells and CD8 T cells preferentially associate with other activated
lymphocytes and MHC-I+ tumor cells. An OPLSDA model was constructed to discriminate activated and
inactivated (IFNγ+/-) NK cells **(A-C)** and CD8 T cells **(E-G)**. A 5-fold cross validation yielded models with
96% (NK cells) and 97% (CD8 T cells) classification accuracy. The model outperformed all of 1000 ran-
domly permuted models (p < 0.001). **(A & E)** The scatter plot shows the X scores on latent variables 1 and
2 (LV1 & LV2), where each point represents one cell. **(B & F)** The bar plot shows the Variable Importance
in Projection (VIP) scores, artificially oriented in the direction of loadings on LV1 and colored according to
their association with activated or inactivated NK cells or CD8 T cells. |VIP score| > 1 indicate a variable
with greater than average influence on the projection. **(C & G)** The violin plots show univariate comparisons
between top-scoring features. White circles indicate the median. Significance was determined by a two-
tailed t-test (* p < $1x10^{-27}$; ** p < $1x10^{-30}$; *** p < $1x10^{-50}$). **(D & H)** Percentage of IFNγ+ NK cells **(D)** and
CD8 T cells **(H)** in the tumor and stroma are shown as pie charts. The violin plots show the number of MHC-
I+ tumor cell neighbors of IFNγ± lymphocytes in tumor and stroma separately. **(I)** Representative mIF image
showing a cluster of activated lymphocytes in an MHC-I-deficient tumor nest. **(J)** The K-cross correlation
value is plotted against increasing radii for IFNγ+ NK cells and IFNγ+ CD8 T cells as the target cell and
either MHC-I$^+$ (blue) or MHC-I$^-$ (pink) center cells. 95% confidence intervals are shown in gray, and the
black line represents a Poisson (null) distribution.