Purging viral latency by a bifunctional HSV-vectored therapeutic vaccine in chronically SIV-infected macaques

Ziyu Wen,[1#] Pingchao Li,[2#] Yue Yuan,[1#] Congcong Wang,[1#] Minchao Li,[1] Haohang Wang,[1] Minjuan Shi,[1] Yizi He,[2] Mingting Cui,[1] Ling Chen,[2*] Caijun Sun,[1,3*]

[1]School of Public Health (Shenzhen), Sun Yat-sen University, Guangdong, China; [2]State Key Laboratory of Respiratory Disease, Guangzhou Institutes of Biomedicine and Health (GIBH), Chinese Academy of Sciences, Guangzhou, China; [3]Key Laboratory of Tropical Disease Control (Sun Yat-sen University), Ministry of Education, Guangzhou, China.

#These authors contributed equally to this work.
* For correspondence:
Caijun Sun, suncaijun@mail.sysu.edu.cn (CS);
Ling Chen, chen_ling@gibh.ac.cn (LC)

Abstract

The persistence of latent viral reservoirs remains the major obstacle to eradicating human immunodeficiency virus (HIV). We herein reported that recombinant herpes simplex virus type I (HSV-1) with ICP34.5 deletion could more effectively reactivate HIV latency than its wild-type counterpart. Mechanistically, HSV-ΔICP34.5 promoted the phosphorylation of HSF1 by decreasing the recruitment of protein phosphatase 1 (PP1α), thus effectively binding to the HIV LTR to reactivate the latent reservoirs. In addition, HSV-ΔICP34.5 enhanced the phosphorylation of IKKα/β through the degradation of IκBα, leading to p65 accumulation in the nucleus to elicit NF-κB pathway-dependent reactivation of HIV latency. Then, we constructed the recombinant HSV-ΔICP34.5 expressing simian immunodeficiency virus (SIV) env, gag, or the fusion antigen sPD1-SIVgag as an HIV therapeutic vaccine, aiming to achieve a functional cure by simultaneously reactivating viral latency and eliciting antigen-specific immune responses. Results showed that these constructs effectively elicited SIV-specific immune responses, reactivated SIV latency, and delayed viral rebound after the interruption of antiretroviral therapy (ART) in chronically SIV-infected rhesus macaques. Collectively, these findings provide insights into the rational design of HSV-vectored therapeutic strategies for pursuing an HIV functional cure.

Keywords: HIV, SIV, HSV, therapeutic vaccine, functional cure

Introduction

The epidemic of acquired immunodeficiency syndrome (AIDS), caused by human immunodeficiency virus type I (HIV-1), is still a huge challenge for global public health, with approximately 39 million people living with HIV-1 as of 2022. To date, there is neither a curable drug nor a prophylactic vaccine for clinical use against HIV-1 infections. Antiretroviral therapy (ART) can effectively control HIV-1 replication to an undetectable level, but the termination of ART usually results in prompt viral rebound from latent viral reservoirs (Archin et al., 2014, Looker et al., 2017, Calistri et al., 2003, Heng et al., 1994). Thus, it is of great priority to explore novel strategies for curing HIV latency. The "Shock and Kill" strategy is considered a promising approach for purging HIV-1 reservoirs, involving the activation of latently infected cells to express viral products (Shock), followed by viral cytopathic effects or specific cytolytic T lymphocytes (CTLs) to eliminate the activated cells (Kill) (Wu et al., 2022, Yang et al., 2019, Kim et al., 2018). Numerous latency-reversing agents (LRAs) (Yang et al., 2019, Wu et al., 2022), including methylation inhibitors, histone deacetylase (HADC) inhibitors (Archin et al., 2017, Lehrman et al., 2005), and bromodomain and extra terminal domain (BET) protein inhibitors (Li et al., 2013, Bisgrove et al., 2007), have been identified to reactivate latent HIV-1 in preclinical studies, but there is no ideal LRA available for clinical patients yet.

Herpes simplex virus (HSV), a human herpesvirus, features a 152-kb double-stranded DNA genome encoding over 80 proteins (Poh, 2016). Owing to its distinctive genetic background, high capacity, broad tropism, thermostability, and excellent safety profile, modified HSV variants have found extensive applications

in gene therapy and oncolytic virotherapy. For example, talimogene laherparepvec (T-VEC), an HSV-1 variant with ICP34.5 deletion and GM-CSF insertion, received FDA approval in 2015 for treating malignant melanoma, showcasing notable safety and efficacy in clinical practice. Recombinant HSV-based constructs have also emerged as efficacious gene delivery vectors against infectious diseases. Early studies indicated that prophylactic vaccines based on HSV, expressing simian immunodeficiency virus (SIV) antigens, could elicit robust antigen-specific CTL responses in mice and monkeys, providing enduring and partial protection against pathogenic SIVmac239 challenges (Kaur et al., 2007, Murphy et al., 2000). Moreover, increasing data suggest the crucial role of HIV-specific CTL in controlling viral replication and eliminating potential HIV reservoirs (Collins et al., 2020, Leitman et al., 2017). Significantly, epidemiological research suggests a synergistic effect between HSV and HIV infections, with HSV infection in HIV patients being associated with increased HIV-1 viral load and disease progression (Looker et al., 2017, Calistri et al., 2003, Heng et al., 1994). Some studies have further unveiled the ability of HSV to activate HIV latent reservoirs (Amici et al., 2004, Amici et al., 2001, Pierce et al., 2023). Given the potential of HSV to simultaneously induce antigen-specific immune responses and reactivate latent viral reservoirs, we propose a proof-of-concept strategy to achieve an HIV functional cure using a modified bifunctional HSV-vectored therapeutic vaccine.

Results

The modified HSV-ΔICP34.5-based constructs reactivated HIV latency more efficiently than wild-type HSV counterparts

The J-Lat 10.6 cell line, originating from Jurkat T cells with latent HIV-1 provirus, was infected with wild-type HSV-1 17 strain (HSV-wt) at different multiplicities of infection (MOI) to assess its capability to reactivate HIV latency. Flow cytometry analysis showed that green fluorescent protein (GFP) expression was increased in a dose-dependent manner (*Figure 1A*), and the mRNA levels of HIV-1 LTR, Tat, Gag, Vif, and Vpr were also significantly increased in response to HSV stimulation (*Figure 1B*), demonstrating that the latent HIV provirus can be reactivated to a certain extent by wild-type HSV. In addition, different HSV-1 strains, including HSV-1 Mckrae and HSV-1 F strain, can also reactivate HIV latency (*Figure 1 - figure supplement 1*). Remarkably, we found for the first time that the HSV-1 17 strain with ICP34.5 deletion (HSV-ΔICP34.5) could reactivate HIV latency more efficiently than HSV-wt. Specifically, the mRNA levels of HIV genes (LTR, Tat, Gag, Vpr, Vif) were substantially increased in HSV-ΔICP34.5-infected J-Lat 10.6 cells than in HSV-wt treated cells, although there was a weaker replication ability of HSV-ΔICP34.5 in these cells than that of HSV-wt, as indicated by the mRNA level of HSV UL27 (*Figure 1C and D*). Furthermore, this finding was verified in ACH-2 cells, derived from T cells latently infected with replication-competent HIV-1. A significantly higher level of p24 protein, a key indicator of HIV replication, was found in the HSV-ΔICP34.5-infected ACH-2 cells than in the HSV-wt treated cells

(*Figure 1E*), and the mRNA levels of HIV-1-related genes (LTR, Tat, Gag, Vpr, Vif)

were also significantly increased (*Figure 1F*). Subsequently, we generated J-Lat

10.6 cells stably expressing ICP34.5-Flag-Tag (J-Lat 10.6-ICP34.5) using the

recombinant lentivirus system and confirmed the expression of the ICP34.5 protein

(*Figure 1G*). Our research revealed that HSV-ΔICP34.5 displayed reduced

reversal capacity for the latent HIV reservoir in J-Lat 10.6-ICP34.5 cells compared

to J-Lat 10.6 cells (*Figure 1G*). Furthermore, in J-Lat 10.6-ICP34.5 cells, the

potency of latent reversal agents like phorbol 12-myristate 13-acetate (PMA) and

TNF-α was notably reduced when contrasted with J-Lat 10.6 cells (*Figure 1H*).

These findings indicate that HSV ICP34.5 can effectively inhibit the reactivation of

the HIV latency, and HSV constructs lacking ICP34.5 potentially reactivate HIV

latency with high efficiency.

117 **The modified HSV-based constructs effectively reactivated HIV**

118 **latency by modulating the IKKα/β-NF-κB pathway and PP1-HSF1**

119 **pathway**

Next, the mechanism of potentially reactivating viral latency by the modified HSV-

ΔICP34.5-based constructs was explored. J-Lat10.6 cells were infected with HSV-

wt and HSV-ΔICP34.5, and our results showed that HSV-ΔICP34.5 significantly

enhanced the phosphorylation of IKKα/β, promoted the degradation of IKBα, and

thus led to the accumulation of p65 in the nucleus (*Figure 2A*). NF-κB is a well-

known host transcription factor that exists in the form of the NF-κB-IκB complex in

resting cells, but IκB can degrade and release the NF-κB dimer to enter the

nucleus and promote gene transcription in response to external stimulation. Using

the coimmunoprecipitation (Co-IP) assay, we verified that the ICP34.5 protein had

a specific interaction with IKKα/β, and then ICP34.5 could dephosphorylate IKKα/β.

Moreover, the overexpression of ICP34.5 effectively inhibited lipopolysaccharide

(LPS) -induced NF-κB pathway activation by inhibiting p65 entry into the nucleus

(*Figure 2B-C*).

To further clarify the underlying mechanism, we next performed the IP-MS assay

to identify other potential molecules contributing to this reactivation in J-Lat 10.6-

ICP34.5 overexpressing cells, and we found that ICP34.5 can also interact with

heat shock 1 protein (HSF1) (*table supplement* 1). HSF1 has been reported as a

transcription factor correlated with the reactivation of HIV latency (Xu et al., 2022,

Lin et al., 2018, Zeng et al., 2017). To test whether HSF1 contributes to the

reactivation of HIV latency by HSV-ΔICP34.5-based constructs, KRIBB11, an

inhibitor of HSF1, was administered to HSV-ΔICP34.5-infected J-Lat 10.6 cells.

The results indicated that the reactivation ability of HSV-ΔICP34.5 was significantly

inhibited by KRIBB11 treatment in a dose-dependent manner (*Figure 2D and E*).

Furthermore, a significant enhancement of the binding of HSF1 to the HIV LTR

was observed upon HSV-ΔICP34.5 infection, leading to an increase in the

reactivation of HIV latency (*Figure 2F*). The direct interaction between ICP34.5 and

HSF1 was also identified by Co-IP assay. Importantly, HSF1 was effectively

dephosphorylated at Ser320 as a result of the overexpression of ICP34.5, while

no influence on the mRNA or protein expression of HSF1 was observed (*Figure 2,*

G and H, Figure 2-figure supplement 2). Considering that protein phosphatase 1

(PP1) can interact with ICP34.5 and dephosphorylate eIF2α (Li et al., 2011), we

then investigated the interaction between PP1α and ICP34.5 (*Figure 2I*).

Additionally, a direct interaction between PP1α and HSF1 was found (*Figure 2J-*

L), allowing for the dephosphorylation of HSF1 and then affecting its ability to

reactivate HIV latency.

Collectively, these findings demonstrated that our modified HSV-ΔICP34.5-

based constructs effectively reactivated HIV latency by modulating the IKKα/β-NF-

κB pathway and PP1-HSF1 pathway.

Construction of recombinant HSV-1 expressing exogenous SIV

genes elicited robust immune responses in mice

Prior research suggested that HSV-ΔICP34.5 holds significant promise in

reactivating latent HIV efficiently. Subsequently, we explored the potential of

advancing HSV-ΔICP34.5 as a bifunctional therapeutic vector to revive latent viral

reservoirs and inducing antigen-specific immune responses against chronic HIV

infection. To achieve this, the HSV-1 ICP34.5 gene was selectively knocked out,

and exogenous antigens were introduced using the bacterial artificial chromosome

(BAC)/galactokinase (galK) selection system. Additionally, the ICP47 gene was

ablated to augment the immunogenicity of the HSV vector (*Figure 3A, Figure 3-*

figure supplement 3). A series of recombinant HSV-ΔICP34.5ΔICP47-based

vectors expressing SIV gag and env antigen were constructed, and the antigen

expression of these constructs was confirmed by Western blotting assay (*Figure*

3B and C). Furthermore, to improve the immunogenicity of the targeted antigen,

we fused the SIV gag with soluble PD1 (sPD1), enabling it to competitively bind

with PD-L1 and thereby block the PD1/PD-L1 immune inhibitory pathway (*Figure*

3D). Consistent with the above findings, these HSV-ΔICP34.5ΔICP47-based SIV

vaccines also efficiently reactivated latent HIV proviruses (*Figure 3-figure*

supplement 4). Then, the immunogenicity of the above modified HSV-

ΔICP34.5ΔICP47-based SIV vaccine was assessed in mice (*Figure 3E*). Our

results showed that these constructs effectively elicited SIV antigen-specific T cell

immune responses using the interferon γ (IFN-γ) ELISpot assay and the

intracellular cytokine staining (ICS) assay (*Figure 3F-M*). Of note, the frequency of

SIV Gag-specific IFN-γ -secreting spot-forming cells (SFCs) in the HSV-sPD1-

SIVgag group (1350 SFCs per 10^6 splenocytes) was significantly higher than that

in the HSV-SIVgag group (498 SFCs per 10^6 splenocytes) (*Figure 3F*). The

frequency of SIV Env2-specific IFN-γ -secreting SFCs in the HSV- SIVenv group

was significantly higher than the HSV-empty group (*Figure 3G*). Furthermore, the

functionality of antigen-specific T cell subsets in response to SIV antigen

stimulation was confirmed using the ICS assay (*Figure 3H-M*). Consistently, the

HSV-sPD1-SIVgag group showed a significantly higher frequency of SIV-specific

CD3+ T, CD4+ T, and CD8+ T cell subsets secreting IFN-γ, IL-2, and TNF-α

cytokines compared to the HSV-SIVgag group (*Figure 3I-K*). Notably, a heightened

proportion of Gag-specific effector memory T cells (Tem) of CD8+ T cell subset

was observed in comparison to the HSV-Gag group (*Figure 3L*). In addition, a

higher frequency of SIV- Env2-specific CD4+ T cells secreting IFN-γ was observed

in the HSV-SIVenv group than in the HSV-empty group (*Figure 3M*). These data

indicated that the vaccines constructed in this study elicit a T cell immune response

in mice. Moreover, the blockade of PD1/PDL1 signaling pathway effectively

enhanced vaccine-induced T cell immune responses, which was consistent with

our and other previous studies (Zhou et al., 2013, Wu et al., 2022, Pan et al., 2018,

Xiao et al., 2014).

The modified HSV-based constructs efficiently elicited SIV-specific immune responses in chronically SIV-infected macaques

Next, the immunogenicity of these HSV-vectored SIV vaccines was further

investigated in chronically SIV_{mac239}-infected rhesus macaques (RMs). To mimic

chronically infected HIV patients in clinic practice, all RMs used in this study were

chronically infected with SIV and received ART treatment for several years, as

reported in our previous studies (Pan et al., 2018, Yang et al., 2019, Wu et al.,

2021, Wu et al., 2022, He et al., 2023). Based on sex, age, viral load, and CD4

count, nine RMs were assigned into three groups: ART+saline group (n=3),

ART+HSV-empty group (n=3), and ART+HSV-sPD1-SIVgag/SIVenv group (n=3).

All RMs received ART (FTC/6.7 mg/kg/once daily, PMPA/10 mg/kg/once daily)

treatment to avoid the interference of free SIV particles. On day 33 and day 52,

these RMs were immunized with saline, HSV-empty, or HSV-sPD1-

SIVgag/SIVenv respectively. On day 70, ART treatment in all RMs was

discontinued to evaluate the time interval of viral rebound. Samples were collected

at different time points to monitor virological and immunological parameters (Figure

4A, *table supplement* 2). To reduce the impact of individual RM variations, the

difference in SIV-specific IFN-γ-secreting SFCs between post-immunization and

pre-immunization (∆SFCs) was used to evaluate the immune response induced by

HSV-vectored SIV vaccines. The results showed that SIV Gag-specific ∆SFCs in

the ART+HSV-sPD1-SIVgag/SIVenv group were greatly increased when

compared with those in the ART+HSV-empty group and ART+saline group (*Figure*

4B-D). A similar enhancement of SIV Gag-specific TNF-α -secreting CD4$^+$ T and

CD8$^+$ T subsets was also verified by ICS assay (*Figure 4E*). Collectively, these

data demonstrated that the HSV-sPD1-SIVgag/SIVenv construct elicited robust

SIV-specific T cell immune responses in ART-treated, SIV-infected RMs.

The modified HSV-based constructs effectively reactivated SIV

latency in vivo and delayed viral rebound in chronically SIV-

infected, ART-treated macaques

Finally, we investigated the therapeutic efficacy of HSV-sPD1-SIVgag/SIVenv in

chronically SIV-infected, ART-treated RMs. Consistent with our previous studies,

the plasma viral load (VL) in these RMs was effectively suppressed during ART

treatment, but rebounded after ART discontinuation. The VL in the ART+saline

group promptly rebounded after ART discontinuation, with an average 8.63-fold

increase in the rebounded peak VL compared with the pre-ART VL (*Figure 5A, D*

and E). However, plasma VL in the ART+HSV-empty group and the ART+HSV-

sPD1-SIVgag/SIVenv group exhibited a delayed rebound interval (*Figure 5B-D*).

Remarkably, there was a lower rebounded peak VL than pre-ART VL in the

ART+HSV-sPD1-SIVgag/SIVenv group (average 12.20-fold decrease), while a

higher rebounded peak VL than pre-ART VL in the ART+HSV-empty group

(average 2.74-fold increase) (*Figure 5E*). Then, we assessed the potential effect

on the latent SIV reservoirs in vivo by administering our modified HSV-based SIV

therapeutic constructs in these RMs. Although there were no obvious viral blips

observed in these RMs, we found significant suppression of integrated SIV DNA

provirus in the ART+HSV-sPD1-SIVgag/SIVenv group. However, the copies of the

SIV DNA provirus were significantly improved in the ART+HSV-empty group and

ART+saline group (*Figure 5F*). More interestingly, we next assessed the

magnitude of SIV Pol antigen-specific immune responses, which could represent

to some extent the level of newly generated virions from the reactivated SIV

reservoirs, because SIV Pol antigen was not included in our designed vaccine

constructs. Specifically, the Pol-specific SFCs amount on Day 70 (511 SFCs/10^6

PBMCs, post-vaccination) was higher than that on Day 33 (315 SFCs/10^6 PBMCs,

pre-vaccination) in the ART+HSV-sPD1-SIVgag/SIVenv group. In addition, there

was a similar observation in the ART+HSV-empty group. In contrast, the Pol-

specific SFCs gradually decreased with the duration of ART treatment in the

ART+saline group (*Figure 5G*). In addition, the CD4$^+$/CD8$^+$ T cell ratio (*Figure 5H*)

and body weight (*Figure 5-figure supplement 5*) after treatment were effectively

ameliorated in the RMs of the ART+HSV-sPD1-SIVgag/SIVenv group, but not in

the ART+HSV-empty group and ART+saline group. Taken together, these findings

suggested that the latent SIV reservoirs might be effectively purged because of the

effect of simultaneously reactivating viral latency and eliciting SIV-specific immune

responses by our modified HSV-based SIV therapeutic constructs, thus resulting

in a delayed viral rebound in chronically SIV-infected, ART-treated macaques.

Discussion

To conquer the continuous epidemic of AIDS, exploring novel strategies to render and eliminate HIV latency should stand as a pivotal pursuit. Currently, numerous strategies, including shock and kill, block and lock, chimeric antigen receptor T-cell therapy, therapeutic vaccination, and gene editing, have been extensively investigated to target the latent HIV reservoirs for an HIV functional cure (Deeks, 2012, Yeh and Ho, 2021, Maldini et al., 2020, Herzig et al., 2019, Dash et al., 2023, Dashti et al., 2023, Walker-Sperling et al., 2022). However, there is no safe and effective approach for clinical use in HIV-1 patients yet. In the present study, we occasionally found that the modified HSV-ΔICP34.5-based constructs could reactivate HIV latency more efficiently than wild-type HSV counterpart, which inspired us to develop a proof-of-concept strategy based on a bifunctional HSV-vectored therapeutic vaccine, aiming to simultaneously reactivate viral latency and elicit HIV/SIV-specific immune responses for HIV functional cure. Our results indicated that these modified HSV-based constructs efficiently elicited antigen-specific immune responses in mice and chronically SIV-infected macaques, and further therapeutic efficacy experiments showed that this strategy effectively reactivated SIV latency in vivo and delayed viral rebound in chronically SIV-infected, ART-treated macaques (*Figure 6*).

The latent HIV proviruses can harbor it into the host genome with a quiescent transcription state, and thus cannot be recognized by immune surveillance or drug killing (Churchill et al., 2016, Pierson et al., 2000). Therefore, it is critical to disrupt viral latency for developing an HIV cure strategy. Based on our experimental data,

the mechanism for efficiently reactivating viral latency by the modified HSV-

ΔICP34.5-based constructs may involve regulating the IKKα/β-NF-κB pathway and

PP1-HSF1 pathway. Indeed, during its replication, HSV can activate the double-

stranded RNA-dependent protein kinase (PKR) pathway, and thus phosphorylate

the protein translation initiation regulator eIF2α, resulting in the initiation of protein

translation (Farassati et al., 2001). Previous studies have shown that the

reactivation potential of HSV might be intertwined with NF-κB, Sp1, and other

unknown transcription factors by ICP0, ICP4, and ICP27 (Chapman et al., 1991,

Amici et al., 2004, Vlach and Pitha, 1992, Mosca et al., 1987b, Golden et al., 1992,

Vlach and Pitha, 1993, Mosca et al., 1987a). However, the underlying mechanism

by which the ICP34.5-deleted HSV construct can greatly improve the reactivation

efficacy of HIV latency remains elusive. ICP34.5 is a neurotoxic factor that can

antagonize innate immune responses, including PKR, TANK binding kinase (TBK1)

signaling, and Beclin1-mediated apoptosis (He et al., 1997, Manivanh et al., 2017,

Orvedahl et al., 2007). ICP34.5 binds to host PP1 and mediates the

dephosphorylation of eIF2α, thus allowing protein synthesis and reversing the

effects of PKR and host antiviral functions (He et al., 1997, He et al., 1998). In this

study, our findings further unveiled an interaction between ICP34.5 and HSF1,

resulting in reduced HSF1 phosphorylation via recruitment of PP1α. Interestingly,

previous reports indicated that HSF1 could positively regulate HIV gene

transcription (Rawat and Mitra, 2011), which is facilitated by its nuclear entry post-

phosphorylation and subsequent recruitment of p300 for self-acetylation, along

with binding to the HIV-1 LTR. Studies have also shown that HSF1 could further

orchestrate p-TEFb recruitment to promote HIV-1 transcriptional elongation (Peng

et al., 2020, Lin et al., 2018, Pan et al., 2016b, Pan et al., 2016a). Under stress-

induced conditions, phosphorylation triggers the formation of the HSF1 trimer, thus

facilitating its nuclear entry to bind to heat shock elements (HSEs) for gene

transcription regulation (Bonner et al., 2000, Timmons et al., 2020). Additionally,

we also demonstrated that ICP34.5 interacted with IKKα and IKKβ, thereby

impeding NF-κB nuclear entry and further curbing HIV latency. Consistently,

previous studies have also suggested that ICP34.5 could disrupt the NF-κB

pathway and possibly affect the maturation of dendritic cells (Jin et al., 2011).

Intriguingly, these findings collectively indicated that ICP34.5 might play an

antagonistic role with the reactivation potential of HSV, and thus our modified HSV-

ΔICP34.5 constructs reactivate HIV/SIV latency through the release of

imprisonment from ICP34.5.

Another observation in this study is that the HSV-sPD1-SIVgag/SIVenv

construct elicited robust and persistent SIV-specific T cell immune responses in

ART-treated, chronically SIV-infected macaques. Increasing evidence has

indicated that HIV-specific cytolytic T lymphocytes (CTLs) can facilitate the

suppression of latent viral reservoirs, and thus, the induction of robust and

persistent HIV-specific CTL responses is essential for achieving long-term

disease-free and transmission-free HIV control (Collins et al., 2020). Featured

polyfunctional CD8[+] T cells may contribute to HIV elite controllers or long-

term non-progressors, a rare proportion of HIV-infected individuals who can

spontaneously control viral replication even without ART treatment (Owen et al.,

2010, Ferre et al., 2009, O'Connell et al., 2009). In addition, both our previous study and others have demonstrated that strong antigen-specific CD8[+] T cell immune responses, especially effector memory CD8[+] T cells, were associated with a lower viral load, and in vivo CD8[+] lymphocyte depletion with intravenous infusion of anti-CD8 monoclonal antibody could lead to dramatical viral rebound in these vaccinated elite macaques (Perdomo-Celis et al., 2022, Pandrea et al., 2011). Notably, previous studies have shown that HSV-based vaccines expressing HIV/SIV antigen did elicit specific CD8[+] T cell immune responses in mice and macaques, but the magnitude was not as strong as other viral vectors (Parker et al., 2007, Murphy et al., 2000, Kaur et al., 2007). Therefore, to further improve its immunogenicity in vivo, some modifications were adopted to optimize the HSV-vectored vaccine. 1) The ICP47 protein, encoded by the US12 gene, can bind to the transporter associated with antigen presentation (TAP) 1/2, inhibiting the transport of viral peptides into the endoplasmic reticulum and the loading of peptides onto nascent major histocompatibility complex (MHC) class I molecules to activate CD8[+] T cells (Orr et al., 2005). Therefore, the ICP47 gene was deleted from our developed HSV vector. 2) Negative immunoregulatory molecules, including PD1, TIM-3, and LAG-3, are usually involved in the pathogenesis of HIV infection, as well as in chronically SIV-infected macaques. Among them, PD1 upregulation can result in the exhaustion and dysfunction of CD8[+] T cells (Trautmann et al., 2006). In addition, PD1 expression on memory CD4[+] T cells might be linked with HIV latency. In this study, we modified the SIV antigen by fusing to soluble PD1 (sPD1), which can block the PD1/PDL1 pathway by

competitively binding with PDL1 and thus improve HIV/SIV vaccine-induced CD8[+]

T cell immune responses (Zhou et al., 2013, Wu et al., 2022, Pan et al., 2018, Xiao

et al., 2014).

Although promising, there are still some limitations to our study. First, this is a

pilot study with relatively small numbers of RMs, and future studies with a larger

number of animals can be conducted to better verify our strategy. Second, the

HSV-sPD1-SIVgag/SIVenv vaccine resulted in delayed viral rebound and a lower

peak viral load post-rebound than pre-ART treatment but did not completely

suppress SIV virus rebound in this study, which may be attributed to suboptimal

doses and treatment, implying that we should further optimize this regimen for

eventually achieving an HIV functional cure in future studies. Taken together, these

findings demonstrated that our modified HSV-ΔICP34.5-based constructs

potentially reactivated HIV/SIV latency by modulating the IKKα/β-NF-κB pathway

and PP1-HSF1 pathway, and thereby we developed a proof-of-concept HIV

functional cure strategy based on a bifunctional HSV-vectored therapeutic vaccine,

which can provide insights into the rational design of novel strategies for pursuing

an HIV functional cure.

Methods

Mouse ethics statement and vaccination

Female BALB/c mice, aged six to eight weeks, were procured from the Experimental Animal Center of Sun Yat-sen University. A total of twenty-five mice were randomly divided into five groups to evaluate the immunogenicity of the recombinant HSV-vector-based SIV vaccine. During the initial week, each mouse was subcutaneously administered a vaccination of 1×10^6 PFU of the respective recombinant HSV-vector vaccines (HSV-empty, HSV-sPD1, HSV-SIVgag, HSV-sPD1-SIVgag, HSV-SIVenv). Following a two-week interval, a booster vaccination was administered via the subcutaneous route using 2×10^6 PFU of recombinant HSV-vector vaccines. The subsequent assessment of the immune response involved ELISpot and intracellular cytokine staining (ICS) assays in accordance with the vaccination schedule.

Ethics statement and vaccination of macaques

A total of 9 chronically SIV_{mac239}-infected rhesus macaques (RMs) were included in this study. All RMs received ART treatment (FTC/6.7 mg/kg/once daily, PMPA/10 mg/kg/once daily) for a duration of 33 days. The RMs were allocated into three groups based on their baseline plasma SIV viral loads (VL): ART + saline (n=3) as the control group, ART+ HSV-empty (n=3) as the sham group, and ART+ HSV-sPD1-SIVgag/SIVenv (n=3) as the vaccinated group. On day 0, all RMs received ART treatment. Once the VL dropped below the limit of detection (1×10^2 copies/mL), the vaccinated group and sham group were primed with a

subcutaneous vaccination of 1×10^7 PFU HSV-sPD1-SIVgag/SIVenv or HSV-

empty, respectively, while the control group received an injection of 0.9% saline.

After three weeks of the prime vaccination, the vaccinated group and sham group

RMs received a subcutaneous booster vaccination of 5×10^7 PFU HSV-sPD1-

SIVgag/SIVenv or HSV-empty, respectively. After two weeks of booster

vaccination, the ART treatment for all RMs was interrupted. The immune response

was evaluated by IFN-γ ELISpot and ICS assays. Plasma viral load was regularly

monitored throughout the studied period.

Peptide pools

The peptide pools, encompassing the complete sequences of SIV Gag, Env, and

Pol proteins, comprising 15 amino acids with 11 overlaps, were generously

provided by the HIV Reagent Program, National Institutes of Health (NIH), USA.

Gag pools comprise 125 peptides, while Env and Pol pools are subdivided into

Env1 (109 peptides) and Env2 (109 peptides) pools, as well as Pol1 (131 peptides)

and Pol2 (132 peptides) pools. These peptide pools were dissolved in dimethyl

sulfoxide (DMSO, Sigma) at a concentration of 0.4 mg/ peptide/mL for subsequent

immunological assays.

Construction of recombinant HSV

413   For the generation of recombinant HSV-vector-based vaccines, we implemented

414   modifications at the ICP34.5 loci through homologous recombination within the

415   BAC-HSV-1 system. Specifically, double copies of the ICP34.5 and ICP47 genes

416   were either deleted or inserted into the respective counterparts: sPD1, SIVgag,

417   sPD1-SIVgag, and SIVenv genes. The sPD1-SIVgag gene was created by fusing

the N-terminal region of mouse soluble PD1 (sPD1) with the C-terminal segment

of SIVgag, connected by a GGGSGGG linker, which was engineered through

overlapping PCR.

IFN-γ ELISpot assay

The IFN-γ- ELISpot assay was conducted in accordance with our previous study

(Sun et al., 2010). In the mouse experiment, $2.5{\times}10^5$ freshly isolated mouse spleen

lymphocytes were simulated with SIV Gag, Env1, and Env2 peptide pools. In the

monkey experiment, $2{\times}10^5$ peripheral blood mononuclear cells (PBMCs) were

seeded and simulated with the SIV Gag, Env, and Pol peptide pools. Mock

stimulation utilized DMSO (Sigma), while concanavalin A (ConA, 10 μg/mL) was

employed as a positive control. Spot quantification was performed using an

ELISpot reader (Mabtech), and peptide-specific spot counts were determined by

subtracting the spots from mock stimulation.

Intracellular cytokine staining (ICS)

The ICS assay was performed as described previously (Wu et al., 2022). In the

mouse experiment, $2{\times}10^6$ freshly isolated mouse spleen lymphocytes were

stimulated with SIV Gag, Env1, and Env2 peptide pools. For the monkey

experiment, $2{\times}10^6$ PBMCs were seeded and simulated with the SIV Gag, Env, and

Pol peptide pools. DMSO and PMA (Thermo Scientific) plus ionomycin were used

as the mock and positive controls, respectively. Data analysis was performed using

FlowJo software (version 10.8.1), and the antibodies used are listed in *table*

439 *supplement* 3. The frequencies of cytokines produced from peptide-specific cells

440 were analyzed by subtracting mock stimulation.

SIV viral RNA and DNA copy assays and absolute T cell count

Absolute T cell count and the levels of plasma mRNA and SIV total DNA were

quantified as described previously (Wu et al., 2022). Plasma viral RNA copy

numbers were determined via SYBR green-based real-time quantitative PCR

(Takara), using SIV gag-specific primers (*table supplement 4*). Viral RNA copy

numbers were calculated based on the standard curve established using SIV_{mac239}

gag standards. The lower limit of detection for this assay was 100 copies/mL of

plasma. Total cellular DNA was extracted from approximately 0.5 to 5 million cells

using a QIAamp DNA Blood Minikit (Qiagen). PCR assays were performed with

200 ng samples of DNA, and SIV viral DNA was quantified using a pair of primers

targeting a conserved region of the SIV gag gene, as previously described.

Quantitation was performed by comparing the results to the standard curve of SIV

gag copies.

Plasmids, cells, and viruses

pVAX-Myc-PDL1, pVAX-Flag-ICP34.5, pVAX-Myc-HSF1, pVAX-HA-PP1α: full-

length mouse PDL1, full-length HSV-1 ICP34.5, and full-length human HSF1 and

human PP1α with N or C-terminal Tag were cloned into the pVAX vector. pVAX-

empty, pcDNA3.1-IKKα, and pcDNA3.1-IKKβ plasmids were stored in our

laboratory. pVAX-empty was used as a mock transfection in our study.

293T cells (from the embryonic kidney of a female human fetus), Vero cells (from

the kidney of a female normal adult African green monkey), and Hela cells (from

the cervical cancer cells of an African American woman) were cultured in complete

Dulbecco's modified Eagle's medium (DMEM, Gibco) containing 10% fetal bovine

serum (FBS, Gibco) and 1% penicillin/streptomycin (Gibco) at 37°C in an

atmosphere of 5% CO_2. The J-Lat 10.6 cells (Jurkat cells contain the HIV-1 full-

length genome whose Env was frameshifted and inserted with GFP in place of Nef)

and the HIV-1 latently infected CD4[+] CEM cells ACH-2 (A3.01 cell integrated HIV-

1 proviral DNA) were cultured in complete RPIM640 (Gibco) containing 10% FBS

and 1% penicillin/streptomycin at 37°C in an atmosphere of 5% CO_2. J-Lat 10.6-

ICP34.5 cells were constructed in our laboratory.

The HSV-1 (McKrae) and HSV-1 (F) strains were stored in our laboratory. HSV-

1 (17 strain) was rescued from pBAC-GFP-HSV in Vero cells. HSV-ΔICP34.5 was

rescued from pBAC-GFP-HSV-ΔICP34.5 (with double copies of ICP34.5 deleted)

in Vero cells.

475   RT-qPCR

476   RT-qPCR was performed as described in our previous study (Zhao et al., 2022).

477   Data were normalized to β-actin. The primer sequences are listed in Supplemental

478   Table 4. Fold changes in the threshold cycle (Ct) values were calculated using the

479    $2^{-\Delta\Delta Ct}$ method.

480   Chromatin immunoprecipitation (ChIP) assay

481   ChIP analysis was conducted using the SimpleChIP Enzymatic Chromatin IP kit

482   (Agarose Beads) (CST), following the manufacturer's instructions. Quantitative

483   real-time PCR was employed for detecting the LTR sequence. The sequences of

484 primers used in the LTR ChIP are listed in Supplemental Table 4. % Input= 2% $\times$

485 $2^{[(ct)2\% \text{ input sample (ct) IP sample}]}$.

Co-immunoprecipitation (Co-IP)

487 Cells were harvested and subjected to Co-IP assay, following the protocol outlined

488 in our previous study (Zhao et al., 2022).

Western blotting analysis

490 Nuclear and cytoplasmic proteins were extracted by kits following the

491 manufacturer's instructions (Beyotime). The Western blotting assay was

492 performed as previously described (Zhao et al., 2022).

Statistical analysis

494 GraphPad Prism 8.0 (GraphPad Software, San Diego, California) was used for

495 statistical analysis. For intragroup direct comparisons, Student's unpaired two-

496 tailed t test was performed to analyze significant differences. For comparisons of

497 multiple groups, one-way ANOVAs were performed. Significance levels are

498 indicated as *P< 0.05, ** P< 0.01, *** P< 0.001, **** P< 0.0001.

Study approval

500 Mice experiment was approved by the Laboratory Animal Ethics Committee

501 guidelines at Sun Yat-sen University (approval number: SYSU-IACUC-2021-

502 000185). Chinese rhesus macaques (Macaca mulatta) were housed at the Landau

503 Animal Experimental Center, Guangdong Landau Biotechnology Co., Ltd.

504 (approval number: N2021101).

Acknowledgements

We appreciate the staff at the Animal Center of GIBH for their excellent technical assistance. We thank all other members of our group for their helpful advice and discussion to improve this project. We also appreciate the NIH AIDS Research and Reference Reagent Program for providing SIV peptide pools.

Additional information

Competing interests

The authors have declared that no conflict of interest exists.

Funding

This project was supported by National Natural Science Foundation of China (81971927, 82271786, 32370171), National Key R&D Program of China (2022YFE0203100, 2021YFC2300103).

Author Contributions

All authors were involved in drafting or critically revising the manuscript for important intellectual content, and all authors approved the final version for publication. CS conceived the project and provided the main funding. CS and LC designed the experiments and reviewed the manuscript. ZW, PL, YY, and CW performed most of the experiments and analyzed the data. ZW, PL, YY, CW, ML, HW, MS, YH, and MC established the animal models and performed animal experiments. ZW and CS interpreted the data and wrote the manuscript.

Additional files

Supplementary files

Figure S1-S5. Table S1-S4.

Data Availability Statement

529   Source data contain the numerical data used to generate the figures.

References

AMICI, C., BELARDO, G., ROSSI, A. & SANTORO, M. G. 2001. Activation of I kappa b kinase by herpes
simplex virus type 1. A novel target for anti-herpetic therapy. *J Biol Chem* **276**: 28759-28766.DOI:
https://doi.org/10.1074/jbc.M103408200, PMID: 11387335
AMICI, C., BELARDO, G., ROZERA, C., BERNASCONI, D. & SANTORO, M. G. 2004. Inhibition of
herpesvirus-induced HIV-1 replication by cyclopentenone prostaglandins: role of IkappaB kinase
(IKK). *Aids* **18**: 1271-1280.DOI: https://doi.org/10.1097/00002030-200406180-00005, PMID:
15362659
ARCHIN, N. M., KIRCHHERR, J. L., SUNG, J. A., CLUTTON, G., SHOLTIS, K., XU, Y., ALLARD, B., STUELKE,
E., KASHUBA, A. D., KURUC, J. D., ERON, J., GAY, C. L., GOONETILLEKE, N. & MARGOLIS, D. M. 2017.
Interval dosing with the HDAC inhibitor vorinostat effectively reverses HIV latency. *J Clin Invest* **127**:
3126-3135.DOI: https://doi.org/10.1172/jci92684, PMID: 28714868
ARCHIN, N. M., SUNG, J. M., GARRIDO, C., SORIANO-SARABIA, N. & MARGOLIS, D. M. 2014.
Eradicating HIV-1 infection: seeking to clear a persistent pathogen. *Nat Rev Microbiol* **12**: 750-
764.DOI: https://doi.org/10.1038/nrmicro3352, PMID: 25402363
BISGROVE, D. A., MAHMOUDI, T., HENKLEIN, P. & VERDIN, E. 2007. Conserved P-TEFb-interacting
domain of BRD4 inhibits HIV transcription. *Proc Natl Acad Sci U S A* **104**: 13690-13695.DOI:
https://doi.org/10.1073/pnas.0705053104, PMID: 17690245
BONNER, J. J., CHEN, D., STOREY, K., TUSHAN, M. & LEA, K. 2000. Structural analysis of yeast HSF
by site-specific crosslinking. *J Mol Biol* **302**: 581-592.DOI:
https://doi.org/10.1006/jmbi.2000.4096, PMID: 10986120
CALISTRI, A., PAROLIN, C. & PALù, G. 2003. Herpes simplex virus type 1 can either suppress or
enhance human immunodeficiency virus type 1 replication in CD4-positive T lymphocytes. *J Med
Virol* **70**: 163-170.DOI: https://doi.org/10.1002/jmv.10350, PMID: 12629659
CHAPMAN, C. J., HARRIS, J. D., COLLINS, M. K. L. & LATCHMAN, D. S. 1991. A recombinant HIV
provirus is synergistically activated by the HIV Tat protein and the HSV IE1 protein but not by the
HSV IE3 protein. *AIDS* **5**,
CHURCHILL, M. J., DEEKS, S. G., MARGOLIS, D. M., SILICIANO, R. F. & SWANSTROM, R. 2016. HIV
reservoirs: what, where and how to target them. *Nat Rev Microbiol* **14**: 55-60.DOI:
https://doi.org/10.1038/nrmicro.2015.5, PMID: 26616417
COLLINS, D. R., GAIHA, G. D. & WALKER, B. D. 2020. CD8(+) T cells in HIV control, cure and
prevention. *Nat Rev Immunol* **20**: 471-482.DOI: https://doi.org/10.1038/s41577-020-0274-9,
PMID: 32051540
DASH, P. K., CHEN, C., KAMINSKI, R., SU, H., MANCUSO, P., SILLMAN, B., ZHANG, C., LIAO, S.,
SRAVANAM, S., LIU, H., WAIGHT, E., GUO, L., MATHEWS, S., SARIYER, R., MOSLEY, R. L.,
POLUEKTOVA, L. Y., CAOCCI, M., AMINI, S., GORANTLA, S., BURDO, T. H., EDAGWA, B., GENDELMAN,
H. E. & KHALILI, K. 2023. CRISPR editing of CCR5 and HIV-1 facilitates viral elimination in
antiretroviral drug-suppressed virus-infected humanized mice. *Proc Natl Acad Sci U S A* **120**:
e2217887120.DOI: https://doi.org/10.1073/pnas.2217887120, PMID: 37126704
DASHTI, A., SUKKESTAD, S., HORNER, A. M., NEJA, M., SIDDIQI, Z., WALLER, C., GOLDY, J., MONROE,
D., LIN, A., SCHOOF, N., SINGH, V., MAVIGNER, M., LIFSON, J. D., DELEAGE, C., TUYISHIME, M.,
FALCINELLI, S. D., KING, H. A. D., KE, R., MASON, R. D., ARCHIN, N. M., DUNHAM, R. M., SAFRIT, J.
T., JEAN, S., PERELSON, A. S., MARGOLIS, D. M., FERRARI, G., ROEDERER, M., SILVESTRI, G. &

CHAHROUDI, A. 2023. AZD5582 plus SIV-specific antibodies reduce lymph node viral reservoirs in antiretroviral therapy-suppressed macaques. *Nat Med*.DOI: https://doi.org/10.1038/s41591-023-02570-7, PMID: 37783968

DEEKS, S. G. 2012. HIV: Shock and kill. *Nature* **487**: 439-440.DOI: https://doi.org/10.1038/487439a, PMID: 22836995

FARASSATI, F., YANG, A. D. & LEE, P. W. 2001. Oncogenes in Ras signalling pathway dictate host-cell permissiveness to herpes simplex virus 1. *Nat Cell Biol* **3**: 745-750.DOI: https://doi.org/10.1038/35087061, PMID: 11483960

FERRE, A. L., HUNT, P. W., CRITCHFIELD, J. W., YOUNG, D. H., MORRIS, M. M., GARCIA, J. C., POLLARD, R. B., YEE, H. F., JR., MARTIN, J. N., DEEKS, S. G. & SHACKLETT, B. L. 2009. Mucosal immune responses to HIV-1 in elite controllers: a potential correlate of immune control. *Blood* **113**: 3978-3989.DOI: https://doi.org/10.1182/blood-2008-10-182709, PMID: 19109229

GOLDEN, M. P., KIM, S., HAMMER, S. M., LADD, E. A., SCHAFFER, P. A., DELUCA, N. & ALBRECHT, M. A. 1992. Activation of human immunodeficiency virus by herpes simplex virus. *J Infect Dis* **166**: 494-499.DOI: https://doi.org/10.1093/infdis/166.3.494, PMID: 1354237

HE, B., GROSS, M. & ROIZMAN, B. 1997. The gamma(1)34.5 protein of herpes simplex virus 1 complexes with protein phosphatase 1alpha to dephosphorylate the alpha subunit of the eukaryotic translation initiation factor 2 and preclude the shutoff of protein synthesis by double-stranded RNA-activated protein kinase. *Proc Natl Acad Sci U S A* **94**: 843-848.DOI: https://doi.org/10.1073/pnas.94.3.843, PMID: 9023344

HE, B., GROSS, M. & ROIZMAN, B. 1998. The gamma134.5 protein of herpes simplex virus 1 has the structural and functional attributes of a protein phosphatase 1 regulatory subunit and is present in a high molecular weight complex with the enzyme in infected cells. *J Biol Chem* **273**: 20737-20743.DOI: https://doi.org/10.1074/jbc.273.33.20737, PMID: 9694816

HE, Y., WU, C., LIU, Z., ZHANG, Y., FENG, F., LIN, Z., WANG, C., YANG, Q., WEN, Z., LIU, Y., ZHANG, F., LIN, Y., ZHANG, H., QU, L., LI, L., CAI, W., SUN, C., CHEN, L. & LI, P. 2023. Arsenic trioxide-induced apoptosis contributes to suppression of viral reservoir in SIV-infected rhesus macaques. *Microbiol Spectr*: e0052523.DOI: https://doi.org/10.1128/spectrum.00525-23, PMID: 37695104

HENG, M. C., HENG, S. Y. & ALLEN, S. G. 1994. Co-infection and synergy of human immunodeficiency virus-1 and herpes simplex virus-1. *Lancet* **343**: 255-258.DOI: https://doi.org/10.1016/s0140-6736(94)91110-x, PMID: 7905094

HERZIG, E., KIM, K. C., PACKARD, T. A., VARDI, N., SCHWARZER, R., GRAMATICA, A., DEEKS, S. G., WILLIAMS, S. R., LANDGRAF, K., KILLEEN, N., MARTIN, D. W., WEINBERGER, L. S. & GREENE, W. C. 2019. Attacking Latent HIV with convertibleCAR-T Cells, a Highly Adaptable Killing Platform. *Cell* **179**: 880-894.e810.DOI: https://doi.org/10.1016/j.cell.2019.10.002, PMID: 31668804

JIN, H., YAN, Z., MA, Y., CAO, Y. & HE, B. 2011. A herpesvirus virulence factor inhibits dendritic cell maturation through protein phosphatase 1 and Ikappa B kinase. *J Virol* **85**: 3397-3407.DOI: https://doi.org/10.1128/jvi.02373-10, PMID: 21248029

KAUR, A., SANFORD, H. B., GARRY, D., LANG, S., KLUMPP, S. A., WATANABE, D., BRONSON, R. T., LIFSON, J. D., ROSATI, M., PAVLAKIS, G. N., FELBER, B. K., KNIPE, D. M. & DESROSIERS, R. C. 2007. Ability of herpes simplex virus vectors to boost immune responses to DNA vectors and to protect against challenge by simian immunodeficiency virus. *Virology* **357**: 199-214.DOI: https://doi.org/10.1016/j.virol.2006.08.007, PMID: 16962628

KIM, Y., ANDERSON, J. L. & LEWIN, S. R. 2018. Getting the "Kill" into "Shock and Kill": Strategies to Eliminate Latent HIV. *Cell Host Microbe* **23**: 14-26.DOI: https://doi.org/10.1016/j.chom.2017.12.004, PMID: 29324227

LEHRMAN, G., HOGUE, I. B., PALMER, S., JENNINGS, C., SPINA, C. A., WIEGAND, A., LANDAY, A. L.,

COOMBS, R. W., RICHMAN, D. D., MELLORS, J. W., COFFIN, J. M., BOSCH, R. J. & MARGOLIS, D. M. 2005. Depletion of latent HIV-1 infection in vivo: a proof-of-concept study. *Lancet* **366**: 549-555.DOI: https://doi.org/10.1016/s0140-6736(05)67098-5, PMID: 16099290

LEITMAN, E. M., THOBAKGALE, C. F., ADLAND, E., ANSARI, M. A., RAGHWANI, J., PRENDERGAST, A. J., TUDOR-WILLIAMS, G., KIEPIELA, P., HEMELAAR, J., BRENER, J., TSAI, M. H., MORI, M., RIDDELL, L., LUZZI, G., JOOSTE, P., NDUNG'U, T., WALKER, B. D., PYBUS, O. G., KELLAM, P., NARANBHAI, V., MATTHEWS, P. C., GALL, A. & GOULDER, P. J. R. 2017. Role of HIV-specific CD8(+) T cells in pediatric HIV cure strategies after widespread early viral escape. *J Exp Med* **214**: 3239-3261.DOI: https://doi.org/10.1084/jem.20162123, PMID: 28983013

LI, Y., ZHANG, C., CHEN, X., YU, J., WANG, Y., YANG, Y., DU, M., JIN, H., MA, Y., HE, B. & CAO, Y. 2011. ICP34.5 protein of herpes simplex virus facilitates the initiation of protein translation by bridging eukaryotic initiation factor 2alpha (eIF2alpha) and protein phosphatase 1. *J Biol Chem* **286**: 24785-24792.DOI: https://doi.org/10.1074/jbc.M111.232439, PMID: 21622569

LI, Z., GUO, J., WU, Y. & ZHOU, Q. 2013. The BET bromodomain inhibitor JQ1 activates HIV latency through antagonizing Brd4 inhibition of Tat-transactivation. *Nucleic Acids Res* **41**: 277-287.DOI: https://doi.org/10.1093/nar/gks976, PMID: 23087374

LIN, J., ZHANG, X., LU, W., XU, X., PAN, X., LIANG, T., DUAN, S., CHEN, Y., LI, L. & LIU, S. 2018. PR-957, a selective immunoproteasome inhibitor, reactivates latent HIV-1 through p-TEFb activation mediated by HSF-1. *Biochem Pharmacol* **156**: 511-523.DOI: https://doi.org/10.1016/j.bcp.2018.08.042, PMID: 30170098

LOOKER, K. J., ELMES, J. A. R., GOTTLIEB, S. L., SCHIFFER, J. T., VICKERMAN, P., TURNER, K. M. E. & BOILY, M. C. 2017. Effect of HSV-2 infection on subsequent HIV acquisition: an updated systematic review and meta-analysis. *Lancet Infect Dis* **17**: 1303-1316.DOI: https://doi.org/10.1016/s1473-3099(17)30405-x, PMID: 28843576

MALDINI, C. R., CLAIBORNE, D. T., OKAWA, K., CHEN, T., DOPKIN, D. L., SHAN, X., POWER, K. A., TRIFONOVA, R. T., KRUPP, K., PHELPS, M., VRBANAC, V. D., TANNO, S., BATESON, T., LESLIE, G. J., HOXIE, J. A., BOUTWELL, C. L., RILEY, J. L. & ALLEN, T. M. 2020. Dual CD4-based CAR T cells with distinct costimulatory domains mitigate HIV pathogenesis in vivo. *Nat Med* **26**: 1776-1787.DOI: https://doi.org/10.1038/s41591-020-1039-5, PMID: 32868878

MANIVANH, R., MEHRBACH, J., KNIPE, D. M. & LEIB, D. A. 2017. Role of Herpes Simplex Virus 1 γ34.5 in the Regulation of IRF3 Signaling. *J Virol* **91**.DOI: https://doi.org/10.1128/jvi.01156-17, PMID: 28904192

MOSCA, J. D., BEDNARIK, D. P., RAJ, N. B., ROSEN, C. A., SODROSKI, J. G., HASELTINE, W. A., HAYWARD, G. S. & PITHA, P. M. 1987a. Activation of human immunodeficiency virus by herpesvirus infection: identification of a region within the long terminal repeat that responds to a trans-acting factor encoded by herpes simplex virus 1. *Proc Natl Acad Sci U S A* **84**: 7408-7412.DOI: https://doi.org/10.1073/pnas.84.21.7408, PMID: 2823260

MOSCA, J. D., BEDNARIK, D. P., RAJ, N. B., ROSEN, C. A., SODROSKI, J. G., HASELTINE, W. A. & PITHA, P. M. 1987b. Herpes simplex virus type-1 can reactivate transcription of latent human immunodeficiency virus. *Nature* **325**: 67-70.DOI: https://doi.org/10.1038/325067a0, PMID: 3025748

MURPHY, C. G., LUCAS, W. T., MEANS, R. E., CZAJAK, S., HALE, C. L., LIFSON, J. D., KAUR, A., JOHNSON, R. P., KNIPE, D. M. & DESROSIERS, R. C. 2000. Vaccine protection against simian immunodeficiency virus by recombinant strains of herpes simplex virus. *J Virol* **74**: 7745-7754.DOI: https://doi.org/10.1128/jvi.74.17.7745-7754.2000, PMID: 10933680

O'CONNELL, K. A., BAILEY, J. R. & BLANKSON, J. N. 2009. Elucidating the elite: mechanisms of control in HIV-1 infection. *Trends Pharmacol Sci* **30**: 631-637.DOI:

https://doi.org/10.1016/j.tips.2009.09.005, PMID: 19837464

ORR, M. T., EDELMANN, K. H., VIEIRA, J., COREY, L., RAULET, D. H. & WILSON, C. B. 2005. Inhibition

of MHC class I is a virulence factor in herpes simplex virus infection of mice. *PLoS Pathog* **1**: e7.DOI:

https://doi.org/10.1371/journal.ppat.0010007, PMID: 16201019

ORVEDAHL, A., ALEXANDER, D., TALLóCZY, Z., SUN, Q., WEI, Y., ZHANG, W., BURNS, D., LEIB, D. A.

& LEVINE, B. 2007. HSV-1 ICP34.5 confers neurovirulence by targeting the Beclin 1 autophagy

protein. *Cell Host Microbe* **1**: 23-35.DOI: https://doi.org/10.1016/j.chom.2006.12.001, PMID:

18005679

OWEN, R. E., HEITMAN, J. W., HIRSCHKORN, D. F., LANTERI, M. C., BISWAS, H. H., MARTIN, J. N.,

KRONE, M. R., DEEKS, S. G. & NORRIS, P. J. 2010. HIV+ elite controllers have low HIV-specific T-cell

activation yet maintain strong, polyfunctional T-cell responses. *Aids* **24**: 1095-1105.DOI:

https://doi.org/10.1097/QAD.0b013e3283377a1e, PMID: 20400885

PAN, E., FENG, F., LI, P., YANG, Q., MA, X., WU, C., ZHAO, J., YAN, H., CHEN, R., CHEN, L. & SUN, C.

2018. Immune Protection of SIV Challenge by PD-1 Blockade During Vaccination in Rhesus

Monkeys. *Front Immunol* **9**: 2415.DOI: https://doi.org/10.3389/fimmu.2018.02415, PMID:

30405615

PAN, X. Y., ZHAO, W., WANG, C. Y., LIN, J., ZENG, X. Y., REN, R. X., WANG, K., XUN, T. R., SHAI, Y. &

LIU, S. W. 2016a. Heat Shock Protein 90 Facilitates Latent HIV Reactivation through Maintaining

the Function of Positive Transcriptional Elongation Factor b (p-TEFb) under Proteasome Inhibition.

J Biol Chem **291**: 26177-26187.DOI: https://doi.org/10.1074/jbc.M116.743906, PMID: 27799305

PAN, X. Y., ZHAO, W., ZENG, X. Y., LIN, J., LI, M. M., SHEN, X. T. & LIU, S. W. 2016b. Heat Shock Factor

1 Mediates Latent HIV Reactivation. *Sci Rep* **6**: 26294.DOI: https://doi.org/10.1038/srep26294,

PMID: 27189267

PANDREA, I., GAUFIN, T., GAUTAM, R., KRISTOFF, J., MANDELL, D., MONTEFIORI, D., KEELE, B. F.,

RIBEIRO, R. M., VEAZEY, R. S. & APETREI, C. 2011. Functional cure of SIVagm infection in rhesus

macaques results in complete recovery of CD4+ T cells and is reverted by CD8+ cell depletion. *PLoS*

Pathog **7**: e1002170.DOI: https://doi.org/10.1371/journal.ppat.1002170, PMID: 21829366

PARKER, S. D., ROTTINGHAUS, S. T., ZAJAC, A. J., YUE, L., HUNTER, E., WHITLEY, R. J. & PARKER, J.

N. 2007. HIV-1(89.6) Gag expressed from a replication competent HSV-1 vector elicits persistent

cellular immune responses in mice. *Vaccine* **25**: 6764-6773.DOI:

https://doi.org/10.1016/j.vaccine.2007.06.064, PMID: 17706843

PENG, W., HONG, Z., CHEN, X., GAO, H., DAI, Z., ZHAO, J., LIU, W., LI, D. & DENG, K. 2020.

Thiostrepton Reactivates Latent HIV-1 through the p-TEFb and NF-κB Pathways Mediated by Heat

Shock Response. *Antimicrob Agents Chemother* **64**.DOI: https://doi.org/10.1128/aac.02328-19,

PMID: 32094131

PERDOMO-CELIS, F., PASSAES, C., MONCEAUX, V., VOLANT, S., BOUFASSA, F., DE TRUCHIS, P.,

MARCOU, M., BOURDIC, K., WEISS, L., JUNG, C., BOURGEOIS, C., GOUJARD, C., MEYER, L., MüLLER-

TRUTWIN, M., LAMBOTTE, O. & SáEZ-CIRIóN, A. 2022. Reprogramming dysfunctional CD8+ T cells

to promote properties associated with natural HIV control. *J Clin Invest* **132**.DOI:

https://doi.org/10.1172/jci157549, PMID: 35380989

PIERCE, C. A., LOH, L. N., STEACH, H. R., CHESHENKO, N., PRESTON-HURLBURT, P., ZHANG, F.,

STRANSKY, S., KRAVETS, L., SIDOLI, S., PHILBRICK, W., NASSAR, M., KRISHNASWAMY, S., HEROLD,

K. C. & HEROLD, B. C. 2023. HSV-2 triggers upregulation of MALAT1 in CD4+ T cells and promotes

HIV latency reversal. *J Clin Invest* **133**.DOI: https://doi.org/10.1172/jci164317, PMID: 37079384

PIERSON, T., MCARTHUR, J. & SILICIANO, R. F. 2000. Reservoirs for HIV-1: mechanisms for viral

persistence in the presence of antiviral immune responses and antiretroviral therapy. *Annu Rev*

Immunol **18**: 665-708.DOI: https://doi.org/10.1146/annurev.immunol.18.1.665, PMID: 10837072

POH, A. 2016. First Oncolytic Viral Therapy for Melanoma. *Cancer Discov* **6**: 6.DOI: https://doi.org/10.1158/2159-8290.Cd-nb2015-158, PMID: 26552414

RAWAT, P. & MITRA, D. 2011. Cellular heat shock factor 1 positively regulates human immunodeficiency virus-1 gene expression and replication by two distinct pathways. *Nucleic Acids Res* **39**: 5879-5892.DOI: https://doi.org/10.1093/nar/gkr198, PMID: 21459854

SUN, C., ZHANG, L., ZHANG, M., LIU, Y., ZHONG, M., MA, X. & CHEN, L. 2010. Induction of balance and breadth in the immune response is beneficial for the control of SIVmac239 replication in rhesus monkeys. *J Infect* **60**: 371-381.DOI: https://doi.org/10.1016/j.jinf.2010.03.005, PMID: 20227437

TIMMONS, A., FRAY, E., KUMAR, M., WU, F., DAI, W., BULLEN, C. K., KIM, P., HETZEL, C., YANG, C., BEG, S., LAI, J., POMERANTZ, J. L., YUKL, S. A., SILICIANO, J. D. & SILICIANO, R. F. 2020. HSF1 inhibition attenuates HIV-1 latency reversal mediated by several candidate LRAs In Vitro and Ex Vivo. *Proc Natl Acad Sci U S A* **117**: 15763-15771.DOI: https://doi.org/10.1073/pnas.1916290117, PMID: 32571938

TRAUTMANN, L., JANBAZIAN, L., CHOMONT, N., SAID, E. A., GIMMIG, S., BESSETTE, B., BOULASSEL, M. R., DELWART, E., SEPULVEDA, H., BALDERAS, R. S., ROUTY, J. P., HADDAD, E. K. & SEKALY, R. P. 2006. Upregulation of PD-1 expression on HIV-specific CD8+ T cells leads to reversible immune dysfunction. *Nat Med* **12**: 1198-1202.DOI: https://doi.org/10.1038/nm1482, PMID: 16917489

VLACH, J. & PITHA, P. M. 1992. Herpes simplex virus type 1-mediated induction of human immunodeficiency virus type 1 provirus correlates with binding of nuclear proteins to the NF-kappa B enhancer and leader sequence. *J Virol* **66**: 3616-3623.DOI: https://doi.org/10.1128/jvi.66.6.3616-3623.1992, PMID: 1316471

VLACH, J. & PITHA, P. M. 1993. Differential contribution of herpes simplex virus type 1 gene products and cellular factors to the activation of human immunodeficiency virus type 1 provirus. *J Virol* **67**: 4427-4431.DOI: https://doi.org/10.1128/jvi.67.7.4427-4431.1993, PMID: 8389940

WALKER-SPERLING, V. E. K., MERCADO, N. B., CHANDRASHEKAR, A., BORDUCCHI, E. N., LIU, J., NKOLOLA, J. P., LEWIS, M., MURRY, J. P., YANG, Y., GELEZIUNAS, R., ROBB, M. L., MICHAEL, N. L., PAU, M. G., WEGMANN, F., SCHUITEMAKER, H., FRAY, E. J., KUMAR, M. R., SILICIANO, J. D., SILICIANO, R. F. & BAROUCH, D. H. 2022. Therapeutic efficacy of combined active and passive immunization in ART-suppressed, SHIV-infected rhesus macaques. *Nat Commun* **13**: 3463.DOI: https://doi.org/10.1038/s41467-022-31196-5, PMID: 35710819

WU, C., HE, Y., ZHAO, J., LUO, K., WEN, Z., ZHANG, Y., LI, M., CUI, Y., LIU, Z., WANG, C., HAN, Z., LI, G., FENG, F., LI, P., CHEN, L. & SUN, C. 2022. Exacerbated AIDS Progression by PD-1 Blockade during Therapeutic Vaccination in Chronically Simian Immunodeficiency Virus-Infected Rhesus Macaques after Interruption of Antiretroviral Therapy. *J Virol* **96**: e0178521.DOI: https://doi.org/10.1128/JVI.01785-21, PMID: 34818070

WU, C., ZHAO, J., LI, R., FENG, F., HE, Y., LI, Y., HUANG, R., LI, G., YANG, H., CHENG, G., CHEN, L., MA, F., LI, P. & SUN, C. 2021. Modulation of Antiviral Immunity and Therapeutic Efficacy by 25-Hydroxycholesterol in Chronically SIV-Infected, ART-Treated Rhesus Macaques. *Virol Sin* **36**: 1197-1209.DOI: https://doi.org/10.1007/s12250-021-00407-6, PMID: 34057681

XIAO, L., WANG, D., SUN, C., LI, P., JIN, Y., FENG, L. & CHEN, L. 2014. Enhancement of SIV-specific cell mediated immune responses by co-administration of soluble PD-1 and Tim-3 as molecular adjuvants in mice. *Hum Vaccin Immunother* **10**: 724-733.DOI: https://doi.org/10.4161/hv.27340, PMID: 24326266

XU, X., LIN, Y., ZENG, X., YANG, C., DUAN, S., DING, L., LU, W., LIN, J., PAN, X., MA, X. & LIU, S. 2022. PARP1 Might Substitute HSF1 to Reactivate Latent HIV-1 by Binding to Heat Shock Element. *Cells* **11**.DOI: https://doi.org/10.3390/cells11152331, PMID: 35954175

YANG, Q., FENG, F., LI, P., PAN, E., WU, C., HE, Y., ZHANG, F., ZHAO, J., LI, R., FENG, L., HU, F., LI, L.,
ZOU, H., CAI, W., LEHNER, T., SUN, C. & CHEN, L. 2019. Arsenic Trioxide Impacts Viral Latency and
Delays Viral Rebound after Termination of ART in Chronically SIV-Infected Macaques. *Adv Sci*
(Weinh) **6**: 1900319.DOI: https://doi.org/10.1002/advs.201900319, PMID: 31380187
YEH, Y. J. & HO, Y. C. 2021. Shock-and-kill versus block-and-lock: Targeting the fluctuating and
heterogeneous HIV-1 gene expression. *Proc Natl Acad Sci U S A* **118**.DOI:
https://doi.org/10.1073/pnas.2103692118, PMID: 33758027
ZENG, X., PAN, X., XU, X., LIN, J., QUE, F., TIAN, Y., LI, L. & LIU, S. 2017. Resveratrol Reactivates
Latent HIV through Increasing Histone Acetylation and Activating Heat Shock Factor 1. *J Agric Food*
Chem **65**: 4384-4394.DOI: https://doi.org/10.1021/acs.jafc.7b00418, PMID: 28471170
ZHAO, J., CHEN, J., WANG, C., LIU, Y., LI, M., LI, Y., LI, R., HAN, Z., WANG, J., CHEN, L., SHU, Y., CHENG,
G. & SUN, C. 2022. Kynurenine-3-monooxygenase (KMO) broadly inhibits viral infections via
triggering NMDAR/Ca2+ influx and CaMKII/ IRF3-mediated IFN-beta production. *PLoS Pathog* **18**:
e1010366.DOI: https://doi.org/10.1371/journal.ppat.1010366, PMID: 35235615
ZHOU, J., CHEUNG, A. K., TAN, Z., WANG, H., YU, W., DU, Y., KANG, Y., LU, X., LIU, L., YUEN, K. Y. &
CHEN, Z. 2013. PD1-based DNA vaccine amplifies HIV-1 GAG-specific CD8+ T cells in mice. *J Clin*
Invest **123**: 2629-2642.DOI: https://doi.org/10.1172/jci64704, PMID: 23635778
778
779

Figure Legends

Figure 1. The modified HSV-ΔICP34.5-based constructs reactivated HIV latency more efficiently than wild-type HSV counterparts. (**A**) J-Lat 10.6 cells (1×10^6) were infected with varying MOIs of wild-type HSV-1 17 strain for 30 h. The increasing expression level of GFP$^+$ cells with increasing MOI of HSV-1 was displayed with the pseudocolor plot (left) and the corresponding bar chart (right). (**B**) The mRNA levels of HIV-1 LTR, Tat, Gag, Vpr, and Vif in J-Lat 10.6 cells following infection with the wild-type HSV-1 17 strain were shown with the histogram. (**C**) J-Lat 10.6 cells (1×10^6) were infected with HSV-wt or HSV-ΔICP34.5 at an MOI of 0.1 for 30 h, and then the mRNA levels of HIV-1 LTR, Tat, Gag, Vpr, Vif, and (**D**) HSV-1 UL27 were shown with the histogram. (**E**) ACH-2 cells (1×10^6) were infected with HSV-wt or HSV-ΔICP34.5 at an MOI of 0.1 for 30 h, and then the p24 protein level was detected using an HIV-1 p24 ELISA kit. (**F**) ACH-2 cells (1×10^6) were infected with HSV-wt or HSV-ΔICP34.5 at an MOI of 0.1 for 30 h, and then the mRNA levels of HIV-1 LTR, Tat, Gag, Vpr and Vif were shown with the histogram. (**G**) J-Lat 10.6 and J-Lat 10.6-ICP34.5 cells were infected with HSV-wt or HSV-ΔICP34.5, and then the mRNA levels of HIV-1 Tat were shown with the histogram (left). The blotting showed that the J-Lat 10.6 cells stably expressing HSV ICP34.5 (J-Lat 10.6-ICP34.5) can appropriately express ICP34.5 protein using Flag-tag antibodies (right). (**H**) J-Lat 10.6 and J-Lat 10.6-ICP34.5 cells were respectively stimulated with PMA (10 ng/mL) and TNF-α (10 ng/mL), and the expression level of GFP$^+$ cells was displayed with the corresponding bar chart. Data shown are mean ± SD. **P<0.01, ***P<0.001, ****P<0.0001. ns: no significance.

Figure 2. The modified HSV-based constructs effectively reactivated HIV latency by modulating the NF-κB pathway and HSF1 pathway. (**A**) J-Lat 10.6 cells were infected with HSV-wt and HSV-ΔICP34.5 at an MOI of 0.1. The cytoplasmic and nuclear proteins were separated to detect the expression level of p65, p-IKKα/β and IkBα. GAPDH and Lamin B1 served as loading controls for cytoplasmic and nuclear proteins, respectively. (**B**) 293T cells were transfected with Flag-ICP34.5, IKKα (left), or IKKβ (right), and the cell lysates and IP complexes were analyzed through Co-IP assays. (**C**) 293T cells were transfected with Flag-ICP34.5 or empty vector (Vec) for 24h, and then treated with LPS (1μg/mL) for 8h. The cytoplasmic and nuclear proteins were analyzed by Western blotting (WB) assay. (**D-E**) J-Lat 10.6 cells were infected with HSV-ΔICP34.5 and then treated with different concentrations of KRIBB11. The relative fold change in LTR and Tat mRNA was analyzed by qPCR. (**F**) J-Lat 10.6 cells were infected with HSV-wt or HSV-ΔICP34.5 at MOI of 0.1 for 36h. ChIP-qPCR was conducted to evaluate the ability of HSF1 to bind to the LTR. IgG and Histone antibody (His) were used as negative and positive controls, respectively. (**G**) 293T cells were transfected with Flag-ICP34.5 and Myc-HSF1, and the cell lysates and IP complexes were analyzed through Co-IP assays. (**H**) 293T cells were transfected with 0, 1, 2, 4 μg of Flag-ICP34.5 and analyzed by WB analysis. (**I**) 293T cells were transfected with Flag-ICP34.5 and HA-PP1α, and the cell lysates and IP

complexes were analyzed through Co-IP assays. (**J**) 293T cells were transfected
with 0, 0.5, 1, 2 μg of HA-PP1α and analyzed by WB analysis. 293T cells were
transfected with Myc-HSF1 with or without Flag-ICP34.5 (**K**), or were transfected
with Myc-HSF1 and HA-PP1α (**L**), and then the cell lysates and IP complexes were
analyzed through Co-IP assays. Data shown are mean ± SD. ***P*<0.01,
*****P*<0.0001. ns: no significance.
831
Figure 3. Recombinant HSV-1 vector-based SIV vaccines induce specific T
cell immune responses in mice. (**A**) The schematic diagram illustrated the
process of constructing recombinant HSV through the BAC/galK selection system.
Firstly, ICP34.5 gene was replaced with galK gene through homologous
recombination. Subsequently, galK was substituted with the target gene
expression cassette containing the hCMV promoter and BGH terminator. Finally,
the ICP47 gene was deleted. (**B**) The image of one clone of the rescued
recombinant HSV with the brightfield (top) and the fluorescence channel (bottom).
(**C**) Vero cells were infected with recombinant HSV constructs, and the protein
expression of targeted genes was detected by using SIV-infected monkey serum.
(**D**) HeLa cells were transfected with Myc-PDL1, followed by infection with HSV-
empty, HSV-SIVgag, or HSV-sPD1-SIVgag at an MOI of 0.1 for 24 h. The cell
lysates were subjected to Co-IP analysis. (**E**) Schematic schedule of vaccinated
mice. Twenty-five mice were randomly allocated to five groups, HSV-empty, HSV-
sPD1, HSV-SIVgag, HSV-sPD1-SIVgag, and HSV-SIVenv. At week 0 and week 2,
mice were injected with the corresponding vaccines. At week 4, mice were
sacrificed and spleen lymphocytes were collected to evaluate immune response.
(**F-G**) Column graphs depicted the Gag or Env1, Env2-specific spot-forming cells
(SFCs) per 10^6 spleen lymphocytes as measured by IFN-γ ELISpot assay. (**H**) The
pseudocolor plot of flow cytometry illustrated the gating strategy. Column graphs
depicted the frequencies of IFN-γ, IL-2, and TNF-α production from gag-specific
CD3$^+$ T (**I**), CD4$^+$ T (**J**), and CD8$^+$ T cells (**K**). (**L**) The bar chart illustrated the
proportion of Tem of CD4$^+$ T and CD8$^+$ T cells under stimulation with the SIV Gag
peptide pools. (**M**) The bar chart illustrated the frequencies of Env2-specific IFN-γ^+
CD4$^+$ T cells. These data were expressed as the mean±SD from five mice samples.
Three independent experiments for the animal immunization were repeated.
***P*<0.05, ***P*<0.01, ****P*<0.001, *****P*<0.0001. ns: no significance.

Figure 4. The modified HSV-based constructs efficiently elicited SIV-specific
immune responses in chronically SIV-infected macaques. (**A**) Schematic
schedule of the macaque experiment. Nine chronically SIV-infected macaques
were assigned into three groups: ART+saline group (n=3), ART+HSV-empty group
(n=3), and ART+HSV-sPD1-SIVgag/SIVenv group (n=3). All SIV-infected
macaques received ART treatment (FTC/6.7 mg/kg/once daily, PMPA/10
mg/kg/once daily) for 33 days. On day 33 and day 52, macaques were immunized
with saline, HSV-empty, and HSV-sPD1-SIVgag/SIVenv respectively. On day 70
after the second vaccination, ART treatment was interrupted in all macaques.
Samples were collected at different time points to monitor virological and
immunological parameters. (**B**) The representative images for the Gag or Env-

specific spots (2.5×10^5 cells per well) of each macaque pre-vaccination (before,
day 33) and post-vaccination (after, day 70) by ELISpot assay. (**C-D**) The
difference in SIV-specific IFN-γ-secreting cells (ΔSFCs) between pre-immunization
and post-immunization for the assessment of the immune response induced by
HSV-vectored SIV vaccines. (**E**) The difference in SIV-specific TNF-α/IFN-γ/IL-2-
secreting CD4$^+$ T and CD8$^+$ T subsets between pre-immunization and post-
immunization was detected by ICS assay.

Figure 5. The modified HSV-based constructs effectively reactivated SIV
latency in vivo in chronically SIV-infected, ART-treated macaques. (**A-D**) The
viral load (VL) change in plasma for each animal was monitored during the whole
experiment by y real-time PCR. The detection limit is 100 copies per mL plasma.
The shadow represented the duration of ART administration. (**E**) The VL change
in plasma between pre-ART and the peak value in the rebound stage after ART
discontinuation. (**F**) The change of total SIV DNA copies between pre-ART and
viral rebound after ART discontinuation. (**G**) The change of the SIV Pol-specific
IFN-γ-secreting cells between pre-immunization (day 33) and post-immunization
(day 70) was detected by ELISpot assay. (**H**) The change of the CD4$^+$ T/ CD8$^+$ T
ratio.
888
Figure 6. Pattern to illustrate the proof-of-concept strategy based on a
bifunctional HSV-vectored therapeutic vaccine for HIV functional cure. In the
present study, the modified HSV-ΔICP34.5-based constructs effectively
reactivated HIV/SIV latency by modulating the IKKα/β-NF-κB pathway and PP1-
HSF1 pathway (Shock) and simultaneously elicited antigen-specific polyfunctional
CD8$^+$ T cells to eliminate cells infected with the reactivated virion (Kill). BAC:
bacterial artificial chromosome; rHSV: recombinant HSV; TCR: T-cell receptor;
PD1: Programmed Cell Death Protein 1; CD40L: CD40 Ligand.
897

898   Figure 1

899
900

Figure 2

Figure 3

908   Figure 4

909
910

911   Figure 5

914   Figure 6

915