Effect of the SAS chromatin-remodeling complex on chromatin accessibility

during flower development

Jing Guo[a], Zhen-Zhen Liu[a], Yin-Na Su[a], Xin-Jian He[a,b*]

[a] National Institute of Biological Sciences, Beijing, 102206, China

[b] Tsinghua Institute of Multidisciplinary Biomedical Research, Tsinghua University,

100084, Beijing, China

[*] Corresponding author: Xin-Jian He (hexinjian@nibs.ac.cn)

Key words: chromatin remodeling, SWI/SNF, SAS, chromatin accessibility, flower

development, transcription

Running title: Effect of SAS on flower development

One-sentence summary: The SAS-type SWI/SNF complex establishes chromatin

accessibility at distal promoter and upstream intergenic regions of flower

development-related genes, thereby contributing to flower development.

Abstract

While the role of transcription factors in flower development is well understood, the

effect of chromatin remodeling on this process remains largely unclear. We conducted

a comprehensive analysis to investigate the genome-wide effects of the SAS, BAS,

and MAS-type SWI/SNF complexes on chromatin accessibility and gene transcription

in both seedlings and inflorescences of *Arabidopsis thaliana*. Our results indicate that

the SAS complex plays a crucial role in promoting the accessibility of genes involved

in flower development. Within inflorescences, the SAS complex binds to numerous

genes related to flower development and is responsible for establishing chromatin

accessibility at distal promoter and upstream intergenic regions of these genes. In

contrast ,the BAS and MAS complexes exhibit minimal involvement in regulating the

accessibility of these genes. Notably, the genomic regions bound by the SAS complex

are enriched with sites occupied by multiple MADS family transcription factors

associated with flower development. This study highlights the significant role of the

SAS complex in modulating chromatin accessibility during flower development,

providing insights into the dynamic regulation of chromatin remodeling during plant

development.

Introduction

Accurate gene expression in specific cells and developmental stages is crucial for the

development of organisms. ATP-dependent chromatin-remodeling complexes play a

vital role in facilitating the access of transcription factors and transcription machinery

to DNA by utilizing the energy derived from ATP hydrolysis, thereby driving dynamic

changes in gene expression. The SWI/SNF (Switch defective/Sucrose non

fermentable) complex, originally discovered in yeast, was the first identified

chromatin-remodeling complex (Cairns et al., 1994; Peterson et al., 1994). In

Drosophila melanogaster, SWI/SNF components were found to be Trithorax group

factors that counteract the repressive effect of Polycomb group proteins on the

transcription of specific homeotic genes, ensuring the accurate expression of these

homeotic genes at precise positions (Kennison and Tamkun, 1988; Tamkun et al.,

1992). Moreover, extensive studies in mammals have underscored the significance of

SWI/SNF complexes in maintaining pluripotency and driving cell fate determination

during development (Ho and Crabtree, 2010; Bieluszewski et al., 2023). Mutations in

SWI/SNF subunits in humans have been closely linked to various types of tumors and

developmental disorders (Cenik and Shilatifard, 2021). In Arabidopsis, the

SWI/SNF-type chromatin-remodeling ATPases consist of BRAHMA (BRM),

SPLAYED (SYD), and MINUSCULE1/2 (MINU1/2). Recent studies utilizing affinity

purification followed by mass spectrometry (AP-MS) analysis have identified three

classes of SWI/SNF complexes in Arabidopsis: the BRM-associated SWI/SNF (BAS)

complex, the SYD-associated SWI/SNF (SAS) complex, and the MINU1/2-associated

SWI/SNF (MAS) complex. Each of these complexes is composed of 8-15 subunits,

with some subunits being specific to a particular complex, while others are shared by

two or three complexes (Guo et al., 2022; Fu et al., 2023). However, the precise roles

of different Arabidopsis SWI/SNF complexes in specific developmental processes

remain elusive.

The effects of SWI/SNF components on plant growth and development have been

extensively studied. Single mutants of *brm* and *syd*, as well as the weak *minu1/2*

double mutant, display pleiotropic developmental defects, indicating the crucial role

of SWI/SNF complexes in plant development. Moreover, the combination of *brm* and

syd mutations, along with the *minu1/2* double knockout mutations, leads to lethality,

further emphasizing the importance of these complexes (Wagner and Meyerowitz,

2002; Farrona et al., 2004; Bezhani et al., 2007; Sang et al., 2012). Notably, the

mutants of BAS, SAS, and MAS components exhibit both overlapping and distinct

phenotypes throughout Arabidopsis development (Bezhani et al., 2007; Guo et al.,

2022). In particular, SAS-specific mutants, such as *syd*, *swi3d*, and *sys1/2/3*,

consistently show severe inflorescence developmental phenotypes, including defects

in shoot meristem maintenance, splayed-opened sepals, dramatic homeotic

transformation of floral organs, and complete sterility (Wagner and Meyerowitz, 2002;

Kwon et al., 2005; Sarnowski et al., 2005; Guo et al., 2022). The BAS mutant *brm*

also exhibits flower defects, including abnormal size and number of floral organs,

homeotic transformation of the second and third flower whorls, fused filaments, and

reduced fertility (Farrona et al., 2004; Hurtado et al., 2006). However, the floral

homeotic defects in *brm* are generally milder compared to SAS-specific mutants (Guo

et al., 2022). Additionally, other BAS-specific mutants, such as *swi3c*, *brip1/2*, and

brd1/2/13, show weaker or nearly normal flower and silique phenotypes in

comparison to *brm* (Archacki et al., 2009; Yu et al., 2020; Jaronczyk et al., 2021; Yu

et al., 2021; Guo et al., 2022). Although MAS-specific mutants, including *minu1/2*

and *pms2a/b*, have been reported to display abnormalities in the internal structures of

flowers, shorter siliques, and reduced fertility, their sepals and petals develop

normally (Sang et al., 2012; Diego-Martin et al., 2022; Guo et al., 2022). These

studies suggest that SAS plays a more important role in flower development

compared to BAS and MAS.

Flower development encompasses multiple stages, including floral induction, floral

meristem formation, and floral organ development, which is a strictly controlled

program regulated by a network of transcription factors, with the LEAFY (LFY)

transcription factor and various MADS-box transcription factors playing crucial roles

(Weigel, 1995). The genetic association between SWI/SNF components and flower

development-related transcription factors was established over two decades ago when

the *syd* mutation was found to enhance the floral homeotic defects in the weak *lfy*

mutant (Wagner and Meyerowitz, 2002; Bieluszewski et al., 2023). In *brm* and *syd*,

the expression of several floral homeotic genes is downregulated (Hurtado et al., 2006;

Wu et al., 2012; Lin et al., 2023). It has been reported that LFY and the MADS-box

transcription factor SEP3 recruit BRM and SYD to counteract Polycomb repression at

the floral homeotic genes *AP3* and *AGAMOUS* (*AG*) during flower development (Wu

et al., 2012). Additionally, the AP-MS results have demonstrated that BRM, SYD, and

certain other SWI/SNF subunits are co-purified with multiple MADS transcription

factors in flowers (Smaczniak et al., 2012), indicating interactions between SWI/SNF components and MADS transcription factors involved in flower development. A recent report has shown that LEAF AND FLOWER RELATED (LFR), a shared subunit of SAS and MAS, collaborates with the SAS catalytic subunit SYD to regulate the chromatin state and transcription of *AG* (Lin et al., 2023). These studies have mainly focused on the regulation of SWI/SNF components on specific genes associated with flower development. However, the genome-wide effects of SWI/SNF complexes on the transcription of flower development-related genes and the specific roles of the SAS-type SWI/SNF complex in flower development compared to the BAS and MAS complexes remain largely unclear.

In this study, we conducted a comprehensive genome-wide analysis to compare the impact of BAS, SAS, and MAS complexes on chromatin accessibility and gene expression between seedlings and inflorescences. Our findings revealed that SAS exerts a stronger effect on chromatin accessibility and gene transcription in inflorescences when compared to seedlings. Our examination of BAS, SAS, and MAS-bound genomic regions revealed that BAS and SAS exhibit a similar distribution pattern across the Arabidopsis genome and that the pattern is different from that of MAS. Notably, SAS is specifically recruited to a substantial subset of flower development-related genes in inflorescences and is responsible for establishing the chromatin accessibility of these genes. We observed a strong association between the recruitment of SAS and the presence of crucial transcription factors involved in flower development, such as SBP and MADS transcription factors. SAS is exclusively required for the accessibility of distal promoter regions and upstream intergenic regions of numerous genes involved in flower development, whereas BAS is dispensable for the accessibility of those regions. These results reveal the specific remodeling function of SAS in flower development and provide insights into the dynamic role of chromatin-remodeling complexes in plant development.

Results

SAS exhibits enhanced impact on gene expression in inflorescences relative to seedlings

134
135

We previously have determined the genome-wide effect of three classes of SWI/SNF complexes on chromatin accessibility and gene expression in the seedlings (Guo et al., 2022). However, whether and how the SWI/SNF complexes dynamically regulate chromatin accessibility and gene expression in different development processes are largely unclear. SAS mutants show several specific developmental phenotypes relative to BAS and MAS mutants (Guo et al., 2022), a prominent phenotype among which is the severe floral developmental defects (Supplemental Fig. S1A). In addition, anthocyanin accumulation was also observed in the inflorescence stem and rosette leaves of SAS mutants but not in BAS or MAS mutants (Supplemental Fig. S1, A and B). To investigate the dynamic role of SWI/SNF complexes in regulating chromatin accessibility and gene expression, we performed RNA deep sequencing (RNA-seq) in the inflorescences and subsequently compared the effects of SWI/SNF mutations on gene expression between the seedlings and the inflorescences. The SWI/SNF mutants used in the RNA-seq analysis contain: the BAS mutant *brm*; the SAS mutants *syd*, *swi3d*, and *sys1/2/3*; and the MAS mutants *minu1/2* and *pms2a/b*.

Multi-dimensional scaling (MDS) analyses of the RNA-seq data revealed a high reproducibility of three biological replicates for the inflorescences of each genotype (Fig. 1A). Based on the gene expression patterns, the mutants from the identical SWI/SNF complex are clustered (Fig. 1B), confirming the reliability of the RNA-seq data. We compared differentially expressed genes (DEGs) detected in the seedlings and inflorescences of SWI/SNF mutants relative to the wild type. Although the SAS mutants had the smaller number of DEGs compared to the BAS and MAS mutants in seedlings (Guo et al., 2022), a significant increase of DEGs was detected in the inflorescences of SAS mutants (Fig. 1C). Except for the weak *syd* mutant allele, the two other SAS mutants *swi3d* and *sys1/2/3* had more DEGs than the BAS and MAS mutants (Fig. 1C; Supplemental Fig. S1C; Supplemental Dataset 1), which is

consistent with the fact that the SAS mutants show more severe defects in the flower

development than the BAS and MAS mutants (Fig. 1D). This suggests that the SAS

complex plays a more important role in inflorescences than in seedlings. We

compared DEGs identified in the seedlings and inflorescences of each SWI/SNF

mutant, and found that only a small portion of DEGs overlap between seedlings and

inflorescences (Fig. 1, E-G; Supplemental Fig. S1, D-F), suggesting that the SWI/SNF

complexes have a dynamic role in regulating gene expression during development.

Comparison of SWI/SNF-dependent accessibility between inflorescences and
seedlings

Considering that the SWI/SNF complexes are mainly responsible for activating gene

expression by promoting chromatin accessibility (Clapier et al., 2017), and given that

the three classes of SWI/SNF complexes have been shown to facilitate genome-wide

chromatin accessibility in the seedlings (Guo et al., 2022), we determined chromatin

accessibility by performing the assay for transposase-accessible chromatin using

sequencing (ATAC-seq) and compared the effects of the SWI/SNF mutations on

chromatin accessibility between seedlings and inflorescences. On the basis of the

principal component analysis (PCA) of ATAC-seq data detected in the inflorescences,

the cluster of SAS mutants is more distant from the wild type than the BAS and MAS

mutants (Fig. 2A). Consistently, in the inflorescences, down-regulated differentially

accessible regions identified in the SAS mutants are more abundant than those in the

BAS and MAS mutants (Fig. 2B; Supplemental Dataset 2). By comparing the

down-regulated differentially accessible regions identified in each SWI/SNF mutant

between seedlings and inflorescences, we found that the inflorescence-specific

down-regulated differentially accessible regions are also more abundant in the SAS

mutants than those in the BAS and the MAS mutants (Fig. 2C). The accessibility

changes in inflorescences are strongly correlated among the mutants of the same

complex and are weakly correlated among the mutants of different complexes (Fig.

2D). Specially, the changes of accessibility in inflorescences are negatively correlated

between the SAS mutants and the MAS mutants (Fig. 2D), which is consistent with

our previous finding that SAS and MAS have antagonistic effects on regulating

chromatin accessibility in seedlings (Guo et al., 2022). These analyses suggest that the

three classes of SWI/SNF complexes regulate chromatin accessibility at different

genomic regions and that SAS complex contributes more to chromatin accessibility

than the other two types of SWI/SNF complexes in inflorescences.

Furthermore, we analyzed the distribution of the down-regulated differentially

accessible regions in both seedlings and inflorescences of each SWI/SNF mutant and

found that the distributions are overall identical between the seedlings and

inflorescences (Supplemental Fig. S2, A and B; Supplemental Dataset 2). While the

BAS and MAS-regulated accessible regions are enriched at the TSS-flanking regions,

the SAS-regulated accessible regions are enriched at distal promoter regions and

intergenic regions, which we hereafter refer to as distal accessible regions

(Supplemental Fig. S2, A and B). In both seedlings and inflorescences, the SAS

complex facilitates the accessibility at ~50% of distal accessible regions, whereas

BAS and MAS mediate the accessibility only at < 10% of them (Supplemental Fig.

S2C). In particular, the SAS complex contributes to chromatin accessibility at ~40%

of the super enhancer regions defined by a previous study in Arabidopsis (Zhao et al.,

2022), whereas the BAS and MAS complexes only show a marginal effect on the

accessibility of the super enhancer regions (Supplemental Fig. S2D). These results

point to the important role of SAS in promoting chromatin accessibility at distal

accessible regions in both seedlings and inflorescences.

The ATAC-seq analysis indicated that the genes with decreased accessibilities are

highly overlapped in different SAS mutants (Fig. 2E), suggesting the SAS

components function together in regulating chromatin accessibility. Although the BAS

and SAS complexes regulate chromatin accessibility primarily at different genomic

regions (Fig. 2D; Supplemental Fig. S2, A and B), a substantial subset (~40%) of

genes with down-regulated accessibility in the BAS mutant are shared by those with

down-regulated accessibility in the SAS mutant (Fig. 2F). Given that the BAS and

SAS can regulate the accessibility of the same set of target genes at proximal and

distal regions, respectively (Supplemental Fig. S2, A-C), the BAS and SAS

complexes are likely to collaborate in regulating the accessibility of different regions

at a subset of their shared target genes.

SAS regulates the accessibility and transcription of genes involved in flower

development

We conducted a gene set enrichment analysis (GSEA) to examine genes with

down-regulated accessibility in the inflorescences of the BAS, SAS, and MAS

mutants. We found that flower-specific expressed genes exhibit enrichment among the

genes with down-regulated accessibility in the SAS mutant, and to a lesser extent in

the BAS mutant, whereas we did not observe this enrichment in the MAS mutant (Fig.

2, G-I; Supplemental Fig. S3). Consistently, the Gene Ontology (GO) analysis

revealed a significant enrichment of flower development-related genes among those

exhibiting down-regulated accessibility in the BAS and SAS mutants, and such

enrichment was not observed in the MAS mutants (Supplemental Fig. S3;

Supplemental Dataset 3). Interestingly, the genes with down-regulated accessibility in

the MAS mutants showed enrichment in terms associated with protein targeting and

metabolic processes (Supplemental Fig. S3; Supplemental Dataset 3). These results

indicate that SAS and BAS complexes but not the MAS complex are responsible for

promoting the accessibility of genes involved in flower development.

Among the genes with down-regulated accessibility in the inflorescences of each

SWI/SNF mutant, the DEGs are most abundant in the SAS mutants, followed by the

BAS mutant and then the MAS mutants (Fig. 2J), confirming that the SAS complex

and to a lesser extent the BAS complex play a more important role in promoting

chromatin accessibility compared to the MAS complex. Moreover, the DEGs with

decreased expression were more abundant than those with increased expression in

each SWI/SNF mutant (Fig. 2J), supporting the notion that SWI/SNF-dependent

accessibility primarily functions in activating transcription rather than repressing it. It

is noteworthy that most of the genes with down-regulated accessibility in SWI/SNF

mutants did not show changes in expression (Fig. 2J), indicating that the

SWI/SNF-dependent accessibility contributes to establishing a priming chromatin

state for transcription but is not sufficient for actual transcription to occur. Taken

together, these results suggest that among the three classes of SWI/SNF complexes,

the SAS complex plays a major role in promoting chromatin accessibility and

facilitating transcriptional activation during flower development.

SAS is responsible for establishing the accessibility of genes involved in flower

development

The dynamic regulation of chromatin accessibility is involved in the regulation of

gene expression during flower development (Yan et al., 2019). To comprehensively

investigate the accessibility of genes involved in flower development at the

whole-genome level, we compared the chromatin accessibility between seedlings and

inflorescences based on our ATAC-seq results in the wild type. The ATAC-seq

analysis identified a total of 1,325 genes (corresponding to 1,842 peaks) with

increased accessibility and 1,344 genes (corresponding to 1,742 peaks) with decreased

accessibility in inflorescences compared to seedlings (Fig. 3A, Supplemental Dataset

4). The genes displaying increased accessibility in inflorescences were found to be

enriched with gene ontology (GO) terms associated with flower development

(Supplemental Fig. S4A). Conversely, the genes exhibiting decreased accessibility

were enriched with GO terms related to stress response (Supplemental Fig. S4B).

Next, we determined to what extent the SWI/SNF complexes contribute to the

accessibility of inflorescence-specific accessible genes, and found that > 50% of the

inflorescence-specific accessible genes were regulated by the SAS components, while

around 20% were regulated by the BAS or MAS components (Fig. 3B). In contrast,

the numbers of seedling-specific accessible genes regulated by BAS, SAS, and MAS

were not markedly different (Fig. 3C). Among the 1,325 inflorescence-specific

accessible genes, 47% (629/1325) of them are overlapped with the 1,778 genes with decreased accessibility in all the three tested SAS mutants (Fig. 3D). The GO analysis indicated that the 629 SAS-regulated, inflorescence-specific accessible genes (type I) are closely associated with flower development, whereas the remaining 696 inflorescence-specific accessible genes (type II) not regulated by SAS were enriched with the term 'maintenance of cell number', which is not directly related to flower development (Fig. 3E), confirming that SAS is specifically responsible for establishing the accessibility of flower development-related genes.

To characterize the accessible regions regulated by three classes of SWI/SNF complexes in both seedlings and inflorescences, we drew the average accessible signals at the 1,325 inflorescence-specific accessible genes and the 1,344 seedling-specific accessible genes in both wild type plants and SWI/SNF mutants. For the inflorescence-specific accessible genes, we observed not only a TSS-flanking accessible region but also a distal accessible region upstream of the TSS (Fig. 3F, Supplemental Fig. S5A). This characteristic aligns with the typical feature of SAS-regulated accessible genes (Guo et al., 2022). However, the seedling-specific accessible genes do not exhibit this distinct 'SAS feature' (Fig. 3G; Supplemental Fig. S5B). SAS is the primary regulator required for the accessibility of inflorescence-specific accessible genes, and that BAS, SAS, and MAS collectively contribute to the accessibility of seedling-specific genes (Fig. 3G; Supplemental Fig. S5B). It is worth noting that SAS weakly contributes to the accessibility of inflorescence-specific accessible genes in seedlings (Fig. 3F; Supplemental Fig. S5A). However, in wild type plants, the accessibility of these genes is markedly lower in seedlings than in inflorescences (Fig. 3F; Supplemental Fig. S5A). Therefore, SAS contributes to chromatin accessibility at the distal region of inflorescence-specific accessible genes.

Comparison of the binding patterns of BAS, SAS, and MAS in inflorescences and seedlings

To investigate the molecular mechanism underlying the different effects of SWI/SNF

complexes on chromatin accessibility between seedlings and inflorescences, we

conducted chromatin immunoprecipitation followed by sequencing (ChIP-seq) for the

BAS subunit SWI3C, the SAS subunit SWI3D, and the MAS subunit PMS2B. We

observed that the replicates of the same sample are more correlated than different

samples (Supplemental Fig. S6A), indicating the reliability of our ChIP-seq results.

The results indicated that the binding peaks of the BAS and SAS subunits in both

seedlings and inflorescences formed a cluster with strong correlation, while

displaying weak correlation with the binding peaks of the MAS subunit (Fig. 4A).

However, upon closer examination, we found that the binding peaks of BAS and SAS

subunits within the same developmental stage exhibited higher correlation than one

subunit across different developmental stages (Fig. 4A), suggesting that BAS and

SAS exhibited similar distribution patterns in either seedlings or inflorescences, and

their distributions showed differences between the two developmental stages.

Analysis of the binding peaks called from the ChIP-seq data revealed that the three

SWI/SNF complexes displayed a similar number of binding peaks between seedlings

and inflorescences (Fig. 4B; Supplemental Dataset 5). Previous studies have indicated

that, in seedlings, BAS and SAS share the majority of their binding peaks, while most

of the binding peaks of MAS are different from those of BAS and SAS (Guo et al.,

2022; Fu et al., 2023). The overlapping patterns of BAS, SAS, and MAS peaks in

inflorescences were similar to those in seedlings (Supplemental Fig. S6B). However,

whether and how the binding of SWI/SNF complexes to chromatin is altered during

plant development is largely unclear.

By comparing the binding peaks of each SWI/SNF subunit between seedlings and

inflorescences, we identified 1,023 BAS peaks, 1,415 SAS peaks, and 291 MAS

peaks exhibiting increased accumulation in inflorescences relative to seedlings, which

we termed inflorescence-specific binding peaks; and revealed 549 BAS peaks, 712

SAS peaks, and 679 MAS peaks with decreased accumulation in inflorescences

compared to seedlings, which we termed seedling-specific binding peaks (Fig. 4C;

Supplemental Fig. S6, C-E; Supplemental Dataset 6). Although the three SWI/SNF

complexes showed similar numbers of seedling-specific binding peaks, the number of

inflorescence-specific binding peaks was significantly higher for SAS and BAS

compared to MAS (Fig. 4B). This suggests that the binding of SAS and BAS to

chromatin undergoes dynamic establishment during flower development. We assigned

the binding peaks to nearby genes and then defined the binding genes for the three

SWI/SNF complexes (Fig. 4, B and C), and found that the BAS and SAS-binding

genes were significantly overlapped among either inflorescence- or seedlings-specific

binding genes, whereas only a small portion of MAS-binding genes were overlapped

with BAS or SAS-binding genes (Supplemental Fig. 6, F and G). This was

exemplified by the binding signals of BAS, SAS, and MAS in seedlings and

inflorescences at their random target genomic regions (Fig. 4D). These results suggest

that the binding of BAS and SAS to chromatin is altered during development at a

substantial subset of their target loci.

Furthermore, we determined whether the distribution of the binding peaks for each

SWI/SNF complex across the genic region is altered in inflorescences compared to

that in seedlings, and found that the overall distribution of total binding peaks for each

SWI/SNF complex is similar between inflorescences and seedlings (Supplemental Fig.

S6H). However, the proportion of the SAS and BAS-binding peaks in intergenic and

distal promoter regions exhibited a slight increase in inflorescences relative to

seedlings (Supplemental Fig. S6H). Upon comparing the distributions of

inflorescence-specific SAS-binding regions and seedling-specific SAS-binding

regions, we observed that SAS has a higher tendency to bind intergenic and distal

promoter regions in inflorescences than in seedlings (Fig. 4E). The GO analysis

revealed that the inflorescence-specific SAS-binding genes are primarily associated

with flower development, while the seedling-specific SAS-binding genes are enriched

in terms of stress response (Fig. 4, F and G). These results suggest that the SAS is

specifically recruited to a subset of genes involved in flower development during the

reproductive phase, rather than maintaining a constant association with these genes

between the vegetative and reproductive processes.

Recruitment of SAS establishes the accessibility of flower-development genes

To determine the impact of BAS, SAS, and MAS binding on gene accessibility in

inflorescences and seedlings, we examined the genes specifically bound by each

complex in either inflorescences or seedlings and compared them to genes that

exhibited decreased accessibility in the corresponding complex mutant. Interestingly,

we found that inflorescence-specific SAS-binding genes showed the highest

proportion of accessibility changes compared to the inflorescence-specific BAS- and

MAS-binding genes (Fig. 5A). It is worth noting that although the numbers of

inflorescence-specific binding genes were comparable for BAS and SAS, the

proportion of genes with down-regulated accessibility was significantly lower in the

BAS mutant (95/805) than in the SAS mutant (382/1,071). This discrepancy in

accessibility changes between the BAS and SAS mutants might be attributed to the

different remodeling activities of the two complexes at their target regions. Further

examination of SAS-specific binding regions in inflorescences revealed that the

regions with decreased accessibility exhibited higher binding and accessibility signals

compared to regions with maintained accessibility (Fig. 5, B and C). This suggests

that the enhanced binding affinity of SAS at specific binding regions in inflorescences

promotes the remodeling activity, leading to increased accessibility.

To identify inflorescence-specific accessible genes that are specifically bound and

remodeled by SAS, we conducted an overlap analysis between the 382 SAS-specific

binding genes with decreased accessibility in the SAS mutant inflorescences (Fig. 5A)

and the 1,325 inflorescence-specific accessible genes (Fig. 3A). Remarkably, we

found that 61% (234/382) of the genes specifically bound and remodeled by SAS in

inflorescences were inflorescence-specific accessible genes, accounting for 18%

(234/1,325) of the total inflorescence-specific accessible genes (Fig. 5D). This subset

includes crucial flower development-related genes, such as *LFY* (Schultz and Haughn,

1991), *SEP2/3/4* (Pelaz et al., 2000; Ditta et al., 2004), *SAUR*s (Stortenbeker and

Bemer, 2019), and *SPL*s (Chen et al., 2010) (Fig. 5D). Snapshots illustrating the

association between SWI/SNF binding and SWI/SNF-dependent accessibility at

representative genes are provided (Fig. 5E). These analyses highlight the contribution

of SAS recruitment to the establishment of accessibility for flower

development-related genes during flower development. Additionally, we discovered

that 16% (211/1,325) of the inflorescence-specific accessible genes exhibited

SAS-dependent accessibility but were bound by SAS in both inflorescences and

seedlings (Fig. 5E), suggesting that the involvement of SAS in inflorescence-specific

accessibility is also regulated after SAS is recruited to its target genes.

Furthermore, we investigated whether the SAS complex is involved in the expression

of flower development-related genes in the inflorescences. We found that the

accessibility of inflorescence-specific accessible regions is significantly lower in the

SAS mutant than in the wild type in the inflorescences (Supplemental Fig. S7A),

confirming that the SAS complex is crucial for facilitating the accessibility of the

flower development-related genes specifically in inflorescences. As determined by our

RNA-seq data, the expression levels of inflorescence-specific accessible genes are

significantly higher in inflorescences than in seedlings, and their expression levels are

reduced in the SAS mutant compared to the wild type in inflorescences (Supplemental

Fig. S7B). In the SAS mutant, the inflorescence-specific accessible genes with

decreased expression are more abundant than those with increased expression

(Supplemental Fig. S7C). These results suggest that the SAS complex contributes to

chromatin accessibility and thereby facilitates the transcription of flower

development-related genes. However, we also found that the expression levels of a

substantial subset of inflorescence-specific accessible genes are not significantly

reduced in the SAS mutant (Supplemental Fig. S7C). Therefore, the SAS-dependent

chromatin accessibility observed in inflorescences may result in transcription

occurring exclusively in specific cells and developmental stages, and these localized

transcriptional events may not be detectable in the RNA-seq analysis of mixed

inflorescence tissues.

SAS-dependent accessible regions and transcription factors

Similar to seedlings, where most SWI/SNF-regulated accessible regions are direct targets of SWI/SNF complexes (Guo et al., 2022), we observed that in the inflorescences of SWI/SNF mutants, the majority of accessible regions with decreased accessibility are also directly bound by their corresponding SWI/SNF complexes (Fig. 6A). Out of the 3,972 accessible regions with decreased accessibility detected in *swi3d* inflorescences, 60% of them are bound by SAS. Specifically, 14% of these regions are exclusively bound by SAS in inflorescences, while 46% are bound by SAS in both seedlings and inflorescences (Fig. 6B). Comparison of the inflorescence-specific SAS-binding regions with the shared SAS-binding regions in seedlings and inflorescences revealed that the inflorescence-specific regions exhibit higher SAS-binding signals and lower accessibility signals (Fig. 6, C and D). This observation aligns with the fact that distal promoter and upstream intergenic regions generally display lower chromatin accessibility compared to regions near the transcription start site, thus supporting the specific recruitment of the SAS complex to the distal promoter and upstream intergenic regions of flower development-related genes in inflorescences. It is important to note that the inflorescence-specific binding of SAS to its target loci only accounts for a small subset of SAS-dependent accessible regions (Fig. 6B). This further supports the notion that the regulation of SAS-dependent accessibility in inflorescences involves additional downstream steps following the binding of SAS to its target loci.

Previous studies have demonstrated that transcription factors either recruit SWI/SNF complexes to specific target genes or collaborate with SWI/SNF to regulate chromatin accessibility and transcriptional activation (Bieluszewski et al., 2023). In order to investigate how SAS-dependent chromatin accessibility is regulated during flower development, we examined whether the inflorescence-specific SAS-binding regions are enriched with any transcription factor-binding DNA motifs. Interestingly, we found that the inflorescence-specific SAS-binding regions are significantly enriched

with the DNA motifs bound by MADS and SBP transcription factors (Fig. 6E;

Supplemental Dataset 7). Given that the MADS and SBP transcription factors are

important for flower development (Chen et al., 2010; Theißen et al., 2016), it is likely

that these transcription factors either facilitate the recruitment of SAS to flower

development-related genes or collaborate with SAS to regulate the accessibility and/or

transcription of these genes. To further explore the relationship between SAS and the

MADS-family transcription factors, we compared the inflorescence-specific

SAS-binding regions with the MADS-binding regions previously identified in

inflorescences or seedlings (Chen et al., 2018; van Mourik et al., 2023). The analysis

indicated that the inflorescence-specific SAS-binding regions exhibited a high level of

overlap (15%~50%) with the regions bound by multiple MADS transcription factors

in inflorescences, whereas the overlap (<10%) was markedly reduced when they are

compared with the regions bound by the MADS transcription factors detected in

seedlings (Fig. 6F). These findings support the idea that MADS transcription factors

collaborate with the SAS complex to regulate the transcription of genes involved in

flower development.

Discussion

SWI/SNF complexes play crucial roles in the regulation of diverse developmental

processes in both metazoans and plants. Recent studies have illustrated the functional

variations among the three Arabidopsis SWI/SNF complexes: BAS, SAS, and MAS

(Guo et al., 2022; Fu et al., 2023). However, the precise mechanisms by which these

complexes exert specific functions during different developmental stages remain

largely unknown. In this study, we performed a comprehensive analysis to compare

the impact of BAS, SAS, and MAS on chromatin accessibility throughout the entire

genome between vegetative and reproductive development processes. Our results

demonstrate that while the three SWI/SNF complexes contribute to a comparable

amount of accessible regions in seedlings, there is a significantly higher proportion of

inflorescence-specific accessible regions that depend on SAS (Figure 7A). This

indicates that SAS exhibits a stronger influence on inflorescences compared to BAS

and MAS. The findings highlight the specific role of SAS in inflorescences, which is

consistent with the more severe defects in floral development observed in SAS

mutants relative to BAS and MAS mutants.

Our analysis, combining ATAC-seq with ChIP-seq data, implicates two mechanisms

through which the SAS complex contributes to establishing the accessibility of genes

involved in flower development. Firstly, SAS is recruited to genes involved in flower

development, thereby establishing the accessibility of these genes (Fig. 7B). In

Arabidopsis, LFY has been identified as a pioneer transcription factor that binds to

nucleosomal DNA and recruit SWI/SNF components to establish the accessibility of

genes responsible for determining the floral organ identity (Jin et al., 2021; Lai et al.,

2021). Certain MADS transcription factors, such as AP1 and SEP3, have also been

suggested as potential pioneer factors involved in flower development (Lai et al.,

2018). Given the demonstrated interaction between SYD and LFY or MADS

transcription factors (Wu et al., 2012), it is likely that multiple pioneering

transcription factors collaborate to recruit the SAS complex to different sets of flower

development-related genes. Secondly, the SAS complex exhibits constant binding to

genes involved in flower development during both the vegetative and reproductive

processes, with an enhanced chromatin remodeling activity specifically during the

reproductive process (Fig. 7B). The change in the chromatin remodeling activity of

SAS at different developmental stages can be attributed to either its differential

recognition ability to histone modifications or its interactions with different

transcription factors. Further studies are needed to investigate how histone

modifications and/or transcription factors regulate the recruitment of SAS to

chromatin or its chromatin remodeling activity during the reproductive process.

A previous study have indicated that establishing the accessibility of distal accessible

regions is critical for the activation of genes involved in flower development during

reproductive development (Yan et al., 2019). However, the specific mechanisms

responsible for establishing accessibility in the distal regions during the reproductive

process remain largely unclear. Our findings show that the SAS complex is recruited

to numerous genes involved in flower development during the reproductive process

and plays a crucial role in establishing their accessibility. While BAS and SAS target

similar genomic regions, SAS exhibits a greater ability to regulate the accessibility of

distal accessible regions. This suggests that the impact of SAS on chromatin

accessibility is not only regulated by its binding to chromatin but also by its chromatin

remodeling activity. Notably, distal accessible regions generally exhibit lower levels

of histone acetylation compared to proximal promoter regions (Yan et al., 2019),

indicating that differential histone modifications may regulate the remodeling

activities of BAS and SAS. These findings are consistent with studies conducted in

mammals, which have demonstrated the impact of histone modifications on the

genomic targeting and remodeling activity of various SWI/SNF complexes (Mashtalir

et al., 2021). In contrast to SAS, MAS primarily contributes to the accessibility of

genes unrelated to flower development, further highlighting the functional

specificities of different SWI/SNF complexes in Arabidopsis.

In mammals, there are three classes of SWI/SNF complexes: BAF, PBAF, and ncBAF.

Among these, the BAF complex has been shown to specifically target distal enhancer

regions, which play a crucial role in facilitating the transcription of genes involved in

development (Alver et al., 2017). Similarly, in Arabidopsis, distal accessible regions

have been identified as predictive markers for intergenic enhancers, exhibiting highly

stage-specific patterns during flower development (Zhu et al., 2015; Yan et al., 2019).

In this study, we present evidence that the SAS complex specifically contributes to the

accessibility of distal accessible regions for numerous genes involved in flower

development. This suggests that the SAS complex serves as the primary chromatin

remodeler responsible for establishing the stage-specific accessibility of genes

involved in flower development. Thus, it is likely that the SAS complex functions as

the counterpart of the human BAF complex in regulating the accessibility of distal

enhancer regions in genes related to flower development.

Transcription factors are known to have cell-type-specific functions, while chromatin remodeling complexes are generally considered to have broader regulatory roles. It is likely that the SAS complex collaborates with transcription factors to regulate gene transcription during reproductive development. In mammals, the abundance of chromatin remodelers compared to transcription factors has facilitated the identification of chromatin remodelers co-purified with transcription factors. However, the identification of transcription factors co-purified with chromatin remodelers has proven challenging (Gourisankar et al., 2023). Similar difficulties have been encountered in plants, where MADS transcription factors have been used as baits to successfully identify SWI/SNF components, while attempts to identify MADS transcription factors using SWI/SNF components as baits have been unsuccessful (Smaczniak et al., 2012; Guo et al., 2022). In a recent online study, the interaction between transcription factors and components of chromatin-remodeling complexes during stomatal development was identified using proximity labeling methods (Liu et al., 2023). In the future, investigating the specific interaction between SWI/SNF chromatin complexes and different transcription factors during flower development will contribute to our understanding of the molecular mechanisms underlying the specific functions of the SAS complex in flower development.

Materials and methods

Plant materials

Arabidopsis were in the Columbia-0 (Col-0) ecotype. SWI/SNF mutants related to this study include: *brm-1* (SALK_030046C), *syd* (CS822017), *swi3d* (SALK_100310), *minu1* (SALK_015562), *minu2* (SALK_057856), *pms2a* (SALK_141512), *pms2b* (SALK_010411), and *sys1/2/3* (generated by CRISPR-Cas9 system). Double mutants are generated by crossing. Detailed information on genotyping primers and CRISPR-Cas9 targets for SWI/SNF mutants can be found in the previous study (Guo et al., 2022). Full length genomic sequences of genes encoding SWI/SNF subunits driven by 1.5-2 kb native promoter were cloned into modified pCAMBIA-1305-GFP

for ChIP-seq or pCAMBIA-1305-3×Flag for AP-MS to generate C-terminal tagged

transgenes. The primers used for these constructions can be found in the previous

study (Guo et al., 2022). Arabidopsis seeds were sown on the Murahsige and Skoog

(MS) medium and stratified at 4 ℃ for 2 days. Subsequently, they were grown under

long-day conditions (16 h of light at 23 ℃ and 8 h of darkness at 22 ℃). Whole plants

of 12-day-old seedlings grown on MS medium and inflorescences of 4~5-week-old

plants in soil were used for RNA-seq, ATAC-seq, and ChIP-seq experiments.

RNA-seq and data analysis

About 0.1 gram of the inflorescence tips from the SWI/SNF mutants and the wild type

plants were harvested and ground in liquid nitrogen. For each genotype, three

independent samples were collected as biological replicates. RNA was isolated by

TRIzol reagent (Invitrogen, 15596018). Library construction and sequencing were

carried out by BGI (Wuhan, China) using the DNBSEQ platform (sequencing method:

SE50). The raw reads of RNA-seq were processed with the removal of adaptor

sequences and the filter of low-quality reads. The clean reads were then mapped to the

Arabidopsis genome using HISAT2 (version 2.2.1) (Kim et al., 2015) with default

parameters. After counting the reads mapped on the exon by featureCounts (version

2.0.1)(Liao et al., 2014), edgeR (version 3.32.1) (Robinson et al., 2010) was used to

evaluate the expression level, fold change and False Discovery Rate (FDR) of each

gene. DEGs were filtered based on the criteria of $|log2FC| \geq 1$ and $FDR < 0.05$.

ATAC-seq and data analysis

About 0.1-0.2 gram of the inflorescence tips from the SWI/SNF mutants and the wild

type plants were used for ATAC-seq. Two biological replicates were set for each

genotype. The ATAC-seq experiments were performed according to the previous

study (Guo et al., 2022). After extracting and assessing the quality of the nuclei, about

10000~50000 nuclei of each sample were collected for Tn5 tagmentation-based

library construction using TruePrep DNA library Prep Kit V2 for Illumina (Vazyme,

TD501). The ATAC-seq libraries were sent to Novogene and sequenced by an

Illumina-NovaSeq 6000 (sequencing method: PE150).

The raw data were processed with Trim Galore (version 0.6.6) (Krueger, 2015) to

remove adaptors and low-quality reads. Clean reads were then mapped to the

Arabidopsis genome using bowtie2 (version 2.4.2) (Langmead and Salzberg, 2012)

with the multi-mapped reads or PCR duplicates filtered out. The unique mapped reads

were converted to bigwig files by bamCoverage in Deeptools (version 3.5.1)(Ramírez

et al., 2016) with the parameter 'binSize 10' and 'normalizeUsing RPKM' and then

visualized by the Integrative Genomics Viewer (IGV) (version 2.4.13) for snapshots.

Accessible peaks were called using macs2 (version 2.2.7.1) (Zhang et al., 2008) with

the following parameter: macs2 callpeak -t sample.bam -f BAMPE -g 1.2e+8

-nomodel --bdg --keep-dup all. Only the peaks that present in both replicates

(irreproducible discovery rate < = 0.05) were retained for further analysis.

Differentially accessible regions between the two samples were identified using the

method 'DESEQ2' in DiffBind (version 3.0.15) (Rossinnes et al., 2011), with criteria

of |log2FC| ≥0.5 and FDR < 0.05. Peaks were assigned to genes with the same rules as

described in the previous study (Guo et al., 2022). Briefly, a peak that overlapped with

the gene body including 1 kb upstream region were annotated to the corresponding

gene, and if a peak was overlapped with more than one gene, it was annotated to the

gene harboring the closest TSS to the midpoint of the peak.

GSEA and GO analyses

The GSEA analysis was performed based on the gene sets generated from ATAC-seq

analysis in this study and the RNA-seq data of various Arabidopsis tissues from the

published study (Mergner et al., 2020). The expression level in flower and the average

expression level across different tissues of the total Arabidopsis genes were used as

the input expression dataset and the GSEA analysis was performed using GSEA

software (version 4.3.2) according to the manual (Subramanian et al., 2005). The

parameter 'Metric for ranking genes' was set as 'Diff_of_Classes'. The term 'flower

specificity' represents the expression level of a gene in the flower minus the average

expression level of that gene across all the tissues. The GO analysis was performed by

ClusterProfiler (version 4.2.2) (Yu et al., 2012) using the 'org.At.tair.db' database.

ChIP-seq and data analysis

ChIP-seq were performed following the protocol from the previous studies with minor

changes (Gendrel et al., 2005; Guo et al., 2022). Briefly, 4 grams of fresh seedlings or

the inflorescence tips were collected and cross-linked in the buffer containing 1%

formaldehyde under vacuum. The nuclei were extracted, washed, and sonicated by the

Bioruptor 2000 in the sonication buffer containing 0.33% SDS. Subsequently, the

fragmented chromatin was separated by centrifugation and the SDS concentration was

diluted to 0.1%. For each sample, 0.1% of the total volume was set aside as the input

sample, while the remaining diluted chromatin fragments were incubated overnight

with the GFP antibody (Abcam, ab290) at a 1:600 dilution. The Dynabeads Protein A

(Invitrogen, 100.01D) were added into the solution and incubated for 2-3 h, and then

washed sequentially with low-salt buffer, high-salt buffer, and LiCl buffer, followed

by two washes with TE buffer. The DNA-protein complexes were then eluted from

the beads and de-crosslinked at 65℃ overnight. After the digestion of RNA and

protein, the DNA was purified for library construction using NEBnext® Ultra^{TM} II

DNA Library Prep Kit for Illumina (E7654S). The ChIP-seq libraries were sent to

Novogene (Tianjin, China) for sequencing using an Illumina-NovaSeq 6000

(sequencing method: PE150).

After removal of adaptors from the raw reads, the clean reads were mapped to the

Arabidopsis genome using Bowtie2 (version 2.4.2) with the same mapping method as

for the ATAC-seq data. The binding peaks were called by macs2 (version 2.2.7.1) with

the following parameters: macs2 callpeak -c input.bam -t IP.bam -f BAMPE -g 1.2e+8.

Peaks were filtered and visualized using the same methods and standards as described

in the ATAC-seq data analysis. Differential binding peaks of SWI/SNF subunits

between seedlings and inflorescences were identified using the 'DESEQ2' method in

DiffBind (version 3.0.15), with criteria of |log2FC| ≥0.5 and FDR < 0.05. The TF

motif analysis was conducted by the findMotifsGenome.pl program in Homer

(version 4.11)(Heinz et al., 2010) with default parameters using different sets of

regions as input files.

Plotting methods

The MDS plot were generated using the plotMDS function of the R package edgeR

(version 3.32.1) with default parameters. The scatter plot was created using the

ggscatter function of the R package ggpubr (version 0.4.0)(Kassambara, 2020). The

PCA plot were generated using DiffBind (version 3.0.15) with default parameters.

The heatmap based on Pearson correlation coefficients was generated using the

heatmap.2 function of the R package gplots (version 3.1.1)(Warnes et al., 2014). Bar

charts and the pie chart were created using Graphpad Prism (version 8.0). Box plots

were generated using the boxplot function in R (version 4.1.0). The heatmap

representing the enrichment of TF motifs was generated using Pheatmap (version

1.0.12) (Kolde, 2012). Venn diagrams, butterfly bar plots, the volcano plot, and the

bubble plot containing multiple GO results of different samples were created using an

online   platform   for   data   analysis   and   visualization   at

https://www.bioinfomatics.com.cn. The bubble plot showing the overlap of TF

binding regions and the SAS-specific binding regions were plotted using ggplot2

(version 3.3.6).

Accession numbers

The accession numbers of genes reported in this study are as follows: AT2G46020

(*BRM*), AT2G28290 (*SYD*), AT4G34430 (*SWI3D*), AT5G07940 (*SYS1*), AT5G07970

(*SYS2*), AT5G07980 (*SYS3*), AT3G06010 (*MINU1*), AT5G19310 (*MINU2*),

AT3G08020 (*PMS2A*) and AT3G52100 (*PMS2B*).

Data availability

The raw data for ATAC-seq, ChIP-seq, and RNA-seq experiments have been deposited in the Gene Expression Omnibus (GEO) database under the accession numbers GSE254654, GSE254655, and GSE254656, respectively. The access tokens are as follows: inapmusabpcvvar for GSE254654, wbahscwojrotpqn for GSE254655, and kfqbwguylhudhmh for GSE254656.

Author contribution

JG and XJH conceived the project and designed the experiments. JG and ZZL performed the experiments. JG and YNS performed the bioinformatic analysis. JG and XJH wrote the manuscript.

Acknowledgements

This work was supported by the National Natural Science Foundation of China to X. J. He (32025003).

References

Alver, B.H., Kim, K.H., Lu, P., Wang, X., Manchester, H.E., Wang, W., Haswell, J.R., Park, P.J., and Roberts, C.W. (2017). The SWI/SNF chromatin remodelling complex is required for maintenance of lineage specific enhancers. Nat commun. **8,** 1-10.

Archacki, R., Sarnowski, T.J., Halibart-Puzio, J., Brzeska, K., Buszewicz, D., Prymakowska-Bosak, M., Koncz, C., and Jerzmanowski, A. (2009). Genetic analysis of functional redundancy of BRM ATPase and ATSWI3C subunits of Arabidopsis SWI/SNF chromatin remodelling complexes. Planta. **229,** 1281-1292.

Bezhani, S., Winter, C., Hershman, S., Wagner, J.D., Kennedy, J.F., Kwon, C.S., Pfluger, J., Su, Y., and Wagner, D. (2007). Unique, shared, and redundant roles for the Arabidopsis SWI/SNF chromatin remodeling ATPases BRAHMA and SPLAYED. Plant Cell **19,** 403-416.

Bieluszewski, T., Prakash, S., Roulé, T., and Wagner, D. (2023). The Role and

Activity of SWI/SNF Chromatin Remodelers. Annu Rev Plant Biol. **74,** 139-163.

Cairns, B.R., Kim, Y.-J., Sayre, M.H., Laurent, B.C., and Kornberg, R.D. (1994).

A multisubunit complex containing the SWI1/ADR6, SWI2/SNF2, SWI3, SNF5, and

SNF6 gene products isolated from yeast. Proc Natl Acad Sci U S A. **91,** 1950-1954.

Cenik, B.K., and Shilatifard, A. (2021). COMPASS and SWI/SNF complexes in

development and disease. Nat Rev Genet. **22**, 38-58

Chen, D., Yan, W., Fu, L.Y., and Kaufmann, K. (2018). Architecture of gene

regulatory networks controlling flower development in Arabidopsis thaliana. Nat

Commun. **9,** 4534.

Chen, X., Zhang, Z., Liu, D., Zhang, K., Li, A., and Mao, L. (2010). SQUAMOSA

promoter-binding protein-like transcription factors: star players for plant growth and

development. J Integr Plant Biol. **52,** 946-951.

Clapier, C.R., Iwasa, J., Cairns, B.R., and Peterson, C.L. (2017). Mechanisms of

action and regulation of ATP-dependent chromatin-remodelling complexes. Nat Rev

Mol Cell Biol. **18,** 407-422.

Diego-Martin, B., Pérez-Alemany, J., Candela-Ferre, J., Corbalán-Acedo, A.,

Pereyra, J., Alabadí, D., Jami-Alahmadi, Y., Wohlschlegel, J., and

Gallego-Bartolomé, J. (2022). The TRIPLE PHD FINGERS proteins are required for

SWI/SNF complex-mediated +1 nucleosome positioning and transcription start site

determination in Arabidopsis. Nucleic Acids Res. **50,** 10399-10417.

Ditta, G., Pinyopich, A., Robles, P., Pelaz, S., and Yanofsky, M.F. (2004). The

SEP4 gene of Arabidopsis thaliana functions in floral organ and meristem identity.

Curr Biol. **14,** 1935-1940.

Farrona, S., Hurtado, L., Bowman, J.L., and Reyes, J.C. (2004). The *Arabidopsis*

thaliana SNF2 homolog AtBRM controls shoot development and flowering.

Development. **131,** 4965-4975.

Fu, W., Yu, Y., Shu, J., Yu, Z., Zhong, Y., Zhu, T., Zhang, Z., Liang, Z., Cui, Y.,

Chen, C., et al (2023). Organization, genomic targeting, and assembly of three

distinct SWI/SNF chromatin remodeling complexes in Arabidopsis. Plant Cell. **35,**

2464-2483.

Gendrel, A.-V., Lippman, Z., Martienssen, R., and Colot, V. (2005). Profiling
histone modification patterns in plants using genomic tiling microarrays. Nat methods.
2, 213-218.

Gourisankar, S., Krokhotin, A., Wenderski, W., and Crabtree, G.R. (2023).
Context-specific functions of chromatin remodellers in development and disease. Nat
Rev Genet. Published online November 24, 2023. doi:10.1038/s41576-023-00666-x

Guo, J., Cai, G., Li, Y.Q., Zhang, Y.X., Su, Y.N., Yuan, D.Y., Zhang, Z.C., Liu,
Z.Z., Cai, X.W., Guo, J.,et al. (2022). Comprehensive characterization of three
classes of Arabidopsis SWI/SNF chromatin remodelling complexes. Nat Plants. **8,**
1423-1439.

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y.C., Laslo, P., Cheng, J.X.,
Murre, C., Singh, H., and Glass, C.K. (2010). Simple combinations of
lineage-determining transcription factors prime cis-regulatory elements required for
macrophage and B cell identities. Mol Cell. **38,** 576-589.

Ho, L., and Crabtree, G.R. (2010). Chromatin remodelling during development.
Nature. **463,** 474-484.

Hurtado, L., Farrona, S., and Reyes, J.C. (2006). The putative SWI/SNF complex
subunit BRAHMA activates flower homeotic genes in *Arabidopsis thaliana*. Plant
Mol Biol. **62,** 291-304.

Jaronczyk, K., Sosnowska, K., Zaborowski, A., Pupel, P., Bucholc, M., Malecka,
E., Siwirykow, N., Stachula, P., Iwanicka-Nowicka, R., and Koblowska, M. (2021).
Bromodomain-containing subunits BRD1, BRD2, and BRD13 are required for proper
functioning of SWI/SNF complexes in Arabidopsis. Plant Commun. **2,** 100174.

Jin, R., Klasfeld, S., Zhu, Y., Garcia, M.F., Xiao, J., Han, S.-K., Konkol, A., and
Wagner, D. (2021). LEAFY is a pioneer transcription factor and licenses cell
reprogramming to floral fate. Nat Commun. **12,** 1-14.

Kassambara, A. (2020). ggpubr:'ggplot2'based publication ready plots. R package
version 0.4. 0.

Kennison, J.A., and Tamkun, J.W. (1988). Dosage-dependent modifiers of
polycomb and antennapedia mutations in Drosophila. Proc Natl Acad Sci U S A. **85,**

8136-8140.

Kim, D., Langmead, B., and Salzberg, S.L. (2015). HISAT: a fast spliced aligner

with low memory requirements. Nat methods. **12,** 357-360.

Kolde, R. (2012). Pheatmap: pretty heatmaps. R package version 1.

Krueger, F. (2015). Trim galore. A wrapper tool around Cutadapt and FastQC to

consistently apply quality and adapter trimming to FastQ files **516**.

Kwon, C.S., Chen, C.B., and Wagner, D. (2005). *WUSCHEL* is a primary target for

transcriptional regulation by SPLAYED in dynamic control of stem cell fate in

Arabidopsis. Genes Dev. **19,** 992-1003.

Lai, X., Verhage, L., Hugouvieux, V., and Zubieta, C. (2018). Pioneer Factors in

Animals and Plants-Colonizing Chromatin for Gene Regulation. Molecules. **23,** 1914.

Lai, X., Blanc-Mathieu, R., GrandVuillemin, L., Huang, Y., Stigliani, A., Lucas,

J., Thévenon, E., Loue-Manifel, J., Turchi, L., Daher, H.,et al. (2021). The LEAFY

floral regulator displays pioneer transcription factor properties. Mol Plant. **14,**

829-837.

Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie

2. Nat Methods. **9,** 357.

Liao, Y., Smyth, G.K., and Shi, W. (2014). featureCounts: an efficient general

purpose program for assigning sequence reads to genomic features. Bioinformatics.

30, 923-930.

Lin, X., Yuan, T., Guo, H., Guo, Y., Yamaguchi, N., Wang, S., Zhang, D., Qi, D.,

Li, J., Chen, Q.,et al. (2023). The regulation of chromatin configuration at

AGAMOUS locus by LFR-SYD-containing complex is critical for reproductive organ

development in Arabidopsis. Plant J. **116,** 478-496.

Liu, A., Mair, A., Matos, J.L., Vollbrecht, M., Xu, S.L., and Bergmann, D.C.

(2023). Cell Fate Programming by Transcription Factors and Epigenetic Machinery in

Stomatal Development. bioRxiv. doi:10.1101/2023.08.23.554515

Mashtalir, N., Dao, H.T., Sankar, A., Liu, H., Corin, A.J., Bagert, J.D., Ge, E.J.,

D'Avino, A.R., Filipovski, M., and Michel, B.C. (2021). Chromatin landscape

signals differentially dictate the activities of mSWI/SNF family complexes. Science.

373, 306-315.

Mergner, J., Frejno, M., List, M., Papacek, M., Chen, X., Chaudhary, A., Samaras, P., Richter, S., Shikata, H., Messerer, M., Lang, et al. (2020). Mass-spectrometry-based draft of the Arabidopsis proteome. Nature. **579,** 409-414.

Pelaz, S., Ditta, G.S., Baumann, E., Wisman, E., and Yanofsky, M.F. (2000). B and C floral organ identity functions require *SEPALLATA* MADS-box genes. Nature. **405,** 200-203.

Peterson, C.L., Dingwall, A., and Scott, M.P. (1994). Five SWI/SNF gene products are components of a large multisubunit complex required for transcriptional enhancement. Proc Natl Acad Sci U S A. **91,** 2905-2908.

Ramírez, F., Ryan, D.P., Grüning, B., Bhardwaj, V., Kilpert, F., Richter, A.S., Heyne, S., Dündar, F., and Manke, T. (2016). deepTools2: a next generation web server for deep-sequencing data analysis. Nucleic Acids Res. **44,** W160-W165.

Robinson, M.D., McCarthy, D.J., and Smyth, G.K. (2010). edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. Bioinformatics. **26,** 139-140.

Rossinnes, C.S., Stark, R., Teschendorff, A.E., Holmes, K.A., Ali, H.R., Dunning, M.J., Brown, G.D., Gojis, O., Ellis, I.O., and Green, A.R. (2011). Differential oestrogen receptor binding is associated with clinical outcome in breast cancer. Nature. **481,** 389-393.

Sang, Y., Silva-Ortega, C.O., Wu, S., Yamaguchi, N., Wu, M.-F., Pfluger, J., Gillmor, C.S., Gallagher, K.L., and Wagner, D. (2012). Mutations in two non-canonical Arabidopsis SWI2/SNF2 chromatin remodeling ATPases cause embryogenesis and stem cell maintenance defects. Plant J. **72,** 1000-1014.

Sarnowski, T.J., Rios, G., Jasik, J., Swiezewski, S., Kaczanowski, S., Li, Y., Kwiatkowska, A., Pawlikowska, K., Kozbial, M., Kozbial, P., et al. (2005). SWI3 subunits of putative SWI/SNF chromatin-remodeling complexes play distinct roles during Arabidopsis development. Plant Cell. **17,** 2454-2472.

Schultz, E.A., and Haughn, G.W. (1991). *LEAFY*, a Homeotic Gene That Regulates Inflorescence Development in Arabidopsis. Plant Cell. **3,** 771-781.

Smaczniak, C., Immink, R.G., Muiño, J.M., Blanvillain, R., Busscher, M.,

Busscher-Lange, J., Dinh, Q.D., Liu, S., Westphal, A.H., Boeren, S., et al. (2012).

Characterization of MADS-domain transcription factor complexes in Arabidopsis

flower development. Proc Natl Acad Sci U S A. **109,** 1560-1565.

Stortenbeker, N., and Bemer, M. (2019). The *SAUR* gene family: the plant's toolbox

for adaptation of growth and development. J Exp Bot. **70,** 17-27.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette,

M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene

set enrichment analysis: a knowledge-based approach for interpreting genome-wide

expression profiles. Proc Natl Acad Sci U S A. **102,** 15545-15550.

Tamkun, J.W., Deuring, R., Scott, M.P., Kissinger, M., Pattatucci, A.M.,

Kaufman, T.C., and Kennison, J.A. (1992). brahma: a regulator of Drosophila

homeotic genes structurally related to the yeast transcriptional activator SNF2/SWI2.

Cell. **68,** 561-572.

Theißen, G., Melzer, R., and Rümpler, F. (2016). MADS-domain transcription

factors and the floral quartet model of flower development: linking plant development

and evolution. Development. **143,** 3259-3271.

van Mourik, H., Chen, P., Smaczniak, C., Boeren, S., Kaufmann, K., Bemer, M.,

Angenent, G.C., and Muino, J.M. (2023). Dual specificity and target gene selection

by the MADS-domain protein FRUITFULL. Nat Plants. **9,** 473-485.

Wagner, D., and Meyerowitz, E.M. (2002). SPLAYED, a novel SWI/SNF ATPase

homolog, controls reproductive development in Arabidopsis. Curr Biol. **12,** 85-94.

Warnes, G.R., Bolker, B., Bonebakker, L., Gentleman, R., Liaw, W.H.A., Lumley,

T., Maechler, M., Magnusson, A., Moeller, S., and Schwartz, M. (2014). Gplots:

various R programming tools for plotting data. R package version 2.

Weigel, D. (1995). The genetics of flower development: from floral induction to

ovule morphogenesis. Annu Rev Genet. **29,** 19-39.

Wu, M.-F., Sang, Y., Bezhani, S., Yamaguchi, N., Han, S.-K., Li, Z., Su, Y.,

Slewinski, T.L., and Wagner, D. (2012). SWI2/SNF2 chromatin remodeling ATPases

overcome polycomb repression and control floral organ identity with the LEAFY and

SEPALLATA3 transcription factors. Proc Natl Acad Sci U S A. **109,** 3576-3581.

Yan, W., Chen, D., Schumacher, J., Durantini, D., Engelhorn, J., Chen, M.,

Carles, C.C., and Kaufmann, K. (2019). Dynamic control of enhancer activity

drives stage-specific gene expression during flower morphogenesis. Nat Commun. **10,**

1-16.

Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). clusterProfiler: an R package for

comparing biological themes among gene clusters. Omics. **16,** 284-287.

Yu, Y., Liang, Z., Song, X., Fu, W., Xu, J., Lei, Y., Yuan, L., Ruan, J., Chen, C.,

and Fu, W. (2020). BRAHMA-interacting proteins BRIP1 and BRIP2 are core

subunits of Arabidopsis SWI/SNF complexes. Nat Plants**. 6,** 1-12.

Yu, Y., Fu, W., Xu, J., Lei, Y., Song, X., Liang, Z., Zhu, T., Liang, Y., Hao, Y., and

Yuan, L. (2021). Bromodomain-containing proteins BRD1, BRD2, and BRD13 are

core subunits of SWI/SNF complexes and vital for their genomic targeting in

Arabidopsis. Mol Plant. **14,** 888-904.

Zhang, Y., Liu, T., Meyer, C.A., Eeckhoute, J., Johnson, D.S., Bernstein, B.E.,

Nusbaum, C., Myers, R.M., Brown, M., and Li, W. (2008). Model-based analysis

of ChIP-Seq (MACS). Genome Biol. **9,** R137.

Zhao, H., Yang, M., Bishop, J., Teng, Y., Cao, Y., Beall, B.D., Li, S., Liu, T., Fang,

Q., Fang, C., et al. (2022). Identification and functional validation of super-enhancers

in Arabidopsis thaliana. Proc Natl Acad Sci U S A. **119,** e2215328119.

Zhu, B., Zhang, W., Zhang, T., Liu, B., and Jiang, J. (2015). Genome-Wide

Prediction and Validation of Intergenic Enhancers in Arabidopsis Using Open

Chromatin Signatures. Plant Cell. **27,** 2415-2426.

Figure legends

Figure 1. Comparison of DEGs identified by RNA-seq in SWI/SNF mutants between

inflorescences and seedlings. **A)** MDS plot of the inflorescence RNA-seq samples.

Clustering of BAS, SAS and MAS mutants are marked with green, blue, and red

circles, respectively. **B)** Heat map showing the pairwise Pearson correlation

coefficients based on the expression fold changes in inflorescences for the total DEGs

of the displayed SWI/SNF mutants. **C)** The number of DEGs in SWI/SNF mutants

relative to the wild type in both seedlings and inflorescences. **D)** The scatter plot

showing the correlation between the average petal number (Guo et al., 2022) and the

number of DEGs in inflorescences of the SWI/SNF mutants. A fitted regression line

and a 95% confidence interval band are indicated. R represents the Pearson

correlation coefficient. *P* value was determined by two-tailed *t* test. **E-G)** The overlap

of up-regulated genes and down-regulated DEGs in seedlings and inflorescences of

brm (**E**), *syd* (**F**) and *minu1/2* (**G**). The purple bar 'a' and 'b' represent genes shared

by other sets and genes specifically belonging to one set of the venn diagrams,

respectively.

Figure 2. Analysis of differentially accessible regions in inflorescences of SWI/SNF

mutants relative to the wild type. **A)** Principal component analysis (PCA) of the

inflorescence ATAC-seq data of the wild type and SWI/SNF mutants. The clusters of

BAS, SAS and MAS mutants are marked with green, blue, and red circles,

respectively. **B)** The number of up-regulated and down-regulated differentially

accessible regions in inflorescences of the SWI/SNF mutants identified by ATAC-seq.

The number of genes annotated by the differentially accessible regions is indicated in

brackets with brown font. DARs, differentially accessible regions. **C)** The proportion

and number of inflorescent-specific, seedling-specific and shared differentially

accessible regions among total differentially accessible regions identified in the

inflorescences and seedlings of each SWI/SNF mutant compared to the wild type.

DARs, differentially accessible regions. **D)** Heat map showing the pairwise Pearson

correlation coefficient based on the accessibility changes at the total genes with

changed accessibility in inflorescences of the displayed SWI/SNF mutants. **E)** Venn

diagram displaying the overlap of genes with down-regulated accessibility in

inflorescences of SAS mutants *syd*, *swi3d* and *sys1/2/3* compared to the wild type. **F)**

Venn diagram showing the overlap of genes with down-regulated accessibility in

inflorescences of *brm*, *syd* and *minu1/2* compared to the wild type. **G-I)** Line plots

depicting the enrichment of flower-specific expressed genes among the genes with

down-regulated accessibility in *brm* (**G**), *syd* (**H**) and *minu1/2* (**I**) as determined by

GSEA analysis. The grey regions represent the ranked list of the differences between

the expression in flowers and the average expression in all tissues of each *Arabidopsis*

gene. The normalized enrichment scores were labeled. *P* values were determined by

permutation. **J)** The proportion of genes showing up-regulated, down-regulated, and

unchanged expression among genes with down-regulated accessibility in the

inflorescences of the indicated SWI/SNF mutants relative to the wild type.

Figure 3. Effect of BAS, SAS, and MAS mutations on the accessibility of

seedling-specific and inflorescence-specific accessible regions. **A)** M (log ratio)

versus A (log average) (MA) plot showing the differential analysis results of

ATAC-seq peaks in inflorescences versus in seedlings of the wild type plants. The plot

shows the $\log_2$ (fold change) and $\log_2$ (average reads concentration) of the ATAC-seq

peaks. The red points represent the significant changed peaks with FDR<0.05 and

$|\log_2$ (fold change) $| \geq 0.5$. **B)** The percentage of genes with down-regulated

accessibility in the inflorescences of indicated SWI/SNF mutants among the

inflorescence-specific accessible genes. **C)** The percentage of genes with

down-regulated accessibility in the seedlings of indicated SWI/SNF mutants among

the seedling-specific accessible genes. **D)** Venn diagram showing the overlap of the

genes with consistently down-regulated accessibility in the inflorescences of SAS

mutants (*syd*, *swi3d*, and *sys1/2/3*) compared to the wild type and the

inflorescence-specific accessible genes. Gene set I represents the

inflorescence-specific accessible genes regulated by SAS, and gene set II represents

the inflorescence-specific accessible genes not regulated by SAS. **E)** Bubble plot

showing the enriched GO terms for gene set I and gene set II defined in D**.** The size of

the bubble represents the gene number, and the color of the bubble represents the

adjusted *P* value of the enriched term. **F, G)** Profile plots depicting the ATAC-seq

signals of the SWI/SNF mutants and the wild type plants in inflorescences and in

seedlings at the genic regions with inflorescence-specific accessibility (**F**) and the

genic regions with seedling-specific accessibility (**G**).

Figure 4. Specifically binding regions of BAS, SAS, and MAS in seedlings and

inflorescences. **A)** Heat map showing the pairwise Spearman correlation coefficients

based on the ChIP-seq read counts of the representative subunit of BAS (SWI3C),

SAS (SWI3D), and MAS (PMS2B) in seedlings (SDL) and inflorescences (IFL). **B)**

Peak number determined by ChIP-seq for the subunits of BAS, SAS and MAS in

seedlings and inflorescences. The number of genes annotated by the peaks is provided

in   brackets   with   brown   font.   **C)**   Number   of   seedling-specific   and

inflorescence-specific binding peaks for the BAS, SAS and MAS subunits. The

number of genes annotated by the peaks is provided in brackets with brown font. **D)**

Snapshot showing the ChIP-seq signals of BAS, SAS and MAS subunits in

inflorescences and seedlings at a genomic region. Blue peaks represent ChIP-seq

signal in seedlings, and pink peaks represent ChIP-seq signal in inflorescences. Peaks

marked with brown boxes indicate differential binding regions of BAS/SAS subunits

between seedlings and inflorescences. Peaks marked with purple boxes indicate

differential binding regions between the MAS subunit and the BAS/SAS subunits.

The scale of normalized reads is shown. **E)** The proportion of regions annotated to

specified chromatin features. The left two bars represent regions specifically bound by

SAS in seedlings and inflorescences. Random: 8,000 randomly selected genomic sites.

F, G) Bubble plots showing the top 10 enriched GO terms in genes specifically bound

by SAS in inflorescences (**F**) and in seedlings (**G**).

Figure 5. Specific binding of SAS in inflorescence contributes to establishing the

accessibility of flower development-related genes. **A)** The proportion of genes with

down-regulated accessibility and maintained accessibility in the corresponding

mutants among genes specifically bound by BAS, SAS and MAS in inflorescences

(IFL) and in seedlings (SDL). **B, C)** Boxplots showing the enrichment of SAS in

inflorescences (**B**) and the accessibility in the inflorescences of the wild type plants

(**C**) at the regions specifically bound by SAS in inflorescences with down-regulated

accessibility (green) or maintained accessibility (pink) in the SAS mutant. The center

lines and box edges of the box plots represents medians and the interquartile range

(IQR), respectively. Whiskers extent within 1.5 times the IQR. *P* values were

determined by two-tailed Mann-Whitney *U* test. **D)** Venn diagram showing the

overlap between three gene sets: (I) genes specifically bound by SAS in

inflorescences with decreased accessibility in the SAS mutant, (II) genes bound by

SAS in both seedlings and inflorescences with decreased accessibility in the SAS

mutant, and (III) inflorescence-specific accessible genes. Development-related genes

selected from the overlap between gene sets I and III are listed below the Venn

diagram. **E)** Snapshots displaying the ChIP-seq signal of the three SWI/SNF

complexes and the ATAC-seq signal of SWI/SNF mutants in inflorescences and in

seedlings at the genic region of genes involved in flower development. Blue

represents the signal in seedlings, and pink represents the signal in inflorescences.

Differential peaks of ChIP-seq and ATAC-seq in seedlings and inflorescences are

marked by shadows. The scale of normalized reads is shown in each snapshot.

Figure 6. Characterization of SAS-dependent accessible regions in inflorescences. **A)**

The proportion of the BAS/SAS/MAS bound and unbound regions among regions

with down-regulated accessibility in inflorescences of the core enzyme mutants of

BAS, SAS and MAS. **B)** Pie chart showing the proportion of the peaks not bound by

SAS (I), commonly bound by SAS in both seedlings and inflorescences (II) and

specifically bound by SAS in inflorescences (III) among the total down-regulated

peaks in the inflorescences of *swi3d* relative to the wild type. **C**, **D)** Boxplots showing

the enrichment of SWI3D (**C**) and the accessibility in the inflorescences of the

wild-type plants (**D**) at the regions I, II and III defined in Fig. 5F. The center lines and

box edges of the box plots represents medians and the interquartile range (IQR),

respectively. Whiskers extend within 1.5 times the IQR. *P* values were determined by

two-tailed Mann-Whitney *U* test. **E)** Heatmap exhibiting the different enrichment of

the different enrichment of various motifs at the indicated regions. **F)** Bubble plot

showing the ratio of the transcription factor (TF) binding peaks collected by the previous studies (Chen et al., 2018; van Mourik et al., 2023) among the 1,415 specific binding peaks of SWI3D. The binding peaks of TFs in inflorescences and seedlings are marked in purple and green, respectively.

Figure 7. The function of SAS complex in inflorescence-specific accessibility. **A)** Different contributions of BAS, SAS, and MAS to chromatin accessibility in seedlings and inflorescences. Ellipses represent three SWI/SNF complexes, with the size of the ellipse representing the extent of contribution to seedling- or inflorescence-specific accessibility by each SWI/SNF complex. **B)** Schematics showing the mechanisms of two subsets of inflorescence-specific accessible genes regulated by SAS. Grey shadows represent the binding of SAS to chromatin; pink peaks represent chromatin accessibility; blue arrows represent transcription. 'IFL-spe gene': inflorescence-specific accessible genes. Yellow hexagons represent transcription factors (TFs). Inflorescence-specific accessible genes are divided into two types based on the binding specificity of SAS in inflorescences compared to seedlings. For type I inflorescence-specific accessible genes, SAS binds to the distal regions of these genes specifically in the inflorescences, and this specific binding contributes to increased accessibility and transcription of these genes. MADS and SBP TFs are typically co-targeted with SAS. For type II inflorescence-specific accessible genes, SAS binds the cis-regulatory regions of these genes in both seedling and inflorescences, but exhibits enhanced remodeling activity in inflorescences. This enhanced activity results in increased accessibility and transcription of these genes.

Supplemental information

Supplemental Figure S1. Inflorescence phenotype and different gene expression analysis in SWI/SNF mutants.

Supplemental Figure S2. Proximal and distal accessible regions regulated by SWI/SNF complexes in seedlings and in inflorescences.

Supplemental Figure S3. GO enrichment of genes with down-regulated accessibility

in inflorescences of SWI/SNF mutants.

Supplemental Figure S4. GO enrichment of seedling-specific and

inflorescence-specific accessible genes.

Supplemental Figure S5. Effect of SWI/SNF subunit mutations on chromatin

accessibility at inflorescence-specific accessible genes and seedling-specific

accessible genes.

Supplemental Figure S6. Comparison of the binding regions of BAS, SAS and MAS

in seedlings and inflorescences.

Supplemental Figure S7. Effect of *swi3d* mutation on accessibility and expression of

inflorescence-specific accessible genes.

Supplemental Dataset 1. Differentially expressed genes identifed by RNA-seq in

SWI/SNF mutants relative to wild type in inflorescences.

Supplemental Dataset 2. Differentially accessible regions of SWI/SNF mutants

identified by ATAC-seq in inflorescences.

Supplemental Data set 3. GO analysis of genes with decreased accessibilily in

SWI/SNF mutants identified by ATAC-seq.

Supplemental Data set 4. Differentially accessible regions of wild-type plants

between inflorescences and seedlings.

Supplemental Dataset 5. ChIP-seq peaks of SWI/SNF subunits in inflorescences and

seedlings.

Supplemental Dataset 6. Differentially binding regions of SWI/SNF subunits

identified by ChIP-seq between inflorescence and seedlings.

Supplemental Data set 7. Transcription factor-binding motif analysis.

Figure 1. Comparison of DEGs identified by RNA-seq in SWI/SNF mutants between inflorescences and seedlings. **A)** MDS plot of the inflorescence RNA-seq samples. Clustering of BAS, SAS and MAS mutants are marked with green, blue, and red circles, respectively. **B)** Heat map showing the pairwise Pearson correlation coefficients based on the expression fold changes in inflorescences for the total DEGs of the displayed SWI/SNF mutants. **C)** The number of DEGs in SWI/SNF mutants relative to the wild type in both seedlings and inflorescences. **D)** The scatter plot showing the correlation between the average petal number (Guo et al., 2022) and the number of DEGs in inflorescences of the SWI/SNF mutants. A fitted regression line and a 95% confidence interval band are indicated. R represents the Pearson correlation coefficient. *P* value was determined by two-tailed *t* test. **E-G)** The overlap of up-regulated genes and down-regulated DEGs in seedlings and inflorescences of *brm* (**E**), *syd* (**F**) and *minu1/2* (**G**). The purple bar 'a' and 'b' represent genes shared by other sets and genes specifically belonging to one set of the venn diagrams, respectively.

Figure 2. Analysis of differentially accessible regions in inflorescences of SWI/SNF mutants relative to the wild type. **A)** Principal component analysis (PCA) of the inflorescence ATAC-seq data of the wild type and SWI/SNF mutants. The clusters of BAS, SAS and MAS mutants are marked with green, blue, and red circles, respectively. **B)** The number of up-regulated and down-regulated differentially accessible regions in inflorescences of the SWI/SNF mutants identified by ATAC-seq. The number of genes annotated by the differentially accessible regions is indicated in brackets with brown font. DARs, differentially accessible regions. **C)** The proportion and number of inflorescent-specific, seedling-specific and shared differentially accessible regions among total differentially accessible regions identified in the inflorescences and seedlings of each SWI/SNF mutant compared to the wild type. DARs, differentially accessible regions. **D)** Heat map showing the pairwise Pearson correlation coefficient based on the accessibility changes at the total genes with changed accessibility in inflorescences of the displayed SWI/SNF mutants. **E)** Venn diagram displaying the overlap of genes with down-regulated accessibility in inflorescences of SAS mutants *syd*, *swi3d* and *sys1/2/3* compared to the wild type. **F)** Venn diagram showing the overlap of genes with down-regulated accessibility in inflorescences of *brm*, *syd* and *minu1/2* compared to the wild type. **G-I)** Line plots depicting the enrichment of flower-specific expressed genes among the genes with down-regulated accessibility in *brm* (**G**), *syd* (**H**) and *minu1/2* (**I**) as determined by GSEA analysis. The grey regions represent the ranked list of the differences between the expression in flowers and the average expression in all tissues of each Arabidopsis gene. The normalized enrichment scores were labeled. *P* values were determined by permutation. **J)** The proportion of genes showing up-regulated, down-regulated, and unchanged expression among genes with down-regulated accessibility in the inflorescences of the indicated SWI/SNF mutants relative to the wild type.

Figure 3. Effect of BAS, SAS, and MAS mutations on the accessibility of seedling-specific and inflorescence-specific accessible regions. **A)** M (log ratio) versus A (log average) (MA) plot showing the differential analysis results of ATAC-seq peaks in inflorescences versus in seedlings of the wild type plants. The plot shows the $\log_2$ (fold change) and $\log_2$ (average reads concentration) of the ATAC-seq peaks. The red points represent the significant changed peaks with FDR<0.05 and |$\log_2$ (fold change) | ≥ 0.5. **B)** The percentage of genes with down-regulated accessibility in the inflorescences of indicated SWI/SNF mutants among the inflorescence-specific accessible genes. **C)** The percentage of genes with down-regulated accessibility in the seedlings of indicated SWI/SNF mutants among the seedling-specific accessible genes. **D)** Venn diagram showing the overlap of the genes with consistently down-regulated accessibility in the inflorescences of SAS mutants (*syd*, *swi3d*, and *sys1/2/3*) compared to the wild type and the inflorescence-specific accessible genes. Gene set I represents the inflorescence-specific accessible genes regulated by SAS, and gene set II represents the inflorescence-specific accessible genes not regulated by SAS. **E)** Bubble plot showing the enriched GO terms for gene set I and gene set II defined in (**D**). The size of the bubble represents the gene number, and the color of the bubble represents the adjusted *P* value of the enriched term. **F, G)** Profile plots depicting the ATAC-seq signals of the SWI/SNF mutants and the wild type plants in inflorescences and in seedlings at the genic regions with inflorescence-specific accessibility (**F**) and the genic regions with seedling-specific accessibility (**G**).

Figure 4. Specifically binding regions of BAS, SAS, and MAS in seedlings and inflorescences. **A)** Heat map showing the pairwise Spearman correlation coefficients based on the ChIP-seq read counts of the representative subunit of BAS (SWI3C), SAS (SWI3D), and MAS (PMS2B) in seedlings (SDL) and inflorescences (IFL). **B)** Peak number determined by ChIP-seq for the subunits of BAS, SAS and MAS in seedlings and inflorescences. The number of genes annotated by the peaks is provided in brackets with brown font. **C)** Number of seedling-specific and inflorescence-specific binding peaks for the BAS, SAS and MAS subunits. The number of genes annotated by the peaks is provided in brackets with brown font. **D)** Snapshot showing the ChIP-seq signals of BAS, SAS and MAS subunits in inflorescences and seedlings at a genomic region. Blue peaks represent ChIP-seq signal in seedlings, and pink peaks represent ChIP-seq signal in inflorescences. Peaks marked with brown boxes indicate differential binding regions of BAS/SAS subunits between seedlings and inflorescences. Peaks marked with purple boxes indicate differential binding regions between the MAS subunit and the BAS/SAS subunits. The scale of normalized reads is shown. **E)** The proportion of regions annotated to specified chromatin features. The left two bars represent regions specifically bound by SAS in seedlings and inflorescences. Random: 8,000 randomly selected genomic sites. **F, G)** Bubble plots showing the top 10 enriched GO terms in genes specifically bound by SAS in inflorescences (**F**) and in seedlings (**G**).

Figure 5. Specific binding of SAS in inflorescence contributes to establishing the accessibility of flower development-related genes. **A)** The proportion of genes with down-regulated accessibility and maintained accessibility in the corresponding mutants among genes specifically bound by BAS, SAS and MAS in inflorescences (IFL) and in seedlings (SDL). **B, C)** Boxplots showing the enrichment of SAS in inflorescences (**B**) and the accessibility in the inflorescences of the wild type plants (**C**) at the regions specifically bound by SAS in inflorescences with down-regulated accessibility (green) or maintained accessibility (pink) in the SAS mutant. The center lines and box edges of the box plots represents medians and the interquartile range (IQR), respectively. Whiskers extent within 1.5 times the IQR. *P* values were determined by two-tailed Mann-Whitney *U* test. **D)** Venn diagram showing the overlap between three gene sets: (I) genes specifically bound by SAS in inflorescences with decreased accessibility in the SAS mutant, (II) genes bound by SAS in both seedlings and inflorescences with decreased accessibility in the SAS mutant, and (III) inflorescence-specific accessible genes. Development-related genes selected from the overlap between gene sets I and III are listed below the Venn diagram. **E)** Snapshots displaying the ChIP-seq signal of the three SWI/SNF complexes and the ATAC-seq signal of SWI/SNF mutants in inflorescences and in seedlings at the genic region of genes involved in flower development. Blue represents the signal in seedlings, and pink represents the signal in inflorescences. Differential peaks of ChIP-seq and ATAC-seq in seedlings and inflorescences are marked by shadows.

Figure 6. Characterization of SAS-dependent accessible regions in inflorescences. **A)** The proportion of the BAS/SAS/MAS bound and unbound regions among regions with down-regulated accessibility in inflorescences of the core enzyme mutants of BAS, SAS and MAS. **B)** Pie chart showing the proportion of the peaks not bound by SAS (I), commonly bound by SAS in both seedlings and inflorescences (II) and specifically bound by SAS in inflorescences (III) among the total down-regulated peaks in the inflorescences of *swi3d* relative to the wild type. **C, D)** Boxplots showing the enrichment of SWI3D (**C**) and the accessibility in the inflorescences of the wild-type plants (**D**) at the regions I, II and III defined in Fig. 5F. The center lines and box edges of the box plots represents medians and the interquartile range (IQR), respectively. Whiskers extend within 1.5 times the IQR. *P* values were determined by two-tailed Mann-Whitney *U* test. **E)** Heatmap exhibiting the different enrichment of the different enrichment of various motifs at the indicated regions. **F)** Bubble plot showing the ratio of the transcription factor (TF) binding peaks collected by the previous studies (Chen et al., 2018; van Mourik et al., 2023) among the 1,415 specific binding peaks of SWI3D. The binding peaks of TFs in inflorescences and seedlings are marked in purple and green, respectively.

Figure 7. The function of SAS complex in inflorescence-specific accessibility. **A)** Different contributions of BAS, SAS, and MAS to chromatin accessibility in seedlings and inflorescences. Ellipses represent three SWI/SNF complexes, with the size of the ellipse representing the extent of contribution to seedling- or inflorescence-specific accessibility by each SWI/SNF complex. **B)** Schematics showing the mechanisms of two subsets of inflorescence-specific accessible genes regulated by SAS. Grey shadows represent the binding of SAS to chromatin; pink peaks represent chromatin accessibility; blue arrows represent transcription. 'IFL-spe gene': inflorescence-specific accessible genes. Yellow hexagons represent transcription factors (TFs). Inflorescence-specific accessible genes are divided into two types based on the binding specificity of SAS in inflorescences compared to seedlings. For type I inflorescence-specific accessible genes, SAS binds to the distal regions of these genes specifically in the inflorescences, and this specific binding contributes to increased accessibility and transcription of these genes. MADS and SBP TFs are typically co-targeted with SAS. For type II inflorescence-specific accessible genes, SAS binds the cis-regulatory regions of these genes in both seedling and inflorescences, but exhibits enhanced remodeling activity in inflorescences. This enhanced activity results in increased accessibility and transcription of these genes.

Supplemental Figure S1. Inflorescence phenotype and different gene expression analysis in SWI/SNF mutants. **A)** Photographs of inflorescences and flowers from 5~6-week-old SWI/SNF mutants. To show the inner stamens and pistil of a flower, some sepals and petals are removed in the second row of pictures. **B)** Images of rosette leaves from 8-week-old SWI/SNF mutants. To show the anthocyanin accumulation in rosette leaves, shoots and inflorescences were removed from the plant. **C)** The number of up-regulated and down-regulated DEGs in seedlings of inflorescences of SWI/SNF mutants, as determined by RNA-seq. **D-F)** Venn diagrams showing the overlap of up-regulated DEGs and down-regulated DEGs in seedlings and inflorescences of *swi3d* (**D**), *sys1/2/3* (**E**), and *pms2a/b* (**F**). The purple bar 'a' and 'b' represent genes shared by other sets and genes specifically belonging to one set of the venn diagrams, respectively.

Supplemental Figure S2. Proximal and distal accessible regions regulated by SWI/SNF complexes in seedlings and in inflorescences. **A)** The illustration of specified genomic features used for the peak annotation. ARs, accessible regions. **B)** Proportion of down-regulated differentially accessible regions annotated to specified features to the total down-regulated differentially accessible regions in seedlings or inflorescences of the indicated SWI/SNF mutants. The proportion of accessible regions annotated to specified genomic features to total accessible regions identified by ATAC-seq in the seedlings and inflorescences of wild type plants is also presented. Random: 8,000 random genomic sites. DARs, differentially accessible regions. **C)** Proportion of downregulated differentially accessible regions in each SWI/SNF mutant to the total accessible regions annotated to specified genomic features in seedlings and in inflorescences. **D)** The proportion of super enhancers including accessible regions regulated by SWI/SNF components in seedlings and in inflorescences to total super enhancers identified in a previous study (Zhao et al., 2022).

Supplemental Figure S3. GO enrichment of genes with down-regulated accessibility in inflorescences of SWI/SNF mutants. Bubble plots show the enriched GO terms in genes with down-regulated accessibility in inflorescences of SWI/SNF mutants.

Supplemental Figure S4. GO enrichment of seedling-specific and inflorescence-specific accessible genes. **A, B)** Bubble plots showing the top 20 enriched GO terms in specific accessible genes in inflorescences (**A**) and seedlings (**B**) of wild type plants.

Supplemental Figure S5. Effect of SWI/SNF subunit mutations on chromatin accessibility at inflorescence-specific accessible genes and seedling-specific accessible genes. **A, B)** Boxplots showing the chromatin accessibility levels in seedlings and inflorescences of the wild type and SWI/SNF mutants at the 400~1500 upstream region of TSS and at the -400~200 bp TSS-flanking region. Analysis was conducted separately for inflorescence-specific genes (**A**) and seedling-specific genes (**B**). The center lines and box edges of the box plots represents medians and the interquartile range (IQR), respectively. Whiskers extend within 1.5 times the IQR. *P* values were determined by two-tailed Wilcoxon signed rank test between the mutants and the wild-type control in each group.

Supplemental Figure S6. Comparison of the binding regions of BAS, SAS and MAS in seedlings and inflorescences. **A)** Heat map showing the pairwise Spearman correlation coefficients based on the ChIP-seq read counts of the two replicates of BAS (SWI3C), SAS (SWI3D), and MAS (PMS2B) subunits in seedlings or in inflorescences. **B)** Venn diagram illustrating the overlap of the genes bound by BAS, SAS and MAS in inflorescences. **C-E)** MA plots showing the up-regulated and down-regulated binding peaks of the BAS (**C**), SAS (**D**) and MAS (**E**) subunits in inflorescences compared to seedlings. The red points represent significant changed peaks with FDR<0.05 and |log$_2$ (fold change) | ≥ 0.5. **F, G)** Venn diagrams showing the overlap of genes specifically bound by BAS, SAS and MAS in inflorescences (**F**) and in seedlings (**G**). **H)** The proportion of regions bound by BAS, SAS and MAS annotated to specified chromatin features in seedlings and in inflorescences. Random: 8,000 random genomic sites.

Supplemental Figure S7. Effect of *swi3d* mutation on accessibility and expression of inflorescence-specific accessible genes. **A)** Boxplots showing the accessibility of inflorescence-specific accessible regions in seedlings and inflorescences of the wild type and *swi3d* mutant. **B)** Boxplots displaying the expression levels of inflorescence-specific accessible genes in seedlings and inflorescences of the wild type and *swi3d* mutant. The center lines and box edges of the box plots represent medians and the interquartile range (IQR), respectively. Whiskers extend within 1.5 times the IQR. *P* values were determined by two-tailed Wilcoxon signed rank test. **C)** Volcano plot showing the differential expression of inflorescence-specific accessible genes in the *swi3d* mutant compared to the wild type. The fold change and FDR were calculated using the inflorescence RNA-seq data. Genes that are up-regulated in *swi3d* (log$_2$(fold change)>0, FDR<0.05) are marked in red, genes that are down-regulated (log$_2$(fold change)<0, FDR<0.05) are marked in blue, and genes that are not significantly differentially expressed (FDR<0.05) are marked in grey. Representative points are marked with gene names.