Multi-gradient Permutation Survival Analysis Identifies Mitosis and Immune Signatures

Steadily Associated with Cancer Patient Prognosis

Xinlei Cai[1], Yi Ye[2], Xiaoping Liu[1], Zhaoyuan Fang[4,5], Luonan Chen[1,2,6], Fei Li[3*], Hongbin

Ji[1,2,6*]

[1]Key Laboratory of Systems Health Science of Zhejiang Province, School of Life Science,

Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences,

Hangzhou, 310024, China.

[2]State Key Laboratory of Cell Biology, Shanghai Institute of Biochemistry and Cell Biology,

Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences, Shanghai

200031, China.

[3]Department of Pathology and Frontier Innovation Center, School of Basic Medical Sciences,

Fudan University, Shanghai, China.

[4]Zhejiang University-University of Edinburgh Institute, Zhejiang University School of

Medicine, Haining , 314400, China

[5]The Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou,

310000, China

[6]School of Life Science and Technology, Shanghai Tech University, Shanghai 200120, China.

Running title: Mitosis and immune signatures in predicting cancer patient prognosis

*Corresponding author:

Hongbin Ji, State Key Laboratory of Cell Biology, Shanghai Institute of Biochemistry and

Cell Biology, Center for Excellence in Molecular Cell Science, Chinese Academy of

Sciences, 320 Yue Yang Road, Shanghai, 200031, China, Phone: 86-21-54921108, Fax: 86-

21-54921101, E-mail address: hbji@sibcb.ac.cn;

Fei Li, Department of Pathology and Frontier Innovation Center, School of Basic Medical

Sciences, Fudan University, 131 Dong An Road, Shanghai, 200032, China, E-mail address:

li_fei@fudan.edu.cn.

Abstract

The inconsistency of the association between genes and cancer prognosis is often attributed to many variables that contribute to patient survival. Whether there exist the Genes Steadily Associated with Prognosis (GEARs) and what their functions are remain largely elusive. We have developed a novel method called "Multi-gradient Permutation Survival Analysis " (MEMORY) to screen the GEARs using RNA-seq data from the TCGA database. Then we employed a network construction approach to identify hub genes from GEARs, and utilized them for cancer classification. In the case of LUAD, the GEARs were found to be related to mitosis. Our analysis suggested that LUAD cell lines carrying PIK3CA mutations exhibit increased drug resistance. For BRCA, the GEARs were related to immunity. The analysis revealed that CDH1 mutation might influence immune infiltration through the EMT process in BRCA. We further explored the prognostic relevance of mitosis and immunity through their respective scores. This study offers significant biological insights into GEARs and highlights their potential as robust prognostic indicators across diverse cancer types.

Keywords: Multi-gradient Permutation Survival Analysis (MEMORY), Gene Steadily Associated with Prognosis (GEAR), Hub Genes, Mitosis, Immune

Introduction

Cancer is one of the leading causes of death worldwide, with over 200 types identified. Despite of huge advancements in the field, accurately predicting cancer patient prognosis still remains a significant challenge[1,2]. Previous studies have highlighted the complexity of this task, which is influenced by various factors including cancer type, clinical stage, therapeutic interventions, nursing care, unexpected comorbidities, and other non-cancer related illnesses that may interplay[3-6]. Furthermore, the gene expression plays a crucial role in predicting cancer patient prognosis, such as *HER2*, *VEGF*, *Ki67*, etc[7-12]. Nevertheless, there is still a need for further improvement in prognostic accuracy to better inform treatment decisions and patient outcomes.

Survival analysis is commonly used to assess the correlation between genes and cancer patient prognosis[13]. However, inconsistent findings are frequently observed, even within the same type of cancer. For instance, studies exploring the correlation between *CCND1* and prognosis in NSCLC reported contrasting results, including positive correlation, inverse correlation, or negligible influence[14-16]. These discrepancies can be attributed to various influential factors, such as differences in sample size, cohort characteristics (including cancer subtypes and tumor staging) and variations in therapy approaches[17]. Such observations raise an important question: whether there exist Genes Steadily Associated with Prognosis (GEARs) that consistently correlate with patient outcomes across different conditions, particularly considering the varying sample sizes used in different studies? Affirmative answers to this question would have significant implication not only for the development of more accurate prognostic models but also for enhancing our understanding of cancer biology.

The Cancer Genome Atlas (TCGA) is a comprehensive cancer genomics project initiated in the United States. It consists of transcriptome data, genomic data, and clinical information pertaining to 33 different cancer types, making it the largest cancer clinical sample database currently available. In this study, we developed a novel method called "Multi-gradient Permutation Survival Analysis" (MEMORY) with the utilization of TCGA RNA-seq data, and accessed the potential existence of GEARs across 15 cancer types, each comprising a cohort

of over 200 patients. Furthermore, we also evaluated the prognostic predictive power of these

GEARs and explored their potential biological functions in driving cancer progression.

Results

Multi-gradient permutation survival analysis identifies GEARs associated with mitosis

and immune across multiple cancer types

GEARs are a group of genes consistently and significantly correlate with patient survival,

independent of the sample size. To identify these GEARs, we developed a novel method

called "Multi-gradient Permutation Survival Analysis " (MEMORY). This method allows us to

assess the correlation between a specific gene and cancer patient prognosis using available

transcriptomic dataset (**Figure 1A, Table 1**). Initially, we started with a sampling number at

10% of the cohort size initially, and gradually increased the sampling number with each 10%

increment, which was analyzed with 1000 permutations. By calculating the statistical

probability of each gene's association with patient survival, we can identify a group of

GEARs for further analyses.

In this study, we utilized the TCGA datasets for several reasons, including the diversity of

cancer types, the availability of gene expression profiles, and patient prognosis information.

We set a minimum cohort size of 200 and included 15 eligible cancer types for analysis.

These cancer types include bladder urothelial carcinoma (BLCA), breast invasive carcinoma

(BRCA), cervical squamous cell carcinoma and endocervical Adenocarcinoma (CESC), colon

Adenocarcinoma (COAD), kidney renal papillary cell carcinoma (KIRC), brain lower grade

glioma (LGG), liver hepatocellular carcinoma (LIHC), lung adenocarcinoma (LUAD), lung

squamous cell sarcinoma (LUSC), ovarian serous cystadenocarcinoma (OV), pancreatic

adenocarcinoma (PAAD), prostate adenocarcinoma (PRAD), stomach adenocarcinoma

(STAD), thyroid carcinoma (THCA), and uterine corpus endometrial carcinoma (UCEC). As

the number of samples increases, the survival probability of certain genes gradually

approaches 1. Once this score reaches 0.8 and remained consistent with further sample

gradient increase, the gene is considered a GEAR (**Figure 1B**). Remarkably, we successfully

identified a set of GEARs across all 15 cancer types (**Supplementary Figure 1-2 , Table 2**).

The GEAR counts in most cancer types ranged from 100 to1000, with exceptions of CESC,

KIRC, LGG, and PAAD with over 1000 GEARs, and THCA with only 22 GEARS (**Table 2**).

In LUAD, the top 10 GEARs with the highest significance probabilities were *TLE1*, *GNG7*,

ERO1A, *ANLN*, *DKK1*, *TMEM125*, *S100A16*, *KNL1*, *STEAP1*, and *BEX4*. Most of these

genes are known to promote LUAD malignant progression except for S100A16[18-26]. In other

cancer types, *BEX4* was identified as a common GEAR in KIRC, LGG, PAAD, and STAD

(Table 2). *BEX4* is reported as a proto-oncogene promoting cancer onset and malignant

progression in multiple cancers including LUAD, glioblastoma multiforme and oral squamous

cell carcinoma [25,27,28]. We also identified the most significant GEARs in individual cancer

types, such as *TLL1* (BLCA), *PGK1* (BRCA), *RFXANK* (CESC), *DPP7* (COAD), *VWA8*

(KIRC), *SCMH1* (LGG), *HILPDA* (LIHC), *TLE1* (LUAD), *CD151* (LUSC), *ANKRD13A*

(OV), *SOCS2* (PAAD), *DRG2* (PRAD), *ADAMTS6* (STAD), *PSMB8* (THCA), *ASS1* (UCEC)

(**Table 3).** Many of these genes were previously reported to associate with tumorigenesis[18,29-

40]. For example, TLE1 is known as a transcriptional repressor that promotes cell proliferation,

migration, and inhibits apoptosis in LUAD [41,42]. Additionally, *PGK1*, a key enzyme in the

glycolytic process, has been shown to promote cell proliferation, migration, and invasion in

multiple cancers[43,44]. These findings demonstrate the intricate link between the functionality

of GEARs and the initiation and progression of cancer.

To gain deep insights into the biological functions of these identified GEARs, we conducted

Gene Ontology Enrichment Analysis (GOEA) (**Figure 1C**). Interestingly, we found the

mitosis-related pathways were enriched in LICH, LUAD, LGG, and PAAD, and the immune-

related pathways were enriched in BRCA and UCEC[45]. Additionally, other cancer types

exhibited enrichment in various pathways, such as organic acid metabolism pathway in KIRC,

oxidative phosphorylation pathway in CESC, organ development-related pathways in LUSC,

neurogenesis-related pathways in BLCA, and sustenance metabolism pathways in THCA.

Given the crucial role of mitosis in cancer progression and the significance of the immune

system in cancer-host interactions, we specifically focused on the mitosis and immune-related

GERAs, which accounted for approximately 40% of all the analyzed cancers.

Identification of hub genes in mitosis and immune-related cancers

To further identify crucial genes within the GEARs, we conducted a significant survival

network (SSN) and extracted the higher-ranked edges to construct the core survival network

(CSN) (**Figure 2A, Table 4**). We created 6 CSNs with varying numbers of edges (250, 500,

1000, 2000, 4000, 8000). The top 10 GEARs with highest degree in these networks were

defined as hub genes [46]. We then classified the samples using the hub genes derived from

these networks and evaluated their predictive power for patient survival. We determined that

the optimal number of edges for the network ranged from 1000 to 2000, as it demonstrated

the best clustering effect of hub genes (**Figure 2B, Supplementary Figure 3A-O,**

Supplementary Figure 4A-O). Furthermore, we conducted GOEA on the hub genes selected

from the network with 1000 edges and found that the results were consistent with the GEAR

analysis (**Table 5**). Specifically, the hub genes in LIHC, LGG, and LUAD were enriched in

mitosis-related pathways, whereas the hub genes in BRCA and UCEC were enriched in

immune-related pathways (**Figure 2C**). For instance, the 9 hub genes in LUAD were

associated with functions related to mitosis, whereas the 8 hub genes of BRCA were

associated with immune-related functions (**Figure 2D-E**).

Furthermore, we conducted survival-dependent analyses on the hub genes of LGG, LIHC, and

LUAD. These analyses revealed that mitosis-related hub genes are closely associated with

cancer cell viability, especially those hub genes that are correlated with multiple cancer types

such as CDC20, TOP2A, BIRC5 and TPX2[47-50] (**Supplementary Figure 5A-C**). The

importance of these genes is well established. For instance, TPX2, a hub gene in all three

cancers, is known to play a crucial role in normal spindle assembly during mitosis and is

essential for cell proliferation[51]. The significance of these hub genes for the survival of cancer

cells suggests that the expression levels of these hub genes can be used to screen for inhibitors

of tumor cell growth. By integrating the Genomics of Drug Sensitivity in Cancer (GDSC) and

Connectivity Map (cMAP) databases, we identified 76 individual compounds that are able to

effectively suppress the expression of the 10 hub genes in LUAD cell lines (**Figure 2F,**

Supplementary Figure 5D-E). These compounds also significantly inhibit LUAD cell

growth and may serve as potential therapeutic agents for the treatment of LUAD.

We then analyzed the LUAD dataset, which had a larger sample size compared to LGG and

LIHC. Based on the expression of the hub genes, we classified the LUAD samples into three

subgroups: mitosis low (ML), mitosis medium (MM), and mitosis high (MH) (**Figure 2G,**

Table 6). Previous study has reported three classic subgroups of LUAD known as the

terminal respiratory unit (TRU), the proximal-proliferative (PP), and the proximal-

inflammatory (PI) subgroups[52]. Our analysis revealed that the ML subgroup was primarily

enriched with TRU subgroup and a small number of samples from the PP subgroup, but not

the PI subgroup. The MH subgroup showed high expression of mitosis-related genes and

predominantly encompassed PI and PP subgroups, whereas the MM group exhibited

intermediate levels of mitosis-related gene expression (**Figure 2G-H**). This suggests that the

new categorization method can help identify new factors that influence patient prognosis

(**Supplementary Figure 5F-G**).

Distinct genetic mutation landscapes characterize different clusters of LUAD

The expression of hub genes provided crucial indicators for distinguishing various subgroups

of LUAD. However, the underlying mechanisms driving these variations in expression

patterns remain unknown. In our subsequent research, we aimed to explore the genomic

differences, particularly in terms of gene mutations, among different LUAD subgroups.

Initially, we analyzed the variations of genetic mutation landscape of different LUAD

subgroups by assessing the tumor mutation burden (TMB). Interestingly, we found significant

differences in TMB among the various subgroups of LUAD (**Figure 3A**). Analysis of

common driver genes further revealed distinct proportions of tumors with ALK and ROS1

fusions in the favorable prognosis ML and MM subgroups, compared to tumors with

mutations in KRAS, EGFR, BRAF, and ERBB2 (**Figure 3B**). Furthermore, we observed

unique mutation characteristics in each of these 3 subgroups (**Figure 3C-E**). For example,

TP53 mutation, a prevalent tumor suppressor gene mutation, was frequently observed in the

MH subgroup with a mutation rate of 73%, compared to mutation rates of 14% in the ML

subgroup and 39% in the MM subgroup. Additionally, genes such as *CSMD3*, *RP1L1*, *ZFHX4*

exhibited similar trend in mutation frequency across the three subgroups. These findings

indicate substantial genomic differences among these three LUAD subgroups based on the

hub genes.

We identified gene mutations that showed significant changes through gene dependence

analysis (**Figure 4A**). To further the functional implications of these mutations, we enriched

them using a pathway system called Nested Systems in Tumors (NeST)[53]. The results

revealed notable differences in multiple functional pathways, including androgen receptor,

cell cycle, *TP53*, *EGFR*, IL-6, and *PIK3CA*-related pathways (**Figure 4B**). To gain further

insights into the functional implications of these different mutations, we assessed the gene

dependency of cells with these mutations using the DepMap database. Importantly, we

observed that the pathway differences between the MM and MH clusters were primarily

associated with *PIK3CA*-related pathways. Previous studies have reported that *PIK3CA*

mutation confers resistance in colorectal cancer, lung cancer, and breast cancer[54-56]. Therefore,

we aimed to validate the role of PIK3CA mutations in drug resistance using public databases.

We further analyzed A549 cells (a lung adenocarcinoma cell line with KRAS mutation) and

SW1573 cells (a lung adenocarcinoma cell line with both KRAS and PIK3CA mutations)

using their respective GDSC data and identified five potentially effective inhibitors (BMS-

345541, Dactinomycin, Epirubicin, Irinotecan, and Topotecan) from the previously screened

set of 76 LUAD cell growth inhibitors. Strikingly, SW1573 cells exhibited increased

resistance to all these 5 inhibitors when compared to A549 cells (**Figure 4C**). In line with

this, clinical data also supports the notion that concurrent *PIK3CA* mutation is a poor

prognostic factor for LUAD patients[57]. These findings suggest that *PIK3CA* mutation may

contribute to drug resistance in LUAD.

Distinct clusters of BRCA exhibit different immune infiltration landscapes

We further focused on the immune-related cancer types, with BRCA standing out as a

representative immune-related cancer. This prompted us to conduct a more comprehensive

analysis (**Figure 1C**). We hierarchically grouped the BRCA subgroups into three distinct

clusters based on the expression of hub genes: Immune low (IL), Immune medium (IM), and

Immune high (IH) (**Figure 5A, Table 7**). Interestingly, all the classifications based on the

widely used PAM50 classification contained samples from the IL, IM, and IH subgroups,

although there were minor differences in proportions[58] (**Figure 5B, Supplementary Figure**

6A-B). In basal-type breast cancer, which generally has a poor prognosis, a set of samples

exhibited high immune infiltration and consequently better outcomes. Conversely, within the

HER2 subgroups, known for a better prognosis, there were samples with poor outcomes.

Integrating our method with traditional classification may enable a more detailed stratification

of breast cancer samples. Mutation analysis revealed distinct patterns of gene mutations

among the three subgroups, with significant differences in mutation frequencies of genes such

as *TP53*, *CDH1*, and *LRP2* (**Figure 5C-E**). We conducted a comparison of the proportions of

immune cells across these three subgroups and observed a strong association between the

overall immune cell proportion and the clustering results based on hub genes (**Figure 5F,**

Supplementary Figure 6C-L). Specifically, the IH subgroup exhibited significantly higher

proportions of CD8$^+$ T cells and Treg cells, with mean percentages at 5.0% and 6.0%,

respectively, compared to 1.4% and 3.2% in the IM subgroup, and 0.3% and 1.6% in the IL

subgroup. These findings suggest the presence of distinct immune responses in these three

subgroups, indicating a potential association between genetic mutation pattern and the

immune microenvironment landscape.

Next, we explored the specific genomic factors influencing immune infiltration in BRCA.

Mutation analysis revealed that *CDH1* gene mutation ranked as the second most prevalent

genetic alteration in BRCA samples, following TP53 mutation (**Figure 5C-E**). Previous study

has indicated that *CDH1* is involved in mechanisms regulating cell-cell adhesions, mobility,

and proliferation of epithelial cells[59]. We found that *CDH1*-mutant samples exhibited

significantly lower *CDH1* expression compared to *CDH1*-wildtype samples, and *CDH1*

expression showed a close correlation with immune cell infiltration (**Figure 5G,**

Supplementary Figure 6M). These results were consistent with clinical observations of

CDH1 mutation and high immune infiltration in invasive lobular carcinoma of the breast [60].

We further investigated how *CDH1* mutation influenced immune infiltration. In BRCA, we

observed a significant association between the expression of *CDH1* and the expression of

EMT marker genes *VIM* and *TWIST2* (**Supplementary Figure 6N-O**)[61,62]. This suggests that

the regulation of *CDH1* is intricately linked with EMT process. Through calculating the EMT

score, we observed a positive correlation between the EMT score and the proportion of

immune infiltration (**Figure 5H**). Consequently, *CDH1* might influence immune infiltration

through the EMT process.

Mitotic and immune signatures predict patient prognosis at pan-cancer level

We further sought to explore the prognostic relevance of mitosis and immune by calculating

their scores and assessing their correlation with patient outcomes at the pan-cancer level. The

scores for these biological processes were computed using RNA-seq data from the TCGA

database, and the median score was used as a cut-off to categorize patients into different

groups. Out of the 33 cancer types analyzed, the prognosis of 17 cancer types showed a

significant correlation with at least one of these two pathways (**Figure 6**). Specifically, 10

cancer types were exclusively associated with the mitosis score, 2 cancer types were

exclusively associated with the immune score, and 5 cancer types exhibited a correlation with

both mitosis and immune scores simultaneously. Overall, the identification of mitosis and

immune-related biological processes as significant prognostic factors at the pan-cancer level

suggests their potential utility as valuable biomarkers for predicting patient prognosis.

Discussion

Due to the multifaceted nature of variables affecting cancer patient prognosis, it remains

uncertain whether there exist a set of genes steadily associated with cancer prognosis ,

regardless of sample size and other factors. Here, we utilized the MEMORY method to

address this question and discovered that all the cancer types have GEARs. We observed

significant variation in the number of GEARs among different cancer types, indicative of

cancer type-specific patterns. The substantial heterogeneity in driver gene and mortality rates

among various cancer types could potentially explain this phenomenon. For example, THCA,

known for its low malignancy, displays the fewest genetic expression alterations among all

the studied cancer types. This observation could be correlated with the relatively high five-

year survival rate in THCA, which exceeds 50% even in advanced stage[63]. In contrast, PRAD,

despite of a favorable prognosis similar to THCA, exhibits a significantly higher number of

genetic expression alterations. We hypothesize that the positive prognosis in PRAD cases

might be largely attributed to early diagnosis, which enables timely treatment for the majority

of PRAD patients [64]. This discrepancy between the number of genetic alterations and

prognosis in THCA and PRAD highlights the intricate nature of cancer genetics and

emphasizes the importance of personalized considerations in cancer treatment and prognosis

evaluation. Furthermore, certain genes are commonly found in the GEARs of various cancer

types, indicating their potential significance in cancer development. For instance, BEX4 is

present in the GEARs of LUAD, KIRC, LGG, PAAD, and STAD, and known to play

oncogenic role in inducing carcinogenic aneuploid transformation via modulating the

acetylation of α-tubulin[65,66]. Aneuploidy is a hallmark characteristic of cancer and can lead to

alterations in the dosage of oncogenes or tumor suppressor genes, thereby influencing tumor

initiation and progression [67-70]. The regulation of aneuploid transformation by BEX4 may

represent a common mechanism through which this gene impacts prognosis across different

cancer types. Subsequently, we discovered that GEARs across different cancers displayed

distinct functional characteristics. Notably, a recurrent theme was the prominence of mitosis-

related and immune-related features. Specifically, the GEARs of LGG, LIHC, and LUAD

were enriched in mitosis-related pathways, whereas BRCA and UCEC showed the

enrichment of immune-related pathways. Mitosis and immune processes have been widely

recognized as having a significant impact on patient prognosis[71-75]. However, current clinical

guidelines do not yet recommend the use of transcriptomic data to assess scores related to

mitotic and immune pathways for predicting patient outcomes, despite the well-established

association between these pathways and prognosis in cancer[76-81]. Cancers enriched with

mitosis pathways often exhibit heterogeneous tumor growth kinetics across individuals, with

tumor size being one of the crucial factors influencing patient survival [82-84]. For instance, in

the case of LGG, the growth rate of tumors directly impacts the patient's prognosis due to the

primary location in the brain. Consequently, alterations in the expression levels of genes

related to mitosis are often indicative of the prognosis of LGG [85]. Cancer types that are

enriched in immune-related pathways, such as BRCA and UCEC, are closely associated with

deficiencies in DNA mismatch repair (MMR) [86,87]. Cancers enriched in immune-related

pathways often exhibit heterogeneous tumor growth kinetics across individuals, where tumor

size is closely correlated with patient survival.

To explore the underlying differences between samples associated with distinct mitotic and

immune-related pathways, we employed GEAR analysis to identify hub genes for further molecular classification and mutation analysis. Specifically, we focused on LUAD and BRCA, two representative cancer types exhibiting enriched mitosis and immune signatures in GEARs. By classifying the hub gene, we divided LUAD into three subgroups: ML, MM and MH, which displayed significant difference in survival outcomes. Subsequently, we utilized NeST to analyze the mutations within these subgroups. Notably, there were significant differences observed in cell cycle and signal transduction-related pathways among ML, MM and MH subgroups. Of particular importance, the *PIK3CA*-related pathway emerged as a key differentiating pathway between the MH and MM subgroups. Our analyses suggest that LUAD cell line carrying *PIK3CA* mutation may exhibit increased drug resistance. This aligns with previous studies demonstrating that targeting the PI3K pathway can overcome drug resistance[88,89]. Of course, future research is warranted to elucidate the exact functional role of *PIK3CA* mutations in LUAD.

To investigate the factors influencing immune infiltration in BRCA, we classified the samples based on hub genes and analyzed the differentially mutated genes. Interestingly, we found that *CDH1* mutation occurred at a high frequency in the IH subgroups. This led us to speculate that *CDH1* plays a crucial role in the regulation of immune infiltration in BRCA. Furthermore, previous studies have reported that *CDH1* inactivation promotes immune infiltration in breast cancer[60]. *CDH1* is a vital gene associated with EMT, and various studies have demonstrated the close relationship between EMT and the tumor immune microenvironment in different cancers[90]. Consistently, our results also supported the correlation between EMT score and immune infiltration in BRCA. These findings suggest that the mutations in *PIK3CA* and *CDH1*, identified through GEAR analysis, have significant impacts on cancer development and hold potential value in improving clinical therapies. This further emphasizes the importance of GEARs in understanding cancer biology and guiding treatment strategies.

Lastly, we investigated the prognostic predictive capabilities of the mitosis score and immune score at the pan-cancer level. Surprisingly, we found that approximately half of the cancer types exhibited significant correlations between these two scores and patient prognosis.

Interestingly, even for cancers originating from the same primary location, their correlations

with these scores could differ, indicating the potential diversity of mechanisms underlying

cancer-related mortality. For instance, both LGG and GBM are brain tumors, but the primary

risk factor for patient prognosis in LGG is closely linked to tumor diameter, whereas the main

risk factor for GBM patients lies in its high invasiveness and challenge of surgical

resection[91,92].

Despite of the strong association between GEAR and patient prognosis, the edges of the CSN

constructed based on GEARs was undirected. As a result, the hub genes identified from

GEAR in the CSN may primarily serve as stable and effective biological markers.

Additionally, through multi-omics analysis, we obtained some functional mutations, but the

therapeutic significance of these mutations remains to be elucidated. For example, further

study is needed to understand the exact role of PIK3CA mutations in promoting tumor cell

proliferation and drug resistance. Similarly, the association of *CDH1* mutations with the

infiltration of multiple immune cell types also warrants additional experimental investigation.

In our future studies, we plan to utilize protein-protein interaction networks and pathway

databases to construct a novel network based on the CSN. This approach will allow us to

directly screen genes from GEARs that could potentially serve as therapeutic targets.

Undoubtedly, future efforts are still required to utilize protein-protein interaction networks

and pathway databases to construct a new network based on CSN.

In conclusion, our study utilized the MEMORY algorithm to identify GEARs in 15 cancer

types, highlighting the significance of mitosis and immunity in cancer prognosis. Our findings

demonstrate that GEARs possess substantial biological significance beyond their role as

prognostic biomarkers. This study provides valuable guidance for establishing standards for

survival analysis evaluation and holds potential for the development of novel therapeutic

strategies.

Methods

Datasets

The gene expression profiles and clinical information of 33 cancers were obtained from the TCGA database and downloaded by the GDC data website (https: //portal.gdc.cancer.gov/). All gene expression data and survival data were integrated and normalized.

Multi-gradient permutation survival analysis

Gene expression data of cancer patients were obtained from the RNA-seq expression matrix of TCGA after TPM normalization. We randomly sampled the gene expression data according to the gradient. The sampling strategy was as follows: Ten gradient increases in sample size were pre-set, ranging from approximately 10% to about 100%, with intervals of 10%. Random samples were taken from total samples of each cancer 1000 times, based on each pre-set sample size. Multiple sampling matrix was obtained after the sampling strategy was performed in each gradient. The survival analysis was performed for every sampling matrix by "survival" and "survminer". "survival" and "survminer" are R packages for survival analysis. The survival analysis method was as follows according to the median expression value of every gene, every sampling matrix was divided into a high and low expression group and performed survival analysis by "survival" and "survminer". We used 1 for significant survival analysis results ($P < 0.05$) and 0 for non-significant results ($P > 0.05$). The survival analysis matrices were obtained after these processes.

Construction of core survival network

The survival analysis matrices were integrated to a significant-probability matrix by a formula:

$$A_{ij} = \frac{\sum_{n=1}^{1000} k_{j \times i \times n}}{1000} \qquad (1)$$

where the A_{ij} is the value from row i and column j in the significant-probability matrix. We defined the sampling size k_j reached saturation when the max value of column j was equal to 1 in a significant-probability matrix. The least value of k_j was selected, and the genes with their corresponding A_{ij} greater than 0.8 were extracted as GEAR.

We also defined survival analysis similarity (*SAS*) as the similarity of the effect on patient

prognosis in two genes. The *SAS* computational formula is as follows:

$$SAS = \frac{A \cap B}{A \cup B - 2(A \cap B) + 1} \qquad (2)$$

Where *A* and *B* are results of *1000* survival analyses at the first sampling gradient, which are

sequences of *0* and *1* of length *1000*. $A \cap B$ means the number of events that are

simultaneously 1 in both A and B. $A \cup B$ means the size of the union of *A* and *B*. The *SAS* of

significant genes with total genes was calculated, and the significant survival network was

constructed. Then some *SAS* values at the top of the rankings were extracted, and the *SAS* was

visualized to a network by Cytoscape[93]. The network was named core survival network

(CSN). The degree of each node was calculated, and the nodes with the top10 degrees were

defined hub genes. The effect of molecular classification by hub genes is indicated that 1000

to 2000 was a range that the result of molecular classification was best. One thousand was

chosen as the number of *SAS*. Therefore, one thousand was selected as the number of *SAS*

values for constructed significant survival network.

Gene ontology enrichment analysis

Gene ontology has been used to classify genes based on functions. The gene functions were

divided three types, includes molecular function (MF), biological process (BP), and cellular

component (CC). ClusterProfiler is an R package for gene set enrichment analysis[94]. The gene

ontology enrichment analysis (GOEA) was processed in significant gene sets and hub genes

by ClusterProfiler.

Hub gene classification

Tumor samples were genotyped. First, the expression matrix of hub genes corresponding to

cancer types was extracted from RNA-seq data. The matrix M is log-transformed after adding

a pseudo count of 1: $M' = log_2(M + 1)$. Finally, M' was grouped by hierarchical clustering.

Calculation of tumor mutation burden (TMB) and differential mutation

The TMB and differential mutation gene analysis were carried out to explore the differences

between different genotyped samples at the genomic level. The data are from the gene

mutation data of TCGA tumor samples, and the grouping information is from the hub gene

hierarchical clustering results. Maftools is an R package for the analysis of somatic variant

data, which can export results in form of charts and graph[95]. The calculation of TMB and

difference mutation analysis were used by the maftools.

Quantifying the effect of gene mutations for tumor cell viability

Gene dependence refers to the extent to which genes are essential for cell proliferation and

survival. The Cancer Dependency Map (Depmap, https://depmap.org/portal/) database

provides genome-wide gene dependence data for large number of tumor cell lines[96]. Gene

mutation data of cell lines were obtained from the CCLE database

(https://sites.broadinstitute.org/ccle). The genes appearing in the gene mutation data of cell

lines were extracted and sorted into mutation list. Each gene in the list was then analyzed for

survival-dependent differences. For each gene in the mutation list, we compared the gene

dependence score (S) between cell lines with and without the gene mutation. Specifically, cell

lines from the CCLE mutation dataset harboring a mutation in a given gene were classified

into one group, while cell lines without the mutation constituted the control group. S for the

mutation (S_m) and wild-type (S_{wt}) cell lines were then extracted from the DepMap database.

A two-sample t-test was used to assess the statistical significance of differences between S_m

and S_{wt}. The test yielded a P-value, with an alpha level set at 0.05, to determine the

significance of the difference in means. D_m was defined as gene mutation dependence, the

computational formula was as follows:

466
$$D_m = |\bar{s}_m - \bar{s}_{wt}| \qquad (3)$$

467   Subsequently, we conducted a standardization assessment of D_m, employing the following

468   formula:

469
$$S_{dm} = \frac{2(\bar{s}_m - \bar{s}_{wt})}{\bar{s}_m + \bar{s}_{wt}} \qquad (4)$$

In this study, we categorized mutations identified in cell lines into three groups based on their

impact on gene dependency. Functional mutations refer to the mutations that exhibit

significant changes in gene dependency scores, with a P-value < 0.05 and $S_{dm} > 0.1$. Non-

functional mutations were those without significant changes in gene dependency scores

compared to wildtype. Subtype-associated functional mutations were a subset of functional

mutations that show statistically significant differences in occurrence frequency across

different LUAD subtypes.

Identification of drugs downregulating hub gene expression

The IC50 data of 76 compounds of LUAD cell lines were obtained from Genomics of Drug

Sensitivity in Cancer (GDSC) database[97]. The hub gene expression data of A549 cell lines

after treatment with 76 compounds was obtained from Connectivity MAP (cMAP,

https://clue.io/) database, and the data were grouped by hierarchical clustering[98]. The drugs

which were presented only in the hub gene expression suppression group were considered to

effective against the LUAD cell.

Immune infiltration analysis

485 Immune infiltration data was calculated by quantiseq method[99]. Immunedeconv was an R

486 package for unified access to computational methods for estimating immune Cell fractions

487 from bulk RNA-seq data[100]. The data were from the gene expression data of TCGA tumor

488 samples after TPM was standardized. Then the immune cell fractions of tumor samples were

489 calculated by quantiseq method invoking immunedeconv.

Biological function score calculation

491 GSVA is an R package for calculate biological function score based on a single sample[101].

492 The data was from TCGA in the calculation process, and the software package was GSVA.

493 The gene set of the immune score was from other study (**Table 8**) [99]. The gene sets that

494 calculated the mitosis score were obtained from gene ontology (GO:0140014).

Acknowledgment

496 This work was supported by the National Key Research and Development Program of China

497 (grants 2022YFA1103900 to H.J., F.L.; 2020YFA0803300 to H.J.); the National Natural

498 Science Foundation of China (grants 82341002 to H.J., 32293192 to H.J., 82030083 to H.J.,

82173340 to L.H., 82273400 to Y.J., 32100593 to X.T., 82203306 to W.H., 82372763 to X.W., 82303916 to C.G., 82303039 to Z.Q., 82141101 to F.L., 2022hwyq16 to F.L., 82372794 to F.L.); the Basic Frontier Scientific Research Program of Chinese Academy of Science (ZDBS-LY-SM006 to H.J.); the Innovative research team of high-level local universities in Shanghai (SSMU-ZLCX20180500 to H.J.); Science and Technology Commission of Shanghai Municipality (21ZR1470300 to L.H.).

Author Contributions

H.J. and F.L. conceived the idea and designed the analysis. X.C. performed all bioinformatics analyses. Y.Y., X.L. and L.C. provided technical assistance and helpful comments. H.J., F.L. and X.C. wrote the manuscript. All authors approved the final version.

Disclosure of conflicts of interest

The authors declare that there is no conflict of interests.

References

1. Tomczak, K., Czerwinska, P., and Wiznerowicz, M. (2015). The Cancer Genome Atlas (TCGA): an immeasurable source of knowledge. Contemp Oncol (Pozn) *19*, A68-77. 10.5114/wo.2014.47136.
2. Cancer Genome Atlas Research, N., Weinstein, J.N., Collisson, E.A., Mills, G.B., Shaw, K.R., Ozenberger, B.A., Ellrott, K., Shmulevich, I., Sander, C., and Stuart, J.M. (2013). The Cancer Genome Atlas Pan-Cancer analysis project. Nat Genet *45*, 1113-1120. 10.1038/ng.2764.
3. Liu, J., Lichtenberg, T., Hoadley, K.A., Poisson, L.M., Lazar, A.J., Cherniack, A.D., Kovatich, A.J., Benz, C.C., Levine, D.A., Lee, A.V., et al. (2018). An Integrated TCGA Pan-Cancer Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. Cell *173*, 400-416 e411. 10.1016/j.cell.2018.02.052.
4. Smeltzer, M.P., Faris, N.R., Ray, M.A., and Osarogiagbon, R.U. (2018). Association of Pathologic Nodal Staging Quality With Survival Among Patients With Non-Small Cell Lung Cancer After Resection With Curative Intent. JAMA Oncol *4*, 80-87. 10.1001/jamaoncol.2017.2993.
5. Miyauchi, E., Morita, S., Nakamura, A., Hosomi, Y., Watanabe, K., Ikeda, S., Seike, M., Fujita, Y., Minato, K., Ko, R., et al. (2022). Updated Analysis of NEJ009: Gefitinib-Alone Versus Gefitinib Plus Chemotherapy for Non-Small-Cell Lung Cancer With Mutated EGFR. J Clin Oncol, JCO2102911. 10.1200/JCO.21.02911.

6.   Teh, S.H., Uong, S., Lin, T.Y., Shiraga, S., Li, Y., Gong, I.Y., Herrinton, L.J., and Li, R.A.
(2021). Clinical Outcomes Following Regionalization of Gastric Cancer Care in a US
Integrated Health Care System. J Clin Oncol *39*, 3364-3376. 10.1200/JCO.21.00480.

7.   Gimotty, P.A., Van Belle, P., Elder, D.E., Murry, T., Montone, K.T., Xu, X., Hotz, S., Raines,
S., Ming, M.E., Wahl, P., and Guerry, D. (2005). Biologic and prognostic significance of
dermal Ki67 expression, mitoses, and tumorigenicity in thin invasive cutaneous melanoma. J
Clin Oncol *23*, 8048-8056. 10.1200/JCO.2005.02.0735.

8.   Wu, Y., Lv, M., Qian, T., and Shen, Y. (2021). Correlation analysis of Ki67 and CK7
expression with clinical characteristics and prognosis of postoperative cervical
adenocarcinoma patients. Ann Palliat Med *10*, 9544-9552. 10.21037/apm-21-1974.

9.   Mouabbi, J.A., Singareeka Raghavendra, A., Bassett, R.L., Jr., Hassan, A., Tripathy, D., and
Layman, R.M. (2023). Survival Outcomes in Patients With Hormone Receptor-Positive
Metastatic Breast Cancer With Low or No ERBB2 Expression Treated With Targeted
Therapies Plus Endocrine Therapy. JAMA Netw Open *6*, e2313017.
10.1001/jamanetworkopen.2023.13017.

10.   Kang, Y.J., Oh, S.J., Bae, S.Y., Kim, E.K., Lee, Y.J., Park, E.H., Jeong, J., Park, H.K., Suh,
Y.J., and Kim, Y.S. (2023). Predictive biological factors for late survival in patients with
HER2-positive breast cancer. Sci Rep *13*, 11008. 10.1038/s41598-023-38200-y.

11.   Cai, C., Wang, X., Fu, Q., and Chen, A. (2022). The VEGF expression associated with
prognosis in patients with intrahepatic cholangiocarcinoma: a systematic review and meta-
analysis. World J Surg Oncol *20*, 40. 10.1186/s12957-022-02511-7.

12.   Zhang, C., Wang, L., Xiong, C., Zhao, R., Liang, H., and Luo, X. (2021). The role of vascular
endothelial growth factor as a prognostic and clinicopathological marker in osteosarcoma: a
systematic review and meta-analysis. J Orthop Surg Res *16*, 738. 10.1186/s13018-021-02888-
3.

13.   Lai, Y., OuYang, G., Sheng, L., Zhang, Y., Lai, B., and Zhou, M. (2021). Novel prognostic
genes and subclasses of acute myeloid leukemia revealed by survival analysis of gene
expression data. BMC Med Genomics *14*, 39. 10.1186/s12920-021-00888-0.

14.   Anton, R.C., Coffey, D.M., Gondo, M.M., Stephenson, M.A., Brown, R.W., and Cagle, P.T.
(2000). The expression of cyclins D1 and E in predicting short-term survival in squamous cell
carcinoma of the lung. Mod Pathol *13*, 1167-1172. 10.1038/modpathol.3880215.

15.   Esposito, V., Baldi, A., De Luca, A., Tonini, G., Vincenzi, B., Santini, D., Persichetti, P.,
Mancini, A., Citro, G., Baldi, F., et al. (2005). Cell cycle related proteins as prognostic
parameters in radically resected non-small cell lung cancer. J Clin Pathol *58*, 734-739.
10.1136/jcp.2004.023531.

16.   Dworakowska, D., Jassem, E., Jassem, J., Boltze, C., Wiedorn, K.H., Dworakowski, R.,
Skokowski, J., Jaskiewicz, K., and Czestochowska, E. (2005). Prognostic value of cyclin D1
overexpression in correlation with pRb and p53 status in non-small cell lung cancer (NSCLC).
J Cancer Res Clin Oncol *131*, 479-485. 10.1007/s00432-004-0661-9.

17.   Freedman, L.S. (1982). Tables of the number of patients required in clinical trials using the
logrank test. Stat Med *1*, 121-129. 10.1002/sim.4780010204.

18.   Yao, X., Ireland, S.K., Pham, T., Temple, B., Chen, R., Raj, M.H., and Biliran, H. (2014).
TLE1 promotes EMT in A549 lung cancer cells through suppression of E-cadherin. Biochem
Biophys Res Commun *455*, 277-284. 10.1016/j.bbrc.2014.11.007.

575 19. Zheng, H., Tian, H., Yu, X., Ren, P., and Yang, Q. (2021). G protein gamma 7 suppresses
576   progression of lung adenocarcinoma by inhibiting E2F transcription factor 1. Int J Biol
577   Macromol *182*, 858-865. 10.1016/j.ijbiomac.2021.04.082.

578 20. Chen, G., Wang, Q., and Wang, K. (2022). MicroRNA-218-5p affects lung adenocarcinoma
579   progression through targeting endoplasmic reticulum oxidoreductase 1 alpha. Bioengineered
580   *13*, 10061-10070. 10.1080/21655979.2022.2063537.

581 21. Xu, J., Zheng, H., Yuan, S., Zhou, B., Zhao, W., Pan, Y., and Qi, D. (2019). Overexpression of
582   ANLN in lung adenocarcinoma is associated with metastasis. Thorac Cancer *10*, 1702-1709.
583   10.1111/1759-7714.13135.

584 22. Zhang, P., Li, S., Lv, C., Si, J., Xiong, Y., Ding, L., Ma, Y., and Yang, Y. (2018). BPI-9016M, a
585   c-Met inhibitor, suppresses tumor cell growth, migration and invasion of lung adenocarcinoma
586   via miR203-DKK1. Theranostics *8*, 5890-5902. 10.7150/thno.27667.

587 23. Wang, T., Zhai, R., Lv, X., Wang, K., and Xu, J. (2020). LINC02418 promotes malignant
588   behaviors in lung adenocarcinoma cells by sponging miR-4677-3p to upregulate KNL1
589   expression. BMC Pulm Med *20*, 217. 10.1186/s12890-020-01229-0.

590 24. Huo, S.F., Shang, W.L., Yu, M., Ren, X.P., Wen, H.X., Chai, C.Y., Sun, L., Hui, K., Liu, L.H.,
591   Wei, S.H., et al. (2020). STEAP1 facilitates metastasis and epithelial-mesenchymal transition
592   of lung adenocarcinoma via the JAK2/STAT3 signaling pathway. Biosci Rep *40*.
593   10.1042/BSR20193169.

594 25. Zhao, Z., Li, J., Tan, F., Gao, S., and He, J. (2018). mTOR up-regulation of BEX4 promotes
595   lung adenocarcinoma cell proliferation by potentiating OCT4. Biochem Biophys Res
596   Commun *500*, 302-309. 10.1016/j.bbrc.2018.04.064.

597 26. Fan, T., Liu, Y., Liu, H., Wang, L., Tian, H., Zheng, Y., Zheng, B., Xue, L., Li, C., and He, J.
598   (2022). Transmembrane Protein-Based Risk Model and H3K4me3 Modification
599   Characteristics in Lung Adenocarcinoma. Frontiers in Oncology *12*.
600   10.3389/fonc.2022.828814.

601 27. Lee, S., Kang, H., Shin, E., Jeon, J., Youn, H., and Youn, B. (2021). BEX1 and BEX4 Induce
602   GBM Progression through Regulation of Actin Polymerization and Activation of YAP/TAZ
603   Signaling. Int J Mol Sci *22*. 10.3390/ijms22189845.

604 28. Gao, W., Li, J.Z., Chen, S.Q., Chu, C.Y., Chan, J.Y., and Wong, T.S. (2016). Decreased brain-
605   expressed X-linked 4 (BEX4) expression promotes growth of oral squamous cell carcinoma. J
606   Exp Clin Cancer Res *35*, 92. 10.1186/s13046-016-0355-6.

607 29. He, Y., Luo, Y., Zhang, D., Wang, X., Zhang, P., Li, H., Ejaz, S., and Liang, S. (2019). PGK1-
608   mediated cancer progression and drug resistance. Am J Cancer Res *9*, 2280-2302.

609 30. Matsuura, K., Sawai, H., Ikeo, K., Ogawa, S., Iio, E., Isogawa, M., Shimada, N., Komori, A.,
610   Toyoda, H., Kumada, T., et al. (2017). Genome-Wide Association Study Identifies TLL1
611   Variant Associated With Development of Hepatocellular Carcinoma After Eradication of
612   Hepatitis C Virus Infection. Gastroenterology *152*, 1383-1394. 10.1053/j.gastro.2017.01.041.

613 31. Poplawski, P., Alseekh, S., Jankowska, U., Skupien-Rabian, B., Iwanicka-Nowicka, R.,
614   Kossowska, H., Fogtman, A., Rybicka, B., Boguslawska, J., Adamiok-Ostrowska, A., et al.
615   (2023). Coordinated reprogramming of renal cancer transcriptome, metabolome and
616   secretome associates with immune tumor infiltration. Cancer Cell Int *23*, 2. 10.1186/s12935-
617   022-02845-y.

618 32. Guo, M., Li, F., Zhao, L., Fang, Z., Yu, H., Songyang, Z., and Xiong, Y. (2023). Pan-cancer

619   investigation of C-to-U editing reveals its important role in cancer development and new
620   targets for cancer treatment. Front Oncol *13*, 1097667. 10.3389/fonc.2023.1097667.

33.   Povero, D., Johnson, S.M., and Liu, J. (2020). Hypoxia, hypoxia-inducible gene 2 (HIG2)/HILPDA, and intracellular lipolysis in cancer. Cancer Lett *493*, 71-79. 10.1016/j.canlet.2020.06.013.

34.   Erfani, S., Hua, H., Pan, Y., Zhou, B.P., and Yang, X.H. (2021). The Context-Dependent Impact of Integrin-Associated CD151 and Other Tetraspanins on Cancer Development and Progression: A Class of Versatile Mediators of Cellular Function and Signaling, Tumorigenesis and Metastasis. Cancers (Basel) *13*. 10.3390/cancers13092005.

35.   Won, M., Park, K.A., Kim, S., Ju, E., Ko, Y., Yoo, H., Ro, H., Lee, J., Oh, J., Lee, E.G., et al. (2022). ANKRD13a controls early cell-death checkpoint by interacting with RIP1 independent of NF-kappaB. Cell Death Differ *29*, 1152-1163. 10.1038/s41418-021-00906-9.

36.   Xu, J., Chen, Q., Tian, K., Liang, R., Chen, T., Gong, A., Mathy, N.W., Yu, T., and Chen, X. (2020). m6A methyltransferase METTL3 maintains colon cancer tumorigenicity by suppressing SOCS2 to promote cell proliferation. Oncol Rep *44*, 973-986. 10.3892/or.2020.7665.

37.   Xu, C., Li, H., Zhang, L., Jia, T., Duan, L., and Lu, C. (2016). MicroRNA-1915-3p prevents the apoptosis of lung cancer cells by downregulating DRG2 and PBX2. Mol Med Rep *13*, 505-512. 10.3892/mmr.2015.4565.

38.   Mead, T.J. (2022). ADAMTS6: Emerging roles in cardiovascular, musculoskeletal and cancer biology. Front Mol Biosci *9*, 1023511. 10.3389/fmolb.2022.1023511.

39.   Yang, B.Y., Song, J.W., Sun, H.Z., Xing, J.C., Yang, Z.H., Wei, C.Y., Xu, T.Y., Yu, Z.N., Zhang, Y.N., Wang, Y.F., et al. (2018). PSMB8 regulates glioma cell migration, proliferation, and apoptosis through modulating ERK1/2 and PI3K/AKT signaling pathways. Biomed Pharmacother *100*, 205-212. 10.1016/j.biopha.2018.01.170.

40.   Keshet, R., Lee, J.S., Adler, L., Iraqi, M., Ariav, Y., Lim, L.Q.J., Lerner, S., Rabinovich, S., Oren, R., Katzir, R., et al. (2020). Targeting purine synthesis in ASS1-expressing tumors enhances the response to immune checkpoint inhibitors. Nat Cancer *1*, 894-908. 10.1038/s43018-020-0106-7.

41.   Palaparti, A., Baratz, A., and Stifani, S. (1997). The Groucho/Transducin-like Enhancer of split Transcriptional Repressors Interact with the Genetically Defined Amino-terminal Silencing Domain of Histone H3. Journal of Biological Chemistry *272*, 26604-26610. 10.1074/jbc.272.42.26604.

42.   Lin, W., Wang, J., Ge, J., Zhou, R., Hu, Y., Xiao, L., Peng, Q., and Zheng, Z. (2023). The activity of cuproptosis pathway calculated by AUCell algorithm was employed to construct cuproptosis landscape in lung adenocarcinoma. Discover Oncology *14*. 10.1007/s12672-023-00755-7.

43.   Li, X., Jiang, Y., Meisenhelder, J., Yang, W., Hawke, D.H., Zheng, Y., Xia, Y., Aldape, K., He, J., Hunter, T., et al. (2016). Mitochondria-Translocated PGK1 Functions as a Protein Kinase to Coordinate Glycolysis and the TCA Cycle in Tumorigenesis. Mol Cell *61*, 705-719. 10.1016/j.molcel.2016.02.009.

44.   Fu, Q., and Yu, Z. (2020). Phosphoglycerate kinase 1 (PGK) in cancer: A promising target for diagnosis and therapy. Life Sci *256*, 117863. 10.1016/j.lfs.2020.117863.

45.   Waldman, A.D., Fritz, J.M., and Lenardo, M.J. (2020). A guide to cancer immunotherapy:

from T cell basic science to clinical practice. Nat Rev Immunol *20*, 651-668. 10.1038/s41577-
020-0306-5.

46.   Zhou, Z., Li, Y., Hao, H., Wang, Y., Zhou, Z., Wang, Z., and Chu, X. (2019). Screening Hub
Genes as Prognostic Biomarkers of Hepatocellular Carcinoma by Bioinformatics Analysis.
Cell Transplant *28*, 76S-86S. 10.1177/0963689719893950.

47.   Hwang, L.H., Lau, L.F., Smith, D.L., Mistrot, C.A., Hardwick, K.G., Hwang, E.S., Amon, A.,
and Murray, A.W. (1998). Budding yeast Cdc20: a target of the spindle checkpoint. Science
279, 1041-1044. 10.1126/science.279.5353.1041.

48.   Uuskula-Reimand, L., and Wilson, M.D. (2022). Untangling the roles of TOP2A and TOP2B
in transcription and cancer. Sci Adv *8*, eadd4920. 10.1126/sciadv.add4920.

49.   Lamers, F., van der Ploeg, I., Schild, L., Ebus, M.E., Koster, J., Hansen, B.R., Koch, T.,
Versteeg, R., Caron, H.N., and Molenaar, J.J. (2011). Knockdown of survivin (BIRC5) causes
apoptosis in neuroblastoma via mitotic catastrophe. Endocr Relat Cancer *18*, 657-668.
10.1530/ERC-11-0207.

50.   Wittmann, T., Wilm, M., Karsenti, E., and Vernos, I. (2000). TPX2, A novel xenopus MAP
involved in spindle pole organization. J Cell Biol *149*, 1405-1418. 10.1083/jcb.149.7.1405.

51.   Kufer, T.A., Sillje, H.H., Korner, R., Gruss, O.J., Meraldi, P., and Nigg, E.A. (2002). Human
TPX2 is required for targeting Aurora-A kinase to the spindle. J Cell Biol *158*, 617-623.
10.1083/jcb.200204155.

52.   Cancer Genome Atlas Research, N. (2014). Comprehensive molecular profiling of lung
adenocarcinoma. Nature *511*, 543-550. 10.1038/nature13385.

53.   Zheng, F., Kelly, M.R., Ramms, D.J., Heintschel, M.L., Tao, K., Tutuncuoglu, B., Lee, J.J.,
Ono, K., Foussard, H., Chen, M., et al. (2021). Interpretation of cancer mutations using a
multiscale map of protein systems. Science *374*, eabf3067. 10.1126/science.abf3067.

54.   Wang, Q., Shi, Y.L., Zhou, K., Wang, L.L., Yan, Z.X., Liu, Y.L., Xu, L.L., Zhao, S.W., Chu,
H.L., Shi, T.T., et al. (2018). PIK3CA mutations confer resistance to first-line chemotherapy in
colorectal cancer. Cell Death Dis *9*, 739. 10.1038/s41419-018-0776-6.

55.   Shibata, T., Kokubu, A., Tsuta, K., and Hirohashi, S. (2009). Oncogenic mutation of PIK3CA
in small cell lung carcinoma: a potential therapeutic target pathway for chemotherapy-resistant
lung cancer. Cancer Lett *283*, 203-211. 10.1016/j.canlet.2009.03.038.

56.   Hanker, A.B., Pfefferle, A.D., Balko, J.M., Kuba, M.G., Young, C.D., Sanchez, V., Sutton,
C.R., Cheng, H., Perou, C.M., Zhao, J.J., et al. (2013). Mutant PIK3CA accelerates HER2-
driven transgenic mammary tumors and induces resistance to combinations of anti-HER2
therapies. Proc Natl Acad Sci U S A *110*, 14372-14377. 10.1073/pnas.1303204110.

57.   Eng, J., Woo, K.M., Sima, C.S., Plodkowski, A., Hellmann, M.D., Chaft, J.E., Kris, M.G.,
Arcila, M.E., Ladanyi, M., and Drilon, A. (2015). Impact of Concurrent PIK3CA Mutations on
Response to EGFR Tyrosine Kinase Inhibition in EGFR-Mutant Lung Cancers and on
Prognosis in Oncogene-Driven Lung Adenocarcinomas. J Thorac Oncol *10*, 1713-1719.
10.1097/JTO.0000000000000671.

58.   Thorsson, V., Gibbs, D.L., Brown, S.D., Wolf, D., Bortone, D.S., Ou Yang, T.H., Porta-Pardo,
E., Gao, G.F., Plaisier, C.L., Eddy, J.A., et al. (2018). The Immune Landscape of Cancer.
Immunity *48*, 812-830 e814. 10.1016/j.immuni.2018.03.023.

59.   Meigs, T.E., Fedor-Chaiken, M., Kaplan, D.D., Brackenbury, R., and Casey, P.J. (2002).
Galpha12 and Galpha13 negatively regulate the adhesive functions of cadherin. J Biol Chem

277, 24594-24600. 10.1074/jbc.M201984200.

60.   An, Y., Adams, J.R., Hollern, D.P., Zhao, A., Chang, S.G., Gams, M.S., Chung, P.E.D., He, X.,
Jangra, R., Shah, J.S., et al. (2018). Cdh1 and Pik3ca Mutations Cooperate to Induce Immune-
Related Invasive Lobular Carcinoma of the Breast. Cell Rep *25*, 702-714 e706.
10.1016/j.celrep.2018.09.056.

61.   Zeisberg, M., and Neilson, E.G. (2009). Biomarkers for epithelial-mesenchymal transitions. J
Clin Invest *119*, 1429-1437. 10.1172/JCI36183.

62.   Yang, J., Mani, S.A., Donaher, J.L., Ramaswamy, S., Itzykson, R.A., Come, C., Savagner, P.,
Gitelman, I., Richardson, A., and Weinberg, R.A. (2004). Twist, a master regulator of
morphogenesis, plays an essential role in tumor metastasis. Cell *117*, 927-939.
10.1016/j.cell.2004.06.006.

63.   Egner, J.R. (2010). AJCC cancer staging manual. Jama *304*, 1726-1727.

64.   Rebello, R.J., Oing, C., Knudsen, K.E., Loeb, S., Johnson, D.C., Reiter, R.E., Gillessen, S.,
Van der Kwast, T., and Bristow, R.G. (2021). Prostate cancer. Nat Rev Dis Primers *7*, 9.
10.1038/s41572-020-00243-0.

65.   Lee, J.K., Lee, J., Go, H., Lee, C.G., Kim, S., Kim, H.S., Cho, H., Choi, K.S., Ha, G.H., and
Lee, C.W. (2016). Oncogenic microtubule hyperacetylation through BEX4-mediated sirtuin 2
inhibition. Cell Death Dis *7*, e2336. 10.1038/cddis.2016.240.

66.   Lee, J.K., Ha, G.H., Kim, H.S., and Lee, C.W. (2018). Oncogenic potential of BEX4 is
conferred by Polo-like kinase 1-mediated phosphorylation. Exp Mol Med *50*, 1-12.
10.1038/s12276-018-0168-0.

67.   Beroukhim, R., Mermel, C.H., Porter, D., Wei, G., Raychaudhuri, S., Donovan, J., Barretina,
J., Boehm, J.S., Dobson, J., Urashima, M., et al. (2010). The landscape of somatic copy-
number alteration across human cancers. Nature *463*, 899-905. 10.1038/nature08822.

68.   Davoli, T., Xu, A.W., Mengwasser, K.E., Sack, L.M., Yoon, J.C., Park, P.J., and Elledge, S.J.
(2013). Cumulative haploinsufficiency and triplosensitivity drive aneuploidy patterns and
shape the cancer genome. Cell *155*, 948-962. 10.1016/j.cell.2013.10.011.

69.   Girish, V., Lakhani, A.A., Thompson, S.L., Scaduto, C.M., Brown, L.M., Hagenson, R.A.,
Sausville, E.L., Mendelson, B.E., Kandikuppa, P.K., Lukow, D.A., et al. (2023). Oncogene-
like addiction to aneuploidy in human cancers. Science *381*, eadg4521.
10.1126/science.adg4521.

70.   Bosco, N., Goldberg, A., Zhao, X., Mays, J.C., Cheng, P., Johnson, A.F., Bianchi, J.J., Toscani,
C., Di Tommaso, E., Katsnelson, L., et al. (2023). KaryoCreate: A CRISPR-based technology
to study chromosome-specific aneuploidy by targeting human centromeres. Cell *186*, 1985-
2001 e1919. 10.1016/j.cell.2023.03.029.

71.   Ha, S.Y., Choi, M., Lee, T., and Park, C.K. (2016). The Prognostic Role of Mitotic Index in
Hepatocellular Carcinoma Patients after Curative Hepatectomy. Cancer Res Treat *48*, 180-
189. 10.4143/crt.2014.321.

72.   Zhou, C., Wang, Y., Liu, X., Liang, Y., Fan, Z., Jiang, T., Wang, Y., and Wang, L. (2018).
Molecular profiles for insular low-grade gliomas with putamen involvement. J Neurooncol
138, 659-666. 10.1007/s11060-018-2837-1.

73.   Makinen, J.M., Laitakari, K., Johnson, S., Makitaro, R., Bloigu, R., Paakko, P., Lappi-Blanco,
E., and Kaarteenaho, R. (2017). Histological features of malignancy correlate with growth
patterns and patient outcome in lung adenocarcinoma. Histopathology *71*, 425-436.

10.1111/his.13236.

74. Byrne, A., Savas, P., Sant, S., Li, R., Virassamy, B., Luen, S.J., Beavis, P.A., Mackay, L.K., Neeson, P.J., and Loi, S. (2020). Tissue-resident memory T cells in breast cancer control and immunotherapy responses. Nat Rev Clin Oncol *17*, 341-348. 10.1038/s41571-020-0333-y.

75. Liu, W., Sun, L., Zhang, J., Song, W., Li, M., and Wang, H. (2021). The landscape and prognostic value of immune characteristics in uterine corpus endometrial cancer. Biosci Rep *41*. 10.1042/BSR20202321.

76. Ducreux, M., Cuhna, A.S., Caramella, C., Hollebecque, A., Burtin, P., Goere, D., Seufferlein, T., Haustermans, K., Van Laethem, J.L., Conroy, T., et al. (2015). Cancer of the pancreas: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Ann Oncol *26 Suppl 5*, v56-68. 10.1093/annonc/mdv295.

77. Cardoso, F., Kyriakides, S., Ohno, S., Penault-Llorca, F., Poortmans, P., Rubio, I.T., Zackrisson, S., Senkus, E., and clinicalguidelines@esmo.org, E.G.C.E.a. (2019). Early breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-updagger. Ann Oncol *30*, 1194-1220. 10.1093/annonc/mdz173.

78. Oaknin, A., Bosse, T.J., Creutzberg, C.L., Giornelli, G., Harter, P., Joly, F., Lorusso, D., Marth, C., Makker, V., Mirza, M.R., et al. (2022). Endometrial cancer: ESMO Clinical Practice Guideline for diagnosis, treatment and follow-up. Ann Oncol *33*, 860-877. 10.1016/j.annonc.2022.05.009.

79. Vogel, A., Cervantes, A., Chau, I., Daniele, B., Llovet, J.M., Meyer, T., Nault, J.C., Neumann, U., Ricke, J., Sangro, B., et al. (2018). Hepatocellular carcinoma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Ann Oncol *29*, iv238-iv255. 10.1093/annonc/mdy308.

80. Stupp, R., Brada, M., van den Bent, M.J., Tonn, J.C., Pentheroudakis, G., and Group, E.G.W. (2014). High-grade glioma: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Ann Oncol *25 Suppl 3*, iii93-101. 10.1093/annonc/mdu050.

81. Postmus, P.E., Kerr, K.M., Oudkerk, M., Senan, S., Waller, D.A., Vansteenkiste, J., Escriu, C., Peters, S., and Committee, E.G. (2017). Early and locally advanced non-small-cell lung cancer (NSCLC): ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Ann Oncol *28*, iv1-iv21. 10.1093/annonc/mdx222.

82. Gui, C., Kosteniuk, S.E., Lau, J.C., and Megyesi, J.F. (2018). Tumor growth dynamics in serially-imaged low-grade glioma patients. J Neurooncol *139*, 167-175. 10.1007/s11060-018-2857-x.

83. Alvarez, M., Benhammou, J.N., Darci-Maher, N., French, S.W., Han, S.B., Sinsheimer, J.S., Agopian, V.G., Pisegna, J.R., and Pajukanta, P. (2022). Human liver single nucleus and single cell RNA sequencing identify a hepatocellular carcinoma-associated cell-type affecting survival. Genome Med *14*, 50. 10.1186/s13073-022-01055-5.

84. Infante, M., Berghmans, T., Heuvelmans, M.A., Hillerdal, G., and Oudkerk, M. (2013). Slow-growing lung cancer as an emerging entity: from screening to clinical management. Eur Respir J *42*, 1706-1722. 10.1183/09031936.00186212.

85. Hoshino, T. (1984). A commentary on the biology and growth kinetics of low-grade and high-grade gliomas. J Neurosurg *61*, 895-900. 10.3171/jns.1984.61.5.0895.

86. Sajjadi, E., Venetis, K., Piciotti, R., Invernizzi, M., Guerini-Rocco, E., Haricharan, S., and Fusco, N. (2021). Mismatch repair-deficient hormone receptor-positive breast cancers:

Biology and pathological characterization. Cancer Cell Int *21*, 266. 10.1186/s12935-021-
01976-y.

87. Doghri, R., Houcine, Y., Boujelbene, N., Driss, M., Charfi, L., Abbes, I., Mrad, K., and
Sellami, R. (2019). Mismatch Repair Deficiency in Endometrial Cancer:
Immunohistochemistry Staining and Clinical Implications. Appl Immunohistochem Mol
Morphol *27*, 678-682. 10.1097/PAI.0000000000000641.

88. Zhang, Q., Sun, T.Y., Kang, P.M., Qian, K., Deng, B., Zhou, J.H., Wang, R.W., Jiang, B., Li,
K., Liu, F., et al. (2016). Combined analysis of rearrangement of ALK, ROS1, somatic
mutation of EGFR, KRAS, BRAF, PIK3CA, and mRNA expression of ERCC1, TYMS,
RRM1, TUBB3, EGFR in patients with non-small cell lung cancer and their clinical
significance. Cancer Chemoth Pharm *77*, 583-593. 10.1007/s00280-016-2969-y.

89. Donev, I.S., Wang, W., Yamada, T., Li, Q., Takeuchi, S., Matsumoto, K., Yamori, T., Nishioka,
Y., Sone, S., and Yano, S. (2011). Transient PI3K inhibition induces apoptosis and overcomes
HGF-mediated resistance to EGFR-TKIs in EGFR mutant lung cancer. Clin Cancer Res *17*,
2260-2269. 10.1158/1078-0432.CCR-10-1993.

90. Wang, G., Xu, D., Zhang, Z., Li, X., Shi, J., Sun, J., Liu, H.Z., Li, X., Zhou, M., and Zheng, T.
(2021). The pan-cancer landscape of crosstalk between epithelial-mesenchymal transition and
immune evasion relevant to prognosis and immunotherapy response. NPJ Precis Oncol *5*, 56.
10.1038/s41698-021-00200-4.

91. Brown, T.J., Bota, D.A., van Den Bent, M.J., Brown, P.D., Maher, E., Aregawi, D., Liau,
L.M., Buckner, J.C., Weller, M., Berger, M.S., and Glantz, M. (2019). Management of low-
grade glioma: a systematic review and meta-analysis. Neurooncol Pract *6*, 249-258.
10.1093/nop/npy034.

92. Chen, L., Ma, J., Zou, Z., Liu, H., Liu, C., Gong, S., Gao, X., and Liang, G. (2022). Clinical
characteristics and prognosis of patients with glioblastoma: A review of survival analysis of
1674 patients based on SEER database. Medicine (Baltimore) *101*, e32042.
10.1097/MD.0000000000032042.

93. Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,
Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for integrated
models of biomolecular interaction networks. Genome Res *13*, 2498-2504.
10.1101/gr.1239303.

94. Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). clusterProfiler: an R package for comparing
biological themes among gene clusters. OMICS *16*, 284-287. 10.1089/omi.2011.0118.

95. Mayakonda, A., Lin, D.C., Assenov, Y., Plass, C., and Koeffler, H.P. (2018). Maftools:
efficient and comprehensive analysis of somatic variants in cancer. Genome Res *28*, 1747-
1756. 10.1101/gr.239244.118.

96. Tsherniak, A., Vazquez, F., Montgomery, P.G., Weir, B.A., Kryukov, G., Cowley, G.S., Gill, S.,
Harrington, W.F., Pantel, S., Krill-Burger, J.M., et al. (2017). Defining a Cancer Dependency
Map. Cell *170*, 564-576 e516. 10.1016/j.cell.2017.06.010.

97. Yang, W., Soares, J., Greninger, P., Edelman, E.J., Lightfoot, H., Forbes, S., Bindal, N., Beare,
D., Smith, J.A., Thompson, I.R., et al. (2013). Genomics of Drug Sensitivity in Cancer
(GDSC): a resource for therapeutic biomarker discovery in cancer cells. Nucleic Acids Res *41*,
D955-961. 10.1093/nar/gks1111.

98. Subramanian, A., Narayan, R., Corsello, S.M., Peck, D.D., Natoli, T.E., Lu, X., Gould, J.,

Davis, J.F., Tubelli, A.A., Asiedu, J.K., et al. (2017). A Next Generation Connectivity Map:
L1000 Platform and the First 1,000,000 Profiles. Cell *171*, 1437-1452 e1417.
10.1016/j.cell.2017.10.049.

99.   Finotello, F., Mayer, C., Plattner, C., Laschober, G., Rieder, D., Hackl, H., Krogsdam, A.,
Loncova, Z., Posch, W., Wilflingseder, D., et al. (2019). Molecular and pharmacological
modulators of the tumor immune contexture revealed by deconvolution of RNA-seq data.
Genome Med *11*, 34. 10.1186/s13073-019-0638-6.

100.   Sturm, G., Finotello, F., and List, M. (2020). Immunedeconv: An R Package for Unified
Access to Computational Methods for Estimating Immune Cell Fractions from Bulk RNA-
Sequencing Data. Methods Mol Biol *2120*, 223-232. 10.1007/978-1-0716-0327-7_16.

101.   Hanzelmann, S., Castelo, R., and Guinney, J. (2013). GSVA: gene set variation analysis for
microarray and RNA-seq data. BMC Bioinformatics *14*, 7. 10.1186/1471-2105-14-7.

 Figure Legends

Figure 1: MEMORY uncovers the enrichment of mitosis and immune signatures in multiple

cancers. (A) Sample sizes, ranging from 10% to 100% with 10% intervals ("gradient"), were

used for 1000 permutations of survival analyses of all 15 cancer types Each matrix was

divided into high and low expression groups based on median gene expression values, and

survival analyses were performed. Log-rank test significance results were coded as 1 for

significant (P < 0.05) and 0 for non-significant (P > 0.05) outcomes, forming survival analysis

matrices and the summarized significant probability matrix, which allowed the identification

of GEARs. (B) The maximum of significant probability for each gradient sample size of

every cancer type. Sample number ratio refers to the percentage of samples in each sampling

gradient compared to the total number of samples. (C) The pathways were enriched by GOEA

based on the GEARs of each cancer type. The displayed pathways represent the top 5 most

significant pathways for each cancer type. Mitosis-related pathways were marked in red

whereas immune-related pathways were marked in blue.

Figure 2: Identification of hub genes in mitosis or immune-related cancers. (A) The SAS of

all GEARs was calculated, and CSN was constructed. (B) The P-value of survival analysis

was determined based on hub gene**s** grouped by edge numbers (250, 500, 1000, 2000, 4000

and 8000 edges) from six CSNs. (C) The left nodes represent five different cancers, while the

right nodes detail the functional types of hub-gene pathways. The height of the edges in the

middle represents the proportion of hub-gene pathways corresponding to a specified

functional type. (D-E) The CSN of LUAD and BRCA. The enlarged section represents the

hub gene. (F) LUAD clustering based on hub gene expression, compared with inhibitor-based

classification[52]. (G) LUAD clustering based on hub gene expression in comparison with the

classic classification.

Figure 3: Different LUAD subgroups were characterized by unique genetic mutation profiles.

(A) TMB analysis in three groups of LUAD samples. ***P < 0.0005. (B) Proportion of

different oncogenic drivers including KRAS, EGFR, BRAF, ERBB2 mutations, and ALK,

ROS1 fusions in three LUAD subgroups. (C-E) Comparison of top gene mutations in three

LUAD clusters, including ML vs. MM (C), MM vs. MH (D), ML vs MH (E).

Figure 4: PIK3CA mutation associates with mitosis and drug resistance in LUAD. (A)
Cancer-related gene mutations were annotated based on gene dependency scores data from
Depmap database. Functional mutations were indicated in green; functional (subtype-
associated) mutations were highlighted in red; non-functional mutations were indicated in
gray. (B) The analysis of the NeST differential pathways across three groups including ML vs.
MM, MM vs. MH, ML vs MH. (C) The heat map showed the hub gene expression of A549
after compounds treatment from cMAP database[98]. Blue annotations meant 76 compounds
that can inhibit hub gene expression. (D) Comparison of the IC50 z-score of five tumor cell
growth inhibitors for A549 (*KRAS* mutation) and SW1573 (*KRAS* and *PIK3CA* mutation).

Figure 5: The hub gene classification of BRCA revealed a significant association between
EMT and immune infiltration. (A) The hub gene expression heatmap containing the result of
hub gene classification and PAM50 classification of BRCA[58]. (B) Comparison of molecular
classification based on hub genes with PAM50 classification. (C-E) Comparison of gene
mutations in three groups of BRCA samples, including IL vs. IM (C), IM vs. IH (D), IL vs. IH
(E). (F) Comparison of the total immune cell rate in three BRCA clusters. ***P < 0.0005. (G)
Comparison of *CDH1* expression in BRCA samples with wildtype or mutant *CDH1*. ***P <
0.0005. (H) A schematic diagram illustrating the correlation between the EMT score and
immune cell rate. The red line represents the fitting curve. The correlation analysis was
performed using Pearson correlation coefficient.

Figure 6: Mitosis and immune signatures predict patient prognosis at the pan-cancer level.
The mitosis and immune-related pathway scores of 33 cancer types were analyzed, and the
median score was utilized as a threshold to categorize patients for survival analysis. Among
the 33 cancer types examined, 10 cancer types were exclusively associated with the mitosis
score, 2 cancer types were exclusively associated with the immune score, and 5 cancer types
showed a correlation with both mitosis and immunity scores concurrently.

Figure 2

A

C

B

Figure 6

Multi-gradient Permutation Survival Analysis Identifies Mitosis and Immune Signatures Steadily Associated with Cancer Patient Prognosis

Xinlei Cai[1], Yi Ye[2], Xiaoping Liu[1], Zhaoyuan Fang[4,5], Luonan Chen[1,2,6], Fei Li[3*], Hongbin Ji[1,2,6*]

[1]Key Laboratory of Systems Health Science of Zhejiang Province, School of Life Science, Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou, 310024, China.

[2]State Key Laboratory of Cell Biology, Shanghai Institute of Biochemistry and Cell Biology, Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences, Shanghai 200031, China.

[3]Department of Pathology and Frontier Innovation Center, School of Basic Medical Sciences, Fudan University, Shanghai, China.

[4]Zhejiang University-University of Edinburgh Institute, Zhejiang University School of Medicine, Haining , 314400, China

[5]The Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, 310000, China

[6]School of Life Science and Technology, Shanghai Tech University, Shanghai 200120, China.

Running title: Mitosis and immune signatures in predicting cancer patient prognosis

*Corresponding author:

Hongbin Ji, State Key Laboratory of Cell Biology, Shanghai Institute of Biochemistry and Cell Biology, Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences, 320 Yue Yang Road, Shanghai, 200031, China, Phone: 86-21-54921108, Fax: 86-21-54921101, E-mail address: hbji@sibcb.ac.cn;

Fei Li, Department of Pathology and Frontier Innovation Center, School of Basic Medical

Sciences, Fudan University, 131 Dong An Road, Shanghai, 200032, China, E-mail address:

li_fei@fudan.edu.cn.

Supplementary Figure 1: (A) to (O) The number of genes significantly related to prognosis which were obtained through MEMORY was positively correlated with sample size. The vertical value of the dotted line is 0.8.

Supplementary Figure 2: (A) to (O) The significant probability of GEAR at each gradient. The vertical value of the dotted line is 0.8.

Supplementary Figure 3: (A) to (O) The core survival networks of 15 cancer types. Red nodes are hub genes.

Supplementary Figure 4: (A) to (O) 15 cancer types were genotyped through hierarchical clustering based on the expression of hub genes, and the samples of each cancer were divided into three clusters

Supplementary Figure 5: (A-C) The gene dependency analysis of hub genes for LGG (A), LIHC (B), LUAD (C). (D) The number of compounds associated with A549 in the cMAP and GDSC databases. (E) Relative sensitivity of A549 cell line to all compounds from the GDSC database, calculated based on IC50 values. (F) Kaplan-Meyer overall survival curves for LUAD classic subgroups. (G) Kaplan-Meyer overall survival curves for LUAD hub gene subgroups.

Supplementary Figure 6: (A) Kaplan-Meyer overall survival curves for BRCA classic subgroups. (B) Kaplan-Meyer overall survival curves for BRCA PAM50 subgroups. (C-L) Comparison of infiltration of different types of immune cells in three group BRCA samples. ***P < 0.0005, **P < 0.005 and *P < 0.05. (M) Comparison of the proportion of immune

cells in the total cells in BRCA samples with high *CDH1* expression and low *CDH1* expression. ***P < 0.0005. (N) Schematic diagram of the correlation between *CDH1* expression and *VIM* expression. The red line was a fitting curve. Pearson correlation coefficient was used for correlation analysis. (O) Schematic diagram of the correlation between *CDH1* expression and *TWIST2* expression. The red line was a fitting curve. Pearson correlation coefficient was used for correlation analysis.