Multi-gradient Permutation Survival Analysis Identifies Mitosis and Immune Signatures Steadily Associated with Cancer Patient Prognosis

Xinlei Cai[1], Yi Ye[2], Xiaoping Liu[1], Zhaoyuan Fang[4,5], Luonan Chen[1,2,6], Fei Li[3*], Hongbin Ji[1,2,6*]

[1]Key Laboratory of Systems Health Science of Zhejiang Province, School of Life Science, Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou, 310024, China.

[2]State Key Laboratory of Cell Biology, Shanghai Institute of Biochemistry and Cell Biology, Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences, Shanghai 200031, China.

[3]Department of Pathology and Frontier Innovation Center, School of Basic Medical Sciences, Fudan University, Shanghai, China.

[4]Zhejiang University-University of Edinburgh Institute, Zhejiang University School of Medicine, Haining，314400, China

[5]The Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, 310000, China

[6]School of Life Science and Technology, Shanghai Tech University, Shanghai 200120, China.

Running title: Mitosis and immune signatures in predicting cancer patient prognosis

*Corresponding author:

Hongbin Ji, State Key Laboratory of Cell Biology, Shanghai Institute of Biochemistry and Cell Biology, Center for Excellence in Molecular Cell Science, Chinese Academy of Sciences, 320 Yue Yang Road, Shanghai, 200031, China, Phone: 86-21-54921108, Fax: 86-21-54921101, E-mail address: hbji@sibcb.ac.cn;

Fei Li, Department of Pathology and Frontier Innovation Center, School of Basic Medical

29   Sciences, Fudan University, 131 Dong An Road, Shanghai, 200032, China, E-mail address:

30   li_fei@fudan.edu.cn.

59   li_fei@fudan.edu.cn.

Abstract

The inconsistency of the association between genes and cancer prognosis is often attributed to many variables that contribute to patient survival. Whether there exist the Genes Steadily Associated with Prognosis (GEARs) and what their functions are remain largely elusive. We have developed a novel method called "Multi-gradient Permutation Survival Analysis " (MEMORY) to screen the GEARs using RNA-seq data from the TCGA database. Then we employed a network construction approach to identify hub genes from GEARs, and utilized them for cancer classification. In the case of LUAD, the GEARs were found to be related to mitosis. Our analysis suggested that LUAD cell lines carrying *PIK3CA* mutations exhibit increased drug resistance. For BRCA, the GEARs were related to immunity. The analysis revealed that *CDH1* mutation might influence immune infiltration through the EMT process in BRCA. We further explored the prognostic relevance of mitosis and immunity through their respective scores. This study offers significant biological insights into GEARs and highlights their potential as robust prognostic indicators across diverse cancer types.

Keywords: Multi-gradient Permutation Survival Analysis (MEMORY), Gene Steadily Associated with Prognosis (GEAR), Hub Genes, Mitosis, Immune

Introduction

Cancer is one of the leading causes of death worldwide, with over 200 types identified. Despite of huge advancements in the field, accurately predicting cancer patient prognosis still remains a significant challenge[1,2]. Previous studies have highlighted the complexity of this task, which is influenced by various factors including cancer type, clinical stage, therapeutic interventions, nursing care, unexpected comorbidities, and other non-cancer related illnesses that may interplay[3-6]. Furthermore, the gene expression plays a crucial role in predicting cancer patient prognosis, such as *HER2*, *VEGF*, *Ki67*, etc[7-12]. Nevertheless, there is still a need for further improvement in prognostic accuracy to better inform treatment decisions and patient outcomes.

Survival analysis is commonly used to assess the correlation between genes and cancer patient prognosis[13]. However, inconsistent findings are frequently observed, even within the same type of cancer. For instance, studies exploring the correlation between *CCND1* and prognosis in NSCLC reported contrasting results, including positive correlation, inverse correlation, or negligible influence[14-16]. These discrepancies can be attributed to various influential factors, such as differences in sample size, cohort characteristics (including cancer subtypes and tumor staging) and variations in therapy approaches[17]. Such observations raise an important question: whether there exist Genes Steadily Associated with Prognosis (GEARs) that consistently correlate with patient outcomes across different conditions, particularly considering the varying sample sizes used in different studies? Affirmative answers to this question would have significant implication not only for the development of more accurate prognostic models but also for enhancing our understanding of cancer biology.

The Cancer Genome Atlas (TCGA) is a comprehensive cancer genomics project initiated in the United States. It consists of transcriptome data, genomic data, and clinical information pertaining to 33 different cancer types, making it the largest cancer clinical sample database currently available. In this study, we developed a novel method called "Multi-gradient Permutation Survival Analysis" (MEMORY) with the utilization of TCGA RNA-seq data, and accessed the potential existence of GEARs across 15 cancer types, each comprising a cohort of over 200 patients. Furthermore, we also evaluated the prognostic predictive power of these

GEARs and explored their potential biological functions in driving cancer progression.

Results

Multi-gradient permutation survival analysis identifies GEARs associated with mitosis

and immune across multiple cancer types

GEARs are a group of genes consistently and significantly correlate with patient survival,

independent of the sample size. To identify these GEARs, we developed a novel method called

"Multi-gradient Permutation Survival Analysis " (MEMORY). This method allows us to assess

the correlation between a specific gene and cancer patient prognosis using available

transcriptomic dataset (**Figure 1A, Table 1**). Initially, we started with a sampling number at 10%

of the cohort size initially, and gradually increased the sampling number with each 10%

increment, which was analyzed with 1000 permutations. By calculating the statistical

probability of each gene's association with patient survival, we can identify a group of GEARS

for further analyses.

In this study, we utilized the TCGA datasets for several reasons, including the diversity of

cancer types, the availability of gene expression profiles, and patient prognosis information.

We set a minimum cohort size of 200 and included 15 eligible cancer types for analysis. These

cancer types include bladder urothelial carcinoma (BLCA), breast invasive carcinoma (BRCA),

cervical squamous cell carcinoma and endocervical Adenocarcinoma (CESC), colon

Adenocarcinoma (COAD), kidney renal papillary cell carcinoma (KIRC), brain lower grade

glioma (LGG), liver hepatocellular carcinoma (LIHC), lung adenocarcinoma (LUAD), lung

squamous cell sarcinoma (LUSC), ovarian serous cystadenocarcinoma (OV), pancreatic

adenocarcinoma (PAAD), prostate adenocarcinoma (PRAD), stomach adenocarcinoma

(STAD), thyroid carcinoma (THCA), and uterine corpus endometrial carcinoma (UCEC). As

the number of samples increases, the survival probability of certain genes gradually approaches

1. Once this score reaches 0.8 and remained consistent with further sample gradient increase,

the gene is considered a GEAR (**Figure 1B**). Remarkably, we successfully identified a set of

GEARs across all 15 cancer types (**Supplementary Figure 1-2，Table 2**). The GEAR counts

in most cancer types ranged from 100 to1000, with exceptions of CESC, KIRC, LGG, and

PAAD with over 1000 GEARs, and THCA with only 22 GEARS (**Table 2**). In LUAD, the top

10 GEARs with the highest significance probabilities were *TLE1*, *GNG7*, *ERO1A*, *ANLN*,

DKK1, *TMEM125*, *S100A16*, *KNL1*, *STEAP1*, and *BEX4*. Most of these genes are known to

promote LUAD malignant progression except for S100A16[18-26]. In other cancer types, *BEX4*

was identified as a common GEAR in KIRC, LGG, PAAD, and STAD (Table 2). *BEX4* is

reported as a proto-oncogene promoting cancer onset and malignant progression in multiple

cancers including LUAD, glioblastoma multiforme and oral squamous cell carcinoma [25,27,28].

We also identified the most significant GEARs in individual cancer types, such as *TLL1*

(BLCA), *PGK1* (BRCA), *RFXANK* (CESC), *DPP7* (COAD), *VWA8* (KIRC), *SCMH1* (LGG),

HILPDA (LIHC), *TLE1* (LUAD), *CD151* (LUSC), *ANKRD13A* (OV), *SOCS2* (PAAD), *DRG2*

(PRAD), *ADAMTS6* (STAD), *PSMB8* (THCA), *ASS1* (UCEC) (**Table 3**). Many of these genes

were previously reported to associate with tumorigenesis[18,29-40]. For example, *TLE1* is known

as a transcriptional repressor that promotes cell proliferation, migration, and inhibits apoptosis

in LUAD [41,42]. Additionally, *PGK1*, a key enzyme in the glycolytic process, has been shown to

promote cell proliferation, migration, and invasion in multiple cancers[43,44]. These findings

demonstrate the intricate link between the functionality of GEARs and the initiation and

progression of cancer.

To gain deep insights into the biological functions of these identified GEARs, we conducted

Gene Ontology Enrichment Analysis (GOEA) (**Figure 1C**). Interestingly, we found the

mitosis-related pathways were enriched in LICH, LUAD, LGG, and PAAD, and the immune-

related pathways were enriched in BRCA and UCEC[45]. Additionally, other cancer types

exhibited enrichment in various pathways, such as organic acid metabolism pathway in KIRC,

oxidative phosphorylation pathway in CESC, organ development-related pathways in LUSC,

neurogenesis-related pathways in BLCA, and sustenance metabolism pathways in THCA.

Given the crucial role of mitosis in cancer progression and the significance of the immune

system in cancer-host interactions, we specifically focused on the mitosis and immune-related

GERAs, which accounted for approximately 40% of all the analyzed cancers.

Identification of hub genes in mitosis and immune-related cancers

To further identify crucial genes within the GEARs, we conducted and extracted the higher-

ranked edges to construct the core survival network (CSN) (**Figure 2A, Table 4**). The top 10

GEARs with highest degree in these networks were defined as hub genes [46]. We then classified

the samples using the hub genes derived from these networks and evaluated their clinical relevance (**Supplementary Figure 3A-O, Supplementary Figure** 4A-O). A certain degree of correlation with cancer stages (TNM stages) is observed in most cancer types except for COAD, LUSC and PRAD（**Supplementary Figure 5A-K**）. Furthermore, we conducted GOEA on the hub genes selected from CSNs and found that the results were consistent with the GEAR analysis (**Table 5**). Specifically, the hub genes in LIHC, LGG, and LUAD were enriched in mitosis-related pathways, whereas the hub genes in BRCA and UCEC were enriched in immune-related pathways (**Figure 2B**). For instance, the 9 hub genes in LUAD were associated with functions related to mitosis, whereas the 8 hub genes of BRCA were associated with immune-related functions (**Figure 2C-D**).

Furthermore, we conducted survival-dependent analyses on the hub genes of LGG, LIHC, and LUAD. These analyses revealed that mitosis-related hub genes are closely associated with cancer cell viability, especially those hub genes that are correlated with multiple cancer types such as *CDC20*, *TOP2A*, *BIRC5* and *TPX2*[47-50] (**Supplementary Figure 6A-C**). The importance of these genes is well established. For instance, *TPX2*, a hub gene in all three cancers, is known to play a crucial role in normal spindle assembly during mitosis and is essential for cell proliferation[51]. The significance of these hub genes for the survival of cancer cells suggests that the expression levels of these hub genes can be used to screen for inhibitors of tumor cell growth. By integrating the Genomics of Drug Sensitivity in Cancer (GDSC) and Connectivity Map (cMAP) databases, we identified 76 individual compounds that are able to effectively suppress the expression of the 10 hub genes in LUAD cell lines (**Figure 2E, Supplementary Figure 6D-E**). These compounds also significantly inhibit LUAD cell growth and may serve as potential therapeutic agents for the treatment of LUAD.

We then analyzed the LUAD dataset, which had a larger sample size compared to LGG and LIHC. Based on the expression of the hub genes, we classified the LUAD samples into three subgroups: mitosis low (ML), mitosis medium (MM), and mitosis high (MH) (**Figure 2F, Table 6**). Previous study has reported three classic subgroups of LUAD known as the terminal respiratory unit (TRU), the proximal-proliferative (PP), and the proximal-inflammatory (PI)

subgroups[52]. Our analysis revealed that the ML subgroup was primarily enriched with TRU

subgroup and a small number of samples from the PP subgroup, but not the PI subgroup. The

MH subgroup showed high expression of mitosis-related genes and predominantly

encompassed PI and PP subgroups, whereas the MM group exhibited intermediate levels of

mitosis-related gene expression (**Figure 2F-G**). This suggests that the new categorization

method can help identify new factors that influence patient prognosis (**Supplementary Figure**

6F-G).

Distinct genetic mutation landscapes characterize different clusters of LUAD

The expression of hub genes provided crucial indicators for distinguishing various subgroups

of LUAD. However, the underlying mechanisms driving these variations in expression patterns

remain unknown. In our subsequent research, we aimed to explore the genomic differences,

particularly in terms of gene mutations, among different LUAD subgroups. Initially, we

analyzed the variations of genetic mutation landscape of different LUAD subgroups by

assessing the tumor mutation burden (TMB). Interestingly, we found significant differences in

TMB among the various subgroups of LUAD (**Figure 3A**). Analysis of common driver genes

further revealed distinct proportions of tumors with *ALK* and *ROS1* fusions in the favorable

prognosis ML and MM subgroups, compared to tumors with mutations in *KRAS*, *EGFR*, *BRAF*,

and *ERBB2* (**Figure 3B**). Moreover, low tumor mitotic activity is associated with better

prognosis in LUAD subgroups with *EGFR* mutations or pan-negative (no oncogenic alteration

in genes including *KRAS*, *EGFR*, *BRAF*, *ERBB2*, *PIK3CA*, *ALK* and *ROS1*) but not in those

with *KRAS* or *BRAF* mutations (**Figure 3C-D, Supplementary Figure 6H-J**). Furthermore,

we observed unique mutation characteristics in each of these 3 subgroups (**Figure 3E-G**). For

example, TP53 mutation, a prevalent tumor suppressor gene mutation, was frequently observed

in the MH subgroup with a mutation rate of 73%, compared to mutation rates of 14% in the ML

subgroup and 39% in the MM subgroup. Additionally, genes such as *CSMD3*, *RP1L1*, *ZFHX4*

exhibited similar trend in mutation frequency across the three subgroups. These findings

indicate substantial genomic differences among these three LUAD subgroups based on the hub

genes.

We identified gene mutations that showed significant changes through gene dependence

analysis (**Figure 4A**). To further the functional implications of these mutations, we enriched

them using a pathway system called Nested Systems in Tumors (NeST)[53]. The results revealed

notable differences in multiple functional pathways, including androgen receptor, cell cycle,

TP53, *EGFR*, IL-6, and *PIK3CA*-related pathways (**Figure 4B**). To gain further insights into

the functional implications of these different mutations, we assessed the gene dependency of

cells with these mutations using the DepMap database. Importantly, we observed that the

pathway differences between the MM and MH clusters were primarily associated with

PIK3CA-related pathways. Previous studies have reported that *PIK3CA* mutation confers

resistance in colorectal cancer, lung cancer, and breast cancer[54-56]. Therefore, we aimed to

validate the role of *PIK3CA* mutations in drug resistance using public databases. We further

analyzed A549 cells (a lung adenocarcinoma cell line with *KRAS* mutation) and SW1573 cells

(a lung adenocarcinoma cell line with both *KRAS* and *PIK3CA* mutations) using their respective

GDSC data and identified five potentially effective inhibitors (BMS-345541, Dactinomycin,

Epirubicin, Irinotecan, and Topotecan) from the previously screened set of 76 LUAD cell

growth inhibitors. Strikingly, SW1573 cells exhibited increased resistance to all these 5

inhibitors when compared to A549 cells (**Figure 4C**). In line with this, clinical data also

supports the notion that concurrent *PIK3CA* mutation is a poor prognostic factor for LUAD

patients[57]. These findings suggest that *PIK3CA* mutation may contribute to drug resistance in

LUAD.

Distinct clusters of BRCA exhibit different immune infiltration landscapes

We further focused on the immune-related cancer types, with BRCA standing out as a

representative immune-related cancer. This prompted us to conduct a more comprehensive

analysis (**Figure 1C**). We hierarchically grouped the BRCA into three distinct immune

subgroups based on the expression of hub genes: Immune low (IL), Immune medium (IM), and

Immune high (IH) (**Figure 5A, Table 7**). Interestingly, all the classifications based on the

widely used PAM50 classification contained samples from the IL, IM, and IH subgroups,

although there were minor differences in proportions[58] (**Figure 5B, Supplementary Figure**

7A-B). Next, we investigated the relationship among PAM50 classification, immune subtypes,

 Integrating our method with traditional classification may enable a more detailed stratification of breast cancer samples. Mutation analysis revealed distinct patterns of gene mutations among the three subgroups, with significant differences in mutation frequencies of genes such as *TP53*, *CDH1*, and *LRP2* (**Figure 5C-E**). We conducted a comparison of the proportions of immune cells across these three subgroups and observed a strong association between the overall immune cell proportion and the clustering results based on hub genes (**Figure 5F, Supplementary Figure 8A-J**). Specifically, the IH subgroup exhibited significantly higher proportions of CD8$^+$ T cells and Treg cells, with mean percentages at 5.0% and 6.0%, respectively, compared to 1.4% and 3.2% in the IM subgroup, and 0.3% and 1.6% in the IL subgroup. These findings suggest the presence of distinct immune responses in these three subgroups, indicating a potential association between genetic mutation pattern and the immune microenvironment landscape.

Next, we explored the specific genomic factors influencing immune infiltration in BRCA. Mutation analysis revealed that *CDH1* gene mutation ranked as the second most prevalent genetic alteration in BRCA samples, following TP53 mutation (**Figure 5C-E**). Previous study has indicated that *CDH1* is involved in mechanisms regulating cell-cell adhesions, mobility, and proliferation of epithelial cells[59]. We found that *CDH1*-mutant samples exhibited significantly lower *CDH1* expression compared to *CDH1*-wildtype samples, and *CDH1* expression showed a close correlation with immune cell infiltration (**Figure 5G, Supplementary Figure 8K**). These results were consistent with clinical observations of *CDH1* mutation and high immune infiltration in invasive lobular carcinoma of the breast [60].

We further investigated how *CDH1* mutation influenced immune infiltration. In BRCA, we observed a significant association between the expression of *CDH1* and the expression of EMT marker genes *VIM* and *TWIST2* (**Supplementary Figure 8L-M**)[61,62]. This suggests that the regulation of *CDH1* is intricately linked with EMT process. Through calculating the EMT score, we observed a positive correlation between the EMT score and the proportion of immune infiltration (**Figure 5H**). Consequently, *CDH1* might influence immune infiltration through the

EMT process.

Mitotic and immune signatures predict patient prognosis at pan-cancer level

We further analyzed the association of *CDH1* and *PIK3CA* with specific biological processes

at pan-cancer level. Our data showed that *PIK3CA* level was positively correlated with mitotic

scores in most cancer types, whereas the correlation between *CDH1* expression and the

proportion of immune cell infiltration was positive in PRAD, LGG, OV, Uveal Melanoma

(UVM), THCA, LIHC and LUAD but negative in BRCA, Testicular Germ Cell Tumors

(TGCT), Thymoma (THYM), LUSC, BLCA, Head and Neck squamous cell carcinoma

(HNSC), Sarcoma (SARC), Esophageal carcinoma (ESCA), PAAD and STAD (Figure 6A-B).

Finally, we sought to explore the prognostic relevance of mitosis and immune by calculating

their scores and assessing their correlation with patient outcomes at the pan-cancer level. The

scores for these biological processes were computed using RNA-seq data from the TCGA

database, and the median score was used as a cut-off to categorize patients into different groups.

Out of the 33 cancer types analyzed, the prognosis of 19 cancer types showed a significant

correlation with at least one of these two pathways (**Figure 6C**). Specifically, 10 cancer types

were exclusively associated with the mitosis score, 4 cancer types were exclusively associated

with the immune score, and 5 cancer types exhibited a correlation with both mitosis and

immune scores simultaneously. Overall, the identification of mitosis and immune-related

biological processes as significant prognostic factors at the pan-cancer level suggests their

potential utility as valuable biomarkers for predicting patient prognosis.

Discussion

Due to the multifaceted nature of variables affecting cancer patient prognosis, it remains

uncertain whether there exist a set of genes steadily associated with cancer prognosis,

regardless of sample size and other factors. Here, we utilized the MEMORY method to address

this question and discovered that all the cancer types have GEARs. We observed significant

variation in the number of GEARs among different cancer types, indicative of cancer type-

specific patterns. The substantial heterogeneity in driver gene and mortality rates among various

cancer types could potentially explain this phenomenon. For example, THCA, known for its

low malignancy, displays the fewest genetic expression alterations among all the studied cancer

types. This observation could be correlated with the relatively high five-year survival rate in

THCA, which exceeds 50% even in advanced stage[63]. In contrast, PRAD, despite of a favorable

prognosis similar to THCA, exhibits a significantly higher number of genetic expression

alterations. We hypothesize that the positive prognosis in PRAD cases might be largely

attributed to early diagnosis, which enables timely treatment for the majority of PRAD patients

[64]. This discrepancy between the number of genetic alterations and prognosis in THCA and

PRAD highlights the intricate nature of cancer genetics and emphasizes the importance of

personalized considerations in cancer treatment and prognosis evaluation. Furthermore, certain

genes are commonly found in the GEARs of various cancer types, indicating their potential

significance in cancer development. For instance, *BEX4* is present in the GEARs of LUAD,

KIRC, LGG, PAAD, and STAD, and known to play oncogenic role in inducing carcinogenic

aneuploid transformation via modulating the acetylation of α-tubulin[65,66]. Aneuploidy is a

hallmark characteristic of cancer and can lead to alterations in the dosage of oncogenes or tumor

suppressor genes, thereby influencing tumor initiation and progression [67-70]. The regulation of

aneuploid transformation by *BEX4* may represent a common mechanism through which this

gene impacts prognosis across different cancer types. Subsequently, we discovered that GEARs

across different cancers displayed distinct functional characteristics. Notably, a recurrent theme

was the prominence of mitosis-related and immune-related features. Specifically, the GEARs

of LGG, LIHC, and LUAD were enriched in mitosis-related pathways, whereas BRCA and

UCEC showed the enrichment of immune-related pathways. Mitosis and immune processes

have been widely recognized as having a significant impact on patient prognosis[71-75]. However,

current clinical guidelines do not yet recommend the use of transcriptomic data to assess scores

related to mitotic and immune pathways for predicting patient outcomes, despite the well-

established association between these pathways and prognosis in cancer[76-81]. Cancers enriched

with mitosis pathways often exhibit heterogeneous tumor growth kinetics across individuals,

with tumor size being one of the crucial factors influencing patient survival [82-84]. For instance,

in the case of LGG, the growth rate of tumors directly impacts the patient's prognosis due to

the primary location in the brain. Consequently, alterations in the expression levels of genes

related to mitosis are often indicative of the prognosis of LGG [85]. Cancer types that are enriched

in immune-related pathways, such as BRCA and UCEC, are closely associated with

deficiencies in DNA mismatch repair (MMR) [86,87]. Cancers enriched in immune-related

pathways often exhibit heterogeneous tumor growth kinetics across individuals, where tumor

size is closely correlated with patient survival.

To explore the underlying differences between samples associated with distinct mitotic and

immune-related pathways, we employed GEAR analysis to identify hub genes for further

molecular classification and mutation analysis. Specifically, we focused on LUAD and BRCA,

two representative cancer types exhibiting enriched mitosis and immune signatures in GEARs.

By classifying the hub gene, we divided LUAD into three subgroups: ML, MM and MH, which

displayed significant difference in survival outcomes. Subsequently, we utilized NeST to

analyze the mutations within these subgroups. Notably, there were significant differences

observed in cell cycle and signal transduction-related pathways among ML, MM and MH

subgroups. Of particular importance, the *PIK3CA*-related pathway emerged as a key

differentiating pathway between the MH and MM subgroups. Our analyses suggest that LUAD

cell line carrying *PIK3CA* mutation may exhibit increased drug resistance. This aligns with

previous studies demonstrating that targeting the PI3K pathway can overcome drug

resistance[88,89]. Of course, future research is warranted to elucidate the exact functional role of

PIK3CA mutations in LUAD.

To investigate the factors influencing immune infiltration in BRCA, we classified the samples

based on hub genes and analyzed the differentially mutated genes. Interestingly, we found that

CDH1 mutation occurred at a high frequency in the IH subgroups. This led us to speculate that

CDH1 plays a crucial role in the regulation of immune infiltration in BRCA. Furthermore,

previous studies have reported that *CDH1* inactivation promotes immune infiltration in breast

cancer[60]. *CDH1* is a vital gene associated with EMT, and various studies have demonstrated the

close relationship between EMT and the tumor immune microenvironment in different

cancers[90]. Consistently, our results also supported the correlation between EMT score and

immune infiltration in BRCA. These findings suggest that the mutations in *PIK3CA* and *CDH1*,

identified through GEAR analysis, have significant impacts on cancer development and hold

potential value in improving clinical therapies. This further emphasizes the importance of

GEARs in understanding cancer biology and guiding treatment strategies.

Lastly, we investigated the prognostic predictive capabilities of the mitosis score and immune

score at the pan-cancer level. Surprisingly, we found that approximately half of the cancer

types exhibited significant correlations between these two scores and patient prognosis.

Interestingly, even for cancers originating from the same primary location, their correlations

with these scores could differ, indicating the potential diversity of mechanisms underlying

cancer-related mortality. For instance, both LGG and GBM are brain tumors, but the primary

risk factor for patient prognosis in LGG is closely linked to tumor diameter, whereas the main

risk factor for GBM patients lies in its high invasiveness and challenge of surgical resection[91,92].

Despite of the strong association between GEAR and patient prognosis, the edges of the CSN

constructed based on GEARs was undirected. As a result, the hub genes identified from GEAR

in the CSN may primarily serve as stable and effective biological markers. Additionally,

through multi-omics analysis, we obtained some functional mutations, but the therapeutic

significance of these mutations remains to be elucidated. For example, further study is needed

to understand the exact role of *PIK3CA* mutations in promoting tumor cell proliferation and

drug resistance. Similarly, the association of *CDH1* mutations with the infiltration of multiple

immune cell types also warrants additional experimental investigation. In our future studies,

we plan to utilize protein-protein interaction networks and pathway databases to construct a

novel network based on the CSN. This approach will allow us to directly screen genes from

GEARs that could potentially serve as therapeutic targets. Undoubtedly, future efforts are still

required to utilize protein-protein interaction networks and pathway databases to construct a

new network based on CSN.

In conclusion, our study utilized the MEMORY algorithm to identify GEARs in 15 cancer types,

highlighting the significance of mitosis and immunity in cancer prognosis. Our findings

demonstrate that GEARs possess substantial biological significance beyond their role as

prognostic biomarkers. This study provides valuable guidance for establishing standards for

survival analysis evaluation and holds potential for the development of novel therapeutic

strategies.

Methods

Datasets

The gene expression profiles and clinical information of 33 cancers were obtained from the TCGA database and downloaded by the GDC data website (https: //portal.gdc.cancer.gov/). All gene expression data and survival data were integrated and normalized.

Multi-gradient permutation survival analysis

Gene expression data of cancer patients were obtained from the RNA-seq expression matrix of TCGA after TPM normalization. We randomly sampled the gene expression data according to the gradient. The sampling strategy was as follows: Ten gradient increases in sample size were pre-set, ranging from approximately 10% to about 100%, with intervals of 10%. Random samples were taken from total samples of each cancer 1000 times, based on each pre-set sample size. Multiple sampling matrix was obtained after the sampling strategy was performed in each gradient. The survival analysis was performed for every sampling matrix by "survival" and "survminer". "survival" and "survminer" are R packages for survival analysis. The survival analysis method was as follows according to the median expression value of every gene, every sampling matrix was divided into a high and low expression group and performed survival analysis by "survival" and "survminer". We used 1 for significant survival analysis results (P < 0.05) and 0 for non-significant results (P > 0.05). The survival analysis matrices were obtained after these processes.

Construction of core survival network

The survival analysis matrices were integrated to a significant-probability matrix by a formula:

$$A_{ij} = \frac{\sum_{n=1}^{1000} k_{j \times i \times n}}{1000} \qquad (1)$$

where the A_{ij} is the value from row i and column j in the significant-probability matrix. We defined the sampling size k_j reached saturation when the max value of column j was equal to 1 in a significant-probability matrix. The least value of k_j was selected, and the genes with their

corresponding A_{ij} greater than 0.8 were extracted as GEAR.

We also defined survival analysis similarity (*SAS*) as the similarity of the effect on patient

prognosis in two genes. The *SAS* computational formula is as follows:

$$SAS = \frac{A \cap B}{A \cup B - 2(A \cap B) + 1} \qquad (2)$$

Where *A* and *B* are results of *1000* survival analyses at the first sampling gradient, which are

sequences of *0* and *1* of length *1000*. $A \cap B$ means the number of events that are simultaneously

1 in both A and B. $A \cup B$ means the size of the union of *A* and *B*. The *SAS* of significant genes

with total genes was calculated, and the significant survival network was constructed. Then

some *SAS* values at the top of the rankings were extracted, and the *SAS* was visualized to a

network by Cytoscape[93]. The network was named core survival network (CSN). The degree of

each node was calculated, and the nodes with the top10 degrees were defined hub genes. The

effect of molecular classification by hub genes is indicated that 1000 to 2000 was a range that

the result of molecular classification was best. One thousand was chosen as the number of *SAS*.

Therefore, one thousand was selected as the number of *SAS* values for constructed significant

survival network.

Gene ontology enrichment analysis

Gene ontology has been used to classify genes based on functions. The gene functions were

divided three types, includes molecular function (MF), biological process (BP), and cellular

component (CC). ClusterProfiler is an R package for gene set enrichment analysis[94]. The gene

ontology enrichment analysis (GOEA) was processed in significant gene sets and hub genes by

ClusterProfiler.

Hub gene classification

Tumor samples were genotyped using RNA-seq data. The following steps outline the

genotyping and clustering methods used in this study. The expression matrix of hub genes

corresponding to various cancer types was extracted from the RNA-seq data. This involved

filtering the RNA-seq data to retain only the expression levels of the identified hub genes. To

normalize the expression data and reduce the impact of extreme values, a pseudo count of 1

was added to each expression value, and the resulting matrix M was log-transformed using the

formula: $M' = log_2(M + 1)$. The log-transformed matrix M' was then subjected to

hierarchical clustering. This clustering was performed using the Ward's method to minimize

the variance within clusters. The distance metric used was the Euclidean distance. The number

of clusters was determined based on the dendrogram cut-off.

Calculation of tumor mutation burden (TMB) and differential mutation

The TMB and differential mutation gene analysis were carried out to explore the differences

between different genotyped samples at the genomic level. The data are from the gene mutation

data of TCGA tumor samples, and the grouping information is from the hub gene hierarchical

clustering results. Maftools is an R package for the analysis of somatic variant data, which can

export results in form of charts and graph[95]. The calculation of TMB and difference mutation

analysis were used by the maftools.

Quantifying the effect of gene mutations for tumor cell viability

Gene dependence refers to the extent to which genes are essential for cell proliferation and

survival. The Cancer Dependency Map (Depmap, https://depmap.org/portal/) database

provides genome-wide gene dependence data for large number of tumor cell lines[96]. Gene

mutation data of cell lines were obtained from the CCLE database

(https://sites.broadinstitute.org/ccle). The genes appearing in the gene mutation data of cell

lines were extracted and sorted into mutation list. Each gene in the list was then analyzed for

survival-dependent differences. For each gene in the mutation list, we compared the gene

dependence score (S) between cell lines with and without the gene mutation. Specifically, cell

lines from the CCLE mutation dataset harboring a mutation in a given gene were classified into

one group, while cell lines without the mutation constituted the control group. S for the mutation

(S_m) and wild-type (S_{wt}) cell lines were then extracted from the DepMap database.

A two-sample t-test was used to assess the statistical significance of differences between S_m and

S_{wt}. The test yielded a P-value, with an alpha level set at 0.05, to determine the significance of

the difference in means. D_m was defined as gene mutation dependence, the computational

formula was as follows:

$$D_m = |\bar{s}_m - \bar{s}_{wt}| \qquad (3)$$

Subsequently, we conducted a standardization assessment of D_m, employing the following formula:

$$S_{dm} = \frac{2(\bar{s}_m - \bar{s}_{wt})}{\bar{s}_m + \bar{s}_{wt}} \qquad (4)$$

In this study, we categorized mutations identified in cell lines into three groups based on their impact on gene dependency. Functional mutations refer to the mutations that exhibit significant changes in gene dependency scores, with a P-value < 0.05 and $S_{dm} > 0.1$. Non-functional mutations were those without significant changes in gene dependency scores compared to wildtype. Subtype-associated functional mutations were a subset of functional mutations that show statistically significant differences in occurrence frequency across different LUAD subtypes.

Identification of drugs downregulating hub gene expression

The IC50 data of 76 compounds of LUAD cell lines were obtained from Genomics of Drug Sensitivity in Cancer (GDSC) database[97]. The hub gene expression data of A549 cell lines after treatment with 76 compounds was obtained from Connectivity MAP (cMAP, https://clue.io/) database, and the data were grouped by hierarchical clustering[98]. The drugs which were presented only in the hub gene expression suppression group were considered to effective against the LUAD cell.

Immune infiltration analysis

Immune infiltration data was calculated by quantiseq method[99]. Immunedeconv was an R package for unified access to computational methods for estimating immune Cell fractions from bulk RNA-seq data[100]. The data were from the gene expression data of TCGA tumor samples after TPM was standardized. Then the immune cell fractions of tumor samples were calculated by quantiseq method invoking immunedeconv.

Biological function score calculation

GSVA is an R package for calculate biological function score based on a single sample[101]. The

data was from TCGA in the calculation process, and the software package was GSVA. The immune score was T-cell infiltration rate of that calculated by quantiseq [99]. The gene sets that calculated the mitosis score were obtained from gene ontology (GO:0140014).

Acknowledgment

This work was supported by the National Key Research and Development Program of China (grants 2022YFA1103900 to H.J., F.L.; 2020YFA0803300 to H.J.); the National Natural Science Foundation of China (grants 82341002 to H.J., 32293192 to H.J., 82030083 to H.J., 82173340 to L.H., 82273400 to Y.J., 32100593 to X.T., 82203306 to W.H., 82372763 to X.W., 82303916 to C.G., 82303039 to Z.Q., 82141101 to F.L., 2022hwyq16 to F.L., 82372794 to F.L.); the Basic Frontier Scientific Research Program of Chinese Academy of Science (ZDBS-LY-SM006 to H.J.); the Innovative research team of high-level local universities in Shanghai (SSMU-ZLCX20180500 to H.J.); Science and Technology Commission of Shanghai Municipality (21ZR1470300 to L.H.).

Author Contributions

H.J. and F.L. conceived the idea and designed the analysis. X.C. performed all bioinformatics analyses. Y.Y., X.L. and L.C. provided technical assistance and helpful comments. H.J., F.L. and X.C. wrote the manuscript. All authors approved the final version.

Disclosure of conflicts of interest

The authors declare that there is no conflict of interests.

References

1. Tomczak, K., Czerwinska, P., and Wiznerowicz, M. (2015). The Cancer Genome Atlas (TCGA): an immeasurable source of knowledge. Contemp Oncol (Pozn) *19*, A68-77. 10.5114/wo.2014.47136.
2. Cancer Genome Atlas Research, N., Weinstein, J.N., Collisson, E.A., Mills, G.B., Shaw, K.R.,

Ozenberger, B.A., Ellrott, K., Shmulevich, I., Sander, C., and Stuart, J.M. (2013). The Cancer
Genome Atlas Pan-Cancer analysis project. Nat Genet *45*, 1113-1120. 10.1038/ng.2764.

3.   Liu, J., Lichtenberg, T., Hoadley, K.A., Poisson, L.M., Lazar, A.J., Cherniack, A.D., Kovatich,
A.J., Benz, C.C., Levine, D.A., Lee, A.V., et al. (2018). An Integrated TCGA Pan-Cancer
Clinical Data Resource to Drive High-Quality Survival Outcome Analytics. Cell *173*, 400-416
e411. 10.1016/j.cell.2018.02.052.

4.   Smeltzer, M.P., Faris, N.R., Ray, M.A., and Osarogiagbon, R.U. (2018). Association of
Pathologic Nodal Staging Quality With Survival Among Patients With Non-Small Cell Lung
Cancer After Resection With Curative Intent. JAMA Oncol *4*, 80-87.
10.1001/jamaoncol.2017.2993.

5.   Miyauchi, E., Morita, S., Nakamura, A., Hosomi, Y., Watanabe, K., Ikeda, S., Seike, M., Fujita,
Y., Minato, K., Ko, R., et al. (2022). Updated Analysis of NEJ009: Gefitinib-Alone Versus
Gefitinib Plus Chemotherapy for Non-Small-Cell Lung Cancer With Mutated EGFR. J Clin
Oncol, JCO2102911. 10.1200/JCO.21.02911.

6.   Teh, S.H., Uong, S., Lin, T.Y., Shiraga, S., Li, Y., Gong, I.Y., Herrinton, L.J., and Li, R.A. (2021).
Clinical Outcomes Following Regionalization of Gastric Cancer Care in a US Integrated Health
Care System. J Clin Oncol *39*, 3364-3376. 10.1200/JCO.21.00480.

7.   Gimotty, P.A., Van Belle, P., Elder, D.E., Murry, T., Montone, K.T., Xu, X., Hotz, S., Raines, S.,
Ming, M.E., Wahl, P., and Guerry, D. (2005). Biologic and prognostic significance of dermal
Ki67 expression, mitoses, and tumorigenicity in thin invasive cutaneous melanoma. J Clin
Oncol *23*, 8048-8056. 10.1200/JCO.2005.02.0735.

8.   Wu, Y., Lv, M., Qian, T., and Shen, Y. (2021). Correlation analysis of Ki67 and CK7 expression
with clinical characteristics and prognosis of postoperative cervical adenocarcinoma patients.
Ann Palliat Med *10*, 9544-9552. 10.21037/apm-21-1974.

9.   Mouabbi, J.A., Singareeka Raghavendra, A., Bassett, R.L., Jr., Hassan, A., Tripathy, D., and
Layman, R.M. (2023). Survival Outcomes in Patients With Hormone Receptor-Positive
Metastatic Breast Cancer With Low or No ERBB2 Expression Treated With Targeted Therapies
Plus Endocrine Therapy. JAMA Netw Open *6*, e2313017.
10.1001/jamanetworkopen.2023.13017.

10.   Kang, Y.J., Oh, S.J., Bae, S.Y., Kim, E.K., Lee, Y.J., Park, E.H., Jeong, J., Park, H.K., Suh, Y.J.,
and Kim, Y.S. (2023). Predictive biological factors for late survival in patients with HER2-
positive breast cancer. Sci Rep *13*, 11008. 10.1038/s41598-023-38200-y.

11.   Cai, C., Wang, X., Fu, Q., and Chen, A. (2022). The VEGF expression associated with prognosis
in patients with intrahepatic cholangiocarcinoma: a systematic review and meta-analysis. World
J Surg Oncol *20*, 40. 10.1186/s12957-022-02511-7.

12.   Zhang, C., Wang, L., Xiong, C., Zhao, R., Liang, H., and Luo, X. (2021). The role of vascular
endothelial growth factor as a prognostic and clinicopathological marker in osteosarcoma: a
systematic review and meta-analysis. J Orthop Surg Res *16*, 738. 10.1186/s13018-021-02888-
3.

13.   Lai, Y., OuYang, G., Sheng, L., Zhang, Y., Lai, B., and Zhou, M. (2021). Novel prognostic genes
and subclasses of acute myeloid leukemia revealed by survival analysis of gene expression data.
BMC Med Genomics *14*, 39. 10.1186/s12920-021-00888-0.

14.   Anton, R.C., Coffey, D.M., Gondo, M.M., Stephenson, M.A., Brown, R.W., and Cagle, P.T.
(2000). The expression of cyclins D1 and E in predicting short-term survival in squamous cell

carcinoma of the lung. Mod Pathol *13*, 1167-1172. 10.1038/modpathol.3880215.

15.   Esposito, V., Baldi, A., De Luca, A., Tonini, G., Vincenzi, B., Santini, D., Persichetti, P., Mancini,
A., Citro, G., Baldi, F., et al. (2005). Cell cycle related proteins as prognostic parameters in
radically resected non-small cell lung cancer. J Clin Pathol *58*, 734-739.
10.1136/jcp.2004.023531.

16.   Dworakowska, D., Jassem, E., Jassem, J., Boltze, C., Wiedorn, K.H., Dworakowski, R.,
Skokowski, J., Jaskiewicz, K., and Czestochowska, E. (2005). Prognostic value of cyclin D1
overexpression in correlation with pRb and p53 status in non-small cell lung cancer (NSCLC).
J Cancer Res Clin Oncol *131*, 479-485. 10.1007/s00432-004-0661-9.

17.   Freedman, L.S. (1982). Tables of the number of patients required in clinical trials using the
logrank test. Stat Med *1*, 121-129. 10.1002/sim.4780010204.

18.   Yao, X., Ireland, S.K., Pham, T., Temple, B., Chen, R., Raj, M.H., and Biliran, H. (2014). TLE1
promotes EMT in A549 lung cancer cells through suppression of E-cadherin. Biochem Biophys
Res Commun *455*, 277-284. 10.1016/j.bbrc.2014.11.007.

19.   Zheng, H., Tian, H., Yu, X., Ren, P., and Yang, Q. (2021). G protein gamma 7 suppresses
progression of lung adenocarcinoma by inhibiting E2F transcription factor 1. Int J Biol
Macromol *182*, 858-865. 10.1016/j.ijbiomac.2021.04.082.

20.   Chen, G., Wang, Q., and Wang, K. (2022). MicroRNA-218-5p affects lung adenocarcinoma
progression through targeting endoplasmic reticulum oxidoreductase 1 alpha. Bioengineered *13*,
10061-10070. 10.1080/21655979.2022.2063537.

21.   Xu, J., Zheng, H., Yuan, S., Zhou, B., Zhao, W., Pan, Y., and Qi, D. (2019). Overexpression of
ANLN in lung adenocarcinoma is associated with metastasis. Thorac Cancer *10*, 1702-1709.
10.1111/1759-7714.13135.

22.   Zhang, P., Li, S., Lv, C., Si, J., Xiong, Y., Ding, L., Ma, Y., and Yang, Y. (2018). BPI-9016M, a
c-Met inhibitor, suppresses tumor cell growth, migration and invasion of lung adenocarcinoma
via miR203-DKK1. Theranostics *8*, 5890-5902. 10.7150/thno.27667.

23.   Wang, T., Zhai, R., Lv, X., Wang, K., and Xu, J. (2020). LINC02418 promotes malignant
behaviors in lung adenocarcinoma cells by sponging miR-4677-3p to upregulate KNL1
expression. BMC Pulm Med *20*, 217. 10.1186/s12890-020-01229-0.

24.   Huo, S.F., Shang, W.L., Yu, M., Ren, X.P., Wen, H.X., Chai, C.Y., Sun, L., Hui, K., Liu, L.H.,
Wei, S.H., et al. (2020). STEAP1 facilitates metastasis and epithelial-mesenchymal transition
of lung adenocarcinoma via the JAK2/STAT3 signaling pathway. Biosci Rep *40*.
10.1042/BSR20193169.

25.   Zhao, Z., Li, J., Tan, F., Gao, S., and He, J. (2018). mTOR up-regulation of BEX4 promotes
lung adenocarcinoma cell proliferation by potentiating OCT4. Biochem Biophys Res Commun
500, 302-309. 10.1016/j.bbrc.2018.04.064.

26.   Fan, T., Liu, Y., Liu, H., Wang, L., Tian, H., Zheng, Y., Zheng, B., Xue, L., Li, C., and He, J.
(2022). Transmembrane Protein-Based Risk Model and H3K4me3 Modification Characteristics
in Lung Adenocarcinoma. Frontiers in Oncology *12*. 10.3389/fonc.2022.828814.

27.   Lee, S., Kang, H., Shin, E., Jeon, J., Youn, H., and Youn, B. (2021). BEX1 and BEX4 Induce
GBM Progression through Regulation of Actin Polymerization and Activation of YAP/TAZ
Signaling. Int J Mol Sci *22*. 10.3390/ijms22189845.

28.   Gao, W., Li, J.Z., Chen, S.Q., Chu, C.Y., Chan, J.Y., and Wong, T.S. (2016). Decreased brain-
expressed X-linked 4 (BEX4) expression promotes growth of oral squamous cell carcinoma. J

Exp Clin Cancer Res *35*, 92. 10.1186/s13046-016-0355-6.

29.   He, Y., Luo, Y., Zhang, D., Wang, X., Zhang, P., Li, H., Ejaz, S., and Liang, S. (2019). PGK1-
mediated cancer progression and drug resistance. Am J Cancer Res *9*, 2280-2302.

30.   Matsuura, K., Sawai, H., Ikeo, K., Ogawa, S., Iio, E., Isogawa, M., Shimada, N., Komori, A.,
Toyoda, H., Kumada, T., et al. (2017). Genome-Wide Association Study Identifies TLL1 Variant
Associated With Development of Hepatocellular Carcinoma After Eradication of Hepatitis C
Virus Infection. Gastroenterology *152*, 1383-1394. 10.1053/j.gastro.2017.01.041.

31.   Poplawski, P., Alseekh, S., Jankowska, U., Skupien-Rabian, B., Iwanicka-Nowicka, R.,
Kossowska, H., Fogtman, A., Rybicka, B., Boguslawska, J., Adamiok-Ostrowska, A., et al.
(2023). Coordinated reprogramming of renal cancer transcriptome, metabolome and secretome
associates with immune tumor infiltration. Cancer Cell Int *23*, 2. 10.1186/s12935-022-02845-y.

32.   Guo, M., Li, F., Zhao, L., Fang, Z., Yu, H., Songyang, Z., and Xiong, Y. (2023). Pan-cancer
investigation of C-to-U editing reveals its important role in cancer development and new targets
for cancer treatment. Front Oncol *13*, 1097667. 10.3389/fonc.2023.1097667.

33.   Povero, D., Johnson, S.M., and Liu, J. (2020). Hypoxia, hypoxia-inducible gene 2
(HIG2)/HILPDA, and intracellular lipolysis in cancer. Cancer Lett *493*, 71-79.
10.1016/j.canlet.2020.06.013.

34.   Erfani, S., Hua, H., Pan, Y., Zhou, B.P., and Yang, X.H. (2021). The Context-Dependent Impact
of Integrin-Associated CD151 and Other Tetraspanins on Cancer Development and Progression:
A Class of Versatile Mediators of Cellular Function and Signaling, Tumorigenesis and
Metastasis. Cancers (Basel) *13*. 10.3390/cancers13092005.

35.   Won, M., Park, K.A., Kim, S., Ju, E., Ko, Y., Yoo, H., Ro, H., Lee, J., Oh, J., Lee, E.G., et al.
(2022). ANKRD13a controls early cell-death checkpoint by interacting with RIP1 independent
of NF-kappaB. Cell Death Differ *29*, 1152-1163. 10.1038/s41418-021-00906-9.

36.   Xu, J., Chen, Q., Tian, K., Liang, R., Chen, T., Gong, A., Mathy, N.W., Yu, T., and Chen, X.
(2020). m6A methyltransferase METTL3 maintains colon cancer tumorigenicity by suppressing
SOCS2 to promote cell proliferation. Oncol Rep *44*, 973-986. 10.3892/or.2020.7665.

37.   Xu, C., Li, H., Zhang, L., Jia, T., Duan, L., and Lu, C. (2016). MicroRNA-1915-3p prevents the
apoptosis of lung cancer cells by downregulating DRG2 and PBX2. Mol Med Rep *13*, 505-512.
10.3892/mmr.2015.4565.

38.   Mead, T.J. (2022). ADAMTS6: Emerging roles in cardiovascular, musculoskeletal and cancer
biology. Front Mol Biosci *9*, 1023511. 10.3389/fmolb.2022.1023511.

39.   Yang, B.Y., Song, J.W., Sun, H.Z., Xing, J.C., Yang, Z.H., Wei, C.Y., Xu, T.Y., Yu, Z.N., Zhang,
Y.N., Wang, Y.F., et al. (2018). PSMB8 regulates glioma cell migration, proliferation, and
apoptosis through modulating ERK1/2 and PI3K/AKT signaling pathways. Biomed
Pharmacother *100*, 205-212. 10.1016/j.biopha.2018.01.170.

40.   Keshet, R., Lee, J.S., Adler, L., Iraqi, M., Ariav, Y., Lim, L.Q.J., Lerner, S., Rabinovich, S., Oren,
R., Katzir, R., et al. (2020). Targeting purine synthesis in ASS1-expressing tumors enhances the
response to immune checkpoint inhibitors. Nat Cancer *1*, 894-908. 10.1038/s43018-020-0106-
7.

41.   Palaparti, A., Baratz, A., and Stifani, S. (1997). The Groucho/Transducin-like Enhancer of split
Transcriptional Repressors Interact with the Genetically Defined Amino-terminal Silencing
Domain of Histone H3. Journal of Biological Chemistry *272*, 26604-26610.
10.1074/jbc.272.42.26604.

42. Lin, W., Wang, J., Ge, J., Zhou, R., Hu, Y., Xiao, L., Peng, Q., and Zheng, Z. (2023). The activity
of cuproptosis pathway calculated by AUCell algorithm was employed to construct cuproptosis
landscape in lung adenocarcinoma. Discover Oncology *14*. 10.1007/s12672-023-00755-7.
43. Li, X., Jiang, Y., Meisenhelder, J., Yang, W., Hawke, D.H., Zheng, Y., Xia, Y., Aldape, K., He,
J., Hunter, T., et al. (2016). Mitochondria-Translocated PGK1 Functions as a Protein Kinase to
Coordinate Glycolysis and the TCA Cycle in Tumorigenesis. Mol Cell *61*, 705-719.
10.1016/j.molcel.2016.02.009.
44. Fu, Q., and Yu, Z. (2020). Phosphoglycerate kinase 1 (PGK1) in cancer: A promising target for
diagnosis and therapy. Life Sci *256*, 117863. 10.1016/j.lfs.2020.117863.
45. Waldman, A.D., Fritz, J.M., and Lenardo, M.J. (2020). A guide to cancer immunotherapy: from
T cell basic science to clinical practice. Nat Rev Immunol *20*, 651-668. 10.1038/s41577-020-
0306-5.
46. Zhou, Z., Li, Y., Hao, H., Wang, Y., Zhou, Z., Wang, Z., and Chu, X. (2019). Screening Hub
Genes as Prognostic Biomarkers of Hepatocellular Carcinoma by Bioinformatics Analysis. Cell
Transplant *28*, 76S-86S. 10.1177/0963689719893950.
47. Hwang, L.H., Lau, L.F., Smith, D.L., Mistrot, C.A., Hardwick, K.G., Hwang, E.S., Amon, A.,
and Murray, A.W. (1998). Budding yeast Cdc20: a target of the spindle checkpoint. Science *279*,
1041-1044. 10.1126/science.279.5353.1041.
48. Uuskula-Reimand, L., and Wilson, M.D. (2022). Untangling the roles of TOP2A and TOP2B in
transcription and cancer. Sci Adv *8*, eadd4920. 10.1126/sciadv.add4920.
49. Lamers, F., van der Ploeg, I., Schild, L., Ebus, M.E., Koster, J., Hansen, B.R., Koch, T., Versteeg,
R., Caron, H.N., and Molenaar, J.J. (2011). Knockdown of survivin (BIRC5) causes apoptosis
in neuroblastoma via mitotic catastrophe. Endocr Relat Cancer *18*, 657-668. 10.1530/ERC-11-
0207.
50. Wittmann, T., Wilm, M., Karsenti, E., and Vernos, I. (2000). TPX2, A novel xenopus MAP
involved in spindle pole organization. J Cell Biol *149*, 1405-1418. 10.1083/jcb.149.7.1405.
51. Kufer, T.A., Sillje, H.H., Korner, R., Gruss, O.J., Meraldi, P., and Nigg, E.A. (2002). Human
TPX2 is required for targeting Aurora-A kinase to the spindle. J Cell Biol *158*, 617-623.
10.1083/jcb.200204155.
52. Cancer Genome Atlas Research, N. (2014). Comprehensive molecular profiling of lung
adenocarcinoma. Nature *511*, 543-550. 10.1038/nature13385.
53. Zheng, F., Kelly, M.R., Ramms, D.J., Heintschel, M.L., Tao, K., Tutuncuoglu, B., Lee, J.J., Ono,
K., Foussard, H., Chen, M., et al. (2021). Interpretation of cancer mutations using a multiscale
map of protein systems. Science *374*, eabf3067. 10.1126/science.abf3067.
54. Wang, Q., Shi, Y.L., Zhou, K., Wang, L.L., Yan, Z.X., Liu, Y.L., Xu, L.L., Zhao, S.W., Chu,
H.L., Shi, T.T., et al. (2018). PIK3CA mutations confer resistance to first-line chemotherapy in
colorectal cancer. Cell Death Dis *9*, 739. 10.1038/s41419-018-0776-6.
55. Shibata, T., Kokubu, A., Tsuta, K., and Hirohashi, S. (2009). Oncogenic mutation of PIK3CA
in small cell lung carcinoma: a potential therapeutic target pathway for chemotherapy-resistant
lung cancer. Cancer Lett *283*, 203-211. 10.1016/j.canlet.2009.03.038.
56. Hanker, A.B., Pfefferle, A.D., Balko, J.M., Kuba, M.G., Young, C.D., Sanchez, V., Sutton, C.R.,
Cheng, H., Perou, C.M., Zhao, J.J., et al. (2013). Mutant PIK3CA accelerates HER2-driven
transgenic mammary tumors and induces resistance to combinations of anti-HER2 therapies.
Proc Natl Acad Sci U S A *110*, 14372-14377. 10.1073/pnas.1303204110.

57.   Eng, J., Woo, K.M., Sima, C.S., Plodkowski, A., Hellmann, M.D., Chaft, J.E., Kris, M.G., Arcila,
M.E., Ladanyi, M., and Drilon, A. (2015). Impact of Concurrent PIK3CA Mutations on
Response to EGFR Tyrosine Kinase Inhibition in EGFR-Mutant Lung Cancers and on Prognosis
in Oncogene-Driven Lung Adenocarcinomas. J Thorac Oncol *10*, 1713-1719.
10.1097/JTO.0000000000000671.

58.   Thorsson, V., Gibbs, D.L., Brown, S.D., Wolf, D., Bortone, D.S., Ou Yang, T.H., Porta-Pardo,
E., Gao, G.F., Plaisier, C.L., Eddy, J.A., et al. (2018). The Immune Landscape of Cancer.
Immunity *48*, 812-830 e814. 10.1016/j.immuni.2018.03.023.

59.   Meigs, T.E., Fedor-Chaiken, M., Kaplan, D.D., Brackenbury, R., and Casey, P.J. (2002).
Galpha12 and Galpha13 negatively regulate the adhesive functions of cadherin. J Biol Chem
277, 24594-24600. 10.1074/jbc.M201984200.

60.   An, Y., Adams, J.R., Hollern, D.P., Zhao, A., Chang, S.G., Gams, M.S., Chung, P.E.D., He, X.,
Jangra, R., Shah, J.S., et al. (2018). Cdh1 and Pik3ca Mutations Cooperate to Induce Immune-
Related Invasive Lobular Carcinoma of the Breast. Cell Rep *25*, 702-714 e706.
10.1016/j.celrep.2018.09.056.

61.   Zeisberg, M., and Neilson, E.G. (2009). Biomarkers for epithelial-mesenchymal transitions. J
Clin Invest *119*, 1429-1437. 10.1172/JCI36183.

62.   Yang, J., Mani, S.A., Donaher, J.L., Ramaswamy, S., Itzykson, R.A., Come, C., Savagner, P.,
Gitelman, I., Richardson, A., and Weinberg, R.A. (2004). Twist, a master regulator of
morphogenesis, plays an essential role in tumor metastasis. Cell *117*, 927-939.
10.1016/j.cell.2004.06.006.

63.   Egner, J.R. (2010). AJCC cancer staging manual. Jama *304*, 1726-1727.

64.   Rebello, R.J., Oing, C., Knudsen, K.E., Loeb, S., Johnson, D.C., Reiter, R.E., Gillessen, S., Van
der Kwast, T., and Bristow, R.G. (2021). Prostate cancer. Nat Rev Dis Primers *7*, 9.
10.1038/s41572-020-00243-0.

65.   Lee, J.K., Lee, J., Go, H., Lee, C.G., Kim, S., Kim, H.S., Cho, H., Choi, K.S., Ha, G.H., and
Lee, C.W. (2016). Oncogenic microtubule hyperacetylation through BEX4-mediated sirtuin 2
inhibition. Cell Death Dis *7*, e2336. 10.1038/cddis.2016.240.

66.   Lee, J.K., Ha, G.H., Kim, H.S., and Lee, C.W. (2018). Oncogenic potential of BEX4 is conferred
by Polo-like kinase 1-mediated phosphorylation. Exp Mol Med *50*, 1-12. 10.1038/s12276-018-
0168-0.

67.   Beroukhim, R., Mermel, C.H., Porter, D., Wei, G., Raychaudhuri, S., Donovan, J., Barretina, J.,
Boehm, J.S., Dobson, J., Urashima, M., et al. (2010). The landscape of somatic copy-number
alteration across human cancers. Nature *463*, 899-905. 10.1038/nature08822.

68.   Davoli, T., Xu, A.W., Mengwasser, K.E., Sack, L.M., Yoon, J.C., Park, P.J., and Elledge, S.J.
(2013). Cumulative haploinsufficiency and triplosensitivity drive aneuploidy patterns and shape
the cancer genome. Cell *155*, 948-962. 10.1016/j.cell.2013.10.011.

69.   Girish, V., Lakhani, A.A., Thompson, S.L., Scaduto, C.M., Brown, L.M., Hagenson, R.A.,
Sausville, E.L., Mendelson, B.E., Kandikuppa, P.K., Lukow, D.A., et al. (2023). Oncogene-like
addiction to aneuploidy in human cancers. Science *381*, eadg4521. 10.1126/science.adg4521.

70.   Bosco, N., Goldberg, A., Zhao, X., Mays, J.C., Cheng, P., Johnson, A.F., Bianchi, J.J., Toscani,
C., Di Tommaso, E., Katsnelson, L., et al. (2023). KaryoCreate: A CRISPR-based technology
to study chromosome-specific aneuploidy by targeting human centromeres. Cell *186*, 1985-
2001 e1919. 10.1016/j.cell.2023.03.029.

71.   Ha, S.Y., Choi, M., Lee, T., and Park, C.K. (2016). The Prognostic Role of Mitotic Index in
Hepatocellular Carcinoma Patients after Curative Hepatectomy. Cancer Res Treat *48*, 180-189.
10.4143/crt.2014.321.

72.   Zhou, C., Wang, Y., Liu, X., Liang, Y., Fan, Z., Jiang, T., Wang, Y., and Wang, L. (2018).
Molecular profiles for insular low-grade gliomas with putamen involvement. J Neurooncol *138*,
659-666. 10.1007/s11060-018-2837-1.

73.   Makinen, J.M., Laitakari, K., Johnson, S., Makitaro, R., Bloigu, R., Paakko, P., Lappi-Blanco,
E., and Kaarteenaho, R. (2017). Histological features of malignancy correlate with growth
patterns and patient outcome in lung adenocarcinoma. Histopathology *71*, 425-436.
10.1111/his.13236.

74.   Byrne, A., Savas, P., Sant, S., Li, R., Virassamy, B., Luen, S.J., Beavis, P.A., Mackay, L.K.,
Neeson, P.J., and Loi, S. (2020). Tissue-resident memory T cells in breast cancer control and
immunotherapy responses. Nat Rev Clin Oncol *17*, 341-348. 10.1038/s41571-020-0333-y.

75.   Liu, W., Sun, L., Zhang, J., Song, W., Li, M., and Wang, H. (2021). The landscape and
prognostic value of immune characteristics in uterine corpus endometrial cancer. Biosci Rep *41*.
10.1042/BSR20202321.

76.   Ducreux, M., Cuhna, A.S., Caramella, C., Hollebecque, A., Burtin, P., Goere, D., Seufferlein,
T., Haustermans, K., Van Laethem, J.L., Conroy, T., et al. (2015). Cancer of the pancreas: ESMO
Clinical Practice Guidelines for diagnosis, treatment and follow-up. Ann Oncol *26 Suppl 5*, v56-
68. 10.1093/annonc/mdv295.

77.   Cardoso, F., Kyriakides, S., Ohno, S., Penault-Llorca, F., Poortmans, P., Rubio, I.T., Zackrisson,
S., Senkus, E., and clinicalguidelines@esmo.org, E.G.C.E.a. (2019). Early breast cancer:
ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-updagger. Ann Oncol
30, 1194-1220. 10.1093/annonc/mdz173.

78.   Oaknin, A., Bosse, T.J., Creutzberg, C.L., Giornelli, G., Harter, P., Joly, F., Lorusso, D., Marth,
C., Makker, V., Mirza, M.R., et al. (2022). Endometrial cancer: ESMO Clinical Practice
Guideline for diagnosis, treatment and follow-up. Ann Oncol *33*, 860-877.
10.1016/j.annonc.2022.05.009.

79.   Vogel, A., Cervantes, A., Chau, I., Daniele, B., Llovet, J.M., Meyer, T., Nault, J.C., Neumann,
U., Ricke, J., Sangro, B., et al. (2018). Hepatocellular carcinoma: ESMO Clinical Practice
Guidelines for diagnosis, treatment and follow-up. Ann Oncol *29*, iv238-iv255.
10.1093/annonc/mdy308.

80.   Stupp, R., Brada, M., van den Bent, M.J., Tonn, J.C., Pentheroudakis, G., and Group, E.G.W.
(2014). High-grade glioma: ESMO Clinical Practice Guidelines for diagnosis, treatment and
follow-up. Ann Oncol *25 Suppl 3*, iii93-101. 10.1093/annonc/mdu050.

81.   Postmus, P.E., Kerr, K.M., Oudkerk, M., Senan, S., Waller, D.A., Vansteenkiste, J., Escriu, C.,
Peters, S., and Committee, E.G. (2017). Early and locally advanced non-small-cell lung cancer
(NSCLC): ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. Ann
Oncol *28*, iv1-iv21. 10.1093/annonc/mdx222.

82.   Gui, C., Kosteniuk, S.E., Lau, J.C., and Megyesi, J.F. (2018). Tumor growth dynamics in
serially-imaged low-grade glioma patients. J Neurooncol *139*, 167-175. 10.1007/s11060-018-
2857-x.

83.   Alvarez, M., Benhammou, J.N., Darci-Maher, N., French, S.W., Han, S.B., Sinsheimer, J.S.,
Agopian, V.G., Pisegna, J.R., and Pajukanta, P. (2022). Human liver single nucleus and single

cell RNA sequencing identify a hepatocellular carcinoma-associated cell-type affecting survival.
Genome Med *14*, 50. 10.1186/s13073-022-01055-5.

84.   Infante, M., Berghmans, T., Heuvelmans, M.A., Hillerdal, G., and Oudkerk, M. (2013). Slow-
growing lung cancer as an emerging entity: from screening to clinical management. Eur Respir
J *42*, 1706-1722. 10.1183/09031936.00186212.

85.   Hoshino, T. (1984). A commentary on the biology and growth kinetics of low-grade and high-
grade gliomas. J Neurosurg *61*, 895-900. 10.3171/jns.1984.61.5.0895.

86.   Sajjadi, E., Venetis, K., Piciotti, R., Invernizzi, M., Guerini-Rocco, E., Haricharan, S., and Fusco,
N. (2021). Mismatch repair-deficient hormone receptor-positive breast cancers: Biology and
pathological characterization. Cancer Cell Int *21*, 266. 10.1186/s12935-021-01976-y.

87.   Doghri, R., Houcine, Y., Boujelbene, N., Driss, M., Charfi, L., Abbes, I., Mrad, K., and Sellami,
R. (2019). Mismatch Repair Deficiency in Endometrial Cancer: Immunohistochemistry
Staining and Clinical Implications. Appl Immunohistochem Mol Morphol *27*, 678-682.
10.1097/PAI.0000000000000641.

88.   Zhang, Q., Sun, T.Y., Kang, P.M., Qian, K., Deng, B., Zhou, J.H., Wang, R.W., Jiang, B., Li, K.,
Liu, F., et al. (2016). Combined analysis of rearrangement of ALK, ROS1, somatic mutation of
EGFR, KRAS, BRAF, PIK3CA, and mRNA expression of ERCC1, TYMS, RRM1, TUBB3,
EGFR in patients with non-small cell lung cancer and their clinical significance. Cancer
Chemoth Pharm *77*, 583-593. 10.1007/s00280-016-2969-y.

89.   Donev, I.S., Wang, W., Yamada, T., Li, Q., Takeuchi, S., Matsumoto, K., Yamori, T., Nishioka,
Y., Sone, S., and Yano, S. (2011). Transient PI3K inhibition induces apoptosis and overcomes
HGF-mediated resistance to EGFR-TKIs in EGFR mutant lung cancer. Clin Cancer Res *17*,
2260-2269. 10.1158/1078-0432.CCR-10-1993.

90.   Wang, G., Xu, D., Zhang, Z., Li, X., Shi, J., Sun, J., Liu, H.Z., Li, X., Zhou, M., and Zheng, T.
(2021). The pan-cancer landscape of crosstalk between epithelial-mesenchymal transition and
immune evasion relevant to prognosis and immunotherapy response. NPJ Precis Oncol *5*, 56.
10.1038/s41698-021-00200-4.

91.   Brown, T.J., Bota, D.A., van Den Bent, M.J., Brown, P.D., Maher, E., Aregawi, D., Liau, L.M.,
Buckner, J.C., Weller, M., Berger, M.S., and Glantz, M. (2019). Management of low-grade
glioma: a systematic review and meta-analysis. Neurooncol Pract *6*, 249-258.
10.1093/nop/npy034.

92.   Chen, L., Ma, J., Zou, Z., Liu, H., Liu, C., Gong, S., Gao, X., and Liang, G. (2022). Clinical
characteristics and prognosis of patients with glioblastoma: A review of survival analysis of
1674 patients based on SEER database. Medicine (Baltimore) *101*, e32042.
10.1097/MD.0000000000032042.

93.   Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., Amin, N.,
Schwikowski, B., and Ideker, T. (2003). Cytoscape: a software environment for integrated
models of biomolecular interaction networks. Genome Res *13*, 2498-2504. 10.1101/gr.1239303.

94.   Yu, G., Wang, L.G., Han, Y., and He, Q.Y. (2012). clusterProfiler: an R package for comparing
biological themes among gene clusters. OMICS *16*, 284-287. 10.1089/omi.2011.0118.

95.   Mayakonda, A., Lin, D.C., Assenov, Y., Plass, C., and Koeffler, H.P. (2018). Maftools: efficient
and comprehensive analysis of somatic variants in cancer. Genome Res *28*, 1747-1756.
10.1101/gr.239244.118.

96.   Tsherniak, A., Vazquez, F., Montgomery, P.G., Weir, B.A., Kryukov, G., Cowley, G.S., Gill, S.,

835   Harrington, W.F., Pantel, S., Krill-Burger, J.M., et al. (2017). Defining a Cancer Dependency
836   Map. Cell *170*, 564-576 e516. 10.1016/j.cell.2017.06.010.

837 97. Yang, W., Soares, J., Greninger, P., Edelman, E.J., Lightfoot, H., Forbes, S., Bindal, N., Beare,
838   D., Smith, J.A., Thompson, I.R., et al. (2013). Genomics of Drug Sensitivity in Cancer (GDSC):
839   a resource for therapeutic biomarker discovery in cancer cells. Nucleic Acids Res *41*, D955-961.
840   10.1093/nar/gks1111.

841 98. Subramanian, A., Narayan, R., Corsello, S.M., Peck, D.D., Natoli, T.E., Lu, X., Gould, J., Davis,
842   J.F., Tubelli, A.A., Asiedu, J.K., et al. (2017). A Next Generation Connectivity Map: L1000
843   Platform and the First 1,000,000 Profiles. Cell *171*, 1437-1452 e1417.
844   10.1016/j.cell.2017.10.049.

845 99. Finotello, F., Mayer, C., Plattner, C., Laschober, G., Rieder, D., Hackl, H., Krogsdam, A.,
846   Loncova, Z., Posch, W., Wilflingseder, D., et al. (2019). Molecular and pharmacological
847   modulators of the tumor immune contexture revealed by deconvolution of RNA-seq data.
848   Genome Med *11*, 34. 10.1186/s13073-019-0638-6.

849 100. Sturm, G., Finotello, F., and List, M. (2020). Immunedeconv: An R Package for Unified Access
850   to Computational Methods for Estimating Immune Cell Fractions from Bulk RNA-Sequencing
851   Data. Methods Mol Biol *2120*, 223-232. 10.1007/978-1-0716-0327-7_16.

852 101. Hanzelmann, S., Castelo, R., and Guinney, J. (2013). GSVA: gene set variation analysis for
853   microarray and RNA-seq data. BMC Bioinformatics *14*, 7. 10.1186/1471-2105-14-7.

Figure Legends

Figure 1: MEMORY uncovers the enrichment of mitosis and immune signatures in multiple cancers. (A) Sample sizes, ranging from 10% to 100% with 10% intervals ("gradient"), were used for 1000 permutations of survival analyses of all 15 cancer types Each matrix was divided into high and low expression groups based on median gene expression values, and survival analyses were performed. Log-rank test significance results were coded as 1 for significant (P < 0.05) and 0 for non-significant (P > 0.05) outcomes, forming survival analysis matrices and the summarized significant probability matrix, which allowed the identification of GEARs. (B) The maximum of significant probability for each gradient sample size of every cancer type. Sample number rate refers to the percentage of samples in each sampling gradient compared to the total number of samples. (C) The pathways were enriched by GOEA based on the GEARs of each cancer type. The displayed pathways represent the top 5 most significant pathways for each cancer type. Mitosis-related pathways were marked in red whereas immune-related pathways were marked in blue.

Figure 2: Identification of hub genes in mitosis or immune-related cancers. (A) The SAS of all GEARs was calculated, and CSN was constructed. (B) The left nodes represent five different cancers, while the right nodes detail the functional types of hub-gene pathways. The height of the edges in the middle represents the proportion of hub-gene pathways corresponding to a specified functional type. (C-D) The CSN of LUAD and BRCA. The enlarged section represents the hub gene. (E) LUAD clustering based on hub gene expression, compared with inhibitor-based classification[52]. (F-G) LUAD clustering based on hub gene expression in comparison with the classic classification.

Figure 3: Different LUAD subgroups were characterized by unique genetic mutation profiles. (A) TMB analysis in three groups of LUAD samples. ***P < 0.0005. (B) Proportion of different oncogenic drivers including *KRAS*, *EGFR*, *BRAF*, *ERBB2* mutations, and *ALK*, *ROS1* fusions in three LUAD subgroups. (C) Kaplan-Meyer overall survival curves for EGFR-mutation samples of LUAD by hub gene subgroups. (D) Kaplan-Meyer overall survival curves for pan-negative LUAD samples by hub gene subgroups. (E-G) Comparison of top gene mutations in

three LUAD clusters, including ML vs. MM (C), MM vs. MH (D), ML vs MH (E).

Figure 4: PIK3CA mutation associates with mitosis and drug resistance in LUAD. (A) Cancer-

related gene mutations were annotated based on gene dependency scores data from Depmap

database. Functional mutations were indicated in green; functional (subtype-associated)

mutations were highlighted in red; non-functional mutations were indicated in gray. (B) The

analysis of the NeST differential pathways across three groups including ML vs. MM, MM vs.

MH, ML vs MH. (C) The heat map showed the hub gene expression of A549 after compounds

treatment from cMAP database[98]. Blue annotations meant 76 compounds that can inhibit hub

gene expression. (D) Comparison of the IC50 z-score of five tumor cell growth inhibitors for

A549 (*KRAS* mutation) and SW1573 (*KRAS* and *PIK3CA* mutation).

Figure 5: The hub gene classification of BRCA revealed a significant association between EMT

and immune infiltration. (A) The hub gene expression heatmap containing the result of hub

gene classification and PAM50 classification of BRCA[58]. (B) Comparison of molecular

classification based on hub genes with PAM50 classification. (C-E) Comparison of gene

mutations in three groups of BRCA samples, including IL vs. IM (C), IM vs. IH (D), IL vs. IH

(E). (F) Comparison of the total immune cell rate in three BRCA clusters. ***P < 0.0005. (G)

Comparison of *CDH1* expression in BRCA samples with wildtype or mutant *CDH1*. ***P <

0.0005. (H) A schematic diagram illustrating the correlation between the EMT score and

immune cell rate. The red line represents the fitting curve. The correlation analysis was

performed using Pearson correlation coefficient.

Figure 6: Mitosis and immune signatures predict patient prognosis at the pan-cancer level. (A)

The correlation of mitosis scores of 33 TCGA cancer types with *PIK3CA* expression was

analyzed. (B) The correlation of immune cell infiltration rate of 33 TCGA cancer types with

CDH1 expression was analyzed. (C) The mitosis and immune-related pathway scores of 33

cancer types were analyzed, and the median score was utilized as a threshold to categorize

patients for survival analysis. Among the 33 cancer types examined, 10 cancer types were

exclusively associated with the mitosis score, 2 cancer types were exclusively associated with

the immune score, and 5 cancer types showed a correlation with both mitosis and immunity

scores concurrently.

A

B

A

B

C