**Bycatch mitigation of endangered marine life**

Mireia Villafáfila García[1], Antonio Carpio[2,3*], Marga L Rivas[1]

[1] Department of Biology, Institute of Marine Science INMAR, University of Cádiz, Spain.

[2] Department of Zoology, Campus of Rabanales, University of Cordoba, 14071 Córdoba, Spain.

[3] Grupo de Sanidad y Biotecnología (SaBio), Instituto de Investigación en Recursos Cinegéticos,

IREC (UCLM-CSIC-JCCM), Ronda Toledo 12, 13071 Ciudad Real, Spain.

Corresponding author: Antonio J Carpio: a.carpio.camargo@gmail.com

**Author Contributions:** Conceptualization, A.J.C., M.L.R.: data cleansing and writing—
original draft preparation, M.V.; writing—review and supervision, A.J.C., M.L.R. All
authors have read and agreed to the published version of the manuscript.

**Acknowledgments:** M.V.G. is supported by the 'Plan propio de estímulo y apoyo a la
Investigación y Transferencia. INICIA-INV: Iniciación a la investigación (2022-2023)' of
the University of Cadiz. A.J.C. is supported by a 'Juan de la Cierva' contract (IJC2020-
042629-I) funded by MCIN/AEI/10.13039/501100011033 and by the European Union
Next GenerationEU/PRTR. M.L.R. was supported by Postdoctoral Research Contracts of
the Andalusian government (Spain) and FEDER EU funds. This work received financial
support from the Agencia Andaluza de Cooperación Internacional para el Desarrollo
AACID (projects EMEP 2022UI007_2022).

**ABSTRACT**

The fishing gear deployed by fishermen in seas and oceans throughout the world not only captures target species but also unintentionally ensnares non-target species, a phenomenon known as "by-catch". This unintended capture of marine life can represent significant challenges for the fishing industry, with adverse impacts on both the environment and species such as sea turtles, marine mammals, seabirds and elasmobranchs, which may be injured or even killed. To address this problem, the fishing industry has implemented regulations and mitigation measures. In this literature review, we have examined 389 articles published between 2010 and 2022 that assess the effectiveness of these measures. It has been demonstrated that the most effective measures are 'pingers' for marine mammals, 'TEDs' (Turtle Excluder Devices) for sea turtles, and 'BSLs' (Bird Scaring Lines), more commonly known as 'tori lines', for seabirds. The most complex case is that of elasmobranchs, and the most effective measure has yet to be discovered. This complexity arises from the ongoing targeted fishing of these species, resulting in less monitoring of their catches and, therefore, fewer surveys. Overall, we encourage the global implementation of these measures by the fishing industry in order to reduce by-catch in an attempt to ensure the future of many endangered species.

Keywords: Incidental fishing · cetaceans · LEDs · marine turtles · marine birds · rays · sea turtles · sharks

## 1. INTRODUCTION

Fishing poses a significant threat to marine vertebrates, including sea turtles, marine mammals, seabirds and elasmobranchs. This threat stems from the impact of the fishing industry on these species, both as a direct target and incidentally. The underlying cause of this issue is the extensive presence of fishing fleets on the various seas and oceans. In 2020, the global fishing fleet was estimated to consist of approximately 4.1 million vessels, with Asia being the predominant continent, accounting for a total of 2.68 million vessels (FAO, 2022). These vessels utilize a range of fishing gear, depending on their target species. The most common gear types include gillnets, longlines, purse seines with and without purse lines, trawl nets and traps (FAO, 2022).

This fishing gear has a substantial impact on oceans and their ecosystems, resulting in a series of adverse effects. These include an overexploitation of species, such as overfishing, which can lead to the decline and even collapse of populations, ultimately affecting marine biodiversity and ecosystems (Crowder et al. 2008). Certain fishing gear also destroys habitats, as occurs with that used for bottom trawling since it damages coral reefs, seagrass beds and seabeds (Crowder et al. 2008). Furthermore, the disruption of the food web caused by the decline of specific species as a result of overfishing can affect other species dependent on them as a food source, leading to a cascading effect known as the 'top-down' or 'bottom-up' effect (Crowder et al. 2008). Additionally, non-target or by-catch species are often captured alongside the intended target species when non-selective fishing methods are employed, and a wide variety of species is captured, including those not intended for capture or trade (Crowder et al. 2008). This includes species that are commercially undesirable owing to their lower value or failure to comply with size or weight regulations, along with protected or endangered

species (which may have serious implications for their conservation) and non-

commercial species that are not of interest for trade (Agardy, 2000). In some cases, these

non-target species are not only captured but also injured or even killed, and these

species include marine mammals, sea turtles, seabirds and elasmobranchs (Crowder et

al. 2008).

Certain management measures with which to mitigate the impacts on these species have

been implemented, including catch and net restrictions and the deployment of specific

fishing technologies tailored to the groups affected (Lucas and Berggren, 2023). In the

case of sea turtles, physical and visual techniques have been tested, such as the Turtle

Excluder Device (TED) (Warden, 2011) and Light Emitting Diodes (LEDs), which could also

be a good alternative as regards mitigating sea turtle by-catch. In the case of marine

mammals, acoustic, physical, visual and echolocation measures have also been studied,

and acoustic deterrent devices such as pingers have been used. However, some studies

have demonstrated that certain species may become habituated to pingers over time

(Moan and Bjørge, 2021). With regard to seabirds, olfactory, physical and visual

measures are used to mitigate by-catch, with Bird Scaring Lines (BSLs), more commonly

known as tori lines, being the visual measure that has been most widely used in longline

fisheries (Domingo et al. 2017). In the case of elasmobranchs, all types of measures have

been tried, i.e., acoustic, olfactory, physical, visual, echolocation and electrosensory.

However, the knowledge regarding their effectiveness is limited, as in the case of

employing tori lines to reduce shark by-catch (Seidu et al. 2022; Jiménez et al. 2019).

In order to obtain a global overview, the principal objective of this study is to conduct a

comprehensive review of the existing scientific literature focused on assessing the

effectiveness of specific mitigation measures as regards reducing the by-catch of endangered species groups, including sea turtles, marine mammals, seabirds and elasmobranchs. As secondary objectives, this study aims to: (i) identify the mitigation measures most commonly employed to minimize by-catch on a global scale; (ii) analyze the geographic distribution of by-catch in fisheries by country in order to evaluate variations, and (iii) provide an overview of the worldwide mitigation measurements that have proven to have the greatest effectiveness as regards reducing by-catch for each group. The overall objective of this research is to contribute with valuable insights into mitigating the effect of by-catch on endangered marine species and to shed light on measures that have proven to be the most successful, along with areas in which further research is required.

## 2. MATERIALS AND METHODS

### 2.1. Literature review

This review was carried out using the Scopus, Web of Science and Google Scholar search engines to search for publications containing citation indices spanning from 2010 to 2022. The keywords used were a combination of "bycatch" AND "fisheries" AND "mitigation" AND "measures" AND "sea turtles" or "marine mammals" or "seabirds" or "elasmobranchs". These four groups of marine vertebrates were selected as the objective of this study. The data collected were classified according to the animal class, species name in English, scientific name, UICN category, the mitigation measure used, the percentage of reduction in by-catch achieved, and finally, the reference of each paper (Table 1 SI).

**2.2.    Species and areas**

We collected global data and information from scientific literature concerning the by-catch of the four marine vertebrate groups, such as sea turtles, marine mammals, seabirds and elasmobranchs, by fisheries in the different seas and oceans worldwide. Species of sea turtles included in the review belonged to the Cheloniidae and Dermochelyidae families. These species are distributed in seas and oceans throughout the world (except those in the Arctic and Antarctic), with the exception of Kemp's ridley turtle (*Lepidochelys kempii*), which is solely a resident of the Gulf of Mexico and the northeastern coast of North America (Wibbels and Bevan, 2019).

Marine mammals such as odontocetes and mysticetes (families: Phocoenidae, Balaenopteridae, Delphinidae, Iniidae/Pontoporiidae), Sirenian and Pinniped (families: Dugongidae, Phocidae and Otariidae) were also included in the review. This group is, as a whole, widely distributed, as there are species that inhabit all the oceans in the world (including those in the Arctic and Antarctic), as in the case of the humpback whale (*Megaptera novaeangliae*) (Cooke, 2018). However, the majority of species are concentrated in the Atlantic, Pacific and Indian oceans, as is the case of the common dolphin (*Delphinus delphis*) (Braulik et al. 2021) and bottlenose dolphin (*Tursiops truncatus*) (Wells et al. 2019).

The seabird species included in our review belong to families Diomedeidae, Procellariidae, Spheniscidae, Anatidae, Phalacrocoracidae, Laridae, Sulidae, Stercorariidae and Oceanitidae. These birds are distributed across all continents, oceans and seas worldwide. Notably, the Mediterranean Sea serves as a prominent breeding and feeding area for several endangered species, such as the Balearic shearwater

(*Puffinus mauretanicus*) (BirdLife International. 2018a) and the Yelkouan shearwater

(*Puffinus yelkouan*) (BirdLife International. 2018b).

The review also encompassed elasmobranchs, which include sharks and rays from

diverse families including Sphyrnidae, Triakidae, Carcharhinidae, Myliobatidae,

Squalidae, Rajidae, Somniosidae, Rhinobatidae, Brachaeluridae, Lamnidae,

Torpedinidae, Scyliorhinidae, Alopiidae, Pseudocarcharidae, Dasyatidae, Mobulidae,

Glaucostegidae and Pentanchidae. Many of the species within this group are commonly

found in coastal areas and inhabit the oceanic zone delimited by both tropics (Cancer

and Capricorn). However, there are exceptions, such as the Greenland shark (*Somniosus*

*microcephalus*), which inhabits the Arctic Circle (Kulka et al. 2020).

Overall, most of the sea turtle, marine mammal, seabird and elasmobranch species

included in the study are classified by the UICN Red List of Threatened Species (UICN,

2023) as being Least concern (LC), Near threatened (NT), Vulnerable (VU), Endangered

(EN), Critically Endangered (CR) and Data Deficient (DD) (Methods, Table 2 SI).

**2.3.    Statistical analysis**

In order to verify the nature of the data obtained and compiled (Table 1 SI), the normality

and homoscedasticity of the data were analyzed by employing R commander (version

2.7-0) using the Shapiro-Wilk test and Bartlett test, respectively. The data did not meet

these criteria in any of the cases, and we therefore employed a non-parametric Kruskal-

Wallis's test. This test was used to examine whether there was a relationship (at the

global level) between the percentage of reduction in by-catch and fishing areas, along

with the percentage of reduction in by-catch per species group. In both cases, statistical

significance was considered when $p < 0.01$. In order to carry out an individual analysis of

the relationship between the efficiency of mitigation measures among the fisheries

involved by group, a Goodness-of-Fit G-test was carried out by employing the R Studio

program (version 4.0.2), using an R Script (Mangiafico, 2015) and the "DescTools" (Andri

et al. 2021) and "RVAideMemoire" packages (Hervé, 2023).

**3. RESULTS AND DISCUSSION**

**3.1.    Overview of published literature**

Of a total of 389 studies, 316 were excluded on the basis of specific criteria (Figure 1 SI).

A total of 73 papers were eventually selected in order to analyze by-catch by year and

fishery area. These are shown in Table 1 SI as follows: sea turtles in green, marine

mammals in blue, seabirds in orange and elasmobranchs in purple. However, when

attempting to assess the number of trials of fishing gear, it was possible to find only 31

studies that provided information on the number of trials (8140). Furthermore, studies

related to proofs of concept or carried out in laboratories were not taken into account.

The number of papers published each year that were found thanks to the literature

search are shown in (Table 3 SI), with a maximum of 9 in 2018 and 2020. In contrast, only

3 studies were found in 2022 and 2 in 2010.

The number of studies assessing by-catch by continent is shown in Figure 2 SI. The

continent for which most studies had been conducted was America, with 28 studies,

followed by Europe with 22. Within America, the United States carried out the largest

number of studies, with 7 in total. In Europe, studies were conducted by 14 different

countries, unlike that which occurred with Oceania, where all the studies were carried

out in Australia. This figure also shows that there is a lack of studies in developing nations

or locations in which small-scale fisheries (SSFs) are widespread, as is the case of Africa,

where only 8 studies have been conducted. With regard to the number of trials carried

out with each type of fishing gear, gillnets dominated the count with a total of 7636 trials,

distantly followed by set net with 273 trials (Figure 3 SI).

**3.2. Worldwide mitigation measures**

Several mitigation measures with which to minimize the by-catch of these megafauna

species have been tested in different fisheries around the world. A summary of the

mitigation measures, categorized by the sensory system involved for each group, is

provided in Table 1. In the case of sea turtles, the most commonly used olfactory

measure involved using an alternative bait type. With regard to visual measures, LED

lights emerged as a promising alternative, while in the case of physical measures, various

escape options such as TEDs were frequently employed. In the case of odoncetes,

mysticetes and other marine mammals such as the harbour seal (*Phoca vitulina*) and the

Californian sea lion (*Zalophus californianus),* the measures most frequently used were

pingers, a type of Acoustic Deterrent Devices (ADDs). With regard to seabirds, the

mitigation measure that predominated was the tori line, a type of seabird scaring device

(Gilman et al. 2021) consisting of a line towed from a high point at the aft of a vessel,

from which several streamers are attached to scare seabirds and prevent their access to

the critical area where baited hooks sink (Domingo et al. 2017). The main mitigation

measure used for elasmobranchs was electrosensory devices. However, there was no

single predominant measure, with SMART hooks (Grant et al. 2018; O'Connell et al.

2014), rare earth (Porsmoguer et al. 2015; Westlake et al. 2018), ferrite magnets

(Richards et al. 2018) and Electropositive Metal (EPM) Alloy (Godin et al. 2013;

Hutchinson et al. 2012) all being employed. However, it seems that tori lines have

obtained good results for some of the species in this group (Jiménez, Forselledo, et al.

2019).

## 3.3.    Mitigation measures by group

Overall, it was found that, globally, there were significant differences among the

percentage of reduction in by-catch per fishery area ($\chi$=36.33, df=19, p<0.01, n=20,

Kruskal-Wallis's test) (Figure 1) and in the percentage of success in by-catch per group

($\chi$=14.67, df=3, p<0.01, n=4, Kruskal-Wallis's test) (Figure 2A).

However, no significant differences were found as regards the percentage of reduction

in by-catch resulting from different mitigation types ($\chi$=10.17, df=4, p=0.04, n=6, Kruskal-

Wallis's test) (Figure 2B).

### 3.3.1.  Sea turtles

In the case of sea turtles, there was no significant difference among the percentage of

reduction in by-catch according to the fishery area (G=9.15, X-squared df=10, p=0.52,

n=11, G-test). Notably, the US Mid-Atlantic bottom trawl gear (Warden, 2011) and the

industrial trawling fishery of Gabon (Casale et al. 2017) attained the highest values of

reduction in by-catch per individual (mean ± SD, 0.97 ± NA and 1.0 ± NA, respectively).

Similarly, no significant differences were observed among sea turtle species (G=7.46, X-

squared df=4, p=0.11, n=24, G-test). However, the species with the highest percentages

of reduction in by-catch reduction were the loggerhead turtle (*Caretta caretta*) (n=112)

and the olive ridley turtle (*Lepidochelys olivacea*) (n=114) (97% and 100%, respectively).

This was achieved using physical measures, specifically TEDs (Figure 1).

### 3.3.1.1. Olfactory

One olfactory measure was investigated by testing the use of alternative bait types to mitigate incidental sea turtle by-catch in longline fisheries. The best results were attained by replacing squid bait with mackerel bait (Figure 3A), reducing by-catch by 88% and 85% for all turtles species (Coelho et al. 2012). In addition, by combining the effects of the mackerel by exchanging the traditional J-style hooks for two circle hooks (one non-offset and one with 10° offset), it was possible to obtain a 50-59% reduction in the by-catch (n=223) (Coelho et al. 2015).

### 3.3.1.2. Physical

Tori lines were used in a Uruguayan pelagic longline fishery to reduce the by-catch. A reduction of 18.1% was obtained for the loggerhead turtle (n=83), while one of 73.3% was obtained for the leatherback turtle (*Dermochelys coriacea*) (n=15) (Jiménez et al. 2019) (Figure 3A).

TEDs have also been tested in numerous trawl fisheries, as occurred in the US Mid-Atlantic, with a reduction in by-catch of 97% for the loggerhead turtle (n=112) (Warden, 2011). A reduction in by-catch of 100% was documented for Indian fisheries (Raghu et al. 2016), and a similar reduction of 100% was achieved for four sea turtle species (n=131) in Gabonese fisheries (Casale et al. 2017) (Figure 1).

### 3.3.1.3. Visual

Of all the possible visual mitigations methods for sea turtles, those that have been studied most are LED lights. However, there are also others, such as chemical lightsticks, physical models (predator cut-outs) and buoyless nets (Figure 3A).

The implementation of 500 nm green LEDs in an Indonesian small-scale coastal gillnet

fishery led to a reduction in multi-species sea turtle by-catch of 61.4% (n=10), and

specifically 59.5% of green turtle (*Chelonia mydas*) (n=14) (Gautama et al. 2022). This

same measure was also used in a Mexican gillnet fishery, obtaining a reduction of 50%

for the loggerhead turtle (n=17) (Senko et al., 2022) and 59% (n=85), 63.9% (n=125) and

48.8% (n=41) for green turtles (Wang et al. 2010; Ortiz et al. 2016; Kakai, 2019). Chemical

lightsticks, meanwhile, obtained a reduction of 40% for the green turtle (n=85) (Wang et

al. 2010). The implementation of 100-400 nm LED lights in a small-scale gillnet fishery

led to a decrease in green turtle by-catch of 93% (n=13) (Darquea et al. 2020). In Italian

and Mexican fisheries, the use of LEDs led to a reduction in loggerhead turtle by-catch

of 100% (n=18) (Lucchetti et al. 2019; Virgili et al. 2018), and a reduction in green turtle

by-catch of 39.7% (n=209) when using UV net illumination (Wang et al. 2013). Although

TEDs have attained the best results, more trials on LEDs should be included, as they seem

to be a good alternative by which to reduce the by-catch of these species.

**3.3.2. Marine mammals**

In the case of marine mammals, there were no significant differences among the

percentages of reduction in by-catch according to fishery areas (G=5.00, X-squared df=5,

p=0.41, n=6, G-test). Notably, the highest values of by-catch reduction per individual

were attained by the small set net Japanese fishery (Amano et al. 2017) and Norwegian

commercial fisheries (Moan and Bjørge, 2021) (mean ± SD, 10.0 ± NA and 13.0 ± 9.8,

respectively). Similarly, no significant differences were found among marine mammal

species (G=0.57, X-squared df=9, p=0.99, n=11, G-test). However, the narrow-ridged

finless porpoise (*Neophocaena asiaeorientalis*) (n=10) and harbour porpoise (*Phocoena*

*phocoena*) (n=20) were the species with the highest percentage of reduction in by-catch

(100% and 96.9%, respectively) when using acoustic measures, specifically pingers

(Figure 1).

### 3.3.2.1. Acoustic

Sensory technologies, specifically acoustic reflectors and pingers, were designed in the

late 1970s and 1980s to deter marine mammals in gillnet fisheries (Dawson, 1991).

The AQUAmark 100 pinger, which operates at between 20 and 160kHz, achieved a

reduction in narrow-ridged finless porpoise by-catch of 100% (n=10) (Amano et al. 2017)

(Figure 1). The long-term effectiveness of the Dukane Netmark 1000 Pinger, which

operates at 12-100 kHz harmonics, was assessed in a gillnet fishery (Carretta and Barlow,

2011). However, only the by-catch of two species decreased, specifically the common

dolphin by 47.4% and the Northern elephant seal (*Mirounga angustirostris*) by 80.8%

(n=164). When used in a driftnet fishery, the same pinger achieved reductions of

between 18.2% and 100%, depending on the species (Mangel et al. 2013).

The evaluation of two pingers in gillnet fisheries, i.e., the Banana pinger by Fishtek

Marine Industries (operating at 50-120 kHz, 154 dB) and the Dolphin pinger by Future

Oceans (operating at 70 kHz, 132 dB), attained positive results, with reductions of

96.90% and 33.50% (Moan and Bjørge, 2021) (Figure 1). Furthermore, the

implementation of these devices did not have a major negative impact on their daily

fishing operations and contributed to the reduction in marine mammal by-catch.

### 3.3.2.2. Physical

Berninsone et al. (2020) replaced gillnets with longlines in order to minimize the by-catch

of franciscana (*Pontoporia blainvillei*), which it is considered the most threatened

cetacean in the South Western Atlantic (Negri et al. 2012; Bordino and Albareda, 2004).

This study decreased the by-catch of this species by 90% (n=85) (Figure 3B), thus showing

that this method is an excellent alternative. However, only one study was carried out,

which makes it difficult to generalize its effectiveness to other areas.

**3.3.2.3.   Visual**

Unlike that which occurs with sea turtles, visual mitigation measures are not commonly

used to prevent marine mammal by-catch. Tori lines and Bird Line Weighting (BLW) have

been tested in Uruguayan pelagic longline fisheries with no significant results (Jiménez

et al. 2019). In order to evaluate the real effect of these measures on marine mammals,

further research is consequently necessary.

**3.3.2.4.   Echolocation reflection**

Mysticetes and odontocetes use echolocation, which helps them to determine the

location of objects in the sea. A mitigation measure consisting of adding acrylic glass

spheres to a gillnet has consequently been developed in order to reduce the by-catch of

the harbour porpoise (n=5), obtaining a reduction of 60% (Kratzer et al. 2021). Another

measure tested was the modification of two types of nets, a barium sulfate net and a

stiff nylon net (Bordino et al. 2013), but a reduction of only 7.4% (n=54) was obtained

(Figure 3B).

**3.3.3.  Seabirds**

With regard to seabirds, the percentage of reduction in by-catch was not significantly

different among fishery areas (G=7.84, X-squared df=5, p=0.16, n=6, G-test). Similarly, no

significant differences were found among seabird species (G=1.58, X-squared df=16, p=1,

n=20, G-test). Notably, the species with the highest percentage of by-catch reduction

were the Atlantic yellow-nosed albatross (*Thalassarche chlororhynchos*) (n=43), the

black-browed albatross (*Thalassarche melanophris*) (n=22) and the white-chinned petrel

(*Procellaria aequinoctialis*) (n=486) (100%, 100% and 97.7%, respectively) when using

visual measures, and specifically tori lines (Figure 1).

### 3.3.3.1. Olfactory

The main olfactory mitigation measures used to minimize seabird by-catch were offal

discard management (Kuepfer et al. 2022; Collins et al. 2021; Rollinson et al. 2017),

thawed bait (Collins et al. 2021; Rollinson et al. 2017), blue-dyed bait (Gilman et al.

2021), artificial bait (Cortés and González-Solís, 2018) and replacing squid with mackerel

as bait (Gonzalez et al. 2012; Li et al. 2012) (**¡Error! No se encuentra el origen de la r**

**eferencia.**). These measures were, in certain instances, reinforced with non-sensory

methods, such as night setting (Collins et al. 2021; Rollinson et al. 2017), seasonal

closures (Collins et al. 2021), hook management (Collins et al. 2021) and the limitation

of by-catch rates per year (Rollinson et al. 2017).

Despite the variety of existing measures, there is limited information regarding their

effectiveness as regards reducing seabird by-catch and their effect on commercial

catches. For instance, the evaluation of artificial bait demonstrated a reduction in target

catches of 77% when compared to control lines (Cortés and González-Solís, 2018), but

sample sizes were not included in the study.

### 3.3.3.2. Physical

In order to reduce the by-catch rate of seabirds by using physical measures, the increase

in the sink rate of baited hooks by reducing the distance between the hook and the

weight of the branch lines (65g) was tested in a pelagic longline fishery (Jiménez et al.

2019), obtaining a reduction of 42.5%. Others studies propose the introduction of BLW

as a mitigation measure, such as that by Paterson et al. (2019), which was carried out in

a demersal longline fishery where a reduction in the by-catch was from 90.9% to 100%.

**3.3.3.3.    Visual**

The measures most commonly used in the case of seabirds are those of visual mitigation

and include techniques such as LEDs (Bielli et al. 2020; Mangel et al. 2018; Field et al.

2019), high contrast panels (Field et al. 2019; Oliveira et al. 2021), buoys with looming

eyes (Rouxel et al. 2021), night setting (Cortés and González-Solís, 2018) and tori lines

(Cortés and González-Solís, 2018; Gilman et al. 2021) (**¡Error! No se encuentra el origen de**

**la referencia.**).

The implementation of 500 nm LEDs was positive as regards reducing the by-catch of 4

species (n=46), with a reduction of 84% (Bielli et al. 2020), while green LEDs led to a

reduction of 85.1% (Mangel et al. 2018). However, in the study carried out by Field et al.

(2019), the efficacy of two types of 500 nm LEDs (constant green lights and flashing white

LED lights) achieved a reduction of only 32.6% (n=43), although the use of high contrast

panels reduced the by-catch of species by 50.8% (n=65). New devices such as the

"Looming eyes buoy" (LEB) have also emerged, leading to a decrease in the by-catch of

seabirds species of 22% (n=5724) (Rouxel et al. 2021) (Figure 3C).

Another measure is that of night setting, which has been tested with the artisanal

demersal longliners of the Western Mediterranean. Although the sample sizes were

limited, the results obtained showed a reduction of 83.3% and 100% (n=19) (Figure 3C),

and an increase in sample testing is, therefore, recommended in order to ensure the

efficiency of this measure.

Finally, the most widely used measure with which to reduce the by-catch of seabirds is

that of tori lines, which have been shown to provide a significant reduction in the by-

catch of all birds species, from 97.7% to 100% (Domingo et al. 2017; Paterson et al. 2019)

(Figure 1).

### 3.3.4. Elasmobranchs

With regard to elasmobranchs, the percentage of reduction in by-catches was

significantly different according to the study site (G=11.83, X-squared df=2, p<0.01, n=3,

G-test), with a Uruguayan longline fishery attaining the highest values as regards a

reduction in by-catch (Jiménez et al. 2019). No significant differences were found among

elasmobranch species (G= 1.60, X-squared df=17, p=1, n=21, G-test), although the

species that attained the highest percentage of reduction in by-catch were the night

shark (*Carcharhinus signatus)* (n=38) and the smooth hammerhead (*Sphyrna zygaena*)

(n=190) (89.5% and 86.3%, respectively) when using visual measures such as tori lines

(Figure 1).

### 3.3.4.1. Acoustic

381 During the study of the effectiveness of pingers at reducing the by-catch of certain

382 species of marine mammals, their effect was also analyzed for elasmobranchs. No

383 significant differences in captures were attained when using Aquamark 100 and 200

384 pingers (Bilgin and Kose, 2018; Mangel et al. 2013).

### 3.3.4.2. Olfactory

386 Only one olfactory mitigation measure with which to reduce the by-catch of different

387 species of sharks and rays has been studied over a 13-year period (Coelho et al. 2012):

388  that of replacing squid with mackerel bait. However, the effectiveness of this measure

389  was limited.

### 3.3.4.3.  Physical

Physical measures by which to mitigate the by-catch of elasmobranchs include BLW

(Jiménez et al. 2019), Bycatch Reduction Devices (BRDs) (Gupta et al. 2020) and the use

of a 'tickler' (Kynoch et al. 2015), i.e., a piece of chain placed in front of the bottom gear

of the trawler that is considered effective as regards catching skates and rays that may

escape under the net. The inclusion of BLW in a Uruguayan longline fishery led to a

reduction in the by-catch of the scalloped hammerhead (*Sphyrna lewini*) (n=2) of 100%,

while the figure for the pelagic stingray (*Pteroplatytrygon violacea*) (n=18) was 27.8%

(Figure 3D). However, in the case of the "tickler", the number of species captured

increased for all species with the exception of the lesser-spotted dogfish (*Scylorhinus*

*canicular*) (n=1525), which attained a decrease of 2.3% (Kynoch et al. 2015) (Figure 3D).

### 3.3.4.4.  Visual

The use of LED lights as a mitigation measure has been tested for all four groups (sea

turtles, marine mammals, seabirds and elasmobranchs), with an uncertain effect on

elasmobranchs (Mangel et al. 2018). However, in a Mexican gillnet fishery, there was a

reduction in the elasmobranch by-catch of 95% (Senko et al. 2022). Tori lines have also

obtained good results for this group, reducing the by-catch rate for the porbeagle

(*Lamna nasus*) (n=34), copper shark (*Carcharhinus brachyurus*) (n = 8), night shark (n=38)

and smooth hammerhead (n=190) by 41.2%, 87.5%, 89.5% and 86.3%, respectively

(Jiménez et al. 2019) (Figure 1).

### 3.3.4.5. Echolocation

Rays do not use echolocation, and the by-catch data obtained after the implementation of the acrylic glass spheres by Kratzer et al. (2021) in a Turkish commercial fishery confirm this. More thornback skate (*Raja clavata*) individuals were caught in the modified gillnet (n=97) than in the standard one (n=41).

### 3.3.4.6. Electrosensory

Sharks have a complex and extensive electrosensory system, which includes the ampullae of Lorenzini located around the snout or rostral area (Kajiura and Holland, 2002). The use of SMART hooks in a longline fishery in the Gulf of Maine (USA) led to a reduction in the number of shark species caught, from 25% to 100% (O'Connell et al. 2014) (Figure 3D).

The last sensory type measure found was the use of hooks made from a neodymium-praseodymium alloy, use by longlines in USA and Ecuador with the scalloped hammerhead (n=52), leading to a reduction of 61.5% (Hutchinson et al. 2012) (Figure 3D).

### 3.4. Limitations of this review

The objective of this literature review was to provide a comprehensive overview of the most effective measures used to date in order to reduce the by-catch of sea turtles, marine mammals, seabirds and elasmobranchs. However, it was difficult to carry out the global standardization of data because many studies were incomplete owing to a lack of sample sizes (Königson et al. 2022; Kuepfer et al. 2022; Godin et al. 2013), the existence of small sizes (O'Connell et al. 2014; Domingo et al. 2017; Jiménez et al. 2019) or the absence of the name of the species being studied (*Diomedea spp., Procellaria spp.,*

*Delphinus spp., Globicephala spp.*) (Yokota et al. 2011; Mangel et al. 2013). Moreover, in

many cases the effectiveness of different mitigation measures for the species involved

was not included (Kynoch et al. 2015; Porsmoguer et al. 2015; Basran et al. 2020), making

it difficult to make comparisons among studies.

**3.5.     Recommendations for future research**

A limited number of complete studies on by-catch mitigation measures were found.

These included TEDs for sea turtles (Warden, 2011; Casale et al. 2017), pingers for

marine mammals (Amano et al. 2017; Moan and Bjørge, 2021) and tori lines for seabirds

(Domingo et al. 2017; Paterson et al. 2019). However, the available research on this topic

is still lacking in many aspects.

There is a need for more studies that quantitatively assess the actual amount of by-catch

(Basran et al. 2020; Culik et al. 2015; Westlake et al. 2018). Moreover, most of the results

obtained often vary according to geographical areas, species and fishing practices, thus

highlighting the importance of conducting further research into effective strategies by

which to mitigate by-catch, particularly in regions in which SFFs are prevalent, such as

Asia, Africa and South America.

It is also essential to establish standardized reporting practices, define study parameters,

specify research locations and context, and examine unintended impacts on animal

populations so as to attain accurate comparisons (Kynoch et al. 2015; Porsmoguer et al.

2015; Grant et al. 2018). Consistency in measurement metrics is crucial, focusing on the

number of individuals captured per unit of effort (Senko et al. 2022; Berninsone et al.

2020; Gautama et al. 2022). Thorough documentation should encompass specifics such

as gear type, study locale, the technology employed, and technical specifications (Senko

et al. 2022; Mangel et al. 2013). It is, therefore, also recommended that studies explicitly

detail sample and effect sizes (O'Connell et al. 2014; Domingo et al. 2017).

Furthermore, we believe that it is necessary to increase exploration into the combination

of sensory deterrents in order to reduce by-catch across various taxonomic groups

(Coelho et al. 2015; Gilman et al. 2021). Future research should prioritize the use of cost-

efficient technologies that are straightforward to implement, as these are more likely to

gain the support and compliance of the fishing industry. This will make it possible to work

toward preserving the future of many endangered species and reducing the impact of

by-catch.

**4.   REFERENCES**

Agardy, T. (2000). Effects of fisheries on marine ecosystems: A conservationist's

perspective.      ICES      Journal      of      Marine      Science,      57(3),      761–765.

https://doi.org/10.1006/jmsc.2000.0721

Amano, M., Kusumoto, M., Abe, M., Akamatsu, T. (2017). Long-term effectiveness of

pingers on a small population of finless porpoises in Japan. Endangered Species

Research, 32(1), 35–40. https://doi.org/10.3354/esr00776

Andri, S., Ken, A., Andreas, A., Nanina, A., Tomas, A., Chandima, A., ..., Ben, B. (2021).

DescTools: Tools for descriptive statistics. R package version, 0.99, 43.

Basran, C. J., Woelfing, B., Neumann, C., Rasmussen, M. H. (2020). Behavioural

Responses of Humpback Whales (*Megaptera novaeangliae*) to Two Acoustic Deterrent

Devices in a Northern Feeding Ground off Iceland. Aquatic Mammals, 46(6), 584–602.

https://doi.org/10.1578/AM.46.6.2020.584

Berninsone, L. G., Bordino, P., Gnecco, M., Foutel, M., Mackay, A. I., Werner, T. B. (2020).

Switching Gillnets to Longlines: An Alternative to Mitigate the Bycatch of Franciscana

Dolphins (*Pontoporia blainvillei*) in Argentina. Frontiers in Marine Science, 7(August), 1–

19. https://doi.org/10.3389/fmars.2020.00699

Bielli, A., Alfaro-Shigueto, J., Doherty, P. D., Godley, B. J., Ortiz, C., Pasara, A., Wang, J. H.,

Mangel, J. C. (2020). An illuminating idea to reduce bycatch in the Peruvian small-scale

gillnet fishery. Biological Conservation, 241(August 2019), 108277.

https://doi.org/10.1016/j.biocon.2019.108277

Bilgin, S., Kose, O. (2018). Testing two types of acoustic deterrent devices (pingers) to

reduce harbour porpoise, *Phocoena phocoena* (Cetacea: Phocoenidae), by catch in

turbot (Psetta maxima) set gillnet fishery in the Black Sea, Turkey. Cahiers de Biologie

Marine, 59(5), 473–479. https://doi.org/10.21411/CBM.A.D5B58D5B

BirdLife International (2018a). *Puffinus mauretanicus*. The IUCN Red List of Threatened

Species. https://doi.org/10.2305/IUCN.UK.2018-2.RLTS.T22728432A132658315.en

BirdLife International (2018b). *Puffinus yelkouan*. The IUCN Red List of Threatened

Species. https://doi.org/10.2305/IUCN.UK.2018-2.RLTS.T22698230A132637221.en.

Bordino, P., Albareda, D. (2004). Incidental mortality of franciscana dolphin *Pontoporia*

*blainvillei* in coastal gillnet fisheries in Northern Buenos Aires, Argentina. Paper

SC/56/SM11. INT. WHAL. COMMN. MEETING, Sorrento, Italy.

Bordino, P., Mackay, A., Werner, T., Northridge, S., Read, A. (2013). Franciscana bycatch

is not reduced by acoustically reflective or physically stiffened gillnets. Endangered

Species Research, 21(1), 1–12. https://doi.org/10.3354/esr00503

Braulik, G., Jefferson, T. A., Bearzi, G. (2021). *Delphinus delphis*. The IUCN Red List of

Threatened                    Species.                    https://doi.org/10.2305/IUCN.UK.2021-

2.RLTS.T134817215A199893039.e

Carretta, J. V., Barlow, J. (2011). Long-term effectiveness, failure rates, and "dinner bell"

properties of acoustic pingers in a gillnet fishery. Marine Technology Society Journal,

45(5), 7–19. https://doi.org/10.4031/MTSJ.45.5.3

Casale, P., Abitsi, G., Aboro, M. P., Agamboue, P. D., Agbode, L., Allela, N. L., Angueko, D.,

Bibang Bi Nguema, J. N., Boussamba, F., Cardiec, F., Chartrain, E., Ciofi, C., Emane, Y. A.,

Fay, J. M., Godley, B. J., Kouerey-Oliwiwina, C. K., de Dieu Lewembe, J., Leyoko, D.,

MbaAsseko, G., … Formia, A. (2017). A first estimate of sea turtle bycatch in the industrial

trawling    fishery    of    Gabon.    Biodiversity    and    Conservation,    26(10),    2421–2433.

https://doi.org/10.1007/s10531-017-1367-z

Coelho, R., Santos, M. N., Fernandez-Carvalho, J., Amorim, S. (2012). Effects of hook and

bait on sea turtle catches in an equatorial Atlantic pelagic longline fishery. Bull Marine

Sci, 88(3) (https://doi.org/10.5343/bms.2011.1065), 449–467.

Coelho, R., Santos, M. N., Fernandez-Carvalho, J., Amorim, S. (2015). Effects of hook and

bait in a tropical northeast Atlantic pelagic longline fishery: Part I-Incidental sea turtle

bycatch.             Fisheries             Research,             164(3),             302–311.

https://doi.org/10.1016/j.fishres.2014.11.008

Collins, M. A., Hollyman, P. R., Clark, J., Soeffker, M., Yates, O., Phillips, R. A. (2021).

Mitigating the impact of longline fisheries on seabirds: Lessons learned from the South

Georgia Patagonian toothfish fishery (CCAMLR Subarea 48.3). Marine Policy, 131(June),

104618. https://doi.org/10.1016/j.marpol.2021.104618

Cooke, J. G. (2018). *Megaptera novaeangliae*. The IUCN Red List of Threatened Species.

https://doi.org/10.2305/IUCN.UK.2018-2.RLTS.T13006A50362794.en.

Cortés, V., González-Solís, J. (2018). Seabird bycatch mitigation trials in artisanal

demersal longliners of the Western Mediterranean. PLoS ONE, 13(5), 1–21.

https://doi.org/10.1371/journal.pone.0196731

Crowder, L. B., Hazen, E. L., Avissar, N., Bjorkland, R., Latanich, C., Ogburn, M. B. (2008).

The impacts of fisheries on marine ecosystems and the transition to ecosystem-based

management. Annual Review of Ecology, Evolution, and Systematics, 39, 259–278.

https://doi.org/10.1146/annurev.ecolsys.39.110707.173406

Culik, B., VonDorrien, C., Müller, V., Conrad, M. (2015). Synthetic communication signals

influence wild harbour porpoise (*Phocoena phocoena*) behaviour. Bioacoustics, 24(3),

201–221. https://doi.org/10.1080/09524622.2015.1023848

Darquea, J. J., Ortiz-Alvarez, C., Córdova-Zavaleta, F., Medina, R., Bielli, A., Alfaro-

Shigueto, J., Mangel, J. C. (2020). Trialing net illumination as a bycatch mitigation

measure for sea turtles in a small-scale gillnet fishery in Ecuador. Latin American Journal

of Aquatic Research, 48(3), 446–455. https://doi.org/10.3856/vol48-issue3-fulltext-

2428

Dawson, S. M. (1991). Modifying gillnets to reduce entanglement of cetaceans. Marine

Mammal Science, 7(3), 274–282. https://doi.org/https://doi.org/10.1111/j.1748-

7692.1991.tb00102.x

Domingo, A., Jiménez, S., Abreu, M., Forselledo, R., Yates, O. (2017). Effectiveness of tori

line use to reduce seabird bycatch in pelagic longline fishing. PLoS ONE, 12(9), 1–15.

https://doi.org/10.1371/journal.pone.0184465

FAO (2022). El estado mundial de la pesca y la acuicultura 2022. Hacia la transformación

azul. Roma, FAO, 1–257. https://doi.org/10.4060/cc0461es

Field, R., Crawford, R., Enever, R., Linkowski, T., Martin, G., Morkūnas, J., Morkūnė, R.,

Rouxel, Y., Oppel, S. (2019). High contrast panels and lights do not reduce bird bycatch in

Baltic Sea gillnet fisheries. Global Ecology and Conservation, 18.

https://doi.org/10.1016/j.gecco.2019.e00602

Gautama, D. A., Susanto, H., Riyanto, M., Wahju, R. I., Osmond, M., Wang, J. H. (2022).

Reducing sea turtle bycatch with net illumination in an Indonesian small-scale coastal

gillnet fishery. November, 1–12. https://doi.org/10.3389/fmars.2022.1036158

Gilman, E., Chaloupka, M., Ishizaki, A., Carnes, M., Naholowaa, H., Brady, C., Ellgen, S.,

Kingma, E. (2021). Tori lines mitigate seabird bycatch in a pelagic longline fishery.

Reviews in Fish Biology and Fisheries, 31(3), 653–666. https://doi.org/10.1007/s11160-

021-09659-7

Godin, A. C., Wimmer, T., Wang, J. H., Worm, B. (2013). No effect from rare-earth metal

deterrent on shark bycatch in a commercial pelagic longline trial. Fisheries Research,

143, 131–135. https://doi.org/10.1016/j.fishres.2013.01.020

Gonzalez, A., Vega, R., Barbieri, M. A., Yanez, E. (2012). Determinación de los factores

que inciden en la captura incidental de aves marinas en la flota palangrera pelágica

chilena. Latin American Journal of Aquatic Research, 40(3), 786–799.

https://doi.org/10.3856/vol40-issue3-fulltext-25

Grant, S. M., Sullivan, R., Hedges, K. J. (2018). Greenland shark (*Somniosus*

*microcephalus*) feeding behavior on static fishing gear, effect of SMART (Selective

Magnetic and Repellent-Treated) hook deterrent technology, and factors influencing

entanglement in bottom longlines. PeerJ, 2018(5). https://doi.org/10.7717/peerj.4751

Gupta, T., Booth, H., Arlidge, W., Rao, C., Manoharakrishnan, M., Namboothri, N.,

Shanker, K., Milner-Gulland, E. J. (2020). Mitigation of Elasmobranch Bycatch in Trawlers:

A Case Study in Indian Fisheries. Frontiers in Marine Science, 7(July), 1–17.

https://doi.org/10.3389/fmars.2020.00571

Hervé, M (2023). RVAideMemoire: testing and plotting procedures for bioestatistics. R

package version 0.9-83–3.

Hutchinson, M., Wang, J. H., Swimmer, Y., Holland, K., Kohin, S., Dewar, H., Wraith, J.,

Vetter, R., Heberer, C., Martinez, J. (2012). The effects of a lanthanide metal alloy on

shark catch rates. Fisheries Research, 131–133(December 2011), 45–51.

https://doi.org/10.1016/j.fishres.2012.07.006

Jiménez, S., Domingo, A., Forselledo, R., Sullivan, B. J., Yates, O. (2019). Mitigating

bycatch of threatened seabirds: the effectiveness of branch line weighting in pelagic

longline fisheries. Animal Conservation, 22(4), 376–385.

https://doi.org/10.1111/acv.12472

Jiménez, S., Forselledo, R., Domingo, A. (2019). Effects of best practices to reduce seabird

bycatch in pelagic longline fisheries on other threatened, protected and bycaught

megafauna species. Biodiversity and Conservation, 28(13), 3657–3667. https://doi.org/10.1007/s10531-019-01842-4

Kajiura, S. M., Holland, K. N. (2002). Electroreception in juvenile scalloped hammerhead and sandbar sharks. Journal of Experimental Biology, 205(23), 3609–3621. https://doi.org/10.1242/jeb.205.23.3609

Kakai, T (2019). Assessing the effectiveness of LED lights for the reduction of sea turtle bycatch in an artisanal gillnet fishery - a case study from the north coast of Kenya. Marine Science, 18(2), 37–44. https://doi.org/10.1038/278097a0

Königson, S., Naddafi, R., Hedgärde, M., Pettersson, A., Östman, Ö., Benavente Norrman, E., Amundin, M. (2022). Will harbour porpoises (*Phocoena phocoena*) be deterred by a pinger that cannot be used as a "dinner bell" by seals? Marine Mammal Science, 38(2), 469–485. https://doi.org/10.1111/mms.12880

Kratzer, I. M. F., Brooks, M. E., Bilgin, S., Özdemir, S., Kindt-Larsen, L., Larsen, F., Stepputtis, D. (2021). Using acoustically visible gillnets to reduce bycatch of a small cetacean: first pilot trials in a commercial fishery. Fisheries Research, 243(December 2020). https://doi.org/10.1016/j.fishres.2021.106088

Kuepfer, A., Sherley, R. B., Brickle, P., Arkhipkin, A., Votier, S. C. (2022). Strategic discarding reduces seabird numbers and contact rates with trawl fishery gears in the Southwest Atlantic. Biological Conservation, 266(January), 109462. https://doi.org/10.1016/j.biocon.2022.109462

Kulka, D. W., Cotton, C. F., Anderson, B., Derrick, D., Herman, K., Dulvy, N. K. (2020).

*Somniosus microcephalus*. The IUCN Red List of Threatened Species.

https://doi.org/10.2305/IUCN.UK.2020-3.RLTS.T60213A124452872.en.

Kynoch, R. J., Fryer, R. J., Neat, F. C. (2015). A simple technical measure to reduce bycatch

and discard of skates and sharks in mixed-species bottom-trawl fisheries. ICES Journal of

Marine Science, 72(6), 1861–1868. https://doi.org/10.1038/278097a0

Li, Y., Browder, J. A., Jiao, Y. (2012). Hook effects on seabird bycatch in the United States

Atlantic pelagic longline fishery. Bulletin of Marine Science, 88(3), 559–569.

https://doi.org/10.5343/bms.2011.1039

Lucas, S., Berggren, P. (2023). A systematic review of sensory deterrents for bycatch

mitigation of marine megafauna. Reviews in Fish Biology and Fisheries, 33(1), 1-33.

https://doi.org/10.1007/s11160-022-09736-5

Lucchetti, A., Bargione, G., Petetta, A., Vasapollo, C., Virgili, M. (2019). Reducing sea

turtle bycatch in the mediterranean mixed demersal fisheries. Frontiers in Marine

Science, 6(JUL), 1–12. https://doi.org/10.3389/fmars.2019.00387

Mangel, J. C., Alfaro-Shigueto, J., Witt, M. J., Hodgson, D. J., Godley, B. J. (2013). Using

pingers to reduce bycatch of small cetaceans in Peru's small-scale driftnet fishery. Oryx,

47(4), 595–606. https://doi.org/10.1017/S0030605312000658

Mangel, J. C., Wang, J., Alfaro-Shigueto, J., Pingo, S., Jimenez, A., Carvalho, F., Swimmer,

Y., Godley, B. J. (2018). Illuminating gillnets to save seabirds and the potential for multi-

taxa bycatch mitigation. Royal Society Open Science, 5(7), 4–7.

https://doi.org/10.1098/rsos.180254

Mangiafico, S. S (2015). An R Companion for the Handbook of Biological Statistics,

version 1.3.9, revised 2023. rcompanion.org/rcompanion/

Moan, A., Bjørge, A. (2021). Pinger trials in Norwegian commercial fisheries confirm that

pingers reduce harbour porpoise bycatch rates and demonstrate low level of pinger-

associated negative impacts on day-to-day fishing operations. IWC Scientific Committee,

68, 1–18.

Negri, M. F., Denuncio, P., Panebianco, M. V., Cappozzo, H. L. (2012). Bycatch of

franciscana dolphins *Pontoporia blainvillei* and the dynamic of artisanal fisheries in the

species' southernmost area of distribution. Brazilian Journal of Oceanography, 60(2),

149–158. https://doi.org/10.1590/S1679-87592012000200005

O'Connell, C. P., He, P., Joyce, J., Stroud, E. M., Rice, P. H. (2014). Effects of the SMART

TM (Selective Magnetic and Repellent-Treated) hook on spiny dogfish catch in a longline

experiment in the Gulf of Maine. Ocean and Coastal Management, 97, 38–43.

https://doi.org/10.1016/j.ocecoaman.2012.08.002

Oliveira, N., Almeida, A. N. A., Alonso, H., Constantino, E., Ferreira, A., Gutiérrez, I.,

Santos, A. N. A., Silva, E., Andrade, J. (2021). A contribution to reducing bycatch in a high

priority area for seabird conservation in Portugal. Bird Conservation International, 31(4),

553–572. https://doi.org/10.1017/S0959270920000489

Ortiz, N., Mangel, J. C., Wang, J., Alfaro-Shigueto, J., Pingo, S., Jimenez, A., Suarez, T.,

Swimmer, Y., Carvalho, F., Godley, B. J. (2016). Reducing green turtle bycatch in small-

scale fisheries using illuminated gillnets: The cost of saving a sea turtle. Marine Ecology

Progress Series, 545, 251–259. https://doi.org/10.3354/meps11610

Paterson, J. R. B., Yates, O., Holtzhausen, H., Reid, T., Shimooshili, K., Yates, S., Sullivan,

B. J., Wanless, R. M. (2019). Seabird mortality in the Namibian demersal longline fishery

and recommendations for best practice mitigation measures. Oryx, 53(2), 300–309.

https://doi.org/10.1017/S0030605317000230

Porsmoguer, S. B., Bănaru, D., Boudouresque, C. F., Dekeyser, I., Almarcha, C. (2015).

Hooks equipped with magnets can increase catches of blue shark (*Prionace glauca*) by

longline         fishery.         Fisheries         Research,         172,         345–351.

https://doi.org/10.1016/j.fishres.2015.07.016

Raghu, R., Boopendranath, M. R., Vinod, M. (2016). Performance Evaluation of Turtle

Excluder Device off Dhamra in Bay of Bengal Energy analysis of fishing systems View

project Responsible fishing View project. Fishery Technology, 53(10), 183–189.

https://www.researchgate.net/publication/307594951

Richards, R. J., Raoult, V., Powter, D. M., Gaston, T. F. (2018). Permanent magnets reduce

bycatch of benthic sharks in an ocean trap fishery. Fisheries Research, 208(March), 16–

21. https://doi.org/10.1016/j.fishres.2018.07.006

Rollinson, D. P., Wanless, R. M., Ryan, P. G. (2017). Patterns and trends in seabird bycatch

in the pelagic longline fishery off South Africa. African Journal of Marine Science, 39(1),

9–25. https://doi.org/10.2989/1814232X.2017.1303396

Rouxel, Y., Crawford, R., Cleasby, I. R., Kibel, P., Owen, E., Volke, V., Schnell, A. K., Oppel,

S. (2021). Buoys with looming eyes deter seaducks and could potentially reduce seabird

bycatch       in       gillnets.       Royal       Society       Open       Science,       8(5).

https://doi.org/10.1098/rsos.210225

Seidu, I., Brobbey, L. K., Danquah, E., Oppong, S. K., van Beuningen, D., Seidu, M., Dulvy,

N. K. (2022). Fishing for survival: Importance of shark fisheries for the livelihoods of

coastal communities in Western Ghana. Fisheries Research, 246(January 2021), 106157.

https://doi.org/10.1016/j.fishres.2021.106157

Senko, J. F., Peckham, S. H., Aguilar-Ramirez, D., Wang, J. H. (2022). Net illumination

reduces fisheries bycatch, maintains catch value, and increases operational efficiency.

Current Biology, 32(4), 911-918.e2. https://doi.org/10.1016/j.cub.2021.12.050

UICN (2023). IUCN Red List of Threatened Species. https://www.iucnredlist.org

Virgili, M., Vasapollo, C., Lucchetti, A. (2018). Can ultraviolet illumination reduce sea

turtle bycatch in Mediterranean set net fisheries? Fisheries Research, 199 (May 2017),

1–7. https://doi.org/10.1016/j.fishres.2017.11.012

Wang, J., Barkan, J., Fisler, S., Godinez-Reyes, C., Swimmer, Y. (2013). Developing

ultraviolet illumination of gillnets as a method to reduce sea turtle bycatch. Biology

Letters, 9(5), 3–6. https://doi.org/10.1098/rsbl.2013.0383

Wang, J. H., Fisler, S., Swimmer, Y. (2010). Developing Visual deterrents to reduce sea

turtle bycatch in gill net fisheries. Marine Ecology Progress Series, 408, 241–250.

https://doi.org/10.3354/meps08577

Warden, M. L. (2011). Modeling loggerhead sea turtle (*Caretta caretta*) interactions with

US Mid-Atlantic bottom trawl gear for fish and scallops, 2005-2008. Biological

Conservation, 144(9), 2202–2212. https://doi.org/10.1016/j.biocon.2011.05.012

Wells, R. S., Natoli, A., Braulik, G. (2019). *Tursiops truncatus*. The IUCN Red List of

Threatened Species. https://doi.org/10.2305/IUCN.UK.2019-

1.RLTS.T22563A156932432.en.

Westlake, E. L., Williams, M., Rawlinson, N. (2018). Behavioural responses of

draughtboard sharks (*Cephaloscyllium laticeps*) to rare earth magnets: Implications for

shark bycatch management within the Tasmanian southern rock lobster fishery. Fisheries

Research, 200(January), 84–92. https://doi.org/10.1016/j.fishres.2018.01.001

Wibbels, T., Bevan, E. (2019). *Lepidochelys kempii*. The IUCN Red List of Threatened

Species. https://doi.org/10.2305/IUCN.UK.2019-2.RLTS.T11533A155057916.en

Yokota, K., Minami, H., Kiyota, M. (2011). Effectiveness of tori-lines for further reduction

of incidental catch of seabirds in pelagic longline fisheries. Fisheries Science, 77(4), 479–

485. https://doi.org/10.1007/s12562-011-0357-4

Table 1. Mitigation measures proposed by different countries depending on the sensory system involved for each megafauna group, with their references (Methods SI).

| MITIGATION TYPE | | | | Sea turtles | Cetaceans | Other mammals | Seabirds | Elasmobranchs | Reference |
|---|---|---|---|---|---|---|---|---|---|
| | Acoustic | Acoustic Deterrent Device | Pinger | | X | X | | X | (1) |
| | | Acoustic Harassment Device | Seal scarer | | X | | | | (2) |
| | Olfactory | Alternative bait type | Mackerel bait replacing squid | X | | | X | X | (3) |
| | | Offal discard management | --- | | | | X | | (4) |
| | Physical | Structural changes | Changes of fishing nets | | X | | | | (5) |
| | | | Hook change | X | | | | | (6) |
| | | | Branch Line Weighting | X | | X | X | X | (7) |
| | | Escape option | Bycatch Reduction Device | X | | | | | (8) |
| | | | Turtle Excluder Device | X | | | | X | (9) |
| | | | Bird exclusion device | | | | X | | (10) |
| | Visual | Lights | LED lights | X | | | X | X | (11) |
| | | | UV-LED | X | | | | | (12) |
| | | | Chemical lightsticks | X | | | | | (13) |
| | | Physical model | Predator cut-outs | X | X | | X | X | (14) |
| | | | Looming eyes | X | | | X | | (15) |
| | | | High contrast panels | | | | X | | (16) |
| | | | Tori lines | X | | X | X | X | (17) |
| | | Gear colour | Colour net/hook | | | | X | | (18) |
| | | Bait colour | Blue dye | | | | X | | (19) |
| | Echolocation | Acoustic reflection device | Net material alteration | | X | | | | (20) |
| | | | Acrylic spheres | | X | | | X | (21) |
| | Electrosensory | Modified hook | SMART hook | | | | | X | (22) |
| | | Magnet | Neodymium-iron-boron | | | | | X | (23) |
| | | | Ferrite | | | | | X | (24) |
| | | Electropositive Metal Alloy | Neodymium-praseodymium | | | | | X | (25) |

**Figure 1.** World map showing the percentage of reduction in by-catch per group (sea turtles in red, marine mammals in light blue, seabirds in green and elasmobranchs in dark blue) by continent (America divided into North (1) and South (2) America, Africa (3), Europe (4), Asia (5) and Oceania (6)), obtained by the most frequently used mitigation measure: TEDs for sea turtles, pingers for marine mammals and tori lines for seabirds and elasmobranchs.

**Figure 2. A.** Violin plot illustrating the percentage of reduction in by-catch obtained for the species groups (marine turtles, marine mammals, seabirds and elasmobranchs). **B.** Percentage of reduction in by-catch obtained with the different mitigation types (acoustic, visual, echolocation, electrosensory, physical and olfactory). The horizontal lines inside each box correspond to the mean, while the vertical lines at the ends of each box refer to standard deviation (SD).

**Figure 3.** Success in reducing by-catch (%) of **A.** Sea turtles; **B.** Marine mammals; **C.** Seabirds, and **D**. Elasmobranchs, using different by-catch mitigation measures. The horizontal lines inside each box correspond to the mean, while the vertical lines at the ends of each box refer to SD.