RUNNING HEAD: *Bacillus cereus* **group exposure assessment model**

Exposure assessment suggests some cytotoxic *Bacillus cereus* **group genotypes**

can grow over 3 logs in HTST milk throughout the shelf life at temperature

abuse conditions

Jun Su[1]*, Tyler Chandross-Cohen[2]*, Chenhao Qian[1], Laura Carroll[3,4,5,6], Kayla Kimble[2],
Mackenna Yount[2], Martin Wiedmann[1], Jasna Kovac[2#]

[1]Department of Food Science, Cornell University, Ithaca, NY 14850

[2]Department of Food Science, The Pennsylvania State University, University Park, PA 16802

[3]Department of Clinical Microbiology, SciLifeLab, Umeå University, Umeå, Sweden

[4]Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University, Umeå,
Sweden

[5]Umeå Centre for Microbial Research (UCMR), Umeå University, Umeå, Sweden

[6]Integrated Science Lab (IceLab), Umeå University, Umeå, Sweden

*Authors contributed equally.

[#]Corresponding author.

ABSTRACT

Cytotoxic *Bacillus cereus* group strains are common causes of foodborne illness including

diarrhea. However, our ability to assess food safety risks associated with the exposure to

cytotoxic *B. cereus* group strains *via* contaminated food is limited due to the lack of predictive

tools. In this study, we experimentally quantified the growth of 17 cytotoxic *B. cereus* group

strains, representing six phylogenetic groups, in skim milk broth and used the growth data to

develop an exposure assessment model. While none of the tested strains showed detectable

growth in HTST milk at 4 or 6°C, 15 of the 17 strains showed growth at 10°C, 1 of the 17 strains

showed growth at 8°C, and all strains grew at $\geq$14°C. Growth data for 16 strains allowed us to

generate linear secondary growth models, which were then used to develop the exposure

assessment model. We simulated a five-stage supply chain with up to 35 consumer storage days,

as that was the timing when distinguishable variations in percent milk containers over 10^5

CFU/m with different *B. cereus* genotypes were observed. When the initial contamination level

of the HTST milk is set at an average of 100 CFU/mL, the model predicts that, on consumer

home storage day 21 and 35, 2.81±0.66 and 4.13±2.53 % (mean ± standard deviation) of the milk

containers would exceed *B. cereus* group concentrations of 10^5 CFU/mL; these data represent the

average across all strains. Sensitivity analysis showed that variation in the input parameter Q0,

the initial physiological state of cells, has the largest effect on model's prediction for 1 of 4 group

II isolates, 1 of 6 group IV isolates and both group V isolates, suggesting the need to better

characterize the growth parameters of these isolates. What-if scenario analysis showed that

increased mean and variability in storage temperature at the consumer's home both have

substantial influence on final predicted *B. cereus* group concentration in milk containers. This

model introduces an initial tool designed to facilitate risk-based food safety decision making for

products that are contaminated with low *B. cereus* group levels.

Keywords: *Bacillus cereus* group, exposure assessment, microbial risk management, precision

food safety

INTRODUCTION

The *Bacillus cereus* group, also referred to as *B. cereus sensu lato* (*s.l.*), is a complex composed of closely related species that can be categorized into eight phylogenetic groups (Carroll et al., 2020). This group of bacteria raises concerns within the food industry due to their ability to form endospores that withstand heat treatments, such as high-temperature short time (**HTST**) pasteurization of milk (Buss da Silva et al., 2022; Tirloni et al., 2022). Certain strains within this group possess the potential to induce foodborne illness through emetic intoxication, a consequence of the production of heat-stable toxin cereulide within a food matrix. On the other hand, other strains from this group can cause diarrheal intoxication. Upon ingestion by the human host, these bacteria, while growing in the small intestine, produce pore-forming enterotoxins, leading to cell death (Jessberger et al., 2020). Notably, the estimated annual number of foodborne illnesses caused by *B. cereus* group members in the US is reported to be 63,623, with a hospitalization rate of 0.4% (CDC, 2018). Moreover, it is important to acknowledge that foodborne illnesses caused by *B. cereus* group members are often underreported due to their short-lived nature and mild symptoms (Stenfors Arnesen et al., 2008). Most foodborne illnesses caused by the *B. cereus* group have been associated with 10^5 to 10^8 cells/spores per gram of food. However, it has been proposed that any food that contains $>10^3$ *B. cereus* group cells/spores per gram cannot be considered safe for consumption (EFSA, 2005).

Members of *B. cereus* group are common biological hazards along the dairy production chain, and they often persist in dairy processing environment biofilms (Tirloni et al., 2022). These hazards are particularly relevant to perishable HTST milk, which requires refrigeration along the entire supply chain to ensure product quality and safety. Despite this requirement, HTST milk can be subject to temperature abuses due to an inefficient cold chain. For instance, although efficient temperature control of food products inside refrigerated trucks can usually be achieved, increases in temperature, sometimes to >10°C, during ground operations at the beginning and end of transportation have been frequently reported (Ndraha et al., 2018). Significant

temperature abuses have also been observed during consumer transportation from the retail

establishment to the home, and inappropriate home storage of food products, which elevate the

food safety risk (Mercier et al., 2017). It has been reported that many home refrigerators

worldwide are running at higher than recommended temperatures, and this inadequate

refrigeration is frequently cited as a factor in incidents of food poisoning (James, Onarinde, and

James, 2017). In recent (2003 to 2016) foodborne outbreaks caused by *B. cereus* group strains,

most cases were attributed to temperature abuse of food, such as refrigeration temperatures as

high as 14°C or long-term storage of food at room temperature (Choi and Kim, 2020).

As *B. cereus* group strains can enter the supply chain at multiple entry points and survive heat

treatment, occurrence of low levels of this microorganism in pasteurized milk is not unexpected

(Berthold-Pluta et al., 2019; Bartoszewicz, Hansen, and Swiecicka, 2008; Svensson et al., 2004).

While the number of *B. cereus* group infections linked to milk and dairy product consumption is

unknown, *B. cereus* group-related outbreaks and recalls have been reported. In July 2010, a *B.*

cereus linked diarrhea outbreak occurred among workers in a Korean company, with an attack

rate of 20.3%. The contamination source was found to be underground water (Choi et al., 2011).

Additionally, in 2018, Australian Capital Territory Department of Health had investigated a *B.*

cereus related outbreak associated with a multi-course dinner after four people reported

gastrointestinal illnesses to a restaurant in Canberra. The beef retained by the restaurant for

microbiological analysis showed an unsatisfactory level of *B. cereus* (i.e., a count of 19,000

CFU/g) (Thirkell et al., 2019). Therefore, to facilitate better food safety management decisions

about mitigation strategies for the *B. cereus* group, industry needs an exposure assessment tool to

quantitatively estimate the dose of *B. cereus* group strains at the point of human consumption,

given that a food product is contaminated with a specific concentration and *B. cereus* group

genotype (Notermans et al., 1997).

In this project, we modeled the growth of selected cytotoxic *B. cereus* group strains from

different phylogenetic groups in contaminated HTST milk along a simulated supply chain and

used the data to predict the *B. cereus* group concentrations in a product upon human

consumption. We hypothesized that different *B. cereus* group strains have different growth

capabilities in HTST milk. Therefore, the percentage of HTST milk containers that contain *B.*

cereus group concentrations above a given threshold (i.e., 10^3 and 10^5 CFU/mL) on a given

storage day differs by the *B. cereus* group strain that contaminates the product. While a more

realistic HTST milk shelf life is usually 18-21 days, we modeled a five-stage supply chain with

up to 35 consumer storage days, as that was the timing when noticeable variations in percent

milk containers over 10^5 CFU/ml with different *B. cereus* genotypes were observed.

<div align="center">MATERIALS AND METHODS</div>

Selection of Bacillus cereus Group Isolates

A total of 17 *B. cereus* group strains were included in this study spanning 6 of the 8 *B. cereus*

phylogenetic groups (Carroll et al., 2020). Isolates were selected from a collection of over 300 *B.*

cereus group isolates with available genome assemblies to represent (i) diverse virulence gene

clusters (i.e., considering the presence and absence of *ces, nhe, hbl, cytk-1, cytk-2, sph, cap, has,*

bps), and (ii) phylogenetic groups, which were assigned using BTyper3 v3.3.3 (Carroll et al.,

2020). Specifically, BTyper3 was used to calculate average nucleotide identity (ANI) values

relative to the species type strain genomes of all validly published and effective *B. cereus* group

species published at the time (*n* = 28, accessed March 20, 2023). Using 95 ANI (i.e., the

threshold typically used to delineate bacterial species) (Jain et al., 2018), the 17 *B. cereus* group

strains used in this study most closely resembled the species type strain genomes of *B.*

pseudomycoides (group I isolates PS00125 and PS00135), *albus* (group II isolate PS00193),

tropicus (group II isolate PS00457), *mobilis* (group II isolate PS00518), *pacificus* (group III

isolate PS00474), *cereus* (group IV isolates PS00402, PS00407, PS00433 and PS00495),

thuringiensis (group IV isolates PS00413 and PS00649), *toyonensis* (group V isolates PS00570

and PS00638) and *cytotoxicus* (group VII isolates PS00194 and PS00536). One strain (PS00564)

did not share >95 ANI with any *B. cereus* group species type strain genome; however, using

panC group assignment, PS00564 belonged to group II (i.e., the phylogenetic group that contains

B. tropicus), shared >94.0 ANI with the *B. tropicus* type strain genome, and was thus grouped

with isolate PS00457 for this study. These selected isolates were grouped into 11 clusters

containing unique virulence gene profiles, and further sub-clustered by phylogenetic group.

Consequently, we generated 10 sub-clusters that contained cytotoxic isolates (61 cytotoxic

isolates in total). Up to two cytotoxic isolates from each virulence-phylogenetic subcluster were

selected for inclusion in this study by using a random number generator (Table 1).

[Table 1]

Selection of Temperatures and Sampling Points

Three temperatures, which represent three industrially relevant scenarios, were selected for the

growth experiments, which were conducted with 17 cytotoxic *B. cereus* group strains. These

include 4°C, the food refrigeration temperature recommended by the Food and Drug

Administration (FDA, 2023), 10°C, the refrigeration abuse temperature, and 22°C, the room

temperature. For two strains (PS00125 and PS00135), which showed growth at 22°C but not at

10°C, we also collected growth data at 16°C.

In addition, we also conducted growth/no growth experiments at 6 and 8°C for 15 strains that

showed growth at 10°C but not at 4°C to better characterize their growth. Two strains (PS00125

and PS00135) that showed growth at 22°C but not at 10°C were also tested at 14 and 18°C. The

sampling points for growth experiments at each temperature were selected based on preliminary

growth experiments in SMB, which informed the selection of sampling time points to ensure that

at least 3 data points were obtained for each growth phase (i.e., lag, exponential growth, and

stationary phase).

Preparation of Spore Suspensions

All strains were streaked from -80°C cryostocks onto brain heart infusion (BHI) agar (BD, cat.

No. 241830, Franklin Lakes, NJ), followed by incubating at 30°C for 24 h. A single isolated

colony was then inoculated in 5 ml of BHI broth (BD, cat. no. 237500, Franklin Lakes, NJ) in a

15 ml conical tube (VWR, cat. no. 89039-670, Radnor, PA). Inoculated broths were incubated at

30°C for 24 h without shaking. Two separate AK Agar #2 plates (Sporulating Agar) (BD, cat. No.

210912, Franklin Lakes, NJ) were spread-plated with 100 µl of overnight culture and incubated

at 30°C for 5 days. To confirm the growth on the AK #2 agar, a wet mount on a microscopy slide

was prepared and examined using a phase contrast microscope. After the presence of spores was

confirmed, the plates were flooded with 8 ml of chilled sterile deionized water and biomass was

scraped with an L-shaped spreader to suspend the spores. The spores were then transferred into

new 15 ml conical tubes, and the suspensions were centrifuged (Avanti Centrifuge J-26 XPI) at

10,000 g for 15 minutes. The supernatant was removed, and the spores were resuspended in 3.5

ml of sterile deionized water. Centrifugation was repeated two more times for a total of three

washes. After the final wash, the water was removed, and the spores were resuspended in 5 ml of

50% ethanol and incubated at 4°C in a tube rotator (VWR Mini Tube Rotator) running at 15 rpm

for 12 h at room temperature to inactivate remaining vegetative cells. After this incubation, the

suspension was centrifuged at 10,000 g for 15 minutes. The supernatant was removed, and the

spores were resuspended in 3.5 ml of sterile deionized water. Centrifugation was repeated an

additional two times for three total washes. After the last centrifugation, the spores were

resuspended in 3 ml of sterile deionized water and stored at 4°C for up to 6 months.

To determine the spore suspension concentrations, 100 µl of each spore suspension was

transferred to a 1.5 ml microcentrifuge tube (VWR, cat. no. 89000-028, Radnor, PA) and

incubated in a water bath at 80°C for 12 minutes, followed by cooling on ice. Using the 1X

phosphate buffer saline (PBS), serial dilutions of the spore suspensions were prepared and spread

plated, in duplicate, onto BHI agar, followed by a 24 h incubation at 30°C. Colonies on each

plate were manually counted, and the CFU/mL of the heat-treated spore suspension was

calculated.

Growth Experiment in Skim Milk Broth

Growth experiment was initiated by preparing a spore suspension at a concentration of 10^3

CFU/mL in 40 ml of skim milk broth (SMB) (BD, cat. No. 232100, Franklin Lakes, NJ),

followed by heat treatment at 80°C for 12 minutes. The growth experiments were conducted in

two independent biological replicates, each with two technical replicates, for 17 selected *B.*

cereus group strains at 4°C, 10°C and 22°C. At 4°C, samples were collected at time points 0, 96,

144, 192, 240, 288, 384, and 576 h. At 10°C, samples were collected at time points 0, 6, 24, 48,

96, 192, 240, 288, 384, 504 h. At 22°C, data was collected in three experiments with

complementary sampling time points, as well as overlapping time points to ensure

reproducibility. Specifically, in one experiment, samples were collected at time points 0, 1, 2, 4,

6, 8, and 24 h. In the second experiment, samples were collected at time points 0, 14, 15, 16, 18,

20, and 22 h. In the third experiment, samples were collected at time points 0, 24, 27, and 30 h,

with a 32 h time point being added for isolates that had not reached the stationary phase at 30 h.

Growth experiments conducted at 16°C for two strains (PS00125 and PS00135) were also

conducted in two biological replicates, each with two technical replicates, with data collected at

time points 0, 1, 2, 4, 6, 8, 14, 18, 24, 48, and 96 h.

At each sampling time point, 1 ml of the sample was serially 10-fold diluted using a 1X PBS

buffer and 100 μl of dilutions 10^{-1}, 10^{-3}, & 10^{-5} were plated on BHI agar plates, in duplicates. The

BHI agar plates were incubated at 30°C for 12 to 24 h before colony enumeration. Isolates with

filamentous colony morphology were pour-plated to prevent colony spreading and were

incubated at 30°C for 12 to 24 h. The colonies on countable plates, with 30 to 300 colonies, were

manually counted. Plating of 100 μl SMB and 100 μl 1X PBS were included in each experiment

as negative controls. Experiments were not considered valid if the negative controls had

observable colony growth.

Growth/No Growth Experiments

For growth/no growth experiments, spore suspensions were inoculated in SMB at a

concentration of 10^3 CFU/mL, and incubated at 6°C or 8°C for 504 h. The data was collected at

time points 0 and 504 h. Experiments were conducted with two independent biological

replicates, each with two technical replicates. For two strains that showed growth at 22°C but not

at 10°C, the growth/no growth experiments were also conducted at 14°C and 18°C. Spore

suspensions of these strains were inoculated in SMB at a concentration of 10^3 CFU/mL, and

incubated for 384 h. The data was collected at time points 0 and 384 h. Experiments were

conducted with two independent biological replicates, each with two technical replicates.

Microbial concentration was determined as described in the previous section. An increase of cell

concentration by 1 log CFU/mL during the growth/no growth experiment was considered

growth.

Primary Model Fitting

To estimate the growth parameters, a Baranyi model (Baranyi and Roberts, 1994) was fitted to

growth data for 17 selected *B. cereus* group strains collected at 22°C, 16°C and 10°C using the

packages "nlsMicrobio" version 0.0-3 (Baty, 2022) and "minpack.lm" version 1.2-3 (Elzhov,

2022) in R version 4.1.2 (R Core Team, 2021). The Baranyi model was selected as the primary

model since it was the one used for microbial growth prediction under dynamic environmental

conditions in R package "biogrowth" version 1.0.2 (Garre, 2022). This primary model was used

for exposure assessment model development to facilitate prediction of growth under dynamic

temperature conditions. The equation (Eq. 1) is presented below:

$$N_t = N_0 + \mu_{max}A(t) - \ln\left(1 + \frac{e^{\mu_{max}A(t)} - 1}{e^{(N_{max} - N_0)}}\right) \qquad (1)$$

with: $A(t) = t + \frac{1}{\mu_{max}}\ln\left[e^{-\mu_{max}t} + e^{-\mu_{max}\lambda} - e^{-\mu_{max}t - \mu_{max}\lambda}\right]$

where μ_{max} is the maximum specific growth rate in ln/h, λ is lag time in h, N_0 is the initial

population in $\log_{10}$ CFU/mL and N_{max} is the maximum population reached in $\log_{10}$ CFU/mL.

The last three data points of isolate PS00413 replicate 1 at 10°C were excluded when fitting

Baranyi model since cells had entered the death phase (Supplemental Material Figure S1).

Secondary Model Fitting

A reduced Ratkowsky model was selected as the secondary model to calculate T_{min} and b

(Ratkowsky et al., 1982), since our available data was confined to growth under sub-optimal

temperatures. The unit of μ_{max} was converted to $\log_{10}$/day before fitting the secondary model in R

version 4.1.2 (R Core Team, 2021). The equation (Eq. 2) is presented below:

$\sqrt{\mu_{max}} = b(T - T_{\min})$ (2)

where T is temperature represented in °C, b is the slope and T_{min} is the theoretical minimum

growth temperature represented in °C.

Growth/no growth experiment data at 6°C, 8°C, and 10°C were also used as input data to

construct the secondary model. Specifically, for strains that showed no growth at 6°C, 8°C, or

10°C, μ_{max} was assumed to be 0 at the given temperature and this μ_{max} value was used as an input

when fitting the secondary growth model.

Statistical Analysis

Kruskal-Wallis tests were performed on the λ, μ_{max} and N_{max} of all strains whose growth data

were obtained from both 22°C and 10°C growth experiments. A Bonferroni-corrected Dunn's test

was used to identify the pairs of strains that showed significant differences in lag time and

maximum growth rate. The non-parametric tests were applied because the growth parameters for

these strains failed to meet normality and equal variance assumptions.

Exposure Assessment Model Overview

The exposure assessment model simulates the growth of selected cytotoxic *B. cereus* group

strains in one lot of HTST milk transported along a five-stage supply chain that includes (i)

processing facility storage, (ii) transportation from the processing facility to retail, (iii) retail

storage, (iv) transportation from retail to the consumer's home, and (v) consumer storage for up

to 35 days. Each hypothetical lot of HTST milk was contaminated by one *B. cereus* group strain

and had 100 half-gallon milk containers represented by 100 iterations. This model predicts *B.*

cereus group concentrations in milk containers on days 14, 21, and 35 of consumer storage.

Model outputs were also used to compute the percentage of milk containers with *B. cereus* group

concentrations that exceeded 10^3 and 10^5 CFU/mL at days 14, 21, and 35.

Exposure Assessment Model Parameters

The input parameters for the exposure assessment model include initial contamination levels,

growth parameters of selected *B. cereus* group strains, and the temperature profiles for each stage

of the supply chain. The initial contamination levels (N_0) for 100 iterations were generated from

a Poisson distribution in R version 4.1.2 (R Core Team, 2021) with a lambda (i.e., mean) value of

100 CFU/mL. The seed for the initial contamination level generation was set to 42, for

consistency.

The initial physiological state of cells (Q0) was calculated as shown in Eq. 3 and Eq. 4 and

described in the Baranyi model (Baranyi and Roberts, 1994):

$h_0 = \mu_{max} \lambda$ (3)

where μ_{max} is the specific maximum growth rate in ln/h, λ is the lag time in h.

h_0 was averaged across 2 replicates at 22°C and 2 replicates at 10°C for each strain to obtain h_{av}

which was used to calculate Q0 specific to each strain:

$Q0 = \dfrac{1}{\exp(h_{av}) - 1}$ (4)

The maximum microbial population (N_{max}) for each given *B. cereus* group strain was estimated

from the Baranyi model and averaged across all replicates from growth experiments at different

temperatures for a given *B. cereus* group strain.

The maximum growth rate at the optimum growth temperature (μ_{opt}) was calculated as shown in

Eq. 5:

$\mu_{opt} = \left(b \cdot \left(T_{opt} - T_{min} \right) \right)^2$ (5)

where T_{min} and b were calculated from the secondary model. T_{opt} is the optimum growth

temperatures by clade previously reported (Nakamura, 1998, Baranyi et al., 2017). The unit of

μ_{opt} is in $\log_{10}$/day.

A unique time and temperature profile for every milk container in each lot was generated by

randomly drawing a time-temperature combination from the corresponding time and temperature

distributions (see Table 2) at each stage of the supply chain as described in a previous study

(Qian et al., 2023). The seed was set to 1, for consistency.

[Table 2]

Sensitivity Analysis

A sensitivity analysis was performed to assess the impact of input parameters, specifically Q0

and N_{max}, on the model's outcome, which is the percent milk containers exceeding *B. cereus*

group concentrations of 5 log on consumer storage day 35. In addition to the average h_0, both the

minimum and maximum h_0 were employed to determine the corresponding maximum and

minimum Q0. The input parameters for running the model on selected strains included the

minimum, average, and maximum values for Q0 (representing the physiological state of the cell,

calculated as the product of growth rate and lag time). Similarly, the minimum, average, and

maximum values for N_{max} were used as input parameters for running the model on the selected

strains.

What-If Scenario

Previous studies (Ndraha et al., 2018) suggest that temperature abuse along the HTST milk

supply chain is most likely to occur during product transportation (from processing facility to

retail or from retail to the consumer's home) and during consumer home storage. Here, we

simulated four possible scenarios that could occur along an HTST milk supply chain including:

(i) a delivery delay by 3 days when HTST milk products were transported from processing

facility to retail, with no temperature abuse (e.g., amid a severe snowstorm in New York, milk

delivery was delayed by 3 days due to impassable roads, safety concerns, and the time required

for road clearing and recovery operations; the products were kept in a well-controlled

distribution center during the delay), (ii) a temperature abuse simulating a consumer's

transportation of HTST milk products from retail to home in a hot summer, which results in an

average of 33.5°C (i.e., 25°C higher than the average temperature of the base model), (iii) a

slight temperature deviation in a consumer's home refrigerator, modelled as a 1°C temperature

increase (i.e., an average home storage temperature of 5.45°C), and (iv) an increase in the

variability of refrigeration temperatures due to unstable power supply (e.g., rolling blackout in

California during summer), modelled as a 1°C increase in the dispersion parameter of the

distribution of consumer home storage temperature. Percent milk containers that had *B. cereus*

group concentrations $>10^5$ CFU/ml on consumer storage day 21 and 35 were calculated for all

four scenarios and compared to the base model prediction (Table 3).

[Table 3]

Model Programming and Data Availability

The model was developed in R version 4.1.2 (R Core Team, 2021) with the modified

"biogrowth" package version 1.0.2 (Garre, 2022). The seed for Monte Carlo simulation was set

to 1, for consistency. The raw data used for model construction and R code for the entire study

are available at https://github.com/FSL-MQIP/Bacillus-cereus-exposure-assessment-model.

317　　　　　　　　　　　　　　　　　　**RESULTS**

Bacillus cereus Group Strains Differ in Growth Characteristics, Including the Observed

Minimum Growth Temperature

The 17 selected *B. cereus* group strains (representing six phylogenetic groups) differed in terms

of their minimum growth temperature in SMB. While no strain showed detectable growth (i.e., >

1 log increase in CFU/ml) at 4°C, 15 strains showed growth at 10°C; 14 of these strains did not

grow at 6 or 8°C, but one strain (PS00638 from group V) grew at 8°C. The two strains that did

not show growth at 10°C showed growth at 14 and 16°C, as determined based on the growth/no

growth experiments at these two temperatures. Hence, the experimentally obtained growth

boundaries for the 17 strains assessed here were (i) 8°C (1 strain), (ii) 10°C (14 strains), and (iii)

14°C (2 strains, which represented the only two group I strains characterized in this study).

Due to different minimum growth temperatures associated with the 17 tested strains, the

following datasets were available to estimate growth parameters (i.e., lag time, maximum growth

rate, maximum microbial population): (i) 22°C for all 17 strains; (ii) 10°C for 15 strains, and (iii)

16°C for 2 strains. For 22°C, parameter estimates obtained after fitting the growth data with the

Baranyi model were (i) lag time ranging from 0 h to 22.04 h, (ii) maximum growth rate ranging

from 0.31 ln/h to 2.02 ln/h; and (iii) maximum microbial population ranging from 5.24 $\log_{10}$

CFU/mL to 7.61 $\log_{10}$ CFU/mL. By comparison, for 10°C, parameter estimates were (i) lag times

from 10.02 h to 194.05 h, (ii) maximum growth rates from 0.03 ln/h to 0.34 ln/h and (iii)

maximum microbial population from 5.01 $\log_{10}$ CFU/mL to 7.37 $\log_{10}$ CFU/mL. At 16°C,

parameter estimates were (i) lag times from 9.12 h to 14.02 h, (ii) maximum growth rates from

0.35 ln/h to 0.95 ln/h and (iii) maximum microbial populations from 6.17 $\log_{10}$ CFU/mL to 6.28

$\log_{10}$ CFU/mL (Supplemental Material Table S1).

B. cereus Phylogenetic Groups Differ in Growth Characteristics, including the Maximum

Growth Rate at 22°C and Hypothetical Minimum Growth Temperature

To further evaluate *B. cereus* group growth characteristics, the growth parameters obtained for

different strains were analyzed based on phylogenetic groups (i.e., group I, II, IV, V, and VII; see

Fig. 1). Group III was excluded from these analyses since it only contained one strain. The

Kruskal-Wallis test for 22°C growth parameters indicated that both lag time and maximum

growth rate differed significantly among phylogenetic groups (p-values of 0.040 and 0.036,

respectively), while maximum microbial population did not differ significantly (p-values > 0.1).

Group I strains showed the highest median maximum growth rate (0.92 ln/h), which was

significantly (p = 0.024) higher than that for group IV strains (0.48 ln/h). With regard to the lag

time, group VII strains had the longest median lag time (20.30 h), whereas group IV strains had

the shortest median lag time (8.43 h).

By comparison, the Kruskal-Wallis test for 10°C growth parameters showed no significant

differences (p-values > 0.1) for any of the three parameters (i.e., lag time, maximum growth rate,

maximum microbial population) among phylogenetic groups. Numerically, group V strains had

the highest median maximum growth rate (0.10 ln/h), whereas group IV strains had the lowest

median maximum growth rate (0.06 ln/h). Group II strains had the longest median lag time of

89.11 h, while group V strains had the shortest median lag time of 53.94 h (Fig. 1).

[Figure 1]

To further assess differences in secondary growth parameters between groups, we fitted

maximum growth rates for all 17 strains with a secondary growth model (i.e., reduced

Ratkowsky model). This model was converged using data from all strains except isolate

PS00135 (group I) that had a non-linear relationship between maximum growth rate and

temperature under sub-optimal growth conditions (Supplemental Material Fig. S2). We therefore

were not able to calculate T_{min} (i.e., hypothetical minimum growth temperature) for this strain.

For the rest of the strains, T_{min} ranged from 4.26°C to 9.22°C. The average T_{min} of strains from

groups II, III, IV, V and VII were similar, ranging from 5.47°C to 6.98°C. The only strain from

group I (PS00125) showed a substantially higher T_{min} (9.22°C) compared to the strains from

other groups (Table 4).

[Table 4]

The Base Model Predicts a Higher Percent of Containers Exceeding B. cereus Group

Concentrations of 10^5 and 10^3 CFU/ml for One Group IV Strain and One Group V Strain

The growth data for the 16 *B. cereus* group strains that showed linear secondary models were

used for the development of an exposure assessment model. The base model underwent a "sanity

check" to assess the growth predictions for 16 *B. cereus* group strains in HTST milk products

stored at 3°C for 90 days, with an initial contamination level of 100 CFU/mL. This temperature

was selected because it was about 1°C below the lowest T_{min} of all strains calculated from the

secondary model. As expected, the model predicted no growth for any of the 16 strains at 3°C,

which aligns with the experimental outcomes that none of the strains had detectable growth at

4°C.

The exposure assessment model was subsequently used to predict growth in HTST milk for each

of the 16 included strains up to day 35 of consumer storage, using temperature profiles that

include different time and temperature distributions for 5 stages of the supply chain (Table 2).

The resulting data were used to compute the percentage of simulated milk containers that had B.

$cereus$ group concentrations $>10^5$ CFU/ml and 10^3 CFU/ml on consumer home storage day 14,

21, 35, respectively. The percentage of milk containers exceeding 10^5 CFU/ml on day 14, 21 and

35 were 1.94±0.85 (mean ± standard deviation), 2.81±0.66 and 4.13±2.53, respectively, while the

percentage of milk containers exceeding 10^3 CFU/ml on the same days were 3.00±0.82,

3.88±2.25 and 5.63±3.67.

Model predictions for individual strains (Fig. 2), showed limited differences among strains with

regard to percentage of containers containing above 10^5 CFU/ml (Fig. 2a) at consumer storage

days 14 and 21. However, at day 35, four strains yielded $>5\%$ of containers that exceeded 10^5

CFU/ml, including two strains that yielded approximately 10% of containers that exceeded 10^5

CFU/ml (isolates PS00413 [group IV] and PS00638 [group V]). The same four strains that

showed a higher percentage of containers above 10^5 CFU/ml also showed a higher percentage of

containers above 10^3 CFU/ml (Fig. 2b).

Our analyses identified five strains for which the percentage of containers above 10^3 CFU/ml

(i.e., 3%) was the same at consumer storage days 14, 21, and 35 (i.e., isolates PS00193 and

PS00518 from group II, isolate PS00407 from group IV, and isolates PS00194 and PS00536

from group VII). These five strains, as well as four additional strains (i.e., isolate PS00457 from

group II, isolate PS00474 from group III, isolates PS00495 and PS00649 from group IV) also

showed the same percentage (i.e., 3%) of containers above 10^5 CFU/ml at consumer storage days

21 and 35. We further evaluated the conditions for strains that had the same model predictions at

different consumer storage time to verify our model is performing reasonably. An analysis of the

100 model simulations for these nine strains (which represented 3 group II strains, 1 group III

strain, 3 group IV strains, and 2 group VII strains) showed that for each of these strains, only 3

out of 100 simulated temperature profiles had a consumer home storage temperature $>10°C$. For

the other 97 simulations, the consumer storage temperature was either (i) below T_{min} of these

nine strains (and thus resulted in no predicted growth) or (ii) above T_{min} but below $10°C$, a

temperature at which our model predicts that these isolates grew so slowly that they would not

exceed 10^5 CFU/ml by consumer storage day 35.

[Figure 2]

Sensitivity Analysis Revealed that Variation in the Input Parameter Q0 Significantly

Influenced the Model Predictions for Certain Strains

In the sensitivity analysis, we evaluated the impact of both N_{max} and Q0 on the percent milk

containers exceeding *B. cereus* group concentrations of 10^5 CFU/ml on consumer storage day 35

for 16 strains. This combination of consumer storage day and concentration cut-off was used

because 10^5 CFU/ml represents the *B. cereus* regulatory threshold in several countries and day 35

showed some of the largest variations in model predictions between strains. The sensitivity

analysis for N_{max} predicted an average of 4.06±2.54 (mean ± standard deviation) and 4.13±2.53%

milk containers exceeding *B. cereus* group concentrations of 10^5 CFU/ml on consumer storage

day 35 when using the minimum and maximum N_{max} as the input parameters, respectively, as

compared to 4.13±2.53% when using the average N_{max}. Only one out of the 16 strains (PS00402;

group IV) showed detectable changes in the percent milk containers over 10^5 CFU/ml. For this

strain, the percent milk containers over 5 logs dropped by 1% point when using the minimum

N_{max} as the input as compared to using the average N_{max} (Supplement al Material Table S2). Our

data thus indicate that model outcomes showed limited sensitivity to changes in N_{max}.

In contrast, the model outcomes were more sensitive to the parameter Q0. Using the minimum

and maximum Q0 as the input parameters in the sensitivity analysis resulted in an average of

3.00±2.42 (mean ± standard deviation) and 6.31±4.14% milk containers exceeding *B. cereus*

group concentrations of 10^5 CFU/ml on consumer storage day 35, respectively, as compared to

4.13±2.53% when using the average Q0. This analysis also revealed that the sensitivity of model

outcomes to variation in Q0 differed substantially by strain. The four strains where model

outcomes were most sensitive to changes in Q0 were isolates PS00402 (group IV), PS00638

(group V), PS00564 (group II), and PS00570 (group V). On the other hand, the model outcomes

were not sensitive to the change in Q0 for three strains from group IV and the same predictions

were generated with varying Q0 values (Fig. 3). This is because (i) one strain (i.e., isolate

PS00433 from group IV) had lower variability in the parameter Q0 compared to other strains,

and (ii) our model predicted that the other two strains (i.e., isolates PS00407 and PS00495 from

group IV) grew very slowly and would not reach >10^5 CFU/ml on consumer storage day 35

under storage temperature of <10°C even in the worst-case situation (i.e., highest experimentally

determined Q0). Overall, the sensitivity analysis indicated that additional growth data collection

for strains with great uncertainty in Q0 is necessary to improve the prediction precision.

[Figure 3]

Small Temperature Increases and Increased Temperature Fluctuations in Consumer

Refrigerators Can Substantially Increase the Percentage of Milk Containers Predicted to

Contain B. cereus Group Concentrations >10^5 CFU/ml

What-if scenario analyses were used to evaluate the impact of four different practical scenarios

on possible consumer exposure to milk with *B. cereus* group concentrations >10^5 CFU/ml at

consumer storage day 21 (representing a more realistic HTST milk shelf life) and 35 (the timing

when we start to observe distinguishable variations in the percentage of milk containers

containing over 10^5 CFU/ml among *B. cereus* group strains). Our results show that a three-day

delay on HTST milk product delivery from processing facility to retail without temperature

abuse (scenario i) did not alter the percentage of milk containers exceeding 10^5 CFU/ml of *B.*

cereus group concentrations on consumer home storage day 21 and 35 (i.e., 2.81 and 4.13%,

identical to the base model prediction). Similarly, a temperature abuse simulation of consumer

transportation of HTST milk products from retail to home in a hot summer (average milk

temperature of 33.5°C with an average time of 1 h; scenario ii) led to 3.06 and 4.44% milk

containers exceeding 10^5 CFU/ml on consumer home storage day 21 and 35, which are slightly

higher than the base model predictions (2.81 and 4.13%). Under scenario ii), the model also

predicted that three strains had higher percentages of milk containers containing *B. cereus* group

concentrations >10^5 CFU/ml compared to the baseline on consumer storage day 21. Specifically,

5 and 3% of the milk containers contaminated with isolates PS00413 and PS00433 from group

IV and 5% of the milk containers contaminated with isolate PS00638 from group V had *B.*

cereus group concentrations >10^5 CFU/ml, which is higher compared to 4, 2, and 3% predicted

by the base model, respectively. In contrast, two strains (PS00413 from group IV and PS00570

from group V) were predicted to exceed 10^5 CFU/ml in a higher percentage (11% and 10%,

respectively) of milk containers on consumer storage day 35, compared to the prediction by the

baseline model (i.e., 10% and 6%, respectively).

The first two scenarios, which represent practical events during transportation, did not change

model predictions for most, if not all, strains. However, the last two scenarios, which represent

consumer behavior or ability to manage refrigeration temperature, resulted in a more profound

impact. A slight temperature deviation at prolonged consumer home storage (scenario iii) had a

much higher predicted impact on the percentage of milk containers containing >10^5 CFU/ml than

the two scenarios detailed above. The model predicted 4.56±3.05 (mean ± standard deviation)

and 8.88±3.84 % milk containers with *B. cereus* group concentrations >10^5 CFU/ml on consumer

storage day 21 and 35, respectively (as compared to the base model prediction of 2.81 and 4.13

%). Under scenario iii, 1 of 6 group IV strains and both group V strains were predicted to exceed

10^5 CFU/ml on consumer storage day 21 in more than 10% milk containers, whereas the base

model predicted that none of the strains would exceed 10^5 CFU/ml in more than 10% milk

containers. In contrast, 2 of 4 group II strains, 1 of 1 group III strain, 2 of 6 group IV strains and

2 of 2 group V strains were predicted to exceed 10^5 CFU/ml in more than 10% milk containers

on consumer storage day 35 (compared to 1 of 6 group IV strains and 1 of 2 group V strains

predicted by the base model). Increasing the variability in consumer storage temperature has an

even more substantial effect on the percentage of milk containers exceeding *B. cereus* group

concentrations of 10^5 CFU/ml than a consistently higher average storage temperature (scenario

iii). The model predicts that 7.56±3.20 and 10.81±3.15 % milk containers had *B. cereus* group

concentrations over 10^5 CFU/ml on consumer storage day 21 and 35. Under this scenario, 1 of 4

group II strains, 1 of 6 group IV strains and both group V strains were predicted to have more

than 10% of milk containers exceeding 10^5 CFU/ml on consumer storage day 21, while 15 of 16

strains were predicted to have more than 10% of milk containers exceeding 10^5 CFU/ml on

consumer storage day 35 (compared to 0 and 2 strains predicted to have more than 10% of milk

containers with over 10^5 CFU/ml on consumer storage day 21 and 35 by the base model) (Table

5). The more profound impact from scenarios iii and iv can be explained by the fact that (i)

consumer storage is the longest time in the entire supply chain, and therefore any temperature

change would have a more extended effect, and (ii) changing deviation and variability increases

the number of abnormal temperature conditions that exceed the growth boundaries of more

strains.

[Table 5]

DISCUSSION

This study collected detailed growth data for a genomically diverse set of *B. cereus* group strains

and used these data to develop an exposure assessment model to facilitate industry risk

assessments for diarrheal *B. cereus* group members in HTST milk. Our experimental data

indicates that *B. cereus* group strains show a range of growth characteristics, including

differences in minimum growth temperature, which impact the exposure through consumption of

contaminated HTST milk. We specifically found that (i) *B. cereus* group strains from different

phylogenetic groups differ in key cardinal values (i.e., minimum growth temperatures) and (ii)

there are differences in growth parameters within a given phylogenetic group. Our data thus

supports that the accuracy of *B. cereus* group exposure assessments can be improved by

accounting for genetic diversity within the *B. cereus* group. Importantly, what-if scenarios run

with the initial exposure assessment reported here already provide important insights into *B.*

cereus group risk assessment and risk management and specifically show that absolute

temperature and temperature variation during consumer storage have a large impact on predicted

B. cereus group exposure. While our model can be useful for industry risk assessments of

diarrheal *B. cereus* group members, advances in the understanding of *B. cereus* group virulence

are needed to develop full risk assessment models that include public health metrics.

Bacillus cereus Group Members Show a Range of Growth Characteristics, including

Differences in Minimum Growth Temperatures and Maximum Growth Rates, which Impact

the Risk Associated with the Contamination of HTST Milk

We found that the 16 tested *B. cereus* group strains showed practically relevant variations in both

experimental and theoretical minimum growth temperatures. The experimentally obtained

growth boundaries for most tested *B. cereuses* group strains (i.e., 14 strains) in our study were at

10°C, and two strains each had a lower or higher minimum growth temperature (8°C and 14°C).

This is consistent with the finding of Lott et al. (2023) that some cytotoxic sporeformers in the *B.*

cereus group may grow at abuse temperatures (e.g., 10°C), but not at lower temperatures. By

comparison, one study reported that experimentally observed minimal growth temperatures of 30

diarrheal enterotoxin producing *B. cereus* group strains ranged from ≤5°C to 11°C (Dufrenne et

al., 1994). Similarly, a growth study of 11 psychrotolerant *B. cereus* strains (Borge et al., 2001)

reported that the lowest temperatures with observed growth (in brain heart infusion medium and

commercial 1.5% UHT milk) were 4°C for one strain, 6°C for two strains and 7°C for seven

strains. Guinebretière et al. (2008) reported phylogenetic group-specific minimum growth

temperatures: (i) 10°C for group I (28 strains tested), (ii) 7°C for group II (33 strains tested), (iii)

15°C for group III (96 strains tested), 10°C for group IV (101 strains tested), 8°C for group V (17

strains tested), and 20°C for group VII (2 strains tested). This is generally consistent with our

findings except that the group III and group VII isolates in our study had lower experimental

minimum growth temperatures (i.e., 10°C) as compared to data reported by Guinebretière et al

(2008).

The experimental minimum growth temperatures for all strains in our study were consistent with

their theoretical minimum growth temperatures (T_{min}) predicted by the secondary model.

Specifically, the one group I strain with a minimum growth temperature of 14°C also had a

substantially higher theoretical minimum growth temperature (T_{min}) of 9.2°C compared to the

other 15 strains that showed detectable growth at 10°C (T_{min} ranging from 4.26°C to 7.06°C).

Notably, Carlin et al (2013) reported T_{min} for strains from group II-V are comparable to our

study; however, two group VII strains in their study were reported to have substantially higher

T_{min} (i.e., 12.6°C and 19.1°C) than the T_{min} in our study (i.e., 6.89°C and 7.06°C). Our growth

experiment also showed that *B. cereus* group members have a wide range of estimated maximum

growth rates at 22°C (0.31 ln/h to 2.02 ln/h) and 10°C (0.03 ln/h to 0.34 ln/h). This is comparable

to the reported range of *B. cereus* group maximum growth rates (12 strains, growth experiment

conducted in brain heart infusion supplemented with yeast extract at $2gL^{-1}$ and glucose at $3gL^{-1}$)

of $0.27\ h^{-1}$ to $1.17\ h^{-1}$ at 22°C and $0.02\ h^{-1}$ to $0.17\ h^{-1}$ at 10°C (Carlin et al., 2013). Overall, the

high variability of growth characteristics of *B. cereus* group members, including minimum

growth temperatures and maximum growth rates, suggests that it is crucial to evaluate a diverse

set of strains when predicting *B. cereus* group growth. This is particularly important, as accurate

predictions of the growth of different *B. cereus* strains are essential to correctly predict the

exposure risk if milk is contaminated with a specific *B. cereus* group strain.

Quantitative B. cereus Group Exposure Assessments Can Be Improved by Accounting for

Genetic Diversity of B. cereus Group Members and Common HTST Milk Microbiota

Very few of the previous quantitative exposure assessments conducted in pasteurized fluid milk

have addressed how the genetic diversity of *B. cereus* group members might affect human

exposure (Notermans et al., 1997, Ačai et al., 2014, Lewin et al., 2019). Consistent with Carlin et

al. (2013), who reported the variation of cardinal values for *B. cereus* group members with

respect to their phylogenetic affiliation, our study further emphasizes the need to account for

genetic variability when conducting *B. cereus* group quantitative exposure assessments. This is

specifically supported by the fact that group I isolate PS00125, which has the highest T_{min}

(9.22°C) in our study, was indeed predicted to have the lowest percentage of milk containers

exceeding 10^5 CFU/ml on consumer storage day 35. Importantly, model input parameters other

than T_{min} (e.g., the initial physiological state of cells, Q0) may also impact exposure risk. This is

supported by our exposure assessment model predictions that the risks of *B. cereus* group

exposure on a given consumer storage day (especially consumer storage day 35) varies among

strains and phylogenetic groups that show similar T_{min}. For instance, isolates PS00413 and

PS00402, both from group IV, had similar T_{min} (i.e., 4.88°C and 4.26°C); however, they showed

substantial variations in another model input parameter, Q0 (i.e., 1.26E-02 and 1.91E-09). As a

result, the model predicted that on consumer storage day 35, as high as 10% of the milk

containers with isolate PS00413 exceeded 10^5 CFU/ml, whereas only 4% of the milk containers

with isolate PS00402 exceeded the same threshold.

Additionally, HTST milk is known to have a rich microbiota that consists of thermoduric bacteria

(e.g., *Microbacterium* spp.) and gram negative bacteria associated with post-pasteurization

contamination (e.g., *Pseudomonas*) (Lott, 2023). This suggests the need to account for the effect

of these competing microorganisms on the growth of *B. cereus* group strains in HTST milk.

Previous studies have been conducted to model the interactions between two microorganisms in

the same system (Martens et al., 1999; Courtin and Rul, 2004). Similarly, future studies might be

conducted to quantify the microbial interactions between *B. cereus* group strains and other

microbial contaminants in the HTST milk, thus improving the accuracy of the exposure

assessment model.

What-if Scenarios Suggest That Absolute Temperature and Temperature Variation during

Consumer Storage Have a Large Impact on the Predicted B. cereus Group Exposure

The exposure assessment reported here was used to run selected what-if scenarios to assess the

importance of selected supply chain disruptions and changes, as well as different consumer

storage temperature scenarios that may mimic practices in different locations, seasons, or

infrastructure conditions. The two what-if scenarios that modelled disruption to distribution

included scenarios that mimicked (i) extended storage in a distribution center (without

temperature abuse) and (ii) temperature abuse during consumer transport from retail to home.

Both what-if scenarios showed limited impacts on risk metrics (2.81 and 3.06% of containers

that exceed 10^5 CFU/ml, as compared to 2.81% for the base scenario on consumer storage day

21). This result is not surprising, as scenario (ii) was modelled as a relatively short time period of

temperature abuse (median of 1 h; range of 14 min to 6 h). Similarly, Huang et al (2019) found

that an acute temperature abuse (i.e., 35°C for 2 h) did not substantially increase the growth of

foodborne pathogens (i.e., *Salmonella enterica* and *Listeria monocytogenes*) on cut fruit and

radishes compared to those stored constantly at 4°C. Our results also support the

recommendations for consumer time-temperature control, which is often called the "2-h rule and

1-h rule". This rule suggests that foods should be kept outside the danger zone (4.4–60.0 °C) for

less than 2 h (for temperature between 4.4 and 32.2 °C) or 1 h (for temperatures >32.2 °C) (Cho

et al., 2020). Importantly, these what-if scenarios also illustrate how this model could be used to

assess predicted *B. cereus* group exposures if supply chains might experience certain "worst

case" disruptions (e.g., lack of refrigeration for extended time periods).

The two what-if scenarios that modelled the impact of different temperature conditions during

home storage included scenarios that mimicked (i) a slight temperature deviation in a consumer's

home refrigerator, and (ii) an increase in the variability of refrigeration temperatures in a

consumer's home. These scenarios showed that even what may be considered small differences

in refrigeration conditions (e.g., a change of mean temperature by 1°C) can substantially impact

likely *B. cereus* group exposure. This finding aligns with our existing knowledge that prolonged

temperature abuse during consumer home storage accelerates the growth of many cytotoxic *B.*

cereus group isolates (Ceuppens et al., 2011). It leads to a notable occurrence, with more than

10% of milk containers exceeding 10^5 CFU/ml on consumer home storage day 35. Of particular

interest, even changes in variability of home refrigeration temperatures had substantial impacts.

This type of higher variability may arise from the use of older refrigerators, more commonly

prevalent in households with lower income, and/or in regions or seasons (e.g., hot summers)

where rolling blackouts, i.e., planned short-term power outages, are common. Our findings are

consistent with previous studies (Jofré et al., 2019) that refrigeration temperature fluctuation

increases the risk of exposure to foodborne pathogens, thus negatively affecting the safety of

food products. Overall, the scenarios detailed here (as well as a similar scenario that can easily

be run with the code available) will help assess *B. cereus* group exposures that are likely to occur

in different distribution areas that may differ in average refrigeration temperatures. Examples of

distribution areas that differ in refrigeration conditions include European countries such as

Germany that have a recommended refrigeration temperature of 7°C (DAAD, 2023), as

compared to 4°C, the standard US refrigeration temperature recommended by FDA.

While Our Model Will Facilitate Industry Risk Assessments for Diarrheal B. cereus Group

Strains, Advances in the Understanding of B. cereus Group Virulence Are Needed to Develop

Full Risk Assessments that Include Public Heath Metrics

The model reported here will provide industry with an initial tool that can facilitate risk-based

food safety decision-making for products that are contaminated with low *B. cereus* group levels.

For example, if an HTST fluid milk product is found to contain low levels of a specific *B. cereus*

group strain, our model can be used to predict, for different time points, which proportion of

products are expected to exceed a certain concentration threshold (e.g., 3 or 5 $\log_{10}$ CFU/ml).

Risk managers could then use this information to inform their decision making. For example, a

risk manager may decide that a product with a 21-day shelf life and a predicted 2% of containers

exceeding 10^5 CFU/ml at day 21 is considered low risk, particularly in a country with no

regulatory threshold and a "low risk" supply chain (e.g., a well-developed cold chain with

products typically consumed substantially before the end of the shelf life); additional

considerations will likely impact these decisions (e.g., risk tolerance, lot size). In addition, this

model could also be adopted to assess regulatory risk, such as the likelihood that a lot with low

B. cereus group levels may yield a test result that exceeds a regulatory threshold, using

approaches like those described previously (Chen et al., 2022). We, however, do appreciate that

many users may be hesitant to use this initial model for risk management related decisions, for

instance, because we currently lack the ability to predict the number of human cases that may be

caused by a given lot. For example, a prediction that the risk of causing a single human case is

substantially below 1% may be more valuable than the type of exposure risk that is provided as

an output by our model. This type of approach has previously been used in a risk assessment that

was designed to help with decision making on lots of produce that may be contaminated with

Listeria monocytogenes (Zoellner et al., 2019). However, using the exposure assessment reported

here, prediction of human illness cases that may be caused by a lot is currently impossible due to

the lack of a universally applicable dose-response relationship that will allow us to predict the

likelihood of *B. cereus* toxico-infection per serving and estimate the number of cases within the

population (Notermans et al., 1997, Ačai et al., 2014, Lewin et al., 2019). Nevertheless, our

model could be easily adapted into a full risk assessment if *B. cereus* group dose-response data

become available.

ACKNOWLEDGMENTS

657 This work was supported by the USDA NIFA grant 12686965, Hatch Appropriations PEN04853

658 and Accession 7005519, and the Multistate Capacity Grant 4666.

REFERENCES

REFERENCES

Ačai, P., Ľ. Valík, and D. Liptakova. 2014. Quantitative risk assessment of *Bacillus cereus* in

pasteurised milk produced in the Slovak Republic. Czech J. Food Sci. 32(2):122-131.

https://doi.org/10.17221/106/2013-CJFS

Baranyi, J., N. Buss da Silva, and M. Ellouze. 2017. Rethinking Tertiary Models: Relationships

between Growth Parameters of *Bacillus cereus* Strains. Front. Microbiol. 8:1890.

https://doi.org/10.3389/fmicb.2017.01890

Baranyi, J. and T. A. Roberts. 1994. A dynamic approach to predicting bacterial growth in food.

Int. J. Food Microbiol. 23(3-4):277-294. https://doi.org/10.1016/0168-1605(94)90157-0

Bartoszewicz, M., B. M. Hansen, and I. Swiecicka. 2008. The members of the *Bacillus cereus*

group are commonly present contaminants of fresh and heat-treated milk. Food Microbiol. 25(4),

588-596. https://doi.org/10.1016/j.fm.2008.02.001

Baty, F., M.-L. Delignette-Muller, and A. Siberchicot. 2022. Package 'nlsMicrobio'. Accessed

Dec. 12, 2023. https://cran.r-project.org/web/packages/nlsMicrobio/nlsMicrobio.pdf

Beno, S. M., R. H. Orsi, R. A. Cheng, D. J. Kent, J. Kovac, D. R. Duncan, N. H. Martin, and M.

Wiedmann. 2019. Genes associated with psychrotolerant *Bacillus cereus* group isolates. Front.

Microbiol. 10:662. https://doi.org/10.3389/fmicb.2019.00662

Berthold-Pluta, A., A. Pluta, M. Garbowska, and I. Stefańska. 2019. Prevalence and toxicity

characterization of *Bacillus cereus* in food products from Poland. Foods. 8(7), 269.

https://doi.org/10.3390/foods8070269

Borge, G. I. A., M. Skeie, T. Sørhaug, T. Langsrud, and P. E. Granum. 2001. Growth and toxin

profiles of *Bacillus cereus* isolated from different food sources. Int. J. Food Microbiol.

69(3):237-246. https://doi.org/10.1016/s0168-1605(01)00500-1

Buss da Silva, N., M. Ellouze, K. Rouzeau-Szynalski, N. Johnson, M. H. Zwietering, and H. M.

W. den Besten. 2022. Predicting *B. cereus* growth and cereulide production in dairy mix. Int. J.

Food Microbiol. 364:109519. https://doi.org/10.1016/j.ijfoodmicro.2021.109519

Carlin, F., C. Albagnac, A. Rida, M.-H. Guinebretière, O. Couvert, and C. Nguyen-The. 2013.

Variation of cardinal growth parameters and growth limits according to phylogenetic affiliation

in the *Bacillus cereus* Group. Consequences for risk assessment. Food Microbiol. 33(1):69-76.

https://doi.org/10.1016/j.fm.2012.08.014

Carroll, L. M., R. A. Cheng, and J. Kovac. 2020. No assembly required: using BTyper3 to assess

the congruency of a proposed taxonomic framework for the *Bacillus cereus* group with historical

typing methods. Front. Microbiol. 11:580691. https://doi.org/10.3389/fmicb.2020.580691

CDC. 2018. Estimates of foodborne illness in the United States. Accessed Dec. 12, 2023.

https://www.cdc.gov/foodborneburden/index.html

Ceuppens, S., A. Rajkovic, M. Heyndrickx, V. Tsilia, T. Van De Wiele, N. Boon, & M.

Uyttendaele. 2011. Regulation of toxin production by *Bacillus cereus* and its food safety

implications. Crit Rev Microbiol. 37(3), 188-213.

https://doi.org/10.3109/1040841X.2011.558832

Chen, R., R. H. Orsi, V. Guariglia-Oropeza, and M. Wiedmann. 2022. Development of a

modeling tool to assess and reduce regulatory and recall risks for cold-smoked salmon due to

listeria monocytogenes contamination. J. Food Prot. 85(9), 1335-1354.

https://doi.org/10.4315/JFP-22-025

Cho, T. J., S. A. Kim, H. W. Kim, S. M. Park, and M. S. Rhee. 2020. Changes in consumers'

food purchase and transport behaviors over a decade (2010 to 2019) following health and

convenience food trends. Int. J. Environ. Res. Public Health. 17(15), 5448.

https://doi.org/10.3390/ijerph17155448

Choi, W. and S.-S. Kim. 2020. Outbreaks, germination, and inactivation of *Bacillus cereus* in

food products: A review. J. Food Prot. 83(9):1480-1487. https://doi.org/ 10.4315/0362-

028X.JFP-19-429

Choi, K. B., H. S., Lim, K., Lee, G. Y., Ha, K. H., Jung and C. K., Sohn. 2011. Epidemiological

investigation for outbreak of food poisoning caused by *Bacillus cereus* among the workers at a

local company in 2010. J Prev Med Public Health. 44(2), 65-73.

https://doi.org/10.3961/jpmph.2011.44.2.65

Courtin, P. and F. Rul. 2004. Interactions between microorganisms in a simple ecosystem: yogurt

bacteria as a study model. Lait. 84(1-2), 125-134. https://doi.org/10.1051/lait:2003031

DAAD (German Academic Exchange Service). 2023. Helpful tips for saving energy. Accessed

Dec. 20, 2023. https://www.daad.de/en/study-and-research-in-germany/plan-your-studies/saving-

energy/

Dufrenne, J., P. Soentoro, S. Tatini, T. Day, and S. Notermans. 1994. Characteristics of *Bacillus*

cereus related to safe food production. Int. J. Food Microbiol. 23(1):99-109.

https://doi.org/10.1016/0168-1605(94)90225-9

EFSA. 2005. Opinion of the Scientific Panel on biological hazards (BIOHAZ) on *Bacillus cereus*

and other *Bacillus* spp in foodstuffs. EFSA J. 3(4):175. https://doi.org/10.2903/j.efsa.2005.175

Elzhov, T. V., K. M. Mullen, A.-N. Spiess, and B. Bolker. 2022. Package 'minpack.lm'.

Accessed Dec. 12, 2023. https://cran.r-project.org/web/packages/minpack.lm/minpack.lm.pdf

FDA (Food and Drug Administration) and Health Canada. 2015. Joint FDA/Health Canada

quantitative assessment of the risk of *Listeriosis* from soft-ripened cheese consumption in the

United States and Canada. Accessed Sep. 22, 2023. https:/ / www.fda .gov/media/90488/

download

FDA (Food and Drug Administration). 2023. Are You Storing Food Safely? Accessed Dec. 12,

2023. https://www.fda.gov/consumers/consumer-updates/are-you-storing-food-safely

Garre, A., J. Koomen, H. den Besten, and M. Zwietering. 2022. Package 'biogrowth'. Accessed

Dec. 12, 2023. https://cran.r-project.org/web/packages/biogrowth/biogrowth.pdf

Guinebretière, M. H., F. L. Thompson, A. Sorokin, P. Normand, P. Dawyndt, M.

Ehling□Schulz, B. Svensson, V. Sanchis, C. Nguyen□The, and M. Heyndrickx. 2008.

Ecological diversification in the *Bacillus cereus* group. Environ. Microbiol. 10(4):851-865.

https://doi.org/10.1111/j.1462-2920.2007.01495.x

Huang, J., Y. Luo, B. Zhou, J. Zheng, and X. Nou. 2019. Growth and survival of *Salmonella*

enterica and *Listeria monocytogenes* on fresh-cut produce and their juice extracts: Impacts and

interactions of food matrices and temperature abuse conditions. Food Control. 100, 300-304.

https://doi.org/10.1016/j.foodcont.2018.12.035

Jain, C., L. M., Rodriguez-R, A. M., Phillippy, K. T., Konstantinidis, and S. Aluru. 2018. High

throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat.

Commun. 9(1), 5114. https://doi.org/10.1038/s41467-018-07641-9

James, C., B. A. Onarinde, and S. J. James. 2017. The use and performance of household

refrigerators: A review. Compr. Rev. Food. Sci. Food. Saf. 16(1), 160-179.

https://doi.org/10.1111/1541-4337.12242

Jeßberger, N., V. M. Krey, C. Rademacher, M.-E. Böhm, A.-K. Mohr, M. Ehling-Schulz, S.

Scherer, and E. Märtlbauer. 2015. From genome to toxicity: a combinatory approach highlights

the complexity of enterotoxin production in *Bacillus cereus*. Front. Microbiol. 6:560.

https://doi.org/10.3389/fmicb.2015.00560

Jessberger, N., R. Dietrich, P. E. Granum, and E. Märtlbauer. 2020. The *Bacillus cereus* food

infection as multifactorial process. Toxins. 12(11), 701. https://doi.org/10.3390/toxins12110701

Jofré, A., M. L. Latorre-Moratalla, M. Garriga, and S. Bover-Cid. 2019. Domestic refrigerator

temperatures in Spain: Assessment of its impact on the safety and shelf-life of cooked meat

products. Food Res Int. 126, 108578. https://doi.org/10.1016/j.foodres.2019.108578

Kovac, J., H. den Bakker, L. M. Carroll, and M. Wiedmann. 2017. Precision food safety: a

systems approach to food safety facilitated by genomics tools. TrAC-Trend Anal. Chem. 96:52-

61. https://doi.org/10.1016/j.trac.2017.06.001

Lewin, A., G. Delage, F. Bernier, and M. Germain. 2019. Banked human milk and quantitative

risk assessment of *Bacillus cereus* infection in premature infants: a simulation study. Can. J.

Infect. Dis. Med. Microbiol. 2019. https://doi.org/10.1155/2019/6348281

Lott, T. T. 2023. Gram-Negative Bacteria and *Microbacterium* Spp. Represent Greater

Processing Challenges than Psychrotolerant Spore-Forming Bacteria Do for Extending the Shelf-

Life of High Temperature, Short Time (HTST) Pasteurized Fluid Milk. Doctoral dissertation.

Department of Food Science, Cornell University, Ithaca.

Lott, T. T., M. Wiedmann, and N. H. Martin. 2023. Shelf-life storage temperature has a

considerably larger effect than high-temperature, short-time pasteurization temperature on the

growth of spore-forming bacteria in fluid milk. J. Dairy Sci. 106(6), 3838-3855. https://doi.org/

10.3168/jds.2022-22832

Martens, D. E., C. Béal, P. Malakar, M. H. Zwietering, and K. Van'T Riet. 1999. Modelling the

interactions between *Lactobacillus curvatus* and *Enterobacter cloacae*: I. Individual growth

kinetics. Int. J. Food Microbiol. 51(1), 53-65. https://doi.org/10.1016/S0168-1605(99)00095-1

Mercier, S., S. Villeneuve, M. Mondor, and I. Uysal. 2017. Time–temperature management

along the food cold chain: A review of recent developments. Compr. Rev. Food Sci. Food Saf.

16(4):647-667. https://doi.org/10.1111/1541-4337.12269

Nakamura, L. 1998. *Bacillus pseudomycoides* sp. nov. Int. J. Syst. Bacteriol. 48(3):1031-1035.

10.1099/00207713-48-3-1031

Ndraha, N., H.-I. Hsiao, J. Vlajic, M.-F. Yang, and H.-T. V. Lin. 2018. Time-temperature abuse

in the food cold chain: Review of issues, challenges, and recommendations. Food Control.

89:12-21. https://doi.org/10.1016/j.foodcont.2018.01.027

Notermans, S., J. Dufrenne, P. Teunis, R. Beumer, M. Te Giffel, and P. P. Weem. 1997. A risk

assessment study of *Bacillus cereus* present in pasteurized milk. Food Microbiol. 14(2):143-151.

https://doi.org/10.1006/fmic.1996.0076

Pouillot, R., M. B. Lubran, S. C. Cates, and S. Dennis. 2010. Estimating parametric distributions

of storage time and temperature of ready-to-eat foods for U.S. households. J. Food Prot. 73:312–

321. https://doi.org/10.4315/0362-028X-73.2.312

Qian, C., S. Murphy, T. Lott, N. Martin, and M. Wiedmann. 2023. Development and deployment

of a supply-chain digital tool to predict fluid milk spoilage due to psychrotolerant sporeformers.

J. Dairy Sci. https://doi.org/10.3168/jds.2023-23673

Ratkowsky, D. A., J. Olley, T. McMeekin, and A. Ball. 1982. Relationship between temperature

and growth rate of bacterial cultures. J. Bacteriol. 149(1):1-5. https://doi.org/10.1128/jb.149.1.1-

5.1982

Riol, C. D., R. Dietrich, E. Märtlbauer, and N. Jessberger. 2018. Consumed foodstuffs have a

crucial impact on the toxic activity of enteropathogenic *Bacillus cereus*. Front. Microbiol.

9:1946. https://doi.org/10.3389/fmicb.2018.01946

R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. Accessed Dec 12, 2023. https://www.R-project.org

Stenfors Arnesen, L. P., A. Fagerlund, and P. E. Granum. 2008. From soil to gut: *Bacillus cereus*

and its food poisoning toxins. FEMS Microbiol. Rev. 32(4), 579-606.

https://doi.org/10.1111/j.1574-6976.2008.00112.x

Svensson, B., K. Ekelund, H. Ogura, and A. Christiansson. 2004. Characterisation of *Bacillus*

cereus isolated from milk silo tanks at eight different dairy plants. Int. Dairy J. 14(1), 17-27.

https://doi.org/10.1016/S0958-6946(03)00152-3

Thirkell, C., T. Sloan-Gardner, M. Kaczmarek, and B. Polkinghorne. 2019. An outbreak of

Bacillus cereus toxin-mediated emetic and diarrhoeal syndromes at a restaurant in Canberra,

Australia 2018. Accessed Jan 16, 2024. https://openresearch-

repository.anu.edu.au/handle/1885/202456

Tirloni, E., S. Stella, F. Celandroni, D. Mazzantini, C. Bernardi, and E. Ghelardi. 2022. *Bacillus*

cereus in Dairy Products and Production Plants. Foods. 11(17):2572.

https://doi.org/10.3390/foods11172572

Zoellner, C., M. Wiedmann, and R. Ivanek. 2019. An assessment of listeriosis risk associated

with a contaminated production lot of frozen vegetables consumed under alternative consumer

handling scenarios. J. Food Prot. 82(12), 2174-2193. https://doi.org/10.4315/0362-028X.JFP-19-

092

Figure 1. Boxplot that shows the within and between group variability of growth parameters
(i.e., lag and μ_{max} at 22°C for (A, D) 16 *B. cereus* group strains representing 5 phylogenetic
groups [group I (n=4), group II (n=8), group IV (n=12), group V (n=4), group VII (n=4)], (B, E)
14 *B. cereus* group strains at 10°C, representing 4 phylogenetic groups, and (C, F) 2 *B. cereus*
group strains from group I at 16°C (group I strains did not grow at 10°C)). Group III was
excluded from this plot since it only had one strain. ***P < 0.05 based on post hoc Dunn's test
performed following a Kruskal-Wallis test that found μ_{max} to be significantly different between
group I and group IV strains when grown at 22°C.

Figure 2. Base model prediction results. The percentage of simulated milk containers
contaminated with a given *B. cereus* group (i.e., I, II, III, IV, V, VII) that exceeded 5 $\log_{10}$
CFU/ml (A) and 3 $\log_{10}$ CFU/ml (B) on consumer home storage day 14, 21, or 35.

Figure 3. Tornado diagram for sensitivity analysis of Q0 that shows the percentage point
increases (when using the maximum Q0 as the input) and decreases (when using the minimum
Q0 as the input) from the base model prediction results (i.e., the percent of milk containers over
5 $\log_{10}$ CFU/ml on consumer storage day 35 when using the average Q0 as the input parameter)
for 16 *B. cereus* group strains.

837 **Table 1.** Number of all *B. cereus* group isolates (A) and cytotoxic *B. cereus* group isolates (B) in
838 all clusters and sub-clusters

(A) Number of all *B. cereus* group isolates								
Phylogenetic group[2]								
Cluster[1]	I	II	III	IV	V	VI	VII	VIII	Total
1	0	0	0	0	0	0	8	0	8
2	0	0	0	0	1	0	0	0	1
3	0	5	19	1	0	0	0	0	25
4	0	0	0	1	0	0	0	0	1
5	0	5	1	74	3	0	0	0	83
6	0	1	0	2	0	0	0	0	3
7	0	0	20	0	0	0	0	0	20
8	0	16	21	0	0	0	0	0	37
9	0	19	3	4	36	36	0	4	103
10	5	0	0	0	1	3	0	0	9
11	0	0	0	0	1	0	0	0	1
(B) Number of cytotoxic *B. cereus* group isolates only								
Phylogenetic group								
Cluster	I	II	III	IV	V	VI	VII	VIII	Total
1	0	0	0	0	0	0	1	0	1
2	0	0	0	0	0	0	0	0	0
3	0	3	1	0	0	0	0	0	4
4	0	0	0	0	0	0	0	0	0
5	0	1	0	42	0	0	0	0	43
6	0	0	0	2	0	0	0	0	2
7	0	0	0	0	0	0	0	0	0
8	0	0	0	0	0	0	0	0	0
9	0	1	0	2	4	0	0	0	7
10	4	0	0	0	0	0	0	0	4
11	0	0	0	0	0	0	0	0	0

839 [1]Isolates are clustered based on unique virulence gene profiles.
840 [2]Phylogenetic groups assigned by BTyper3 are used for sub-clustering.

842 **Table 2.** Variables used in the time-temperature distributions of the five-stage supply chain

Symbol	Variable name	Distribution	Unit	Reference
	Stage 1. Processing plant storage			
t_F	Storage time of milk containers	Uniform (1,2)	d	(Qian et al., 2023)
T_F	Storage temperature of milk containers	Uniform (3.5, 4.5)	°C	(Qian et al., 2023)
	Stage 2. Transportation from processing plant to retail			
t_T	Storage time of milk containers	Triangular (1, 10, 5)	d	(FDA and Health Canada, 2015)
T_T	Storage temperature of milk containers	Triangular (1.7, 10.0, 4.4)	°C	(FDA and Health Canada, 2015)
	Stage 3. Retail storage			
t_S	Storage time of milk containers	TruncNormal (0.042, 10.0, 1.821, 3.3)[1]	d	(Qian et al., 2023)
T_S	Storage temperature of milk containers	TruncNormal (-1.4, 5.4, 2.3, 1.8)	°C	(FDA and Health Canada, 2015)
	Stage 4. Transportation from retail to consumer's home			
t_{T2}	Storage time of milk containers	TruncNormal (0.01, 0.24, 0.04, 0.02)	d	Expert opinion
T_{T2}	Storage temperature of milk containers	TruncNormal (0, 10.0, 8.5, 1.0)	°C	Expert opinion
	Stage 5. Consumer storage			
t_H	Storage time of milk containers	1-35	d	
T_H	Storage temperature of milk containers	TruncLaplace (-1, 15, 4.06, 2.31)[2]	°C	(Pouillot et al., 2010)

843 [1] In a TruncNormal (a, b, c, d) distribution, a = min, b = max, c = mean, and d = standard deviation.
844 [2] In a TruncLaplace (a, b, c, d) distribution, a = min, b = max, c = location parameter, and d = dispersion.
845

 Table 3. Implementation of what-if scenarios and outcomes

Scenario	Implementation[1]	Modified time/temperature distribution[1]	Percent milk containers exceeding 5 logs on consumer storage day 35 (mean ± standard deviation)
Baseline	N/A	N/A	4.13±2.53
Scenario (i) A three-day delay on HTST milk product delivery from processing facility to retail without temperature abuse	Shift in the distribution of transportation time from processing plant to retail 3 units towards the right	t_T ~ Triangular (4, 13, 8)	4.13±2.53
Scenario (ii) A temperature abuse when a consumer transported the HTST milk products from retail back home in a hot summer	Shift in the distribution of transportation temperature from retail to consumer's home 25 units towards the right	T_{T2} ~ TruncNormal (25, 35, 33.5, 1.0)	4.44±3.03
Scenario (iii) A slight temperature deviation in consumer's home refrigerator	Shift in the distribution of consumer home storage temperature 1 unit towards the right	T_H ~ TruncLaplace (0, 16, 5.06, 2.31)	8.88±3.84
Scenario (iv) An increase in the variability of refrigeration temperatures at consumer's home	Increase in the dispersion parameter of the distribution of consumer home storage temperature by 1 unit	T_H ~ TruncLaplace (-1, 15, 4.06, 3.31)	10.81±3.15

847 [1]Details of variables used in the time-temperature distributions of the five-stage supply chain can be found in Table
848 2.

849 **Table 4.** The slope of secondary model and theoretical minimum growth temperature for 16
850 isolates

Isolate	Phylogenetic group	b^1	T_{min} $(°C)^2$
PS00125	I	0.30	9.22
PS00193	II	0.17	6.79
PS00457	II	0.14	5.88
PS00564	II	0.20	6.28
PS00518	II	0.20	6.86
PS00474	III	0.15	6.41
PS00413	IV	0.13	4.88
PS00433	IV	0.12	5.83
PS00495	IV	0.17	6.79
PS00407	IV	0.15	6.71
PS00402	IV	0.13	4.26
PS00649	IV	0.15	6.46
PS00570	V	0.14	5.42
PS00638	V	0.15	5.52
PS00194	VII	0.21	6.89
PS00536	VII	0.21	7.06

851 [1] b, slope of secondary model.
852 [2] T_{min}, theoretical minimum growth temperature.

853 **Table 5.** Percent milk containers over 5 $\log_{10}$ CFU/ml at consumer storage day 35 and 21

Isolate	Group	Base model		Scenario (i)[1]		Scenario (ii)[2]		Scenario (iii)[3]		Scenario (iv)[4]	
		d35	d21	d35	d21	d35	d21	d35	d21	d35	d21
Avg. of all isolates[5]		4.13	2.81	4.13	2.81	4.44	3.06	8.88	4.56	10.81	7.56
PS00125	I	1	1	1	1	1	1	3	2	2	2
PS00193	II	3	3	3	3	3	3	6	3	10	5
PS00457	II	3	3	3	3	3	3	10	3	12	9
PS00518	II	3	3	3	3	3	3	7	3	10	9
PS00564	II	5	3	5	3	5	3	12	7	13	10
PS00474	III	3	3	3	3	3	3	10	3	10	7
PS00402	IV	4	2	4	2	4	2	9	3	11	5
PS00407	IV	3	3	3	3	3	3	5	3	10	5
PS00413	IV	10	4	10	4	11	5	17	11	17	14
PS00433	IV	3	2	3	2	3	3	6	3	10	4
PS00495	IV	3	3	3	3	3	3	7	3	10	9
PS00649	IV	3	3	3	3	3	3	10	3	10	9
PS00570	V	6	3	6	3	10	3	14	10	14	10
PS00638	V	10	3	10	3	10	5	14	10	14	12
PS00194	VII	3	3	3	3	3	3	6	3	10	6
PS00536	VII	3	3	3	3	3	3	6	3	10	5

predicted by base model compared to what-if scenarios

[1]**Scenario (i),** a three-day delay on HTST milk product delivery from processing facility to retail without
temperature abuse.
[2] **Scenario (ii),** a temperature abuse when a consumer transported the HTST milk products from retail back home in
a hot summer.
[3]**Scenario (iii),** a slight temperature deviation in consumer's home refrigerator.
[4]**Scenario (iv),** an increase in the variability of refrigeration temperatures at consumer's home.
[5]Percent milk containers (%) over 5 $\log_{10}$ CFU/ml at consumer storage day 35 (d35) and day 21 (d21) averaged
across all 16 isolates.