Testing the evolutionary potential of an alpine plant: Phenotypic plasticity

in response to growth temperature far outweighs parental environmental

effects and other genetic causes of variation

Pieter A. Arnold[1,*], Shuo Wang[1,2], Rocco F. Notarnicola[1,3], Adrienne B. Nicotra[1],

Loeske E. B. Kruuk[1,4]

[1] Division of Ecology and Evolution, Research School of Biology, The Australian National

University, Canberra, ACT, Australia

[2] Liaoning Key Laboratory for Biological Invasions and Global Changes, College of

Bioscience and Biotechnology, Shenyang Agricultural University, Shenyang, Liaoning

Province 110866, China

[3] Evolutionary Biology Group, Faculty of Biology, Adam Mickiewicz University, Poznań,

Poland

[4] Institute of Ecology and Evolution, School of Biological Sciences, University of Edinburgh,

Edinburgh, EH9 3FL, UK.

[*]Corresponding author: Pieter A. Arnold, Email address: pieter.arnold@anu.edu.au

ORCIDs

PAA: 0000-0002-6158-7752

SW: 0000-0002-1700-1136

RFN: 0000-0001-9860-6497

ABN: 0000-0001-6578-369X

LEBK: 0000-0001-8588-1123

Running title: Plasticity and selection in response to warming

Keywords: Adaptive plasticity, genotype-by-environment, global change, heritability,

parental effects, phenotypic selection, thermal plasticity, thermal tolerance

Highlight

We found strong plasticity to growth environment in many phenotypic traits, but little effect of parental environment, revealing capacity to respond rapidly to climate warming, and potential for evolutionary change.

Abstract

Phenotypic plasticity and rapid evolution are fundamental processes by which organisms can maintain their function and fitness in the face of environmental changes. Here we quantified the plasticity and evolutionary potential of an alpine herb *Wahlenbergia ceracea*. Utilising its mixed-mating system, we generated outcrossed and self-pollinated families that were grown in either cool or warm environments, and that had parents that had also been grown in either cool or warm environments. We then analysed the contribution of environmental and genetic factors to variation in a range of phenotypic traits including phenology, leaf mass per area, photosynthetic function, thermal tolerance, and reproductive fitness. The strongest effect was that of current growth temperature, indicating strong phenotypic plasticity. All traits except thermal tolerance were plastic, whereby warm-grown plants flowered earlier, grew larger, produced more reproductive stems compared to cool-grown plants. Flowering onset and biomass were heritable and under selection, with early flowering and larger plants having higher relative fitness. There was little evidence for transgenerational plasticity, maternal effects, or genotype-by-environment interactions. Inbreeding delayed flowering and reduced reproductive fitness and biomass. Overall, we found that *W. ceracea* has the capacity to respond rapidly to climate warming via plasticity, and the potential for evolutionary change.

Introduction

Climate change is exposing organisms to ever increasing mean temperatures and more frequent extreme events (Harris *et al.*, 2018). Temperature is a universal, pervasive environmental variable that limits a species' occupiable niche (Walther, 2003; Tattersall *et al.*, 2012; Nievola *et al.*, 2017). It is therefore unsurprising that steadily warming mean temperatures have been observed to affect biological processes, such as phenology, globally (Walther *et al.*, 2002; Parmesan and Yohe, 2003). For example, in warmer years the date of flowering onset in plants occurs considerably earlier, and has been shown to be tightly related to the pattern of mean climatic warming (Parmesan and Hanley, 2015), especially in high elevation plants (Giménez-Benavides *et al.*, 2011; Dorji *et al.*, 2020). Indirect reproductive effects due to phenological changes from warming, as well as direct effects on fertility and reproductive traits, can also have consequences for individual- and population-level fitness (Anderson, 2016). To determine the realised impact of warming on natural selection and evolutionary processes, we must assess how warming affects phenotypic responses and fitness, and how these vary among individuals in a population.

There are several mechanisms, not mutually exclusive, by which plants can tolerate climatic and environmental changes. Plants could already have a high natural resilience to warmer temperatures, thereby partially or completely avoiding thermal stress (De Kort *et al.*, 2020; Andrew *et al.*, 2023). Alternatively, phenotypic plasticity – the capacity of a single genotype to exhibit multiple phenotypes based on changes in the environment (Bradshaw, 1965, 2006) – could facilitate short-term changes to their phenotype, to limit exposure to stress effects or to mitigate damage (Nicotra *et al.*, 2010; Fox *et al.*, 2019; Brooker *et al.*, 2022). For instance, exposure to increased mean temperatures can benefit some plants by acclimating (or priming) them to face more stressful conditions in a more prepared state, so that extreme events have a lesser impact (Hilker *et al.*, 2016). Plants generally have limited capacity to move to escape unfavourable temperatures due to the constraints and costs of dispersal, therefore responding to local conditions through plasticity is likely to be evolutionarily favoured (Edelaar *et al.*, 2017).

The widespread occurrence of phenotypic plasticity generates assumptions that it is an evolutionary adaptation to environmental heterogeneity (Hendry, 2016). In other words, selection will favour plasticity because it improves an individual's performance or fitness in a particular environment and would therefore be considered adaptive (Bonser, 2021). Here, Genotype × Environment interactions (G×E) specifically refers to variation in plastic

responses among genotypes within a population. That is, the expression of phenotypes in

response to the environment (E) varies depending on the genotype (G). Although phenotypic

plasticity is at times seen as equivalent to G×E, plasticity can result simply in response to

variation in E, without there being significant variation attributable to G. For plasticity to

evolve, however, there has to be genetic variation in plasticity on which selection can act,

hence a significant G×E interaction is a requisite for demonstrating the adaptive significance

of plasticity (Josephs, 2018). Yet, evidence suggests that plasticity being costly is about as

common as it is being beneficial; the exact balance will depends on the severity of the stress

and the relative strengths of the costs and benefits of plasticity (van Buskirk and Steiner,

2009; Auld *et al.*, 2010; Hendry, 2016). Indeed, there is emerging evidence that trait

canalisation (i.e., no plasticity, also called robustness) in response to warming may be

favoured when costs of plasticity are high or plasticity is maladaptive (Stinchcombe *et al.*,

2004; Arnold *et al.*, 2019b; Svensson *et al.*, 2020).

While plasticity in response to warming is common in plants (Nicotra *et al.*, 2010),

the degree to which different traits respond can differ markedly. From a global meta-analysis,

plasticity in leaf morphology, plant biomass, and several physiological traits (including

chlorophyll content and photosynthetic efficiency: F_V/F_M and ϕPSII) all indicate strong

responses to increasing mean annual temperature (Stotz *et al.*, 2021). However, within a

species there may be substantial inter-trait variation in responses. For example, the alpine

herb *Wahlenbergia ceracea* exhibits different patterns of plasticity in response to

temperature, depending on trait type. Warmer, but not stressful, growth temperatures increase

leaf mass per area (LMA), heat tolerance traits, and reproductive output (flowers and

capsules), and also induce earlier flowering, decreased photosynthetic efficiency, and

lowered biomass (Arnold *et al.*, 2022). Along with the variation in plastic responses among

traits of different types, there is also substantial intraspecific variation in plastic responses to

temperature (Arnold *et al.*, 2022).

The patterns of selection acting on plant traits may also be affected by climate and

environmental factors. Climate change has already had widespread effects on plant traits,

especially the timing of reproductive events, where earlier flowering under warmer

conditions almost ubiquitously increases fitness and is therefore under strong directional

selection (Franks *et al.*, 2007; Anderson *et al.*, 2012; Anderson, 2016; Wadgymar *et al.*,

2018b; Ehrlén and Valdés, 2020). The relationship between functional traits and fitness

depends on the environment, yet other than phenology, empirical tests of selection on

functional traits due to climate warming are uncommon (Geber and Griffen, 2003; Kimball *et*
al., 2012). Totland (1999) found that *Ranunculus acris* were under selection for more flowers
in both control and warmed field conditions, but selection for larger leaf size was only
apparent in control conditions. Using an urban (warmer and drier) environment as a climate
change analogue to test for its effect on selection, Lambrecht *et al.* (2016) found strong
evidence for selection on functional traits, including increased plant size, leaf number,
specific leaf area, and later senescence in urban conditions in *Crepis sancta*. Establishing
which phenotypic traits are correlated with fitness (i.e., under selection) under benign and
stressful conditions is a crucial component for understanding the potential for rapid
evolutionary responses (Anderson, 2016). For evolutionary change to keep pace with climate
change, the phenotypic trait under selection must also have heritable variation on which
selection can act (Scheiner *et al.*, 2020). However, there are also alternative mechanisms of
affecting phenotype change across generations.

Transgenerational plasticity or parental environmental effects can be demonstrated
when the conditions to which a parent is exposed shape offspring phenotype function and
fitness in their environment (Mousseau and Fox, 1998; Bonduriansky, 2021), most cases of
which are mediated by maternal effects (Herman and Sultan, 2011). A factorial design of at
least two offspring and two parental environments that match or mismatch allows
transgenerational and within-generation plasticity to be tested simultaneously (Uller *et al.*,
2013). Adaptive transgenerational plasticity (also called 'anticipatory parental effects')
theory posits that selection on parental responses to their environment confers benefits to the
offspring when their environment matches that of the parents, particularly when the
environmental conditions are stressful (Uller, 2008; Herman and Sultan, 2011). Conversely,
when the parent and offspring environments are not stable or are unpredictable, or when they
mismatch (e.g., due to change in season or annual change), there may be a cost to the
offspring of producing a phenotype that reflects the parental environment rather than the
current environment (Engqvist and Reinhold, 2016).

A previous meta-analysis has revealed little evidence for transgenerational plasticity
conferring a clear benefit in plants, especially when close proxies for fitness are used (Uller
et al., 2013). Yet, there are cases of strong parental effects in response to temperature in
plants. For example, Whittle *et al.* (2009) found *Arabidopsis thaliana* plants substantially
increased reproductive output under relatively hot conditions (30°C) when prior generations
had also been grown in the same hot conditions. There is also generally stronger evidence for

transgenerational plasticity affecting early life traits like seed germination. For example, Wadgymar *et al.* (2018a) showed greater and more variable transgenerational plasticity than within-generation plasticity in germination of *Boechera stricta* plants across an elevation gradient. The effects of parental temperature in *W. ceracea* have also been found to affect germination and dormancy patterns, but to a lesser extent than the temperatures in which seeds germinated (Wang *et al.*, 2021; Notarnicola *et al.*, 2023b).

Climatic warming is expected to increase rates of self-pollination in plants with mixed-mating strategies (i.e., those that can facultatively self-pollinate in the absence of cross-pollination), due to phenological mismatches with pollinators (Hegland *et al.*, 2009). However, while self-pollination is a contingency strategy that may assure reproductive success in the face of climate warming, inbreeding depression may be worsened in stressful environments (Armbruster and Reed, 2005) and it also reduces adaptive potential (Hamann *et al.*, 2021). While the effects of inbreeding under some stressors associated with climate change (drought, herbivory, nutrient deficiency) have been studied in mixed-mating species (Hamann *et al.*, 2021), to our knowledge only one study has investigated the effects of inbreeding with warming in a mixed-mating species (Wang *et al.*, 2021).

In the current study, we used a large-scale glasshouse experiment to measure a suite of phenotypic traits on outcrossed and self-pollinated *W. ceracea* plants, from families that were grown in either cool or warm environments that had parents that were grown in either cool or warm environments. We addressed the following questions (Q):

1) To what extent do a suite of traits respond to growth temperature through phenotypic plasticity?
2) Is there any evidence of transgenerational plasticity or benefits for offspring that are grown under conditions that match the conditions in which their parents were grown?
3) Are the phenotypic traits heritable, and are there either maternal effects or G×E interactions?
4) What is the direction and strength of selection on the traits and how does it vary with growth temperature?
5) Are there effects of inbreeding on the suite of traits, and does any inbreeding depression vary with temperature?

For Q1, we hypothesised that the warmer growth temperature would be stressful, reducing plant function and biomass, but that plants would also respond by flowering earlier, which may compensate to result in equal or higher fitness, and that acclimatory processes

would improve heat tolerance. For Q2, we hypothesised that the parental environment would

have a small effect on offspring phenotype (i.e., development of seed under parental

temperatures that matched the offspring growth temperature would benefit fitness compared

to mismatched offspring). For Q3, we hypothesised that the heritable traits would be

flowering onset, biomass, LMA, and heat tolerance, based on previous findings of highly

plastic responses to temperature and intraspecific variation in these responses (Arnold *et al.*,

2022). For Q4, we hypothesised that traits under selection would be the same as those

heritable in Q3. For Q5, we predicted that there would be evidence for inbreeding depression

primarily in fitness. Addressing this series of questions together will elucidate the

evolutionary potential of functional responses to warmer growth temperatures that are

expected under future climate change.

Materials and methods

Species description, seed source, and F0-F2 generations

Wahlenbergia ceracea Lothian (Campanulaceae) waxy bluebell is a short-lived, protandrous,

and facultatively autogamous biennial herb that is sparsely distributed across south-eastern

Australia and Tasmania (Nicotra *et al.*, 2015). For this study, seeds were collected between

1590 m and 2100 m a.s.l. from Kosciuszko National Park, NSW, Australia (36.43°S,

148.33°E) in 2015 and 2016 (see Notarnicola *et al.*, 2021 for further details), and brought to

The Australian National University, Canberra, ACT, Australia. These seeds formed the F0

generation: we describe below their rearing and subsequent breeding design for F1, F2, and

F3 generations, with the F3 plants then being used for the analyses here of phenotypic

plasticity, transgenerational plasticity, heritable genetic variance, and inbreeding depression.

F0, F1, and F2 generations

Detailed descriptions of the conditions and breeding design for producing plants for

generations F0 to F3 in this study have been given previously (Notarnicola *et al.*, 2021; Wang

et al., 2021; Arnold *et al.*, 2022). Briefly, F0 plants grown from field-collected seeds were

raised in glasshouses that mimicked average alpine summer temperatures (25/15°C during

germination and growth, which was reduced to 20/15°C at peak flowering). F1 plants were

produced by crossing 48 F0 plants as pollen donors with 48 F0 plants as ovule donors (96

plants from 63 unique F0 families) to produce 48 F1 families by hand-pollination. F1 plants

were raised under the same conditions as the F0 plants (Supplementary Fig. S1). The F2

generation was produced using a partial-diallel (maximising the number of parents used to

generate families) half-sib breeding design in which 12 F1 plants were used as pollen donors

and each outcrossed with at least four unrelated pollen receivers (48 pollen receivers in total).

F2 plants were grown in glasshouses under two temperature regimes ('*cool*': 20/15°C

day/night conditions and '*warm*' 30/25°C day/night conditions) that hereafter constitute the

'*parental temperature*'. These two temperature regimes represent the cool conditions to

which these alpine plants are already well-adapted, and the warm conditions simulate

substantial warming that could occur under an extreme climate change scenario.

F3 breeding design and pedigree

Within each parental temperature treatment, 12 F2 lines were assigned as pollen donors and

these were each outcrossed with four unrelated pollen receivers, such that cool and warm F2

lines were full-sibs of the same F1 parentage. Each of the F2 parent plants in each treatment

were also self-pollinated to generate 12 F3 inbred maternal and 11 inbred paternal lines, in

addition to the 12 outcrossed F3 lines that were reciprocally outcrossed from paired F2 pollen

donors from each parental temperature treatment (Supplementary Fig. S1). In total, there

were 96 outcrossed F3 families and 46 inbred F3 families, all controlled by hand-pollination

and bagging flowers. The capsules that were formed after hand-pollination of the F2 plants

were removed after they had opened, dried, and the seeds had browned. All seeds taken from

these capsules were stored in a desiccator for at least seven weeks at ~20°C and 15-20%

relative humidity before the beginning of the experiment.

F3 growth experiment and temperature treatments

For each F3 cross, 20-30 seeds from a single capsule were sown across two 50 mm Petri

dishes containing 1% agar, each corresponding to a growth temperature treatment. The dishes

were sealed and moved to an incubator for cold stratification at 5°C in darkness for six weeks

to release seeds from dormancy, thereby improving subsequent germination success (Wang *et*

al., 2021). At least eight healthy seedlings per dish were transplanted into punnets containing

seed raising mix (Debco Pty Ltd, VIC, Australia) and moved into ideal common glasshouse

conditions (25/18°C) to grow for two months. The 25/18°C temperature regime was chosen

based on the daytime temperature that increased growth rates of *W. ceracea* seedlings

(Arnold *et al.*, 2022), while maintaining natural cooling at night. Liquid fertiliser (Thrive

Soluble All Purpose Plant Food; Yates, NSW, Australia) at a concentration of 0.5 g L^{-1} was

added regularly to promote growth and seedlings were watered twice daily. Up to eight

healthy seedlings (6-40 mm in diameter) from each family were transplanted into individual

plastic pots (125 mm diameter, ~600 mL) filled with soil suitable for Australian natives

combined with 3 g L^{-1} of low phosphorus slow-release fertiliser (Scotts Osmocote Plus Trace

Elements: Native Gardens; Evergreen Garden Care Australia, NSW, Australia).

The potted F3 plants were moved to their *growth temperature* treatment seven days

after transplantation (commenced 7 November 2019). For the *cool* treatment, plants were

placed in a large glasshouse room set to 20/15°C (day/night) under natural photoperiod and

for the *warm* treatment, plants were placed in an adjacent glasshouse room set to 30/20°C

(day/night). An automatic shade screen was active between 12:00-14:30 and when external

temperatures exceeded 30°C for the cool treatment and 33°C for the warm treatment, to

prevent excess solar radiation and overheating of the glasshouses, otherwise plants received

natural light. Plants were watered daily initially until growing well and then watered to

weight to ensure that the soil did not dry out and water was not limiting. As such, the plants

in the warm treatment received more frequent watering than did the cool treatment. Pest

treatments (VectoBac larvicide for treating fungus gnats and sulphur evaporation for treating

powdery mildew) were conducted as required and liquid fertiliser 1 g L^{-1} was applied

approximately fortnightly as required for maintaining healthy growth. Within each treatment,

we randomised the distribution of one plant from each family to each of four blocks

consisting of 29 columns and five rows, which was replicated across both rooms. Not all

families had eight healthy plants, but each had at least five plants that were distributed

randomly across both treatments. Empty pots were used in place of a missing plant, so that

the position layout was preserved across all blocks. In total, there were 1,024 plants (out of an

ideal 1,152) at the beginning of the growth experiment (512 in each growth treatment).

F3 phenotypic trait measurement

We measured a range of phenotypic traits on the F3 plants grown in the two temperature

treatments, where chosen traits characterise many important plant functions, which have

potential to respond to environmental factors. The date of the first flower produced by every

plant was recorded throughout the experiment, which was checked at least every 2-3 days,

and this was converted to the day of *flowering onset* since the beginning of the growth

temperature treatments. After four weeks in the treatments (9 January 2020), we began

phenotyping leaves for thermal tolerance limits and photosynthetic parameters using

chlorophyll fluorescence. We used a Pulse Amplitude Modulated (PAM) chlorophyll

fluorescence imaging system (MAXI-Imaging-PAM, Heinz Walz GmbH, Effeltrich,

Germany) to measure the *photosystem II (PSII) operating efficiency (ϕPSII)* and the

maximum quantum efficiency of PSII photochemistry (F_V/F_M). We also measured the heat and

cold tolerance limits of leaves by measuring the temperature-dependent change in minimal

chlorophyll fluorescence (T-F_0) using controllable thermoelectric Peltier plates (plate: CP-

121HT; controller: TC-36-25; TE-Technology, Inc., Michigan, USA) in conjunction with

imaging fluorimeters. We determined the critical threshold temperatures at which F_0

increased rapidly to indicate the impairment of PSII, as measures of physiological *heat*

tolerance ($T_{crit-hot}$) and *cold tolerance ($T_{crit-cold}$)*.

For the PSII and heat tolerance assays, an array of 30 leaves was taped to paper and

placed on the Peltier plate surface with heavy glass on top to maximise thermal transfer

between the plate and leaves. Every second leaf had a calibrated type-T thermocouple

attached to the underside of the leaf in a checkerboard pattern, which were interfaced with a

DataTaker datalogger (DT80; Lontek, NSW, Australia) recording every 5 s, so that leaf

temperature could be measured or approximated directly. The Peltier plate was held at 25°C

and leaves were initially light-adapted under 280 μmol photons m^{-2} s^{-1} for 30 min to obtain

stable minimal fluorescence (F_0') values, after which we applied three saturating pulses at

~4,000 μmol photons m^{-2} s^{-1} for 720 ms each at 2 min intervals to determine the maximal

fluorescence (F_M') when PSII reaction centres were closed. Variable fluorescence in the light

($F_V' = F_M' - F_0'$) was then used to calculate ϕPSII (equivalent to F_V'/F_M') (Baker, 2008) from

the mean of the three measurements. These leaves were then dark-adapted for 25 min to

reach a stable minimal chlorophyll fluorescence in the dark (F_0) prior to applying a single

saturating pulse to measure F_V/F_M, which is the equivalent of ϕPSII in the dark (Baker,

2008). After 1 min, we began recording F_0 values continuously at 20 s intervals using a weak

blue pulse modulated measuring light (0.5 μmol photons m^{-2} s^{-1}) at 1 Hz to measure

chlorophyll fluorescence without driving PSII photochemistry. We measured the T-F_0 while

heating the Peltier plate to 65°C at a rate of 30°C h^{-1}.

Another similar system was used to measure cold tolerance limits in a cool room set

to 6±2°C. The system was the same as described above except all 30 leaves had

thermocouples attached to leaf undersides and the Peltier plate was set to 6°C to match the

ambient cool room temperature. Leaves were dark adapted for 25 min before the temperature

was reduced to –20°C at 15°C h^{-1}, while recording F_0 values continuously at 20 s intervals,

and thermocouple (leaf) temperature every 5 s with a DataTaker datalogger (DT85-S4).

The T-F_0 curve shows a stable, slow rise of F_0 values until a critical temperature

threshold occurs where F_0 rises rapidly. The critical limit for both heat tolerance ($T_{\text{crit-hot}}$) and

cold tolerance ($T_{\text{crit-cold}}$) is the temperature at the inflection point of intersecting regression

lines for each of the T-F_0 rise phases of relative F_0, which is used as an indication of the

thermal tolerance limits of photochemistry (Arnold *et al.*, 2021). In both cases, the inflection

point was calculated using a break-point regression analysis of the leaf temperature and

relative F_0 values using the *segmented* R package (Muggeo, 2017) in the R Environment for

Statistical Computing v4.3.1 (R Core Team, 2023).

The hailstorm and measurements at harvest

Unfortunately, on 20 January 2020, one of the most severe hailstorms in recorded history in

Canberra damaged the glasshouses containing the plants (Rickards and Watson, 2020). There

was no direct damage to the study plants because a shade screen was present, but the two

controlled temperature treatments were lost due to damaged glasshouse infrastructure.

Cooling systems were restored 21 January 2020, but heating could not be restored. To ensure

we did not therefore lose the investment in the long-term experiment, we took measurements

on all plants immediately following the hailstorm. Specifically, we measured *chlorophyll*

content using a handheld chlorophyll meter (SPAD-502; Konica Minolta Inc., Osaka, Japan)

and *leaf mass per area (LMA)* on all plants within four days following the storm. Three

healthy leaves were removed from each plant, immediately measured for chlorophyll content,

and then scanned for leaf area before being placed in a drying oven at 60°C for > 72 h for

weighing and calculating LMA. We had initially planned to continue the experiment until

autumn (a further eight weeks), reduce temperatures to induce senescence, and then measure

lifetime fitness and biomass. However, since controlled senescence was not possible, and

since different plants were at various stages of flowering and producing seed at the time of

the storm, we used a proxy for lifetime fitness across all plants: the *total number of*

reproductive stems on an individual (i.e., flowers, capsules, and hardened, brown stems that

clearly indicated a capsule had matured on the stem). We confirmed that the total number of

reproductive stems was a suitable index of fitness by estimating its correlation with the

harvested capsule mass × number of capsules weighed from a representative subset of 100

plants. We found a strong correlation (Pearson's $r = 0.873 \pm 0.056$), which did not differ

among treatments (Supplementary Table S2, Supplementary Fig. S2). On 24-25 January

2020, we counted the number of reproductive stems on all plants, as well as collecting mature

capsules. However, due to the hailstorm alternative controlled temperature growth space was

limited and we elected to systematically harvested half of all plants (blocks 2 and 4)

following the reproductive stem count. We harvested these plants and measured *dry above-*

ground biomass in bags for drying at 60°C for > 72 h and subsequent weighing.

Plants from the blocks that were not harvested immediately following the hailstorm

(blocks 1 and 3) were moved to four controlled environment Growth Capsules (Photon

Systems Instruments, Brno, Czech Republic) run by the Australian Plant Phenomics Network

at ANU where we aimed to continue the experiment. The Growth Capsules were set to match

the glasshouse conditions as best possible, however the plants did not thrive in the Growth

Capsules due to lower light, reduced air flow, and higher humidity. After 10 days in Growth

Capsule conditions, we therefore measured thermal tolerance and chlorophyll fluorescence

traits on the plants that had not yet been measured, and then ceased the experiment on 13-14

February 2020. We repeated the reproductive stem count and added new stems to the count

from 24-25 January 2020, and then harvested these plants for biomass as above.

The sample sizes for each trait were $n = 1,023$ for number of reproductive stems,

$n = 1,003$ for flowering onset, $n = 988$ for above-ground biomass, $n = 984$ for chlorophyll

content, $n = 975$ for LMA, $n = 717$ for ϕPSII, $n = 717$ for F_V/F_M, $n = 685$ for $T_{\text{crit-hot}}$, and

$n = 707$ for $T_{\text{crit-cold}}$. The inherent differences between the pre- and post-hailstorm

measurements and harvesting are explicitly accounted for in our statistical analyses.

Statistical analyses

For all analyses of the F3 phenotypic traits, models were fit using the R package *brms*

(Bürkner, 2018) in the R Environment for Statistical Computing v4.3.1 (R Core Team, 2023).

All *brm* models were run using four chains, each with 4000 iterations, 2000 of which were

sampling, with *adapt_delta* ≥ 0.99 and *max_treedepth* = 15 so that the majority of $\hat{R} \leq 1.005$,

indicating that chains had effectively mixed. All response variable distributions exhibited

some skewness, therefore we set *skew_normal* distributions for the univariate *brm* models,

which are an extension of the normal (Gaussian) distribution family that also estimate a skew

parameter. We verified that skew-normal models were a good fit to the data and that they

were a better fit than models using a Gaussian distribution with posterior predictive checking

(Gabry *et al.*, 2019). To facilitate model convergence, ϕPSII and F_V/F_M were both scaled by a

factor of ten to avoid very small parameter estimates.

To test the main effects of growth temperature, parental temperature, and inbreeding

on each trait and its plasticity, we initially fit univariate random regression mixed models

(RRMMs; Arnold *et al.*, 2019a) that included a structured pedigree (often called an 'animal

model'; Kruuk, 2004; Wilson *et al.*, 2010), following the R *brms* form:

$$y \sim Growth\ temp. \times Parental\ temp. + Inbreed + Hail +$$
$$(1|Block) + (1|Maternal) + (1 + Growth\ temp.|A)$$

y is the phenotypic trait. Fixed effects were *Growth temp.*, a two-level factor of growth

temperature; *Parental temp.*, a two-level factor of parental temperature; and their interaction;

Inbreed, a two-level factor of inbreeding (outcrossed or self-pollinated); and *Hail*, a two-level

factor of whether the measurement was taken before or after the hailstorm. Random

intercepts were: *Block*, a four-level factor of experimental block; *Maternal*, an identification

term for the F2 ovule donor to quantify maternal effects with 102 levels; and *A*, the additive

genetic component with covariance structure defined by a pedigree of relatedness values

among individuals, which was converted into an inverse *A* matrix using the *MCMCglmm*

package (Hadfield, 2010). We included the random slope term *Growth temp.* with the *A* term

to test for a genotype × environment (G×E) interaction. To evaluate whether the G×E term

was important, we compared models with and without the *Growth temp.* slope term using

leave-one-out cross validation (LOO-CV) to estimate predictive accuracy of each candidate

model (Vehtari *et al.*, 2017). We then calculated Bayesian stacking weights, which evaluate

the average performance of the combined posterior predictive distribution of candidate

models (Yao *et al.*, 2018). We report the full model including the slope term given that our

models were not overparameterized relative to our sample sizes. We include R^2 values for

mixed-effects models: marginal R^2 (mR^2) to estimate variance explained by fixed effects and

the difference between mR^2 and conditional R^2 (cR^2) to estimate variance explained by

random effects (Nakagawa and Schielzeth, 2013) using the *performance* package (Lüdecke *et*

al., 2021).

To calculate narrow-sense heritability h^2, we took the posterior distribution of the

additive genetic variance V_A from the animal model and divided it by the total phenotypic

variance V_P, where $V_P = (V_A + V_B + V_M + V_R)$, and V_B is block variance, V_M is maternal

variance, and V_R is residual variance. Since we included a maternal effect term in the models,

we also estimated the contribution of direct maternal effects m^2 as V_M/V_P.

To test for linear (directional) and quadratic (stabilising or disruptive) selection on

traits, we fit multiple regression models of the trait and fitness, similarly to Noble *et al.*

(2013). We estimated standardised selection gradients by converting the number of

reproductive stems, which was our proxy for fitness, to relative fitness (*w*, by dividing by the

mean of each growth treatment) and each trait was mean-centred and standardised to unit

variance (Lande and Arnold, 1983), for the overall (all plants), cool-grown plants, and warm-

grown plants separately. For the overall model we also included linear and non-linear

interaction terms with the trait and temperature to determine if selection varied with

temperature. Linear selection gradients (β) came from regression models without quadratic

and the growth temperature × parental temperature interaction terms, whereas quadratic

selection gradients (γ) come from models including these terms (Lande and Arnold, 1983).

Quadratic terms and their 95% credible intervals (95% CI) were doubled prior to reporting,

such that they can be interpreted as stabilising or disruptive selection gradients (Stinchcombe

et al., 2008). Positive linear selection coefficients can be interpreted as directional selection

where individuals with larger phenotypic trait values have higher relative fitness on average.

Positive quadratic selection coefficients can be interpreted as individuals with trait values at

the edges of the trait distribution have higher relative fitness on average (convex function

shape; disruptive selection). Negative quadratic selection coefficients can be interpreted as

individuals with trait values in the centre of the trait distribution have higher relative fitness

on average (concave shape; stabilising selection). A trait that has both significant linear and

quadratic coefficients indicates an overarching directional change with a non-linear shape.

Note on the effects of the hailstorm and later harvest date or later trait measurements

The impact of the hailstorm (which caused a delay in measurement of some traits and later

harvest time for a subset of plants) was evident in some traits. Importantly, all models in our

analyses included a term to account for this effect. As expected, plants that were harvested

later had a greater number of reproductive stems and greater biomass than plants harvested

immediately after the hailstorm. The plants that had traits measured post-hailstorm also had

slightly higher chlorophyll content and F_V/F_M, but ϕPSII and the thermal tolerance traits

($T_{\text{crit-hot}}$ and $T_{\text{crit-cold}}$) were unaffected by the hailstorm (Tables 1 and 2).

Results

Q1: To what extent do a suite of phenotypic traits respond to growth temperature through phenotypic plasticity?

To test for plasticity in trait responses to temperature, we compared the change in mean trait value between cool and warm growth temperatures among parental temperature and cross type (self-pollinated *vs* outcrossed) groups. We present the overall mean effects in Fig. 1 and mean treatment- and family-level reaction norms in Supplementary Fig. S3, as well as summary statistics for each trait in Supplementary Table S1. Warm growth temperature had a significant positive effect on reproductive fitness, biomass, chlorophyll content, LMA, F_V/F_M, and ϕPSII (Fig. 1A,C-G, Tables 1, 2). Flowering onset also occurred significantly earlier (7.8 days on average) in warm-grown plants (Fig. 1B, Table 1). Seven traits showed significant phenotypic plasticity to growth temperature, with extensive variation around these average effects (Supplementary Fig. S3). There was no evidence of growth temperature effects on either heat or cold tolerance of PSII. Although the warm-grown plants exhibited a slightly higher $T_{crit\text{-}hot}$ than the cool-grown plants, as would be expected with a thermal acclimation response, this difference was not significant and the average response was canalised (Fig. 1H, Table 2). Similarly, $T_{crit\text{-}cold}$ did not differ significantly between treatments and was also, on average, canalised (Fig. 1I, Table 2).

Q2: Is there any evidence of transgenerational plasticity or benefits for offspring that are grown under conditions that match the conditions in which their parents were grown?

In this experiment, we used a factorial design to separate the effects of growing parental plants under relatively cool and warm temperatures and the subsequent effects of their offspring growing under the same (matched) or opposite (mismatched) temperature regimes. We hypothesised that development of seed under the parental temperature that matched the offspring growth temperature (e.g., cool × cool, or warm × warm) would exhibit phenotypes that performed better than mismatched offspring (e.g., cool × warm, or warm × cool). However, we found no evidence that the parental temperature had any effect on any of the measured traits (Fig. 1, Supplementary Fig. S4), nor of any significant interactions between growth and parental temperatures (Tables 1 and 2). Therefore, there was no evidence in any trait that matching parent-offspring environments was beneficial, nor was there any evidence that mismatching was detrimental (Supplementary Figs S5 and S6).

Q3: Are the phenotypic traits heritable, and are there either maternal effects or G×E

interactions?

Estimates of heritability were relatively low across all traits (ranging from 0.01 to 0.14), with

flowering onset, biomass, and chlorophyll content being the only traits for which there was

support for a non-zero heritability (Table 3). While maternal effects were included in all

models, estimates of their variance components were small and the credible intervals of the

estimates were not clearly distinct from zero (Table 3).

Flowering onset had a clear negative correlation between V_A intercept and slope,

indicating that families that flowered early on average were those with the lowest plasticity.

However, we only found support for G×E (here specifically referring to V_A depending on

growth environment) being important for two traits: chlorophyll content and LMA (Table 1).

Q4: What is the direction and strength of selection on the traits and how does it vary with

growth temperature?

Our estimates of selection varied substantially among the traits. We found evidence for both

linear (β; directional) and non-linear (γ; stabilising or disruptive) selection gradients for

flowering onset, biomass, and LMA (Table 4; Fig. 2), but no clear evidence of selection on

any other phenotypic trait (Table 4).

Relative fitness was higher in individuals with earlier flowering onset: individuals

flowering later had generally very low fitness (Fig. 2A). The flowering onset of warm-grown

plants had a stronger signal of selection than cool-grown plants (i.e., a more negative linear

selection coefficient and a large quadratic coefficient; Table 4; Fig. 2B,C), where relative

fitness was lower in warm-grown plants that had intermediate to high values of flowering

onset (Fig. 2C; see Supplementary Tables S3–S6 for full models including the interaction

between selection and temperature). Relative fitness was lowest in low biomass individuals,

but there was a strong positive linear selection coefficient (increased fitness as biomass

increased) for all plants combined and under both growing temperatures separately (Table 4;

Fig. 2D,E).

Selection patterns differed between cool-grown and warm-grown plants

(Supplementary Tables S3 and S5). Although there were no significant non-linear selection

patterns in cool-grown plants (Table 4), we present the predicted non-linear fits for direct

comparison to the warm-grown plants (Fig. 2). In warm-grown plants, there was a

significantly negative quadratic selection coefficient for biomass, where the relationship

between relative fitness and biomass tapered off at very high values of biomass (Fig. 2F).

LMA was under stabilising selection for all plants combined, such that plants with

intermediate values of LMA had higher relative fitness (Table 4; Fig. 2G-I). Cool-grown

plants had a relatively small positive linear selection coefficient and a non-significant

negative quadratic coefficient (Fig. 2H), whereas warm-grown plants were not under linear

selection but showed a stabilising selection response that favoured intermediate LMA values

(Fig. 2I).

Q5: Are there effects of inbreeding on the suite of traits, and does any inbreeding depression

vary with temperature?

We predicted that plants that were the result of outcrossing as opposed to self-pollination

would have higher fitness due to inbreeding depression in the latter. Self-pollination had a

significant negative effect on reproductive fitness (Table 1), such that these plants produced a

mean of 7.7 fewer reproductive stems than plants that were outcrossed (19.3% reduction;

Fig. 3A). Self-pollination delayed mean flowering onset by 2.1 days (5.1% reduction;

Fig. 3B) and reduced mean biomass by 1.35 g (13.2% reduction; Fig. 3C). None of the other

leaf traits, photosystem or thermal tolerance traits were affected significantly by self-

pollination, and the effects of inbreeding depression were consistent across both growth

temperatures (Supplementary Figs S7 and S8, Tables 1 and 2).

Discussion

In this study we tested the effects of growth temperature, parental temperature, and

inbreeding on the multivariate phenotypes of an alpine plant with a mixed-mating system.

We found strong phenotypic plasticity for most traits, with even 10°C warmer average

growth temperatures having largely positive effects on fitness. There was substantial among-

family variation in trait values in each environment, as well as in the direction and magnitude

of reaction norms. Coupled with strong selection gradients and heritability of flowering

onset, biomass, and chlorophyll content, we have evidence for plasticity in response to

climate warming as well as evolutionary responses in *W. ceracea*, with limited indication that

plasticity itself is adaptive.

Growth temperature induces plastic responses in all traits except thermal tolerance

Phenotypic plasticity is a nearly ubiquitous response to warming conditions for functional traits that are limited by thermally-dependent reaction rates, or for traits that respond to abiotic cues associated with seasonal changes, such as photoperiod and temperature (Stotz *et al.*, 2021). While prolonged or chronic warming certainly can be limiting for plants (Nievola *et al.*, 2017; Lippmann *et al.*, 2019), alpine plant growth and reproduction is typically restricted to a relatively short growing season that follows the release from cold temperature constraints (Körner, 2003; Dolezal *et al.*, 2020). It is therefore reasonable to propose that in the non-limiting, well-watered, and warmer growing conditions, *W. ceracea* was stimulated to both grow and reproduce more than the cool-grown plants. This supports our hypothesised responses, except for that of heat tolerance. Warm-grown plants that had higher chlorophyll content and LMA, coupled with higher photosynthetic efficiency traits clearly allowed higher biomass production than cool-grown plants. Then, faster development and growth under warm conditions permitted earlier flowering onset that also increased the length of the reproductive period while allowing greater investment in reproduction. Our results indicate that these plastic responses are inducing an adaptive shift in the direction of higher fitness (Radchuk *et al.*, 2019), and our results are consistent with empirical field research that finds warming in colder climate plant species stimulates growth and reproduction (Dolezal *et al.*, 2020). For example, temperature enhancement using open top chambers in the field in Germany stimulated both growth and reproduction significantly during the growing season in alpine grassland species, although herbaceous perennials were less responsive than graminoids or shrubs (Kudernatsch *et al.*, 2008).

The capacity for plants to increase their physiological thermal tolerance has been predicted to be a key response to climate warming (Geange *et al.*, 2021). For example, $T_{\text{crit-hot}}$ is well known to increase rapidly by 4°C or more within hours to days during an acute heat stress event (e.g., Andrew *et al.*, 2023; Zhu *et al.*, 2024). Long-term exposure to warm growth temperatures can also increase $T_{\text{crit-hot}}$, for example by ~0.16°C per 1°C of growth temperature (Zhu *et al.*, 2018). We have previously observed plasticity in heat tolerance ($T_{\text{crit-hot}}$) in the F1 and F2 generations of our *W. ceracea* experiments to long-term warming of 28-30°C. In those experiments, warm-grown plants increased their critical heat tolerance limits by 0.7-3.2°C relative to cool growing conditions (Notarnicola *et al.*, 2021; Arnold *et al.*, 2022), while also having a range of important effects on phenotypic and reproductive traits. It is worth noting that both these previous studies constrained pot sizes and used different, more

confined controlled growth environments, which limits the value of making comparisons to

our current glasshouse study (see Karitter *et al.*, 2023 for a discussion of differences in

phenotypic expression among common-environment experiments). We have also shown that

moderate warming to 30°C can result in upregulation of genes related to post-transcriptional

processes and downregulation of genes related to photosynthesis-related processes in

W. ceracea (Notarnicola *et al.*, 2023a). These suites of genetic changes that differ between

cool to warm growing conditions may not correspond directly to changes in the phenotypic

trait $T_{crit-hot}$, which increased by a mean of only ~0.26°C in the current study. Therefore, this

canalisation of $T_{crit-hot}$ to warming that we observed suggests that the warm-grown plants

(while being well-watered) were not severely stressed, despite 30°C being far warmer than

typical growing conditions for alpine *W. ceracea*. The warm treatment may have also

alleviated temperature limitation on growth processes that can occur in alpine environments,

which aligns with other trait responses that we observed. However, we encourage future

studies to also embark on detailed examinations of thermal tolerance responses using thermal

performance curves to determine pivotal temperature thresholds, as well as plasticity and

selection thereof (Wooliver *et al.*, 2022).

Strong non-linear selection on heritable traits, especially under warm conditions

Warming has the potential to drive rapid evolutionary change in plant phenotypes, provided

that phenotypic traits have a relationship with fitness and are therefore under selection, and

that variation in the traits is heritable (Anderson and Song, 2020; Scheiner *et al.*, 2020). Here

our hypothesis that flowering onset, biomass, LMA, and heat tolerance would be heritable

and under selection was partially supported. We found that flowering onset, biomass, and

chlorophyll content were clearly heritable in *W. ceracea*. Chlorophyll content is proportional

to the concentration of photosynthetic pigments and nitrogen in a leaf (Ling *et al.*, 2011); it is

heritable in wheat (Rosyara *et al.*, 2010; Said *et al.*, 2022), and has a relatively strong

relationship with flowering in other crop species (Senger *et al.*, 2014). To the best of our

knowledge, there has not been another study reporting the heritability of chlorophyll content

in a wild species, but our findings here suggest that variation in photosynthetic pigment

concentration at least has a genetic basis. The low heritability of total reproductive stems

(i.e., our best measure of fitness) may reflect depletion of genetic variance for fitness as

expected from evolutionary theory (Falconer and Mackay, 1996; Kruuk *et al.*, 2000), as well

as large other sources of variance.

Here we found that both flowering onset and biomass were heritable as well as under

relatively strong selection. There was significant negative directional selection (β) on

flowering onset in all cases, where earlier flowering is favoured, and relative fitness in warm-

grown plants also declined non-linearly (non-zero γ) with later flowering. These observed

patterns of selection on flowering phenology align exactly with the findings from the relative

cool and warm ends of the spectrum from a natural geothermal heating experiment on

selection on flowering phenology in the short-lived perennial herb *Cerastium fontanum* in

Iceland (Valdés *et al.*, 2019). In our study, biomass was under strong positive directional

selection across all environments, where larger plants had higher relative fitness, although

fitness tapered off for larger warm-grown plants. Biomass can be a reasonable proxy for

fitness (Younginger *et al.*, 2017), where selection can favour larger individual size to

facilitate plant performance (Aspi *et al.*, 2003). Nevertheless, the plateau in relative fitness at

larger sizes in warm-grown plants may be because the largest individuals would have

relatively higher water demands and stronger resource allocation trade-offs than their smaller

conspecifics under warming. Glasshouse studies can exacerbate effects of water limitation,

and indeed we observed anecdotally that at their peak size, larger individuals in the warm-

grown treatment began to wilt toward the end of hot, sunny days during the experiment.

Heritability of biomass in these F3 plants aligns with previous findings of high V_A in both

early growth rate and biomass in F2 *W. ceracea* plants (Arnold *et al.*, 2022).

LMA was under positive directional selection overall and in cool-grown plants, but

under stabilising selection in warm-grown plants, with intermediate to high but not extreme

LMA values being favoured. LMA is an estimate of the density of carbon and nutrients in a

set area of leaf tissue (i.e., the cost of tissue production for light interception) and is part of a

trait complex that determines photosynthetic capacity, and nitrogen and water use efficiency

(Poorter *et al.*, 2009; Funk *et al.*, 2021). Investment in high LMA may improve resource gain

but only without critical water deficit (Ivanova *et al.*, 2018), and potentially at the cost of

reinvesting the acquired resources into vegetative rather than reproductive tissues (Flores *et*

al., 2014). Thus, within a species, relatively low and high LMA values represent inefficient

resource acquisition-use strategies that trade-off with reproduction, hence extreme values of

LMA are selected against (Flores *et al.*, 2014). Despite previously finding high intraspecific

variation in $T_{crit-hot}$ (Arnold *et al.*, 2022), here it was neither heritable nor under selection,

perhaps because photosynthetic heat tolerance does not affect fitness directly.

Taken together, our results suggest that traits contributing to light interception,

growth, biomass, and flowering phenology are the key traits for ecological and evolutionary

responses in plants to temperature. The capacity for these plants to tolerate and respond to

warmer-than-typical temperatures demonstrates tolerance to moderate warming through a

combination of phenotypic plasticity and intrinsic resilience. Growth, size, and reproductive

traits respond to temperature over longer timescales (weeks to months) and contribute

directly to fitness, whereas physiological traits regulate essential functions on shorter

timescales (hours to days), but do not contribute directly to fitness. We highlight the need for

future studies to take a demographic approach to studying plant responses to environmental

stressors, integrating across the life cycle. Early life stages that are critical for establishment,

growth, and survival, and reproductive stages that may be sensitive to temperature extremes

and which directly affect fitness are typically less often studied than seeds or young adult

plants. Finally, considering that warm-grown plants have different (non-linear) patterns of

selection to cool-grown plants, future studies should concentrate on investigating novel or

edge conditions to determine tipping points or sensitivity for ecological and evolutionary

responses.

No evidence of transgenerational plasticity via parental temperature or maternal effects

By applying a reciprocally matched-mismatched parent-offspring environments design,

combined with the breeding structure, we could test for transgenerational plasticity via the

parent environment effects and maternal effects through a pedigree (Uller *et al.*, 2013).

Evidence for matching parent-offspring environments benefitting offspring is relatively weak

overall in plants (Uller *et al.*, 2013). Based on earlier results with *W. ceracea* (Wang *et al.*,

2021; Notarnicola *et al.*, 2023b), we hypothesised that there could be a small benefit for

offspring performance and fitness when matching their parent environment. However, we

found no convincing evidence for any form of transgenerational plasticity, beneficial or not.

Using the same breeding design for F2 and F3 families as in the current experiment,

Wang *et al.* (2021) tested for transgenerational plasticity in early life traits in *W. ceracea*.

Seeds from parent plants grown in warm conditions had delayed germination (extended

dormancy) and reduced germination success irrespective of their germination temperature,

but none of these effects persisted to affect seedling growth (Wang *et al.*, 2021). In a

comprehensive reciprocal transplant experiment with *Boechera stricta* across an elevation

gradient, Wadgymar *et al.* (2018a) found transgenerational plasticity in the early life traits of

seed viability, germination, and dormancy. Transgenerational plasticity interplayed with

within-generation plasticity across elevations and the effects of both were complex and

context-specific, but parental environmental effects largely did not persist to later life

(Wadgymar *et al.*, 2018a). However, in a reciprocal environment experiment on *Lupinus*

angustifolius under well-watered and drought stress treatments, Matesanz *et al.* (2022) found

significant transgenerational plasticity that affected functional traits and reproduction of the

offspring. Parental effects altered individual seed mass, flowering onset, and growth rate of

the offspring, but these effects were not always beneficial, and offspring environment effects

far outweighed the parental effects for specific leaf area, F_V/F_M, and lifetime reproductive

output (Matesanz *et al.*, 2022). Seed provisioning through maternal resource allocation

affects seed viability, which in turn affects the probability of germination success (Haig and

Westoby, 1988). Unsuitable, limiting, or stressful parental environments can also have direct

adverse effects on reproductive tissues. For example, exposure to high temperature can

disrupt reproductive development and reduce pollen viability, leading to smaller and/or less

viable seeds, which have altered germination responses (Herman and Sultan, 2011; Sehgal *et*

al., 2018; Goel *et al.*, 2023).These examples highlight that transgenerational plasticity can

certainly affect seed traits, but that persistent effects into adult phenotypes of the offspring

might be less common or weaker (Herman and Sultan, 2011; Wang *et al.*, 2021; Notarnicola

et al., 2023b), although there are notable exceptions (e.g., Whittle *et al.*, 2009).

Inbreeding impairs reproduction and biomass but not physiological function

We predicted that inbreeding would significantly reduce plant performance and fitness

compared to crossing, due to inbreeding depression. *Wahlenbergia ceracea* is protandrous

and facultatively autogamous with a mixed-mating system (Nicotra *et al.*, 2015), where self-

pollination of the same flower is delayed by several days following flower opening, which

provides reproductive assurance in the absence of external pollination (Goodwillie *et al.*,

2005). Many alpine species with a mixed-mating strategy depend on external pollination to

achieve their maximum potential seed set (Scheffknecht *et al.*, 2007). It is therefore

unsurprising that we found that self-pollination in *W. ceracea* caused marginally delayed

flowering and slightly reduced total reproductive stems and biomass. The magnitude of the

inbreeding depression effect on these fitness-related traits is comparable to the expected

range from a meta-analysis of inbreeding effects on plant fitness (Angeloni *et al.*, 2011).

Inbreeding effects can be exacerbated in stressful environments (Armbruster and Reed,

2005), however we did not observe this effect, nor did we find any inbreeding effect on

functional traits. We suggest that inbreeding (particularly in mixed-mating species) might

affect fitness directly rather than indirectly through traits that mediate resource acquisition.

However, we do not yet know whether negative effects of inbreeding on function and fitness

would be exacerbated under more challenging conditions (e.g., heat coupled with drought) or

extreme events (e.g., heatwaves).

Conclusions and future directions

The capacity for plants to alter their phenotype in response to climate warming is frequently

thought to be adaptive. Here we show through comprehensive analyses that in this alpine

species, warming may alleviate restrictions on growth and reproduction, thereby improving

fitness under warming through plasticity. The exception to this conclusion was thermal

tolerance, which is likely already at sufficient levels. Only flowering onset and biomass were

both heritable and clearly under selection, which differed between the cool and warm growth

environments, and only chlorophyll content and LMA had any evidence for G×E. The effect

of growth environment far exceeded any influence of parental environment; we found little

evidence for substantial maternal effects or transgenerational plasticity in adult traits. Further,

the effect of inbreeding by self-pollination was relatively small, providing reproductive

assurance at low cost. We can conclude that the mixed-mating alpine herb *W. ceracea* clearly

has capacity to respond rapidly to climate warming via phenotypic plasticity as well as the

potential for evolutionary change across generations.

As the climate of alpine ecosystems changes, the duration of the growing season will

extend, generating both new opportunities and new challenges for its inhabitants. Our

experiment found that substantially warmer daytime temperatures (30°C) can still facilitate

growth and reproduction in an alpine herb when water is not limiting. However, climate

change is also expected to progressively dry some alpine ecosystems. Climate projections for

some areas with seasonal snowpack have forecast reduced winter snowfall, earlier snowmelt

in spring, and potentially decreases in summer and autumn precipitation events (Gobiet *et al.*,

2014; Harris *et al.*, 2016). Reductions in water supply have a clearly detrimental effect on

most alpine plants (Sumner and Venn, 2021), and interactions between warming and water

limitation are undoubtedly relevant for future climate scenarios in alpine plant communities

(De Boeck *et al.*, 2016; Winkler *et al.*, 2016). Therefore, an essential next step in building an

understanding of the importance of eco-evolutionary responses to climate change will be to

test the role of water limitation in altering plastic and evolutionary responses to temperature.

Heat stress events are predicted to become more frequent, intense, and longer duration

(Trancoso *et al.*, 2020), on top of a background of mean climate warming (Harris *et al.*,

2018). Extreme events have the potential to change fitness drastically and could be stronger

selective events than gradual environmental change, which will alter evolutionary dynamics

of populations in future (Gutschick and BassiriRad, 2003). The role of extreme events in the

eco-evolutionary dynamics of alpine plants remains largely unexplored, despite alpine

ecosystems being among the most vulnerable to and already impacted by climate change

(Verrall and Pickering, 2020). Using genomic approaches to study climate change responses

in natural populations could reveal the genomic architecture of traits exhibiting plasticity and

under selection, improving our understanding of the mechanisms behind stress responses and

their evolutionary potential (Notarnicola *et al.*, 2023a). Employing a multifaceted research

effort to strengthen our understanding of the roles of plasticity and evolutionary responses to

realistic climate scenarios and extreme events is necessary to evaluate the potential for alpine

plants to respond to future conditions.

Supplementary data

Supplementary data

Table S1: Summary statistics for all phenotypic traits.

Table S2: Linear regression model output testing fitness index suitability.

Table S3: Model outputs for linear selection by growth treatment 1.

Table S4: Model outputs for linear selection by growth treatment 2.

Table S5: Model outputs for quadratic selection by growth treatment 1.

Table S6: Model outputs for quadratic selection by growth treatment 2.

Fig. S1: Summary of the multi-generational breeding design.

Fig. S2: Correlation between total reproductive stems and reproductive mass.

Fig. S3: Reaction norms of phenotypic traits.

Fig. S4: Isolated parental temperature effects for phenotypic traits.

Fig. S5: Matched/mismatched parent-offspring environment effects on traits 1.

Fig. S6: Matched/mismatched parent-offspring environment effects on traits 2.

Fig. S7: Tests for inbreeding depression in phenotypic traits 1.

Fig. S8: Tests for inbreeding depression in phenotypic traits 2.

Acknowledgements

We respectfully acknowledge the traditional custodians of the land from which plant seeds

were originally collected and the land on which this research was conducted: the Ngunnawal,

Ngambri, Ngarigo, and Walgalu people. We sincerely thank The Australian National

University Plant Services staff: Steve Dempsey, Christine Larsen, Darren Marsh, Gavin

Pritchard, and Jenny Rath for their extraordinary contributions to maintaining the glasshouse

plants, and with hailstorm recovery efforts. We thank Richard Poiré for providing a second

Imaging-PAM system and for offering space in the Growth Capsules after the hailstorm. The

phenotyping and harvesting of the plants, particularly in the aftermath of the hailstorm, would

not have been possible without the efforts of a substantial team of volunteers and colleagues,

for whom we are very grateful: Monica Arnold, Verónica Briceño, Zachary Brown,

Stephanie Courtney Jones, Jemimah Hamilton, Ophélie Lasne, Patrick Meir, Tenzin Norzin,

Helen Osmond, Abigail Ryan, Natasha Salisbury, Alice Schacher, and Kaitlyn Spooner.

Further thanks to Alexandra Catling, Kelli Gowland, Pamudika Kiridena, and Melissa

Wakem for their contributions to generating the F3 lines.

Author contributions

PAA – conceptualisation, experimental design, performed the experiments, analysed the data, wrote the first draft.

SW – performed the experiments, editing.

RFN – performed the experiments, editing.

ABN – conceptualisation, experimental design, funding acquisition, supervision, editing.

LEBK – conceptualisation, experimental design, funding acquisition, supervision, editing.

Funding

Australian Research Council Discovery Projects DP170101681 and DP200101382.

Conflict of interest

The authors declare no conflicts of interest.

Data availability

Data and R code will be available in the Dryad repository doi:10.5061/dryad.34tmpg4s9.

References

Anderson JT. 2016. Plant fitness in a rapidly changing world. New Phytologist **210**, 81-87.

Anderson JT, Inouye DW, McKinney AM, Colautti RI, Mitchell-Olds T. 2012. Phenotypic plasticity and adaptive evolution contribute to advancing flowering phenology in response to climate change. Proceedings of the Royal Society B: Biological Sciences **279**, 3843-3852.

Anderson JT, Song B-H. 2020. Plant adaptation to climate change — Where are we? Journal of Systematics and Evolution **58**, 533-545.

Andrew SC, Arnold PA, Simonsen AK, Briceño VF. 2023. Consistently high heat tolerance acclimation in response to a simulated heatwave across species from the broadly distributed *Acacia* genus. Functional Plant Biology **50**, 71-83.

Angeloni F, Ouborg NJ, Leimu R. 2011. Meta-analysis on the association of population size and life history with inbreeding depression in plants. Biological Conservation **144**, 35-43.

Armbruster P, Reed DH. 2005. Inbreeding depression in benign and stressful environments. Heredity **95**, 235-242.

Arnold PA, Briceño VF, Gowland KM, Catling AA, Bravo LA, Nicotra AB. 2021. A high-throughput method for measuring critical thermal limits of leaves by chlorophyll imaging fluorescence. Functional Plant Biology **48**, 634-646.

Arnold PA, Kruuk LEB, Nicotra AB. 2019a. How to analyse plant phenotypic plasticity in response to a changing climate. New Phytologist **222**, 1235-1241.

Arnold PA, Nicotra AB, Kruuk LEB. 2019b. Sparse evidence for selection on phenotypic plasticity in response to temperature. Philosophical Transactions of the Royal Society B: Biological Sciences **374**, 20180185.

Arnold PA, Wang S, Catling AA, Kruuk LEB, Nicotra AB. 2022. Patterns of phenotypic plasticity along a thermal gradient differ by trait type in an alpine plant. Functional Ecology **36**, 2412-2428.

Aspi J, Jäkäläniemi A, Tuomi J, Siikamäki P. 2003. Multilevel phenotypic selection on morphological characters in a metapopulation of *Silene tatarica*. Evolution **57**, 509-517.

Auld JR, Agrawal AA, Relyea RA. 2010. Re-evaluating the costs and limits of adaptive phenotypic plasticity. Proceedings of the Royal Society B: Biological Sciences **277**, 503-511.

Baker NR. 2008. Chlorophyll fluorescence: a probe of photosynthesis in vivo. Annual Review of Plant Biology **59**, 89-113.

Bonduriansky R. 2021. Plasticity across generations. In: Pfennig DW, ed. *Phenotypic Plasticity & Evolution*. Boca Raton, FL, USA: CRC Press, 327-348.

Bonser SP. 2021. Misinterpreting the adaptive value of phenotypic plasticity in studies on plant adaptation to new and variable environments. Plant Biology **23**, 683-685.

Bradshaw AD. 1965. Evolutionary significance of phenotypic plasticity in plants. Advances in Genetics **13**, 115-155.

Bradshaw AD. 2006. Unravelling phenotypic plasticity – why should we bother? New Phytologist **170**, 644-648.

Brooker R, Brown LK, George TS, Pakeman RJ, Palmer S, Ramsay L, Schöb C, Schurch N, Wilkinson MJ. 2022. Active and adaptive plasticity in a changing climate. Trends in Plant Science **27**, 717-728.

Bürkner P-C. 2018. Advanced Bayesian multilevel modeling with the R package brms. The R Journal **10**, 395-411.

De Boeck HJ, Bassin S, Verlinden M, Zeiter M, Hiltbrunner E. 2016. Simulated heat waves affected alpine grassland only in combination with drought. New Phytologist **209**, 531-541.

De Kort H, Panis B, Helsen K, Douzet R, Janssens SB, Honnay O. 2020. Pre☐adaptation to climate change through topography☐driven phenotypic plasticity. Journal of Ecology **108**, 1465-1474.

Dolezal J, Kurnotova M, Stastna P, Klimesova J. 2020. Alpine plant growth and reproduction dynamics in a warmer world. New Phytologist **228**, 1295-1305.

Dorji T, Hopping KA, Meng F, Wang S, Jiang L, Klein JA. 2020. Impacts of climate change on flowering phenology and production in alpine plants: The importance of end of flowering. Agriculture, Ecosystems & Environment **291**, 106795.

Edelaar P, Jovani R, Gomez-Mestre I. 2017. Should I change or should I go? Phenotypic plasticity and matching habitat choice in the adaptation to environmental heterogeneity. The American Naturalist **190**, 506-520.

Ehrlén J, Valdés A. 2020. Climate drives among-year variation in natural selection on flowering time. Ecology Letters **23**, 653-662.

Engqvist L, Reinhold K. 2016. Adaptive trans-generational phenotypic plasticity and the lack of an experimental control in reciprocal match/mismatch experiments. Methods in Ecology and Evolution **7**, 1482-1488.

Falconer DS, Mackay TFC. 1996. *Introduction to quantitative genetics*. Harlow, UK: Longmans Green.

Flores O, Garnier E, Wright IJ, Reich PB, Pierce S, Dìaz S, Pakeman RJ, Rusch GM, Bernard-Verdier M, Testi B, Bakker JP, Bekker RM, Cerabolini BEL, Ceriani RM, Cornu G, Cruz P, Delcamp M, Dolezal J, Eriksson O, Fayolle A, Freitas H, Golodets C, Gourlet-Fleury S, Hodgson JG, Brusa G, Kleyer M, Kunzmann D, Lavorel S, Papanastasis VP, Pérez-Harguindeguy N, Vendramini F, Weiher E. 2014. An evolutionary perspective on leaf economics: phylogenetics of leaf mass per area in vascular plants. Ecology and Evolution **4**, 2799-2811.

Fox RJ, Donelson JM, Schunter C, Ravasi T, Gaitán-Espitia JD. 2019. Beyond buying time: the role of plasticity in phenotypic adaptation to rapid environmental change. Philosophical Transactions of the Royal Society B: Biological Sciences **374**, 20180174.

Franks SJ, Sim S, Weis AE. 2007. Rapid evolution of flowering time by an annual plant in response to a climate fluctuation. Proceedings of the National Academy of Sciences **104**, 1278.

Funk JL, Larson JE, Vose G. 2021. Leaf traits and performance vary with plant age and water availability in *Artemisia californica*. Annals of Botany **127**, 495-503.

Gabry J, Simpson D, Vehtari A, Betancourt M, Gelman A. 2019. Visualization in Bayesian workflow. Journal of the Royal Statistical Society Series A: Statistics in Society **182**, 389-402.

Geange SR, Arnold PA, Catling AA, Coast O, Cook AM, Gowland KM, Leigh A, Notarnicola RF, Posch BC, Venn SE, Zhu L, Nicotra AB. 2021. The thermal tolerance of photosynthetic tissues: a global systematic review and agenda for future research. New Phytologist **229**, 2497-2513.

Geber MA, Griffen LR. 2003. Inheritance and natural selection on functional traits. International Journal of Plant Sciences **164**, S21-S42.

Giménez-Benavides L, García-Camacho R, Iriondo JM, Escudero A. 2011. Selection on flowering time in Mediterranean high-mountain plants under global warming. Evolutionary Ecology **25**, 777-794.

Gobiet A, Kotlarski S, Beniston M, Heinrich G, Rajczak J, Stoffel M. 2014. 21st century climate change in the European Alps—A review. Science of The Total Environment **493**, 1138-1151.

Goel K, Kundu P, Sharma P, Zinta G. 2023. Thermosensitivity of pollen: a molecular perspective. Plant Cell Reports **42**, 843-857.

Goodwillie C, Kalisz S, Eckert CG. 2005. The evolutionary enigma of mixed mating systems in plants: Occurrence, theoretical explanations, and empirical evidence. Annual Review of Ecology, Evolution, and Systematics **36**, 47-79.

Gutschick VP, BassiriRad H. 2003. Extreme events as shaping physiology, ecology, and evolution of plants: toward a unified definition and evaluation of their consequences. New Phytologist **160**, 21-42.

Hadfield JD. 2010. MCMC methods for multi-response generalized linear mixed models: the MCMCglmm R package. Journal of Statistical Software **33**, 1-22.

Haig D, Westoby M. 1988. Inclusive fitness, seed resources, and maternal care. In: Lovett Doust J, Lovett Doust L, eds. *Plant Reproductive Ecology*. Oxford: Oxford University Press, 60-79.

Hamann E, Denney D, Day S, Lombardi E, Jameel MI, MacTavish R, Anderson JT. 2021. Review: Plant eco-evolutionary responses to climate change: Emerging directions. Plant Science **304**, 110737.

Harris RMB, Beaumont LJ, Vance TR, Tozer CR, Remenyi TA, Perkins-Kirkpatrick SE, Mitchell PJ, Nicotra AB, McGregor S, Andrew NR, Letnic M, Kearney MR, Wernberg T, Hutley LB, Chambers LE, Fletcher MS, Keatley MR, Woodward CA, Williamson G, Duke NC, Bowman DMJS. 2018. Biological responses to the press and pulse of climate trends and extreme events. Nature Climate Change **8**, 579-587.

Harris RMB, Remenyi T, Bindoff NL. 2016. The potential impacts of climate change on Victorian alpine resorts. A report to the alpine resorts co-ordinating council. Antarctic Climate and Ecosystems Cooperative Research Centre, Hobart, Australia, 1-203.

Hegland SJ, Nielsen A, Lázaro A, Bjerknes A-L, Totland Ø. 2009. How does climate warming affect plant-pollinator interactions? Ecology Letters **12**, 184-195.

Hendry AP. 2016. Key questions on the role of phenotypic plasticity in eco-evolutionary dynamics. Journal of Heredity **107**, 25-41.

Herman J, Sultan S. 2011. Adaptive transgenerational plasticity in plants: case studies, mechanisms, and implications for natural populations. Frontiers in Plant Science **2**, 102.

Hilker M, Schwachtje J, Baier M, Balazadeh S, Bäurle I, Geiselhardt S, Hincha DK, Kunze R, Mueller-Roeber B, Rillig MC, Rolff J, Romeis T, Schmülling T, Steppuhn A, van Dongen J, Whitcomb SJ, Wurst S, Zuther E, Kopka J. 2016. Priming and memory of stress responses in organisms lacking a nervous system. Biological Reviews **91**, 1118-1133.

Ivanova LA, Zolotareva NV, Ronzhina DA, Podgaevskaya EN, Migalina SV, Ivanov LA. 2018. Leaf functional traits of abundant species predict productivity in three temperate herbaceous communities along an environmental gradient. Flora **239**, 11-19.

Josephs EB. 2018. Determining the evolutionary forces shaping $G \times E$. New Phytologist **219**, 31-36.

Karitter P, March-Salas M, Ensslin A, Rauschkolb R, Godefroid S, Poorter H, Scheepens JF. 2023. Garden, greenhouse or climate chamber? Experimental conditions

influence whether genetic differences are phenotypically expressed. bioRxiv, 2023.2012.2006.570376.

Kimball S, Gremer JR, Angert AL, Huxman TE, Venable DL. 2012. Fitness and physiology in a variable environment. Oecologia **169**, 319-329.

Körner C. 2003. *Alpine plant life*. New York, USA: Springer.

Kruuk LEB. 2004. Estimating genetic parameters in natural populations using the 'animal model'. Philosophical Transactions of the Royal Society of London B: Biological Sciences **359**, 873-890.

Kruuk LEB, Clutton-Brock TH, Slate J, Pemberton JM, Brotherstone S, Guinness FE. 2000. Heritability of fitness in a wild mammal population. Proceedings of the National Academy of Sciences **97**, 698-703.

Kudernatsch T, Fischer A, Bernhardt-Römermann M, Abs C. 2008. Short-term effects of temperature enhancement on growth and reproduction of alpine grassland species. Basic and Applied Ecology **9**, 263-274.

Lambrecht SC, Mahieu S, Cheptou P-O. 2016. Natural selection on plant physiological traits in an urban environment. Acta Oecologica **77**, 67-74.

Lande R, Arnold SJ. 1983. The measurement of selection on correlated characters. Evolution **37**, 1210-1226.

Ling Q, Huang W, Jarvis P. 2011. Use of a SPAD-502 meter to measure leaf chlorophyll concentration in *Arabidopsis thaliana*. Photosynthesis Research **107**, 209-214.

Lippmann R, Babben S, Menger A, Delker C, Quint M. 2019. Development of wild and cultivated plants under global warming conditions. Current Biology **29**, R1326-R1338.

Lüdecke D, Ben-Shachar MS, Patil I, Waggoner P, Makowski D. 2021. performance: An R Package for Assessment, Comparison and Testing of Statistical Models. Journal of Open Source Software **6**, 3139.

Matesanz S, Ramos-Muñoz M, Rubio Teso ML, Iriondo JM. 2022. Effects of parental drought on offspring fitness vary among populations of a crop wild relative. Proceedings of the Royal Society B: Biological Sciences **289**, 20220065.

Mousseau TA, Fox CW. 1998. The adaptive significance of maternal effects. Trends in Ecology & Evolution **13**, 403-407.

Muggeo VMR. 2017. Interval estimation for the breakpoint in segmented regression: a smoothed score-based approach. Australian & New Zealand Journal of Statistics **59**, 311-322.

Nakagawa S, Schielzeth H. 2013. A general and simple method for obtaining R^2 from generalized linear mixed-effects models. Methods in Ecology and Evolution **4**, 133-142.

Nicotra AB, Atkin OK, Bonser SP, Davidson AM, Finnegan EJ, Mathesius U, Poot P, Purugganan MD, Richards CL, Valladares F, van Kleunen M. 2010. Plant phenotypic plasticity in a changing climate. Trends in Plant Science **15**, 684-692.

Nicotra AB, Segal DL, Hoyle GL, Schrey AW, Verhoeven KJF, Richards CL. 2015. Adaptive plasticity and epigenetic variation in response to warming in an Alpine plant. Ecology and Evolution **5**, 634-647.

Nievola CC, Carvalho CP, Carvalho V, Rodrigues E. 2017. Rapid responses of plants to temperature changes. Temperature **4**, 371-405.

Noble DWA, Wechmann K, Keogh JS, Whiting MJ. 2013. Behavioral and morphological traits interact to promote the evolution of alternative reproductive tactics in a lizard. The American Naturalist **182**, 726-742.

Notarnicola RF, Arnold PA, Feng Z-P, Hamilton J, Jones A, W. , Loke S, Nicotra AB, Schwessinger B, Ganguly DR. 2023a. Transcriptional acclimation to warming temperatures of the Australian alpine herb *Wahlenbergia ceracea*. bioRxiv **2023.09.27.559694**.

Notarnicola RF, Kruuk LEB, Nicotra AB, Arnold PA. 2021. Tolerance of warmer temperatures does not confer resilience to heatwaves in an alpine herb. Frontiers in Ecology and Evolution **9**, 615119.

Notarnicola RF, Nicotra AB, Kruuk LEB, Arnold PA. 2023b. Effects of warming temperatures on germination responses and trade-offs between seed traits in an alpine plant. Journal of Ecology **111**, 62-76.

Parmesan C, Hanley ME. 2015. Plants and climate change: complexities and surprises. Annals of Botany **116**, 849-864.

Parmesan C, Yohe G. 2003. A globally coherent fingerprint of climate change impacts across natural systems. Nature **421**, 37-42.

Poorter H, Niinemets Ü, Poorter L, Wright Ian J, Villar R. 2009. Causes and consequences of variation in leaf mass per area (LMA): a meta‐analysis. New Phytologist **182**, 565-588.

R Core Team. 2023. R: A language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing.

Radchuk V, Reed T, Teplitsky C, van de Pol M, Charmantier A, Hassall C, Adamík P, Adriaensen F, Ahola MP, Arcese P, Miguel Avilés J, Balbontin J, Berg KS, Borras A, Burthe S, Clobert J, Dehnhard N, de Lope F, Dhondt AA, Dingemanse NJ, Doi H, Eeva T, Fickel J, Filella I, Fossøy F, Goodenough AE, Hall SJG, Hansson B, Harris M, Hasselquist D, Hickler T, Joshi J, Kharouba H, Martínez JG, Mihoub J-B, Mills JA, Molina-Morales M, Moksnes A, Ozgul A, Parejo D, Pilard P, Poisbleau M, Rousset F, Rödel M-O, Scott D, Senar JC, Stefanescu C, Stokke BG, Kusano T, Tarka M, Tarwater CE, Thonicke K, Thorley J, Wilting A, Tryjanowski P, Merilä J, Sheldon BC, Pape Møller A, Matthysen E, Janzen F, Dobson FS, Visser ME, Beissinger SR, Courtiol A, Kramer-Schadt S. 2019. Adaptive responses of animals to climate change are most likely insufficient. Nature Communications **10**, 3109.

Rickards L, Watson JEM. 2020. Research is not immune to climate change. Nature Climate Change **10**, 180-183.

Rosyara UR, Subedi S, Duveiller E, Sharma RC. 2010. Photochemical efficiency and SPAD value as indirect selection criteria for combined selection of spot blotch and terminal heat stress in wheat. Journal of Phytopathology **158**, 813-821.

Said AA, Moursi YS, Sallam A. 2022. Association mapping and candidate genes for physiological non-destructive traits: Chlorophyll content, canopy temperature, and specific leaf area under normal and saline conditions in wheat. Frontiers in Genetics **13**, 980319.

Scheffknecht S, Dullinger S, Grabherr G, Hülber K. 2007. Mating systems of snowbed plant species of the northeastern Calcareous Alps of Austria. Acta Oecologica **31**, 203-209.

Scheiner SM, Barfield M, Holt RD. 2020. The genetics of phenotypic plasticity. XVII. Response to climate change. Evolutionary Applications **13**, 388-399.

Sehgal A, Sita K, Siddique KHM, Kumar R, Bhogireddy S, Varshney RK, HanumanthaRao B, Nair RM, Prasad PVV, Nayyar H. 2018. Drought or/and heat-stress effects on seed filling in food crops: Impacts on functional biochemistry, seed yields, and nutritional quality. Frontiers in Plant Science **9**, 1705.

Senger E, Peyrat A, Martin M, Montes JM. 2014. Genetic variation in leaf chlorophyll content of *Jatropha curcas* L. Industrial Crops and Products **58**, 204-211.

Stinchcombe JR, Agrawal AF, Hohenlohe PA, Arnold SJ, Blows MW. 2008. Estimating nonlinear selection gradients using quadratic regression coefficients: double or nothing? Evolution **62**, 2435-2440.

Stinchcombe JR, Dorn LA, Schmitt J. 2004. Flowering time plasticity in *Arabidopsis thaliana*: a reanalysis of Westerman & Lawrence (1970). Journal of Evolutionary Biology **17**, 197-207.

Stotz GC, Salgado-Luarte C, Escobedo VM, Valladares F, Gianoli E. 2021. Global trends in phenotypic plasticity of plants. Ecology Letters **24**, 2267-2281.

Sumner E, Venn S. 2021. Plant responses to changing water supply and availability in high elevation ecosystems: A quantitative systematic review and meta-analysis. Land **10**, 1150.

Svensson EI, Gomez-Llano M, Waller JT. 2020. Selection on phenotypic plasticity favors thermal canalization. Proceedings of the National Academy of Sciences **117**, 29767.

Tattersall GJ, Sinclair BJ, Withers PC, Fields PA, Seebacher F, Cooper CE, Maloney SK. 2012. Coping with thermal challenges: physiological adaptations to environmental temperatures. Comprehensive Physiology **2**, 2151-2202.

Totland Ø. 1999. Effects of temperature on performance and phenotypic selection on plant traits in alpine *Ranunculus acris*. Oecologia **120**, 242-251.

Trancoso R, Syktus J, Toombs N, Ahrens D, Wong KK-H, Pozza RD. 2020. Heatwaves intensification in Australia: A consistent trajectory across past, present and future. Science of The Total Environment **742**, 140521.

Uller T. 2008. Developmental plasticity and the evolution of parental effects. Trends in Ecology & Evolution **23**, 432-438.

Uller T, Nakagawa S, English S. 2013. Weak evidence for anticipatory parental effects in plants and animals. Journal of Evolutionary Biology **26**, 2161-2170.

Valdés A, Marteinsdóttir B, Ehrlén J. 2019. A natural heating experiment: Phenotypic and genotypic responses of plant phenology to geothermal soil warming. Global Change Biology **25**, 954-962.

van Buskirk J, Steiner UK. 2009. The fitness costs of developmental canalization and plasticity. Journal of Evolutionary Biology **22**, 852-860.

Vehtari A, Gelman A, Gabry J. 2017. Practical Bayesian model evaluation using leave-one-out cross-validation and WAIC. Statistics and Computing **27**, 1413-1432.

Verrall B, Pickering CM. 2020. Alpine vegetation in the context of climate change: A global review of past research and future directions. Science of The Total Environment **748**, 141344.

Wadgymar SM, Mactavish RM, Anderson JT. 2018a. Transgenerational and within-generation plasticity in response to climate change: insights from a manipulative field experiment across an elevational gradient. The American Naturalist **192**, 698-714.

Wadgymar SM, Ogilvie JE, Inouye DW, Weis AE, Anderson JT. 2018b. Phenological responses to multiple environmental drivers under climate change: insights from a long‐term observational study and a manipulative field experiment. New Phytologist **218**, 517-529.

Walther G-R. 2003. Plants in a warmer world. Perspectives in Plant Ecology, Evolution and Systematics **6**, 169-185.

Walther G-R, Post E, Convey P, Menzel A, Parmesan C, Beebee TJC, Fromentin J-M, Hoegh-Guldberg O, Bairlein F. 2002. Ecological responses to recent climate change. Nature **416**, 389-395.

Wang S, Gowland KM, Kruuk LEB, Nicotra AB, Arnold PA. 2021. Decoupling the effects of parental and offspring warming on seed and seedling traits. Alpine Botany **131**, 105-115.

Whittle CA, Otto SP, Johnston MO, Krochko JE. 2009. Adaptive epigenetic memory of ancestral temperature regime in *Arabidopsis thaliana*. Botany **87**, 650-657.

Wilson AJ, Réale D, Clements MN, Morrissey MM, Postma E, Walling CA, Kruuk LEB, Nussey DH. 2010. An ecologist's guide to the animal model. Journal of Animal Ecology **79**, 13-26.

Winkler DE, Chapin KJ, Kueppers LM. 2016. Soil moisture mediates alpine life form and community productivity responses to warming. Ecology **97**, 1553-1563.

Wooliver R, Vtipilthorpe EE, Wiegmann AM, Sheth SN. 2022. A viewpoint on ecological and evolutionary study of plant thermal performance curves in a warming world. AoB Plants **14**, plac016.

Yao Y, Vehtari A, Simpson D, Gelman A. 2018. Using stacking to average Bayesian predictive distributions (with discussion). Bayesian Analysis **13**, 917-1007.

Younginger BS, Sirová D, Cruzan MB, Ballhorn DJ. 2017. Is biomass a reliable estimate of plant fitness? Applications in Plant Sciences **5**, 1600094.

Zhu L, Bloomfield KJ, Hocart CH, Egerton JJG, O'Sullivan OS, Penillard A, Weerasinghe LK, Atkin OK. 2018. Plasticity of photosynthetic heat tolerance in plants adapted to thermally contrasting biomes. Plant, Cell & Environment **41**, 1251-1262.

Zhu L, Scafaro AP, Vierling E, Ball MC, Posch BC, Stock F, Atkin OK. 2024. Heat tolerance of a tropical–subtropical rainforest tree species *Polyscias elegans*: time-dependent dynamic responses of physiological thermostability and biochemistry. New Phytologist **241**, 715-731.

Table 1: Model output summaries for five phenotypic traits representing reproductive fitness, phenology, biomass, and two leaf traits.

Response variable:	Total reproductive stems (n = 989)	Flowering onset (n = 1,003)	Dry biomass (n = 987)	Chlorophyll content (n = 983)	LMA (n = 974)
Fixed effects					
Estimate [95% CI]					
Intercept (cool Growth temp., cool Parental temp., outcrossed)	**35.907 [28.982, 42.255]**	**45.421 [43.721, 47.094]**	**7.550 [6.732, 8.343]**	**32.910 [31.322, 34.666]**	**1.044 [0.946, 1.134]**
Growth temp. (warm)	**3.946 [0.843, 7.060]**	**-7.899 [-8.861, -6.913]**	**0.745 [0.173, 1.344]**	**3.186 [1.908, 4.436]**	**0.100 [0.047, 0.155]**
Parental temp. (warm)	-1.393 [-5.438, 2.449]	1.153 [-0.336, 2.676]	0.104 [-0.561, 0.758]	-0.204 [-1.468, 1.074]	0.025 [-0.024, 0.073]
Growth temp. × Parental temp.	-0.543 [-4.781, 3.743]	-0.592 [-1.866, 0.626]	-0.153 [-0.915, 0.608]	0.265 [-1.282, 1.816]	-0.059 [-0.127, 0.011]
Cross type (self-pollinated)	**-7.830 [-10.957, -4.789]**	**1.492 [0.458, 2.525]**	**-1.297 [-1.833, -0.744]**	0.785 [-0.227, 1.793]	-0.002 [-0.041, 0.037]
Harvest date (later)	**6.496 [4.119, 8.877]**	--	**4.416 [4.007, 4.820]**	**1.173 [0.376, 1.941]**	**0.155 [0.120, 0.189]**
Random effects: variance components					
Estimate (SD) [95% CI]					
V_B intercept (block)	4.587 [1.216, 14.030]	0.409 [0.012, 1.709]	0.375 [0.020, 1.401]	0.990 [0.085, 3.328]	0.071 [0.018, 0.242]
V_A intercept (additive genetic)	4.271 [0.743,7.317]	2.808 [1.948, 3.588]	0.787 [0.336, 1.205]	1.416 [0.274, 2.297]	0.028 [0.002, 0.063]
V_A slope (G×E) *	1.830 [0.070, 5.025]	1.130 [0.232, 2.021]	0.664 [0.179, 1.147]	1.538 [0.219, 2.732]	0.062 [0.014, 0.106]
V_A intercept-slope correlation	-0.023 [-0.927, 0.929]	-0.793 [-0.993, -0.242]	0.575 [-0.178, 0.984]	0.124 [-0.678, 0.926]	0.294 [-0.735, 0.967]
V_M intercept (maternal)	4.343 [0.744, 7.101]	0.964 [0.044, 2.059]	0.596 [0.051, 1.119]	1.018 [0.072, 2.015]	0.028 [0.002, 0.063]
V_R (residual)	21.415 [20.277, 22.567]	5.268 [5.009, 5.546]	2.788 [2.653, 2.930]	5.663 [5.391, 5.949]	0.285 [0.271, 0.301]
Model stacking weights					
Model with G×E *	0.000	0.396	0.453	**0.594**	**0.716**
Model without G×E *	**1.000**	**0.604**	**0.547**	0.406	0.284
R^2					
mR^2 (fixed effects)	0.247	0.582	0.546	0.228	0.026
cR^2 - mR^2 (random effects)	0.561	0.241	0.171	0.387	0.032

Estimates are posterior modes with [95% CIs]; bold represents fixed effects that have 95% CIs that are distinct from zero; Harvest date refers to plants that were harvested either immediately following the hailstorm or later, see *Methods* for details; * Model outputs reported are full models that include G×E term, however please see the Model stacking weights for whether there is statistical support (bold) for the G×E term.

Table 2: Model output summaries for four phenotypic traits representing photosynthetic physiology and thermal tolerance.

Response variable:	F_V/F_M (n = 717)	ϕPSII (n = 717)	$T_{\text{crit-hot}}$ (°C) (n = 685)	$T_{\text{crit-cold}}$ (°C) (n = 707)
Fixed effects				
Estimate [95% CI]				
Intercept (cool Growth temp., cool Parental temp., outcrossed)	**7.277 [6.969, 7.585]**	**2.574 [2.134, 2.981]**	**44.459 [43.201, 45.691]**	**-13.256 [-14.312, -12.301]**
Growth temp. (warm)	**0.089 [0.029, 0.150]**	**0.332 [0.097, 0.560]**	0.256 [-0.158, 0.675]	0.345 [-0.052, 0.750]
Parental temp. (warm)	0.054 [-0.005, 0.115]	0.037 [-0.196, 0.270]	-0.039 [-0.451, 0.376]	0.186 [-0.217, 0.591]
Growth temp. × Parental temp.	-0.033 [-0.116, 0.047]	-0.023 [-0.339, 0.290]	0.059 [-0.503, 0.627]	-0.158 [-0.720, 0.398]
Cross type (self-pollinated)	0.036 [-0.008, 0.082]	0.082 [-0.091, 0.254]	0.003 [-0.316, 0.326]	-0.186 [-0.490, 0.119]
Hail (measured post-hail)	**-0.124 [-0.221, -0.031]**	-0.028 [-0.312, 0.253]	0.395 [-0.208, 1.034]	-0.101 [-0.623, 0.491]
Random effects: variance components				
Estimate (SD) [95% CI]				
V_B intercept (block)	0.182 [0.008, 0.938]	0.245 [0.007, 1.167]	0.880 [0.157, 3.091]	0.663 [0.083, 2.414]
V_A intercept (additive genetic)	0.027 [0.001, 0.075]	0.124 [0.006, 0.310]	0.228 [0.012, 0.521]	0.182 [0.007, 0.462]
V_A slope (G×E) *	0.042 [0.002, 0.110]	0.174 [0.007, 0.423]	0.307 [0.017, 0.696]	0.334 [0.025, 0.710]
V_A intercept-slope correlation	-0.268 [-0.976, 0.902]	-0.201 [-0.969, 0.902]	-0.136 [-0.951, 0.921]	-0.083 [-0.942, 0.932]
V_M intercept (maternal)	0.022 [0.001, 0.059]	0.117 [0.006, 0.267]	0.203 [0.009, 0.462]	0.141 [0.007, 0.367]
V_R (residual)	0.336 [0.317, 0.358]	1.015 [0.958, 1.074]	1.793 [1.696, 1.901]	2.252 [2.123, 2.390]
Model stacking weights				
Model with G×E *	0.041	0.001	0.000	0.309
Model without G×E *	**0.959**	**0.999**	**1.000**	**0.691**
R^2				
mR^2 (fixed effects)	0.011	0.023	0.018	0.011
cR^2 - mR^2 (random effects)	0.094	0.094	0.335	0.199

Estimates are posterior modes with [95% CIs]; bold represents fixed effects that have 95% CIs that are distinct from zero; Hail refers to plants that were measured for these traits before or after the hailstorm, see *Methods* for details; * Model outputs reported are full models that include G×E term, however please see the Model stacking weights for whether there is statistical support (bold) for the G×E term.

Table 3: Summary of heritability and maternal effects on each phenotypic trait.

Phenotypic trait	h^2 (V_A/V_P)	m^2 (V_M/V_P)
Total reproductive stems	0.040 [<0.001, 0.085]	0.040 [<0.001, 0.085]
Flowering onset	**0.142 [0.056, 0.226]**	0.050 [<0.001, 0.136]
Biomass	**0.114 [0.043, 0.189]**	0.042 [<0.001, 0.107]
Chlorophyll content	**0.078 [0.018, 0.137]**	0.029 [<0.001, 0.082]
LMA	0.025 [<0.001, 0.054]	0.015 [<0.001, 0.045]
F_V/F_M	0.005 [<0.001, 0.019]	0.005 [<0.001, 0.020]
ϕPSII	0.016 [<0.001, 0.052]	0.016 [<0.001, 0.051]
$T_{crit\text{-}hot}$	0.017 [<0.001, 0.051]	0.014 [<0.001, 0.046]
$T_{crit\text{-}cold}$	0.010 [<0.001, 0.032]	0.005 [<0.001, 0.020]

Estimates are posterior modes with [95% CIs]. V_A is additive genetic variance; V_P is total phenotypic variance; V_M is maternal variance. Note that *brms* models are parameterised such that variance components cannot be exactly zero. The minimum value for determining whether a credible interval is distinguishable from zero is 0.001. As such, any value > 0.001 (shown in bold) indicates 95% CIs that are clearly distinct from zero, providing evidence for either heritability or maternal effects on that phenotypic trait.

Table 4: Linear (β) and quadratic (γ) selection coefficients on each of the phenotypic traits (excluding total reproductive stems because it is used to calculated relative fitness) under both growing conditions (all plants), and cool-grown and warm-grown plants alone.

Phenotypic trait	β : All plants	γ : All plants
Flowering onset	**-0.215 [-0.270, -0.162]** *	-0.024 [-0.843, 0.744] *
Biomass	**0.451 [0.399, 0.502]** *	0.321 [-0.045, 0.683] *
Chlorophyll content	-0.017 [-0.060, 0.026]	-0.562 [-1.355, 0.252]
LMA	**0.055 [0.012, 0.100]**	-0.518 [-1.104, 0.010]
F_V/F_M	0.023 [-0.025, 0.076]	-0.394 [-3.368, 2.221]
ϕPSII	0.026 [-0.019, 0.073]	-0.111 [-0.533, 0.341]
$T_{crit-hot}$	-0.030 [-0.083, 0.023]	1.107 [-2.455, 4.733]
$T_{crit-cold}$	-0.056 [-0.111, 0.001]	0.399 [-0.368, 1.151]
Phenotypic trait	β : Cool-grown plants	γ : Cool-grown plants
Flowering onset	**-0.159 [-0.206, -0.117]**	0.067 [-0.611, 0.754]
Biomass	**0.385 [0.337, 0.433]**	0.283 [-0.008, 0.565]
Chlorophyll content	-0.022 [-0.062, 0.018]	-0.459 [-1.118, 0.230]
LMA	**0.039 [0.001, 0.076]**	-0.321 [-0.754, 0.107]
F_V/F_M	0.009 [-0.036, 0.057]	-0.638 [-3.608, 1.972]
ϕPSII	0.006 [-0.040, 0.053]	-0.173 [-0.601, 0.259]
$T_{crit-hot}$	-0.021 [-0.070, 0.029]	1.050 [-2.334, 4.488]
$T_{crit-cold}$	-0.042 [-0.092, 0.007]	0.362 [-0.391, 1.118]
Phenotypic trait	β : Warm-grown plants	γ : Warm-grown plants
Flowering onset	**-0.276 [-0.327, -0.227]**	**0.978 [0.467, 1.471]**
Biomass	**0.386 [0.334, 0.438]**	**-0.600 [-0.905, -0.292]**
Chlorophyll content	0.001 [-0.047, 0.044]	-0.568 [-1.358, 0.192]
LMA	0.020 [-0.028, 0.064]	**-0.733 [-1.356, -0.139]**
F_V/F_M	-0.010 [-0.074, 0.047]	1.257 [-0.970, 3.122]
ϕPSII	0.009 [-0.038, 0.058]	-0.329 [-0.873, 0.169]
$T_{crit-hot}$	-0.044 [-0.092, 0.007]	0.759 [-1.809, 3.380]
$T_{crit-cold}$	-0.028 [-0.079, 0.022]	-0.567 [-1.305, 0.171]

Values in bold represent selection coefficients that have 95% CIs that are distinct from zero. The models for all plants include interaction terms between trait × growth temperature treatment for β and trait² × growth temperature treatment for γ to test whether selection differs by growth treatment. A * symbol denotes that the respective interaction term has 95% CIs that are distinct from zero, which indicates that selection varies depending on growth temperature treatment (i.e., that selection on cool-grown and warm-grown plants differs for that trait). Full model outputs are shown in Supplementary Table S3-S6.

Figure legends

Fig. 1: Mean population-level reaction norms of phenotypic traits: (A) total reproductive stems, (B) flowering onset, (C) biomass, (D) chlorophyll content, (E) LMA, as well as photosystem traits: (F) F_V/F_M and (G) ϕPSII, and thermal tolerance traits: (H) $T_{\text{crit-hot}}$ and (I) $T_{\text{crit-cold}}$ in response to growth temperature treatments. Within each cool and warm growth temperature treatment, plants were grown under an environment that was either cool (blues) or warm (oranges) and were offspring plants were from either outcrossed (solid lines) or self-pollinated (dotted lines) parents. Each parental × growth temperature combination is coloured as follows: parental plants grown under a cool environment that had offspring grown in *i*) a cool environment (dark blue) or *ii*) a warm environment (light orange), and parental plants grown under a warm environment that had offspring grown in *iii*) a cool environment (light blue) or *iv*) a warm environment (dark orange). Reaction norms are drawn based on connections between a shared parental environment and cross type (e.g., parental cool × growth cool and outcrossed is connected to parental cool × growth warm and outcrossed). Points and error bars represent means ± S.E. of the raw data.

Fig. 2: Relationships between relative fitness (*w*; calculated as the number of reproductive stems produced divided by the mean of each growth treatment) and three standardised (within each temperature treatment) phenotypic traits (A-C: flowering onset, D-F: biomass, G-I: LMA) that demonstrated non-zero selection. Left panels are all plants combined, middle panels are cool-grown plants, and right panels are warm-grown plants. Offspring plants were either outcrossed (circles) or self-pollinated (triangles) and were derived from parents that had a parental environment that was either warm or cool. The overall quadratic model fit (posterior predictions) is plotted on each panel. Note that scaling for standardising trait values on the *x*-axis is applied in each data subset and therefore individual data point positions differ along the *x*-axis between all plants, cool-grown plants, and warm-grown plants. Linear (β) and quadratic (γ) selection coefficients and 95% CIs are given in Table 4.

Fig. 3. Tests for inbreeding depression. Mean differences between outcrossed and self-pollinated plants for fitness and phenotypic traits: (A) total reproductive stems, (B) flowering onset, and (C) biomass. Points and error bars represent means ± S.E. of the raw data. All phenotypic traits are shown in Supplementary Fig. S7. Model output is shown in Table 1.

Parental × Growth temperature
- ● Cool × Cool
- ● Warm × Cool
- ● Cool × Warm
- ● Warm × Warm

Cross type
- ●— Outcrossed
- ▲⋯ Selfed