Metavinculin mediates differentiation cues into myogenic gene expression

Abinayaselvi Murugan[1], Nandeesh Bevinahalli Nanjegowda[2], Mathivanan Jothi[1*]

[1]Laboratory of Molecular Therapeutics, Department of Human Genetics, National Institute of

Mental Health and Neurosciences (NIMHANS), Bengaluru, Karnataka, India.

[2]Department of Neuropathology, National Institute of Mental Health and Neurosciences

(NIMHANS), Bengaluru, Karnataka, India.

[*]**Corresponding author**

Mathivanan Jothi, Laboratory of Molecular Therapeutics, Department of Human Genetics,

National Institute of Mental Health and Neurosciences (NIMHANS), Bengaluru 560029,

Karnataka, India

Phone: +918026995125/5124

Fax: +918026564830, +918026562121

Email: mjothi@nimhans.ac.in

Running title: Metavinculin promotes skeletal muscle differentiation

Abstract

Skeletal muscle differentiation is a highly orchestrated process where muscle progenitors fuse to give mature contractile muscle. Ubiquitous to muscle-specific isoform switching helps muscle cells to adapt to various physiological needs. However, how they decipher differentiation signals into myogenic transcription remains unexplored. Metavinculin, a muscle-specific isoform of ubiquitous cytoskeletal scaffold protein vinculin, is involved in actin remodeling and force transduction along with vinculin in muscle cells. Although it is exclusively expressed in muscle cells, its muscle-specific role has yet to be identified. Here, we report the novel function of metavinculin in switching canonical to non-canonical Wnt-signaling to execute skeletal muscle differentiation. Specifically, we have found that metavinculin expression is absent in proliferating myoblasts and is expressed during skeletal muscle differentiation and in vivo regeneration. Splicing factor Rbfox1 is responsible for the metavinculin-specific exon retention by direct binding to its downstream intron during muscle differentiation. Further, metavinculin depletion dampens the skeletal muscle differentiation program while its ectopic expression induces differentiation-associated genes even in growth-promoting conditions. Additionally, metavinculin-depletion downregulates muscle development and upregulates stress-responsive genes with deregulated Wnt-pathway. Subsequent experiments showed that pharmacological canonical Wnt-activation impedes muscle differentiation program, whereas its inhibition stimulates differentiation in growth conditions. Canonical Wnt-inhibition or non-canonical Wnt7b ectopic expression can restore muscle differentiation programs in metavinculin-depleted cells. Overall, these evidences reveal the dynamic interplay of Rbfox1-generated metavinculin in switching canonical to non-canonical Wnt-signaling through Wnt7b to establish skeletal muscle differentiation.

Keywords

Metavinculin, Costameres, Alternative splicing, Rbfox proteins, Wnt Signaling pathway, Wnt7b, Myogenesis, Skeletal muscle.

Introduction

Differentiation cues establish skeletal muscle differentiation through a cascade of cellular events, which ultimately converge to alter the epigenome for differentiation-specific transcriptional output.[1–8] It is tightly controlled by transcriptional factors like Myod1, Myf5, Myog, and Mrf4, collectively known as myocyte regulatory factors (MRFs) and myocyte enhancing factors (MEFs).[9,10] Additionally, splicing factors are also involved in establishing myogenic gene expression as muscle is one of the human tissues with high differentially expressed alternative exons.[11–13] Several proteins, switch ubiquitous to muscle-specific isoforms by distinct gene expression or alternative splicing that enable muscle cells adaptions in physiological conditions.[14–18] Isoform switching is mainly observed in cytoskeletal proteins though it reported in other proteins.[15,16,19,20] Cytoskeletal complex actively and constantly supports contracting muscle cells by modifying cellular response when they adhere to the extracellular matrix or contact with other cells.[21] The cytoskeletal dynamics are influenced by various cell signaling pathways and vice versa during differentiation.[22–24] Particularly, Serum responsive factor (SRF) function is regulated during muscle differentiation by cytoskeletal and sarcomeric dynamics through Myocardin related transcription factor (MRTF) and Striated muscle activator of Rho signaling.[25–27] Further, altered cytoskeleton activates TGF-β signaling to enhance myofibroblasts conversion in β-actin knockout mouse embryonic fibroblasts.[28] However, how isoform switching in cytoskeletal proteins contribute myogenic signaling during differentiation remains unexplored.

Vinculin is a ubiquitous cytoskeletal scaffold protein localized to focal adhesions and adherent junctions.[29] It involved in cell morphology control, integrin clustering, motility, force transduction and adhesion strength.[29] In addition, it may also have regulatory or signaling roles which largely unexplored.[30] Vinculin knockout embryos are not viable and showed retarded somites and limbs in addition to severe cardiac and neuronal defects.[31] Metavinculin, a longer muscle-specific splice variant of vinculin generated by alternative

splicing of exon 19 that encodes 68-amino acid residues, is co-localized with vinculin at the

costameres, sarcolemma, I band in the sarcomere and intercalated discs.[32–37] Its expression

was reported in smooth, cardiac, skeletal muscles and platelets.[38–42] Metavinculin loss or

point mutations at its unique exon is linked with dilated and hypertrophic

cardiomyopathies.[39,43–48] Cholinergic receptor stimulation recruits metavinculin into the

integrin binding complex in airway smooth muscle.[49] Several proteins activate vinculin by

disrupting its auto-inhibitory head and tail interaction and metavinculin forms heterodimer

with activated vinculin.[39,49–51] Upon activation vinculin binds with actin filaments (F-actin),

facilitating vinculin dimerization for its bundling. Conversely, metavinculin binding to F-actin

inhibits vinculin dimerization, thereby preventing actin bundling.[52–56] Further, metavinculin-

specific insert sequestered as a globular subdomain protruded from the F-actin surface also

prevents its assembly.[53] Contrastingly, metavinculin reduces F-actin bending persistence,

increasing susceptibility to breakage.[57] In addition, it also increases cell adhesion complex

stability by altered force propagation with enhanced shear force resistance.[40] These studies

linked metavinculin with actin organization and force transduction and the significance of its

unique expression in muscle cells left unexplored from its discovery.[35,52–57]

Here, we have observed differentiation-specific metavinculin isoform expression in skeletal

muscle cells. Several splicing factors regulate alternative splicing in muscle cells, however,

the factor/s responsible for metavinculin generation has remained unknown.[35,36,58,59] Here,

we report Rbfox1 regulates metavinculin splicing through direct binding at its downstream

intronic region during muscle differentiation. Further, depletion, ectopic expression and

transcriptomic analysis showed its importance of metavinculin in executing skeletal muscle

differentiation. Subsequently, muscle differentiation defects observed in metavinculin-

depleted cells is restored by either canonical Wnt-inhibition or non-canonical Wnt7b ectopic

expression. Together, all our evidences suggest that metavinculin act as a critical mediator

in switching canonical to non-canonical Wnt-signaling to establish skeletal muscle

differentiation.

Results

Metavinculin is induced during skeletal muscle differentiation and regeneration

Metavinculin expression was found in smooth and cardiac and skeletal muscles.[32–34,42,60] Here, we verified the sub-stoichiometric expression of the 124 kDa metavinculin isoform in heart, intestine, and tibialis anterior (TA) muscle tissues from C57BL/6J mice (Figure 1A). We have collected 50% confluent C2C12 (Mouse), L6 (Rat), and HSMM (Human) cells grown in high serum containing media as GM and after reaching 80% confluency they are switched to low serum (DM) media to induce differentiation. We confirmed the cell cycle inhibitor Cdkn1a induction in all muscle cells cultured in differentiation media as growth arrest is a prerequisite for differentiation (Figure S1A).[9,61] In vitro skeletal muscle differentiation was confirmed by the expression of early myogenic markers Myod1, Myog, and late marker Myh (Myosin heavy chain) at RNA (Figure S1A) and protein levels (Figure 1B). Further, we have also estimated the fusion index in these cells using Myh immunofluorescence images (Figure S1B). We have found metavinculin protein expression in C2C12, L6, and HSMM cells cultured under differentiation-promoting condition not in the growth-promoting conditions whereas vinculin expression remains unaltered (Figure 1B). Similar results are observed at transcript levels (Figure 1C). Further, we could detect metavinculin protein expression as early as twelve hours of differentiation induction and it is increased with incubation time (Figure 1D). Thus, these results showed metavinculin isoform is specifically expressed during skeletal muscle differentiation (Figure 1B-D).

Mouse muscle injury models are well-established and extensively used systems to study muscle regeneration. Here, we have utilized in vivo mouse cryoinjury model to analyze the metavinculin expression during skeletal muscle regeneration.[62] Specifically, we have created cryoinjury in the hind limb TA muscle of three-month-old male C57BL/6 mice using a metal rod precooled in dry ice. We have obtained the regenerating tissues and uninjured contralateral TA muscle as control from the same animal at different post-injury time points.

First, muscle regeneration process are evaluated by histopathological examination using Hematoxylin and eosin (H&E) (Data not shown) and Masson's trichrome staining (MTS). We have observed typical histological features of muscle regeneration after muscle injury (Figure 1E). Specifically, the three days post-injured muscle showed active myonecrosis and myophagocytosis in the injured area. The muscle regeneration appeared from the viable zone on day five post-injured muscle that showed resolving myonecrosis and inflammation. We have noticed more regenerating fibers infiltrate the injured area with focal inflammation on day ten and these fibers completely occupied the injured area on day fifteen. At twenty days post-injury the damaged muscle completely regenerated (Figure 1E). RT-qPCR showed induction of metavinculin transcript at day five and decreased gradually until completion of regeneration (Figure 1F). Similarly, metavinculin protein expression is increased in ten days post-injured muscle and decreased in later time points (Figure 1G). Strikingly, we have noticed a transient metavinculin protein reduction in three days post-injured muscle compared to uninjured and marker of proliferation Ki-67 also appeared at this time point (Figure 1G). This further supported the negative correlation of metavinculin expression with proliferation. Hence, our in vitro and in vivo experiments revealed metavinculin expresses in differentiating not in proliferating cells (Figure 1B-1G).

Splicing regulator Rbfox1 retains metavinculin-specific exon during skeletal muscle differentiation

Muscle-specific metavinculin expression was identified more than thirty years ago however the splicing factor regulates vinculin alternative splicing is not identified yet.[35,36] Although several splicing factors include CELF, MBNL, Fox, and PTB families regulate alternative splicing in the muscle cells,[58,59] they have very low tissue specificity score except Rbfox family (Table S3). Notably, skeletal muscle-specific enhanced expression was observed only for Rbfox1 with the tissue specificity Tau index 0.82 (Table S3). Here, we have found differentiation-associated induction of Rbfox1, no change in Rbfox2, and variable expression of Rbfox3 mRNAs in C2C12 and L6 cells (Figure 2A and 2B). Rbfox1 and Rbfox2 are

associated with skeletal muscle and Rbfox3 with neuronal development.[15,16,58,63] Therefore,

we have focused on Rbfox1 and Rbfox2 in metavinculin generation and excluded Rbfox3.

Rbfox family recognizes auxiliary cis element 5'-GCAUG on the target RNA and regulates

exon inclusion or exclusion depending on its location.[58] We have analyzed the putative

Rbfox binding element within the 1kb intronic region that flanks metavinculin-specific exon

using RBPmap Ver. 1.2 and found two potential Rbfox binding sites in the upstream and

eight in the downstream intron region (Figure S2A). To validate Rbfox1/2 binding in

metavinculin-flanking introns we have utilized GST-pulldown assay using recombinant GST-

Rbfox1, GST-Rbfox2, and control GST (Figure S2B). Here, GST-tagged proteins are

incubated with the total RNAs isolated from C2C12 cells grown in GM and DM media. The

bound RNAs are analyzed by RT-qPCR using specific primers directs putative Rbfox binding

sites (Table S1). We have noted GST-Rbfox1 and GST-Rbfox2 binding to the 5'-intronic

flank in both GM and DM conditions (Figure 2C). Further, we have observed the

differentiation-specific enrichment at the 3'-intronic flank by both the proteins, however, GST-

Rbfox1 enriched more than GST-Rbfox2 (Figure 2D). To identify the Rbfox protein that is

responsible for metavinculin generation we have created stable C2-shRbfx1, C2-shRbfx2,

and control C2-Scr cells using lentivirus-mediated transduction shRNAs targeting Rbfox1,

Rbfox2 and scramble shRNA (Table S1). Rbfoxs depletion in these cells are confirmed by

RT-qPCR analysis (Figure 2E). Surprisingly, we have observed that Rbfox1-depletion

significantly increased the Rbfox2 mRNA expression whereas Rbfox2-depletion reduced

Rbfox1 mRNA expression (Figure 2E). Metavinculin mRNA and protein levels are reduced in

C2-shRbfx1 cells compared to C2-shRbfx2 and scramble C2-Scr (Figure 2F and 2G).

Further, we have found that Rbfox1-depletion completely inhibits the expression of early and

late myogenic genes at RNA and protein levels (Figure S2C and S2D). The differentiation

defect in these cells is also noted by reduced myotube formation and fusion index (Figure

S2E and S2F). Although Rbfox2-depletion inhibited skeletal muscle differentiation it is not as

effective as Rbfox1-depletion (Figure S2C-S2F). However, Rbfox2-depletion affects skeletal

muscle differentiation while retaining metavinculin expression (Figure 2F and 2G, S2C-S2F).

These results suggest that Rbfox1 is needed for metavinculin expression during muscle

differentiation (Figure 2E-2G).

To further confirm Rbfox1 in metavinculin generation, we have performed crosslink-RNA

immunoprecipitation (crosslink-RIP). Here, we have used the C2-Scr and C2-shRbfx1 cells

cultured in DM conditions and C2-Scr in GM as an additional control to analyze Rbfox1

bound RNAs using anti-Rbfox1 and isotype control antibodies. RT-qPCR analysis of

crosslink-RIP RNAs showed metavinculin flanking intron regions were enriched only with

anti-Rbfox1 not in the NRIgG (Figure 2H and 2I). Specifically, metavinculin 5'-intron is

enriched in anti-Rbfox1 crosslink-RIP under growth conditions and it is decreased during

differentiation conditions (Figure 2H). In contrast, 3'-intron is enriched in RIP under

differentiation conditions compared to the growth (Figure 2I). The presence of Rbfox1 is

confirmed in input and immunoprecipitate by western blot analysis (Figure 2J). As expected,

Rbfox1-depletion abrogates these enrichments (Figure 2H and 2I). These results further

confirmed that Rbfox1 specifically switches its binding from the 5'- metavinculin intronic

region to the 3'- intronic region during skeletal muscle differentiation thereby retains

metavinculin-specific exon (Figure 2H and 2I).

We have generated stable C2-Rbfox1 and C2-Vec cells that ectopically express Rbfox1

cDNA or vector in C2C12 cells by retroviral transduction to further validate Rbfox1 in

metavinculin generation. Ectopic expression of Rbfox1 is confirmed in RT-qPCR (Figure 2K)

and western blot analysis (Figure 2L). Rbfox1-mediated Mef2d splicing pattern was also

observed in C2-Rbfox1 cells as reported earlier (Figure S2G).[16] Ectopic Rbfox1 induced

metavinculin mRNA compared to vector control, however, it is not reflected in the protein

level (Figure 2M and 2N). The myogenic differentiation genes are also induced only at RNA

levels in C2-Rbfox1 cells compared to C2-Vec (Figure 2O and 2P). Nevertheless, these

results also support the involvement of Rbfox1 in metavinculin exon inclusion in growth-

promoting conditions. Collectively, these evidences strongly suggest that Rbfox1 occupy

downstream flanking intronic region to include metavinculin-specific exon during skeletal

muscle differentiation (Figure 2).

Metavinculin expression is required for skeletal muscle cell differentiation

Metavinculin expression found only in differentiating myoblasts (Figure 1).[32,38,40] We have

designed and cloned two shRNAs (shmVcl1 and shmVcl2) targeting metavinculin-specific

exon region and generated cell lines C2-shmVcl1 and C2-shmVcl2 that stably express these

shRNAs and scramble control C2-Scr by lentiviral mediated transduction to identify

differentiation-specific role of metavinculin. The effective metavinculin depletion is found by

shmVcl2 compared to shmVcl1 shRNA at RNA and protein levels and vinculin levels are not

altered by these shRNAs (Figure 3A and 3B). Therefore, we have used shmVcl2 shRNA to

generate L6-shmVcl2 from L6 rat myoblast and scramble control L6-Scr cells as mentioned

above. RT-qPCR results showed the inhibition of muscle-specific genes Myod1, Myog, Myh,

and growth inhibitor Cdkn1a in metavinculin-depleted C2C12 and L6 cells cultured under DM

media (Figure 3C and 3D). Similarly, differentiation-associated myogenic proteins are also

repressed by metavinculin depletion in these cells (Figure 3E). The inhibition of myogenic

gene activation by metavinculin depletion severely affected myotube formation as shown by

Myh immunofluorescence and fusion index (Figure 3F). Myod1 occupancy to its target gene

Myog and Myh promoters is also inhibited by metavinculin depletion in C2C12 cells (Figure

S3A and S3B). These experiments revealed that metavinculin expression is necessary to

induce myogenic gene expression during skeletal muscle differentiation (Figure 3C-3F).

Therefore, we have verified its ability to induce these genes at growth-promoting conditions.

For this we have transiently expressed GFP-tagged metavinculin in C2C12 and L6

myoblasts and analyzed for myogenic gene expression after 48 hours of post-transfection.

The ectopic expression of metavinculin is confirmed using western blot analysis (Figure 3G).

Metavinculin ectopic expression is sufficient to induce myogenic marker genes Myod, Myog,

and Myh even at growth-promoting conditions in C2C12 and L6 cells (Figure 3G). These

results revealed the importance of metavincluin in inducing myogenic gene expression.

Further, we have analyzed the whole transcriptomic changes by metavinculin depletion

using RNA-Seq analysis. We have isolated total RNA from C2-shmVcl2 and C2-Scr cells

cultured under differentiation conditions using the Trizol method in biological triplicates. The

libraries are synthesized from these RNAs and paired-end sequencing is performed. We

have verified the clustering of two distinct groups in a principal compound analysis (PCA)

(Figure S4). Here, we have found 1,387 differentially expressed genes (DEGs) (FDR <0.05)

includes 809 upregulated and 578 downregulated genes in metavinculin-depleted C2-

shmVcl2 compared to control C2-Scr cells using edgeR (Figure 4A and 4B). Randomly

picked upregulated (Klhl13, Heatr9, Cxcl10, Ccl5, Acod1, Slurp1) and downregulated (Ccn2,

Tagln, Actg2, Myl9, Car3) DEGs are further validated and analyzed for their differentiation-

specific expression changes by RT-qPCR analysis. Here, we have found decreased level of

Klhl13, Ccl5, Acod1, Slurp1 and increased level of Heatr9, Cxcl10 during skeletal muscle

differentiation and they are further increased by metavinculin depletion in C2-shmVcl2 cells

(Figure 4C). In other hand downregulated DEGs Ccn2, Tagln, Actg2, Myl9, Car3

expressions are upregulated during normal skeletal muscle differentiation and metavinculin

depletion inhibits the same (Figure 4D). Gene ontology (GO) of the differentially expressed

genes was analyzed into various functional categories including Biological Process (BP) and

Cellular Components (CC) using the online tool ShinyGO 0.80. Upregulated genes are

mostly associated with the various stress-responsive immune modulatory GO terms under

the BP category (Figure 4E). Further, the CC category of upregulated genes showed the

association of DNA replication fork, preinitiation, MCM, and CMG complexes, which suggest

DNA replication during the cell cycle (Figure S5A). Similar results were also observed in the

KEGG eukaryotic DNA replication complex (Figure S5B). In addition, G1/S-specific cyclin

Ccnd1 and G2/M-specific cyclin Ccnb1 are also appeared in upregulated DEGs and further

analysis of FPKM data revealed more than two-fold upregulation of Ccna2, Ccnb2, Ccne2

and Ccnf cyclins (Figure S5C, Table S4). These findings further suggest that induction of cell

cycle in metavinculin-depleted cells under differentiation-promoting conditions. Besides, BP

analysis of downregulated genes showed that all the top ten statistically significant GO terms

enriched with muscle contraction, differentiation, and development, in particular striated

muscle (Figure 4E). Actin filament-based process and regulation, actin cytoskeleton

regulation are also associated with the downregulated DEGs (Figure 4E). Additionally, tube

morphogenesis and development was also significant GO terms associate with

downregulated genes (Figure 4E). Further, the downregulated DEGs are associated with

typical muscle components like myofilament, actin filament bundle, actomyosin, myofibril, I

band, and Z disc (Figure S6A). Focal adhesion, adherence junction, cell-substrate junction,

and anchoring junction are notable GO terms associated with downregulated genes (Figure

S6A). Hence, GO analysis of downregulated DEGs showed that metavinculin depletion

severely impedes the genes associated with muscle differentiation and functions in addition

to cytoskeletal network genes. In a parallel KEGG analysis, downregulated DEGs are

associated with dilated, hypertrophic cardiomyopathies and Adrenergic, Wnt and MAPK

signaling (Figure S6B). Interestingly, KEGG Wnt-signaling pathway enriched DEGs are

mostly involved in β-catenin independent non-canonical Planar Cell Polarity (PCP) and

Wnt/Ca^{2+} pathways (Figure S7A). Further, we have also observed a significant reduction of

FPKM of non-canonical Wnt7b in C2-shmVcl2 cells compared to C2-Scr cells after removing

the outlier and another non-canonical Wnt11 also showed more than 2-fold reduction of

FPKM however it is not statistically significant (Figure S7B). Interestingly, Wnt10a, a

canonical Wnt-pathway ligand appeared in upregulated DEGs (Table S4). Moreover, these

observations indicate that metavinculin-depletion promotes canonical and inhibits non-

canonical Wnt-signaling under differentiation conditions (Figure S7, Table S4). Overall, the

transcriptome changes in C2-shmVcl2 cells suggest that metavinculin depletion fails to

induce growth arrest and subsequent skeletal muscle differentiation perhaps due to

disrupted Wnt-pathways (Figure 4).

Canonical and non-canonical Wnt-signaling differentially regulates skeletal muscle differentiation

Studies showed the importance of Wnts in muscle development however most of them are focused on the muscle progenitor maintenance with limited evidence in skeletal muscle differentiation.[64-73] Here, metavinculin depletion severely disrupts skeletal muscle differentiation with an altered Wnt-signaling pathway (Figures 3, 4 and S7, Table S4). To confirm the significance of Wnt-activation in muscle differentiation, first we have analyzed the expression of Wnt-target genes in growth and differentiation conditions. RT-qPCR results revealed that canonical Wnt-targets Myc, Ccnd1, Ascl2, Id2, Tcf4 and Mmp9 are reduced in cells cultured under DM conditions compared to GM (Figure 5A). Then we have analysed β-catenin (Ctnnb) expression in C2-shmVcl2 and C2-Scr cells under various culturing conditions. In scramble control cells, β-catenin expression present in GM condition is diminished in DM condition whereas β-catenin level is retained in metavinculin-depleted cells (Figure 5B). Since vinculin directly binds with β-catenin to stabilize E-cadherin at the epithelial cell surface[74] we have verified the similar vinculin/β-catenin interaction during skeletal muscle differentiation by immunoprecipitation analysis. The results revealed vinculin direct interaction with β-catenin in growth conditions and it is absent during differentiation probably due to decreased β-catenin level (Figure 5B and 5C). This interaction is retained in metavinculin-depleted C2-shmVcl2 cells that showed differentiation defect (Figure 3 and 5C). To further explore canonical Wnt-signaling mediated differentiation inhibition, we have treated the cells with the reported canonical Wnt-activator BML-284[75] and analyzed its effect on differentiation potential. BML-284 treatment activated the canonical Wnt-pathway target genes Myc, Ccnd1, Id2, Tcf4, and Mmp9 in C2C12 cells (Figure 5D). Further, BML-284 treatment completely inhibited the induction of growth arrest gene Cdkn1a and myogenic genes Myh, Myog, Myod1, in C2C12 cells during differentiation (Figure 5E). The levels of Myod1, Myog, and Myh proteins are also inhibited by BML-284 treatment (Figure 5F). Wnt-activation also reduced Myh positive cells and myotube formation in C2C12 cells under DM

conditions compared to vehicle-treated cells (Figure 5G). Since canonical Wnt-activation

negatively regulates skeletal muscle differentiation the observed differentiation defect in

metavinculin depleted cells could be due to sustained canonical Wnt-signaling by β-catenin

stabilisation (Figure 5).

Wnt-activation inhibits the Cdkn1a expression which is necessary to induce growth arrest

during skeletal muscle differentiation (Figure 5E).[9,61] As Wnt-activation prevents skeletal

muscle differentiation its inhibition may induce differentiation (Figure 5E-5G). To confirm this,

we have inhibited the canonical Wnt-pathway using a potent Wnt-inhibitor KY02111[76] in

C2C12 cells (Figure 6A). KY02111-treatment induced Cdkn1a transcripts along with Myod1,

Myog, and Myh under the growth promoting conditions compared to vehicle treatment

(Figure 6B). We have also confirmed the induction of myogenic genes at the protein level

(Figure 6C). KY02111-treatment increased Myh-positive C2C12 cells at growth condition,

however, they did not show any fusion (Figure 6D). Since canonical Wnt-inhibition is

essential for skeletal muscle differentiation (Figure 6A-6D) it may recover differentiation

defect in metavinculin-depleted cells. As expected, KY02111-treatment upregulates Cdkn1a,

Myod1, Myog, and Myh mRNAs expression in C2-shmVcl2 cells compared to vehicle treated

cell (Figure 6E). Myh, Myog, and Myod1 levels are also induced in these cells by KY02111

(Figure 6F). Increased myotube formation and fusion index in metavinculin-depleted cells

are also noted in KY02111 treated cells (Figure 6G). Thus, these results revealed that

canonical Wnt-inhibition can induces skeletal muscle differentiation and rescue

differentiation in metavinculin depleted cells (Figure 6).

Metavinculin-depletion inhibits the non-canonical Wnt-pathway while activating the canonical

pathway (Figure S7A, Table S4). Inhibition of non-canonical Wnt7b in metavinculin-depleted

cells (Figure S7B) is further confirmed by RT-qPCR (Figure 7A). Antagonism between non-

cantonical and canonical Wnt-signaling is reported in several studies and we have also

found induction of non-canonical Wnt7b during canonical Wnt-inhibition in C2C12 cells

(Figure 7B).[77–79] Further, we have observed induction of Wnt7b protein in C2C12 and L6

cells during differentiation (Figure 7C). Differentiation-specific Wnt7b induction is inhibited in

metavinculin-depleted C2-shmVcl2 and L6-shmVcl2 cells (Figure 7A and 7C). Since

canonical Wnt-inhibition can activate non-canonical Wnt-signaling we have analyzed the

status of non-canonical Wnt7b in KY02111-treated metavinculin-depleted cells (Figure

7B).[77–80] KY02111-treatment induced Wnt7b mRNA and protein levels in C2-shmVcl2

compared to vehicle while restoring skeletal muscle differentiation (Figure 6E-6G, 7D and

7E). Similarly, ectopic Wnt7b expression also likely to restore skeletal muscle differentiation

these cells (Figure 7F and 7G). As expected, retroviral-mediated ectopic Wnt7b expression

restored of myogenic genes Myh, Myog, Myod1 and Cdnk1a in C2-shmVcl2 cells, however,

the induction of late myogenic marker Myh is less compared to early myogenic genes

(Figure 7H). Further, the restoration of myogenic genes are also reflected in their protein

levels (Figure 7I). Ectopic Wnt7b expression is also improved myotube formation and fusion

index in metavinculin-depleted cells compared to vector control (Figure 7J). Collectively,

these data revealed that canonical Wnt-inhibition or non-canonical activation can restore

skeletal muscle differentiation in metavinculin depleted cells (Figure 6 and 7).

Discussion

Metavinculin is differentially expressed during myoblast proliferation and
differentiation

Metavinculin express along with vinculin at sub-stoichiometric in skeletal muscle as well as

smooth, cardiac muscles (Figure 1A).[32–34,40,42,60] However, proliferation dependent reduction

of metavinculin level reported in primary cells isolated from human aorta and chicken

gizzards and completely absent when they sub-cultured.[32,38,40] Similarly, we have found

metavinculin expression only in differentiated myotubes not in the proliferating myoblasts

while vinculin level remains unaltered (Figure 1B-1D). Proliferating myoblasts first exit from

the cell cycle and undergoes terminal muscle differentiation by activation of various

myogenic genes.[9,61] Differentiation signal leads to master myogenic regulator Myod1

activation which in turn activates its targets Cdkn1a (a growth arrest gene) followed by Myog

by altering their epigenome.[6,7] Later, Myod1 and Myog collaborate to activate terminal

myogenic genes.[3,8] We have found metavinculin isoform expression gradually increased

during in vitro skeletal muscle differentiation with no change in vinculin level (Figure 1D).

After muscle injury, activated resident satellite cells extensively proliferate and undergo

differentiation program to complete muscle regeneration process.[81,82] Here, we have found a

direct correlation of metavinculin expression with skeletal muscle differentiation during

regeneration in the mouse cryoinjury model (Figure 1E-1G); however, its expression is

transiently reduced at the early stages of regeneration where excessive cell division takes

place (Figure 1G). Hence, it appears that metavinculin isoform switching is controlled based

on the status of muscle cell proliferation or differentiation (Figure 1B-1G).[32,34,40]

Rbfox1 regulates metavinculin splicing during skeletal muscle differentiation

Alternative splicing is also major contributor for the diverse muscle proteome in addition to

myogenic transcription factors.[1,11,13,59] Alternative transcript usage gives spatiotemporal

expression of unique isoforms during skeletal muscle differentiation and it strongly correlates

with the various splicing regulators, including Rbfox, Celf, Mbnl and Ptb expression.[58,59]

Among these factors, Rbfox family may collaborate with other splicing regulators to establish

muscle-specific splicing events during muscle differentiation as they exhibit high tissue

specificity (Table S1).[16,63] Rbfox1 and Rbfox2 are the major Rbfox family members

expressed in muscle cells. Rbfox1 expression is induced during myotube formation while

Rbfox2 expression remains unaltered and the same is also reported by others (Figure 2A

and 2B).[16,63] We did not find consistent change in Rbfox3 level during muscle differentiation

as it is involved in neuronal development (Figure 2A and 2B). Rbfox family include or

exclude exons depending on the location of its cis-regulatory elements, in specific, upstream

location excludes and the downstream location includes the associated cassette exon.[83]

Metavinculin is a larger vinculin isoform specifically expressed in differentiated muscle cells

and its expression is reduced in proliferating cells which coincides with Rbfox1 expression

(Figure 1A-1C, 2A and 2B).[32,34,40] Although both Rbfox1 and Rbfox2 can bind in vitro to

metavinculin flanking introns the binding is enriched at downstream intron in myotubes

(Figure 2C and 2D). Moreover, Rbfox1 regulates metavinculin expression over Rbfox2 as

shown by depletion experiments (Figure 2E-2G). Further, crosslink-RIP analysis confirmed

the differentiation-specific switching of Rbfox1 from 5'- to 3'- metavinculin flanking intron

(Figure 2H-2J). While ectopic Rbfox1 expression showed induction of metavinculin

transcripts under growth-promoting conditions however it is not reflected at protein level

which suggested the unstable metavinculin protein in proliferating cells (Figure 1B, 1D and

2K-2N).[32,38,40]

It was reported that Rbfox1 or Rbfox2 depletion does not affect early stages of muscle

differentiation and it affected the myotube formation.[16,63] In contrast, we have observed

Rbfox1-depletion severely inhibited the expression of early myogenic genes Myod1, Myog in

addition to terminal differentiation marker Myh whereas certain level of these myogenic

genes are retained in Rbfox2-depleted cells and this difference observation is probably due

to transient or stable nature of depletion (Figure S2C and S2D).[16,63] Although both Rbfoxs

are necessary for skeletal muscle differentiation Rbfox1 contributes more during

differentiation as its ectopic expression can induce differentiation associated genes even at

growth promoting conditions (Figure S2E, S2F, 2O and 2P). Interestingly, we have found

Rbfox1-depletion negatively and Rbfox2-depletion positively correlated with each other's

expression (Figure 2E). Thus, it appears that there is a negative feedback inhibition between

Rbfox2 and Rbfox1 expression during skeletal muscle differentiation and it will be interesting

to explore further. Nevertheless, these results suggest that induced Rbfox1 expression

critically regulates metavinculin alternative splicing during differentiation by preferentially

binding to its downstream flanking intron (Figure 2).

Metavinculin critically regulates muscle cell proliferation and differentiation

Vinculin links the integrin adhesion complex to the filamentous actin network and evidences

suggest that metavinculin play a similar role along with vinculin with enhanced force

transduction, however, its muscle-specific remains to be identified.[29,37,39,52,55,57,84] Here,

downregulated DEGs in metavinculin-depleted cells are associated with focal adhesion,

adherens, anchoring and cell-substrate junctions in skeletal muscle cells as metavinculin is

necessary to stabilize these complexes similar to vinculin (Figure S6A). The major functional

difference reported between vinculin and metavinculin lays in the actin bundling.[37,52,55–57]

Metavinculin may fine-tune actin organisation in skeletal muscles as downregulated DEGs in

metavinculin-depleted cells associates with actin bundling terms (Figure S6A).[52,53,55,57]

Moreover, we have identified the unique role metavinculin in skeletal muscle differentiation

through depletion and ectopic expression studies. Specifically, metavinculin depletion

inhibited early and late genes involved in muscle differentiation (Figure 3A-3F). Interestingly,

ectopic expression of metavinculin is able to induce these genes even in growth-promoting

conditions (Figure 3G). Transcriptome analysis in metavinculin-depleted cells also confirmed

the downregulation of genes associated with muscle development and contraction (Figure

4C). In particular, downregulated DEGs in metavinculin-depleted cells associate with cellular

compounds actomyosin, myofibril, contractile fiber, sarcomere, I band, and Z disc (Figure

S6A). Vinculin knockout is embryonic lethal due to multiple defects in the heart, neural tube,

somites, and limb.[31] In agreement with this metavinculin depletion also affected the several

genes involved in muscle and neural tube development (Figure 4E). Mutations at the

metavinculin-specific exon or loss of expression reported to be involved in various

cardiomyopathies.[39,43–48,84,85] Downregulated DEGs by metavinculin depletion is also

associate with these cardiomyopathies (Figure S6B). Surprisingly, the heart muscle sections

from muscle-specific metavinculin knockout mice do not show any defect in cardiac tissue

architecture, and the expression/localization of the costamere, intercalated disc, and gap

junction proteins also remains unaltered although they report metavinculin bears higher

molecular forces.[84] This discrepancy in their observation is because they targeted the

upstream region which will not disrupt metavinculin-specific exon 19, as shown in their

supplementary figure 10.[84] However, the evidence from our study supports the metavinculin

play a crucial role in establishing myogenic gene expression during differentiation and

muscle function (Figure 3 and 4).

Metavinculin isoform switching occurs during muscle differentiation and is absent in

proliferating cells (Figure 1B and 1C).[33,38,40] Moreover, metavinculin depletion inhibits the

expression of growth arrest gene Cdkn1a which is necessary to induce cell cycle exit during

skeletal muscle differentiation (Figure 3A-3D).[9,61] Further, metavinculin depletion

upregulated the genes involved in various stress-responsive pathways, DNA replication

complexes, and several cyclins (Figure 4E, S5). Thus, it appears that metavinculin

expression is needed for growth arrest prior to skeletal muscle differentiation as its absence

retains the cells in a proliferative stage even under differentiation-promoting conditions.

Taken together, these results indicate that metavinculin is critical for muscle cells

proliferation and differentiation decisions.

Metavinculin switches canonical to non-canonical Wnt-signaling during muscle

differentiation

Wnt-signaling plays a crucial role in stem cell proliferation, differentiation, and self-renewal

during embryonic development as knockout of Wnt-pathway genes showed embryonic

lethality with multiple tissue defects including muscle.[64,65,67,69,71,72] In the current study,

metavinculin-depletion inhibited the genes associated with muscle development along with

Wnt-pathway deregulation (Figure 3, 4F, S6 and S7, Table S4). Several Wnt ligands controls

MRFs expression during muscle progenitor cell proliferation and differentiation.[64,69–71]

Specifically, canonical Wnt1, 3a, 4, and 6 maintain the expression of Pax3 and Pax7 in

muscle progenitors.[86] Wnt1 and Wnt3a double knockout mouse embryos failed to form

proper dermomyotome due to reduced expression of Myf5.[69] Further, Wnt1 regulates Myf5

expression and non-canonical Wnt7a activates Myod1 in epaxial and hypaxial muscle

respectively.[64,66] Canonical Wnt mediator β-catenin inhibition is required for proper skeletal

muscle differentiation as its activation by BML-284 prevents muscle differentiation (Figure 5,

S8). We have found direct binding of β-catenin with vinculin during myoblast proliferation and

it diminished during differentiation (Figure 5C). Restored β-catenin level by vinculin/β-catenin

interaction may responsible for the differentiation defects observed in metavinculin-depleted

cells (Figure 3, 4F, 5B and 5C). Further, canonical Wnt inhibition by KY02111 induce

myogenic genes even in growth-promoting conditions and restores myogenic differentiation

in metavinculin-depleted cells (Figure 6). Hence, these evidences suggest that metavinculin

destabilizes β-catenin by interfering β-catenin/vinculin association during skeletal muscle

differentiation.

Non-canonical Wnt5a, Wnt5b, Wnt7a, Wnt7b are important for various stages of muscle

regeneration.[65,87,88] Moreover, Wnt11-mediated Wnt/PCP pathway is necessary for

embryonic muscle fibre formation.[68] A recent study demonstrates that Wnt7b expression in

muscle stem cells is repressed by the cis-acting long noncoding RNA Lnc-Rewind and it is

relived during differentiation.[89] We have also found the induction of non-canonical Wnt7b

during skeletal muscle differentiation and metavinculin-depletion inhibited the same (Figure

7A and 7C). KY02111 treatment activates non-canonical Wnt7b and restored myogenic

differentiation in metavinculin-depleted cells (Figure 6E-6G, 7B, 7D and 7E). Similarly,

Wnt7b induction is observed during myogenic differentiation in canonical Wnt10b[-/-]

myoblasts.[87] Non-canonical Wnt ligands can alter canonical Wnt-signaling in a spacio-

temporal manner.[65,87] Specifically, non-canonical $WNT/Ca2^+$ pathway-mediated activation of

NLK and HIPK2 inhibits the transcriptional activity of endpoint Wnt-signaling mediator

TCF/LEF by phosphorylation.[78] In addition, non-canonical Wnt5a is known to inhibit Wnt3a-

mediated canonical signaling during hematopoietic stem cell differentiation.[77] Here, ectopic

Wnt7b is sufficient to improve myogenic induction and the fusion index in metavinculin-

depleted cells (Figure 7F-7J). In contrast to these observations involvement of canonical

Wnt-signalling in inducing muscle differentiation is reported in various studies.[65,70,71,88,90,91]

However, these studies further insist its role in muscle progenitor maintenance not terminal

differentiation. Hence, it appears that canonical Wnt-signaling regulates muscle progenitor

proliferation and maintenance whereas non-canonical Wnt signals involve in terminal muscle

differentiation.

To conclude, vinculin/β-catenin association sustains canonical Wnt-signaling in proliferating

myoblasts. During skeletal muscle differentiation, upregulated Rbfox1 generates

metavinculin isoform which in turn switches canonical to non-canonical Wnt-signaling in

establishing skeletal muscle differentiation program (Figure 8).

Materials and Methods

Animals

All mouse protocols were in accordance with NIMHANS Institutional Animal Ethics

Committee (IAEC) guidelines (Approval number: AEC/74/487/H.G.) which is registered

under Committee for Control and Supervision of Experiments on Animals (CCSEA), Ministry

of Fisheries, Animal Husbandry and Dairying, Government of India (Registration number:

12/GO/ReBi/S/1999/CPCSEA). Three months old C57BL/6J male mice were used for

cryoinjury muscle regeneration model. Cryoinjury was created for 15 seconds in a right

tibialis muscle using 5mm sterilized metal rod precooled in dry ice and uninjured

contralateral muscle was used as a control. Animal weights were monitored on daily basis,

uninjured and regenerating muscle tissues were harvested at 3, 5, 10, 15 and 20 days of

post injury after sacrifice for further analysis.

Cell lines

The C2C12, L6, HSMM myoblasts and 293T cells were used in this study. These cells were

cultured in a Dulbecco's modified Eagle's medium (DMEM) containing either 20% or 10%

Fetal Bovine Serum (FBS) (Gibco, USA) and antibiotic-antimycotic solution (Gibco, USA).

Myoblast cultured in DMEM media contains 20% Fetal Bovine Serum (FBS). These cells

were induced to differentiate into myotubes by switching the cells into differentiation media

containing 2% horse serum (Gibco, USA) for 48 hours (C2C12) or 10days (L6, HSMM). Wnt

activator BML-284 (5 mM) (AdooQ, USA) or inhibitor KY02111 (10 mM) (AdooQ, USA) and

vehicle was treated in indicated culturing conditions and refreshed during media change.

Stable polyclonal population of C2-shmVcl1, C2-shmVcl2, C2-shRbfox1, C2-shRbfox2, C2-

shScr, L6-shScr, L6-shmVcl2 were generated through lentiviral transduction followed by

puromycin (Sigma, USA) selection. Similarly C2-Rbfox1 cells were generated using retroviral

transduction. All the cells were maintained at 37°C, 5% CO_2 in a humidified atmosphere.

Molecular cloning

shRNA oligos targeting Rbfox1, Rbfox2, metavinculin and scramble were designed in our

laboratory and synthesized from the Integrated DNA Technologies (IDT), USA along with

antisense oligos the sequence details were given in Table S1. The sense and antisense

oligos were annealed and cloned into the lentiviral pLKO.1-TRC or pLSL vector. Specifically,

Rbfox1, Rbfox2 and Scramble shRNAs were cloned between EcoRI and AgeI sites in

pLKO.1-TRC vector. Metavinculin targeted two shRNAs (shmVcl1 and shmVcl2) and

scramble were cloned into pLSL into BamH1 and EcoR1 sites. For GST fused Rbfox1 and

Rbfox2 recombinant proteins, we have subcloned respective cDNAs fragments digested

from pEGFP-Rbfox1 and pEGFP-Rbfox2 with BglII and SalI into pGEX6P-1 plasmid

between BamHI and SalI sites. Wnt7b coding sequence was PCR amplified from cDNAs

generated from C2C12 cells using specific primers contains BamHI or EcoRI restriction sites

and cloned into T vector pMD20. Further, Wnt7b cDNAs was subcloned into pBabe-Puro

vector at BamHI and EcoRI sites. Similarly, Rbfox1 cDNA was subcloned from pEGFP-

Rbfox1 plasmid. All the inserts in the cloned plasmids and their orientations were confirmed

by Sanger's sequencing.

Western blot analysis

Twenty milligrams of heart, intestine and tibialis anterior muscle tissues from C57BL/6J

mouse and regenerating tissue obtained at various time points were homogenized in a cell

lysis buffer containing with 1X protease and phosphatase inhibitors (Roche, USA) and

sonicated to get a tissue extracts. Similarly, total extracts were prepared from cell lines using

same lysis buffer. Forty microgram of the obtained tissue or cellular extracts were used for western blot analysis. Briefly, these extracts were separated on SDS-PAGE, transferred to PVDF membrane (Merck, USA), incubated with specific primary antibodies and probed with respective secondary antibodies. The primary and secondary antibodies used in a western blot analysis are given in Table S2. The signal was detected and visualized by ChemiDoc XRS+ (Biorad, USA) or Alliance Q9 (UVITEC, UK) using Immobilon® western HRP substrate (Merck, USA).

Quantitative RT-PCR

Two microgram of total RNAs isolated from tissues or cells using TRIzol (Invitrogen, USA) method were converted to cDNA using RevertAid first strand cDNA synthesis Kit (Thermo Scientific, USA) according to the manufacturer instructions. qRT-PCR from these cDNAs were performed using Maxima SYBR green/ROX qPCR master mix (Thermo Scientific, USA) in a QuantStudio™ 6 Flex Real-Time PCR System (Applied Biosystems, USA) using gene specific PCR primers given in Table S1. The results were analyzed by $2^{-\Delta\Delta}$ CT method. The gene expressions were normalized with endogenous control Actb.

Tissue preparation and histopathological analysis

Muscle regeneration in cryoinjured tissues were analyzed by routine standard histological analysis. Concisely, formalin fixed injured and control muscle tissues were processed in an automatic tissue processor (Tissue-Tek® VIP™ Vacuum Infiltration Processor, Sakura Finetek USA, Inc.) and embedded in paraffin blocks. The tissues in paraffin blocks were subjected to sectioning (3-4 μM) by rotary type of microtome (Leica RM2255 Rotary Microtome, Leica Biosystems, India). The tissue sections were fixed in a glass slide and stained with hematoxylin and eosin (H&E) and Masson's trichrome stain separately for histological analysis after deparaffinization and rehydration. The histological features were analyzed from the stained images captured from Olympus BX53 microscope (Olympus, Japan).

Immunofluorescence

Cells cultured in 35-mm dishes at various conditions were fixed with 4% paraformaldehyde,

permeabilized with 0.1% Triton X-100 in PBS and blocked with 3 % control serum. After

washing, these cells were incubated overnight with anti-Myh antibody. Then, primary

antibody solution was removed, the cells were washed and incubated with Alexa-Fluor 488

(green) conjugated secondary antibody to detect the bound primary antibody. Nuclei were

counter stained with DAPI (Sigma, USA) and the images were acquired from fluorescence

microscope IX73 (Olympus, Japan). The fusion index from the fluorescence images were

calculated using Myotube Analyzer tool.[92]

Recombinant protein purification

GST-Rbfox1, GST-Rbfox2 and GST proteins were purified as mentioned earlier.[6] Briefly,

transformed BL21(DE3) cells with respective pGEX6P-1 plasmids were cultured until OD

reaches 0.6 and the heterologous protein expression was induced with 0.25 mM IPTG for 3

hr. Then the bacteria were lysed with lysis buffer (50 mM TrisCl pH 7.5, 120 mM NaCl, 1 mM

EDTA 0.5 % NP40 and 1 mM PMSF). The clarified supernatant was incubated with GST-

agarose beads (Sigma, USA) overnight. Then the recombinant proteins were eluted from

washed beads with buffer contains 50 mM Tris pH 8.0, 1 mM DTT and 10 mM reduced

glutathione. The recombinant proteins were used for GST-pulldown assay after dialysis.

GST-pulldown assay

Total RNAs isolated form cells grown under GM and DM were treated with Purelink^{TM}

DNAse (Invitrogen, USA). The RNAs then denatured and renatured by heating 2 min at

90°C followed by gradual cooling to room temperature. Two micrograms of total RNA per

pull down were pre-cleared with Glutathione-Agarose beads (Sigma, USA) in the assay

binding buffer (PBS containing 2 mM MgCl2, 0.2 mM ZnCl2, 1mM DTT, 100 U/ml RNAse

Inhibitor, 0.1 mg/ml yeast tRNA (Sigma, USA), 0.05% BSA, and 0.2% NP40). 1/10 of this

material was used as an input. After pre-clearing the RNAs were incubated with two

micrograms of GST-Rbfox1 or GST-Rbfox2 or control GST proteins at room temperature for

1 hour in the assay binding buffer. Twenty microliter Glutathione-Agarose beads was added

and incubated overnight at 4°C to pulldown protein/RNA complex. Then, the beads were

washed four times in the same assay buffer supplemented with 150 mM NaCl at room

temperature. The RNAs from the bead-protein-RNA complex and inputs were recovered by

TRIzol method followed by on-column purification, and the eluted RNAs were converted into

cDNA and analyzed by qPCR with specific primers given in Table S1.

Lenti and retroviral transduction

The viral protocols are reviewed and approved by Institutional Biosafety Committee (IBSC),

NIMHANS with the approval number NIMHANS/DO/INSC MEETING/2017 S.No.5 dated

27.07.2017)   Retro and lenti viruses were produced in 293T packaging cell lines as

described previously.[93] Here, viral plasmids containing cDNA or shRNA along with

packaging plasmids were transiently transfected into 293T cell by calcium phosphate co-

precipitation method. Forty eight hours of post-transfection the viral supernatants were

passed through 0.45 μm syringe and the diluted viruses were used to infect cells in growth

media supplement with 8 μg/ml of fresh polybrene (Sigma, USA) for 8 hr for three

consecutive times. After 48 hr post-transduction the cells were selected with puromycin until

the untransduced cells died or used as transient infection. Viral titers were assessed by GFP

florescence in 293T cells transduced with serial dilution of GFP viral stocks.

Crosslink-RNA immunoprecipitation

The cells grown in appropriate conditions were crosslinked with 4% paraformaldehyde and

collected in PBS. The nuclei from these cells were extracted using nuclear extraction buffer

(1.28M Sucrose, 40mM Tris-Cl pH: 7.5, 20Mm MgCl2, 4% Triton X-100) and the pelleted

nuclei were resuspended in RIP buffer (150 mM KCl, 25 mM Tris, pH 7.4, 5 mM EDTA, 0.5

mM DTT, 0.5% NP40, and 100 U/mL RNase inhibitor). The chromatin was sheared by

sonication and clarified by centrifugation. For RIP, 1 μg of anti-Rbfox1 antibody or control

NRIgG was incubated overnight at 4 °C with gentile rotation and 2 % of these samples taken

as input prior to antibody addition. Then the samples were incubated with protein A-agarose

beads (SCBT, USA) for 2 h at 4 °C to pulldown immune complex. The beads were washed

three times with RIP buffer and one time with PBS. Then it was divided into two part for

protein and RNA analysis. The presence of Rbfox1 protein in the immune complex and

input were analyzed by western blot. The RNAs were isolated from the beads and input

using TRIzol method followed by on column purification. These RNAs were used in RT-

qPCR analysis as mentioned elsewhere. The primer and antibody details are given in the

Table S1 and S2.

Chromatin immunoprecipitation (ChIP)

We have followed ChIP protocol as mentioned earlier.[94] Briefly, the cells grown in

appropriate conditions were fixed with 37 % formaldehyde and collected by scraping in ice

cold PBS with PMSF. The cells were suspended in SDS lysis buffer with protease inhibitor

cocktails and sonicated 15 S each at 50 amplitude for 25 times in a probe sonicator

(Qsonica, USA). The fragment size was verified from fifty microliter of sonicated samples.

The chromatin (100 µg) was precleared with protein A agarose/Salmon sperm DNA beads

(Sigma, USA) and 2 % taken as input. Then the chromatin was incubated with anti-Myod

antibody and NRIgG antibodies overnight at 4 °C. Next the immune complex was recovered

by protein A agarose/Salmon sperm DNA beads incubation for 4 hr at RT and the beads

were serially washed with low salt, high salt, LiCl and TE buffers. Then the beads were

treated with RnaseA, proteinase K. DNA fragments from the beads and input samples were

purified using PureLink™ Quick Gel Extraction and PCR Purification combo kit (Thermo

Scientific, USA) after decrosslink. Myod target Myog and Myh promoter regions were

detected in these samples by qPCR using primers given in Table S1.

RNA-sequencing

Total RNAs isolated from C2-shScr and C2-shmVcl2 cells were used for RNA-sequencing

analysis. RNA integrity was assessed using an Agilent 2200 TapeStation system (Agilent,

USA) with an RNA Integrity Number (RIN) value greater than or equal to 7. The libraries

were synthesized from the these RNAs using TruSeq Stranded Total RNA LT sample Prep

Kit (Human Mouse Rat) (Illumina, USA) according to manufacturer protocol (TruSeq

Stranded Total RNA Sample Prep Guide, Part #15031048 Rev. E). The paired-end

sequencing was performed in the Illumina NovaSeq 6000 System (Illumina, USA) at

Macrogen, Inc, South Korea and ~100 M reads per sample raw data was obtained. Quality

of the reads is checked with the software FASTQC and then trimming is performed with the

software Trimmomatic[95] setting a minimum base quality of 25 and a minimum length of 35

bp. Trimmed reads are then mapped on the reference genome using STAR[96] and

featureCounts[97] is used to perform gene expression quantification. Lowly expressed genes

are removed using the algorithm HTSFilter[98] and finally differentially expressed genes are

identified using edgeR[99] setting 0.05 as the threshold of FDR significance. The RNA-Seq

data analysis was done using artificial intelligence RNA-Seq (A.I.R.), Sequentia Biotech SL,

Barcelona. The genes showed LogFC more than 1 for upregulated and less than -1 for

downregulated DEGs were used for GO enrichment analysis in ShinyGO 0.80.[100]

Co-immunoprecipitation

The cells were washed with ice cold PBS and collected. The total cellular extracts were

prepared form these cells using RIPA lysis buffer. Total 500 μg of cellular extract was

precleared with protein A-agarose beads and was incubated with anti-vinculin antibody

overnight at 4°C in a rotator. The next day protein A-agarose beads were added and

incubated for 3-4 hr at 4°C to bring down the immune complex. The beads were washed

with four times with cold PBS. This immune complex was further analyzed by western blot

analysis.

Data analysis

Unpaired two-tailed student's t test analyses was performed to ascertain statistical

significance of comparing two mean value. Replicated samples were mentioned in figure

legends as 'n='. Data were shown as mean value ±SEM, p value * ≤0.05; **≤0.01; ***≤0.001;

****≤0.0001. and p < 0.05 was considered to be significant.

Acknowledgements

This work is supported by grants from Science and Engineering Research Board (SERB),

India, file number EEQ/2017/000322, ECR/2016/000265 and University Grants Commission,

India UGC/FRP F.NO 4-5 (95)/2014 (BSR)(FRP) to MJ. The authors thank Dr. Ashok Arasu,

a former graduate student, for his assistance in cloning the metavinculin shRNAs. The

authors greatly acknowledge Dr. Asoke Mal, Dr. Sam J Mathew, Dr. Davide Gabellini for

providing crucial reagents. We thanks Central Animal Research Facility, NIMHANS. AM is a

Senior Research fellow (ICMR-SRF), 45/4/2020-HUM/BMS, from the Indian Council of

Medical Research (ICMR). Graphical figure created with BioRender.com.

Author Contributions

M.J. conceived the project. A.M. and M.J. designed the experiments; A.M., N.B., M.J.

performed the experiments. A.M., N.B., M.J. validated and analysed the data. A.M., M.J.

wrote the manuscript; A.M., M.J., N.B. edited the manuscript; M.J. acquired funding. M.J.

supervised the work. All authors read, corrected and approved the manuscript.

Conflict of Interest

The authors declare no competing interests.

References

1.   Parker, M. H., Seale, P. & Rudnicki, M. A. Looking back to the embryo: defining

transcriptional networks in adult myogenesis. *Nat. Rev. Genet.* **4**, 497–507 (2003).

2.   McKinsey, T. A., Zhang, C. L. & Olson, E. N. Signaling chromatin to make muscle.

Curr. Opin. Cell Biol. **14**, 763–772 (2002).

3.   Penn, B. H., Bergstrom, D. A., Dilworth, F. J., Bengal, E. & Tapscott, S. J. A MyoD-

generated feed-forward circuit temporally patterns gene expression during skeletal

muscle differentiation. *Genes Dev.* **18**, 2348–2353 (2004).

4.   Guasconi, V. & Puri, P. L. Chromatin: the interface between extrinsic cues and the

epigenetic regulation of muscle regeneration. *Trends Cell Biol.* **19**, 286–294 (2009).

5.   Chatterjee, B., Wolff, D. W., Jothi, M., Mal, M. & Mal, A. K. p38α MAPK disables

KMT1A-mediated repression of myogenic differentiation program. *Skelet. Muscle* **6**,

(2016).

6.   Mal, A. K. Histone methyltransferase Suv39h1 represses MyoD-stimulated myogenic

differentiation. *EMBO J.* **25**, 3323–3334 (2006).

7.   Mal, A. A role for histone deacetylase HDAC1 in modulating the transcriptional activity

of MyoD: inhibition of the myogenic program. *EMBO J.* **20**, 1739–1753 (2001).

8.   Bergstrom, D. A. *et al.* Promoter-Specific Regulation of MyoD Binding and Signal

Transduction Cooperate to Pattern Gene Expression. *Mol. Cell* **9**, 587–600 (2002).

9.   Olson, E. N. Interplay between proliferation and differentiation within the myogenic

lineage. *Dev. Biol.* **154**, 261–272 (1992).

10.   Arnold, H. H. & Winter, B. Muscle differentiation: more complexity to the network of

myogenic regulators. *Curr. Opin. Genet. Dev.* **8**, 539–544 (1998).

11.   Castle, J. C. *et al.* Expression of 24,426 human alternative splicing events and

predicted cis regulation in 48 tissues and cell lines. *Nat. Genet. 2008 4012* **40**, 1416–

1425 (2008).

12.   Melé, M. *et al.* The human transcriptome across tissues and individuals. *Science (80-.*

). **348**, 660–665 (2015).

13.   Tapial, J. *et al.* An atlas of alternative splicing profiles and functional associations

738  reveals new regulatory programs and genes that simultaneously express multiple

739  major isoforms. *Genome Res.* **27**, 1759–1768 (2017).

740 14. Gómez-Oca, R. *et al.* Differential impact of ubiquitous and muscle dynamin 2 isoforms

741  in muscle physiology and centronuclear myopathy. *Nat. Commun. 2022 131* **13**, 1–20

742  (2022).

743 15. Sebastian, S. *et al.* Tissue-specific splicing of a ubiquitously expressed transcription

744  factor is essential for muscle differentiation. *Genes Dev.* **27**, 1247–1259 (2013).

745 16. Runfola, V., Sebastian, S., Dilworth, F. J. & Gabellini, D. Rbfox proteins regulate

746  tissue-specific alternative splicing of *Mef2D* required for muscle differentiation. *J. Cell*

747  *Sci.* (2015) doi:10.1242/jcs.161059.

748 17. Cocchiararo, I. *et al.* Identification of a muscle-specific isoform of VMA21 as a potent

749  actor in X-linked myopathy with excessive autophagy pathogenesis. *Hum. Mol. Genet.*

750  **32**, 3374–3389 (2023).

751 18. Merkin, J., Russell, C., Chen, P. & Burge, C. B. Evolutionary dynamics of gene and

752  isoform regulation in mammalian tissues. *Science (80-.).* **338**, 1593–1599 (2012).

753 19. Louis, H. A., Pino, J. D., Schmeichel, K. L., Pomiès, P. & Beckerle, M. C. Comparison

754  of Three Members of the Cysteine-rich Protein Family Reveals Functional

755  Conservation and Divergent Patterns of Gene Expression. *J. Biol. Chem.* **272**, 27484–

756  27491 (1997).

757 20. Freiburg, A. *et al.* Series of Exon-Skipping Events in the Elastic Spring Region of Titin

758  as the Structural Basis for Myofibrillar Elastic Diversity. *Circ. Res.* **86**, 1114–1121

759  (2000).

760 21. Hohmann & Dehghani. The Cytoskeleton—A Complex Interacting Meshwork. *Cells* **8**,

761  362 (2019).

762 22. Müller, P., Langenbach, A., Kaminski, A. & Rychly, J. Modulating the Actin

Cytoskeleton Affects Mechanically Induced Signal Transduction and Differentiation in

Mesenchymal Stem Cells. *PLoS One* **8**, e71283 (2013).

23.   Ray, P. & Chapman, S. C. Cytoskeletal Reorganization Drives Mesenchymal

Condensation and Regulates Downstream Molecular Signaling. *PLoS One* **10**,

e0134702 (2015).

24.   Cui, X. *et al.* F□actin cytoskeleton reorganization is associated with hepatic stellate

cell activation. *Mol. Med. Rep.* **9**, 1641–7 (2014).

25.   Miralles, F., Posern, G., Zaromytidou, A.-I. & Treisman, R. Actin dynamics control

SRF activity by regulation of its coactivator MAL. *Cell* **113**, 329–42 (2003).

26.   Kuwahara, K., Barrientos, T., Pipes, G. C. T., Li, S. & Olson, E. N. Muscle-specific

signaling mechanism that links actin dynamics to serum response factor. *Mol. Cell.*

Biol. **25**, 3173–81 (2005).

27.   Arai, A., Spencer, J. A. & Olson, E. N. STARS, a striated muscle activator of Rho

signaling and serum response factor-dependent transcription. *J. Biol. Chem.* **277**,

24453–9 (2002).

28.   Xie, X. & Percipalle, P. Elevated transforming growth factor β signaling activation in β-

actin-knockout mouse embryonic fibroblasts enhances myofibroblast features. *J. Cell.*

Physiol. **233**, 8884–8895 (2018).

29.   Ziegler, W. H., Liddington, R. C. & Critchley, D. R. The structure and regulation of

vinculin. *Trends Cell Biol.* **16**, 453–60 (2006).

30.   Tozeren, A. *et al.* Vinculin and cell-cell adhesion. *Cell Adhes. Commun.* **5**, 49–59

(1998).

31.   Xu, W., Baribault, H. & Adamson, E. D. Vinculin knockout results in heart and brain

defects during embryonic development. *Development* **125**, 327–37 (1998).

32.   Belkin, A. M., Ornatsky, O. I., Kabakov, A. E., Glukhova, M. A. & Koteliansky, V. E.

Diversity of vinculin/meta-vinculin in human tissues and cultivated cells. Expression of

muscle specific variants of vinculin in human aorta smooth muscle cells. *J. Biol.*

Chem. **263**, 6631–5 (1988).

33.   Belkin, A. M., Ornatsky, O. I., Glukhova, M. A. & Koteliansky, V. E.

Immunolocalization of meta-vinculin in human smooth and cardiac muscles. *J. Cell*

Biol. **107**, 545–53 (1988).

34.   Feramisco, J. R., Smart, J. E., Burridge, K., Helfman, D. M. & Thomas, G. P. Co-

existence of vinculin and a vinculin-like protein of higher molecular weight in smooth

muscle. *J. Biol. Chem.* **257**, 11024–31 (1982).

35.   Saga, S., Hamaguchi, M., Hoshino, M. & Kojima, K. Expression of meta-vinculin

associated with differentiation of chicken embryonal muscle cells. *Exp. Cell Res.* **156**,

45–56 (1985).

36.   Byrne, B. J., Kaczorowski, Y. J., Coutu, M. D. & Craig, S. W. Chicken vinculin and

meta-vinculin are derived from a single gene by alternative splicing of a 207-base pair

exon unique to meta-vinculin. *J. Biol. Chem.* **267**, 12845–50 (1992).

37.   Moiseyeva, E. P. *et al.* Organization of the human gene encoding the cytoskeletal

protein vinculin and the sequence of the vinculin promoter. *J. Biol. Chem.* **268**, 4318–

25 (1993).

38.   Glukhova, M. A. *et al.* Meta-vinculin distribution in adult human tissues and cultured

cells. *FEBS Lett.* **207**, 139–41 (1986).

39.   Witt, S., Zieseniss, A., Fock, U., Jockusch, B. M. & Illenberger, S. Comparative

biochemical analysis suggests that vinculin and metavinculin cooperate in muscular

adhesion sites. *J. Biol. Chem.* **279**, 31533–43 (2004).

40.   Thoss, F. *et al.* Metavinculin: New insights into functional properties of a muscle

adhesion protein. *Biochem. Biophys. Res. Commun.* **430**, 7–13 (2013).

41.   Zieseniss, A. *et al.* Raver1 is an integral component of muscle contractile elements.
814   *Cell Tissue Res.* **327**, 583–94 (2007).

815   42.   Chopard, A., Arrighi, N., Carnino, A. & Marini, J. F. Changes in dysferlin, proteins from
816   dystrophin glycoprotein complex, costameres, and cytoskeleton in human soleus and
817   vastus lateralis muscles after a long-term bedrest with or without exercise. *FASEB J.*
818   **19**, 1722–4 (2005).

819   43.   Zemljic-Harpf, A. E. *et al.* Heterozygous Inactivation of the Vinculin Gene Predisposes
820   to Stress-Induced Cardiomyopathy. *Am. J. Pathol.* **165**, 1033–1044 (2004).

821   44.   Maeda, M., Holder, E., Lowes, B., Valent, S. & Bies, R. D. Dilated cardiomyopathy
822   associated with deficiency of the cytoskeletal protein metavinculin. *Circulation* **95**, 17–
823   20 (1997).

824   45.   Olson, T. M. *et al.* Metavinculin Mutations Alter Actin Interaction in Dilated
825   Cardiomyopathy. *Circulation* **105**, 431–437 (2002).

826   46.   Vasile, V. C., Edwards, W. D., Ommen, S. R. & Ackerman, M. J. Obstructive
827   hypertrophic cardiomyopathy is associated with reduced expression of vinculin in the
828   intercalated disc. *Biochem. Biophys. Res. Commun.* **349**, 709–715 (2006).

829   47.   Vasile, V. C., Ommen, S. R., Edwards, W. D. & Ackerman, M. J. A missense mutation
830   in a ubiquitously expressed protein, vinculin, confers susceptibility to hypertrophic
831   cardiomyopathy. *Biochem. Biophys. Res. Commun.* **345**, 998–1003 (2006).

832   48.   VASILE, V. *et al.* Identification of a metavinculin missense mutation, R975W,
833   associated with both hypertrophic and dilated cardiomyopathy. *Mol. Genet. Metab.*
834   **87**, 169–174 (2006).

835   49.   Kim, H. R., Hoque, M. & Hai, C.-M. Cholinergic receptor-mediated differential
836   cytoskeletal recruitment of actin- and integrin-binding proteins in intact airway smooth

muscle. *Am. J. Physiol. Physiol.* **287**, C1375–C1383 (2004).

50.   Bois, P. R. J., Borgon, R. A., Vonrhein, C. & Izard, T. Structural dynamics of alpha-
actinin-vinculin interactions. *Mol. Cell. Biol.* **25**, 6112–22 (2005).

51.   Rangarajan, E. S. & Izard, T. The Cryogenic Electron Microscopy Structure of the Cell
Adhesion Regulator Metavinculin Reveals an Isoform-Specific Kinked Helix in Its
Cytoskeleton Binding Domain. *Int. J. Mol. Sci.* **22**, 645 (2021).

52.   Kim, L. Y. *et al.* The Structural Basis of Actin Organization by Vinculin and
Metavinculin. *J. Mol. Biol.* **428**, 10–25 (2016).

53.   Krokhotin, A. *et al.* Distinct Binding Modes of Vinculin Isoforms Underlie Their
Functional Differences. *Structure* **27**, 1527-1536.e3 (2019).

54.   Rangarajan, E. S., Lee, J. H., Yogesha, S. D. & Izard, T. A Helix Replacement
Mechanism Directs Metavinculin Functions. *PLoS One* **5**, e10679 (2010).

55.   Lee, H. T. *et al.* Vinculin and metavinculin exhibit distinct effects on focal adhesion
properties, cell migration, and mechanotransduction. *PLoS One* **14**, e0221962 (2019).

56.   Janssen, M. E. W., Liu, H., Volkmann, N. & Hanein, D. The C-terminal tail domain of
metavinculin, vinculin's splice variant, severs actin filaments. *J. Cell Biol.* **197**, 585–
593 (2012).

57.   Oztug Durer, Z. A. *et al.* Metavinculin Tunes the Flexibility and the Architecture of
Vinculin-Induced Bundles of Actin Filaments. *J. Mol. Biol.* **427**, 2782–2798 (2015).

58.   Kuroyanagi, H. Fox-1 family of RNA-binding proteins. *Cell. Mol. Life Sci.* **66**, 3895–
3907 (2009).

59.   Nakka, K., Ghigna, C., Gabellini, D. & Dilworth, F. J. Diversification of the muscle
proteome through alternative splicing. *Skelet. Muscle* **8**, 8 (2018).

60.   Chopard, A., Pons, F. & Marini, J.-F. Vinculin and meta-vinculin in fast and slow rat

skeletal muscle before and after hindlimb suspension. *Pflugers Arch.* **444**, 627–33

(2002).

61.   Zhang, P. *et al.* p21CIP1 and p57KIP2 control muscle differentiation at the myogenin

step. *Genes Dev.* **13**, 213–224 (1999).

62.   Endo, Y., Karvar, M. & Sinha, I. Muscle Cryoinjury and Quantification of Regenerating

Myofibers in Mice. *BIO-PROTOCOL* **11**, (2021).

63.   Singh, R. K. *et al.* Rbfox2-Coordinated Alternative Splicing of Mef2d and Rock2

Controls Myoblast Fusion during Myogenesis. *Mol. Cell* **55**, 592–603 (2014).

64.   Borello, U. *et al.* The Wnt/β-catenin pathway regulates Gli-mediated *Myf5* expression

during somitogenesis. *Development* **133**, 3723–3732 (2006).

65.   Brack, A. S., Conboy, I. M., Conboy, M. J., Shen, J. & Rando, T. A. A Temporal

Switch from Notch to Wnt Signaling in Muscle Stem Cells Is Necessary for Normal

Adult Myogenesis. *Cell Stem Cell* **2**, 50–59 (2008).

66.   Brunelli, S., Relaix, F., Baesso, S., Buckingham, M. & Cossu, G. Beta catenin-

independent activation of MyoD in presomitic mesoderm requires PKC and depends

on Pax3 transcriptional activity. *Dev. Biol.* **304**, 604–614 (2007).

67.   Chen, A. E., Ginty, D. D. & Fan, C.-M. Protein kinase A signalling via CREB controls

myogenesis induced by Wnt proteins. *Nature* **433**, 317–322 (2005).

68.   Gros, J., Serralbo, O. & Marcelle, C. WNT11 acts as a directional cue to organize the

elongation of early muscle fibres. *Nature* **457**, 589–593 (2009).

69.   Ikeya, M. & Takada, S. Wnt signaling from the dorsal neural tube is required for the

formation of the medial dermomyotome. *Development* **125**, 4969–4976 (1998).

70.   Polesskaya, A., Seale, P. & Rudnicki, M. A. Wnt Signaling Induces the Myogenic

Specification of Resident CD45+ Adult Stem Cells during Muscle Regeneration. *Cell*

113, 841–852 (2003).

71.   Rochat, A. *et al.* Insulin and Wnt1 Pathways Cooperate to Induce Reserve Cell

887   Activation in Differentiation and Myotube Hypertrophy. *Mol. Biol. Cell* **15**, 4544–4555

888   (2004).

889   72.   VANAMERONGEN, R. & BERNS, A. Knockout mouse models to study Wnt signal

890   transduction. *Trends Genet.* **22**, 678–689 (2006).

891   73.   von Maltzahn, J., Chang, N. C., Bentzinger, C. F. & Rudnicki, M. A. Wnt signaling in

892   myogenesis. *Trends Cell Biol.* **22**, 602–609 (2012).

893   74.   Peng, X., Cuff, L. E., Lawton, C. D. & DeMali, K. A. Vinculin regulates cell-surface E-

894   cadherin expression by binding to β-catenin. *J. Cell Sci.* **123**, 567–577 (2010).

895   75.   Liu, J. *et al.* A Small-Molecule Agonist of the Wnt Signaling Pathway. *Angew. Chemie*

896   *Int. Ed.* **44**, 1987–1990 (2005).

897   76.   Minami, I. *et al.* A Small Molecule that Promotes Cardiac Differentiation of Human

898   Pluripotent Stem Cells under Defined, Cytokine- and Xeno-free Conditions. *Cell Rep.*

899   **2**, 1448–1460 (2012).

900   77.   Nemeth, M. J., Topol, L., Anderson, S. M., Yang, Y. & Bodine, D. M. Wnt5a inhibits

901   canonical Wnt signaling in hematopoietic stem cells and enhances repopulation. *Proc.*

902   *Natl. Acad. Sci.* **104**, 15436–15441 (2007).

903   78.   Ishitani, T. *et al.* The TAK1-NLK Mitogen-Activated Protein Kinase Cascade Functions

904   in the Wnt-5a/Ca $^{2+}$ Pathway To Antagonize Wnt/β-Catenin Signaling. *Mol. Cell. Biol.*

905   **23**, 131–139 (2003).

906   79.   van Amerongen, R., Fuerer, C., Mizutani, M. & Nusse, R. Wnt5a can both activate

907   and repress Wnt/β-catenin signaling during mouse embryonic development. *Dev. Biol.*

908   **369**, 101–114 (2012).

909   80.   Kühl, M. *et al.* Antagonistic regulation of convergent extension movements in

910   Xenopus by Wnt/β-catenin and Wnt/Ca2+ signaling. *Mech. Dev.* **106**, 61–76 (2001).

911   81.   Relaix, F., Rocancourt, D., Mansouri, A. & Buckingham, M. A Pax3/Pax7-dependent

912   population of skeletal muscle progenitor cells. *Nature* **435**, 948–953 (2005).

913   82.   Kassar-Duchossoy, L. *et al.* Pax3/Pax7 mark a novel population of primitive myogenic

914   cells during development. *Genes Dev.* **19**, 1426–1431 (2005).

915   83.   Sun, S., Zhang, Z., Fregoso, O. & Krainer, A. R. Mechanisms of activation and

916   repression by the alternative splicing factors RBFOX1/2. *RNA* **18**, 274–283 (2012).

917   84.   Kanoldt, V. *et al.* Metavinculin modulates force transduction in cell adhesion sites.

918   *Nat. Commun.* **11**, 6403 (2020).

919   85.   Sarker, M. *et al.* Cardiomyopathy Mutations in Metavinculin Disrupt Regulation of

920   Vinculin-Induced F-Actin Assemblies. *J. Mol. Biol.* **431**, 1604–1618 (2019).

921   86.   Otto, A., Schmidt, C. & Patel, K. Pax3 and Pax7 expression and regulation in the

922   avian embryo. *Anat. Embryol. (Berl).* **211**, 293–310 (2006).

923   87.   Vertino, A. M. *et al.* Wnt10b Deficiency Promotes Coexpression of Myogenic and

924   Adipogenic Programs in Myoblasts. *Mol. Biol. Cell* **16**, 2039–2048 (2005).

925   88.   Le Grand, F., Jones, A. E., Seale, V., Scimè, A. & Rudnicki, M. A. Wnt7a Activates

926   the Planar Cell Polarity Pathway to Drive the Symmetric Expansion of Satellite Stem

927   Cells. *Cell Stem Cell* **4**, 535–547 (2009).

928   89.   Cipriano, A. *et al.* Epigenetic regulation of Wnt7b expression by the cis-acting long

929   noncoding RNA Lnc-Rewind in muscle stem cells. *Elife* **10**, (2021).

930   90.   Brack, A. S. *et al.* Increased Wnt Signaling During Aging Alters Muscle Stem Cell Fate

931   and Increases Fibrosis. *Science (80-.).* **317**, 807–810 (2007).

91.   van der Velden, J. L. J. *et al.* Inhibition of glycogen synthase kinase-3β activity is

sufficient to stimulate myogenic differentiation. *Am. J. Physiol. Physiol.* **290**, C453–
C462 (2006).

92. Noë, S. *et al.* The Myotube Analyzer: how to assess myogenic features in muscle
stem cells. *Skelet. Muscle* **12**, 12 (2022).

93. Jothi, M., Nishijo, K., Keller, C. & Mal, A. K. AKT and PAX3-FKHR cooperation
enforces myogenic differentiation blockade in alveolar rhabdomyosarcoma cell. *Cell*
Cycle **11**, (2012).

94. Jothi, M., Mal, M., Keller, C. & Mal, A. K. Small molecule inhibition of PAX3-FOXO1
through AKT activation suppresses malignant phenotypes of alveolar
rhabdomyosarcoma. *Mol. Cancer Ther.* **12**, (2013).

95. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

96. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21
(2013).

97. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program
for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930
(2014).

98. Rau, A., Gallopin, M., Celeux, G. & Jaffrézic, F. Data-based filtering for replicated
high-throughput transcriptome sequencing experiments. *Bioinformatics* **29**, 2146–
2152 (2013).

99. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. <tt>edgeR</tt>: a Bioconductor
package for differential expression analysis of digital gene expression data.
Bioinformatics **26**, 139–140 (2010).

100. Ge, S. X., Jung, D. & Yao, R. ShinyGO: a graphical gene-set enrichment tool for
animals and plants. *Bioinformatics* **36**, 2628–2629 (2020).

Supplementary information is available at Cell Research's website

Figure Legends

Fig. 1: Metavinculin isoform expression during skeletal muscle differentiation and
regeneration

a Western blot showing metavinculin and vinculin expression in cardiac, smooth and skeletal
muscle tissues. **b** Expression of vinculin isoforms Vcl and mVcl, myogenic genes Myod1,
Myog and Myh expression in C2C12, L6 and HSMM myoblasts grown under growth (GM)
and differentiation (DM) media. **c** RT-qPCR showing the relative expression of vinculin and
metavinculin transcripts in various myoblasts cultured under growth and differentiation
conditions. **d** Western blot showing the vinculin isoform expression at indicated time points
during differentiation. **e** Regenerating muscle tissue sections showing myonecrosis (Black
arrow head), myophagocytosis (White arrow head), inflammation (White arrow) and

regenerating fibers (Black arrow) in different days of post cryoinjury. Scale bars 50 μM. **f**

Metavinculin mRNA expression during in vivo muscle regeneration is shown in a bar graph.

g Western blot showing vinculin isoforms and marker of cell proliferation Ki-67 expression in

regenerating muscle tissues. Actb acted as loading control in western blot images. Data are

shown as mean value ±SEM, n=3. p value * ≤0.05; **≤0.01; ***≤0.001; ns-not significant.

Fig. 2: Rbfox1 expression regulates metavinculin splicing during skeletal muscle

differentiation

a Relative expression of Rbfox family members during in vitro differentiation of C2C12 and **b**

L6 myoblasts are shown in RT-qPCR, n=3. **c** In vitro GST-pulldown assay using total RNAs

isolated from C2C12 cells grown in GM and DM media showing relative enrichment of

Rbfox1 and Rbfox2 with 5' intron flanking metavinculin-specific exon and **d** 3' intron flanking

metavinculin-specific exon. **e** Rbfox1 and Rbfox2 mRNA expression in Rbfox depleted C2-

shrbfx1, C2-shRbfx2 and control C2-Scr cells shown in a bar graph. **f** Metavinculin

expression in Rbfox depleted cells shown by RT-qPCR and **g** western blot analysis. **h**

Crosslink-RIP showing the enrichment of Rbfox1 at upstream and **i** downstream

metavinculin flanking intron in C2-Scr and C2-shRbfx1 cells grown under mentioned

culturing conditions. **j** Western blot image showing Rbfox1 level in Anti-Rbfox1 or NRIgG

control immunoprecipitate along with input samples. **k** RT-qPCR showing Rbfox1

expression in C2-Vec and C2-Rbfox1 cells having exogenous vector or Rbfox1 cDNA at

mRNA and **l** protein levels. **m** RT-qPCR showing Rbfox1 expression in C2-Vec and C2-

Rbfox1 cells having exogenous vector or Rbfox1 cDNA at mRNA and **n** protein levels. **o**

Induction of muscle differentiation-associated genes in growth conditions by ectopic Rbfox1

is shown in a bar graph. **p** Western blot image of myogenic proteins in C2-Vec and C2-

Rbfox1 cells. Actb served as loading control in western blot images. Data are shown as

mean value ±SEM, n=3. p value * ≤0.05; **≤0.01; ***≤0.001; ****≤0.0001; ns-not significant.

Fig. 3: Metavinculin depletion shows defective muscle differentiation

a Metavinculin depletion in C2-shmVcl1, C2-shmVcl2 and C2-Scr cells that stably express shRNAs targeting metavinculin or scramble is shown in RT-qPCR and **b** western blot analysis. **c** Myogenic genes transcripts are reduced under DM conditions in metavinculin depleted C2-shmVcl2 and **d** L6-shmVcl2 cells compared to control C2-Scr or L6-Scr is shown in a bar graph. **e** Western blot showing metavinculin depletion inhibits myogenic proteins induction in C2-shmVcl2 and L6-shmVcl2 cells. **f** Representative images of Myh immunostaining (green) and DAPI counter stained nuclei (blue) in C2-shmVcl2, C2-Scr, L6-shmVcl2, L6-Scr cells in DM conditions, Scale bars 50 μM and bar graph showing fusion index. **g** Western blot showing Myod1, Myog (early) and Myh (late) myogenic markers under growth promoting conditions in C2C12 and L6 cells having ectopic metavinculin or vector. Actb is a loading control in western blot images. Data are shown as mean value ±SEM, n=3. p value * ≤0.05; **≤0.01; ***≤0.001; ****≤0.0001.

Fig. 4: Altered transcriptomic landscape in metavinculin depleted cell

a Differentially expressed genes between metavinculin depleted C2-shmVcl2 and scramble control C2-Scr cells shown in a bar graph and **b** a color scale in the heat map showing normalized expression values. **c** Bar graph showing differentiation-specific expression and validation of randomly picked up and **d** down regulated DEGs by RT-qPCR analysis. **e** Bar plot showing Gene ontology (GO): Biological Process terms significantly associated with up and down regulated DEGs. Data are shown as mean value ±SEM, n=3. p value **≤0.01; ***≤0.001; ****≤0.0001.

Fig. 5: Canonical Wnt-activation negatively affects skeletal muscle differentiation

a Relative expression of canonical Wnt target genes Myc, Ccnd1, Ascl2, Id2, Tcf4 and Mmp9 in C2C12 cells cultured under various conditions. **b** Western blot showing canonical Wnt mediator Ctnnb (β-catenin) protein level in C2-Scr and C2-shmVcl2 cells cultured as

indicated. **c** Co-immunoprecipitation analysis showing direct vinculin and Ctnnb interaction in

C2-Scr and C2-shmVcl2 cells grown under indicated conditions. **d** Bar graph showing RT-

qPCR validation of Wnt target genes activation by canonical Wnt activator BML-284 in

C2C12 cells. **e** Relative expression of myogenic differentiation associated genes in C2C12

cells treated with BML-284 or vehicle in DM media shown in RT-qPCR. **f** Myogenic protein

expression in BML-284 or vehicle treated C2C12 cells cultured under DM conditions is

shown in western blot. **g** Representative images of Myh stain (green) and DAPI (blue) in

C2C12 cells treated with BML-284 or vehicle, Scale bar 50 µM. Fusion index is shown in a

bar graph. Actb act as a loading control in western blot. Data are shown as mean value

$\pm$SEM, n=3. p value * ≤0.05; **≤0.01; ***≤0.001; ****≤0.0001.

1042

Fig. 6: Canonical Wnt-inhibition induces skeletal muscle differentiation program in

metavinculin depleted cells

a RT-qPCR showing the expression of Wnt target genes in C2C12 cells treated with

KY02111. **b** Relative expression of skeletal muscle differentiation marker genes in C2C12

cells treated with KY02111 in growth condition is shown in a bar graph. **c** Western blot

showing the induction of myogenic proteins by KY02111 treatment in C2C12 under growth

conditions. **d** Representative immunofluorescence images showing Myh positive cells

(green) and nuclei (blue) in KY02111-treated C2C12, Scale bars 50 µM and its percentage

is shown in a bar graph. **e** RT-qPCR analysis of myogenic genes in metavinculin depleted

C2-shmVcl2 cells treated with KY02111 in DM conditions. **f** Western blot showing muscle

specific genes in KY02111-treated C2-shmVcl2 cells under DM. **g** Representative images of

Myh staining (green) and nuclear stain DAPI (blue) in metavinculin-depleted C2C12 cells

treated with KY02111 in DM, Scale bars 50 µM. Fusion index is shown as bar graph. Actb

act as a loading control in western blot. Data are shown as mean value $\pm$ SEM, n=3. p value

* ≤0.05; **≤0.01; ***≤0.001; ****≤0.0001.

Figure 7: Ectopic expression of non-canonical Wnt7b enhances skeletal muscle

differentiation in metavinculin depleted cells

a Bar graph showing relative expression of Wnt7b in C2-Scr and C2-shmVcl2 cells cultured

under various conditions. **b** Relative expression of Wnt7b in KY02111-treated C2C12 cells is

shown in a bar graph. **c** Induced Wnt7b protein during C2C12 and L6 cells differentiation is

inhibited by metavinculin depletion as shown in western blot. **d** KY02111-treatment induced

Wnt7b expression at RNA and **e** protein level in C2C12 cells is shown by RT-qPCR and

western blot respectively. **f** Ectopic expression of Wnt7b is verified in RT-qPCR and **g**

western blot analysis. **h** Bar graph showing increased the expression of myogenic marker

genes in C2-shmVcl2 cells by ectopic Wnt7b expression. **i** Western blot showing

upregulation of myogenic proteins by Wnt7b ectopic expression in C2-shmVcl2 cells

compared to vector control. **j** Representative pictures of Myh stain with DAPI counter stain in

C2-shmVcl2 cells that has ectopic Wnt7b or vector, Scale bar 50 μM. Fusion index in these

cells are represented in a bar graph. Actb act as a loading control in western blot. Data are

shown as mean value $\pm$ SEM, n=3. p value * $\leq$0.05; **$\leq$0.01; ***$\leq$0.001; ****$\leq$0.0001.

Figure 7: Possible role of metavinculin in muscle differentiation

Graphical representation of metavinculin expression and its possible role in switching

canonical to non-canonical Wnt signalling during skeletal muscle differentiation.

Supplementary Information

Document S1. Contains Tables S1 primers and oligos, Table S2 antibody details, Table S3

Tau specificity score, Supplementary figure legends and supplementary Fig. S1-S7

Excel spreadsheet: Table S4. Dataset RNA-Seq C2-shScr vs. C2-shmVcl2.

1080

Availability of Data and Materials

All unique/stable reagents generated in this study are available from the corresponding

author (MJ) with a completed materials transfer agreement. The RNA-Seq data that support

the findings of this study have been deposited on the Gene Expression Omnibus (GEO) and

are publicly available with accession number GSE181789. All data reported in this paper will

be shared by the corresponding author upon request.

Fig. 1

Fig. 2

Fig. 3

a

Number of genes
1600
1400 — 1387
1200
1000
800 — 809
600 — 578
400
200
0
DEG Up Down

b

C2-Scr_3
C2-Scr_2
C2-Scr_1
C2-shmVcl2_3
C2-shmVcl2_2
C2-shmVcl2_1

Zscale
2
0
-1

c

● GM
● C2-Scr(DM)
● C2-ShmVcl2(DM)

Relative expression

Klhl13 Heatr9 Cxcl10 Ccl5 Acod1 Slurp1

** ** **** *** **** ****

d

● GM
● C2-Scr(DM)
● C2-ShmVcl2(DM)

Relative expression

Ccn2 Tagln Actg2 Myl9 Car3

**** ** *** ** ***

e

GO: Biological process

Up-regulated genes

Defense response to virus
Defense response to symbiont
Response to lipopolysaccharide
Response to molecule of bacterial origin
Response to bacterium
Response to cytokine
Reg. of defense response
Cellular response to cytokine stimulus
Reg. of response to external stimulus
Inflammatory response
Response to lipid
Response to other organism
Response to external biotic stimulus
Defense response to other organism
Biological proc. involved in interspecies interaction between organis
Reg. of response to stress
Reg. of immune system proc.
Negative reg. of response to stimulus
Positive reg. of cell communication

N. of Genes
● 50
● 70
● 90

-log10(FDR)
10
12
14

Fold Enrichment
0 1 2 3 4

Down-regulated genes

Reg. of muscle contraction
Muscle organ development
Reg. of muscle system proc.
Muscle contraction
Muscle system proc.
Striated muscle cell differentiation
Muscle tissue development
Muscle structure development
Muscle cell differentiation
Reg. of actin filament-based proc.
Actin filament-based proc.
Heart development
Actin cytoskeleton organization
Tissue morphogenesis
Circulatory system development
Anatomical structure formation involved in morphogenesis
Tube morphogenesis
Tube development
Animal organ morphogenesis
Reg. of cell differentiation

N. of Genes
● 20
● 30
● 40
● 50
● 60

-log10(FDR)
7.5
10.0
12.5
15.0

Fold Enrichment
0 2 4 6

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8