Haplotype-resolved genome and population genomics of the threatened garden dormouse in Europe

Paige A. Byerly[1,2#*], Alina von Thaden[1,2#], Evgeny Leushkin[1,3], Leon Hilgers[1,3], Shenglin Liu[1,3], Sven Winter[4,5], Tilman Schell[1,3], Charlotte Gerheim[1,3], Alexander Ben Hamadou[1,3], Carola Greve[1,3], Christian Betz[6], Hanno J. Bolz[6], Sven Büchner[8], Johannes Lang[8], Holger Meinig[8], Eva Marie Famira-Parcsetich[8], Sarah P. Stubbe[8], Alice Mouton[9], Sandro Bertolino[10], Goedele Verbeylen[11], Thomas Briner[12], Lídia Freixas[13], Lorenzo Vinciguerra[14], Sarah A. Mueller[15], Carsten Nowak[1,2], and Michael Hiller[1,3,7*]

[1] LOEWE Centre for Translational Biodiversity Genomics, Senckenberganlage 25, 60325 Frankfurt, Germany
[2] Conservation Genetics Group, Senckenberg Research Institute and Natural History Museum Frankfurt, Clamecystraße 12, 63571 Gelnhausen, Germany
[3] Senckenberg Research Institute, Senckenberganlage 25, 60325 Frankfurt, Germany
[4] Senckenberg Biodiversity and Climate Research Centre, Senckenberganlage 25, 60325 Frankfurt am Main, Germany
[5] Research Institute of Wildlife Ecology, University of Veterinary Medicine Vienna, Savoyenstr. 1, 1160 Vienna, Austria
[6] Bioscientia Human Genetics, Institute for Medical Diagnostics GmbH, Ingelheim, Germany
[7] Institute of Cell Biology and Neuroscience, Faculty of Biosciences, Goethe University Frankfurt, Max-von-Laue-Str. 9, 60438 Frankfurt, Germany
[8] Justus-Liebig-University Giessen, Clinic for Birds, Reptiles, Amphibians and Fish, Working Group for Wildlife Research, Frankfurter Strasse 114, 35392 Giessen, Germany
[9] Socio-économie, Environnement et Développement (SEED), University of Liege (Arlon Campus Environment), Belgium
[10] Department of Life Sciences and Systems Biology, University of Turin, 10123 Torino, Italy
[11] Natuurpunt Studie vzw, Mammal Working Group, Michiel Coxiestraat 11, 2800 Mechelen, Belgium
[12] Naturmuseum Solothurn, Klosterplatz 2, 4500 Solothurn, Switzerland
[13] BiBio Research Group, Natural Sciences Museum of Granollers, C/ Francesc Macià 51, 08402 Granollers, Catalonia, Spain.
[14] Naturmuseum St.Gallen, Rorschacher Strasse 263, 9016 St. Gallen, Switzerland
[15] Division of Evolutionary Biology, Faculty of Biology, LMU Munich, Germany

# contributed equally to the manuscript
* corresponding authors paige.byerly@senckenberg.de, michael.hiller@senckenberg.de

Abstract

Genomic resources are important for evaluating genetic diversity and supporting conservation efforts. The garden dormouse (*Eliomys quercinus*) is a small rodent that has experienced one of the most severe modern population declines in Europe. We present a high-quality haplotype-resolved reference genome for the garden dormouse, and combine comprehensive short and long-read transcriptomics datasets with homology-based methods to generate a highly complete gene annotation. Demographic history analysis of the genome revealed a sharp population decline since the last interglacial, indicating an association between colder climates and population declines prior to anthropogenic influence. Using our genome and genetic data from 100 individuals, largely sampled in a citizen-science project across the contemporary range, we conducted the first population genomic analysis for this species. We found clear evidence for population structure across the species' core Central European range. Notably, our data shows that the Alpine population, characterized by strong differentiation and reduced genetic diversity, is reproductively isolated from other regions and likely represents a differentiated evolutionary significant unit (ESU). The predominantly declining Eastern European populations also show signs of recent isolation, a pattern consistent with a range expansion from Western to Eastern Europe during the Holocene, leaving relict populations now facing local extinction. Overall, our findings suggest that garden dormouse conservation may be enhanced in Europe through designation of ESUs.

Introduction

As genome sequencing technology advances and costs decrease, high-quality reference genomes are becoming increasingly recognized as an essential resource for conservation genomics (Formenti et al. 2022; Brandies et al. 2019; Theissinger et al. 2023; Paez et al. 2022). When aligned to a reference genome assembly, informative markers such as genome-wide single nucleotide polymorphisms (SNPs) generated via high-throughput genomic data can contribute to the understanding of past and contemporary population demographics (Gautier et al. 2016; du Plessis et al. 2023; Luo et al. 2023; Campana et al. 2016), reveal evolutionary patterns such as adaptive differentiation (Szarmach et al. 2021; Martchenko and Shafer 2023), and inform the current status of wildlife species of conservation concern (Talla et al. 2023; Viluma et al. 2022). While reduced-representation sequencing (RRS) methodologies continue to be the more cost-effective and efficient means of generating SNP datasets for large numbers of samples (Wright et al. 2020; Peterson et al. 2012), alignment of RRS reads to a high-quality reference genome improves both the precision of SNP calls and the quantity of SNPs recovered when compared to *de novo* read alignment without a genome assembly (Rochette et al. 2019; Brandies et al. 2019). Reference genomes are also useful for improving data recovery from closely-related species, with cross-species alignment shown to be highly effective (Takach et al. 2023; DeSaix et al. 2019; Burri et al. 2015; Nieto-Blázquez et al. 2022). Until recently, reference genomes for non-model organisms were often generated using short-read sequencing. However, short-read reference genomes are prone to errors such as sequence gaps, false duplicates, and misassembled reads, all of which can lead to inaccuracies in downstream data processing (Rhie et al. 2021). By contrast, genome assemblies generated from long-read sequencing are largely free from these sources of error. This higher accuracy combined with greater contiguity makes long-read sequencing more suitable for genomic analyses such as linking phenotype to genotype, detection of large structural variants, and gene expression analyses (Jebb et al. 2020; Rhie et al. 2021; Karimi et al. 2022; Wold et al. 2021). Given the value of high-quality reference genome assemblies for population and conservation genomic analyses, long-read sequencing of reference genomes for wildlife groups should be a focus of ongoing conservation efforts.

Highly-informative data sources such as reference genomes derived from long-read sequencing can have particular value for investigating understudied wildlife groups, as genomic data can contribute to baseline knowledge needed for conservation planning such as population size estimates and historical changes in demography. Within mammals, such foundational knowledge is lacking for many small-bodied taxa such as Rodentia and Eulipotyphla, which have a known deficit of research (Verde Arregoitia 2016; Kennerley et al. 2021) even in well-studied regions such as Central Europe (Pérez-Espona 2017). Indeed, general population trends of small mammals are not well understood in Europe (Lang et al. 2022; Pérez-Espona 2017; Gippoliti and Amori 2007;

Bertolino et al. 2015), though many species appear to be declining (Rammou et al. 2022; Gippoliti and Amori 2007; Lang et al. 2022; Reiners et al. 2014). Population genomic analysis could expand our understanding of population structure and the factors contributing to declines in Europe, but with the exception of a few flagship species such as the Eurasian beaver (*Castor fiber*; (Halley et al. 2021)) it remains a limited tool in European rodent conservation.

The garden dormouse (*Eliomys quercinus*) is a small rodent species that exemplifies the small mammal conservation crisis in Europe. Once distributed across the continent, the garden dormouse is now recognized to have experienced one of the most extensive modern population declines on the European continent, with an approximate 51% range contraction since the 1970s (Bertolino 2017). Currently, it is only considered common in five of the 26 countries that once comprised its historical range, and even within these refuge countries its distribution is patchy and highly localized (Bertolino 2017). Despite its known imperiled status, research on the garden dormouse has been limited, even in comparison to the three other European species in its family Gliridae (Lang et al. 2022). In particular, the causes of its population decline are still unclear. Changing climate and intensification of land use have been suggested as possible influences; however, the demographic shift appears to have started prior to major 20th century landscape-level changes (Meinig and Büchner 2012; Bertolino 2017). Despite this possible influence of habitat loss, the garden dormouse also shows signs of being a habitat generalist, and this forest species has also adapted well to urban areas in some parts of its range while essentially vanishing from others (Meinig and Büchner 2012; Büchner et al. 2024), making its decline an ongoing mystery.

Population genetic analysis has been identified as a major research need in garden dormouse conservation (Meinig and Büchner 2012; Büchner et al. 2024), as genetic data could aid in understanding patterns of connectivity and gene flow between regions and uncover past population processes that may help explain current declines. Prior karyotyping and mitochondrial DNA analyses suggested the existence of four genetically distinct clades within Europe (Perez et al. 2013; Libois et al. 2012), although the existence of hybrid individuals indicated gene flow between clades (Perez et al. 2013). A better understanding of the population structure and genetic diversity within the hypothesized clades could help guide future management priorities and inform both regional and European-wide conservation efforts. Here, we present a high-quality long-read-based reference genome for the garden dormouse, representing the first such complete genome for both the species and its genus. We then demonstrate the utility of the genome by conducting the first genome-wide analysis of population differentiation across the contemporary range of the garden dormouse.

Results

Haplotype-resolved genome assembly

To generate a reference-quality assembly for the garden dormouse, we collected tissue from a male individual sourced from Mainz, Germany in 2023. We used DNA from this individual to generate 124.4 Gb of PacBio High-Fidelity (HiFi) long-read data, with 117 Gbp in long-range Hi-C read pairs. Assuming a 2.5 Gb genome size, similar to other dormice (Zoonomia Consortium 2020; Böhne et al. 2023), the HiFi and Hi-C data have an estimated coverage of ~50× and ~47×, respectively. With these data, we obtained two haplotype-resolved genome assemblies (Fig. S1).

The contig N50 values of the two haplotypes are 51.7 and 45.4 Mb, respectively. This contiguity is at least 2.8-fold higher compared with other Sciuromorpha genome assemblies (Fig. 1A,B, Table S1). Scaffold N50 values are 108.1 and 107.5 Mb. Haplotype 1 contains the X Chromosome, which was assembled as a chromosome-level scaffold. The Y Chromosome could not be assembled well. We identified 24 autosomes, implying that the sequenced individual has a karyotype of 50 chromosomes (2×24 + 2 sex chromosomes) (Fig. 1C,D), which matches the range of 48 to 54 chromosomes identified for *Eliomys* so far and corresponds to the expected karyotype of the Northern clade where the specimen was sampled (Perez et al. 2013). Overall, 99.4 and 99.6% of haplotype 1 and 2 are contained in chromosome-level scaffolds. These metrics exceed the standards set by the Vertebrate Genome Project (Rhie et al. 2021).

Figure 1: Contiguity and Hi-C maps of both haplotype assemblies.
(A, B) Visualization of contig (A) and scaffold (B) sizes of the garden dormouse haplotype 1 and 2 (red font) and other Sciuromorpha genome assemblies. The N50 values are given on the right side. Contigs shorter than 1 Mb and scaffolds shorter than 5 Mb are not visualized individually, but shown as the gray portion of each bar.
(C, D) Hi-C density maps for haplotype 1 (C) and haplotype 2 (D).

Assembly quality and completeness

Using the HiFi reads, we estimate that our haplotype 1 and 2 assemblies have a high base accuracy with a QV (consensus quality) value of 60.46 and 60.24, indicating only one error per megabase. To assess gene completeness, we first used compleasm (Huang and Li 2023) with the BUSCO odb10 set of 9,226 near-universally conserved mammalian genes. Counting completely detected genes present in a single copy, our assemblies contained 98.3% such genes for haplotype 1 and 90.16% for haplotype 2 that lacks the X Chromosome. In comparison to other Sciuromorpha genome assemblies, our

garden dormouse haplotype 1 ranks second after the *Sciurus carolinensis* assembly
(Mead et al. 2020) (Fig. 2A, Table S1). Second, we used TOGA (Kirilenko et al. 2023) to
compare the status of 18,430 ancestral placental mammal coding genes across the
garden dormouse and other Sciuromorpha assemblies. TOGA explicitly distinguishes
between the two major assembly issues (incompleteness and base errors) by classifying
genes into those that have an intact reading frame, those that have gene-inactivating
mutations (frameshifts, stop codons, splice site mutations, exon or gene deletions) and
those that have missing sequences due to assembly incompleteness or fragmentation.
We found that 94.1% of the ancestral genes are intact and only 1.1% have missing exons
(Fig. 2B, Table S1). In comparison to other assemblies in the order Sciuromorpha, our
garden dormouse haplotype 1 ranks third in terms of intact genes. Despite being top-
ranked in the compleasm metric, TOGA reveals that the *S. carolinensis* assembly has
substantially more genes with inactivating mutations, indicating a lower base accuracy
(Fig. 2B).
193
194

Figure 2: Comparison of gene completeness between our haplotype-resolved garden dormouse and other Sciuromorpha genomes.

(A) Fractions of detected, fragmented, and missing BUSCO genes (mammalian odb10 dataset, 9,226 genes), as classified by compleasm v.0.2.2 (Huang and Li 2023) across Sciuromorpha assemblies.

(B) TOGA classification of 18,430 ancestral placental mammal genes across Sciuromorpha assemblies.

Assemblies are ranked by the number of completely-detected (A) and by the number of intact (B) genes. For the garden dormouse, we show both haplotypes in red font and next to each other, but the X Chromosome is only contained in haplotype 1. Species and assembly accessions are listed in Table S1.

Gene annotation

To comprehensively annotate genes of the garden dormouse genome, we
generated transcriptomics data from eight different organs (heart, testis, liver, lung,
spleen, muscle, kidney, and bladder). We combined ~4.2 billion short RNA-seq and ~1.6
million long Iso-Seq reads to produce transcript models. Transcript models were joined
with a comprehensive homology-based annotation, obtained by first generating genome
alignments and then applying TOGA with human and mouse as the reference. The
resulting annotation comprised 19,374 coding genes and as estimated by compleasm in
protein mode, completely contains 99.91% of the mammalian BUSCO genes, which
indicates a very high completeness.

Long-term demographic history of the garden dormouse

To gain insight into the demographic history of the garden dormouse and investigate
potential effects of climatic changes in the Pleistocene, we inferred the effective
population size (N_e) using pairwise sequentially Markovian coalescence (PSMC) (Li and
Durbin 2011). We used a generation time of 1.5 years and a mutation rate of 5.7×10^{-9} per
generation (see Methods).

Our results indicate relatively high N_e (> 130,000) and consistent population growth
from 800 kya towards the last interglacial (Fig. 3). Population sizes reached their
maximum of > 200,000 individuals around the last interglacial. Subsequently, N_e gradually
declined towards the last glacial maximum, reaching an estimated minimum N_e of less
than 10,000 individuals around 10 kya. Together, this indicates that the garden dormouse
thrived in warm periods, while a cooling climate with longer winters caused severe
population declines. Consistent with this result, the Gliridae family thrived during earlier
warm periods in the Miocene (23 to 5.3 Mya) (Nadachoswki and Daoud 1995).

Figure 3: Demographic history of the garden dormouse.
Effective population size (Ne) of *E. quercinus* as estimated with PSMC. N_e in ten thousands is shown on the y-axis and time in years ago on the x-axis. Faint lines indicate uncertainty of inferred N_e based on 100 bootstrap replicates. Blue and green bars show the timing of the last glacial maximum and last interglacial, respectively. The most recent N_e estimate with reduced accuracy is displayed by a dotted line on the left.

Genotyping and variant calling for population genomics

To investigate population structure across the European range of the garden dormouse, we obtained *n* = 100 samples collected opportunistically between 1991–2020 for population genomic analyses and performed normalized Genotyping-by-Sequencing (nGBS). After sequencing, reads were aligned to the garden dormouse reference genome, and the resulting alignments were used to call single-nucleotide polymorphisms (SNPs). We assigned *a priori* population identity following in part the clades identified by (Perez et al. 2013) (Fig. 4A).

After removal of one individual with 98% missing data and one duplicate sample, we obtained data for a total of 41,175 SNPs from *n* = 98 samples with a mean per site depth of 14.75× and mean frequency of missingness of 0.11 (Table S2). For analyses dependent upon allele frequencies, such PCA-based analyses, we pruned SNPs based on linkage disequilibrium, resulting in a dataset of 19,022 SNPs.

We first inferred relatedness among individual samples and found six pairwise groups of first-order relations in the Harz population. Because inclusion of related individuals can bias population genomic analyses, we randomly removed one individual

from each pair above a kinship threshold > 0.35, which corresponds to a first-order familial
relationship, for subsequent analyses (Table S2). All Russian samples were identified as
second order relations (kinship > 0.20), indicating half-sibling or grandparent-offspring
relationships. Therefore we removed all but one individual from Russia, retaining the
sample with the lowest missing data (Table S2). As all individuals with kinship greater
than the threshold were collected on the same day from the same locations, these kinship
estimates likely reflect accurate estimates of relatedness between individuals. This left us
with a final dataset of n = 86 samples (Table S2). Because we had only one sample for
Italy and Russia, we omitted these samples from further population genetic analyses,
retaining these samples only for investigating population structure.
270

Assessing genetic diversity

We quantified genetic diversity as both SNP-based (polymorphisms only) and
genome-wide (both fixed and variable sites) heterozygosity and found highest genetic
diversity for the Iberian Peninsula and Central regions (Table 1). Within the Central
European region, genetic diversity was lower for population clusters east of the Rhine
River (Northeast, Alps, Harz Mountains) than those located west of or near the river valley
(Northwest, Central). This finding is consistent with the overall lower sizes of garden
dormouse populations located east of the Rhine River in Central Europe (Bertolino 2017).
279

Region	n	polymorphic sites	H_{obs}	H_{exp}	F_{IS}	π	H_{obs_A}
Alps	9	5975	0.036	0.068	0.149	0.075	0.00028
Central	4	13552	0.257	0.273	0.123	0.321	0.00017
Harz Mountains	12	12992	0.223	0.233	0.053	0.245	0.00034
Iberian Peninsula	4	11389	0.249	0.230	0.056	0.280	0.00067
Northeast	7	13173	0.209	0.244	0.124	0.264	0.00036
Northwest	48	18062	0.286	0.344	0.180	0.360	0.00042

Table 1. Sample size (*n*) and genetic diversity metrics among eight garden dormouse sampling
regions with number of SNP-based polymorphic sites, observed (H_{obs}) and expected (H_{exp})
heterozygosity, allelic richness rarified by sample size (A_R), the inbreeding coefficient F_{IS}, and
nucleotide diversity π estimated from 19,022 SNP loci, and observed autosomal heterozygosity
(H_{obs_A}) estimated from from ~90,412 variant and ~58,406,409 fixed sites. Values for the
Northwest region were generated by running four separate analyses of *n* = 12 and then averaging
the output to avoid biases resulting from the uneven sample size between the Northwest (*n* = 48) and
the other regions.

F_{ST} was significantly high (confidence intervals did not overlap 0) between most
regional pairwise groupings (Table 2), suggesting that dispersal between spatially disjunct
populations may be limited. However, the highest overall F_{ST} values were consistently
recovered for pairwise comparisons of the Alpine group and all other populations.

294
295

	NW	ALP	HARZ	CE	IB	NE
NW	—	0.336, 0.344	0.162, 0.168	0.063, 0.068	0.187, 0.195	0.133, 0.139
ALP	0.340	—	0.564, 0.576	0.556, 0.568	0.472, 0.488	0.560, 0.571
HARZ	0.165	0.570	—	0.208, 0.219	0.377, 0.389	0.154, 0.163
CE	0.065	0.562	0.213	—	0.245, 0.259	0.083, 0.093
IB	0.191	0.480	0.383	0.252	—	0.331, 0.344
NE	0.136	0.565	0.159	0.088	0.337	—

297 **Table 2.** Pairwise F_{ST} between six garden dormouse sampling regions below the diagonal and
298 95% confidence intervals based on 1,000 bootstraps above the diagonal, corrected for sample
299 size. Sampling regions are defined and abbreviated as Northwest (NW), the Alps (ALP), Harz
300 Mountains (HAR), Central (CE), the Iberian Peninsula (IB), and Northeast (NE).

Assessing population structure

303   Phylogenetic analysis using RAxML showed that individuals from Northeastern
304 Central Europe (Harz Mountains, Central, Northeast, Russia) form a single clade,
305 separated from the Northwest, Alps, Italy, and Iberian Peninsula regions (Fig. S2) except
306 for a single Central individual (G190084BW), which clustered with the Northwest group.
307 This tree conformed with the PCA of genetic distance among individuals and sampling

regions. PCA showed a strong separation between the Alpine and other sampling regions, with this differentiation explaining 14.48% of the variation on PC1 (Fig. 5). Further spatial differentiation between the Northwest region and the other Central European sampling locations explained 8.19% of the variation on PC2. These results were consistent with findings of relatively high F_{ST} among regions.

Figure 4. Garden dormouse population structure inferred from 47,115 nuclear SNP loci.
(A) Group assignment via five PCs obtained by a discriminant analysis of principal components. The predicted assignment of genetic clusters (*K*) 1–3 is shown by four different colors and the *a priori* sampling region identity is highlighted by dotted gray lines and text. The clades I–IV that were previously identified based on mitochondrial DNA (Perez et al. 2013) are outlined in black.

321 One sample from Russia is not pictured. The Rhine River is outlined in blue, and the boundary of
322 the study area is outlined in gray in the inset map.
323 (B) Genetic clustering inferred by Bayesian structure analysis for *K* ranging from 2 to 5 in
324 STRUCTURE, with x-axes representing geographic sampling location and y-axes the proportion
325 of group membership.
326
327
We used DAPC and STRUCTURE to infer the number of genetic clusters within
our samples. DAPC identified *K* = 4 as the most probable number of genetic clusters (Fig.
S3A), while the results for STRUCTURE were less clear, suggesting *K* = 2–4 as possible
cluster numbers (Fig. S4). Cluster membership for DAPC was suggestive of relatively
strong differentiation between the Northwest region versus other regions in Germany (Fig.
4A, Fig. S3B,C), while STRUCTURE indicated greater admixture among these regions at
all values of *K* (Fig. 4B; Fig. S5). Both methods were consistent in differentiating the
Alpine samples from all other regions (Fig. 4B; Fig. S5). Patterns observed for the full
dataset were comparable for the STRUCTURE analysis subset by sample size (Fig. S6),
indicating that our results were not influenced by unequal sample size among our
sampling regions for regions where n > 1.
339

340
Figure 5. Principal component analysis (PCA) for the garden dormouse.
The PCA is based on 19,022 nuclear SNP loci, with colors representing the geographical origin.
Results are plotted on the first two principal components, with PC1 explaining 14.48% and PC2
8.19% of variation within the dataset. Regions where *n* = 1 (Italy and Russia) are indicated by
black circles, with region names labeled within the plot. One individual from the Alps clustered
separately from the Alpine region and is marked with a *.

Overall, population genomic analyses consistently revealed little evidence for
contemporary gene flow between the Alpine region and other populations, and the high
degree of differentiation in this region may indicate long-term isolation of the Alpine group.
Despite the clear isolation of the Alpine group, all analyses were consistent in separating
one individual from other samples collected from the Alpine region (G190088CH). This
individual, which was collected north of the main Alpine ridge, instead showed strong
evidence for admixture with the Northwest and Harz Mountains regions (Fig. 4; Fig. 5).
354

Recent demographic history inferred from the site-frequency spectrum

To infer recent demographic histories and N_e for each individual population, we
used coalescent simulations and compared three models (Fig. S7): constant steady
population size (null model), population bottleneck (instantaneous size change model),
and constant population growth or decline (incremental size change model). For all
populations, the instantaneous size change (bottleneck) model showed the strongest
support based on the lowest AIC value, although likelihood differences between models
were not great (Table 3). The timing of the bottlenecks was largely consistent among the
Alps, the Harz, and the Northeast, indicating a bottleneck event approximately ~3,000
years ago. The Northwest model presented a slightly more distant bottleneck, at 7,365
years ago, which aligns with the population minimum ~10,000 years ago inferred by
PSMC. For the more northerly populations Harz, Northeast, and Northwest, differences
between N_e before and during the bottleneck were not extensive (Table 3), indicating that
the inferred population change may have been more gradual, which is in line with the
steady decline suggested by PSMC. Despite this, all populations show contrastingly low
contemporary N_e, with the lowest N_e inferred for the Harz and Northeast populations. By
contrast, the top model indicated that the Alps population experienced a catastrophic
decline during the bottleneck, with N_e reduced by 99.99% before recovering to over 2x
ancestral population size.
374
375

Population	Model	N_eCON	N_eANC	N_eBOT	TBOT	ΔL	AIC	ΔAIC
Alps	BOT	128,117	82,321	81	2,193	37,862	249,495	0
Harz	BOT	355	89,848	64,104	1,535	78,191	513,286	0
Northeast	BOT	749	65,464	48,177	2,626	81,403	484,918	0
Northwest	BOT	1,241	57,891	53,781	4,910	141,472	1,160,715	0

Alps	ISC	9,158	47,203	NA	NA	37,882	249,591	95
Harz	ISC	135	179,897	NA	NA	78,194	513,305	19
Northeast	ISC	2,318	24,407	NA	NA	81,403	484,921	2
Northwest	ISC	5,931	112,707	NA	NA	141,481	1,160,762	47
Alps	Null	26,348	NA	NA	NA	37,961	249,947	452
Harz	Null	34,143	NA	NA	NA	78,713	515,688	2,402
Northeast	Null	50,657	NA	NA	NA	81,587	485,760	841
Northwest	Null	90,174	NA	NA	NA	142,192	1,164,026	3,311

Table 3. Model selection results from demographic inference using the Site Frequency Spectrum (SFS) in *fastsimcoal2*. Three models were run for each regional grouping of Alps, Harz, Northeast, and Northwest (Population). Model refers to one of three model types: Null model of constant population size (Null), Instantaneous Size Change/Bottleneck (BOT), and Incremental Size Change (ISC). Parameters were estimated for contemporary N_e (N_eCON), ancestral N_e (N_eANC), N_e during bottleneck (N_eBOT), and time of bottleneck in number of generations (TBOT). Model fit and selection criterion are shown as delta likelihood, calculated as maximum estimated likelihood maximum observed likelihood (ΔL), AIC, and ΔAIC.

Discussion

By generating a long-read-based high-quality reference genome for the garden dormouse, our study provides a genetic resource that can be used to aid conservation efforts of small mammals in Europe. This study was performed in the frame of the citizen-science focused project *Spurensuche Gartenschläfer* ('In Search of the Garden Dormouse'), which is the first national conservation and research project for this species in Germany (Büchner et al. 2024). The project aims to restore garden dormouse populations, as the species continues to decline across much of its range (Lang et al. 2022). Using our long-read genome assembly and population genomics data from samples across the species' range, we were able to investigate garden dormouse demographic history, as well as contemporary population structure and genetic diversity parameters.

The long-term demographic history inferred with PSMC indicates that garden dormice underwent a substantial population bottleneck prior to modern anthropogenic influence, with a sustained decline and long-term low N_e following the last interglacial. Our data suggest that the warm climate around the last interglacial period supported high effective population sizes, and that hibernation did not save garden dormouse populations from declining during the last ice age. Previous paleontological data suggested that Gliridae diversified during periods of high glaciation (Lu et al. 2021). This observation has been interpreted as a sign that this family thrived during cold periods, likely because hibernation was an evolutionary advantage to survive cooling climates (Lu et al. 2021). In contrast, our results provide evidence that garden dormouse populations declined by more than 90% as the climate cooled down towards the last glacial maximum, raising the possibility that the observed lineage diversification during glaciation may have been more likely due to population fragmentation. This finding is in line with longer winters leading to reduced fitness following later emergence from hibernation (Lane et al. 2012). Similar results of declining populations after the last interglacial period have further been described for other Sciuromorpha, including the hibernating Gunnison's prairie dog *Cynomys gunnisoni* (Tsuchiya et al. 2020) and the southern flying squirrel *Glaucomys volans* (Wolf et al. 2022)

More recent demographic history inferred with our SNP dataset further supports a bottleneck history for the individual populations analyzed. While the populations of Harz, Northwest, and Northeast showed evidence for moderate reductions of their ancestral population sizes during the historical declines, estimates of contemporary population size indicated a significant reduction. It is conceivable that the more recent demographic trajectories of these regions were of a complex nature that our models were not able to approximate; however, the low contemporary population sizes as compared to ancestral do reflect what is known about the extensive garden dormouse declines in the 20th and 21st centuries. By contrast, the Alps population showed signs of a massive and severe bottleneck with a more than 99.99% population reduction, followed by a more than two-

fold modern recovery. While a severe bottleneck may in part explain the lower genetic diversity of this region, the magnitude of the decline seems too extreme and the estimated contemporary N_e inappropriately high for the Alpine region, which may indicate that none of the models tested accurately represented the demography of the Alpine population. While the accuracy and precision of *fastsimcoal2* is known to be affected by the number of SNPs analyzed and the amount of missing data (Nunziata and Weisrock 2018), *fastsimcoal2* is also sensitive to population census size, and may not accurately detect population declines when census size is low (Nunziata and Weisrock 2018). It is unclear how these and other factors, such as sample size and temporal heterogeneity, may influence model output (Marchi et al. 2024). The Alps sample set spans a collection period of 19 years, covers a wide geographic area, and exhibits extremely low genetic diversity. These factors may have influenced parameter output for this region and led to upwardly biased N_e. Future analysis with whole genome resequencing is needed to better explore the demographic history of the species, and of the Alpine region in particular.

      Our findings of high genetic differentiation across the garden dormouse's core Central European range and the existence of four genetic clusters confirm and extend prior genetic research, which also identified four garden dormouse lineages within a similar sampling area based on mitochondrial DNA haplotypes and chromosome number differences (Perez et al. 2013; Forcina et al. 2022). As in Perez et al. 2013, we found the Alpine region south of the main Alpine ridge to be strongly differentiated from all other sampling regions, as evidenced in our study by both model-based (STRUCTURE), and non-model-based (PCA) approaches. This strong signal of genetic differentiation displayed by the Alpine population further confirms prior mtDNA genetic distance estimates and karyotypic divergence between the Alps and other regions across the garden dormouse's range ($2n$ = 52 "eastern Alps" and 54 "western Alps") (Perez et al. 2013). One specimen collected within our Alps boundary showed admixture with more Northern regions, but this specimen was collected north of the main Alpine ridge and likely does not belong to the highly differentiated Alpine population. We additionally found the regions south of the Alpine range to be differentiated from those north of the range, which supports that this mountain range may act as a barrier to garden dormouse dispersal (Perez et al. 2013). In contrast to previous research, our more comprehensive datasets also allowed us to detect additional moderate population structure within the Central European clade (Fig. 4), showing a clear pattern of differentiation between eastern and western Central Europe.

      Combined with previous findings of reciprocal mtDNA monophyly (Perez et al. 2013), the significant differentiation of nuclear loci found in this study supports the designation of the Alpine garden dormouse population as an evolutionary significant unit (ESU) according to the classic conservation genetics definition (Moritz 1994). Within the Alpine region, the garden dormouse is found in both subalpine and montane habitats, where it occupies coniferous as well as deciduous forest habitat (Bertolino 2017). The

strong genetic differentiation and distinct habitat suggests the potential for long-term
genetic separation, possibly arising from adaptation to local habitat conditions or climate.
Future research should focus on investigating potential adaptive differences with whole
genome resequencing. Although the garden dormouse has been largely reported as
demographically stable in the Alps (Bertolino 2017), our findings of low genetic diversity
and high rates of inbreeding are indicative of small population sizes, and may indicate
that population declines in this area have been greater than previously thought. Hence,
the status of this population should be reevaluated. Perez et al. (2013) also found lower
nucleotide diversity for this region via mtDNA analyses, and surmised that reductions in
genetic diversity may have resulted from population expansion following a genetic
bottleneck, which our results also support. The proportional differences in genetic
diversity parameters between the Alpine and other populations and other sampling
regions reported in this study are comparable to those seen in comparisons between
mainland and both oceanic and montane "habitat" island populations of other threatened
small mammalian species (Rick et al. 2023; White and Searle 2007; Ditto and Frey 2007),
providing further support for the Alpine population as an isolated, genetically distinct unit
that may require the implementation of separate conservation strategies.
The range contraction of the garden dormouse is known to be centered in the
eastern part of its range, with the species becoming more common and populations more
contiguous as one moves further west (Bertolino 2017; Meinig and Büchner 2012). Fossil
evidence indicates that the garden dormouse originated on the Iberian Peninsula and
spread out into Central Europe during the Holocene (Mansino et al. 2014), which may
mean that remaining garden dormice in Eastern Europe represent small, relict
populations on the edge of extirpation (Anděra 1986). This is somewhat supported by the
higher autosomal genetic diversity and lower inbreeding estimate we detected for the
westernmost Iberian clade, which reflects a larger and more connected population.
However, while the garden dormouse is still regarded as relatively common in some parts
of the Iberian Peninsula (Bertolino 2017), extensive declines and local extirpations have
been noted in some parts of Portugal and Spain, and the distribution has been
characterized as patchy and scattered across the region (Santoro et al. 2017; Vale-
Gonçalves and Mira 2023; Pita et al. 2021). A comparative analysis using the entire
mitogenome found that the Iberian clade exhibits relatively lower genetic diversity as
compared to that of Central Europe, with the most genetically diverse individuals instead
centered in France, possibly resulting from admixture following emergence from glacial
refugia in central Europe (Forcina et al. 2022). Given that samples for our study were
collected from the northernmost boundary of the Iberian Peninsula, our characterization
of the Iberian clade may therefore be more reflective of this contiguous core population
rather than of the wider peninsula. This sampling distribution likely also explains the
minimal genetic structuring we found between the Iberian Peninsula and the central

European regions, which contrasts with the strong differentiation shown in mitochondrial
studies with broader sampling distributions (Forcina et al. 2022; Perez et al. 2013).
Given the known contractions and declines in the eastern edge of the garden
dormouse's range, it is likely that the lower genetic diversity of the eastern populations in
our study is reflective of genetic drift resulting from modern isolation of these populations.
Support for this is provided by our observation of reduced genetic diversity and greater
rates of inbreeding for the more easterly populations in Germany (Harz mountains and
Northeast), whereas the Central population shows evidence of admixture with the larger
and more contiguous Northwest population, thereby potentially increasing its genetic
diversity through interbreeding. Larger sampling efforts are required to substantiate these
preliminary findings. Notably, a similar east-west pattern of differentiation has also been
observed for the hazel dormouse (*Muscardinus avellanarius*), which overlaps the garden
dormouse range and is distributed as two separate lineages or 'partial ESUs', delineated
along a similar geographical divide to the garden dormouse (Leyhausen et al. 2022;
Mouton et al. 2017). Unlike the hazel dormouse, however, this east-west pattern of
differentiation was not reflected by mtDNA in the garden dormouse (Perez et al. 2013),
indicating that the division among lineages may be a more recent phenomenon resulting
from habitat fragmentation and/or demographic changes such as population declines.
Although the causes of the garden dormouse's east-to-west pattern of range
contraction are not fully understood, landscape level changes driven by land use and
climate change have been partly implicated (Bertolino 2017). However, it is unclear what
specific role climate might play in this contraction. For the ecologically-similar hazel
dormouse, warmer and wetter winters have been found to have a negative effect on adult
survivorship, possibly because warmer temperatures cause animals to wake during
hibernation, thereby expending energy and reducing fat reserves (Combe et al. 2023).
Garden dormice hibernate in regions of harsh winter, and their life histories appear
strongly tied to winter temperatures, as well as the onset and duration of the season
(Bennett and Richard 2021; Mahlert et al. 2018). Timing and duration of activity periods
and hibernation vary widely based on local climate conditions, with Mediterranean
populations hibernating only 1–2 months per year, if at all (Viñals-Domingo et al. 2020),
while Alpine populations may hibernate for 7 months per year (Bertolino et al. 2001).
Further investigation of the role of climate in shaping the garden dormouse's population
structure via local adaptation may be useful in predicting the future viability of the species
with ongoing climate change.
The garden dormouse is currently listed as "Near Threatened" in the International
Union for Conservation of Nature (IUCN) Red List (Bertolino et al. 2008), but its status
will be changed to "Vulnerable"(Bertolino 2017)in the next assessment (Bertolino 2017)
due to the observed continuous decline in the species' range. Our results support this
change in status, and suggest that the garden dormouse in Europe is threatened due to
population isolation and low regional genetic diversity, especially at the eastern edge of

its range. Hence, intervention is needed to mitigate further population extirpations. We
further provide strong evidence for the designation of the Alpine subpopulation of the
garden dormouse as a separate ESU. Further research is needed to determine if local
adaptation is driving the genetic differentiation of this and other isolated garden dormouse
populations, and how these potential adaptations may influence future population viability
in the face of a changing climate. Our annotated genome will enable comparative
analyses addressing open evolutionary questions of the *Eliomys* species complex, as well
as inform specific conservation planning and management for the garden dormouse in
Europe. We hope that the findings outlined in this study can help guide future research
for the garden dormouse, while highlighting the need for further protection within regions
of population isolation and/or low genetic diversity.
557
558
559

Materials and Methods

Sample collection

Tissue samples for nGBS were collected from across the European distribution in
the frame of a larger conservation project on *Eliomys quercinus* in Germany. Samples
originated from Germany (n = 66), other Central European countries (n = 25), and Russia
(n = 7) and were collected between 1991-2020 (see Table S2). All samples were taken
opportunistically from dead-found individuals brought to museums or animal shelters,
during monitoring, or during citizen-science activities, and all in compliance with
respective local and national laws.

Genome sequencing

To sequence a high-quality genome, we used a male individual collected in March
2023 in Mainz that was euthanized by a veterinarian due to injuries. A voucher specimen
for this individual was prepared and is housed at the Zoological Research Museum
Alexander Koenig under catalog number ZFMK MAM 2024.0263. DNA extraction, PacBio
and Hi-C sequencing followed standard protocols that are described in the
Supplementary Methods.

Transcriptome sequencing

580   We generated RNA- and Iso-Seq transcriptomics data from the same individual,
581   following standard protocols that are described in the Supplementary Methods.

Genotyping-by-Sequencing

584   Tissue samples for normalized genotyping-by-sequencing (nGBS) were extracted
585   using the DNeasy Blood & Tissue Kit (Qiagen) with slight protocol modifications to

increase yield of RNA-free genomic DNA as described in (Leyhausen et al. 2022). Extracts were quantified with fluorometry (Qubit) and spectrophotometry (NanoDrop) and screened for DNA integrity using gel electrophoresis to identify samples of sufficient quality for high-throughput sequencing. Selected samples (n = 100) with ≥ 300 ng of high molecular weight DNA were sent for nGBS (Restriction Associated DNA sequencing [RAD-Seq]) at LGC Genomics GmbH (www.biosearchtech.com). Briefly, nGBS methodologies included enzymatic digestion with the enzyme MslI; purification; barcoding, library preparation, and PCR amplification with Illumina TruSeq amplification primers; size selection for 300–500 bp with the BluePippin system; and sequencing on an Illumina NextSeq 500/550. Sequences were then demultiplexed in house by LGC Genomics using proprietary software. Further details on enzymatic digestion, library preparation and nGBS protocols can be found in (Leyhausen et al. 2022; Arvidsson et al. 2016).

Genome assembly

We produced haplotype-specific assemblies from HiFi reads using hifiasm (v.0.19.5), integrating Hi-C reads for phasing (Cheng et al. 2021). Both haplotypes were subject to filtering procedure using Foreign Contamination Screen Tool (FCS) (Astashyn et al. 2024). Contamination free assemblies were then polished to remove unambiguous heterozygous sites. Specifically, we mapped HiFi reads back to each assembly with minimap2 v.2.26 (Li 2018), removed duplicates using Picard MarkDuplicates tool v.3.1.0 (http://broadinstitute.github.io/picard), and called variants using DeepVariant v1.5.0 (Poplin et al. 2018). We then filtered for sites with genotype 1/1 and a 'PASS' filter value, meaning that all or nearly all reads support an alternative sequence at this position and passed DeepVariants internal filters. Finally, we corrected corresponding nucleotide sites in the assembly using BCFtools consensus v.1.13 (Li 2011).

To scaffold the polished haplotype assemblies, we first mapped the Arima Hi-C reads to each of the two assemblies independently using chromap v.0.2.5 (Zhang et al. 2021) and then processed mapped reads with yahs v1.1a (Zhou et al. 2023). Finally, we performed manual curation of the two scaffolded haplotypes jointly analyzing them in PretextView v.0.2.5 (https://github.com/wtsi-hpag/PretextView). To this end, we remapped the Hi-C reads to both haplotype assemblies concatenated together using again chromap but allowing for multi-mapping reads (-q 0) to avoid discarding information in regions identical between two haplotypes. We also identified telomeric sequences with tidk v.0.2.31 (https://github.com/tolkit/telomeric-identifier) and where necessary corrected wrong contigs orientations to have telomeres in the ends of resulting scaffolds. To finalize changes made via manual dual haplotype curation, as recently proposed by the Vertebrate Genome and Darwin Tree of Life Project, we used the rapid curation framework (https://gitlab.com/wtsi-grit/rapid-curation/-/tree/main) from the Genome Reference Informatics Team (Howe et al. 2021).

626   To estimate the assembly base accuracy, we used Merqury (Rhie et al. 2020) with
627   the HiFi reads. It should be noted that the QV estimates represent an upper bound as
628   they are based on the HiFi reads used for assembly. To assess gene completeness, we
629   used compleasm 0.2.2 (Huang and Li 2023) with the BUSCO odb10 set of 9,226 near-
630   universally conserved mammalian genes and TOGA (below).
631

Repeat masking and whole-genome alignment

RepeatModeler version 2.0.4 (default parameters) was used to generate a *de novo*
repeat library for the garden dormouse assembly. RepeatMasker 4.1.0 (parameters -
engine ncbi) was used with the resulting library to soft-mask the genome.
We followed our previous workflow to align the human (hg38 assembly) and the
mouse (mm10) assembly to both garden dormouse haplotype assemblies (Sharma and
Hiller 2017; Blumer et al. 2022). Briefly, we used LASTZ version 1.04.15 with sensitive
parameters (K = 2400, L = 3000, Y = 9400, H = 2000, and the LASTZ default scoring
matrix) (Harris 2007), axtChain (default parameters except linear-Gap=loose) to compute
co-linear alignment chains (Kent et al. 2003), RepeatFiller (default parameters) to capture
additional alignments between repetitive regions (Osipova et al. 2019), and chainCleaner
(default parameters except minBrokenChainScore = 75,000 and -doPairs) to improve
alignment specificity (Suarez et al. 2017).
645

Gene annotation

We used TOGA version 1.1.4 (Kirilenko et al. 2023) with the alignment chains and
the human GENCODE 38 and the mouse GENCODE M25 annotation (Frankish et al.
2021) to annotate coding genes. We ran TOGA for both garden dormouse haplotypes.
Raw RNA-seq data were processed with fastp v0.23.4 (Chen 2023; Chen et al.
2018) to remove adapters and low quality (< Q15) bases. Processed reads were mapped
to a first haplotype assembly with HISAT2 v.2.2.1 (Kim et al. 2019). For Iso-Seq reads a
circular consensus was generated from the subreads using ccs v.6.4.0
(https://github.com/PacificBiosciences/pbbioconda). Adapter trimming and base quality
filtering was performed with lima v2.2.0
(https://github.com/PacificBiosciences/pbbioconda). Processed reads were cleaned from
polyA tails and polymerase switching artifacts using 'refine' command from the Iso-Seq
package v.4.0.0 (https://github.com/PacificBiosciences/IsoSeq/blob/master/isoseq-
clustering.md) and then clustered together and aligned to generate consensus reads
using the 'cluster' command from the Iso-Seq package. Additionally, reads were filtered
for alignment quality using a custom Perl script (Supplementary Code). To combine
transcriptome information from RNA-seq and Iso-Seq, we then used a hybrid option (--
mix) in StringTie v.2.1.2 (Shumate et al. 2022).
To integrate transcriptomics and homology-based transcript evidence, we first
merged human and mouse TOGA, keeping only transcripts with an intact reading frame.

We then added transcripts classified as partially intact and uncertain loss, but only if they
do not overlap intact transcripts. The RNA- and Iso-Seq merged transcriptome data was
then used to add UTRs to TOGA-annotated coding transcripts with a compatible exon-
intron structure. Finally, we added transcriptome data for genomic loci that don't have a
TOGA transcript prediction to incorporate lineage-specific and non-coding transcripts into
the annotation.
672

Demographic history inferred from the reference genome

We used PSMC (Li and Durbin 2011) to infer the demographic history of the garden
dormouse across the past one million years. Minimap2 v.2.26 (Li 2018) was used to map
the PacBio HiFi reads to haplotype 1. Heterozygous sites were called with DeepVariant
v1.2.0 (Poplin et al. 2018) using the PacBio model. A consensus genome sequence
where the heterozygous sites were represented with IUPAC ambiguity codes was
generated using bcftools v1.17 (Danecek et al. 2021). The consensus sequence was
converted to the input format of PSMC by applying a bin size of 50 bp.

For PSMC, we included data from all chromosome-size scaffolds except for the X
chromosome (scaffold 4). PSMC was run using the parameter settings *-N25 -t15 -r5 -p*
"4+25*2+4+6". We also explored including more parameters (*-p "4+45*2+4+6"* and *-p*
"4+65*2+4+6") as well as splitting the first parameter (e.g., *-p "2+2+25*2+4+6"*), as
recommended by (Hilgers et al. 2024). None of these settings changed the general
results, so we chose to use the default parameters.

Finally, we calibrated PSMC results using a generation time of 1.5 years and a mutation
rate of $5.7*10^{-9}$ per generation. This mutation rate is based on the assumption that garden
dormice exhibit a similar mutation rate as mice, for which the rate is well established
(Uchimura et al. 2015; Milholland et al. 2017). The mean generation time (1.57 years)
was calculated based on a yearly reproductive cycle starting at age one, a maximum age
of 5 years, and an average yearly survival rate of 0.38 (Schaub and Vaterlaus-Schlegel
2001).
694

Population genomics
Genotyping, Filtering and Variant Calling

Full methods for processing the raw GBS data, including trimming and alignment,
are described in the Supplementary Methods. Variant calling was performed with the
ref_map.pl pipeline for reference aligned reads in Stacks v2.65 (Rochette et al. 2019) with
additional filtering parameters given for the populations program: minor allele frequency
of 5%, a minimum percentage of individuals per population of 80, and writing one random
SNP per locus to remove strong physical linkage disequilibrium (-X "populations: -r 0.80
--min-maf 0.05 --write-random-snp"). Additional variant filtering was performed with
VCFtools v0.1.16 (Danecek et al. 2011) to remove indels and the X Chromosome and
only keep biallelic SNPs with a depth per individual between 6 and 50 (flags: '--remove-

indels --min-alleles 2 --max-alleles 2 --min-meanDP 6 --max-meanDP 50 --minDP 6 --
maxDP 50'). After filtering, we removed samples with > 80% missing data to generate our
final VCF file.
Past mitochondrial DNA work has classified four distinct garden dormouse clades
within its European range (Iberian, Italian, Western European, and Alpine; (Perez et al.
2013). We followed these clade designations in assigning *a priori* population identity but
expanded the 'Western European' category by delineating individuals within this clade as
Northwest, the Harz Mountains, Northeast, Central, and Russia (Fig. 4A) as aligning to
known garden dormouse distribution within this region (Meinig and Büchner 2012).
Because linkage disequilibrium (LD) can influence some analyses such as principal
components-based analyses (PCAs), we created a pruned SNP set using the +prune
plugin in BCFtools (Danecek et al. 2021) using an LD threshold of $r^2 < 0.5$ and a sliding
window of 1000, which was applied to analyses that rely on allele frequencies for
population inferences. We hereafter refer to the 'full' and 'pruned' SNP sets.
Excessive numbers of related individuals can bias some population genomics
analyses ((O'Connell et al. 2019); (Wang 2018)). Because our sample collection was not
always random, we anticipated that several individuals within our sample set were related,
and therefore estimated pairwise relatedness within sampling regions with the full SNP
set pruned by $r^2 < 0.2$ using the KING-robust method ((Manichaikul et al. 2010) in
SNPRelate. We then removed 1 from each pair with a kinship estimate > 0.35, which
corresponds to a first-order familial relationship (sibling pair, parent-offspring). This
sample set was used for all subsequent analyses.
728
Genetic Diversity
Genetic diversity estimates can be biased by a number of factors, including allele
frequencies, unbalanced sample size, and missing data (Schmidt et al. 2021). We
calculated observed (H_{obs}) and expected (H_{exp}) heterozygosity, nucleotide diversity (π),
number of polymorphic loci, and the inbreeding coefficient F_{IS} with the pruned SNP set
using the 'population' function in stacks (Rochette et al. 2019). Because biases due to
unbalanced sample sizes can confound comparisons among populations, we additionally
report H_{exp} derived from genotype-wide autosomal data, which is not subject to the same
bias (Schmidt et al. 2021; Sopniewski and Catullo 2024) Autosomal heterozygosity was
generated from the aligned Stacks catalog using the 'population' function with a minor
allele count of 2 to omit singletons and with one random SNP selected per locus. To
further minimize the effects of differences in sample size, we calculated SNP-based
genetic diversity parameters for the Northwest population in four separate runs of *n* = 12
samples per run and report the mean values. We investigated genetic differentiation by
calculating pairwise values of Weir and Cockerhams' unbiased estimator of F_{ST} (Weir and
Cockerham 1984) in the R package hierfstat (Goudet and Goudet 2014) using the pruned
SNP set and with 95% confidence intervals generated with 1,000 bootstrap iterations.

Population Structure

We investigated phylogenetic distance among samples and groups using the full SNP set by inferring a phylogeny in RAxML version 8.2.12 (Stamatakis 2014) using the ASC-GTRCAT model with Lewis correction for ascertainment bias with 100 bootstrap replicates. We plotted the tree in the R 4.3 (R Core Team 2018) package GGTREE (Yu et al. 2017) with a customized script formulated for SNP marker data (Severn-Ellis et al. 2020).

To explore population structure, we first used the pruned SNP set to visualize population clustering via PCA in the R package *adegenet* (Jombart and Bateman 2008)). We estimated the most probable number of genetic clusters K within our sample set using the full SNP set. First, we implemented a discriminant analysis of principal components (DAPC) using the 'find.clusters' function in *adegenet* ((Jombart and Bateman 2008)) to estimate optimal number of *K* testing *K* = 1–8. The most probable number of clusters was identified as the one with the lowest Bayesian Information Criterion (BIC) value. To minimize overfitting, we first determined the optimal number of principal components (PCs) using cross-validation and then ran DAPC retaining the optimal number of PCs and using the identified optimal number of genetic clusters as *K* to predict group assignment for each sample without *a priori* population information. Five principal components (PCs) were retained to explain approximately 52% of the total retained variation without our dataset.

We also used STRUCTURE v2.3.4 (Pritchard 2000) within the program *Structure_threader* (Pina-Martins et al. 2017) to estimate the value of *K*. We ran STRUCTURE with the admixture model and no prior population information, with an optimal burn-in of 10^4 steps and 10^4 additional steps, and for values of *K* from 1 to 8 with 10 replicates per iteration of *K*. We used the R package pophelper (Francis 2017) to align runs, plot output, and determine the most likely value of *K* via both ΔK ((Evanno et al. 2005) and mean probability (MeanLn*P(K)*). STRUCTURE is known to be sensitive to differences in sample size in populations ((Puechmaille 2016), which was a factor in our dataset. To eliminate the potential effects of sampling bias on our results, we also ran STRUCTURE using the same conditions for a random subset of *n* = 5–6 individuals per population, omitting the Russian and Italian samples from the dataset, for values of *K* from 1 to 5.

Recent Demographic History Inferred from the Site-frequency Spectrum

We inferred demographic history among our individual populations using a sequential Markovian coalescent approximation approach in *fastSimcoal2* v2.08 (Excoffier and Foll 2011; Excoffier et al. 2021). We omitted the Central and Iberian Peninsula populations and the Italian and Russian samples due to their low sample sizes. To generate the input site frequency spectrum (SFS) files, we used the distribution of

minor allele frequencies in the full SNP set to create a folded SFS for each population with the R program dartR 2.9.7 (Gruber et al. 2018), using nearest-neighbor imputation to replace missing data. While *fastSimcoal2* can be used to infer complex demographic co-histories, we opted to test models for each population separately to avoid model overparameterization, given our small sample sizes (Alps, n = 9; Harz, n = 12; Northeast, n = 7; Northwest, n = 48). For each population, we compared three models: "null", "instantaneous size change" (a.k.a., bottleneck), and "incremental size change". The null model assumed a constant population size over time and inferred only one parameter, contemporary effective population size (N_e). The instantaneous size change model assumed a rapid population decline and inferred four parameters: ancestral N_e, time of the bottleneck (in generations), population size during the bottleneck, and contemporary N_e. The incremental size change model assumed either steady population growth or decline over time and inferred five parameters: ancestral N_e, time of ancestral N_e (in generations), population growth rate (with a negative value indicating a positive growth rate), rate of change of N_e, and contemporary N_e. For all models, we assumed a generation time of 1.5 years and a mutation rate of 5.7×10^{-9} . Parameter ranges were uniformly distributed between 10 and 10^5 We ran 100 independent iterations of each model per population with 70 EMC cycles and 1×10^6 simulations per model run. We selected the best run within each model as the one with the highest likelihood (smallest difference between maximum likelihood and obtained likelihood) using the custom script fsc-selectbestrun.sh (https://raw.githubusercontent.com/speciationgenomics/scripts/master/fsc-selectbestrun.sh). Because maximum likelihood can be influenced by parameter size, model selection within each population was then performed using Akaike Information Criterion (AIC) comparisons, which account for the number of parameters in the selection process, with AIC calculated using the custom script calculateAIC.sh (https://github.com/speciationgenomics/scripts/blob/master/calculateAIC.sh).

Data Access

Genomic and transcriptomics read data, our annotated genome and the trimmed ddRAD reads have been submitted to the NCBI BioProject database (https://www.ncbi.nlm.nih.gov/bioproject/) under accession number PRJNA1142780. The haplotype assemblies, gene and transposable element annotations are available for download at https://genome.senckenberg.de/download/GardenDormouse/. TOGA annotations are available at https://genome.senckenberg.de/download/TOGA/. We also provide a genome browser showing our genomes and all annotations at https://genome.senckenberg.de/.

Code Access

825 Most of our analyses relied on publicly available tools and methods. New scripts are
826 available at https://github.com/pabyerly/TBG_GardenDormouse and as Supplementary
827 Code.
828

Competing interests

The authors have no competing interests.

Acknowledgment

This work was supported by the LOEWE-Centre for Translational Biodiversity Genomics (TBG) funded by the Hessen State Ministry of Higher Education, Research and the Arts (LOEWE/1/10/519/03/03.001(0014)/52). Funding was also provided by the German Federal Agency for Nature Conservation (BfN) with resources from the Federal Ministry for the Environment, Nature Conservation, Nuclear Safety and Consumer Protection (BMUV), as part of a German-wide project led by the Bund für Umwelt und Naturschutz BUND (Friends of the Earth Germany), Justus Liebig University Gießen, and the Senckenberg Gesellschaft für Naturforschung in the Federal Programme for Biological Diversity called 'In Search of the Garden Dormouse'.

We are very grateful to many people who have helped with sample collection: Peter Adamik, Hermann Ansorge, Jonas Astrin, Simon Capt, Stefanie Jessolat, Jeroen van der Kooij, Ines Leonhard, Gernot Segelbacher, David Stille, Adria Viñals Domingo, Frank Zachos, Alain Frantz, Olga Grigoryeva, Maurice La Haye, Roel Baets, Kasper Van Acker, Caren Raditz, Lutz Nielen, Stefanie Erhardt, Jürg Paul Müller, Andrea Krug, Danilo Hartung, Christine Thiel-Bender, Ferry Böhme, Julia Hofmann, Sarah Beer, Jörg und Elli Hacker, Otfried Wüstemann, Sonja Klein, Thomas Doebel, Martina Klotz, Hartmut Schmid, the Schwarzwald National Park, the Gran Paradiso National Park, the Museo di Scienze Naturali dell'Alto Adige and Museo di Storia Naturale La Specola di Firenze, Antoni Arrizabalaga, Natural Sciences Museum of Granollers, as well as numerous students and citizen scientists without whom this work would have not been possible.

Author contributions: A.T., C.N., M.H. designed and supervised the study, labwork and data analysis. C.G., A.B.H., C.G., C.B, and H.J.B. performed labwork. P.A.B., E.L., L.H., S.L., S.W., and T.S. conducted statistical and bioinformatic analysis. S.B., J.L., H.M., E.M.F-P., S.P.S., A.M., S.B., G.V., T.B., L.F., L.V., and S.A.M. contributed samples, supervised sample collection, and/or provided expert species knowledge. P.A.B., A.T., E.L., L.H., and M.H. wrote the manuscript with input from all authors.

References

Anděra M. 1986. Dormice (Gliridae) in Czechoslovakia. Part I. : Glis glis, Eliomys quercinus (Rodentia: Mammalia). *Folia Musei Rerum Naturalium Bohemiae Occidentalis, Plzeň - Zoologica* **24**: 3–47.

Arvidsson S, Fartmann B, Winkler S, Zimmermann W. 2016. Technical note: Efficient high-throughput SNP discovery and genotyping using normalised genotyping by sequencing

(nGBS). https://biosearch-cdn.azureedge.net/assetsv6/efficient-high-throughput-snp-
discovery-genotyping-ngbs-app-note.pdf (Accessed January 6, 2024).

Astashyn A, Tvedte ES, Sweeney D, Sapojnikov V, Bouk N, Joukov V, Mozes E, Strope PK,
Sylla PM, Wagner L, et al. 2024. Rapid and sensitive detection of genome contamination at
scale with FCS-GX. *Genome Biol* **25**: 60.

Bennett D, Richard FJ. 2021. Distribution modelling of the garden dormouse Eliomys quercinus
(Linnaeus, 1766) with novel climate change indicators. *Mamm Biol* **101**: 589–599.

Bertolino S. 2017. Distribution and status of the declining garden dormouse Eliomys quercinus.
Mammal Review **47**: 133–147. http://dx.doi.org/10.1111/mam.12087.

Bertolino S, Amori G, Henttonen H, Zagorodnyuk I, Zima J, Juškaitis R, Meinig H, Kryštufek B.
2008. Eliomys quercinus. The IUCN Red List of Threatened Species.
https://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T7618A12835766.en.

Bertolino S, Colangelo P, Mori E, Capizzi D. 2015. Good for management, not for conservation:
an overview of research, conservation and management of Italian small mammals. *Hystrix*.
http://www.italian-journal-of-mammalogy.it/pdf-77188-13341?filename=Good for
management_ not.pdf.

Bertolino S, Viano C, Currado I. 2001. Population dynamics, breeding patterns and spatial use
of the garden dormouse (Eliomys quercinus) in an Alpine habitat. *J Zool* **253**: 513–521.

Blumer M, Brown T, Freitas MB, Destro AL, Oliveira JA, Morales AE, Schell T, Greve C, Pippel
M, Jebb D, et al. 2022. Gene losses in the common vampire bat illuminate molecular
adaptations to blood feeding. *Sci Adv* **8**: eabm6494.

Böhne A, Thiel-Bender C, Kukowka S, Darwin Tree of Life Barcoding collective, Wellcome
Sanger Institute Tree of Life programme, Wellcome Sanger Institute Scientific Operations:
DNA Pipelines collective, Tree of Life Core Informatics collective, Darwin Tree of Life
Consortium. 2023. The genome sequence of the hazel dormouse, Muscardinus
avellanarius (Linnaeus, 1758). *Wellcome Open Research* **8**.
https://wellcomeopenresearch.org/articles/8-514 (Accessed January 30, 2024).

Brandies P, Peel E, Hogg CJ, Belov K. 2019. The Value of Reference Genomes in the
Conservation of Threatened Species. *Genes* **10**.
https://pubmed.ncbi.nlm.nih.gov/31717707/.

Büchner S, Lang J, Meinig HU, Nava TF, Zistl K. 2024. The garden dormouse as a research
and conservation priority in a German large-scale citizen science project. *Acta Zoologica
Bulgarica* **Suppl 19**: 1–8.

Burri R, Nater A, Kawakami T, Mugal CF, Olason PI, Smeds L, Suh A, Dutoit L, Bureš S,
Garamszegi LZ, et al. 2015. Linked selection and recombination rate variation drive the
evolution of the genomic landscape of differentiation across the speciation continuum of
Ficedula flycatchers. *Genome Res* **25**: 1656–1665.

Campana MG, Parker LD, Hawkins MTR, Young HS, Helgen KM, Szykman Gunther M,
Woodroffe R, Maldonado JE, Fleischer RC. 2016. Genome sequence, population history,
and pelage genetics of the endangered African wild dog (Lycaon pictus). *BMC Genomics*

17: 1013.

Cheng H, Concepcion GT, Feng X, Zhang H, Li H. 2021. Haplotype-resolved de novo assembly
using phased assembly graphs with hifiasm. *Nat Methods* **18**: 170–175.

Chen S. 2023. Ultrafast one-pass FASTQ data preprocessing, quality control, and deduplication
using fastp. *Imeta* **2**. https://onlinelibrary.wiley.com/doi/10.1002/imt2.107.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics **34**: i884–i890.

Combe FJ, Juškaitis R, Trout RC, Bird S, Ellis JS, Norrey J, Al-Fulaij N, White I, Harris WE.
2023. Density and climate effects on age-specific survival and population growth:
consequences for hibernating mammals. *Anim Conserv* **26**: 317–330.

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G,
Marth GT, Sherry ST, et al. 2011. The variant call format and VCFtools. *Bioinformatics* **27**:
2156–2158.

Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, Whitwham A, Keane T,
McCarthy SA, Davies RM. 2021. Twelve years of SAMtools and BCFtools. *Gigascience* **10**:
1–4.

DeSaix MG, Bulluck LP, Eckert AJ, Viverette CB, Boves TJ, Reese JA, Tonra CM, Dyer RJ.
2019. Population assignment reveals low migratory connectivity in a weakly structured
songbird. *Mol Ecol* **28**: 2122–2135.

Ditto AM, Frey JK. 2007. Effects of ecogeographic variables on genetic variation in montane
mammals: implications for conservation in a global warming scenario. *J Biogeogr* **34**: 1136–
1149.

du Plessis SJ, Blaxter M, Koepfli K-P, Chadwick EA, Hailer F. 2023. Genomics reveals complex
population history and unexpected diversity of Eurasian otters (Lutra lutra) in Britain
relative to genetic methods. *Mol Biol Evol* **40**. https://dx.doi.org/10.1093/molbev/msad207.

Evanno G, Regnaut S, Goudet J. 2005. Detecting the number of clusters of individuals using the
software STRUCTURE: A simulation study. *Mol Ecol* **14**: 2611–2620.

Excoffier L, Foll M. 2011. fastsimcoal: a continuous-time coalescent simulator of genomic
diversity under arbitrarily complex evolutionary scenarios. *Bioinformatics* **27**: 1332–1334.

Excoffier L, Marchi N, Marques DA, Matthey-Doret R, Gouy A, Sousa VC. 2021. fastsimcoal2:
demographic inference under complex evolutionary scenarios. *Bioinformatics* **37**: 4882–
4885.

Forcina G, Camacho-Sanchez M, Cornellas A, Leonard JA. 2022. Complete mitogenomes
reveal limited genetic variability in the garden dormouse Eliomys quercinus of the Iberian
Peninsula. *Anim Biodivers Conserv* 107–122.

Formenti G, Theissinger K, Fernandes C, Bista I, Bombarely A, Bleidorn C, Ciofi C, Crottini A,
Godoy JA, Höglund J, et al. 2022. The era of reference genomes in conservation genomics.
Trends in Ecology and Evolution **37**: 197–202. https://www.

Francis RM. 2017. pophelper: an R package and web app to analyse and visualize population
structure. *Mol Ecol Resour* **17**: 27–32.

Frankish A, Diekhans M, Jungreis I, Lagarde J, Loveland JE, Mudge JM, Sisu C, Wright JC,
Armstrong J, Barnes I, et al. 2021. GENCODE 2021. *Nucleic Acids Res* **49**: D916–D923.

Gautier M, Moazami-Goudarzi K, Levéziel H, Parinello H, Grohs C, Rialle S, Kowalczyk R, Flori
L. 2016. Deciphering the Wisent Demographic and Adaptive Histories from Individual
Whole-Genome Sequences. *Mol Biol Evol* **33**: 2801–2814.

Gippoliti S, Amori G. 2007. Beyond threatened species and reintroduction: establishing priorities
for conservation and breeding programmes for European rodents in zoos. *International Zoo*
Yearbook **41**: 194–202.

Goudet AJ, Goudet MJ. 2014. Package " hierfstat ."

Gruber B, Unmack PJ, Berry OF, Georges A. 2018. dartr: An r package to facilitate analysis of
SNP data generated from reduced representation genome sequencing. *Mol Ecol Resour*
18: 691–699.

Halley DJ, Saveljev AP, Rosell F. 2021. Population and distribution of beavers Castor fiber and
Castor canadensis in Eurasia. *Mamm Rev* **51**: 1–24.

Harris RS. 2007. Improved pairwise alignment of genomic DNA. The Pennsylvania State
University, Ann Arbor, United States https://www.proquest.com/dissertations-
theses/improved-pairwise-alignment-genomic-dna/docview/304835295/se-2.

Hilgers L, Liu S, Jensen A, Brown T, Cousins T, Schweiger R, Guschanski K, Hiller M. 2024.
Avoidable false PSMC population size peaks occur across numerous studies. *bioRxiv*
2024.06.17.599025.
https://www.biorxiv.org/content/biorxiv/early/2024/06/18/2024.06.17.599025 (Accessed
August 14, 2024).

Howe K, Chow W, Collins J, Pelan S, Pointon D-L, Sims Y, Torrance J, Tracey A, Wood J.
2021. Significantly improving the quality of genome assemblies through curation.
Gigascience **10**. http://dx.doi.org/10.1093/gigascience/giaa153.

Huang N, Li H. 2023. compleasm: a faster and more accurate reimplementation of BUSCO.
Bioinformatics **39**. http://dx.doi.org/10.1093/bioinformatics/btad595.

Jebb D, Huang Z, Pippel M, Hughes GM, Lavrichenko K, Devanna P, Winkler S, Jermiin LS,
Skirmuntt EC, Katzourakis A, et al. 2020. Six reference-quality genomes reveal evolution of
bat adaptations. *Nature* **583**: 578–584.

Jombart T, Bateman A. 2008. adegenet: a R package for the multivariate analysis of genetic
markers. *Bioinformatics* **24**: 1403–1405.

Karimi K, Do DN, Wang J, Easley J, Borzouie S, Sargolzaei M, Plastow G, Wang Z, Miar Y.
2022. A chromosome-level genome assembly reveals genomic characteristics of the
American mink (Neogale vison). *Commun Biol* **5**: 1381.

Kennerley RJ, Lacher TE, Hudson MA, Long B, McCay SD, Roach NS, Turvey ST, Young RP.
2021. Global patterns of extinction risk and conservation needs for Rodentia and

Eulipotyphla. *Diversity and Distributions* **27**: 1792–1806.

Kent WJ, Baertsch R, Hinrichs A, Miller W, Haussler D. 2003. Evolution's cauldron: duplication,
deletion, and rearrangement in the mouse and human genomes. *Proc Natl Acad Sci U S A*
100: 11484–11489.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019. Graph-based genome alignment and
genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**: 907–915.

Kirilenko BM, Munegowda C, Osipova E, Jebb D, Sharma V, Blumer M, Morales AE, Ahmed A-
W, Kontopoulos D-G, Hilgers L, et al. 2023. Integrating gene annotation with orthology
inference at scale. *Science* **380**: eabn3107.

Lane JE, Kruuk LEB, Charmantier A, Murie JO, Dobson FS. 2012. Delayed phenology and
reduced fitness associated with climate change in a wild hibernator. *Nature* **489**: 554–557.

Lang J, Büchner S, Meinig H, Bertolino S. 2022. Do We Look for the Right Ones? An Overview
of Research Priorities and Conservation Status of Dormice (Gliridae) in Central Europe.
Sustainability (Switzerland) **14**: 9327. https://www.mdpi.com/2071-1050/14/15/9327/htm.

Leyhausen J, Cocchiararo B, Nowak C, Ansorge H, Bertolino S, Büchner S, Fietz J, Foppen R,
Juškaitis R, La Haye M, et al. 2022. Genotyping-by-sequencing based SNP discovery in a
non-model rodent, the endangered hazel dormouse. *Conserv Genet Resour* **14**: 195–201.

Libois R, Ramalhinho MG, Rosoux R. 2012. Evidence for a differentiated chromosomal race
north of classical south European refuge areas in the garden dormouse Eliomys quercinus.
Acta Theriol **57**: 313–320.

Li H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping and
population genetical parameter estimation from sequencing data. *Bioinformatics* **27**: 2987–
2993.

Li H. 2018. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**: 3094–
3100.

Li H, Durbin R. 2011. Inference of human population history from individual whole-genome
sequences. *Nature* **475**: 493–496.

Luo H, Jiang X, Li B, Wu J, Shen J, Xu Z, Zhou X, Hou M, Huang Z, Ou X, et al. 2023. A high-
quality genome assembly highlights the evolutionary history of the great bustard (Otis tarda,
Otidiformes). *Communications Biology* **6**: 1–11.

Lu X, Costeur L, Hugueney M, Maridet O. 2021. New data on early Oligocene dormice
(Rodentia, Gliridae) from southern Europe: phylogeny and diversification of the family. *J*
Syst Palaeontol **19**: 169–189.

Mahlert B, Gerritsmann H, Stalder G, Ruf T, Zahariev A, Blanc S, Giroud S. 2018. Implications
of being born late in the active season for growth, fattening, torpor use, winter survival and
fecundity. *Elife* **7**. http://dx.doi.org/10.7554/eLife.31225.

Manichaikul A, Mychaleckyj JC, Rich SS, Daly K, Sale M, Chen WM. 2010. Robust relationship
inference in genome-wide association studies. *Bioinformatics* **26**: 2867–2873.

Mansino S, García-Alix A, Ruiz-Sánchez FJ, Montoya P. 2014. A New Eliomys from the Late Miocene of Spain and Its Implications for the Phylogeny of the Genus. *acpp* **60**: 577–588.

Marchi N, Kapopoulou A, Excoffier L. 2024. Demogenomic inference from spatially and temporally heterogeneous samples. *Mol Ecol Resour* **24**: e13877.

Martchenko D, Shafer ABA. 2023. Contrasting whole-genome and reduced representation sequencing for population demographic and adaptive inference: an alpine mammal case study. *Heredity* **131**: 273–281.

Mead D, Fingland K, Cripps R, Portela Miguez R, Smith M, Corton C, Oliver K, Skelton J, Betteridge E, Doulcan J, et al. 2020. The genome sequence of the eastern grey squirrel, Sciurus carolinensis Gmelin, 1788. *Wellcome Open Res* **5**: 27.

Meinig H, Büchner S. 2012. The current situation of the garden dormouse (Eliomys quercinus) in Germany. *Peckiana* **8**: 129–134.

Milholland B, Dong X, Zhang L, Hao X, Suh Y, Vijg J. 2017. Differences between germline and somatic mutation rates in humans and mice. *Nat Commun* **8**: 15183.

Moritz C. 1994. Defining " Evolutionarily Significant Units " for conservation. *Trees* **9**: 373–375.

Mouton A, Mortelliti A, Grill A, Sara M, Kryštufek B, Juškaitis R, Latinne A, Amori G, Randi E, Büchner S, et al. 2017. Evolutionary history and species delimitations: a case study of the hazel dormouse, Muscardinus avellanarius. *Conserv Genet* **18**: 181–196.

Nadachoswki A, Daoud A. 1995. Patterns of myoxid evolution in the Pliocene and Pleistocene of Europe. *Hystrix-italian Journal of Mammalogy* **6**. https://doc.rero.ch/record/15906/files/PAL_E3309.pdf (Accessed February 14, 2024).

Nieto-Blázquez ME, Schreiber D, Mueller SA, Koch K, Nowak C, Pfenninger M. 2022. Human impact on the recent population history of the elusive European wildcat inferred from whole genome data. *BMC Genomics* **23**: 1–14.

Nunziata SO, Weisrock DW. 2018. Estimation of contemporary effective population size and population declines using RAD sequence data. *Heredity* **120**: 196–207.

O'Connell KA, Mulder KP, Maldonado J, Currie KL, Ferraro DM. 2019. Sampling related individuals within ponds biases estimates of population structure in a pond-breeding amphibian. *Ecol Evol* **9**: 3620–3636.

Osipova E, Hecker N, Hiller M. 2019. RepeatFiller newly identifies megabases of aligning repetitive sequences and improves annotations of conserved non-exonic elements. *Gigascience* **8**. http://dx.doi.org/10.1093/gigascience/giz132.

Paez S, Kraus RHS, Shapiro B, Gilbert MTP, Jarvis ED. 2022. Reference genomes for conservation. *Science* **377**: 364–366.

Pérez-Espona S. 2017. Conservation genetics in the European Union – Biases, gaps and future directions. *Biol Conserv* **209**: 130–136.

Perez GCL, Libois R, Nieberding CM. 2013. Phylogeography of the garden dormouse Eliomys quercinus in the western Palearctic region. *J Mammal* **94**: 202–217.

Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE. 2012. Double digest RADseq: An
inexpensive method for de novo SNP discovery and genotyping in model and non-model
species. *PLoS One* **7**. http://dx.doi.org/10.1371/journal.pone.0037135.

Pina-Martins F, Silva DN, Fino J, Paulo OS. 2017. Structure_threader: An improved method for
automation and parallelization of programs structure, fastStructure and MavericK on
multicore CPU systems. *Mol Ecol Resour* **17**: e268–e274.

Poplin R, Chang P-C, Alexander D, Schwartz S, Colthurst T, Ku A, Newburger D, Dijamco J,
Nguyen N, Afshar PT, et al. 2018. A universal SNP and small-indel variant caller using
deep neural networks. *Nat Biotechnol* **36**: 983–987.

Pritchard JK. 2000. Inference of population structure using multi-locus genotypes. *Genetics* **155**:
945–959.

Puechmaille SJ. 2016. The program structure does not reliably recover the correct population
structure when sampling is uneven: Subsampling and new estimators alleviate the problem.
Mol Ecol Resour **16**: 608–627.

Rammou D-L, Astaras C, Migli D, Boutsis G, Galanaki A, Kominos T, Youlatos D. 2022.
European ground squirrels at the edge: Current distribution status and anticipated impact of
climate on Europe's southernmost population. *Land (Basel)* **11**: 301.

R Core Team. 2018. R: A Language and Environment for Statistical Computing. *R Foundation*
for Statistical Computing Vienna Austria {ISBN} 3–900051–07–0.
https://scholar.google.com/scholar?cluster=9441913529578809097&hl=en&oi=scholarr.

Reiners TE, Eidenschenk J, Neumann K, Nowak C. 2014. Preservation of genetic diversity in a
wild and captive population of a rapidly declining mammal, the Common hamster of the
French Alsace region. *Mamm Biol* **79**: 240–246.

Rhie A, McCarthy SA, Fedrigo O, Damas J, Formenti G, Koren S, Uliano-Silva M, Chow W,
Fungtammasan A, Kim J, et al. 2021. Towards complete and error-free genome assemblies
of all vertebrate species. *Nature* **592**: 737–746.

Rhie A, Walenz BP, Koren S, Phillippy AM. 2020. Merqury: reference-free quality,
completeness, and phasing assessment for genome assemblies. *Genome Biol* **21**: 245.

Rick K, Byrne M, Cameron S, Cooper SJB, Dunlop J, Hill B, Lohr C, Mitchell NJ, Moritz C,
Travouillon KJ, et al. 2023. Population genomic diversity and structure in the golden
bandicoot: a history of isolation, extirpation, and conservation. *Heredity* **131**: 374–386.

Rochette NC, Rivera-Colón AG, Catchen JM. 2019. Stacks 2: Analytical methods for paired-end
sequencing improve RADseq-based population genomics. *Mol Ecol* **28**: 4737–4754.

Sambrook J, Russell WD. 2001. Protocol: DNA isolation from mammalian tissue. *Molecular*
Cloning: A Laboratory Manual (Sambrook J.

Schaub M, Vaterlaus-Schlegel C. 2001. Annual and seasonal variation of survival rates in the
garden dormouse (Eliomys quercinus). *J Zool* **255**: 89–96.

Schmidt TL, Jasper M-E, Weeks AR, Hoffmann AA. 2021. Unbiased population heterozygosity
estimates from genome-wide sequence data. *Methods Ecol Evol* **12**: 1888–1898.

Severn-Ellis AA, Scheben A, Neik TX, Saad NSM, Pradhan A, Batley J. 2020. Genotyping for
species identification and diversity assessment using double-digest restriction site-
associated DNA sequencing (ddRAD-Seq). In *Methods in Molecular Biology*, Vol. 2107 of,
pp. 159–187, Humana Press Inc.

Sharma V, Hiller M. 2017. Increased alignment sensitivity improves the usage of genome
alignments for comparative gene annotation. *Nucleic Acids Res* **45**: 8369–8377.

Shumate A, Wong B, Pertea G, Pertea M. 2022. Improved transcriptome assembly using a
hybrid of long and short reads with StringTie. *PLoS Comput Biol* **18**: e1009730.

Sopniewski J, Catullo RA. 2024. Estimates of heterozygosity from single nucleotide
polymorphism markers are context-dependent and often wrong. *Mol Ecol Resour* **24**:
e13947.

Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-analysis of
large phylogenies. *Bioinformatics* **30**: 1312–1313.

Suarez HG, Langer BE, Ladde P, Hiller M. 2017. chainCleaner improves genome alignment
specificity and sensitivity. *Bioinformatics* **33**: 1596–1603.

Szarmach SJ, Brelsford A, Witt CC, Toews DPL. 2021. Comparing divergence landscapes from
reduced-representation and whole genome resequencing in the yellow-rumped warbler
(Setophaga coronata) species complex. *Mol Ecol* **30**: 5994–6005.

Takach BV, Sargent H, Penton CE, Rick K, Murphy BP, Neave G. 2023. Population genomics
and conservation management of the threatened black-footed tree-rat (Mesembriomys
gouldii) in northern Australia. 1–11.

Talla V, Mrazek V, Höglund J, Backström N. 2023. Whole genome re-sequencing uncovers
significant population structure and low genetic diversity in the endangered clouded apollo
(Parnasssius mnemosyne) in Sweden. *Conserv Genet*. https://doi.org/10.1007/s10592-023-
01502-9.

Theissinger K, Fernandes C, Formenti G, Bista I, Berg PR, Bleidorn C, Bombarely A, Crottini A,
Gallo GR, Godoy JA, et al. 2023. How genomics can help biodiversity conservation. *Trends
in Genetics* **39**: 545–559. https://doi.org/10.1016/j.tig.2023.01.005.

Tsuchiya MTN, Dikow RB, Cassin-Sackett L. 2020. First Genome Sequence of the Gunnison's
Prairie Dog (Cynomys gunnisoni), a Keystone Species and Player in the Transmission of
Sylvatic Plague. *Genome Biol Evol* **12**: 618–625.

Uchimura A, Higuchi M, Minakuchi Y, Ohno M, Toyoda A, Fujiyama A, Miura I, Wakana S,
Nishino J, Yagi T. 2015. Germline mutation rates and the long-term phenotypic effects of
mutation accumulation in wild-type laboratory mice and mutator mice. *Genome Res* **25**:
1125–1134.

Verde Arregoitia LD. 2016. Biases, gaps, and opportunities in mammalian extinction risk
research. *Mamm Rev* **46**: 17–29.

Viluma A, Flagstad Ø, Åkesson M, Wikenros C, Sand H, Wabakken P, Ellegren H. 2022. Whole-
genome resequencing of temporally stratified samples reveals substantial loss of haplotype

diversity in the highly inbred Scandinavian Wolf population. *Genome Res* **32**: 449–458.

Viñals-Domingo A, Bertolino S, López-Iborra G, Gil-Delgado JA. 2020. Seasonal survival in a
non-hibernating Mediterranean garden dormouse population. *Mamm Biol* **100**: 581–589.

Wang J. 2018. Effects of sampling close relatives on some elementary population genetics
analyses. *Mol Ecol Resour* **18**: 41–54.

Weir BS, Cockerham CC. 1984. Estimating F-Statistics for the Analysis of Population Structure.
Evolution **38**: 1358.

White TA, Searle JB. 2007. Genetic diversity and population size: island populations of the
common shrew, Sorex araneus. *Mol Ecol* **16**: 2005–2016.

Wold J, Koepfli K-P, Galla SJ, Eccles D, Hogg CJ, Le Lec MF, Guhlin J, Santure AW, Steeves
TE. 2021. Expanding the conservation genomics toolbox: Incorporating structural variants
to enhance genomic studies for species of conservation concern. *Mol Ecol* **30**: 5949–5965.

Wolf JF, Bowman J, Keobouasone S, Taylor RS, Wilson PJ. 2022. A de novo genome assembly
and annotation of the southern flying squirrel (Glaucomys volans). *G3* **12**.
http://dx.doi.org/10.1093/g3journal/jkab373.

Wright BR, Farquharson KA, McLennan EA, Belov K, Hogg CJ, Grueber CE. 2020. A
demonstration of conservation genomics for threatened species management. *Mol Ecol*
Resour **20**: 1526–1541.

Yu G, Smith DK, Zhu H, Guan Y, Lam TTY. 2017. ggtree: an r package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods*
Ecol Evol **8**: 28–36.

Zhang H, Song L, Wang X, Cheng H, Wang C, Meyer CA, Liu T, Tang M, Aluru S, Yue F, et al.
2021. Fast alignment and preprocessing of chromatin profiles with Chromap. *Nat Commun*
12: 6566.

Zhou C, McCarthy SA, Durbin R. 2023. YaHS: yet another Hi-C scaffolding tool. *Bioinformatics*
39. http://dx.doi.org/10.1093/bioinformatics/btac808.

Zoonomia Consortium. 2020. A comparative genomics multitool for scientific discovery and
conservation. *Nature* **587**: 240–245.

1167