Cholesterol modifies Na$_v$1.7 - an *in-silico* and *in-vitro* analysis

Simone Albani*[1,2], Vishal Sudha Bhagavath Eswaran*[3], Alessia Piergentili[1,2,7], Paulo Cesar Telles de Souza[4,5], Angelika Lampert[+,3], and Giulia Rossetti[+,1,6,7]

[1]Institute of Neuroscience and Medicine (INM-9)/Institute for Advanced Simulation (IAS-5), Forschungszentrum Jülich GmbH, Wilhelm-Johnen-Straße, 52425 Jülich, Germany
[2]Faculty of Biology, RWTH Aachen University, Aachen, Germany
[3]Institute of Neurophysiology, Uniklinik RWTH Aachen University, Pauwelsstrasse 30, 52074 Aachen, Germany.
[4]Laboratoire de Biologie et Modélisation de la Cellule, CNRS, UMR 5239, Inserm, U1293, Université Claude Bernard Lyon 1, Ecole Normale Supérieure de Lyon, 46 Allée d'Italie, 69364, Lyon, France.
[5]Centre Blaise Pascal de Simulation et de Modélisation Numérique, Ecole Normale, Supérieure de Lyon, 46 All´ee d'Italie, 69364, Lyon, France.
[6]Jülich Supercomputing Centre (JSC), Forschungszentrum Jülich GmbH, Wilhelm-Johnen-Straße, 52425 Jülich, Germany
[7]Department of Neurology, University Hospital Aachen, RWTH Aachen University, Pauwelsstraße 30, 52074 Aachen

*Equally contributed

+Shared correspondence authorship

Abstract

Cholesterol is a major component of plasma membranes and unsurprisingly plays a significant role in actively regulating the functioning of several membrane proteins in humans. Notably, recent studies have shown that cholesterol depletion can also impact transmission of potentially painful signals in the context of peripheral inflammation, via hyperexcitability of the voltage-gated sodium channel (Na$_v$) subtype 1.9, but the structural mechanisms underlying this regulation remain to be elucidated. In this study, we focus on the role of cholesterol depletion on Na$_v$1.7, which is primarily expressed in the peripheral sensory neurons and linked to various chronic inherited pain syndromes. Coarse-grained molecular dynamics simulations shed light on the dynamic changes of the geometry of Na$_v$1.7 upon membrane cholesterol depletion: A loss of rigidity at key structural motifs linked to activation and fast-inactivation is observed, as well as changes in the geometry of drug-binding regions in the channel. Loss of rigidity in cholesterol depleted conditions should allow the channel to transition between different gating states more easily. *In-vitro* whole-cell patch clamp experiments on HEK293t cells expressing Na$_v$1.7 validated these predictions made *in silico* at the functional level. Hyperpolarizing shifts in the voltage-dependence of activation and fast-inactivation were observed along with an acceleration of the time to peak and onset kinetics of fast inactivation. These results underline the critical role of membrane composition, and of cholesterol in particular, in influencing Na$_v$1.7 gating characteristics. Furthermore, our results hint to a key role of the membrane environment in affecting drug effects and in pathophysiological dysregulation, sharpening our approaches for analgesics design.

Introduction

Cholesterol is a major factor in influencing the biophysical properties of plasma membranes, where it constitutes 30–45% of lipid molecules[1,2]. Several studies have shown that cholesterol modulates the function of a wide range of receptors and ion channels embedded in the membrane via specific, i.e., direct, ligand-like interactions, and non-specific indirect mechanisms[3–5]. For instance, cholesterol impacts the biophysical, microstructure and mechanical properties of the membrane, in turn impacting on the trafficking and expression level of several proteins (for a review see ref.[6]). Also, the molecular rearrangements resulting in the gating of ion channels, the transporting processes and the conformational transitions of transmembrane receptors are mediated by the surrounding lipid bilayer. Thus, all of these functions can be modified by cholesterol[6].

Recently, unexpected functions of cholesterol in controlling transmission of potentially painful (nociceptive) signals were uncovered, involving voltage-gated sodium channels (Na$_v$s). In the human body, nine different subtypes of Na$_v$s are expressed (Na$_v$1.1 to Na$_v$1.9), which are responsible for the fast upstroke of the action potential. They open when the membrane potential in their vicinity depolarizes, allowing sodium ions to flow into the cell[7]. Na$_v$1.7, Na$_v$1.8 and Na$_v$1.9, are expressed predominantly in the peripheral nervous system and are involved in the transduction and propagation of nociceptive signals[8]. A recent study suggests that pharmacological depletion of cellular cholesterol entails sensitization of nociceptive neurons potentially mediated via Na$_v$1.9, promoting mechanical and thermal hyperalgesia[9]. A similar effect was observed with inflammatory mediators, which were reported to support the partitioning of Na$_v$1.9 channels from cholesterol-rich microdomains of the plasma membrane (lipid rafts), in which also sphingolipids and gangliosides are accumulated[10], to cholesterol-poor non-raft regions of the membrane. Low-cholesterol environment enhances voltage-dependent activation of Na$_v$1.9 channels leading to enhanced neuronal excitability[9]. Na$_v$1.8 was also shown to activate more easily in cholesterol-depletion conditions, when the channel is disassociated from lipid rafts[11], suggesting that cholesterol can be a powerful modulator of nociception via Na$_v$s. Although several cholesterol interaction sites on Na$_v$1.9 were suggested, the molecular understanding of cholesterol-regulated hyperexcitability is not yet known[9].

In this study, we employed a range of computational and experimental techniques to assess the transferability of cholesterol-dependent effects to the subtype Na$_v$1.7 (**Fig. 1**), which was shown to be involved in human pain perception[12,13].

Not only do we provide evidence that cholesterol depletion causes hyperexcitability of Na$_v$1.7, similar to what was observed for Na$_v$1.9 and Na$_v$1.8, we also provide a molecular interpretation of how cholesterol interaction with Na$_v$1.7 structure impacts the dynamic features of the channel[9,11]. Notably, such structural and dynamic changes significantly impact Na$_v$1.7 interaction sites of known drugs, suggesting an effect on channel druggability in the presence of membrane-lipid modifiers such as inflammatory agents.

Figure 1. Structural overview of Na$_v$1.7 channel. (**a**) *Extracellular view of the α subunit of Na$_v$1.7 channel. It is a single-chain protein composed of four domains, each containing 6 transmembrane helices named from S1 to S6. In each domain, helices S1 to S4 are part of the voltage-sensing domain (VSD), while the helices S5 and S6 are part of the Pore Domain (PD)[14]. (**b**) Side view of domains III and IV. The fourth transmembrane helix (S4) in each domain, due to its positive residues, moves towards the extracellular side when the membrane is depolarized and is the main driver for voltage sensing. When Na$_v$ channels open and conduct sodium ions, the ion flow is interrupted after a few milliseconds by the process called "fast inactivation" [15]. During this process, a hydrophobic group of residues (Ile-Phe-Met, called IFM motif or inactivation particle) in the linker between Domains III and IV binds to a corresponding pocket in between the VSD of DIII and the PD, and allosterically restricts the pore[16,17].*

Results

We conducted coarse-grained molecular dynamics simulations to investigate the effect of membrane composition on the geometry of the α subunit of the human Na$_v$1.7 channel in a fast-inactivated state (**Fig. 1**). Specifically, we simulated the channel in three different membrane compositions: The first condition was designed to mimic a lipid raft in a neuronal membrane with a complex composition of lipids, including cholesterol and the ganglioside GM1 as main components, which was obtained from the MERMAID webserver (see methods); the second condition was composed solely of phosphatidylcholine (POPC, see methods for further details) to mimic a full detachment from the lipid raft; The third condition represented an intermediate simulation, wherein the neuronal membrane was identical to the first condition but the cholesterol was depleted and replaced with phosphatidylcholine. Experimentally, we performed whole-cell voltage clamp experiments in HEK293t cells expressing Na$_v$1.7 under control conditions and during pharmacologically induced cholesterol depletion.

Properties of the membrane and specificity of lipids interaction. The lipid raft-like membrane displayed a smaller area-per-lipid (APL 0.506 ± 0.003 nm^2, **Fig. 2a**) and an increased thickness (38.98 ±0.25 Å, **Fig. 2b**) compared to the cholesterol-depleted (APL 0.642± 0.004 nm^2, and thickness 37.54 ±0.26 Å) and the single-component membrane (APL 0.661 ±0.004 nm^2, and thickness 38.59 ±0.21 Å). The reduced thickness in the cholesterol-depleted membrane is in line with previous computational and experimental studies carried out with simple membrane compositions in different cholesterol

concentrations[18–20]. A tighter packing of the lipids (here expressed as APL) was also expected in the membrane containing cholesterol[21]. Despite these morphological changes, no significant variation in membrane curvature was observed (gaussian curvature in the range of -0.0125 to 0.008 Å^{-2}). These altered physical properties of the membrane could potentially impact the behavior of membrane proteins. Therefore, we tested for potential direct interactions between the membrane and key structural elements of Na$_v$1.7. Specifically, we calculated the occupancy of each membrane component around the channel to determine possible high-occupancy regions (i.e., regions where a given lipid is more frequently present) (**Fig. S1** and **S2**). In the single-component membrane, no specific interactions between the protein and the phosphatidylcholine molecules were identified, whereas in the analysis of the full lipid-raft membrane two membrane components emerged to establish localized interactions with the channel: Cholesterol and Ganglioside molecules. Such interactions were compared with existing knowledge and pertinent literature regarding interaction sites at the level of the membrane.

Table 1: Pairwise alignment of mouse Na$_v$1.9 (uniprot ID: Q9R053) and human Na$_v$1.7 (uniprot ID: Q15858-1) sequences in correspondence of the cholesterol binding sites found by Amsalem et al. (highlighted in red)[9].

Site 1	DI	Na$_v$1.9	195	NWLDFIVIGTAIAPCFLGNKVNNLSTLRTFRVLRALKAISVISGLKVIVG	244
		Na$_v$1.7	189	NWLDFVVIVFAYLTEFV--NLGNVSALRTFRVLRALKTISVIPGLKTIVG	236
Site 2	DIII	Na$_v$1.9	1024	LRKTCYQIVKHSWFESFIIFVILLSSGALIFEDVNLPSRPQVEKLLKCTD	1073
		Na$_v$1.7	1181	IRKTCYKIVEHSWFESFIVLMILLSSGALAFEDIYIERKKTIKIILEYAD	1231
Site 3	DIII	Na$_v$1.9	1122	LKSFRNLRALRPLRALSQFEGMKVVVNALMSAIPAILNVLLVCLIFWLIF	1171
		Na$_v$1.7	1286	IKSLRTLRALRPLRALSRFEGMRVVVNALIGAIPSIMNVLLVCLIFWLIF	1335

For cholesterol, we started by examining possible cholesterol binding sites suggested for Na$_v$1.9, as their sequence is highly conserved compared to Na$_v$1.7 (**Table 1**), and evaluated the occupancy of lipids surrounding the Na$_v$1.7 protein. To this purpose, we looked for so-called "CRAC" and "CARC" motifs, i.e., particular sequences, positioned near the membrane-solvent boundary, serving as reliable indicators for enhanced cholesterol binding affinity[22]. Very little overlap was identified between any of these motifs and the transmembrane (TM) helices that could be associated with the presence of a possible binding site (**Fig. 2c**). We next highlighted the regions around the channel in our simulations that are more frequently occupied by cholesterol molecules (at least 4% of the sampled frames, in 1 Å^3 voxels), and compared these regions with the positioning of cholesterol and its derivatives in structures obtained by CryoEM (**Fig. 2de**). Cholesteryl hemisuccinate is not naturally occurring in cells[23], but it is often used as a detergent and stabilizer of membrane proteins[24], and as such often found interacting with channel regions, most frequently on the surface of VSD$_{III}$ (S3 and S4), and on the fenestration entrance between DII S5 and DIII S5 and S6 (**Fig. 2f**). Only two common placements between simulation and CryoEM experiments were found: on the corner defined by DII S5/S6 and VSDI, and on the surface of VSD$_{III}$ (**Fig. 2ef**). Cholesterol also appears to bind at the entrance of fenestration DI-DIV and in proximity of VSD$_{II}$ S1 both in CryoEM and simulations. However, in the latter, the binding appears to be less stable as no voxel in these binding sites achieves an occupancy of 7% or higher. The other binding sites observed in Molecular Dynamics are often placed in contact with S5 and S6 of D I, II, and IV, hovering the S4-S5 linker of the adjacent domain (**Fig. 2de**). In particular, S5 of DI seems to interact strongly with cholesterol on both of its sides exposed to the membrane, thus rendering the mechanical hinge created by the S4-5 linker of DIV and non-bonded interactions between VSD$_{IV}$ and S5 and 6 of DI particularly stabilized (**Fig. 2d**).

Even though ganglioside molecules, too, have been demonstrated to increase the excitability of Na$_v$ channels[25], there are no known binding sites for these compounds. In our lipid raft-like simulations we

found that ganglioside molecules predominantly occupy regions near the membrane-solvent interface. There is limited penetration into the fenestrations, and their presence seems to primarily influence the surroundings of VSDs (**Fig. 2g**). Notably, higher occupancy of gangliosides is observed in the vicinity of VSD$_{III}$ in the full raft-like membrane simulation, where their positions complement the cholesterol densities (**Fig. 2h**). The same pattern appears to be partially applicable to the cholesterol-depleted membrane. In this case too, gangliosides exhibit interactions with the protein primarily in proximity to the membrane-solvent interface, but here a concentration of gangliosides is also observed at the entrance of the DI-DII fenestration. Intriguingly, in all the simulations in a complex membrane, VSD$_{IV}$ encounters multiple ganglioside clusters close to the S4-5 linker, potentially constraining its range of motion.

Figure 2. Analysis of membrane properties and lipid-protein interactions. a. *Area per lipid (APL) values, a measure of how tightly packed the lipid molecules are within the membrane, calculated over 12 μs per condition (1 sample/ns)* ***b.*** *Membrane thickness, measured as the average distance of phosphates groups between the two leaflets and calculated over 12 μs per condition (1 sample/ns)* ***c.*** *Sequence of CRAC and CARC motifs along protein sequence, generally no matching with TM helices.* ***e.*** *Extracellular view of Na$_v$1.7 and occupancy (average number beads per voxel per frame) of cholesterol in simulations (red isosurface)* ***f.*** *Extracellular view of Na$_v$1.7 and surface of cholesterol and cholesteryl hemisuccinate molecules previously observed in Cryo EM experiments on eukaryotic Na$_v$s (PDB IDs: 6NT3,6NT4, 7W9K, 7W9L, 7W9P, 7W9T, 7XVE, 7XVF) (red surface)[15,26,27]* ***g.*** *Extracellular view of Na$_v$1.7 and occupancy (average number of beads per voxel per frame) for ganglioside GM1 (DPG1) (purple isosurface)* ***h.*** *Cholesterol (red) and GM1 ganglioside (purple) high-occupancy regions as observed from a side view of the fenestration formed by S5 and S6 of DIII and DIV, that can be observed between VSD$_{II}$ and VSD$_{III}$.*

Membrane composition shapes 4-fold symmetry of Na$_v$1.7 *in silico* modeling. Asymmetric rearrangements in the PD have been associated with channel opening and closing. Specifically, monitoring the position of four residues located in the pore at the level of the inactivation gate enables the observation of a shift in the shape of the pore from a square to a rhomboid during the transition from a closed to an open or inactivated state in the rat Na$_v$1.5 channel[17]. We first evaluated whether fluctuations of the square shape of the channel occur naturally, within a timescale shorter than that required for VSD activation or channel opening.

To achieve this aim, we measured changes in the diagonals of such a square in all three simulated environments, specifically, the angle between vectors connecting VSD$_I$ to VSD$_{III}$ and VSD$_{II}$ to VSD$_{IV}$ (**Fig. 3a** and **S3**). Among the deposited human Na$_v$1.7 structures, this corresponds to about 84.5 degrees (84.45, 84.38, 84.65 for structures with PDB IDs 6J8G, 7W9P, 8G1A respectively). We found that this value is only slightly increased during our simulation in a lipid raft-like membrane (86.64 ± 1.48 degrees). The value remains relatively stable in the cholesterol-depleted raft-like membrane with a slightly larger deviation (84.12 ± 2.03 degrees). However, in the single-component membrane, it exhibits a broader range, assuming a bimodal distribution with peaks at 80.78 and 90.23 degrees (**Fig. 3b**). Next, we investigated which conformational changes drive such geometrical transformation.

Figure 3. Structural changes in VSDs. *a. Extracellular view of Na$_v$1.7 structure and representation of the angle to monitor cross shape of the channel b. Distribution of values of the angle between opposite VSDs for the two conditions (blue: cholesterol-rich lipid raft-like membrane; green: cholesterol-depleted lipid raft-like membrane; red: single-component membrane) c. Extracellular view of VSD$_I$ and definition of the angle to monitor VSD rotation around its intracellular-to-extracellular axis (representation of the axis in Figure S4). d. Distribution of angles of rotation of the four VSDs for the two conditions (blue: cholesterol-rich lipid raft-like membrane; green: cholesterol-depleted lipid raft-like membrane; red: single-component membrane). e. Distribution of tilt angle for the 24 transmembrane helices in the lipid raft-like and single-component membrane simulations. The average of the three replicas per condition is represented by black circles. A coloured stroked asterisk marks a significant difference in the tilt of an TM between*

Rotation of VSDs and changes in PD-VSD interface. CryoEM experiments show that the translocation in space of the VSD around PD is accompanied by a rotation of the VSD around its vertical axis, which in turn alters the interaction pattern between VSD and PD[27,28]. This observation could be linked to a process of deactivation of the VSD, as seen in a recent Na$_v$1.7 CryoEM structure. In this study, mutagenesis-induced deactivation of VSD$_I$ resulted in the domain tracing an angle of approximately 18 degrees around the PD, oriented towards VSD$_{IV}$[27]. Alternatively, a more extensive sampling may reveal similar phenomena, as demonstrated in the activated VSD$_I$ of a wild-type Nav1.8, which was captured after a rotation of 12 degrees around the core of the protein[28]. Although this protocol cannot simulate the rearrangements caused by a VSD deactivation, we observed this rotation in all four domains across our simulations and found that it is often - in 96% of the frames - limited to a range of 6 degrees and normally distributed. Some relevant exceptions appear when the channel is simulated in the single-component membrane (**Fig. 3cd** and **S5**): The distribution of angles for VSD$_{II}$ is wide (Min: -29.9, max: 21.9, 1 standard deviation: 7.2 degrees) and skewed to the right, due to a slightly more favorable, albeit still infrequent, placement of this DI in proximity of VSD$_{III}$ in one of the replicas (**Fig. 3d**, and **S5**); Rotations of the VSD$_{IV}$ are wider and much more frequent (Min. : -31.8, max.: 36.6, standard deviation: 10.9 degrees), with angles larger than 7.2 degrees being observed in most of the sampled conformations. For VSD$_I$, VSD$_{II}$, and VSD$_{IV}$, the channel simulated in a cholesterol-depleted raft-like membrane exhibits an intermediate behavior between the full raft-like and single-component membranes. VSD$_{III}$ deviates from this trend, by showing overlapping behavior between the cholesterol-depleted and single-component membrane simulations (**Fig. 3d**).

Tilt of VSD$_{IV}$. A thinner membrane can cause the transmembrane (TM) helices to tilt, to align their vertical component with the thinner double layer. Across 10 of the 24 TM helices, the median tilt is different in the raft-like vs. single-component comparison. In particular, 7 of the TM helices exhibit a greater tilt in the single-component membrane, especially in DIII and DIV (**Fig. 3ef**) Due to the preference for tilting in these domains and the lack of stabilizing effects from cholesterol and gangliosides at the region between PD and VSD$_{IV}$, the latter domain shifts to establish tighter interactions with the DIII-IV linker (**Fig. 3f** and **S7**). Therefore, our simulations strongly indicate that the above-discussed modified Na$_v$1.7 4-fold symmetry observed in the single-component membrane is due to the rotation of VSD$_{IV}$ around the PD, which can move either in the direction of VSD$_I$ or VSD$_{III}$. Given the tight association of VSD$_{IV}$ with the binding site of the IFM motif and its importance during the last stage of activation, this could have a great impact on the channel's dynamics. Additionally, in the comparison between the cholesterol-depleted v. full lipid raft-like membrane 4 TM helices display different tilting, with only 3 being more tilted in the cholesterol-depleted environment (**Fig. 3e** and **S6**). This suggests that cholesterol and membrane thickness have a limited, direct impact on inducing TM helix tilting in our simulations.

Effect on pore and fenestration size. In Na$_v$ channels, the pore cavity just below the selectivity filter is known to be the binding site for local anesthetics and antiarrhythmics[29]. Along the conduction pathway, at the cytosolic site of the inactivation gate, there is also the BIG (below inactivation gate) site, where drugs like carbamazepine bind[30]. Side openings of the channel, referred to as "fenestrations", are not only an access door for many molecules to the channel pore[29,31], they are interaction hotspots for pyrethroids, DDT, and phospholipids [32–34]. We measured the pore size along the conduction pathway and through the fenestration, to test whether membrane composition has an impact on conduction or druggability of the binding sites in the pore and fenestrations. We found that in the comparison lipid raft-like environment versus single-component membrane, a simple composition of the leaflets does not alter pore size at the level of the selectivity filter and only slightly increases the inactivation gate, but it does have a relevant impact on the central cavity which is reduced by 29.3% in volume, from 373.93 (±63.43) Å^3 to 264.45

(±74.19) $Å^3$ (**Fig. 4ab**). In the membrane resembling a cholesterol-depleted raft, the channel adopts intermediate values for the radius of central cavity (**Fig 4b**) and pore volume (310.74 ±83.80 $Å^3$). Fenestration size is always reduced in the single-component membrane, especially in the case of the fenestration formed by DII and DIII which is 43% smaller in volume and 0.67 Å narrower at its choke point (see **Fig.4cd** and **Table S1**). For fenestrations, the membrane resembling a cholesterol-depleted raft exhibits values that fall between those observed in the other two conditions.

Figure 4. Changes in pore and fenestration size. Pore and fenestration size for the three conditions (blue: lipid raft-like membrane; green: cholesterol-depleted raft-like membrane; red: single-component membrane). *a.* Side view of the pore profile along the conduction pathway. Green representation omitted for clarity (see panel b). *b.* Pore radius and bootstrapped 95%CI (n=1000 frames per simulation, representing 3 μs per condition) along conduction pathway *c.* Extracellular view of fenestration profiles Green representation omitted for clarity (see panel d). *d.* Fenestration radius and bootstrapped 95% CI along fenestration egress pathway.

Effects on other binding sites. Traditionally, drug design targeting $Na_v1.7$ involves two additional binding sites considered significant: the resting VSD_{II} and the activated VSD_{IV}[35]. The former has been identified as a binding site for toxins like ProTx2 or m3-Huwentoxin-IV, while the latter can also interact with simpler organic compounds, such as aryl and acyl sulfonamides. As previously discussed, both domains undergo substantial rotations and translations during the unbiased simulation. Since ligand binding predominantly occurs extracellularly on the portion of the domain facing away from the pore domain, we focused on monitoring significant changes in the shape of the binding domain by measuring the residue-to-residue distance between helices S2 and S3. In the VSD_{II}, opposing changes were observed between the lipid raft-like and single-component conditions: in the lipid raft-like membrane, the extracellular ends of S2 and S3 move apart, while they tend to approach each other in the single-component membrane simulations (**Fig. S8**). In the membrane resembling a cholesterol-depleted

raft, there is no consistent trend observed across the three replicas. However, in two of them, these ends slightly diverge. In the VSD_{IV}, changes were more subtle, with only a trend of helices S2 and S3 slightly converging observed in the lipid raft-like membrane (**Fig. S9**).

Summarizing, our simulations showed an increase in the flexibility and tilting of VSD_{IV} in a simple single-component membrane and also an increase in the occupancy of cholesterol with VSD_{III}. VSD_{IV} has been implicated in the fast inactivation process, while some studies have shown that the VSD_{III} can affect both the activation and fast inactivation processes[36].

Cholesterol depletion alters Na$_v$1.7 activation and fast inactivation properties *in vitro*. Our simulation results suggest a possible role for membrane composition in regulating the activation and fast-inactivation processes of Na$_v$1.7. We also observe *in silico* a high occupancy of cholesterol in regions important for gating. Combined with the known effects of cholesterol on gating of ion channels[37,11,9], we hypothesized that cholesterol in the complex membrane likely plays a pivotal role in our observed results. To validate these hypotheses *in vitro*, activation and fast inactivation properties were measured using electrophysiology. To this end, whole-cell patch clamp of HEK293t cells transfected with human Na$_v$1.7 wild type (hWT) plasmids was performed in control conditions (hWT ctrl) and under cholesterol depletion conditions using methyl-ß-cyclodextrin treatment (hWT MßCD). Cholesterol depletion was achieved by incubating the cells for 1h with the compound prior to recordings. MßCD is commonly used for the manipulation of cholesterol in various expression systems[38–40]. This compound has also been used to study voltage-gated sodium channels (e.g. Na$_v$1.8), and in particular Na$_v$1.4 in the same expression system used in this study[37,11].

Cholesterol depletion increases current densities and hyperpolarizes voltage-dependence of activation and steady-state fast inactivation. Under both control (without cholesterol depletion, defined as hWT ctrl) and experimental condition (with cholesterol depletion using MßCD, defined as hWT MßCD), robust inward currents were observed with quick inactivation kinetics (**Fig. 5a**). Patched cells showed a similar range of membrane capacitance values (C_{slow}), which give an indirect measure of cell size (**Fig. S10**). To account for effects of cell size on the magnitude of inward currents, the currents were normalized to C_{slow} to obtain the current densities. The current densities of hWT MßCD are higher than the current densities of hWT ctrl (**Fig. 5b**). The maximal current densities of hWT MßCD (645.8±186 pA/pF, n=15) were significantly higher than hWT ctrl (206.3±89 pA/pF, n=14), with the average value 3.1 times higher than hWT ctrl (**Table 2**, **Fig. 5c; p < 0.05**). Persistent currents were measured as the mean current values between 34ms and 39.6ms of each test pulse and normalized to the peak currents. The maximal value for each cell was used for further analyses and depicted as a percentage of the peak currents. There was a small, but statistically significant decrease in the mean normalized persistent current values of hWT MßCD (1.3±1.0%, n=15) when compared to hWT ctrl (1.8±0.6%, n=14) (Table 2, **Fig. 5d; p < 0.05**). Given the relatively low amplitudes of the non-normalized persistent currents in hWT ctrl and hWT MßCD (**Fig. S11a and b**), we checked if the values are higher than the noise in our system, which we assessed via the leak currents. The average persistent currents amplitudes were lower than the average leak current amplitudes in hWT ctrl and hWT MßCD across all voltage ranges (**Fig. S11c and d**). Thus, while statistical significance is reached, the measure may be affected by the leak currents. Noticeable differences were measured in the voltage-dependence of activation and steady-state fast inactivation. hWT MßCD induced hyperpolarizing shifts of 7.2mV and 7.8mV in the V1/2 of activation (**Fig. 5d and e**) and steady-state fast inactivation compared to hWT ctrl, respectively (**Fig. 5f and g**). However, no differences were observed in the slope of the curves (**Table 2**).

Table 2: Current density, persistent currents, voltage dependence of activation and steady-state fast inactivation of human Na$_v$1.7 wildtype under control (hWT ctrl) and cholesterol depletion (hWT MßCD) conditions. Values are represented as mean±95%

confidence interval of the mean. Values of hWT MßCD that are significantly different from hWT ctrl are represented by a * in the hWT MßCD column for the parameters which underwent a statistical significance test and had a p value is less than 0.05.

	hWT ctrl	hWT MßCD
n	14	15
Average peak current amplitudes (pA)	-1870±1017	-4813±2016
Average peak current densities (pA/pF)	206.3±89	645.8±186 *
Normalized maximal persistent currents (%)	1.5±0.5	0.9±0.6 *
Activation - V1/2 (mV) - kact	-21.1±1.8 9.3±0.7	-28.3±3.4 * 9.0±0.6
Inactivation - V1/2 (mV) - kinac	-90.1±2.0 8.4±1.6	-97.9±3.2 * 7.7±0.8

Cholesterol depletion alters the time to peak and rate of onset of fast inactivation but not recovery from fast inactivation. In addition to voltage dependencies of activation and fast inactivation, the kinetics of these processes were also measured. The traces obtained with the activation protocol (**Fig. 5e** inset) gives us information on the onset kinetics of activation and fast inactivation. The time to peak is measured as the time it takes from the start of the stimulus pulse (**Fig. 6a** blue arrow) to reach the peak current (**Fig. 6a** green arrow). The time to peak was more rapid in hWT MßCD when compared to hWT ctrl for pulses between -50mV and +40mV (**Fig. 6b**; p < 0.05). The rate of onset of fast inactivation was measured by fitting the inactivating phase of the IV traces to a single exponential fit (**Fig. 6c** in red) and obtaining the time constant(s) of the fit(s). Cholesterol depletion decreased the time constants for fast inactivation leading to a quicker rate of onset of fast inactivation between -40mV and +40mV (**Fig. 6d**; p < 0.05).

The time course of recovery from fast inactivation was measured using the protocol shown in Figure 6e. The peak currents obtained in test pulse I2 were normalized against the peak currents obtained in pre pulse I1 and plotted against the duration of the interpulse. This yields a measure of how many channels recovered from fast inactivation (and hence are ready for subsequent activation in the test pulse) as a function of the time given for the channels to recover. Cholesterol depletion did not alter the recovery kinetics of fast inactivation except at an interpulse duration of 1.6s (**Fig 6e,f**).

Figure 5: *Current density and gating properties of human Na$_v$1.7 wild type (hWT) under control and cholesterol depletion conditions (a) Representative traces of hWT control (hWT ctrl, left) and cholesterol depletion (hWT MßCD, right) conditions. Both conditions had robust inward currents (b) Average current density vs voltage for hWT ctrl and hWT MßCD. Error bars represent the 95% confidence interval. (c) Dot plot of peak current densities for hWT ctrl and hWT MßCD. Error bars represent the 95% confidence interval. * indicates a p-value < 0.05. (d) Maximal persistent currents as a percentage of the peak transient currents (Ipers,max/Ipeak) vs voltage (mV). Error bars represent the 95% confidence interval. * indicates a p-value < 0.05. (e) Normalized conductance-voltage relationships of hWT ctrl and hWT MßCD. g is the conductance at a specific voltage and gmax is the maximum conductance. The voltage-dependence of the activation protocol is indicated in the inset. (f) Dot plot of the V-half of activation for hWT ctrl and hWT MßCD. Error bars represent the 95% confidence interval. * indicates a p-value < 0.05. (g) Dot plot of the slope factor kact of the G-V curve for hWT ctrl and hWT MßCD. Error bars represent the 95% confidence interval. (h) Normalized current-voltage relationships of hWT ctrl and hWT MßCD. I is the inward current at a specific voltage and Imax is the maximal inward current. The voltage-dependence of steady state fast inactivation protocol is indicated in the inset. (i) Dot plot of the V-half of steady state fast inactivation for hWT ctrl and hWT MßCD. Error bars represent the 95% confidence interval of the mean. * indicates*

Figure 6: *Kinetics of activation and fast inactivation (a) Calculation of time to peak. The time to peak is calculated as the time taken from the start of impulse (blue arrow) until peak current is reached (green arrow). (b) time to peak vs voltage curves for hWT ctrl and hWT MßCD. * indicates a p-value < 0.05. Error bars represent the 95% confidence interval. (c) Calculation of the time constant for decay of fast inactivation. The inactivating portions of the traces from the activation protocol (highlighted in red) are fit with a single exponential function. The time constant of the fit is used for analysis (d) Time constant vs voltage curves for hWT ctrl and hWT MßCD. * indicates a p-value < 0.05. Error bars represent the 95% confidence interval. (e) Protocol to study the recovery kinetics from fast inactivation (f) Time course of recovery from fast inactivation. Peak currents from the test pulse (I2) were normalized to the peak currents during the pre-pulse (I1) and plotted against the duration of the interpulse. Error bars represent the 95% confidence interval of the mean.*

Limitations of the models

In silico. While all-atom MD simulations might initially appear as a preferable choice, the inherent slow dynamics of lipids, coupled with their intricate interplay within complex membranes and subtle changes in protein structure, pose challenges in obtaining adequate sampling for convergence in such models. Thus, we employed coarse-grained simulations to investigate the influence of membrane composition on the dynamics of the Na$_v$1.7 channel *in silico*. Utilizing the Martini 2.2 force field, renowned for accurately simulating membrane proteins and their interactions with lipids[41,42], we uncovered that membrane composition significantly affects various geometric properties associated with channel activation, conduction pore size, and druggability in addition to the properties of the membrane. These trends were observed despite some limitations of the *in silico* model. Numerous simulation protocols, including the one employed in this study, possess an inherent bias towards the initial protein configuration, due to the presence of an elastic network connecting the beads representing the protein's backbone in the simulations. This means that large conformational changes cannot be observed in unbiased simulations. To avoid introducing artificial differences, we employed a consistent elastic network to preserve channel shape in all simulations. This ensured that observed variations arose from membrane composition changes rather than a different protein parametrization. Another point to note is the deliberate exclusion of the N- and C-terminal regions along with the DI-DII and DII-DIII linkers from the Na$_v$1.7 structure. This was done since these regions are exclusively intracellular and our focus was solely on the transmembrane regions of the channel. Moreover, the Na$_v$1.7 structure used did not have good density maps in these regions due to their inherent flexibility[43]. Modeling intracellular linkers presents substantial challenges, whether through homology modeling or utilizing AlphaFold (alphafold.ebi.ac.uk/entryQ62205). Lastly, it is important to note that we employed Martini 2.2 instead of the latest version, Martini 3, as the coarse-grained model in this study. Although Martini 3 has demonstrated notable improvements, this new version has not undergone comprehensive benchmarking for investigations involving protein-lipid interactions yet. In addition, Martini 3 protein and lipid models are still under development, with some bonded terms still not fully updated and a few important biomolecule models missing[44]. Recently, models for cholesterol[44] and GM3[45] were introduced, but the GM1 lipid model is still pending release. Given these considerations, we opted to conduct the study using Martini 2.2.

In vitro. In the *in-vitro* approach, we used HEK cells to study the interplay between membrane composition and Na$_v$ functionality. The plasma membranes of these cells show higher rigidity at induced high cholesterol concentration[21,46], and, in standard conditions, different levels of sphingomyelin and phospholipids compared to other widely studied cell types[47]. Although a direct comparison of HEK cell membrane composition with sensory neurons is not reported, a recent publication investigates rat hippocampal neurons and suggests, similar to the results reported for HEK cells, that cholesterol enhances membrane rigidity, which may affect peptide insertion into the membrane[46].

Discussion

In the present study we show that Cholesterol affects shape and channel gating of the pain-related sodium channel Na$_v$1.7 by coarse-grain *in silico* modeling and *in-vitro* patch-clamp analysis of heterologously expressed channels. In the presence or absence of cholesterol varying effects on different ion channels are observed. For example, K$_v$1.3 channels show inhibition of function upon increase in cholesterol concentrations, without any significant changes during depletion of cholesterol[48,49]. In contrast, voltage-gated calcium channels showed an increase of current densities and a hyperpolarizing shift in the

voltage dependence of activation[50], upon cholesterol depletion. Effects on membrane protein functioning by altered membrane properties have also been shown for mechanosensitive piezo channels[51].

In this framework, the primary aim of this study was to expand our understanding on how membrane composition, and in particular cholesterol, can influence the functioning and dynamics of $Na_v1.7$ and highlight the possible consequences in the context of pain transmission and drug design.

The following were the major results from the study: **(1)** single component POPC membranes had reduced thickness and increased area per lipid compared to a full lipid raft-like membrane *in silico,* consistently to what observed in other *in-silico* and *in-vitro* studies[18–20] *;* **(2)** *in silico* modeling of a cholesterol-rich lipid raft-like membrane increased the rigidity of the channel compared with the single-component POPC membrane, as emerging from decreased movements of the channel's VSD_{iv} and alteration of various of its geometric properties (i.e., pore and fenestration size); **(3)** *in vitro* cholesterol depletion increases current densities of $Na_v1.7$ currents **(4)** hyperpolarizes the voltage-dependence of activation and steady-state fast inactivation, and accelerates the time to peak and fast inactivation onset kinetics; **(5)** cholesterol in complex membranes is observed to frequently interact with the $Na_v1.7$ protein close to the S4-S5 linkers *in silico.*

The effect of cholesterol on Na_v geometry and gating observed *in silico* and *in vitro*, respectively, could be attributed to indirect effects on the bilayer properties, or direct interactions with the channel protein[6].

Overall, cholesterol usually induces an increase in the stiffness of the bilayer[6,52], a reduction in its fluidity[21,46], and the resulting denser arrangement of the lipids[53]. Our first finding is in agreement with such experimental observation (**Finding 1**).

Cholesterol-driven membrane's stiffness can indirectly impact on the channel dynamics: Na_v channel activation indeed involves a pronounced movement of the S4 helix within the voltage-sensor and, via displacement of the S4-S5 linker, it induces dilation of the activation gate in the pore allowing ion permeation[54–56]. Similarly, fast inactivation is initiated by displacement of the DIII-DIV linker containing the inactivation particle, which finally binds to the side of the channel pore, thereby closing the permeation pathway. Thus, both, activation and fast inactivation, are based on larger movements within the channel protein, and a less rigid membrane surrounding, as giving when cholesterol is depleted, may result in speeding of these processes. Accordingly, larger rototranslational movements of the VSD_{IV} are observed *in silico* when cholesterol is absent, and these allow a tighter association to the inserted DIII-DIV linker (**Finding 2**). In parallel, in our patch-clamp recording in $Na_v1.7$ expressing cholesterol depleted HEK cells, we observed a speeding of time to peak and fast inactivation onset kinetics. Also steady-state fast inactivation and channel activation occurred at more negative potentials, indicating that less energy is needed for these gating processes when cholesterol is depleted. The observed increase in current density may result from quicker, and thus more coordinated channel opening. Alternatively, the reduced rigidity of the membrane may result in a higher rate of channel insertion, resulting from facilitated trafficking. Thus, the observed effects on channel gating are probably due to indirect action of cholesterol depletion on the properties of the membrane (**Finding 3 and 4**).

Conversely, the clustering of cholesterol molecules around the S4-S5 linkers observed *in silico* suggests that cholesterol can also directly impact in the channel's activation by restricting the displacement of the S4-S5 linker that is necessary for the activation of the VSDs and the opening of the activation gate in the pore allowing ion permeation (**Finding 5**).

Comparing our findings with what was previously observed for other Na$_v$s, cholesterol depletion appears to affect these channels in a subtype specific manner. On one hand, unlike our study, cholesterol depletion of Na$_v$1.4 in HEK293 cells did not increase current densities or change activation properties[37]. On the other hand, in agreement with our studies, Na$_v$1.9 showed an increased activity upon cholesterol depletion that was attributed to direct effects of cholesterol upon binding to any of the 17 motifs identified in the study[9]. Cholesterol depletion in Na$_v$1.8 also possibly reduced the channel's surface expression and reduced the number of neurons conducting depolarizations[57].

Changes in membrane properties might be a more general mechanism by which cholesterol depletion alters the gating of Na$_v$s. Other indirect effects such as altered interactions with signaling molecules, e.g. kinases, and direct binding of cholesterol to regions in the channel, may underlie subtype specific differences of cholesterol depletion and present a promising avenue for further research.

Our findings highlight the importance of considering the effects of lipid membrane composition when employing structure-based drug discovery, as they can impact the size and shape of relevant binding sites. In a broader view, our findings open up new paths for rational drug design of novel analgesics that take into account cholesterol modulation and the environmental features of channels' localization.

Material and Methods

Pairwise sequence alignment. Mouse Na$_v$1.9 (uniprot ID: Q9R053) and human Na$_v$1.7 (uniprot ID: Q15858-1) sequences were aligned using the online tool EMBOSS Needle, using a EBLOSUM62 matrix, a gap penalty of 10.0, and an extend penalty of 0.5 (https://www.ebi.ac.uk/Tools/psa/emboss_needle/).

Molecular Dynamics. All simulations were performed using Gromacs 2019.4[58,59] and the Martini 2.2 forcefield with ElNeDyn22 and lipids 2.0[60–63]. The structure of the channel was downloaded from the PDB database (PDB ID: 6J8G)[43], loops were repaired using Modeler 9.10[64,65], aligned vertically to a horizontal membrane bilayer using PPM webserver 2.0[66]. The topology of the protein was created using Martinize[63]. The default settings for the elastic network were used, with a final number of 4385 distance restraints, with minimum and maximum distance of 0.33 nm and 0.9 nm, respectively. The elastic bond force constant for all the bonds was set to 500 kJ/mol/nm^2. The protein was embedded in the membrane using Insane (https://github.com/Tsjerk/Insane). The raft-like membrane was inspired to the MERMAID webserver representation of a human myelin membrane[67] and it is formed by 12% phosphatidylcholine (POPC), 18% phosphatidylethanolamine (POPE), 30% Cholesterol (CHOL), 10% phosphatidylserine (POPS), 1% phosphatidylinositol (POPI), 8% sphingomyelin (POSM) and 21% monosialotetrahexosylganglioside (DPG1). In the raft-like (-chol) membrane, the cholesterol is replaced by POPC, while in the single-component membrane, the bilayer is composed of 100% POPC. The salt (NaCl) concentration was set to 0.15M. The box size containing the system was adjusted with a short 2ns NPT simulation, then the system was equilibrated with 200ns of simulation in an NVT ensemble, followed by 200ns in an NPT ensemble. During equilibration, 1000 Kj/mol restraints were applied to the backbone of the protein. For each condition, three 4-µs replicas were produced. The last 3 µs of each replica were used for the analysis.

Domain rotation analysis. The intervals defining the 24 transmembrane helices were obtained from Uniprot (gene: SCN9a_human, ID: Q15858) and adapted to match isoform 3 of the channel. The center of mass of each VSD was calculated using the masses of the helices S1 to S4. To calculate the rotation of the VSD around its vertical axis, a vector connecting the S1+S2 and S3+S4 centers of mass was defined for each domain.

Pore and fenestration size. The program HOLE 2 was used to measure conduction pore and fenestration size of the coarse-grained channel during the last microsecond of production run (1 frame/ns, 1000 frames per simulation) of all the replicas[68]. The protein was roto-translated to fit the starting configuration using GROMACS 2019.4. The starting point for the sampling was set to the center of mass of the backbone. The conduction pore was measured along the z coordinate of the simulation box, whereas the fenestration direction was defined by the two vectors connecting residue 952 to 1634; and residue 249 to 1441. The limits of the central cavity were defined on the first frame from the z coordinate of the selectivity filter (5.8 Å above the center of mass of the protein) and Tyrosine 1755 (Na$_v$1.7 isoform 1 numbering) (12.4 Å below the center of mass of the protein). The calculations for each fenestration were truncated at a distance of 22.5 Å from the center of mass of the protein. The null values rarely obtained in the calculations, represent an imperfection of the pathfinding algorithm that does not always succeed in finding a complete path from the begin to the end of the otherwise-open pore or fenestrations. Therefore these values were excluded from the calculations, rather than being considered an interruption in the path. Van Der Waals radii were set to 2.35 Å for the standard beads, and 2.15 Å for the smallest ones. The sampled radii were extracted in Python 3.9.7, aligned, and averaged and plotted using the Numpy (v 1.21.2)[69] and Matplotlib (v 3.4.3)[70] libraries. Volumes were approximated as the sum of π x (sampled radii)^2 x h, where h is the sampling interval corresponding to 0.25 Å. Volumes were calculated per each frame, and the average and standard deviation of 3000 frames per condition was reported. The chokepoint was measured as the minimum non-null radius in the fenestration. It was calculated per each frame and the average and standard deviation of 3000 frames per condition was reported. Bootstrapping of the radii was performed with the scipy (v 1.8.0) python library[71] and "BCa" method and n=1000 and reported in the visualization in Figure 4.

Lipid bilayer thickness, area per lipid and cholesterol binding motif analysis. The position of the head of phosphatidyl lipids (beads named PO4) was extracted from the simulations via MDAnalysis (v 2.0.0 and split into upper and lower layer groups by a horizontal plane placed in correspondence of the average *z* positions of the beads. The subdivision in two layers was updated every frame. The distance between the two leaflets was sampled for all simulations with a sample rate of 1 sample/1ns, for a total of 12,000 measurements per membrane-type. The distribution of the area per lipid was measured using Fatslim[72] and by selecting all PO4, ROH, and AM1 beads as lipid heads. In this case too, the subdivision in two layers was updated every frame. All protein beads were treated as "interacting groups". Emboss fuzzpro (www.bioinformatics.nl/cgi-bin/emboss/fuzzpro) was used to search for CRAC and CARC sequences in the SCN9A protein sequence isoform 1 reported in Uniprot (ID: Q15858-1 (www.uniprot.org/uniprotkb/Q15858/entry#Q15858-1), using as a query "[LV]X(1,5)YX(1,5)[RK]" and "[RK]X(1,5)YX(1,5)[LV]".

Analysis of transmembrane helices tilt. We utilized the centers of mass from the first and last four residues of every transmembrane helix to establish a vector that indicates the direction of the helix. TM helices regions were selected as discussed above. These vectors were then compared to the unit vector of the z-coordinate of the simulation box, and the resulting angle between them was recorded for each helix once every nanosecond. The average (last 1μs, 1000 frames) TM helix tilt of the replicas were split into two groups representing the two conditions to be compared with an independent t-test. The significance threshold was set at p<0.05.

Analysis of occupancy of membrane components. The analysis was performed with VMD (1.9.4a55) and the integrated VolMap tool[73]. For the calculation, a grid spacing of 1Å was used and simulated particles were considered point-like. A total of 12,000 frames (1 frame/ns) per membrane composition was used for the analysis, representing 12 μs.

Transfection of HEK293t Cells. HEK293t cells were cultured in were cultured in Dulbecco's modified Eagle's medium (DMEM/F12; Gibco-Life Technologies, Carlsbad, CA, USA), supplemented with 10% fetal bovine serum (FBS; Gibco-Life Technologies, Carlsbad, CA, USA) and incubated at 37°C and 5% CO_2. Human $Na_v1.7$ (hWT) plasmid was contained in a pCMV6Neo vector. 2-5h post seeding of the HEK293t cells, 1.25µg of hWT plasmid and 0.25µg of green fluorescence protein (pMax-GFP) plasmid were co-transfected into the cells using 3µL of the jetPEI® transfection reagent (Polyplus-transfection S.A., Illkirch, France) and incubated overnight at 37°C and 5% CO_2. Transfected cells were patched 24h post transfection.

Manipulation of Cellular Cholesterol Content. Membrane cholesterol content was decreased by exposure to the cholesterol chelating agent methyl-ß-cyclodextrin (MßCD). 258.2g of powdered MßCD (Sigma-aldrich, C4555) was dissolved in 1ml of distilled water and vortexed to produce a 200mM clear MßCD stock solution. The stock solution was mixed with DMEM/F12 medium to create a 5mM MßCD medium. HEK293t cells that were used for cholesterol depleted conditions were incubated in the 5mM MßCD medium for 1h at 37°C and 5% CO_2 before patching.

Whole-cell Voltage Clamp Experiments. Whole-cell voltage clamp recordings of transfected HEK293t cells were performed at room temperature using the HEKA EPC 10 USB amplifier (HEKA Electronics, Lambrecht, Germany).

The recording pipettes were pulled from borosilicate glass capillaries using a DMZ puller (Zeitz Instruments GmbH, Martinsried, Germany). The pipette resistances varied between 1-3MΩ. The external bath solution contained the following (in mM): 40 NaCl, 100 Choline-Cl, 3 KCl, 1 $MgCl_2$, 1 $CaCl_2$, 10 HEPES, 5 glucose and 10 sucrose. pH of the solution was adjusted to 7.4 using CsOH and osmolarity to about 298 mOsm. 1 in 1000 parts dimethylsulfoxide (DMSO) was added to the bath solution. The internal pipette solution contained the following (in mM): 10 NaCl, 140 CsF, 1 EGTA, 19 HEPES and 18 sucrose. pH of the solution was adjusted to 7.3 using CsOH and osmolarity to about 310mOsm.

For all cells recorded and considered for further analyses, series resistance was compensated by at least 70% to ensure that voltage errors did not exceed 5mV. Leak current was subtracted digitally using the P/4 procedure. Unless stated otherwise, the holding potential was maintained at -120mV. No liquid junction potential corrections were performed. After whole-cell configuration is reached, the inward sodium currents generated by the hWT-expressing cells were allowed to stabilize for 3 min during repeated -10mV depolarization steps before starting the protocols.

The voltage-dependence of activation was assessed from the holding potential (Vhold) using 40ms test pulses with varying potentials from -90mV to +40mV in 10mV increments, with a 5s inter-pulse interval (Figure 5e, inset). The inward currents (I) recorded at each voltage step (V) could then be used to calculate the conductance (G) by the equation: G = I/(V-Vrev), where Vrev is the reversal potential for sodium. The conductance-voltage curves could then be fit using a boltzmann equation to obtain the voltage-dependence characteristics of activation: $G=Gmax/(1 + exp[Vm - V1/2)/k_{act}])$, where Gmax is the maximal sodium conductance, Vm is the membrane voltage, V1/2 is the potential of half maximal activation and k_{act} is the slope factor. The time course of fast inactivation, which estimates how rapidly channels fast inactivate, can be obtained for each test pulse by fitting the inactivating portion of the activation trace (Figure 6c) to a single exponential function: $Y=Y0 + A*exp(-KX)$, where Y0 is the steady-state current amplitude, A is the amplitude coefficient, K is the rate constant and X is the time. The time constant τ_{decay}, which is the reciprocal of K, is plotted against the test-pulse voltages from -40mV to +40mV. The time to peak is measured for each test pulse by calculating the time taken between the start of the test pulse and the channels reaching Imax (Figure 6a). The time to peak is plotted against the test-pulse voltages between -50mV and +40mV. Maximal current densities for each cell were calculated

by normalizing the peak currents for each test pulse by the membrane capacitance of the cell. Current density values have the unit pA/pF.

The voltage-dependence of steady-state fast inactivation was measured using a two-step protocol. Firstly, a 500ms pre-pulse from Vhold with potentials ranging from -140mV to -10mV in 10mV increments was used to inactivate the channels. This was followed by a 40ms test pulse to 0mV to gauge the fraction of channels that are still fast inactivated (Figure 5h, inset). The inward currents (I) measured during the test pulse were normalized to the maximal inward current (Imax) of the cell and plotted against the pre-pulse voltages (V) to obtain a normalized current-voltage curve. The curve was fit using the boltzmann equation: $I/Imax = Imin + (Imax-Imin)/(1+exp[(V1/2 - V)]/k_{inac})$, where Imin is the minimal inward current of the cell, V1/2 is the potential of half-maximal channel fast inactivation, and k_{inac} is the slope factor.

Recovery from fast inactivation was also measured using a two-pulse protocol. Cells were first depolarized using a pre-pulse to +10mV for 500ms from Vhold. This was followed up by an inter-pulse to -120mV with increasing durations in each sweep by a factor of 2 - from 0.1ms to 1600ms. The inter-pulse was followed up with another depolarizing test pulse to +10mV for 40ms to obtain the fraction of channels that could recover from fast inactivation (Figure 6e). The current amplitudes at the test pulse (Itest) was normalized to the current amplitude at the pre pulse (Ipre) and plotted against the inter-pulse durations to obtain the time course of recovery from fast inactivation.

Whole-Cell Voltage Clamp Data Analysis and Statistics

Raw data of the recordings were processed using Fitmaster (HEKA Electronics, Lambrecht, Germany) and exported into Igor Pro (Wavemetrics, Portland, OR, USA) for extraction of features from the traces using in-house scripts. Curve fittings and generation of the graphs were done using Graphpad Prism 5 (Graphpad Software Inc., La Jolla, CA, USA). For statistical testing, the control and cholesterol depleted conditions were compared using either a Student's t-test or a mann-whitney U test based on the normality of the data which was tested using the Shapiro-Wilk normality test. The data is always represented as the mean±95% confidence interval of the mean and the error bars in the graphs denote the 95% confidence interval of the mean.

References

1. Yeagle, P. L. Cholesterol and the cell membrane. *Biochim. Biophys. Acta BBA - Rev. Biomembr.* **822**, 267–287 (1985).
2. Yeagle, P. L. Modulation of membrane function by cholesterol. *Biochimie* **73**, 1303–1310 (1991).
3. Li, H. & Papadopoulos, V. Peripheral-Type Benzodiazepine Receptor Function in Cholesterol Transport. Identification of a Putative Cholesterol Recognition/Interaction Amino Acid Sequence and Consensus Pattern. *Endocrinology* **139**, 4991–4997 (1998).
4. Levitan, I., Fang, Y., Rosenhouse-Dantsker, A. & Romanenko, V. Cholesterol and Ion Channels. in *Cholesterol Binding and Cholesterol Transport Proteins: Structure and Function in Health and Disease* (ed. Harris, J. R.) 509–549 (Springer Netherlands, Dordrecht, 2010). doi:10.1007/978-90-481-8622-8_19.
5. Corradi, V. *et al.* A molecular switch controls the impact of cholesterol on a Kir channel. *Proc. Natl. Acad. Sci.* **119**, e2109431119 (2022).
6. Zakany, F., Kovacs, T., Panyi, G. & Varga, Z. Direct and indirect cholesterol effects on membrane proteins with special focus on potassium channels. *Biochim. Biophys. Acta BBA - Mol. Cell Biol. Lipids* **1865**, 158706 (2020).
7. Yu, F. H. & Catterall, W. A. Overview of the voltage-gated sodium channel family. *Genome Biol.* **4**, 207 (2003).
8. Hameed, S. Nav1.7 and Nav1.8: Role in the pathophysiology of pain. *Mol. Pain* **15**, 1744806919858801 (2019).
9. Amsalem, M., Poilbout, C., Ferracci, G., Delmas, P. & Padilla, F. Membrane cholesterol depletion as a trigger of Nav1.9 channel‐mediated inflammatory pain. *EMBO J.* **37**, (2018).
10. Rudajev, V. & Novotny, J. The Role of Lipid Environment in Ganglioside GM1-Induced Amyloid β Aggregation. *Membranes* **10**, 226 (2020).
11. Pristerà, A., Baker, M. D. & Okuse, K. Association between Tetrodotoxin Resistant Channels and Lipid Rafts Regulates Sensory Neuron Excitability. *PLOS ONE* **7**, e40079 (2012).
12. McDermott, L. A. *et al.* Defining the Functional Role of NaV1.7 in Human Nociception. *Neuron* **101**, 905-919.e8 (2019).
13. Shields, S. D. *et al.* Insensitivity to Pain upon Adult-Onset Deletion of Nav1.7 or Its Blockade with Selective Inhibitors. *J. Neurosci.* **38**, 10180–10201 (2018).
14. Ahern, C. A., Payandeh, J., Bosmans, F. & Chanda, B. The hitchhiker's guide to the voltage-gated sodium channel galaxy. *J. Gen. Physiol.* **147**, 1–24 (2015).
15. Clairfeuille, T. *et al.* Structural basis of α-scorpion toxin action on Nav channels. *Science* **363**, (2019).
16. Li, Z. *et al.* Structure of human Nav1.5 reveals the fast inactivation-related segments as a mutational hotspot for the long QT syndrome. *Proc. Natl. Acad. Sci.* **118**, e2100069118 (2021).
17. Jiang, D. *et al.* Open-state structure and pore gating mechanism of the cardiac sodium channel. *Cell* **184**, 5151-5162.e11 (2021).
18. Hung, W.-C., Lee, M.-T., Chen, F.-Y. & Huang, H. W. The Condensing Effect of Cholesterol in Lipid Bilayers. *Biophys. J.* **92**, 3960–3967 (2007).
19. Martinez-Seara, H. *et al.* Interplay of Unsaturated Phospholipids and Cholesterol in Membranes: Effect of the Double-Bond Position. *Biophys. J.* **95**, 3295–3305 (2008).
20. Javanainen, M. *et al.* Quantitative Comparison against Experiments Reveals Imperfections in Force Fields' Descriptions of POPC–Cholesterol Interactions. *J. Chem. Theory Comput.* **19**, 6342–6352 (2023).
21. Ayee, M. A. & Levitan, I. Paradoxical impact of cholesterol on lipid packing and cell stiffness. *Front. Biosci.-Landmark* **21**, 1245–1259 (2016).
22. Fantini, J. & Barrantes, F. J. How cholesterol interacts with membrane proteins: an exploration of cholesterol-binding sites including CRAC, CARC, and tilted domains. *Front. Physiol.* **4**, (2013).
23. Kulig, W. *et al.* Experimental determination and computational interpretation of biophysical properties of lipid bilayers enriched by cholesteryl hemisuccinate. *Biochim. Biophys. Acta BBA - Biomembr.* **1848**, 422–432 (2015).
24. Autzen, H. E., Julius, D. & Cheng, Y. Membrane mimetic systems in CryoEM: keeping membrane proteins in their native environment. *Curr. Opin. Struct. Biol.* **58**, 259–268 (2019).
25. Salazar, B. C., Castaño, S., Sánchez, J. C., Romero, M. & Recio-Pinto, E. Ganglioside GD1a

increases the excitability of voltage-dependent sodium channels. *Brain Res.* **1021**, 151–158 (2004).

26. Huang, G. *et al.* High-resolution structures of human Nav1.7 reveal gating modulation through α-π helical transition of S6IV. *Cell Rep.* **39**, 110735 (2022).

27. Huang, G. *et al.* Unwinding and spiral sliding of S4 and domain rotation of VSD during the electromechanical coupling in Nav1.7. *Proc. Natl. Acad. Sci.* **119**, e2209164119 (2022).

28. Huang, X. *et al.* Structural basis for high-voltage activation and subtype-specific inhibition of human Nav1.8. *Proc. Natl. Acad. Sci.* **119**, e2208211119 (2022).

29. Körner, J. *et al.* Sodium Channels and Local Anesthetics—Old Friends With New Perspectives. *Front. Pharmacol.* **13**, (2022).

30. Wu, Q. *et al.* Structural mapping of Nav1.7 antagonists. *Nat. Commun.* **14**, 3224 (2023).

31. Sait, L. G. *et al.* Cannabidiol interactions with voltage-gated sodium channels. *eLife* **9**, e58593 (2020).

32. Du, Y. *et al.* Molecular evidence for dual pyrethroid-receptor sites on a mosquito sodium channel. *Proc. Natl. Acad. Sci.* **110**, 11785–11790 (2013).

33. Kaczmarski, J. A. & Corry, B. Investigating the size and dynamics of voltage-gated sodium channel fenestrations. *Channels* **8**, 264–277 (2014).

34. Field, L. M., Emyr Davies, T. G., O'Reilly, A. O., Williamson, M. S. & Wallace, B. A. Voltage-gated sodium channels as targets for pyrethroid insecticides. *Eur. Biophys. J.* **46**, 675–679 (2017).

35. Nguyen, P. T. & Yarov-Yarovoy, V. Towards Structure-Guided Development of Pain Therapeutics Targeting Voltage-Gated Sodium Channels. *Front. Pharmacol.* **13**, (2022).

36. Angsutararux, P., Kang, P. W., Zhu, W. & Silva, J. R. Conformations of voltage-sensing domain III differentially define NaV channel closed- and open-state inactivation. *J. Gen. Physiol.* **153**, e202112891 (2021).

37. Lundbæk, J. A. *et al.* Regulation of Sodium Channel Function by Bilayer Elasticity. *J. Gen. Physiol.* **123**, 599–621 (2004).

38. Rodal, S. K. *et al.* Extraction of Cholesterol with Methyl-β-Cyclodextrin Perturbs Formation of Clathrin-coated Endocytic Vesicles. *Mol. Biol. Cell* **10**, 961–974 (1999).

39. Zeidan, A., Broman, J., Hellstrand, P. & Swärd, K. Cholesterol dependence of vascular ERK1/2 activation and growth in response to stretch: role of endothelin-1. *Arterioscler. Thromb. Vasc. Biol.* **23**, 1528–1534 (2003).

40. Balse, E. *et al.* Cholesterol modulates the recruitment of Kv1.5 channels from Rab11-associated recycling endosome in native atrial myocytes. *Proc. Natl. Acad. Sci.* **106**, 14681–14686 (2009).

41. Corradi, V. *et al.* Emerging Diversity in Lipid–Protein Interactions. *Chem. Rev.* **119**, 5775–5848 (2019).

42. Marrink, S. J. *et al.* Computational Modeling of Realistic Cell Membranes. *Chem. Rev.* **119**, 6184–6226 (2019).

43. Shen, H., Liu, D., Wu, K., Lei, J. & Yan, N. Structures of human Nav1.7 channel in complex with auxiliary subunits and animal toxins. *Science* **363**, 1303–1308 (2019).

44. Marrink, S. J. *et al.* Two decades of Martini: Better beads, broader scope. *WIREs Comput. Mol. Sci.* **13**, e1620 (2023).

45. Grünewald, F. *et al.* Martini 3 Coarse-Grained Force Field for Carbohydrates. *J. Chem. Theory Comput.* **18**, 7555–7569 (2022).

46. Fernández-Pérez, E. J. *et al.* Effect of Cholesterol on Membrane Fluidity and Association of Aβ Oligomers and Subsequent Neuronal Damage: A Double-Edged Sword. *Front. Aging Neurosci.* **10**, (2018).

47. Zhang, Y. *et al.* High-Throughput Lipidomic and Transcriptomic Analysis To Compare SP2/0, CHO, and HEK-293 Mammalian Cell Lines. *Anal. Chem.* **89**, 1477–1485 (2017).

48. Hajdú, P., Varga, Z., Pieri, C., Panyi, G. & Gáspár, R. Cholesterol modifies the gating of Kv1.3 in human T lymphocytes. *Pflüg. Arch.* **445**, 674–682 (2003).

49. Jiang, Q.-X. Cholesterol-Dependent Gating Effects on Ion Channels. *Adv. Exp. Med. Biol.* **1115**, 167–190 (2019).

50. Purcell, E. K., Liu, L., Thomas, P. V. & Duncan, R. K. Cholesterol Influences Voltage-Gated Calcium Channels and BK-Type Potassium Channels in Auditory Hair Cells. *PLOS ONE* **6**, e26289 (2011).

51. Haselwandter, C. A. & MacKinnon, R. Piezo's membrane footprint and its contribution to mechanosensitivity. *eLife* **7**, e41968 (2018).

52. Lundbæk, J. A., Birn, P., Girshman, J., Hansen, A. J. & Andersen, O. S. Membrane Stiffness and Channel Function. *Biochemistry* **35**, 3825–3830 (1996).

53. Mouritsen, O. G. & Zuckermann, M. J. What's so special about cholesterol? *Lipids* **39**, 1101–1113

(2004).

54. Lampert, A. *et al.* A Pore-blocking Hydrophobic Motif at the Cytoplasmic Aperture of the Closed-state Nav1.7 Channel Is Disrupted by the Erythromelalgia-associated F1449V Mutation. *J. Biol. Chem.* **283**, 24118–24127 (2008).

55. Payandeh, J., Scheuer, T., Zheng, N. & Catterall, W. A. The crystal structure of a voltage-gated sodium channel. *Nature* **475**, 353–358 (2011).

56. Xiao, J. *et al.* Regulation and drug modulation of a voltage-gated sodium channel: Pivotal role of the S4–S5 linker in activation and slow inactivation. *Proc. Natl. Acad. Sci.* **118**, e2102285118 (2021).

57. Pristerà, A., Baker, M. D. & Okuse, K. Correction: Association between Tetrodotoxin Resistant Channels and Lipid Rafts Regulates Sensory Neuron Excitability. *PLOS ONE* **8**, 10.1371/annotation/55db09f9 (2013).

58. Abraham, M. J. *et al.* GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. *SoftwareX* **1**, 19–25 (2015).

59. Lindahl, Abraham, Hess & Spoel, van der. GROMACS 2019.4 Source code. Zenodo https://doi.org/10.5281/zenodo.3460414 (2019).

60. Marrink, S. J., Risselada, H. J., Yefimov, S., Tieleman, D. P. & de Vries, A. H. The MARTINI Force Field: Coarse Grained Model for Biomolecular Simulations. *J. Phys. Chem. B* **111**, 7812–7824 (2007).

61. Marrink, S. J., de Vries, A. H. & Mark, A. E. Coarse Grained Model for Semiquantitative Lipid Simulations. *J. Phys. Chem. B* **108**, 750–760 (2004).

62. Monticelli, L. *et al.* The MARTINI Coarse-Grained Force Field: Extension to Proteins. *J. Chem. Theory Comput.* **4**, 819–834 (2008).

63. de Jong, D. H. *et al.* Improved Parameters for the Martini Coarse-Grained Protein Force Field. *J. Chem. Theory Comput.* **9**, 687–697 (2013).

64. Šali, A. & Blundell, T. L. Comparative Protein Modelling by Satisfaction of Spatial Restraints. *J. Mol. Biol.* **234**, 779–815 (1993).

65. Fiser, A., Do, R. K. G. & Šali, A. Modeling of loops in protein structures. *Protein Sci.* **9**, 1753–1773 (2000).

66. Lomize, M. A., Pogozheva, I. D., Joo, H., Mosberg, H. I. & Lomize, A. L. OPM database and PPM web server: resources for positioning of proteins in membranes. *Nucleic Acids Res.* **40**, D370–D376 (2012).

67. Damre, M., Marchetto, A. & Giorgetti, A. MERMAID: dedicated web server to prepare and run coarse-grained membrane protein dynamics. *Nucleic Acids Res.* **47**, W456–W461 (2019).

68. Smart, O. S., Goodfellow, J. M. & Wallace, B. A. The pore dimensions of gramicidin A. *Biophys. J.* **65**, 2455–2460 (1993).

69. Harris, C. R. *et al.* Array programming with NumPy. *Nature* **585**, 357–362 (2020).

70. Hunter, J. D. Matplotlib: A 2D Graphics Environment. *Comput. Sci. Eng.* **9**, 90–95 (2007).

71. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **17**, 261–272 (2020).

72. Buchoux, S. FATSLiM: a fast and robust software to analyze MD simulations of membranes. *Bioinformatics* **33**, 133–134 (2017).

73. Humphrey, W., Dalke, A. & Schulten, K. VMD: Visual molecular dynamics. *J. Mol. Graph.* **14**, 33–38 (1996).