The acoustic adaptation hypothesis across terrestrial vertebrates: a meta-analysis

Bárbara Freitas [1,2]*, Pietro B. D'Amelio [3], Borja Milá [1], Christophe Thébaud [2], Tim Janicke [4]

[1] National Museum of Natural Sciences, Spanish National Research Council (CSIC), Madrid, Spain

[2] Centre de Recherche sur la Biodiversité et l'Environnement (UMR 5300 CNRS-IRD-TINPT-UPS),

Université Paul Sabatier, 31062 Toulouse Cedex 9, France

[3] Department of Behavioral Neurobiology, Max Planck Institute for Biological Intelligence, Seewiesen

82319, Germany

[4] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

*Correspondence to:

Bárbara Freitas

Museo Nacional de Ciencias Naturales

Calle José Gutiérrez Abascal 2

Madrid 28006, Spain

E-mail: barbara.freitas@mncn.csic.es

Author ORCIDs

Bárbara Freitas http://orcid.org/0000-0001-6020-6236

Pietro B. D'Amelio http://orcid.org/0000-0002-4095-6088

Borja Milá http://orcid.org/0000-0002-6446-0079

Christophe Thébaud http://orcid.org/0000-0002-8586-1234

Tim Janicke http://orcid. org/0000-0002-1453-6813

ABSTRACT

Acoustic communication plays a prominent role in various ecological and evolutionary processes

involving social interactions. The shape of acoustic signals is thought to be influenced not only by the

interaction between a signaller and a receiver but also by the acoustic characteristics of the

environment through which the signal is transmitted, which is the essence of the so-called "acoustic

adaptation hypothesis" (AAH). Specifically, the AAH posits that habitat structure affects frequency

and temporal parameters of acoustic signals emitted by a signaller depending on their acoustic

degradation properties. To date, this hypothesis has received mixed results, with the level of support

depending on the taxonomic group under study and the methodology used. We conducted a

systematic literature search of empirical studies testing for an effect of vegetation structure on

acoustic signalling and assessed the generality of the AAH using a meta-analytic approach based on

371 effect sizes from 75 studies and 57 taxa encompassing birds, mammals and amphibians. Overall,

our results do not provide consistent evidence for the AAH, neither in within-species comparisons

(suggesting no overall phenotypically plastic response of acoustic signalling to habitat structure) nor

in among-species comparisons (suggesting no overall evolutionary response). However, when

considering birds only, we found a weak support for the AAH in within-species comparisons, which

was mainly driven by studies that measured frequency bandwidth, suggesting that this variable may

exhibit a phenotypically plastic response related to habitat. For among-species comparisons in birds,

we also found support for the AAH, but this effect was not significant after excluding comparative

studies that did not account for phylogenetic non-independence. Collectively, our synthesis does not

support a universal role of vegetation structure in the evolution of acoustic communication. We

highlight the need for more empirical work on currently under-studied taxa such as amphibians,

mammals, and insects. Furthermore, we outline a framework for forthcoming research on the AAH.

Specifically, we advocate for a more detailed and quantitative characterization of habitats allowing

to identify frequencies with the highest detection probability and to determine if frequencies with

greater detection distance are preferentially used. Finally, we stress that research efforts should

focus on signals in which an increased distance of transmission is beneficial to senders and/or

receivers.

Key words: acoustic communication, amphibians, bioacoustics, birds, evolution, habitat, mammals,

vegetation, vocalizations

I. INTRODUCTION

(1) Acoustic signalling and the effect of multiple selective forces

Acoustic signalling is an essential facet of animal communication, spanning a wide array of species and ecosystems. These signals play a pivotal role in various contexts, including mate attraction, territory defence, predation avoidance (alerting others of a threat), rival intimidation, individual recognition, and social communication (Catchpole & Slater, 2008; Bradbury & Vehrencamp, 2011). Thus, acoustic signals are subject to multiple selective forces imposed on the signaller and receiver that shape them to finely tuned instruments of communication (Wiley & Richards, 1982; Catchpole, 1987; Catchpole & Slater, 2008). These multiple selective forces may act in concert or in opposition, which potentially complicates our understanding of the evolution of acoustic communication (i.e., the adaptive significance of the specific structure of a given signal).

One of the most studied selective forces shaping acoustic signals is the morphology of the signaller. For instance, the size of sound-producing organs is often positively associated with a signaller's body size or mass and has been shown to influence signal frequency in anurans and mammals (Fletcher, 2004; Tonini *et al.*, 2020). Similarly, in birds body size has been found to be negatively correlated with signal minimum frequency and positively with frequency bandwidth (Friis *et al.*, 2022). Thus, when individuals emit sounds, they also convey information about their size such that acoustic cues can serve as honest signals of body condition and reproductive performance, influencing mate choice (McLean, Bishop & Nakagawa, 2012; Welklin *et al.*, 2023). While the production of acoustic signals at high amplitude could attract more mates and give competitive advantages, it involves energetic costs (e.g., Gil & Gahr, 2002; Ophir, Schrader & Gillooly, 2010), to the extent that it may become challenging to produce an optimally designed sound for long-distance transmission. In addition, predators and parasites can use the acoustic signals emitted by their prey as cues to locate them (Ryan, Tuttle & Taft, 1981; Garamszegi & Avilés, 2005), potentially driving the development of signals that minimize their detectability, such as high-frequency pure tones in certain alarm calls to reduce locatability (Klump & Shalter, 1984). In vocal learners such as oscine

passerines, hummingbirds, parrots, humans, elephants, seals, and toothed whales, vocalizations are

also shaped by the capability of reproducing heard sounds (Tyack, 2020). This ability might

counterbalance other predicted trends in shaping acoustic signals. Likewise, interactions with other

animals can shape the acoustic signals in multiple ways, for example by selecting for frequencies

that exploit sensory biases to enhance engagement (Cummings & Endler, 2018).

The structure of acoustic signals can also be influenced by the acoustic characteristics of the

environment (Chappuis, 1971; Slabbekoorn & Smith, 2002; Derryberry *et al.*, 2018). While

propagating from signaller to receiver, signals are subject to several alterations in temporal and

frequency parameters. These alterations include reverberation, the delay between the faster waves

(direct) and slowest waves (indirect) causing distortion of the sound (Morton, 1975; Wiley &

Richards, 1978, 1982), and attenuation, the progressive reduction in amplitude which can be

attributed to three main components: (1) geometric attenuation (also known as spreading loss or

spherical attenuation); (2) atmospheric attenuation; and (3) habitat attenuation (Embleton, 1996).

The latter two components are often collectively referred to as degradation, resulting from

scattering, refraction through obstacles, atmospheric absorption, or echoes, with greater effects

typically observed for high frequencies (Aylor, 1972; Marten & Marler, 1977). In this context, natural

selection is expected to have favoured the evolution of accurate and efficient transmission of

acoustic signals intended for long-distance travel allowing for information transfer from signaller to

receiver with respect to the constraints imposed by their environment.

(2) The Acoustic Adaptation Hypothesis

The various selective forces imposed by the environment outlined above form the

theoretical background of the acoustic adaptation hypothesis (AAH), mainly grounded on the work

of Morton (1975) and Hansen (1979) but coined by Rothstein & Fleischer (1987). The AAH argues

that the transmission of acoustic signals is affected by the environment offering predictions

regarding the optimal shape of acoustic signals to ensure accurate transmission in environments

characterized by high degradation, such as densely vegetated habitats (Morton, 1975; Blumenrath &

Dabelsteen, 2004). Animals inhabiting such hereafter called 'closed habitats' are expected to have

signals of longer duration because this is expected to increase the likelihood of detection and to

leverage the amplitude of reverberated sound waves, extending the range of signal propagation

(Nemeth *et al.*, 2006). To increase transmission distance in closed habitats, acoustic signals should

also have lower repetition rates to minimize interference with reverberated waves (Brown &

Handford, 2000; Naguib, 2003) and narrower frequency bandwidths to concentrate energy in

frequencies less susceptible to attenuation (Wiley & Richards, 1978). Additionally, animals in closed

habitats should be prone to emit acoustic signals at lower frequencies including minimum, mean,

maximum, and peak frequency (Chappuis, 1971; Morton, 1975; Marten, Quine & Marler, 1977).

The AAH is currently considered as a component of the broader Sensory Drive Hypothesis,

which predicts that signals, sensory mechanisms, and microhabitat choice evolve together based on

the characteristics of the habitat, ambient noise patterns, the existence of predators and parasites,

and various sensory and physiological factors (Endler, 1992; Cummings & Endler, 2018). Several

authors have also extended the AAH to a wider framework of environmental adaptation, covering

studies on the effect of environmental noise such as tests of the Lombard effect in the context of

urbanization (Job, Kohler & Gill, 2016; Zhao *et al.*, 2021). Here, we focus on the AAH in the context of

the effect of vegetation structure on acoustic signalling as originally tested by Morton (1975).

Playback transmission experiments using either natural or synthetic sounds have been

conducted to assess the physical foundations of the AAH (Wiley & Richards, 1978). In addition,

comparisons between species and populations inhabiting environments of different vegetation

structure have been performed to test the validity of this hypothesis. While studies testing

environmental filtering of sound have provided a robust foundation for the AAH (reviewed in Hardt

& Benedict, 2021), its influence on animal acoustic signal production remains unclear. In fact,

empirical support for the AAH is highly variable, with many studies supporting the hypothesis

(Bertelli & Tubaro, 2002; Nicholls & Goldizen, 2006) whereas many others contest its validity (Hylton

& Godard, 2001; Bosch & De la Riva, 2004; Mikula *et al.*, 2020). In addition, Hardt & Benedict (2021)

uncovered that animal signals are not better transmitted in a species' habitat compared to

environments they are not adapted to, raising further questions about the generality of the AAH.

In a previous meta-analysis limited to birds, Boncoraglio & Saino (2007) compiled 91 effect

sizes from 26 studies of within- (including 12 species) and among-species (spanning 3 orders)

comparisons to test for an overall effect of habitat structure (open versus closed) on acoustic

variables. The authors found support for the AAH in terms of a lower minimum frequency, maximum

frequency, peak frequency, and frequency bandwidth in more closed habitats based on a global

effect size of Fisher's $z = 0.160$, 0.182, 0.346, and 0.230, respectively (Boncoraglio & Saino, 2007).

Interval duration was the only variable that did not differ between habitats. Since its publication, this

study has served as a key reference in support of the AAH in birds. However, Boncoraglio & Saino

(2007) relied on a limited sample size and did not include measurements of mean frequency, total

duration, or repetition rate in their analyses, despite clear predictions for these three variables

within the framework of the AAH. Furthermore, this study did not explicitly control for potential

sources of statistical non-independence beyond shared study identity, such as shared phylogenetic

history. The lack of control for non-independence among effect sizes can result in inaccurate

conclusions, often leading to spurious statistical significance (Nakagawa *et al.*, 2017), and integrating

phylogenetic structure into meta-analyses is considered critical to ensure that unbiased results are

obtained (Lajeunesse, 2009; Chamberlain *et al.*, 2012).

Although the AAH was initially formulated based on avian studies, there has been a growing

trend in testing this hypothesis in other taxonomic groups. Aiming to examine how widespread the

impact of vegetation structure on the evolution of acoustic signals actually is in vertebrates, Ey &

Fischer (2009) qualitatively reviewed 55 studies on birds, anurans and mammals and concluded that

the AAH did not seem to be universally valid across taxa. However, the authors claimed that in birds,

minimum frequency and frequency bandwidth appeared to align with the predictions of the AAH,

whereas repetition rate, mean, peak, and maximum frequencies did not, which does not fully

correspond to the results obtained by Boncoraglio & Saino (2007).

(3) Assessing the AAH across animals

Despite the large body of research aimed at testing the AAH, it remains unclear to what extent this

hypothesis is valid across acoustic variables in a broad range of animal taxa, and whether contrasting

findings from previous studies reflect methodological artifacts and/or actual differences among taxa.

Our study aims to fill this gap by conducting a meta-analysis of empirical studies having tested the

AAH in the context of airborne acoustic signalling in response to vegetation structure. In this regard,

we build on the work by Boncoraglio & Saino (2007) and aim at providing a robust and

comprehensive meta-analysis. More specifically, our main objective is to address two main

questions: (1) Does the AAH hold across different animal taxa and acoustic parameters? and (2) does

the AAH support phenotypically plastic and/or evolutionary responses to vegetations structure

(inferred from within- and among-species comparisons, respectively)? For these purposes, we

compiled 371 effect sizes testing the AAH in animals. We did not have a clear prediction on whether

effect sizes should differ between within- and among-species comparisons. However, it seems

plausible that effect sizes from among-species comparisons might be weaker, because acoustic

parameters likely evolve not only in relation to the habitat itself but also due to other factors such as

interspecific variation in mate preferences and/or genetic drift.

II. METHODS

This meta-analysis was pre-registered at the Open Science Framework (OSF; study details available

at: https://doi.org/10.17605/OSF.IO/E7JRS). Differences between the pre-registered study design

and the eventually implemented approach are mainly due to methodological improvements and are

explained in Text S1.

(1) Literature search

We conducted a systematic literature search to obtain an unbiased sample of studies testing for a relationship between vegetation structure and acoustic communication in animals following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses approach (PRISMA; Moher *et al.*, 2009) in combination with the PRISMA-EcoEvo guidelines (O'Dea *et al.*, 2021). We screened the Scopus, the ISI Web of Science Core Collection (Clarivate Analytics), and the Open Access Theses and Dissertations (OATD; https://oatd.org/) databases on 16th December 2021 with the search terms defined as ((("*song" OR "vocal*") OR (("acoustic$" OR "bioacoustic$") AND "call$")) AND ("habitat$" OR "vegetation")) OR "acoustic adaptation". These searches yielded 3300, 3233, and 372 records, respectively. We also considered studies included in previous reviews on related questions (Ey & Fischer, 2009; Hardt & Benedict, 2021) and a previous meta-analysis on birds (Boncoraglio & Saino, 2007), resulting in the addition of 58 studies. Furthermore, we posted a request for unpublished data on the 'evoldir' mailing list (http://life.mcmaster.ca/evoldir.html), the ResearchGate platform (https://www.researchgate.net/), and Twitter, and directly asked colleagues who work on these topics. This resulted in one additional study (Melo *et al.*, in prep). Finally, we added one unpublished study on the La Palma population of the Canary Islands Chaffinch (*Fringilla canariensis palmae*; Freitas *et al.*, in prep.).

We excluded 45 studies from the OATD search because full texts were not accessible. We also excluded duplicates by following an R workflow adapted from Lagisz (2022) and by manually checking the results of the Zotero deduplication algorithm (Corporation for Digital Scholarship & Roy Rosenzweig Center for History and New Media, 2022). Then, we assigned screening efforts among authors and screened the titles and abstracts of 4558 studies. We selected potentially relevant primary studies using the *metagear* package (Lajeunesse, 2016) based on the following selection criteria:

1. The focus of the empirical study was on non-human animals (excluding domesticated species), for which acoustic signals are transmitted through the air. We did not consider

studies focusing on echolocation because it is used for orientation in space and/or to locate

items rather than for communication. For syntheses on the effect of habitat structure on

echolocation, see Ey & Fischer (2009). We also excluded studies on aquatic habitats due to

the different acoustic properties and difficulties in defining habitat structure based on

underwater vegetation.

2. At least one temporal or frequency parameter for which the AAH predicts an effect was

measured for adult individuals of the same species (within-species comparisons) or multiple

species (among-species comparisons) across different habitat types. Studies testing for

differences among subspecies were considered as among-species comparisons. Studies that

did not replicate habitat type within species or subspecies were excluded (as a conservative

approach to mitigate the possibility of testing subspecies differences rather than differences

due to habitat). For example, a study was excluded if testing for acoustic signal differences

between a subspecies inhabiting one location of habitat type A and another subspecies in

another location of habitat type B.

3. Habitat was described along a continuous gradient or based on a categorical classification of

vegetation structure (from open to closed habitats). If the categorical classification of the

habitat was done in more than the two categories, we pooled different habitat types into

two categories: 'open' and 'closed' (e.g., grassland, open field, and meadow grouped in

'open habitat'). We did not include studies examining differences between urban and rural

habitats.

4. The study was written in English, French, German, Portuguese, or Spanish according to the

languages that the authors were able to understand.

5. The study reported sufficient information to compute effect sizes (e.g., sample size, test-

statistic, means with estimates of uncertainty).

During the title and abstract screening stage, we excluded 4237 studies because they did not

fit at least one of the requirements listed above. Afterwards, we carried out a full-text screening of

the remaining 321 studies. During this stage we excluded 246 studies because they: (i) did not test

the influence of vegetation structure on acoustic variables (N = 116), (ii) used habitat types that

could not be classified as 'open' or 'closed' (N = 17), (iii) did not replicate habitat type (N = 5), (iv)

focused on echolocation (N = 23), (v) focused on playback/transmission experiments (N = 15), (vi)

focused on the effect on urbanization (N = 4), (vii) used acoustic variables for which the AAH does

not generate predictions (N = 3), (viii) did not report sufficient data (N = 27; i.e., multivariate

statistics not providing raw data, missing key data required to compute an effect size even after

contacting the authors), (ix) represented reviews or non-empirical studies (N = 14), (x) were

inaccessible (N = 12), (xi) were written in other languages (N = 4), or (xii) represented duplicates (N =

6; 5 of which represented theses that are already covered by other primary studies). The final

dataset included 75 primary studies (see Fig. 1 for the PRISMA diagram and Table S1 for a reference

list of all primary studies).

(2) Data collection

We extracted means, measures of variance (standard deviation or standard error) and sample sizes,

correlation coefficients and sample sizes, or the statistical test value and degrees of freedom

reported for the relationship among acoustic variables and vegetation structure from the text,

tables, figures or supplementary files. We used the R package *juicr* version 0.1 (Lajeunesse, 2021) to

extract summary statistics from figures. From five studies that used multivariate statistics (such as

principal components analysis), we reanalysed the raw data if published together with the article or

provided by the authors (N = 2), and corrected also for the phylogeny in cases in which phylogenetic

data was provided (N = 1).

All comparisons included in our meta-analysis followed a descriptive approach, meaning that

none of the primary studies manipulated the habitat experimentally to test for a causal effect on

acoustic variables. Thus, we used the Pearson correlation coefficient (r) as an effect size to

synthesize the findings of primary studies. We converted test statistics provided in the primary

studies into *r* and estimated its sampling variance (*Vr*) using formulas and guidelines reported in

Lajeunesse *et al*. (2013) and using the R package *esc* version 0.5.1 (Lüdecke, 2019). To improve

transparency and allow for reproducibility of data extraction, we report the method used to

compute effect sizes (Table S1) following a framework provided in Ivimey-Cook *et al*. (2023). All

effect sizes were assigned to a type of study: within- or among-species comparisons, to allow testing

for a phenotypically plastic or an evolutionary response of acoustic signalling to vegetation structure,

respectively. We interpreted the support for the AAH in within-species comparisons as suggestive of

a phenotypically plastic response rather than adaptive response since we assumed no genetic

divergence between the different populations tested.

(3) Moderator variables

Besides the global test for the support of the AAH across animals, we aimed at explaining

heterogeneity in effect sizes among studies from a biological and methodological perspective.

Specifically, we explored if support for the AAH differed among animal taxa and acoustic variables. If

an acoustic variable was measured for parts and for the full signal duration, we extracted both effect

sizes separately but assigned them as the same acoustic variable (e.g., song duration and note

duration were both considered as duration).

In addition, we tested for four methodological moderators: habitat classification, habitat

contrast, body size correction, and phylogeny correction. Firstly, we explored effects of the habitat

classification (continuous versus categorical) used in primary studies, i.e., if the studies used a

continuous variable or a categorical classification to describe the vegetation structure of the habitat.

Secondly, we scored the magnitude of contrast between closed and open habitats (for studies using

a categorical classification) that were used in the study (habitat contrast), attributing a '0' when

there was a moderate difference in vegetation structure (e.g., temperate broadleaf forest versus

tropical rainforest), or a '1' when there was a substantial difference in vegetation structure (e.g.,

grassland versus forest). Importantly, primary studies investigated various habitat types and used

different methods for classifying them. This diversity in habitat classification methods could have

contributed to differences in study outcomes (suggested by Hardt & Benedict, 2021). By considering

these two moderators, we tried to account for this potential source of heterogeneity among studies.

Thirdly, we recorded whether the analyses corrected for body size differences within and/or among

species, since this trait is known to be allometrically correlated with the morphology of the sound

apparatus, and therefore to affect the production of sound (Pearse *et al.*, 2018). Finally, we recorded

whether studies reporting among-species comparisons accounted for phylogenetic non-

independence.

(4) Phylogeny

In order to account for phylogenetic non-independence and to quantify the phylogenetic signal (also

termed phylogenetic heritability; de Villemereuil & Nakagawa, 2014), we reconstructed a phylogeny

of all sampled taxa from studies of within- (Fig. S1) and among-species comparisons (Fig. S2). We

used the R package *rotl* (Michonneau, Brown & Winter, 2016) to retrieve the phylogenetic

relationships from the Open Tree of Life (opentreeoflife.org/; Hinchliff *et al.*, 2015) and *ape* (Paradis,

Claude & Strimmer, 2004) to obtain the branch lengths for each taxon by applying Grafen's method

(Grafen, 1989).

In analyses of among-species comparisons, the phylogenetic position of some taxonomic

groups could not be assessed unequivocally so that we assigned the group in question to a higher

taxonomic level. Specifically, we considered Parulinae as Parulidae and Cardinalidae as

Passeriformes. Note that some primary studies tested the AAH across higher taxonomic ranks (such

as Anura, Aves and Passeriformes), while others tested it across a lower taxonomic rank, which

sometimes was nested within the higher ones (artificial pairs, e.g., Anura and Hylidae, Aves and

Corvus). In such cases, we retrieved pairwise estimates of divergence times for the taxonomic ranks

that were not nested from the TimeTree database (http://timetree.org/; Kumar *et al.*, 2017). Then,

for the artificial pairs, we attributed a mean estimate between the known divergence time estimates

306 of the nested pairs and the ones more distantly related. For example, for the pair Anura - Hylidae,

307 we computed the divergence time as the mean between the estimate of Hylidae - *Rhinella* (a genus

308 within Anura) and the estimate of Anura - Felidae, since Felidae is the closest outgroup of Anura in

309 our phylogeny. Then, we computed branch lengths using the methods already specified for the

310 within-species comparison.

311 **(5) Data analyses**

312 All analyses were performed in R 4.2.2 (R Core Team, 2022) through RStudio 1.4.1106 (RStudio

313 Team, 2021), using the packages *metafor* (Viechtbauer, 2010) for the meta-regression models, *brms*

314 (Bürkner, 2017) for models with Bayesian approach, *orchaRd* (Nakagawa *et al.*, 2023) for calculation

315 of heterogeneity for Restricted Maximum Likelihood (REML) models, *dplyr* (Wickham *et al.*, 2022) for

316 data manipulation, *ggplot2* (Wickham, 2016) for data visualization, and *ggtree* (Yu *et al.*, 2017) to

317 produce our final phylogenies as shown in Figs S1 and S2. Silhouettes in figures were extracted from

318 'phylopic' (https.phylopic.org) using *rphylopic* (Gearty, Jones & Chamberlain, 2023).

319 **(a) Meta-analyses**

320 To provide a global test of the AAH across animals and to assess the impact of moderator variables,

321 we ran both frequentist General Linear-Mixed Effects Models (GLMMs) using REML method and the

322 correspondent Bayesian models. Bayesian models were run with 4 chains, each with 4000 iterations,

323 a warmup of 2000 iterations, and were checked for convergence. We ran all models for datasets of

324 within- and among-species comparisons separately. First, we obtained a weighted mean effect size

325 (*r*) by fitting a model with no moderator variables but with study identifier, effect size identifier,

326 taxon name and the phylogeny (transformed into a correlation matrix) as random effects to account

327 for statistical nonindependence. This was done because the final dataset included multiple effect

328 sizes from species with shared phylogenetic history, from the same species and/or from the same

329 study. We also fitted separate models for each taxonomic class (i.e., Aves, Mammalia, Amphibia).

Second, we assessed if biological and methodological moderators impacted effect sizes.

Specifically, we ran models with the same random factors as above and defined acoustic variable

(minimum frequency, mean frequency, maximum frequency, frequency bandwidth, peak frequency,

duration, interval duration, and repetition rate), habitat classification (categorical versus

continuous), habitat contrast (0 or 1), body size correction (yes or no), or phylogenetic correction

(yes or no) as fixed factors. The impact of all moderators was tested separately within Aves, which

represents the taxonomic class with the highest number of effect sizes (see Results).

We assessed heterogeneity I^2 by estimating the proportion of variance among effect sizes

that can be attributed to the different levels of random effects (Higgins & Thompson, 2002).

Specifically, we partitioned total heterogeneity (I^2 Total) into variance arising from study identity (I^2

Study), observation (I^2 Observation), taxon (I^2 Taxon), and from phylogenetic relationship (I^2

Phylogeny; Nakagawa & Santos, 2012), which is also known as phylogenetic heritability (H^2), and

equivalent to Pagel's λ (de Villemereuil & Nakagawa, 2014). For REML models with predictor

variables, we calculated the pseudo-R^2 based on the proportional change in the sum of the variance

components, and its 95% confidence interval based on 10000 bootstrap replicates (Viechtbauer,

2023). For models run with the Bayesian approach, we computed the proportion of variance

explained by the entire model (conditional R^2) and by fixed factors only (marginal R^2; Nakagawa &

Schielzeth, 2013).

(b) Publication bias

In order to assess publication bias, we ran a multilevel meta-regression as suggested by Nakagawa *et

al*. (2022). In contrast to Pearson's correlation coefficient *r*, the sampling variance of Fisher's *z*

statistics only depends on the sample size but not on the effect size itself. Thus, we transformed our

effect size *r* into *z* and computed its sampling variance using guidelines by Borenstein *et al*. (2009).

Then, we ran a GLMM with *z* defined as response variable, its standard error, moderators, and year

as fixed effects and study identifier, observation identifier, taxon name and the phylogenetic

correlation matrix as random factors, which is advised when heterogeneity is high (see Results;

Nakagawa *et al.*, 2022). This aimed at testing (i) whether the effect size depends on its standard

error, which, similarly to the Egger's regression (Egger *et al.*, 1997), may indicate that small studies

get published only (or more often) if effect sizes are large enough to provide statistically significant

support for the tested hypothesis (small-study effect), and (ii) for time-lag bias (Boutron *et al.*, 2019;

also known as the bandwagon effect), which suggests that supportive results get published easier in

a newly emerging field but as more data are collected and scepticism about the theoretical

foundations may arise, non-intuitive results increase (e.g., Jennions & Møller, 2002).

As complementary, we used Kendall's rank correlation test (Begg & Mazumdar, 1994) to

quantify funnel plot asymmetry. Studies with low precision and/or high sampling error that do not

support the tested hypothesis might be more prone to go unpublished. Consequently, the lack of

studies meeting these criteria is expected to drive funnel plot asymmetry (Sterne *et al.*, 2011).

III. RESULTS

(1) Dataset summary

In total, we obtained 371 effect sizes from 75 studies and 57 taxa encompassing vertebrate taxa

only, i.e., amphibians, birds and mammals (Fig. 2 and Table S2). Of these, 219 effect sizes were from

studies reporting within-species comparisons (37 species, including 32 birds, 3 mammals, and 2

amphibians, with 189, 16, and 14 effect sizes, respectively) and 152 were from studies of among-

species comparisons (20 taxa, including 14 birds, 3 mammals, and 3 amphibians, with 129, 11, and

12 effect sizes, respectively). We obtained 57 effects from analyses of minimum frequency, 12 of

mean frequency, 55 of maximum frequency, 47 of frequency bandwidth, 74 of peak frequency, 71 of

duration, 20 of interval duration, and 35 of repetition rate (from 39, 8, 38, 31, 46, 39, 19, and 22

studies, respectively). Regarding habitat classification, we obtained 137 effects from studies using a

continuous variable to define the vegetation structure of the habitat (21 studies) and 234 with a

categorical definition (54 studies), of which 163 effects came from 39 studies comparing highly

dissimilar habitats (habitat contrast of 1) and 71 from 16 studies comparing more similar habitats

(habitat contrast of 0). In total, 49 effect sizes were obtained from 15 studies correcting for body size

(from which 41 were retrieved from among-species comparisons) whereas 322 effect sizes from 60

studies did not account for body size. Finally, 73 effect sizes from 19 studies are based on analysis of

among-species-comparisons correcting for phylogenetic non-independence and 79 effects from 18

studies not correcting for it.

Our dataset included 22 studies (approximately 29% of our dataset) that were already

included in a previous meta-analysis focusing on birds by Boncoraglio & Saino (2007). Nonetheless,

we obtained different effect sizes for 10 (45%) of these studies. Out of a total of 78 effect sizes

present in Boncoraglio & Saino (2007), 7 were identical with respect to its absolute value but having

the opposite sign, 11 differed in more than 0.1, with 4 of them having the opposite sign, 2 effect

sizes could not be retrieved, and 9 differed due to a different analytical approach (most probably

because, contrary to Boncoraglio & Saino (2007), we did not pool data from different syllables or

songs). Moreover, we did not include 13 effect sizes (from 4 studies) used in the previous meta-

analysis because the studies did not fit our selection criteria (i.e., habitats were impossible to classify

as open/closed, habitat was not replicated, the study focused on urbanization, or the reported data

were not sufficient to compute effect sizes). We report these inconsistencies in detail in Table S3.

(2) Test of the AAH for within- and among-species comparisons

Overall, we did not find support for the AAH, neither for within- nor among-species comparisons

(Figs 3 and S3, Tables 1 and S5). None of the tested moderator variables explained a significant

fraction of heterogeneity in effect sizes obtained from within- or among-species comparisons (Figs

S4 and S5, and Tables S4 and S6).

When focusing only on the most studied taxonomic class, Aves, we found weak support for

the AAH for within- and among-species comparisons (Figs 3 and S3). Frequency bandwidth showed a

weak support for the AAH in within-species comparisons (Fig. S6), whereas minimum, maximum and

peak frequencies showed significant support in among-species comparisons (Fig. S7). Analyses of

other acoustic variables did not show support for the AAH. Primary studies of among-species

comparisons obtained significantly higher effect sizes when they did not account for phylogenetic

non-independence than when they did (Tables S4 and S6). When focusing on primary studies that

accounted for phylogenetic non-independence, we did not obtain a significantly positive global

effect size for birds and no acoustic variables showed a significant effect either (Fig. S8).

In analyses of Mammalia and Amphibia, we found no support for the AAH neither in within-

nor in among-species comparisons (Figs 3 and S3, Tables 1 and S5). For these analyses, the number

of effect sizes and species was much lower compared to Aves, which did not allow testing for the

impact of moderators in these taxonomic classes.

The effect sizes we retrieved were highly heterogeneous (REML I^2 Total > 0.80 for all models;

Table 1). For studies of within-species comparisons, heterogeneity arose mostly from variance

among observations and heterogeneity arising from other factors varied depending on the model.

For studies of among-species comparisons, heterogeneity stemmed primarily from among-study

differences for the general model and the model including only birds. For the model including only

studies on birds that accounted for phylogenetic non-independence (phylogeny correction),

heterogeneity arose mostly from variance among species. In models including only mammals or

amphibians, phylogeny explained the highest proportion of total heterogeneity (in both within- and

among-species comparisons). These patterns were confirmed by the Bayesian approach, wherein

the heterogeneity estimates were generally higher for among-studies and among-observations

differences than for the other factors, except for models with only mammals or amphibians (Table

S5).

(3) Publication bias

We detected no evidence for the small-study effect either for within- and among-species

comparisons, since small studies (with smaller sample size and lower precision) did not show

stronger effect sizes than large studies (Figs S9A and B). In fact, for studies reporting within-species

comparisons, we detected the opposite trend, i.e., a significant negative relationship between effect

size and its standard error (Fig. S9A).

We also detected no evidence of the time-lag bias since the year of publication did not

predict effect sizes of within-species comparisons (slope = 0.003, 95% CI = [-0.006, 0.012]) or among-

species comparisons (slope = -0.003, 95% CI = [-0.012, 0.006]). We observed a negative relationship

for studies not accounting for phylogenetic non-independence and a slightly positive relationship for

species that accounted for it, but these relationships were not statistically significant (Figs S9G and

H).

Finally, we found no evidence for funnel plot asymmetry (Figs S9C, D, E, and F) inferred from

Kendall's rank correlation test (Kendall's tau [within-species comparisons] = -0.059, P = 0.212;

Kendall's tau [among-species comparisons] = -0.005, P = 0.928).

IV. DISCUSSION

The acoustic adaptation hypothesis (AAH) posits that animals produce acoustic signals that have

evolved in response to the structure of their habitat, but ever since its original formulation, there

has been much uncertainty about the hypothesis' validity and generality within and among species.

We analysed a dataset of 371 effect sizes across 75 studies and 57 vertebrate taxa to demonstrate

that acoustic variables do not differ significantly between habitats of different vegetation structure

in both within- and among-species comparisons. Therefore, current empirical evidence does not

support the idea that the AAH is universally valid. Only comparisons within bird species provided

weak support for this hypothesis indicating that variation in acoustic signals in relation to vegetation

structure may be more complex and context-dependent than previously thought. Below we discuss

these findings along with the limitations of our study and future directions for research.

(1) Limited support for the AAH

(1) Limited support for the AAH

Our phylogenetically informed synthesis provides no overall support for the AHH, offering

quantitative evidence that acoustic signals do not show predictable responses to vegetation

structure across vertebrates. This result might seem unexpected given the established principles of

the physics of sound transmission (Hardt & Benedict, 2021) and questions the broader utility of the

AAH for understanding the evolution of acoustic communication in natural settings.

We acknowledge that our final dataset exhibits a taxonomic bias with an overrepresentation

of birds. While this reflects a limitation of our study, it also underscores the evident need for

additional research examining the influence of vegetation on acoustic signals of taxa beyond birds—

a call that has been made by numerous authors before (e.g., Ey & Fischer, 2009; Goutte *et al.*, 2018;

Mikula *et al.*, 2020) but remains largely unaddressed. As research extends to taxa that are currently

underexplored, the inclusion of reptiles, insects, and more species of anurans and mammals would

likely enable the examination of other potential moderators, such as signal development (i.e.,

innate versus learned acoustic signals), suggested by other authors to influence results, but that

could not be tested in this study (see Text S1). This would ultimately contribute to a more

comprehensive understanding of acoustic adaptation and animal communication.

The substantial variation in effect size across our dataset (REML I^2 Total > 0.80 for all the

models; Table 1), a common pattern of meta-analyses in ecology and evolution (with the average I^2

being as high as 92%; Senior *et al.*, 2016), remained largely unexplained. For within-species

comparisons, most of the heterogeneity originated from variation among observations, while

differences among studies, taxa or phylogeny had less impact. This among-observations variability

may reflect the within-studies contrasting support of the AAH depending on the acoustic variables

analysed, sometimes even within the same acoustic variables when measured for the entire signal

and its fragments (i.e., for studies that measured acoustic variables across the entire song as well as

its syllables). On the other hand, the heterogeneity of among-species comparisons arose mostly

from variation among studies, which could be due to differences among study designs such as using

a different assessment of vegetation structure, analysing different acoustic variables, correcting or

not for body size or phylogenetic non-independence. Remarkably, only a fraction of primary studies

of among-species comparisons accounted for phylogenetic non-independence and our analyses

suggest that this methodological aspect has an impact on the study's outcome, particularly in birds.

In contrast to a qualitative review by Ey & Fischer (2009), our results suggest that studies not

accounting for phylogenetic non-independence obtained stronger support for the AAH than those

that did. This underscores the need for comparative studies to consider phylogenetic relatedness to

avoid misinterpreting the role of vegetation structure in shaping acoustic signals.

Why then does the AAH not seem to apply universally? Several factors may contribute to the

limited support of the AAH. First, the complexity of the habitat and the methodological difficulties

regarding its classification could render the AAH an oversimplified concept if applied too broadly.

Second, the diversity of acoustic signals across taxa and the multiplicity of selective pressures acting

on them may lead to complex and diverse responses that may override a potential effect of the

habitat.

(a) Complexity of the habitat

Although the AAH predicts that selection should favour acoustic signals that propagate well in an

environment, it has been primarily tested in terms of effects of vegetation on sound transmission.

This approach might be too simplistic, as many other elements may influence sound propagation in a

given habitat (Attenborough & Renterghem, 2021). Physical features of the landscape (hills, valleys,

or slopes), type of substrate, weather conditions, as well as background noise (arising from wind,

moving water, or other species) are only a few of the factors that can also affect how sound travels

(Sueur, Krause & Farina, 2019; Guibard *et al.*, 2022). For example, there is increasing evidence that

animals can adjust their acoustic signals to surrounding noise (Slabbekoorn & Peet, 2003; Gil *et al.*,

2015; Gomes *et al.*, 2022). The significant heterogeneity detected in our analyses probably reflects

variation among studies arising not only from methodological aspects including differences in data

collection, analysis, and reporting but also from the physical features of the habitat.

Even when focusing on vegetation only, characterizing the habitat proves to be a challenging

and not yet standardized task. Early work on the AAH used broad habitat categories (forest, edge,

and grassland; Morton, 1975) and general predictions were then formulated for open versus closed

habitats. These classifications of vegetation structure do not only oversimplify specific habitat

characteristics crucial for sound transmission but also tend to be subjective and prone to ambiguity.

Even the use of more quantitative continuous variables does not resolve this issue because a wide

array of both small-scale (measured at a microhabitat level, e.g., number of trees) and large-scale

(representative of macrohabitat characteristics, e.g., tree cover, canopy cover) variables have been

used, which challenges to generalization of findings. The differences in the assessment of vegetation

structure among studies may have contributed to the heterogeneity in our results as suggested by Ey

& Fischer (2009). However, we found that support for the AAH did not significantly differ among

studies using categorical versus continuous characterization of vegetation structure, or among

studies with varying magnitudes of contrast between closed and open habitats. This indicates that

including different classifications of the habitat did not change the overall results, possibly reflecting

the lack of power of the current classification to capture the environment structure. Additionally,

within the same habitat, both season and the time of day can have a significant influence on signal

transmission. For example, an empirical study conducted in two different forests demonstrated that

these factors impacted detection distance by up to a factor of five (Haupert, Sèbe & Sueur, 2022).

(b) Alternative determinants of signal evolution overriding habitat effects

Animal acoustic signals are shaped by a multitude of selective forces, which can give rise to

contrasting responses, potentially reducing the significance of habitat effects, thus challenging the

universality of the AAH. Whilst our results provide no global support for the AAH, we found weak

evidence for the AAH in birds for within-species comparisons. Avian acoustic signals might therefore

present phenotypic plasticity in response to different vegetation types but the weak support

observed in our analysis suggests that habitat structure plays a minor role, if any, in shaping acoustic

signals even in birds. The lack of support for the AAH in Mammalia and Amphibia aligns with the

previous qualitative review (Ey & Fischer, 2009) and may reflect the limited importance of habitat in

explaining global signal variation in these taxa. Nevertheless, these results demand caution due to

the limited number of species tested for these animal classes.

Further complicating the application of the AAH as a universal framework are diverse

constraints associated with each acoustic variable, that also depend on the taxa. Anuran calls, for

instance, exhibit greater morphological constraints on frequency than on temporal parameters

(Escalona Sulbarán *et al.*, 2019; Tonini *et al.*, 2020), and this has been suggested as the reason for

temporal parameters being more susceptible to environmental selection than spectral ones

(Mendoza-Henao *et al.*, 2023). For birds, there seems to be no consensus regarding the acoustic

parameters that are more prone to be selected by environmental factors. While the previous meta-

analysis found that all frequency variables differed between habitats, consistently supporting the

AAH (Boncoraglio & Saino, 2007), Ey & Fischer (2009) identified minimum frequency and frequency

bandwidth as the only acoustic variables presenting global support for the AAH. Recent findings

propose frequency bandwidth as a more labile acoustic variable, in opposition to peak frequency,

which is more constrained by a species' genetic background (Rivera *et al.*, 2023). Although our

results focussing on plastic changes (i.e. within-species comparisons) align with the findings from

Rivera *et al*. (2023), this is not the case for the results obtained for evolutive changes (i.e. among-

species comparisons), which reveal a lack of support by all the acoustic variables when using studies

that accounted for phylogeny.

Even if animals do not adjust acoustic parameters, they can exhibit other adaptive communication

strategies to optimize signal transmission. In particular, they may move to a more suitable

microhabitat and height for broadcasting (Kime, 2000; Seddon, 2005; Chitnis, Rajan & Krishnan,

2020) or for listening (Jensen, Larsen & Attenborough, 2008). In the presence of background noise

from biological, abiotic, or urban sources, animals may also adjust the broadcasting time to minimize

signal overlap and competition for 'signal space' with other sound-producing species (Hart *et al.*,

2015; Chronister, Rhinehart & Kitzes, 2023; Staniewicz *et al.*, 2023). Ultimately, the benefits of long-

distance transmission, such as improved communication over larger distances or enhanced mate

attraction, may be offset by the predatory, competitive, and parasitic costs associated with it (Ryan

et al., 1981; Garamszegi & Avilés, 2005).

(3) Changing the paradigm

As Brown & Handford (1996) concluded: "*With all the possible selective forces acting on signal*

design, investigators looking for specific or finely detailed differences within or among species, across

habitats, should not be surprised to find results that do not always agree with AAH predictions". This

conclusion is echoed in more recent work (e.g., Ey & Fischer, 2009; Goutte *et al.*, 2018; Mikula *et al.*,

2020) but the AAH has remained a focal point of investigation.

The lack of support for the AAH in our analyses calls for a more comprehensive and

integrative approach to study acoustic adaptation, considering the complex interplay of ecological

and evolutionary factors. We propose this field to move beyond the current confines of the AAH by

highlighting promising avenues for future studies.

(a) Standardizing habitat attenuation measurements to refine AAH predictions

We believe that advances in understanding how environmental features affect acoustic signalling

can be achieved through a different approach to how the habitat is characterised and analysed. This

involves refining the description of the habitat and adopting a broader ecosystem perspective,

moving beyond a sole focus on vegetation structure. Ey & Fischer (2009) suggested a detailed

characterization of each habitat, including altitude, humidity, temperature, local animal acoustic

signals, and vegetation structure. However, obtaining and interpreting such an extensive amount of

information at the relevant scale for communication may pose practical challenges due to the

multitude of factors involved and difficulties in comparing studies. Instead of considering multiple

factors of the habitat separately, it may be preferable to use a single variable that integrates the

direct effects of multiple factors on sound propagation.

A promising approach to characterise the acoustic properties of the environment has been

emphasised recently by Haupert *et al*. (2022). By recording ambient sound and broadcasting white

noise and recording it at various distances from the emitter in two forest types, they found that the

coefficient of attenuation (a_0, expressed in dB/kHz/m) can describe with high accuracy the

attenuation characteristic of an environment for large frequency bandwidths (for 5 kHz frequency

bandwidths between 0-20 kHz in their study). This singular parameter is modelled based on acoustic

physics rather than descriptive statistics and summarizes the contribution of various variables,

including temperature, relative humidity, atmospheric pressure, distance, frequency, ambient sound

pressure level, and vegetation, to global attenuation (Haupert *et al.*, 2022). Moreover, Haupert *et al*.

(2022) used the same playback technique to describe the attenuation spectrum (the pattern of

attenuation by frequency) of the two studied forests. Despite having similar a_0 values, both forests

exhibited very different attenuation spectrum profiles. Within the same environment, detection

distance varied by up to a factor of five depending on the frequency considered. These spectrum

profiles potentially reveal frequencies with the highest detection probability and could be used as

starting point to describe the habitat. Subsequently, studies could establish correlations between

the attenuation profile and the species' (or population's, or even individual's) specific power

spectrum of acoustic signals, in order to test whether frequencies with greater detection distance

are preferentially used. Therefore, we suggest a more comprehensive examination of attenuation in

each locality, employing methodologies such as those outlined by Haupert *et al*. (2022). This would

facilitate global comparisons (through a single variable) and offer a more objective and standardized

assessment of the acoustic properties of a given habitat.

Furthermore, we propose to rephrase the predictions of the AAH to take into account the

attenuation spectrum. Specifically, animals inhabiting environments with different attenuation

spectra are predicted to emit acoustic signals at frequencies (minimum, mean, maximum, and peak

frequency) such that detection distance is maximized. This implies that there is no universal pattern

for the expected change in these variables, unless habitats share similar profiles. Instead, the

alterations will be contingent upon the specific characteristics of the environment. This adjustment

also overcomes an early noticed limitation of the AAH that for maximum efficiency, long-range

acoustic communication in any habitat should employ the lowest possible frequencies (Wiley &

Richards, 1982).

(b) Towards a narrower framework to test the AAH

While the AAH offers valuable insights, it may not fully capture the processes underlying the

evolution of acoustic signalling as it does not account for alternative strategies that animals use in

response to multiple selective pressures acting on signal structure or their interactions. To

comprehensively assess the environmental influence on signal evolution, a more comprehensive

approach is necessary.

Considering the probable lack of generality of the AAH, it would be valuable to narrow down

the focus to signals for which an advantage in transmission over long distances is likely. Studies not

selecting for these signals (e.g., Mikula *et al.*, 2020) might reflect this limitation. For example, many

acoustic signals are part of private duets (i.e., do not need long-distance transmission) such that

there is no selection against easily degrading communication (Larsen, 2020). Likewise, for some

species song is not used for distant communication, as reported in zebra finches, where it may be

more pertinent to assess whether their distance call, rather than their song, is adapted for long

distance transmission through the local environment (Loning, Griffith & Naguib, 2022). This

approach requires an in-depth knowledge of the study species, which might not always be readily

available and can be challenging to obtain. Nevertheless, we believe that tests of the AAH are only

meaningful for signals that are meant for long-distance communication.

V. CONCLUSIONS

(1) Our meta-analysis challenges the general validity of the acoustic adaptation hypothesis (AAH) by showing no overall support for the correlation between vegetation structure and acoustic variables.

(2) The weak support for the AAH in within-species comparisons of birds, mostly driven by frequency bandwidth differences, suggests a phenotypically plastic response of this variable to the vegetation structure of the habitat. By contrast, the AHH is not supported by among-species comparisons of studies accounting for phylogenetic non-independence.

(3) Although many years have passed since the formulation of the AAH, major efforts are still required to confirm the predictive ability of the hypothesis. Future research should refine the characterization of the habitat through a standardized quantification of the attenuation, which will lead to habitat- and species-specific predictions. Moreover, we expect that major advances in the field will benefit from broadening the studied taxonomic range while concurrently being selective about the tested signals.

VI. ACKNOWLEDGEMENTS

We are very grateful to the authors of the primary studies for making their research available, and in particular those providing additional data or information, and unpublished datasets, including Emma Greig, Elodie Ey, Darren Proppe, and Martim Melo. T. Michael Keesey, Julián Bayona, and Dennis Murphy designed the silhouettes used in our figures. We thank Sylvain Haupert, Ole Næsbye Larsen, Frédéric Sèbe, and Keith Attenborough for their insightful comments and discussions, which significantly improved the manuscript. We extend our gratitude to Rémi Tournebize for his assistance with troubleshooting R code errors.

Funding: B.F. was funded by the Foundation for Science and Technology FCT, Portugal) through a PhD grant (2020.04569.BD). This research was partly funded by grant PID2021-124501NB-I00 to B.M. from the Spanish Ministry of Science and Innovation, co-financed by the European Union's

Regional Development Fund (ERDF). C.T. benefited from a grant managed by the Agence Nationale

de la Recherche (CEBA: ANR-10-LABX-25-01). T.J. received funds from the Centre National de la

Recherche Scientifique (CNRS).

VII. AUTHOR CONTRIBUTIONS

B.F. Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project

administration, Visualization, Writing – original draft;

P.B.D. Writing – review and editing

B.M. Supervision, Writing – review and editing

C.T. Supervision, Writing – review and editing

T.J. Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project

administration, Supervision, Validation, Visualization, Writing – review and editing

VIII. DATA AVAILABILITY

All data and R scripts associated with this study can be retrieved from the OSF data repository

(available at xxx; it will be made available upon article acceptance).

IX. REFERENCES

ATTENBOROUGH, K. & RENTERGHEM, T. VAN (2021) *Predicting outdoor sound* Second edition. CRC Press,
Boca Raton.

AYLOR, D. (1972) Noise Reduction by Vegetation and Ground. *The Journal of the Acoustical Society of*
America **51**, 197–205.

BEGG, C.B. & MAZUMDAR, M. (1994) Operating Characteristics of a Rank Correlation Test for
Publication Bias. *Biometrics* **50**, 1088.

BERTELLI, S. & TUBARO, P.L. (2002) Body mass and habitat correlates of song structure in a primitive
group of birds. *Biological Journal of the Linnean Society* **77**, 423–430.

BLUMENRATH, S. & DABELSTEEN, T. (2004) Degradation of Great Tit (*Parus major*) Song Before And After
Foliation: Implications for Vocal Communication in a Deciduous Forest. *Behaviour* **141**, 935–
958.

BONCORAGLIO, G. & SAINO, N. (2007) Habitat structure and the evolution of bird song: a meta-analysis
of the evidence for the acoustic adaptation hypothesis. *Functional Ecology* **21**, 134–142.

BORENSTEIN, M., HEDGES, L.V., HIGGINS, J.P.T. & ROTHSTEIN, H.R. (2009) *Introduction to Meta-Analysis*.
John Wiley & Sons, Chichester, West Sussex, UK.

BOSCH, J. & DE LA RIVA, I. (2004) Are frog calls modulated by the environment? An analysis with anuran
species from Bolivia. *Canadian Journal of Zoology* **82**, 880–888.

BOUTRON, I., PAGE, M.J., HIGGINS, J.P., ALTMAN, D.G., LUNDH, A., HRÓBJARTSSON, A., & ON BEHALF OF THE
COCHRANE BIAS METHODS GROUP (2019) Considering bias and conflicts of interest among the
included studies. In *Cochrane Handbook for Systematic Reviews of Interventions* (eds J.P.T.
HIGGINS, J. THOMAS, J. CHANDLER, M. CUMPSTON, T. LI, M.J. PAGE & V.A. WELCH), pp. 177–204, 1st
edition. Wiley.

BRADBURY, J.W. & VEHRENCAMP, S.L. (2011) *Principles of animal communication* 2nd edition. Sinauer
Associates, Sunderland, Mass.

BROWN, T.J. & HANDFORD, P. (1996) Acoustic signal amplitude patterns: A computer simulation
investigation of the acoustic adaptation hypothesis. *The Condor* **98**, 608–623.

BROWN, T.J. & HANDFORD, P. (2000) Sound Design for Vocalizations: Quality in the Woods, Consistency
in the Fields. *The Condor* **102**, 81–92.

BÜRKNER, P.-C. (2017) **brms** : An *R* Package for Bayesian Multilevel Models Using *Stan*. *Journal of
Statistical Software* **80**, 1–28.

CATCHPOLE, C. & SLATER, P.J.B. (2008) *Bird song: biological themes and variations* 2nd edition.
Cambridge University Press, Cambridge.

CATCHPOLE, C.K. (1987) Bird song, sexual selection and female choice. *Trends in Ecology & Evolution* **2**,
94–97.

CHAMBERLAIN, S.A., HOVICK, S.M., DIBBLE, C.J., RASMUSSEN, N.L., VAN ALLEN, B.G., MAITNER, B.S., AHERN,
J.R., BELL-DERESKE, L.P., ROY, C.L., MEZA-LOPEZ, M., CARRILLO, J., SIEMANN, E., LAJEUNESSE, M.J. &
WHITNEY, K.D. (2012) Does phylogeny matter? Assessing the impact of phylogenetic
information in ecological meta-analysis: Phylogenetic meta-analysis. *Ecology Letters* **15**,
627–636.

CHAPPUIS, C. (1971) Un exemple de l'influence du milieu sur les émissions vocales des oiseaux:
l'évolution des chants en forêt équatoriale. *La terre et la vie* **2**, 183–202.

CHITNIS, S.S., RAJAN, S. & KRISHNAN, A. (2020) Sympatric wren-warblers partition acoustic signal space
and song perch height. *Behavioral Ecology* **31**, 559–567.

CHRONISTER, L.M., RHINEHART, T.A. & KITZES, J. (2023) When birds sing at the same pitch, they avoid
singing at the same time. *Ibis* **165**, 1047–1053.

CORPORATION FOR DIGITAL SCHOLARSHIP & ROY ROSENZWEIG CENTER FOR HISTORY AND NEW MEDIA (2022)
Zotero. Fairfax, Virginia, USA. www.zotero.org.

CUMMINGS, M.E. & ENDLER, J.A. (2018) 25 Years of sensory drive: the evidence and its watery bias.
Current Zoology **64**, 471–484.

DERRYBERRY, E.P., SEDDON, N., DERRYBERRY, G.E., CLARAMUNT, S., SEEHOLZER, G.F., BRUMFIELD, R.T. & TOBIAS,
J.A. (2018) Ecological drivers of song evolution in birds: Disentangling the effects of habitat
and morphology. *Ecology and Evolution* **8**, 1890–1905.

EGGER, M., SMITH, G.D., SCHNEIDER, M. & MINDER, C. (1997) Bias in meta-analysis detected by a simple,
graphical test. *BMJ* **315**, 629–634.

EMBLETON, T.F.W. (1996) Tutorial on sound propagation outdoors. *The Journal of the Acoustical
Society of America* **100**, 31–48.

ENDLER, J.A. (1992) Signals, Signal Conditions, and the Direction of Evolution. *The American Naturalist
139, S125–S153.

ESCALONA SULBARÁN, M.D., IVO SIMÕES, P., GONZALEZ-VOYER, A. & CASTROVIEJO-FISHER, S. (2019)
Neotropical frogs and mating songs: The evolution of advertisement calls in glassfrogs.
Journal of Evolutionary Biology **32**, 163–176.

EY, E. & FISCHER, J. (2009) The 'Acoustic Adaptation Hypothesis' - a review of the evidence from birds,
anurans and mammals. *Bioacoustics* **19**, 21–48.

FLETCHER, N.H. (2004) A simple frequency-scaling rule for animal communication. *The Journal of the
Acoustical Society of America* **115**, 2334–2338.

FRIIS, J.I., SABINO, J., SANTOS, P., DABELSTEEN, T. & CARDOSO, G.C. (2022) Ecological adaptation and
birdsong: how body and bill sizes affect passerine sound frequencies. *Behavioral Ecology* **33**,
798–806.

GARAMSZEGI, L.Z. & AVILÉS, J.M. (2005) Brood parasitism by brown-headed cowbirds and the
expression of sexual characters in their hosts. *Oecologia* **143**, 167–177.

GEARTY, W., JONES, L.A. & CHAMBERLAIN, S. (2023) rphylopic: an R package for accessing and plotting
PhyloPic silhouettes. https://cran.r-project.org/web/packages/rphylopic/index.html.

GIL, D. & GAHR, M. (2002) The honesty of bird song: multiple constraints for multiple traits. *Trends in
Ecology & Evolution* **17**, 133–141.

GIL, D., HONARMAND, M., PASCUAL, J., PÉREZ-MENA, E. & MACÍAS GARCIA, C. (2015) Birds living near
airports advance their dawn chorus and reduce overlap with aircraft noise. *Behavioral
Ecology* **26**, 435–443.

GOMES, L., SOLÉ, M., SOUSA-LIMA, R.S. & BAUMGARTEN, J.E. (2022) Influence of Anthropogenic Sounds on
Insect, Anuran and Bird Acoustic Signals: A Meta-Analysis. *Frontiers in Ecology and Evolution
10, 827440.

GOUTTE, S., DUBOIS, A., HOWARD, S.D., MÁRQUEZ, R., ROWLEY, J.J.L., DEHLING, J.M., GRANDCOLAS, P., XIONG,
R.C. & LEGENDRE, F. (2018) How the environment shapes animal signals: a test of the acoustic
adaptation hypothesis in frogs. *Journal of Evolutionary Biology* **31**, 148–158.

GRAFEN, A. (1989) The phylogenetic regression. *Philosophical Transactions of the Royal Society of
London. B, Biological Sciences* **326**, 119–157.

GUIBARD, A., SÈBE, F., DRAGNA, D. & OLLIVIER, S. (2022) Influence of meteorological conditions and
topography on the active space of mountain birds assessed by a wave-based sound
propagation model. *The Journal of the Acoustical Society of America* **151**, 3703–3718.

HANSEN, P. (1979) Vocal learning: its role in adapting sound structures to long-distance propagation,
and a hypothesis on its evolution. *Animal Behaviour* **27**, 1270–1271.

HARDT, B. & BENEDICT, L. (2021) Can you hear me now? A review of signal transmission and
experimental evidence for the acoustic adaptation hypothesis. *Bioacoustics* **30**, 716–742.

HART, P.J., HALL, R., RAY, W., BECK, A. & ZOOK, J. (2015) Cicadas impact bird communication in a noisy
tropical rainforest. *Behavioral Ecology* **26**, 839–842.

HAUPERT, S., SÈBE, F. & SUEUR, J. (2022) Physics-based model to predict the acoustic detection distance
of terrestrial autonomous recording units over the diel cycle and across seasons: Insights
from an Alpine and a Neotropical forest. *Methods in Ecology and Evolution* **14**, 614–630.

HIGGINS, J.P.T. & THOMPSON, S.G. (2002) Quantifying heterogeneity in a meta-analysis. *Statistics in*
Medicine **21**, 1539–1558.

HINCHLIFF, C.E., SMITH, S.A., ALLMAN, J.F., BURLEIGH, J.G., CHAUDHARY, R., COGHILL, L.M., CRANDALL, K.A.,
DENG, J., DREW, B.T., GAZIS, R., GUDE, K., HIBBETT, D.S., KATZ, L.A., LAUGHINGHOUSE, H.D., MCTAVISH,
E.J., ET AL. (2015) Synthesis of phylogeny and taxonomy into a comprehensive tree of life.
Proceedings of the National Academy of Sciences **112**, 12764–12769.

HYLTON, R. & GODARD, R.D. (2001) Song properties of indigo buntings in open and forested habitats.
The Wilson Bulletin **113**, 243–245.

IVIMEY-COOK, E.R., NOBLE, D.W.A., NAKAGAWA, S., LAJEUNESSE, M.J. & PICK, J.L. (2023) Advice for
improving the reproducibility of data extraction in meta-analysis. *Research Synthesis*
Methods **14**, 911–915.

JENNIONS, M.D. & MØLLER, A.P. (2002) Relationships fade with time: a meta-analysis of temporal
trends in publication in ecology and evolution. *Proceedings of the Royal Society of London.*
Series B: Biological Sciences **269**, 43–48.

JENSEN, K.K., LARSEN, O.N. & ATTENBOROUGH, K. (2008) Measurements and predictions of hooded crow
(Corvus corone cornix) call propagation over open field habitats. *The Journal of the*
Acoustical Society of America **123**, 507–518.

JOB, J.R., KOHLER, S.L. & GILL, S.A. (2016) Song adjustments by an open habitat bird to anthropogenic
noise, urban structure, and vegetation. *Behavioral Ecology* **27**, 1734–1744.

KIME, N.M. (2000) The transmission of advertisement calls in Central American frogs. *Behavioral*
Ecology **11**, 71–83.

KLUMP, G.M. & SHALTER, M.D. (1984) Acoustic Behaviour of Birds and Mammals in the Predator
Context; I. Factors Affecting the Structure of Alarm Signals. II. The Functional Significance
and Evolution of Alarm Signals. *Zeitschrift für Tierpsychologie* **66**, 189–226.

KUMAR, S., STECHER, G., SULESKI, M. & HEDGES, S.B. (2017) TimeTree: A Resource for Timelines,
Timetrees, and Divergence Times. *Molecular Biology and Evolution* **34**, 1812–1819.

LAGISZ, L. (2022) Around meta-analysis (14): deduplicating bibliographic records. *Inter-Disciplinary*
Ecology and Evolution Lab. http://www.i-deel.org/blog/around-meta-analysis-13-
deduplicating-bibliographic-records [accessed 17 March 2022].

LAJEUNESSE, M.J. (2009) Meta-Analysis and the Comparative Phylogenetic Method. *The American*
Naturalist **174**, 369–381.

LAJEUNESSE, M.J. (2016) Facilitating systematic reviews, data extraction and meta-analysis with the
METAGEAR package for R. *Methods in Ecology and Evolution* **7**, 323–330.

LAJEUNESSE, M.J. (2021) Automated, semi-automated, and manual extraction of numerical data from
scientific images, plot, charts, and figures. https://CRAN.R-project.org/package=juicr.

LAJEUNESSE, M.J., ROSENBERG, M.S. & JENNIONS, M.D. (2013) Recovering missing or partial data from
studies: a survey of conversions and imputations for meta-analysis. In *Handbook of Meta-*
analysis in Ecology and Evolution pp. 195–206. Princeton University Press, Princeton.

LARSEN, O.N. (2020) To Shout or to Whisper? Strategies for Encoding Public and Private Information
in Sound Signals. In *Coding Strategies in Vertebrate Acoustic Communication* (eds T. AUBIN &
N. MATHEVON), pp. 11–44. Springer International Publishing, Cham.

LONING, H., GRIFFITH, S.C. & NAGUIB, M. (2022) Zebra finch song is a very short-range signal in the wild:
evidence from an integrated approach. *Behavioral Ecology* **33**, 37–46.

LÜDECKE, D. (2019) esc: Effect Size Computation for Meta Analysis. Zenodo.
https://zenodo.org/record/1249217 [accessed 31 October 2023].

MARTEN, K. & MARLER, P.M. (1977) Sound transmission and its significance for animal vocalization. I.
Temperate habitats. *Behavioral Ecology and Sociobiology* **2**, 271–290.

MARTEN, K., QUINE, D. & MARLER, P. (1977) Sound transmission and its significance for animal
vocalization: II. Tropical forest habitats. *Behavioral Ecology and Sociobiology* **2**, 291–302.

MCLEAN, M.J., BISHOP, P.J. & NAKAGAWA, S. (2012) Male quality, signal reliability and female choice:
assessing the expectations of inter-sexual selection: Assessing the expectations of inter-
sexual selection. *Journal of Evolutionary Biology* **25**, 1513–1520.

MENDOZA-HENAO, A.M., ZAMUDIO, K.R., GUAYASAMIN, J.M., ESCALONA, M. & PARRA-OLEA, G. (2023)
Environment rather than character displacement explains call evolution in glassfrogs.
Evolution **77**, 355–369.

MICHONNEAU, F., BROWN, J.W. & WINTER, D.J. (2016) rotl: an R package to interact with the Open Tree
of Life data. *Methods in Ecology and Evolution* **7**, 1476–1481.

MIKULA, P., VALCU, M., BRUMM, H., BULLA, M., FORSTMEIER, W., PETRUSKOVÁ, T., KEMPENAERS, B. &
ALBRECHT, T. (2020) A global analysis of song frequency in passerines provides no support for
the acoustic adaptation hypothesis but suggests a role for sexual selection. *Ecology Letters*
24, 477–486.

MOHER, D., LIBERATI, A., TETZLAFF, J. & ALTMAN, D.G. (2009) Preferred Reporting Items for Systematic
Reviews and Meta-Analyses: The PRISMA Statement. *Journal of Clinical Epidemiology* **62**,
1006–1012.

MORTON, E.S. (1975) Ecological sources of selection on avian sounds. *The American Naturalist* **109**,
17–34.

NAGUIB, M. (2003) Reverberation of rapid and slow trills: Implications for signal adaptations to long-
range communication. *The Journal of the Acoustical Society of America* **113**, 1749–1756.

NAKAGAWA, S., LAGISZ, M., JENNIONS, M.D., KORICHEVA, J., NOBLE, D.W.A., PARKER, T.H., SÁNCHEZ-TÓJAR, A.,
YANG, Y. & O'DEA, R.E. (2022) Methods for testing publication bias in ecological and
evolutionary meta-analyses. *Methods in Ecology and Evolution* **13**, 4–21.

NAKAGAWA, S., LAGISZ, M., O'DEA, R.E., POTTIER, P., RUTKOWSKA, J., SENIOR, A.M., YANG, Y. & NOBLE, D.W.A.
(2023) orchaRd 2.0: An R package for visualising meta-analyses with orchard plots. *Methods*
in Ecology and Evolution **14**, 2003–2010.

NAKAGAWA, S., NOBLE, D.W.A., SENIOR, A.M. & LAGISZ, M. (2017) Meta-evaluation of meta-analysis: ten
appraisal questions for biologists. *BMC Biology* **15**, 18.

NAKAGAWA, S. & SANTOS, E.S.A. (2012) Methodological issues and advances in biological meta-analysis.
Evolutionary Ecology **26**, 1253–1274.

NAKAGAWA, S. & SCHIELZETH, H. (2013) A general and simple method for obtaining R^2 from generalized
linear mixed-effects models. *Methods in Ecology and Evolution* **4**, 133–142.

NEMETH, E., DABELSTEEN, T., PEDERSEN, S.B. & WINKLER, H. (2006) Rainforests as concert halls for birds:
Are reverberations improving sound transmission of long song elements? *The Journal of the*
Acoustical Society of America **119**, 620–626.

NICHOLLS, J.A. & GOLDIZEN, A.W. (2006) Habitat type and density influence vocal signal design in satin
bowerbirds: Habitat-related call variation in bowerbirds. *Journal of Animal Ecology* **75**, 549–
558.

O'DEA, R.E., LAGISZ, M., JENNIONS, M.D., KORICHEVA, J., NOBLE, D.W.A., PARKER, T.H., GUREVITCH, J., PAGE,
M.J., STEWART, G., MOHER, D. & NAKAGAWA, S. (2021) Preferred reporting items for systematic
reviews and meta-analyses in ecology and evolutionary biology: a PRISMA extension.
Biological Reviews **96**, 1695–1722.

OPHIR, A.G., SCHRADER, S.B. & GILLOOLY, J.F. (2010) Energetic cost of calling: general constraints and
species-specific differences. *Journal of Evolutionary Biology* **23**, 1564–1569.

PARADIS, E., CLAUDE, J. & STRIMMER, K. (2004) APE: Analyses of Phylogenetics and Evolution in R
language. *Bioinformatics* **20**, 289–290.

PEARSE, W.D., MORALES-CASTILLA, I., JAMES, L.S., FARRELL, M., BOIVIN, F. & DAVIES, T.J. (2018) Global
macroevolution and macroecology of passerine song. *Evolution* **72**, 944–960.

R CORE TEAM (2022) R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. http://www.R-project.org/.

RIVERA, M., EDWARDS, J.A., HAUBER, M.E. & WOOLLEY, S.M.N. (2023) Machine learning and statistical
classification of birdsong link vocal acoustic features with phylogeny. *Scientific Reports* **13**,
7076.

ROTHSTEIN, S.I. & FLEISCHER, R.C. (1987) Vocal Dialects and Their Possible Relation to Honest Status
867   Signalling in the Brown-Headed Cowbird. *The Condor* **89**, 1–23.

868 RSTUDIO TEAM (2021) RStudio: Integrated Development Environment for R. RStudio, PBC, Boston, MA.
869   http://www.rstudio.com/.

870 RYAN, M.J., TUTTLE, M.D. & TAFT, L.K. (1981) The costs and benefits of frog chorusing behavior.
871   *Behavioral Ecology and Sociobiology* **8**, 273–278.

872 SEDDON, N. (2005) Ecological adaptation and species recognition drives vocal evolution in Neotropical
873   suboscine birds. *Evolution* **59**, 200–215.

874 SENIOR, A.M., GRUEBER, C.E., KAMIYA, T., LAGISZ, M., O'DWYER, K., SANTOS, E.S.A. & NAKAGAWA, S. (2016)
875   Heterogeneity in ecological and evolutionary meta-analyses: its magnitude and implications.
876   *Ecology* **97**, 3293–3299.

877 SLABBEKOORN, H. & PEET, M. (2003) Birds sing at a higher pitch in urban noise. *Nature* **424**, 267–267.

878 SLABBEKOORN, H. & SMITH, T.B. (2002) Habitat-dependent song divergence in the little greenbul: an
879   analysis of environmental selection pressures on acoustic signals. *Evolution* **56**, 1849–1858.

880 STANIEWICZ, A., SOKOŁOWSKA, E., MUSZYŃSKA, A. & BUDKA, M. (2023) Competition for acoustic space in a
881   temperate-forest bird community. *Behavioral Ecology* **34**, 1043–1054.

882 STERNE, J.A.C., SUTTON, A.J., IOANNIDIS, J.P.A., TERRIN, N., JONES, D.R., LAU, J., CARPENTER, J., RUCKER, G.,
883   HARBORD, R.M., SCHMID, C.H., TETZLAFF, J., DEEKS, J.J., PETERS, J., MACASKILL, P., SCHWARZER, G., ET
884   AL. (2011) Recommendations for examining and interpreting funnel plot asymmetry in meta-
885   analyses of randomised controlled trials. *BMJ* **343**, d4002.

886 SUEUR, J., KRAUSE, B. & FARINA, A. (2019) Climate Change Is Breaking Earth's Beat. *Trends in Ecology &*
887   *Evolution* **34**, 971–973.

888 TONINI, J.F.R., PROVETE, D.B., MACIEL, N.M., MORAIS, A.R., GOUTTE, S., TOLEDO, L.F. & PYRON, R.A. (2020)
889   Allometric escape from acoustic constraints is rare for frog calls. *Ecology and Evolution* **10**,
890   3686–3695.

891 TYACK, P.L. (2020) A taxonomy for vocal learning. *Philosophical Transactions of the Royal Society B:*
892   *Biological Sciences* **375**, 20180406.

893 VIECHTBAUER, W. (2010) Conducting Meta-Analyses in *R* with the **metafor** Package. *Journal of*
894   *Statistical Software* **36**, 1–48.

895 VIECHTBAUER, W. (2023) Confidence intervals for the amount of heterogeneity accounted for in meta-
896   regression models. Ghent, Belgium.
897   https://www.wvbauer.com/lib/exe/fetch.php/talks:2023_viechtbauer_eam_ci_for_r2.pdf.

898 DE VILLEMEREUIL, P. & NAKAGAWA, S. (2014) General Quantitative Genetic Methods for Comparative
899   Biology. In *Modern Phylogenetic Comparative Methods and Their Application in Evolutionary*
900   *Biology* (ed L.Z. GARAMSZEGI), pp. 287–303. Springer Berlin Heidelberg, Berlin, Heidelberg.

901 WELKLIN, J.F., BRANCH, C.L., PITERA, A.M., SONNENBERG, B.R., BENEDICT, L.M., HEINEN, V.K., KOZLOVSKY, D.Y.
902   & PRAVOSUDOV, V.V. (2023) Females pair with males larger than themselves in a socially
903   monogamous songbird. *Animal Behaviour* **199**, 35–49.

WICKHAM, H. (2016) *ggplot2: elegant graphics for data analysis* Second edition. Springer, Cham.

WICKHAM, H., FRANÇOIS, R., HENRY, L. & MÜLLER, K. (2022) dplyr: A Grammar of Data Manipulation.
https://CRAN.R-project.org/package=dplyr.

WILEY, R.H. & RICHARDS, D.G. (1978) Physical constraints on acoustic communication in the
atmosphere: Implications for the evolution of animal vocalizations. *Behavioral Ecology and*
Sociobiology **3**, 69–94.

WILEY, R.H. & RICHARDS, D.G. (1982) Adaptations for Acoustic Communication in Birds: Sound
Transmission and Signal Detection. In *Acoustic Communication in Birds* pp. 131–181.
Elsevier.

YU, G., SMITH, D.K., ZHU, H., GUAN, Y. & LAM, T.T. (2017) GGTREE : an R package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods in*
Ecology and Evolution **8**, 28–36.

ZHAO, L., SANTOS, J.C., WANG, J., RAN, J., TANG, Y. & CUI, J. (2021) Noise constrains the evolution of call
frequency contours in flowing water frogs: a comparative analysis in two clades. *Frontiers in*
Zoology **18**, 37.

Fig. 1. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) diagram of

the protocol for screening, including and excluding studies for this meta-analysis and number of

studies identified at each stage. OATD, Open Access Theses and Dissertations.

Fig. 2. Caterpillar plot for (A) within- and (B) among-species comparisons. Effect sizes are displayed

by taxonomic class, ordered by mean estimate, and with horizontal bars showing 95% confidence

intervals (CIs). Symbols of the point estimate illustrate the acoustic variable measured. Positive

Pearson correlation coefficients (*r*) indicate support for the acoustic adaptation hypothesis. Solid

vertical black lines indicate the estimated global effect size obtained from the REML model with

random factors only, and the grey area denotes the 95% CIs (A: *r* = -0.047, 95% CIs: -0.193 to 0.099;

B: *r* = 0.105, 95% CIs: -0.070 to 0.280).

Fig. 3. Orchard plots for (A) within- and (B) among-species comparisons showing effect sizes
(Pearson correlation coefficients, *r*) and number of taxa (k = total number of effect sizes (total
number of taxa) and mean effect size from REML models. Solid horizontal bars are 95% confidence
intervals (CIs) of global effect sizes obtained for each taxon. Solid vertical black lines indicate the
estimated global effect size from the REML model with the random factors only, and grey bands
denote the 95% CIs (A: *r* = -0.047, 95% CIs: -0.193 to 0.099; B: *r* = 0.105, 95% CIs: -0.070 to 0.280).
Circle size reflects effect size precision (log(1/SE)).

941 **Table 1**. Global effect sizes for the acoustic adaptation hypothesis obtained from the restricted maximum likelihood approach. Results of intercept-only

942 General Linear-Mixed Effects Models are shown for the entire dataset (global model) and for subsets with respect to animal class (Aves, Mammalia or

943 Amphibia). For among-species comparisons of Aves, the results are shown for a subset of studies that accounted for phylogenetic non-independence (Aves

944 (phylogeny correction)). Table shows number of effect sizes (k), number of species or taxa (N), estimates of r and heterogeneity I^2 (total, variance arising

945 from phylogenetic affinities, taxon, study identity, and observation).

Type of study	Taxonomic range	k	N	r		z-value	P-value	I^2 Total	I^2 Phylogeny	I^2 Taxon	I^2 Study	I^2 Observation
Within species	Global model	219	37	-0.047	(-0.193,0.099)	-0.628	0.530	0.954	0.129	0.000	0.298	0.528
	Aves (all)	189	32	0.062	(0.021,0.103)	2.965	**0.003**	0.936	0.000	0.000	0.000	0.936
	Mammalia	16	3	-0.205	(-0.847,0.437)	-0.627	0.530	0.979	0.000	0.897	0.000	0.082
	Amphibia	14	2	-0.443	(-1.087,0.202)	-1.346	0.178	0.904	0.284	0.284	0.284	0.050
Among species	Global model	152	20	0.105	(-0.070,0.280)	1.172	0.241	0.950	0.141	0.049	0.572	0.188
	Aves (all)	129	14	0.140	(0.006,0.274)	2.049	**0.040**	0.943	0.010	0.054	0.673	0.206
	Aves (phylogeny correction)	67	11	0.029	(-0.079,0.137)	0.523	0.601	0.841	0.000	0.425	0.269	0.148
	Mammalia	11	3	-0.257	(-0.605,0.091)	-1.448	0.148	0.966	0.722	0.000	0.223	0.021
	Amphibia	12	3	0.229	(-0.119,0.577)	1.292	0.196	0.864	0.554	0.000	0.000	0.310