Meta-analysis of the acoustic adaptation hypothesis reveals no support for the effect of

vegetation structure on acoustic signalling across terrestrial vertebrates

Bárbara Freitas [1,2,3]*, Pietro B. D'Amelio [4], Borja Milá [1], Christophe Thébaud [2], Tim Janicke [5]

[1] National Museum of Natural Sciences, Spanish National Research Council (CSIC), Madrid, Spain

[2] Centre de Recherche sur la Biodiversité et l'Environnement (UMR 5300 CNRS-IRD-TINPT-UPS),

Université Paul Sabatier, 31062 Toulouse Cedex 9, France

[3] Facultad de Ciencias, Universidad Autónoma de Madrid, 28049 Madrid, Spain

[4] Department of Behavioral Neurobiology, Max Planck Institute for Biological Intelligence, Seewiesen

82319, Germany

[5] CEFE, Univ Montpellier, CNRS, EPHE, IRD, Montpellier, France

*Correspondence to:

Bárbara Freitas

Museo Nacional de Ciencias Naturales

Calle José Gutiérrez Abascal 2

Madrid 28006, Spain

E-mail: barbara.freitas@mncn.csic.es

Tel.: +351 965259456

Author ORCIDs

Bárbara Freitas http://orcid.org/0000-0001-6020-6236

Pietro B. D'Amelio http://orcid.org/0000-0002-4095-6088

Borja Milá http://orcid.org/0000-0002-6446-0079

Christophe Thébaud http://orcid.org/0000-0002-8586-1234

Tim Janicke http://orcid. org/0000-0002-1453-6813

ABSTRACT

Acoustic communication plays a prominent role in various ecological and evolutionary processes

involving social interactions. The properties of acoustic signals are thought to be influenced not only

by the interaction between signaller and receiver but also by the acoustic characteristics of the

environment through which the signal is transmitted. This conjecture forms the core of the so-called

"acoustic adaptation hypothesis" (AAH), which posits that vegetation structure affects frequency

and temporal parameters of acoustic signals emitted by a signaller as a function of their acoustic

degradation properties. Specifically, animals in densely vegetated 'closed habitats' are expected to

produce longer acoustic signals with lower repetition rates and lower frequencies (minimum, mean,

maximum, and peak) compared to the ones inhabiting less vegetated 'open habitats'. To date, this

hypothesis has received mixed results, with the level of support depending on the taxonomic group

and the methodology used. We conducted a systematic literature search of empirical studies testing

for an effect of vegetation structure on acoustic signalling and assessed the generality of the AAH

using a meta-analytic approach based on 371 effect sizes from 75 studies and 57 taxa encompassing

birds, mammals and amphibians. Overall, our results do not provide consistent support for the AAH,

neither in within-species comparisons (suggesting no overall phenotypically plastic response of

acoustic signalling to vegetation structure) nor in among-species comparisons (suggesting no overall

evolutionary response). However, when considering birds only, we found a weak support for the

AAH in within-species comparisons, which was mainly driven by studies that measured frequency

bandwidth, suggesting that this variable may exhibit a phenotypically plastic response to vegetation

structure. For among-species comparisons in birds, we also found support for the AAH, but this

effect was not significant after excluding comparative studies that did not account for phylogenetic

non-independence. Collectively, our synthesis does not support a universal role of vegetation

structure in the evolution of acoustic communication. We highlight the need for more empirical

work on currently under-studied taxa such as amphibians, mammals, and insects. Furthermore, we

propose a framework for future research on the AAH. We specifically advocate for a more detailed

and quantitative characterization of habitats to identify frequencies with the highest detection

probability and to determine if frequencies with greater detection distances are preferentially used.

Finally, we stress that empirical tests of the AAH should focus on signals which are selected for

increased transmission distance.

Key words: acoustic communication, amphibians, bioacoustics, birds, evolution, habitat, mammals,

vegetation, vocalizations

CONTENTS

I. INTRODUCTION

(1) Acoustic signalling and the effect of multiple selective forces

Acoustic signalling is an essential facet of animal communication, spanning a wide array of species and ecosystems. These signals play a pivotal role in various contexts, including mate attraction, territory defence, predation avoidance (alerting others of a threat), rival intimidation, individual recognition, and social communication (Catchpole & Slater, 2008; Bradbury & Vehrencamp, 2011). Thus, acoustic signals are subject to multiple selective forces imposed on the signaller and receiver that shape them to finely tuned instruments of communication (Wiley & Richards, 1982; Catchpole, 1987; Catchpole & Slater, 2008). These multiple selective forces may act in concert or in opposition, which potentially complicates our understanding of the evolution of acoustic communication (i.e., the adaptive significance of the specific structure of a given signal).

One of the most studied selective forces shaping acoustic signals is the morphology of the signaller. For instance, the size of sound-producing organs is often positively associated with a signaller's body size or mass and has been shown to influence signal frequency in anurans and mammals (Fletcher, 2004; Tonini *et al.*, 2020). Similarly, in birds body size has been found to be negatively correlated with signal minimum frequency, and positively with frequency bandwidth (Friis *et al.*, 2022) or peak frequency (Mikula *et al.*, 2021). Thus, when individuals emit sounds, they also convey information about their size such that acoustic cues can serve as honest signals of body condition and reproductive performance, influencing mate choice (McLean, Bishop & Nakagawa, 2012; Welklin *et al.*, 2023). Although producing acoustic signals at high amplitude may attract more mates and provide competitive advantages, it incurs energetic costs (e.g., Gil & Gahr, 2002; Ophir, Schrader & Gillooly, 2010), which can make it challenging to produce an optimally designed sound for long-distance transmission. In addition, predators and parasites can use acoustic signals emitted by their prey as cues to locate them (Ryan, Tuttle & Taft, 1981; Garamszegi & Avilés, 2005), potentially driving the development of signals that minimize their detectability, such as high-frequency pure tones in certain alarm calls to reduce locatability (Klump & Shalter, 1984). In well-

documented vocal learners such as oscine passerines, hummingbirds, parrots, humans, elephants,

seals, and toothed whales, vocalisations are also shaped by the capability of reproducing heard

sounds (Tyack, 2020). This ability might counterbalance other predicted trends (mentioned above)

that shape acoustic signals. For example, learning different signals from immigrants or learning with

errors can introduce variations that offset these trends. Likewise, interactions with other animals

can shape acoustic signals in multiple ways, for example by selecting for frequencies that exploit

sensory biases (Cummings & Endler, 2018).

The structure of acoustic signals can also be influenced by the acoustic characteristics of the

environment into which they are emitted and received (Chappuis, 1971; Slabbekoorn & Smith, 2002;

Derryberry *et al.*, 2018). While propagating from signaller to receiver, signals are subject to several

alterations in temporal and frequency parameters. These alterations include reverberation, i.e., the

delay between the faster waves (direct) and slowest waves (indirect) causing distortion of the sound

(Morton, 1975; Wiley & Richards, 1978, 1982), and attenuation, i.e., the progressive reduction in

amplitude which can be attributed to three main components: (1) geometric attenuation (also

known as spreading loss or spherical attenuation), (2) atmospheric attenuation, and (3) habitat

attenuation (Embleton, 1996). The latter two components are often collectively referred to as

degradation, resulting from scattering, refraction through obstacles, atmospheric absorption, or

echoes, with greater effects typically observed for high frequencies (Aylor, 1972; Marten & Marler,

1977). In this context, natural selection is expected to have favoured the evolution of accurate and

efficient transmission of acoustic signals for long-distance travel, enabling information transfer from

signaller to receiver while accounting for environmental constraints.

(2) The Acoustic Adaptation Hypothesis

The various selective forces imposed by the environment outlined above form the

theoretical background of the acoustic adaptation hypothesis (AAH), mainly grounded in the work of

Morton (1975) and Hansen (1979) but coined by Rothstein & Fleischer (1987). The AAH argues that

the transmission of acoustic signals is affected by the environment and offers explicit predictions

regarding the optimal shape of acoustic signals to ensure accurate transmission in environments

characterized by high degradation, such as densely vegetated habitats (Morton, 1975; Blumenrath &

Dabelsteen, 2004). Animals inhabiting such hereafter called 'closed habitats' (as opposed to less

vegetated 'open habitats') are expected to have signals of longer duration because this is expected

to increase the likelihood of detection and to leverage the amplitude of reverberated sound waves,

extending the range of signal propagation (Nemeth *et al.*, 2006). To increase transmission distance in

closed habitats, acoustic signals should also have lower repetition rates to minimize interference

with reverberated waves (Brown & Handford, 2000; Naguib, 2003) and narrower frequency

bandwidths to concentrate energy at frequencies less susceptible to attenuation (Wiley & Richards,

1978). Additionally, animals in closed habitats should be prone to emit acoustic signals at lower

frequencies including minimum, mean, maximum, and peak frequency (Chappuis, 1971; Morton,

1975; Marten, Quine & Marler, 1977; see Table S1 for the definition and predictions for each

acoustic variable).

The AAH is currently considered as a component of the broader Sensory Drive Hypothesis,

which predicts that signals, sensory mechanisms, and microhabitat choice evolve together based on

the characteristics of the habitat, ambient noise patterns, presence of predators and parasites, and

various sensory and physiological factors (Endler, 1992; Cummings & Endler, 2018). Several authors

have also extended the AAH to a wider framework of environmental adaptation, covering studies on

the effect of environmental noise such as tests of the Lombard effect in the context of urbanization

(Job, Kohler & Gill, 2016; Zhao *et al.*, 2021). Here, we focus on the AAH in the context of the effect of

vegetation structure on acoustic signalling as originally tested by Morton (1975).

Playback transmission experiments using either natural or synthetic sounds have been

conducted to assess the physical foundations of the AAH (Wiley & Richards, 1978). In addition,

comparisons between species and populations inhabiting environments with different vegetation

structure have been performed to test the validity of this hypothesis. While studies testing

environmental filtering of sound have provided a robust foundation for the AAH (reviewed in Hardt

& Benedict, 2021), its influence on animal acoustic signal production remains unclear. In fact,

empirical support for the AAH is highly variable, with many studies supporting the hypothesis

(Bertelli & Tubaro, 2002; Nicholls & Goldizen, 2006) whereas many others contest its validity (Hylton

& Godard, 2001; Bosch & De la Riva, 2004; Mikula *et al.*, 2021). In addition, Hardt & Benedict (2021)

uncovered that animal signals are often not better transmitted in a species' habitat compared to

environments they are not adapted to, raising further questions about the generality of the AAH.

In a previous meta-analysis limited to birds, Boncoraglio & Saino (2007) compiled 91 effect

sizes from 26 studies of within- (including 12 species) and among-species (spanning 3 orders)

comparisons to test for an overall effect of habitat structure (open versus closed) on acoustic

variables. The authors found support for the AAH in terms of a lower minimum frequency, maximum

frequency, peak frequency, and frequency bandwidth in closed habitats based on a global effect size

of Fisher's z = 0.160, 0.182, 0.346, and 0.230, respectively (Boncoraglio & Saino, 2007). Interval

duration was the only variable that did not differ between habitats. Since its publication, this study

has served as a key reference in support of the AAH in birds. However, Boncoraglio & Saino (2007)

relied on a limited sample size and did not include measurements of mean frequency, total duration,

or repetition rate in their analyses, despite clear predictions for these three variables within the

framework of the AAH. Furthermore, this study did not explicitly control for potential sources of

statistical non-independence beyond shared study identity, such as shared phylogenetic history. The

lack of statistical control for non-independence among effect sizes can result in inaccurate

conclusions, often leading to spurious statistical significance (Nakagawa *et al.*, 2017), and integrating

phylogenetic structure into meta-analyses is now considered critical to obtain unbiased results

(Lajeunesse, 2009; Chamberlain *et al.*, 2012).

Although the AAH was initially formulated based on avian studies, there has been a growing

interest in testing this hypothesis in other taxonomic groups. Aiming to examine how widespread

the impact of vegetation structure on the evolution of acoustic signals is in vertebrates, Ey & Fischer

(2009) qualitatively reviewed 55 studies on birds, anurans and mammals and concluded that the

AAH did not seem to be universally valid across taxa. However, the authors claimed that in birds,

effects on minimum frequency and frequency bandwidth appeared to align with the predictions of

the AAH, whereas relationships with repetition rate, mean, peak, and maximum frequencies did not,

which does not fully correspond to the results obtained by Boncoraglio & Saino (2007).

(3) Assessing the AAH across animals

Despite the large body of research aimed at testing the AAH, it still remains unclear to what extent

this hypothesis is valid across acoustic variables in a broad range of animal taxa, and whether

contrasting findings from previous studies reflect methodological artifacts and/or actual differences

among taxa. Our study aims to fill this gap by conducting a meta-analysis of empirical studies that

tested the AAH in the context of airborne acoustic signalling in response to vegetation structure. In

this regard, we build on the work by Boncoraglio & Saino (2007) with the aim to provide a robust

and comprehensive meta-analysis. More specifically, our main objective is to address two main

questions: (1) Does the AAH hold across different animal taxa and acoustic parameters in the

context of vegetation structure? and (2) does the AAH support phenotypically plastic and/or

evolutionary responses to vegetation structure (inferred from within- and among-species

comparisons, respectively)? For these purposes, we compiled 371 effect sizes testing the AAH in

animals. We did not have a clear prediction on whether effect sizes differ between within- and

among-species comparisons. However, it seems plausible that effect sizes from among-species

comparisons might be weaker because acoustic parameters likely evolve not only in relation to the

habitat but also due to other factors such as interspecific variation in mate preferences and/or

neutral processes such as genetic drift.

II. METHODS

This meta-analysis was pre-registered at the Open Science Framework (OSF; study details available

at: https://doi.org/10.17605/OSF.IO/E7JRS). Differences between the pre-registered study design

and the eventually implemented approach are mainly due to methodological improvements and are

explained in Text S1.

(1) Literature search

We conducted a systematic literature search to obtain an unbiased sample of studies testing for a

relationship between vegetation structure and acoustic communication in animals following the

Preferred Reporting Items for Systematic Reviews and Meta-Analyses approach (PRISMA; Moher *et*

al., 2009) in combination with the PRISMA-EcoEvo guidelines (O'Dea *et al.*, 2021). We screened the

Scopus, the ISI Web of Science Core Collection (Clarivate Analytics), and the Open Access Theses and

Dissertations (OATD; https://oatd.org/) databases on 16th December 2021 with the search terms

defined as ((("*song" OR "vocal*") OR (("acoustic$" OR "bioacoustic$") AND "call$")) AND

("habitat$" OR "vegetation")) OR "acoustic adaptation". These searches yielded 3300, 3233, and 372

records, respectively. We also considered studies included in previous reviews on related questions

(Ey & Fischer, 2009; Hardt & Benedict, 2021) and a previous meta-analysis on birds (Boncoraglio &

Saino, 2007), resulting in the addition of 58 studies to be screened. Furthermore, we posted a

request for unpublished data on the 'evoldir' mailing list (http://life.mcmaster.ca/evoldir.html), the

ResearchGate platform (https://www.researchgate.net/), and Twitter, and directly asked colleagues

who work on these topics. This resulted in one additional record (Melo *et al*., in prep). Finally, we

added one unpublished work on the La Palma population of the Canary Islands Chaffinch (*Fringilla*

canariensis palmae; Freitas *et al*., in prep.).

We excluded 45 studies from the OATD search because full texts were not accessible. We

also excluded duplicates by following an R workflow adapted from Lagisz (2022) and by manually

checking the results of the Zotero deduplication algorithm (Corporation for Digital Scholarship & Roy

Rosenzweig Center for History and New Media, 2022). Then, we assigned screening efforts among

BF, BM, and TJ, who collectively screened the titles and abstracts of 4558 studies. To ensure

consistency, 212 of these studies were screened in parallel by two authors. We selected potentially

relevant primary studies using the *metagear* package (Lajeunesse, 2016) based on the following

selection criteria:

1. The focus of the empirical study was on non-human animals, for which acoustic signals are transmitted through the air. We did not consider studies focusing on domesticated species as artificial selection can obscure the selection mechanisms that shape communication signals in the wild. We also did not consider studies focusing on echolocation because it is used for orientation in space and/or to locate items rather than for communication and this could confound our results. For syntheses on the effect of vegetation structure on echolocation, see Ey & Fischer (2009). We also excluded studies on aquatic habitats due to the different acoustic properties and difficulties in defining vegetation structure based on underwater vegetation.

2. At least one temporal or frequency parameter for which the AAH predicts an effect was measured for adult individuals of the same species (within-species comparisons) or multiple species (among-species comparisons) across different vegetation types. Studies testing for differences among subspecies were considered as among-species comparisons. Studies that did not replicate habitat type within species or subspecies were excluded (as a conservative approach to mitigate the possibility of testing subspecies differences rather than differences due to habitat). For example, a study was excluded if testing for acoustic signal differences between a subspecies inhabiting one location of habitat type A and another subspecies in another location of habitat type B.

3. Habitat was described based on a continuous variable or on a categorical classification of vegetation structure (from open to closed habitats). If the categorical classification of the habitat was done in more than the two categories, we pooled different habitat types into two categories: 'open' and 'closed' (e.g., grassland, open field, and meadow grouped in 'open habitat'). We did not include studies examining differences between urban and rural habitats.

4. The study was written in English, French, German, Portuguese, or Spanish according to the

languages that the authors were able to understand.

5. The study reported sufficient information to compute effect sizes (e.g., sample size, test-

statistic, means with estimates of uncertainty).

During the title and abstract screening stage, we excluded 4237 studies thatdid not fit at

least one of the requirements listed above. Afterwards, we carried out a full-text screening of the

remaining 321 studies. During this stage we excluded 246 studies because they: (i) did not test the

influence of vegetation structure on acoustic variables (N = 116), (ii) used habitat types that could

not be classified as 'open' or 'closed' (N = 17), (iii) did not replicate habitat type (N = 5), (iv) focused

on echolocation (N = 23), (v) focused on playback/transmission experiments (N = 15), (vi) focused on

the effect on urbanization (N = 4), (vii) used acoustic variables for which the AAH does not generate

predictions (N = 3), (viii) did not report sufficient data (N = 27; i.e., multivariate statistics not

providing raw data, missing key data required to compute an effect size even after contacting the

authors), (ix) represented reviews or non-empirical studies (N = 14), (x) were inaccessible (N = 12),

(xi) were written in other languages (N = 4), or (xii) represented duplicates (N = 6; of which 5

represented theses that are already covered by other primary studies). The final dataset included 75

primary studies (see Fig. 1 for the PRISMA diagram and Table S2 for a reference list of all primary

studies).

(2) Data collection

BF and TJ extracted means, metrics of uncertainty (standard deviation or standard error) and sample

sizes, correlation coefficients and sample sizes, or the statistical test value and degrees of freedom

reported for the relationship among acoustic variables and vegetation structure from the text,

tables, figures or supplementary materials. We used the R package *juicr* version 0.1 (Lajeunesse,

2021) to extract summary statistics from figures. From five studies that used multivariate statistics

(such as principal components analysis), we reanalysed the raw data if published together with the

article or provided by the authors (*N* = 2), and corrected also for the phylogeny in cases in which

phylogenetic data was provided (*N* = 1). Extractions from 27 studies were cross-checked by both

authors to ensure accuracy.

All comparisons included in our meta-analysis followed a descriptive approach, meaning that

none of the primary studies manipulated vegetation structure experimentally to test for a causal

effect on acoustic variables. Thus, we used the Pearson correlation coefficient (*r*) as an effect size to

synthesize the findings of primary studies. We converted test statistics provided in the primary

studies into *r* and estimated its sampling variance (*Vr*) using formulas and guidelines reported in

Lajeunesse *et al*. (2013) and using the R package *esc* version 0.5.1 (Lüdecke, 2019). To improve

transparency and allow for reproducibility of data extraction, we report the method used to

compute effect sizes (Table S2) following a framework provided in Ivimey-Cook *et al*. (2023). For

each acoustic variable, correlations that aligned with the predictions of the AAH (Table S1) were

assigned a positive sign. For studies using a categorical classification of the vegetation structure (see

section 'Moderator variables') of three groups (open, closed, and mixed; *N* = 3) and where effect

sizes were computed from group means, we only used the values from the two extreme categories:

open and closed. All effect sizes were assigned to a type of study: within- or among-species

comparisons, to allow testing for a phenotypically plastic or an evolutionary response of acoustic

signalling to vegetation structure, respectively. Noting that most of the studies on within-species

comparisons focused on nearby populations, we interpreted the support for the AAH in within-

species comparisons as suggestive of a phenotypically plastic response rather than adaptive

response since we assumed no genetic divergence between the different populations tested within

any given species.

(3) Moderator variables

Besides the global test for the support of the AAH across animals, we aimed at explaining

heterogeneity in effect sizes among studies from a biological and methodological perspective.

Specifically, we explored if support for the AAH differed among animal taxa and acoustic variables. If

an acoustic variable was measured for parts and for the full signal duration, we extracted both effect

sizes separately but assigned them as the same acoustic variable (e.g., song duration and note

duration were both considered as duration).

In addition, we tested for four methodological moderators: habitat classification, habitat

contrast, body size correction, and phylogeny correction. Firstly, we explored effects of the habitat

classification (continuous versus categorical) used in primary studies, i.e., if the studies used a

continuous variable or a categorical classification to describe the vegetation structure of the habitat.

When authors of primary studies ran analyses with both types of classification (N = 2), we retrieved

only the estimates obtained for the continuous variable, as we consider it less subjective than a

categorical classification. Secondly, we scored the magnitude of contrast between closed and open

habitats (for studies using a categorical classification) that were used in the study (habitat contrast),

attributing a '0' when there was a moderate difference in vegetation structure (e.g., temperate

broadleaf forest versus tropical rainforest), or a '1' when there was a substantial difference in

vegetation structure (e.g., grassland versus forest). Importantly, primary studies investigated various

habitat types and used different methods for classifying them. This diversity in methods to classify

vegetation structure could have contributed to differences in study outcomes (suggested by Hardt &

Benedict, 2021). By considering these two moderators, we tried to account for this potential source

of heterogeneity among studies. Thirdly, we recorded whether primary studies accounted for body

size differences within and/or among species, since this trait is known to be allometrically correlated

with the morphology of the sound apparatus, and therefore to affect the production of sound

(Pearse *et al.*, 2018). Finally, we recorded whether studies reporting among-species comparisons

accounted for phylogenetic non-independence.

(4) Phylogeny

In order to account for phylogenetic non-independence and to quantify the phylogenetic signal (also termed phylogenetic heritability; de Villemereuil & Nakagawa, 2014), we reconstructed a phylogeny of all sampled taxa from studies of within- (Fig. S1) and among-species comparisons (Fig. S2). We used the R package *rotl* (Michonneau, Brown & Winter, 2016) to retrieve the phylogenetic relationships from the Open Tree of Life (opentreeoflife.org/; Hinchliff *et al.*, 2015) and *ape* (Paradis, Claude & Strimmer, 2004) to obtain the branch lengths of the trees by applying Grafen's method (Grafen, 1989).

In analyses of among-species comparisons, the phylogenetic position of some taxonomic groups could not be assessed unequivocally so that we assigned the group in question to a higher taxonomic level. Specifically, we considered Parulinae as Parulidae and Cardinalidae as Passeriformes. This decision accounts for the fact that a number of genera traditionally placed within Cardinalidae are embedded within other families, and conversely, some birds traditionally placed in other families belong with the cardinals (Winkler, Billerman & Lovette, 2020). Note that some primary studies tested the AAH across higher taxonomic ranks (such as Anura, Aves and Passeriformes), while others tested it across a lower taxonomic rank, which sometimes was nested within the higher ones (artificial pairs, e.g., Anura and Hylidae, Aves and *Corvus*). In such cases, we retrieved pairwise estimates of divergence times for the taxonomic ranks that were not nested from the TimeTree database (http://timetree.org/; Kumar *et al.*, 2017). Then, for the artificial pairs, we attributed a mean estimate between the known divergence time estimates of the nested pairs and the ones more distantly related. For example, for the pair Anura - Hylidae, we computed the divergence time as the mean of the difference between the estimate of Hylidae - *Rhinella* (a genus within Anura) and the estimate of Anura - Felidae, since Felidae is the closest outgroup of Anura in our phylogeny. Then, we computed branch lengths using the methods already specified for the within-species comparison.

(5) Data analyses

All analyses were performed in R 4.2.2 (R Core Team, 2022) through RStudio 1.4.1106 (RStudio

Team, 2021), using the packages *metafor* (Viechtbauer, 2010) for the meta-regression models, *brms*

(Bürkner, 2017) for models with Bayesian approach, *orchaRd* (Nakagawa *et al.*, 2023a) for calculation

of heterogeneity of Restricted Maximum Likelihood (REML) models, *outliers* (Komsta, 2005) for

outlier analyses, *dplyr* (Wickham *et al.*, 2022) for data manipulation, *ggplot2* (Wickham, 2016) for

data visualization, and *ggtree* (Yu *et al.*, 2017) to produce our final phylogenies as shown in Figs S1

and S2. Silhouettes in figures were extracted from 'phylopic' (https.phylopic.org) using *rphylopic*

(Gearty, Jones & Chamberlain, 2023). All data, code, and package versions used are available at

https://osf.io/b8km9.

(a) Meta-analyses

To provide a global test of the AAH across animals and to assess the impact of moderator variables,

we ran both frequentist General Linear-Mixed Effects Models (GLMMs) using REML method and the

corresponding Bayesian models. We consider both approaches as complementary, with the Bayesian

method having the advantage of estimating the precision of the obtained measures of heterogeneity

(Harrer *et al.*, 2021). However, for brevity, we report the model results of the Bayesian approach

only in the Supporting Information. Bayesian models were run with 4 chains, each with 4000

iterations, a warmup of 2000 iterations, and were checked for convergence. We ran all models for

datasets of within- and among-species comparisons separately. First, we obtained a global effect size

(r) from models in which the effect size obtained from primary studies is defined as the response

variable while weighted by the inverse of its sampling variance and with study identifier, effect size

identifier, taxon name and the phylogeny (transformed into a correlation matrix) as random effects

to account for nonindependence (including no moderators as fixed predictor terms). This was done

because the final dataset included multiple effect sizes from species with shared phylogenetic

history, from the same species and/or from the same study. Thereby, global effect sizes are

estimated while giving more weight to more precise estimates (i.e., those with smaller Vr). To

correct for non-independence of sampling variances of effect sizes (Vr) in REML models, such as

shared control statistics, we constructed a variance-covariance matrix quantifying the non-zero

covariance between sampling errors within the same primary studies, following the methods by

Nakagawa et al. (2023b). We imputed this matrix by assuming that sampling errors within the same

study are correlated with a constant $\rho = 0.5$. Additionally, we checked the robustness of our study

results against different values of ρ (0.3, 0.7, and 0.9). Second, each null-model (i.e., including no

fixed predictor term) was fitted separately for each taxonomic class (i.e., Aves, Mammalia,

Amphibia). Third, we assessed if biological and methodological moderators impacted effect sizes.

Specifically, we ran models with the same random factors as above and defined acoustic variable

(minimum frequency, mean frequency, maximum frequency, frequency bandwidth, peak frequency,

duration, interval duration, and repetition rate), habitat classification (categorical versus

continuous), habitat contrast (0 or 1), body size correction (yes or no), or phylogenetic correction

(yes or no) as fixed factors. The impact of all moderators was tested separately within birds, which

represents the taxonomic class with the highest number of effect sizes (see Results).

We assessed heterogeneity I^2 by estimating the proportion of variance among effect sizes

that can be attributed to the different levels of random effects (Higgins & Thompson, 2002).

Specifically, we partitioned total heterogeneity (I^2 Total) into variance arising from study identity (I^2

Study), observation (I^2 Observation), taxon (I^2 Taxon), and phylogenetic relationship (I^2 Phylogeny;

Nakagawa & Santos, 2012), which is also known as phylogenetic heritability (H^2), and equivalent to

Pagel's λ (de Villemereuil & Nakagawa, 2014). For REML models with predictor variables, we

calculated the pseudo-R^2 based on the proportional change in the sum of the variance components,

and its 95% confidence interval based on 1000 bootstrap replicates (Viechtbauer, 2023). For models

run with the Bayesian approach, we computed the proportion of variance explained by the entire

model (conditional R^2) and by fixed factors only (marginal R^2; Nakagawa & Schielzeth, 2013).

(b) Sensitivity analyses and publication bias

We explored the robustness of our results by running several sensitivity analyses. Firstly, we identified potential outliers in effect sizes using Grubbs' test (Grubbs, 1950). Secondly, we verified if any individual studies influenced the results of our analysis disproportionally by conducting a leave-one-out analysis as described by Nakagawa *et al.* (2023b). This method involves repeatedly fitting the model to the dataset with one study removed at a time. Thirdly, we checked whether conclusions remained when using Fisher's *z* as effect size. In comparison to Pearson's correlation coefficient *r*, the sampling variance of Fisher's *z* statistics depends only on the sample size but not on the effect size itself. Thus, we transformed our effect size *r* into *z* and computed its sampling variance using guidelines by Borenstein *et al.* (2009).

In order to assess publication bias, we ran a multilevel meta-regression as suggested by Nakagawa *et al.* (2022). We ran a GLMM with Fisher's *z* defined as response variable, its standard error, moderators, and year as fixed effects and study identifier, observation identifier, taxon name and the phylogenetic correlation matrix as random factors, which is advised when heterogeneity is high (see Results; Nakagawa *et al.*, 2022). This aimed at testing (i) whether the effect size depends on its standard error, which, similarly to the Egger's regression (Egger *et al.*, 1997), may indicate that small studies get published only (or more often) if effect sizes are large enough to provide statistically significant support for the tested hypothesis (small-study effect), and (ii) for time-lag bias (Boutron *et al.*, 2019; also known as the bandwagon effect), which suggests that supportive results get published easier in a newly emerging field but as more data are collected and scepticism about the theoretical foundations may arise, non-intuitive results increase (e.g., Jennions & Møller, 2002). As complementary, we used Kendall's rank correlation test (Begg & Mazumdar, 1994) to quantify funnel plot asymmetry. Studies with low precision and/or high sampling error that do not support the tested hypothesis might be more prone to go unpublished. Consequently, the lack of studies meeting these criteria is expected to drive funnel plot asymmetry (Sterne *et al.*, 2011).

III. RESULTS

(1) Dataset summary

In total, we obtained 371 effect sizes from 75 studies and 57 taxa encompassing vertebrate taxa only, i.e., amphibians, birds and mammals (Fig. 2 and Table S3). Of these, 219 effect sizes were from studies reporting within-species comparisons (37 species, including 32 birds, 3 mammals, and 2 amphibians, with 189, 16, and 14 effect sizes, respectively) and 152 were from studies of among-species comparisons (20 taxa, including 14 birds, 3 mammals, and 3 amphibians, with 129, 11, and 12 effect sizes, respectively; a more detailed overview of the sample sizes with respect to the tested moderators is provided in Table S3).

Our dataset included 22 studies (approximately 29% of our dataset) that were already included in a previous meta-analysis focusing on birds by Boncoraglio & Saino (2007). Nonetheless, we obtained different effect sizes for 10 (45 %) of these studies. Out of a total of 78 effect sizes reported in Boncoraglio & Saino (2007), 7 were identical with respect to their absolute value but had the opposite sign, 11 differed by more than 0.1, with 4 of them having the opposite sign, 2 effect sizes could not be retrieved, and 9 differed due to a different analytical approach (most probably because, contrary to Boncoraglio & Saino (2007), we did not pool data from different syllables or songs). Moreover, we excluded 13 effect sizes (from 4 studies) used in the previous meta-analysis because the studies did not fit our selection criteria (i.e., habitats were impossible to classify as open or closed with regard to vegetation structure, habitat was not replicated, the study focused on the effects of urbanization, or the reported data were not sufficient to compute effect sizes). We report these inconsistencies in detail in Table S4.

(2) Test of the AAH for within- and among-species comparisons

Overall, we did not find support for the AAH, neither for within- nor for among-species comparisons (Figs 3 and S3, Tables 1 and S6). None of the tested moderator variables explained a significant

fraction of heterogeneity in effect sizes obtained from within- or among-species comparisons (Figs

S4 and S5, and Tables S5 and S7).

When focusing only on the most studied taxonomic class, Aves, we found weak support for

the AAH for within- and among-species comparisons (Figs 3 and S3). Frequency bandwidth showed a

weak support for the AAH in within-species comparisons (Fig. S6), whereas minimum, maximum and

peak frequencies showed significant support in among-species comparisons (Fig. S7). Analyses of

other acoustic variables did not support the AAH. Primary studies of among-species comparisons

obtained significantly higher effect sizes when they did not account for phylogenetic non-

independence than when they did (Tables S5 and S7). When focusing on primary studies that

accounted for phylogenetic non-independence, we did not obtain a significantly positive global

effect size for birds, which was also the case when testing individual acoustic variables separately

(Fig. S8). Results remained consistent when using different values of ρ (0.3, 0.7, and 0.9) to construct

the variance-covariance matrix quantifying the non-zero covariance between sampling errors within

the same primary studies (Table S8).

In analyses of mammals and amphibians, we found no support for the AAH neither in within-

nor in among-species comparisons (Figs 3 and S3, Tables 1 and S6). For these analyses, the number

of effect sizes and species was much lower compared to birds, which did not allow testing for the

impact of moderators in these taxonomic classes.

The effect sizes we retrieved were highly heterogeneous (REML I^2 Total > 0.80 for all models;

Table 1). For studies of within-species comparisons, heterogeneity arose mostly from variance

among observations. For studies of among-species comparisons, heterogeneity stemmed primarily

from among-study differences for the general model and the model including only birds. For the

model including only studies on birds that accounted for phylogenetic non-independence (phylogeny

correction), heterogeneity arose mostly from variance among species. In models including only

mammals or amphibians, phylogeny explained the highest proportion of total heterogeneity (in both

within- and among-species comparisons). These patterns were confirmed by the Bayesian approach,

which generally estimated higher heterogeneity for among-study and among-observation

differences compared to other factors, except for models with only mammals or amphibians (Table

S6).

(3) Sensitivity analyses and publication bias

Sensitivity analyses revealed that model estimates are robust and reliable. Outlier analyses for

estimates of r of the complete datasets of within- and among-species comparisons (Grubb's test of r:

G = 3.045, P-value = 0.230; G = 2.922, P-value = 0.234; respectively, Fig S9). This was also the case for

within- and among-species comparisons of birds (G = 3.075, P-value = 0.176; G = 2.86, P-value =

0.236; respectively, Fig. S9). An outlier was detected when focusing on primary studies that

accounted for phylogenetic non-independence (i.e., maximum frequency estimate for Parulidae,

Grubb's test of r G = 3.390, P-value = 0.013). We found that there were no influential studies having

disproportionate effects on the model estimates (Fig. S10). Leave-one-out analysis demonstrated

that removing one study at a time did not change the results of within-species comparisons for both

the global model and the model including only birds. In among-species comparisons, the removal of

four studies (one on birds and three on mammals) resulted in marginally overall positive support for

the AAH (Fig. S10 B). However, when focusing solely on studies on birds, the removal of the study on

birds did not affect the overall results (Fig. S10 D). Using Fisher's z instead of the Pearson correlation

coefficient as an effect size provided very similar results suggesting our analysis to be robust with

respect to the chosen effect size metric (Table S9).

We detected no evidence for the small-study effect either for within- and among-species

comparisons, since small studies (with smaller sample size and lower precision) did not show

stronger effect sizes than large studies (Figs S11A and B). In fact, for studies reporting within-species

comparisons, we detected the opposite trend, i.e., a significant negative relationship between effect

size and its standard error (Fig. S11A). In addition, we found no evidence for funnel plot asymmetry

(Figs S11C, D, E, and F) inferred from Kendall's rank correlation test (Kendall's tau [within-species

comparisons] = -0.059, *P-value* = 0.212; Kendall's tau [among-species comparisons] = -0.005, *P-value*

= 0.938).

Finally, we also detected no evidence for a time-lag bias since the year of publication did not

predict effect sizes of within-species comparisons (slope = 0.001, 95% CI = [-0.006, 0.009], z-value =

0.380, P-value = 0.705) or among-species comparisons (slope = -0.007, 95% CI = [-0.016, 0.001], z-

value = -1.791, P-value = 0.073). We observed a negative relationship for studies not accounting for

phylogenetic non-independence and a slightly positive relationship for studies that accounted for it,

but these relationships were not statistically significant (Figs S11G and H).

IV. DISCUSSION

The acoustic adaptation hypothesis (AAH) posits that animals produce acoustic signals that have

evolved in response to the structure of their habitat, but ever since its original formulation, there

has been much uncertainty about the hypothesis' validity and generality within and among species.

We analysed a dataset of 371 effect sizes across 75 studies and including birds, mammals, and

amphibians to demonstrate that acoustic variables do not differ significantly between habitats with

different vegetation structure in both within- and among-species comparisons. Therefore, current

empirical evidence does not support the idea that the AAH provides a general explanation for the

evolution of acoustic signals. Only within-species comparisons of birds provided weak support for

this hypothesis indicating that variation in acoustic signals in relation to vegetation structure may be

more complex and context-dependent than previously thought. Below, we discuss these findings,

identify the limitations of our study, and propose directions for future research.

(1) Limited support for the AAH

Our phylogenetically informed synthesis provides no overall support for the AAH, offering

quantitative evidence that acoustic signals do not show predictable responses to vegetation

structure across vertebrates. This result might seem unexpected given the established principles of

the physics of sound transmission (Hardt & Benedict, 2021) and questions the broader utility of the

AAH for understanding the evolution of acoustic communication in natural settings.

We acknowledge that our final dataset exhibits a taxonomic bias with an overrepresentation

of birds. While this reflects a limitation of our study, it also underscores the evident need for

additional research examining the influence of vegetation structure on acoustic signals of taxa

beyond birds—a call that has been made by numerous authors before (e.g., Ey & Fischer, 2009;

Goutte *et al.*, 2018; Mikula *et al.*, 2021) but remains largely unaddressed. As research extends to

taxa that are currently understudied, the inclusion of reptiles, insects, and more species of anurans

and mammals would likely enable the examination of other potential moderators, such as signal

development (i.e., innate versus learned acoustic signals), which has been suggested suggested by

other authors to influence results, but that could not be tested in this study (see Text S1). This would

ultimately contribute to a more comprehensive understanding of acoustic adaptation and animal

communication.

The substantial variation in effect sizes across our dataset (REML I^2 Total > 0.80 for all the

models; Table 1), a common pattern of meta-analyses in ecology and evolution (with the average I^2

being as high as 92%; Senior *et al.*, 2016), remained largely unexplained. For within-species

comparisons, most of the heterogeneity originated from variation among observations, while

differences among studies, taxa or phylogeny effects had less impact. This among-observation

variability may reflect the contrasting support of the AAH within studies depending on the acoustic

variables analysed, sometimes even within the same acoustic variables when measured for the

entire signal and its fragments (i.e., for studies that measured acoustic variables across the entire

song as well as its syllables). On the other hand, the heterogeneity of among-species comparisons

arose mostly from variation among studies, which could be due to differences in study design such

as using a different assessment of vegetation structure, a focus on different acoustic variables, or

differences in statistical correction for body size or phylogenetic non-independence. Remarkably,

only a fraction of primary studies of among-species comparisons accounted for phylogenetic non-

independence and our analyses suggest that this methodological aspect has an impact on the study's

outcome, particularly in birds. In contrast to a qualitative review by Ey & Fischer (2009), our results

suggest that studies not accounting for phylogenetic non-independence tend to obtain stronger

support for the AAH than those that do. This underscores the need for comparative studies to

consider phylogenetic relatedness to avoid misinterpreting the role of vegetation structure in

shaping acoustic signals.

Why then does the AAH not seem to apply universally? Several factors may contribute to the

limited support for the AAH. First, the complexity of the habitat and methodological difficulties

regarding its classification could render the AAH an oversimplified concept if applied too broadly.

Second, the diversity of acoustic signals across taxa and the multiplicity of selective pressures acting

on them may lead to complex and diverse responses that may override a potential effect of the

habitat.

(a) Complexity of the habitat

Although the AAH predicts that selection should favour acoustic signals that propagate well in an

environment, it has been primarily tested in terms of effects of vegetation on sound transmission.

This approach might be too simplistic, as many other elements may influence sound propagation in a

given habitat (Attenborough & Renterghem, 2021). Physical features of the landscape (hills, valleys,

or slopes), type of substrate, weather conditions, as well as background noise (arising from wind,

moving water, or other species) are only a few of the factors that can also affect how sound travels

(Sueur, Krause & Farina, 2019; Guibard *et al.*, 2022). For example, there is increasing evidence that

animals can adjust their acoustic signals to surrounding noise, with most of the studies highlighting

within-species adjustments (Slabbekoorn & Peet, 2003; Gil *et al.*, 2015; Gomes *et al.*, 2022). The

significant heterogeneity detected in our analyses probably reflects variation among studies arising

not only from methodological aspects including differences in data collection, analysis, and reporting

but also from the physical features of the habitat.

Even when focusing on vegetation structure only, characterizing the habitat for the purpose

of investigating animal acoustic communication proves to be a challenging and not yet standardized

task. Early work on the AAH used broad habitat categories (forest, edge, and grassland; Morton,

1975) and general predictions were then formulated for open versus closed habitats. These

classifications of vegetation structure do not only oversimplify specific habitat characteristics crucial

for sound transmission but also tend to be subjective and prone to ambiguity. Even the use of more

quantitative continuous variables does not resolve this issue because a wide array of both small-

scale (measured at a microhabitat level, e.g., number of trees) and large-scale (representative of

macrohabitat characteristics, e.g., tree cover, canopy cover) variables have been used, which

challenges the generalization of findings. The differences in the assessment of vegetation structure

among studies may have contributed to the heterogeneity in our results as suggested by Ey &

Fischer (2009). However, we found that support for the AAH did not significantly differ among

studies using categorical versus continuous characterization of vegetation structure, or among

studies with varying magnitudes of contrast between closed and open habitats. This indicates that

including different classifications of the habitat did not change the overall results, possibly reflecting

the lack of power of the current classification to capture vegetation structure. Additionally, within

the same habitat, both season and the time of day can have a significant influence on signal

transmission. For example, an empirical study conducted in two different forests demonstrated that

these factors impacted detection distance by up to a factor of five (Haupert, Sèbe & Sueur, 2022).

(b) Alternative determinants of signal evolution overriding habitat effects

Animal acoustic signals are shaped by a multitude of selective forces, which can give rise to

contrasting responses, potentially reducing the significance of habitat effects, thus challenging the

universality of the AAH. Whilst our results provide no overall evidence for the AAH, we found weak

support for the AAH in birds for within-species comparisons. Avian acoustic signals might therefore

present phenotypically plastic responses to vegetation structure but the weak support observed in

our analysis suggests a minor role, if any, of the vegetation in shaping acoustic signals even in birds.

The lack of support for the AAH in mammals and amphibians aligns with the previous qualitative review (Ey & Fischer, 2009) and may reflect the limited importance of habitat in explaining global signal variation in these taxa. Nevertheless, these results demand caution due to the limited number of species tested for these animal classes.

Further complicating the application of the AAH as a universal framework are diverse taxon-specific constraints associated with each acoustic variable. Anuran calls, for instance, exhibit greater morphological constraints on frequency than on temporal parameters (Escalona Sulbarán *et al.*, 2019; Tonini *et al.*, 2020), and this has been suggested as the reason for temporal parameters being more susceptible to evolve in response to environmental factors compared to spectral ones (Mendoza-Henao *et al.*, 2023). For birds, there seems to be no consensus regarding the acoustic parameters that are more prone to be selected by environmental factors. While the previous meta-analysis found that all frequency variables differed between habitats, consistently supporting the AAH (Boncoraglio & Saino, 2007), Ey & Fischer (2009) identified minimum frequency and frequency bandwidth as the only acoustic variables presenting global support for the AAH. Recent findings propose frequency bandwidth as a more labile acoustic variable, in opposition to peak frequency, which is more constrained by a species' genetic background (Rivera *et al.*, 2023). Although our results focussing on plastic changes (i.e. within-species comparisons) align with the findings from Rivera *et al*. (2023), this is not the case for the results obtained for evolutionary changes (i.e. among-species comparisons), which reveal a lack of support for all the acoustic variables when using studies that accounted for phylogeny.

Even if animals do not adjust acoustic parameters to vegetation structure, they can exhibit other adaptive communication strategies to optimize signal transmission. In particular, they may use aerial displays (Menezes & Santos, 2020) or move to a more suitable microhabitat and height for broadcasting (Kime, 2000; Seddon, 2005; Chitnis, Rajan & Krishnan, 2020) or for listening (Jensen, Larsen & Attenborough, 2008). In the presence of background noise from biological, abiotic, or

urban sources, animals may also adjust the broadcasting time to minimize signal overlap and

competition for 'signal space' with other sound-producing species (Hart *et al.*, 2015; Chronister,

Rhinehart & Kitzes, 2023; Staniewicz *et al.*, 2023). Ultimately, the benefits of long-distance

transmission, such as improved communication over larger distances or enhanced mate attraction,

may be offset by the associated predatory, competitive, and parasitic costs (Ryan *et al.*, 1981;

Garamszegi & Avilés, 2005).

(2) Changing the paradigm

As Brown & Handford (1996) concluded: "*With all the possible selective forces acting on signal*

design, investigators looking for specific or finely detailed differences within or among species, across

habitats, should not be surprised to find results that do not always agree with AAH predictions". This

conclusion is echoed in more recent work (e.g., Ey & Fischer, 2009; Goutte *et al.*, 2018; Mikula *et al.*,

2021) but the AAH has remained a focal point of investigation.

655   The lack of support for the AAH in our analyses calls for a more comprehensive and

integrative approach to study acoustic adaptation, considering the complex interplay of ecological

and evolutionary factors. We propose this field to move beyond the current confines of the AAH by

highlighting promising avenues for future studies.

(a) Standardizing habitat attenuation measurements to refine AAH predictions

We believe that advances in understanding how environmental features affect acoustic signalling

can be achieved through a different approach of how the habitat is characterised and analysed. This

involves refining the description of habitats and adopting a broader perspective, moving beyond a

sole focus on vegetation structure. Ey & Fischer (2009) suggested a detailed characterization of each

habitat, including altitude, humidity, temperature, local animal acoustic signals, and vegetation

structure. However, obtaining and interpreting such an extensive amount of information at the

relevant scale for communication may pose practical challenges due to the multitude of factors

involved and difficulties in comparing studies. Instead of considering multiple factors of the habitat

separately, it may be preferable to use a single variable that integrates the direct effects of multiple

factors on sound propagation.

A promising approach to characterise the acoustic properties of the environment has been

emphasised recently by Haupert *et al*. (2022). By recording ambient sound and broadcasting white

noise and recording it at various distances from the emitter in two forest types, they found that the

coefficient of attenuation (a_0, expressed in dB/kHz/m) can be used to describe accurately the

attenuation characteristic of an environment for large frequency bandwidths (for 5 kHz frequency

bandwidths between 0-20 kHz in their study). This singular parameter is modelled based on acoustic

physics rather than descriptive statistics and summarizes the contribution of various variables,

including temperature, relative humidity, atmospheric pressure, distance, frequency, ambient sound

pressure level, and vegetation, to global attenuation (Haupert *et al.*, 2022). Moreover, Haupert *et al*.

(2022) used the same playback technique to describe the attenuation spectrum (the pattern of

attenuation by frequency) of the two studied forests. Despite having similar a_0 values, both forests

exhibited very different attenuation spectrum profiles. Within the same environment, detection

distance varied by up to a factor of five depending on the frequency considered. These spectrum

profiles potentially reveal frequencies with the highest detection probability and could be used as

starting point to describe the habitat. Subsequently, studies could establish correlations between

the attenuation profile and the species' (or population's, or even individual's) specific power

spectrum of acoustic signals, in order to test whether frequencies with greater detection distance

are preferentially used. New research should also consider that within the same signal, for example

a song, there might be a maximum transmission distance for the detection (i.e., the hearing and

perception of a conspecific through its song) and for decoding of the signal (i.e., deciphering the

conspecific's performance from its song). In summary, we suggest a more comprehensive

examination of attenuation in each locality, employing methodologies such as those outlined by

Haupert *et al*. (2022). These methodologies are applicable in any environment and encompass all

variables that might influence sound, thus offering a more holistic approach. This would facilitate

global comparisons (through a single variable) and offer a more objective and standardized

assessment of the acoustic properties of a given habitat.

Finally, we propose to rephrase the predictions of the AAH to take into account the

attenuation spectrum. Specifically, animals inhabiting environments with different attenuation

spectra are predicted to emit acoustic signals at frequencies (minimum, mean, maximum, and peak

frequency) and times (duration, interval duration, repetition rate) such that detection and/or

decoding distances are maximized. This implies that there is no universal pattern for the expected

change in these variables, unless habitats share similar profiles. Instead, the alterations will be

contingent upon the specific characteristics of the environment. This adjustment also overcomes an

earlier noticed limitation of the AAH that for maximum efficiency, long-range acoustic

communication in any habitat should employ the lowest possible frequencies (Wiley & Richards,

1982).

(b) Towards a narrower framework to test the AAH

While the AAH offers valuable insights, it may not fully capture the processes underlying the

evolution of acoustic signalling as it does not account for alternative strategies that animals use in

response to multiple selective pressures acting on signal structure or their interactions. To

comprehensively assess the environmental influence on signal evolution, a more comprehensive

approach is necessary.

Considering the detected lack of generality of the AAH, it would be valuable to narrow down

the focus to signals (or specific parts of signals) for which an advantage in transmission over long

distances is likely. Studies not specifically selecting for these signals (e.g., Mikula *et al.*, 2021) might

reflect this limitation. For example, many acoustic signals are part of private duets (i.e., do not need

long-distance transmission) such that there is no selection against easily degrading communication

(Larsen, 2020). Likewise, for some species song is not used for distant communication, as reported in

zebra finches, where it may be more pertinent to assess whether their distance call, rather than

their song, is adapted for long distance transmission through the local environment (Loning, Griffith

& Naguib, 2022). This approach requires an in-depth knowledge of the studied species, which might

not always be readily available and can be challenging to obtain. Nevertheless, we believe that tests

of the AAH are only meaningful for signals that are meant for long-distance communication.

V. CONCLUSIONS

(1) Our meta-analysis challenges the general validity of the acoustic adaptation hypothesis

(AAH) by showing no overall support for the correlation between vegetation structure and

acoustic variables.

(2) The weak support for the AAH in within-species comparisons of birds suggests a

phenotypically plastic response of frequency bandwidth to the vegetation structure of the

habitat. By contrast, the AAH is not supported by among-species comparisons of studies

accounting for phylogenetic non-independence.

(3) Although many years have passed since the formulation of the AAH, major efforts are still

required to confirm the predictive ability of the hypothesis. Future research should refine

the characterization of the habitat through a standardized quantification of the attenuation,

which will lead to habitat- and species-specific predictions. Moreover, we expect that major

advances in the field will benefit from broadening the studied taxonomic range while

concurrently being selective about the tested signals.

VI. ACKNOWLEDGEMENTS

We are very grateful to the authors of the primary studies for making their research available, and in

particular those providing additional data or information, and unpublished datasets, including Emma

Greig, Elodie Ey, Darren Proppe, and Martim Melo. T. Michael Keesey, Julián Bayona, and Dennis

Murphy designed the silhouettes used in our figures. We thank Sylvain Haupert, Ole Næsbye Larsen,

Frédéric Sèbe, and Keith Attenborough for their insightful comments and discussions, which

significantly improved the manuscript. We extend our gratitude to Rémi Tournebize for his

assistance with troubleshooting R scripts.

Funding: B.F. was funded by the Foundation for Science and Technology (FCT, Portugal) through a

PhD grant (2020.04569.BD). This research was partly funded by grant PID2021-124501NB-I00 to

B.M. from the Spanish Ministry of Science and Innovation, co-financed by the European Union's

Regional Development Fund (ERDF). C.T. benefited from a grant managed by the Agence Nationale

de la Recherche (CEBA: ANR-10-LABX-25-01). T.J. received funds from the Centre National de la

Recherche Scientifique (CNRS).

VII. AUTHOR CONTRIBUTIONS

B.F. Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project

administration, Visualization, Writing – original draft;

P.B.D. Supervision, Writing – review and editing

B.M. Supervision, Writing – review and editing

C.T. Supervision, Writing – review and editing

T.J. Conceptualization, Data curation, Formal analysis, Investigation, Methodology, Project

administration, Supervision, Validation, Visualization, Writing – review and editing

VIII. DATA AVAILABILITY

All data and R scripts associated with this study can be retrieved from the OSF data repository (it will

be made public upon article acceptance).

IX. REFERENCES

*ACERO-MURCIA, A.C., RAPOSO DO AMARAL, F., DE BARROS, F.C., DA SILVA RIBEIRO, T., MIYAKI, C.Y. &
MALDONADO-COELHO, M. (2021) Ecological and evolutionary drivers of geographic variation in
songs of a Neotropical suboscine bird: The Drab-breasted Bamboo Tyrant (*Hemitriccus diops*,
Rhynchocyclidae). *Ornithology* **138**, ukab003.

*ANDERSON, M.E. & CONNER, R.N. (1985) Northern Cardinal song in three forest habitats in eastern
Texas. *The Wilson Bulletin* **97**, 436–449.

*ARAUJO, D.M.D. (2019) Estrutura do habitat influencia a tolerância climática e frequência dominante
do canto de anúncio do gênero *Rhinella* (Anura: Bufonidae)? Universidade Federal de Goiás,
Goiânia, Brazil.

ATTENBOROUGH, K. & RENTERGHEM, T. VAN (2021) *Predicting outdoor sound*Second edition. CRC Press,
Boca Raton.

AYLOR, D. (1972) Noise Reduction by Vegetation and Ground. *The Journal of the Acoustical Society of*
America **51**, 197–205.

*BADYAEV, A.V. & LEAF, E.S. (1997) Habitat Associations of Song Characteristics in *Phylloscopus* and
Hippolais Warblers. *The Auk* **114**, 40–46.

*BAQUERO LOZANO, A.C. (2008) Divergence of adaptative phenotypic traits and mate recognition
systems (song and color) in response to recent anthropogenetic habitat changes in an
oceanic island (São Tomé, Gulf of Guinea) Endemic passerine (*Speirops lugubris*). MSc Thesis,
Universidad de Los Andes, Bogotá, Colombia.

BEGG, C.B. & MAZUMDAR, M. (1994) Operating Characteristics of a Rank Correlation Test for
Publication Bias. *Biometrics* **50**, 1088.

BERTELLI, S. & TUBARO, P.L. (2002) Body mass and habitat correlates of song structure in a primitive
group of birds. *Biological Journal of the Linnean Society* **77**, 423–430.

*Bicudo, T., Anciães, M., Benchimol, M., Peres, C.A. & Simões, P.I. (2016) Insularization effects on
acoustic signals of 2 suboscine Amazonian birds. *Behavioral Ecology* **27**, 1480–1490.

*Billings, A.C. (2018) The low-frequency acoustic structure of mobbing calls differs across habitat
types in three passerine families. *Animal Behaviour* **138**, 39–49.

BLUMENRATH, S. & DABELSTEEN, T. (2004) Degradation of Great Tit (*Parus major*) Song Before And After
Foliation: Implications for Vocal Communication in a Deciduous Forest. *Behaviour* **141**, 935–
958.

*Blumstein, D.T. & Turner, A.C. (2005) Can the acoustic adaptation hypothesis predict the structure
of Australian birdsong? *Acta ethologica* **8**, 35–44.

BONCORAGLIO, G. & SAINO, N. (2007) Habitat structure and the evolution of bird song: a meta-analysis
of the evidence for the acoustic adaptation hypothesis. *Functional Ecology* **21**, 134–142.

BORENSTEIN, M., HEDGES, L.V., HIGGINS, J.P.T. & ROTHSTEIN, H.R. (2009) *Introduction to Meta-Analysis*.
John Wiley & Sons, Chichester, West Sussex, UK.

BOSCH, J. & DE LA RIVA, I. (2004) Are frog calls modulated by the environment? An analysis with anuran
species from Bolivia. *Canadian Journal of Zoology* **82**, 880–888.

BOUTRON, I., PAGE, M.J., HIGGINS, J.P., ALTMAN, D.G., LUNDH, A., HRÓBJARTSSON, A., & ON BEHALF OF THE
COCHRANE BIAS METHODS GROUP (2019) Considering bias and conflicts of interest among the
included studies. In *Cochrane Handbook for Systematic Reviews of Interventions* (eds J.P.T.
HIGGINS, J. THOMAS, J. CHANDLER, M. CUMPSTON, T. LI, M.J. PAGE & V.A. WELCH), pp. 177–204, 1st
edition. Wiley.

BRADBURY, J.W. & VEHRENCAMP, S.L. (2011) *Principles of animal communication* 2nd edition. Sinauer
Associates, Sunderland, Mass.

BROWN, T.J. & HANDFORD, P. (1996) Acoustic signal amplitude patterns: A computer simulation
investigation of the acoustic adaptation hypothesis. *The Condor* **98**, 608–623.

BROWN, T.J. & HANDFORD, P. (2000) Sound Design for Vocalizations: Quality in the Woods, Consistency
in the Fields. *The Condor* **102**, 81–92.

*BUENO-ENCISO, J., FERRER, E.S., BARRIENTOS, R. & SANZ, J.J. (2016) Habitat structure influences the song
characteristics within a population of Great Tits *Parus major*. *Bird Study* **63**, 359–368.

BÜRKNER, P.-C. (2017) **brms** : An *R* Package for Bayesian Multilevel Models Using *Stan*. *Journal of
Statistical Software* **80**, 1–28.

*BUSKIRK, J.V. (1997) Independent evolution of song structure and note structure in American wood
warblers. *Proceedings of the Royal Society of London. Series B: Biological Sciences* **264**, 755–
761.

CATCHPOLE, C. & SLATER, P.J.B. (2008) *Bird song: biological themes and variations* 2nd edition.
Cambridge University Press, Cambridge.

CATCHPOLE, C.K. (1987) Bird song, sexual selection and female choice. *Trends in Ecology & Evolution* **2**,
94–97.

CHAMBERLAIN, S.A., HOVICK, S.M., DIBBLE, C.J., RASMUSSEN, N.L., VAN ALLEN, B.G., MAITNER, B.S., AHERN,
J.R., BELL-DERESKE, L.P., ROY, C.L., MEZA-LOPEZ, M., CARRILLO, J., SIEMANN, E., LAJEUNESSE, M.J. &
WHITNEY, K.D. (2012) Does phylogeny matter? Assessing the impact of phylogenetic
information in ecological meta-analysis: Phylogenetic meta-analysis. *Ecology Letters* **15**,
627–636.

CHAPPUIS, C. (1971) Un exemple de l'influence du milieu sur les émissions vocales des oiseaux:
l'évolution des chants en forêt équatoriale. *La terre et la vie* **2**, 183–202.

CHITNIS, S.S., RAJAN, S. & KRISHNAN, A. (2020) Sympatric wren-warblers partition acoustic signal space
and song perch height. *Behavioral Ecology* **31**, 559–567.

CHRONISTER, L.M., RHINEHART, T.A. & KITZES, J. (2023) When birds sing at the same pitch, they avoid
singing at the same time. *Ibis* **165**, 1047–1053.

*COLOMBELLI-NÉGREL, D. & SMALE, R. (2018) Habitat explained microgeographic variation in Little
Penguin agonistic calls. *The Auk: Ornithological Advances* **135**, 44–59.

CORPORATION FOR DIGITAL SCHOLARSHIP & ROY ROSENZWEIG CENTER FOR HISTORY AND NEW MEDIA (2022)
Zotero. Fairfax, Virginia, USA. www.zotero.org.

*COSENS, S.E. & FALLS, J.B. (1984) A comparison of sound propagation and song frequency in
temperate marsh and grassland habitats. *Behavioral Ecology and Sociobiology* **15**, 161–170.

*CROUCH, N.M.A. & MASON-GAMER, R.J. (2019) Identifying ecological drivers of interspecific variation
in song complexity in songbirds (Passeriformes, Passeri). *Journal of Avian Biology* **50**,
jav.02020.

CUMMINGS, M.E. & ENDLER, J.A. (2018) 25 Years of sensory drive: the evidence and its watery bias.
Current Zoology **64**, 471–484.

*DERRYBERRY, E.P. (2009) Ecology shapes birdsong evolution: variation in morphology and habitat
explains variation in white-crowned sparrow song. *The American Naturalist* **174**, 24–33.

DERRYBERRY, E.P., SEDDON, N., DERRYBERRY, G.E., CLARAMUNT, S., SEEHOLZER, G.F., BRUMFIELD, R.T. & TOBIAS,
J.A. (2018) Ecological drivers of song evolution in birds: Disentangling the effects of habitat
and morphology. *Ecology and Evolution* **8**, 1890–1905.

*DOUTRELANT, C., LEMAITRE, O. & LAMBRECHTS, M.M. (2001) Song variation in Blue Tit Parus caeruleus
populations from Corsica and mainland southern France. *Ardea* **89**, 375–385.

EGGER, M., SMITH, G.D., SCHNEIDER, M. & MINDER, C. (1997) Bias in meta-analysis detected by a simple,
graphical test. *BMJ* **315**, 629–634.

EMBLETON, T.F.W. (1996) Tutorial on sound propagation outdoors. *The Journal of the Acoustical
Society of America* **100**, 31–48.

ENDLER, J.A. (1992) Signals, Signal Conditions, and the Direction of Evolution. *The American Naturalist*
139, S125–S153.

*ERDTMANN, L.K. (2012) Propagação e reconhecimento específico do canto de anúncio de anuros: no
tempo e no espaço. PhD Thesis, Instituto Nacional de Pesquisas da Amazônia, Manaus,
Brazil.

ESCALONA SULBARÁN, M.D., IVO SIMÕES, P., GONZALEZ-VOYER, A. & CASTROVIEJO-FISHER, S. (2019)
Neotropical frogs and mating songs: The evolution of advertisement calls in glassfrogs.
Journal of Evolutionary Biology **32**, 163–176.

EY, E. & FISCHER, J. (2009) The 'Acoustic Adaptation Hypothesis' - a review of the evidence from birds,
anurans and mammals. *Bioacoustics* **19**, 21–48.

*EY, E. & FISCHER, J. (2011) Keeping in Contact: Flexibility in Calls of Olive Baboons. In *Primates of
Gashaka: Socioecology and conservation in Nigeria's biodiversity hotspot* (eds V. SOMMER &
C. ROSS), pp. 413–436. Springer New York, New York, NY.

*EY, E., RAHN, C., HAMMERSCHMIDT, K. & FISCHER, J. (2009) Wild female olive baboons adapt their grunt
vocalizations to environmental conditions. *Ethology* **115**, 493–503.

*FARNSWORTH, A. & LOVETTE, I.J. (2008) Phylogenetic and ecological effects on interspecific variation in
structurally simple avian vocalizations. *Biological Journal of the Linnean Society* **94**, 155–173.

FLETCHER, N.H. (2004) A simple frequency-scaling rule for animal communication. *The Journal of the
Acoustical Society of America* **115**, 2334–2338.

*FRANCIS, C.D. & WILKINS, M.R. (2021) Testing the strength and direction of selection on vocal
frequency using metabolic scaling theory. *Ecosphere* **12**, e03733.

*FREEMAN, B.G., MONTGOMERY, G.A. & SCHLUTER, D. (2017) Evolution and plasticity: Divergence of song
discrimination is faster in birds with innate song than in song learners in Neotropical
passerine birds. *Evolution* **71**, 2230–2242.

*FRIIS, J.I., DABELSTEEN, T. & CARDOSO, G.C. (2021) Contingency and determinism in the evolution of
bird song sound frequency. *Scientific Reports* **11**, 11600.

FRIIS, J.I., SABINO, J., SANTOS, P., DABELSTEEN, T. & CARDOSO, G.C. (2022) Ecological adaptation and
birdsong: how body and bill sizes affect passerine sound frequencies. *Behavioral Ecology* **33**,
798–806.

GARAMSZEGI, L.Z. & AVILÉS, J.M. (2005) Brood parasitism by brown-headed cowbirds and the
expression of sexual characters in their hosts. *Oecologia* **143**, 167–177.

*GARCÍA-NAVAS, V. & BLUMSTEIN, D.T. (2016) The effect of body size and habitat on the evolution of
alarm vocalizations in rodents. *Biological Journal of the Linnean Society* **118**, 745–751.

GEARTY, W., JONES, L.A. & CHAMBERLAIN, S. (2023) rphylopic: an R package for accessing and plotting
PhyloPic silhouettes. https://cran.r-project.org/web/packages/rphylopic/index.html.

GIL, D. & GAHR, M. (2002) The honesty of bird song: multiple constraints for multiple traits. *Trends in*
Ecology & Evolution **17**, 133–141.

GIL, D., HONARMAND, M., PASCUAL, J., PÉREZ-MENA, E. & MACÍAS GARCIA, C. (2015) Birds living near
airports advance their dawn chorus and reduce overlap with aircraft noise. *Behavioral*
Ecology **26**, 435–443.

*GOGOLEVA, S.S. (2020) Effects of ecological factors on the acoustic parameters of passerine species
in a tropical lowland forest in southern Vietnam. *Bioacoustics* **29**, 45–60.

GOMES, L., SOLÉ, M., SOUSA-LIMA, R.S. & BAUMGARTEN, J.E. (2022) Influence of Anthropogenic Sounds on
Insect, Anuran and Bird Acoustic Signals: A Meta-Analysis. *Frontiers in Ecology and Evolution*
10, 827440.

*GORETSKAIA, M.I., BEME, I.R., POPOVA, D.V., AMOS, N., BUCHANAN, K.L., SUNNUCKS, P. & PAVLOVA, A.
(2018) Song parameters of the fuscous honeyeater *Lichenostomus fuscus* correlate with
habitat characteristics in fragmented landscapes. *Journal of Avian Biology* **49**, jav-01493.

GOUTTE, S., DUBOIS, A., HOWARD, S.D., MÁRQUEZ, R., ROWLEY, J.J.L., DEHLING, J.M., GRANDCOLAS, P., XIONG,
R.C. & LEGENDRE, F. (2018) How the environment shapes animal signals: a test of the acoustic
adaptation hypothesis in frogs. *Journal of Evolutionary Biology* **31**, 148–158.

GRAFEN, A. (1989) The phylogenetic regression. *Philosophical Transactions of the Royal Society of*
London. B, Biological Sciences **326**, 119–157.

*GREIG, E.I., PRICE, J.J. & PRUETT-JONES, S. (2013) Song evolution in Maluridae: influences of natural and
sexual selection on acoustic structure. *Emu - Austral Ornithology* **113**, 270–281.

GRUBBS, F.E. (1950) Sample Criteria for Testing Outlying Observations. *The Annals of Mathematical*
Statistics **21**, 27–58.

GUIBARD, A., SÈBE, F., DRAGNA, D. & OLLIVIER, S. (2022) Influence of meteorological conditions and
topography on the active space of mountain birds assessed by a wave-based sound
propagation model. *The Journal of the Acoustical Society of America* **151**, 3703–3718.

HANSEN, P. (1979) Vocal learning: its role in adapting sound structures to long-distance propagation,
and a hypothesis on its evolution. *Animal Behaviour* **27**, 1270–1271.

*Hadfield, J.D. (2010) MCMC Methods for Multi-Response Generalized Linear Mixed Models: The
MCMCglmm *R* Package. *Journal of Statistical Software* **33**, 1–22.

*Hasiniaina, A.F., Radespiel, U., Kessler, S.E., Rina Evasoa, M., Rasoloharijaona, S., Randrianambinina,
B., Zimmermann, E., Schmidt, S. & Scheumann, M. (2020) Evolutionary significance of the
variation in acoustic communication of a cryptic nocturnal primate radiation (*Microcebus*
spp.). *Ecology and Evolution* **10**, 3784–3797.

Hardt, B. & Benedict, L. (2021) Can you hear me now? A review of signal transmission and
experimental evidence for the acoustic adaptation hypothesis. *Bioacoustics* **30**, 716–742.

Harrer, M., Cuijpers, P., Furukawa, T.A. & Ebert, D.D. (2021) *Doing meta-analysis with R: a hands-on
guide*First edition. CRC Press, Boca Raton London New York.

Hart, P.J., Hall, R., Ray, W., Beck, A. & Zook, J. (2015) Cicadas impact bird communication in a noisy
tropical rainforest. *Behavioral Ecology* **26**, 839–842.

Haupert, S., Sèbe, F. & Sueur, J. (2022) Physics-based model to predict the acoustic detection distance
of terrestrial autonomous recording units over the diel cycle and across seasons: Insights
from an Alpine and a Neotropical forest. *Methods in Ecology and Evolution* **14**, 614–630.

*Hedwig, D., Verahrami, A.K. & Wrege, P.H. (2019) Acoustic structure of forest elephant rumbles: a
test of the ambiguity reduction hypothesis. *Animal Cognition* **22**, 1115–1128.

Higgins, J.P.T. & Thompson, S.G. (2002) Quantifying heterogeneity in a meta-analysis. *Statistics in
Medicine* **21**, 1539–1558.

*Hill, S.D., Pawley, M.D.M. & Ji, W. (2017) Local habitat complexity correlates with song complexity
in a vocally elaborate honeyeater. *Austral Ecology* **42**, 590–596.

Hinchliff, C.E., Smith, S.A., Allman, J.F., Burleigh, J.G., Chaudhary, R., Coghill, L.M., Crandall, K.A.,
Deng, J., Drew, B.T., Gazis, R., Gude, K., Hibbett, D.S., Katz, L.A., Laughinghouse, H.D., McTavish,
E.J., et al. (2015) Synthesis of phylogeny and taxonomy into a comprehensive tree of life.
Proceedings of the National Academy of Sciences **112**, 12764–12769.

*Hunter, M.L. & Krebs, J.R. (1979) Geographical variation in the song of the great tit (*Parus major*) in
relation to ecological factors. *Journal of Animal Ecology* **48**, 759–785.

Hylton, R. & Godard, R.D. (2001) Song properties of indigo buntings in open and forested habitats.
The Wilson Bulletin **113**, 243–245.

Ivimey-Cook, E.R., Noble, D.W.A., Nakagawa, S., Lajeunesse, M.J. & Pick, J.L. (2023) Advice for
improving the reproducibility of data extraction in meta-analysis. *Research Synthesis
Methods* **14**, 911–915.

Jennions, M.D. & Møller, A.P. (2002) Relationships fade with time: a meta-analysis of temporal
trends in publication in ecology and evolution. *Proceedings of the Royal Society of London.
Series B: Biological Sciences* **269**, 43–48.

Jensen, K.K., Larsen, O.N. & Attenborough, K. (2008) Measurements and predictions of hooded crow
(Corvus corone cornix) call propagation over open field habitats. *The Journal of the
Acoustical Society of America* **123**, 507–518.

JOB, J.R., KOHLER, S.L. & GILL, S.A. (2016) Song adjustments by an open habitat bird to anthropogenic
noise, urban structure, and vegetation. *Behavioral Ecology* **27**, 1734–1744.

KIME, N.M. (2000) The transmission of advertisement calls in Central American frogs. *Behavioral*
Ecology **11**, 71–83.

*KIRSCHEL, A.N.G., BLUMSTEIN, D.T., COHEN, R.E., BUERMANN, W., SMITH, T.B. & SLABBEKOORN, H. (2009)
Birdsong tuned to the environment: green hylia song varies with elevation, tree cover, and
noise. *Behavioral Ecology* **20**, 1089–1095.

*KIRSCHEL, A.N.G., SEDDON, N. & TOBIAS, J.A. (2019) Range-wide spatial mapping reveals convergent
character displacement of bird song. *Proceedings of the Royal Society B: Biological Sciences*
286, 20190443.

*KIRSCHEL, A.N.G., SLABBEKOORN, H., BLUMSTEIN, D.T., COHEN, R.E., DE KORT, S.R., BUERMANN, W. & SMITH,
T.B. (2011) Testing alternative hypotheses for evolutionary diversification in an African
songbird: rainforest refugia versus ecological gradients. *Evolution* **65**, 3162–3174.KLUMP,
G.M. & SHALTER, M.D. (1984) Acoustic Behaviour of Birds and Mammals in the Predator
Context; I. Factors Affecting the Structure of Alarm Signals. II. The Functional Significance
and Evolution of Alarm Signals. *Zeitschrift für Tierpsychologie* **66**, 189–226.

KLUMP, G.M. & SHALTER, M.D. (1984) Acoustic Behaviour of Birds and Mammals in the Predator
Context; I. Factors Affecting the Structure of Alarm Signals. II. The Functional Significance
and Evolution of Alarm Signals. *Zeitschrift für Tierpsychologie* **66**, 189–226.

KOMSTA, L. (2005) outliers: Tests for Outliers. https://CRAN.R-project.org/package=outliers [accessed
8 August 2024].

*KOPUCHIAN, C., LIJTMAER, D.A., TUBARO, P.L. & HANDFORD, P. (2004) Temporal stability and change in a
microgeographical pattern of song variation in the rufous-collared sparrow. *Animal*
Behaviour **68**, 551–559

KUMAR, S., STECHER, G., SULESKI, M. & HEDGES, S.B. (2017) TimeTree: A Resource for Timelines,
Timetrees, and Divergence Times. *Molecular Biology and Evolution* **34**, 1812–1819.

*LACHLAN, R.F. & SERVEDIO, M.R. (2004) Song learning accelerates allopatric speciation. *Evolution* **58**,
2049–2063.

LAGISZ, L. (2022) Around meta-analysis (14): deduplicating bibliographic records. *Inter-Disciplinary*
Ecology and Evolution Lab. http://www.i-deel.org/blog/around-meta-analysis-13-
deduplicating-bibliographic-records [accessed 17 March 2022].

*LAIOLO, P. & ROLANDO, A. (2003) The evolution of vocalisations in the genus *Corvus*: effects of
phylogeny, morphology and habitat. *Evolutionary Ecology* **17**, 111–123.

LAJEUNESSE, M.J. (2009) Meta-Analysis and the Comparative Phylogenetic Method. *The American*
Naturalist **174**, 369–381.

LAJEUNESSE, M.J. (2016) Facilitating systematic reviews, data extraction and meta-analysis with the
METAGEAR package for R. *Methods in Ecology and Evolution* **7**, 323–330.

LAJEUNESSE, M.J. (2021) Automated, semi-automated, and manual extraction of numerical data from
scientific images, plot, charts, and figures. https://CRAN.R-project.org/package=juicr.

LAJEUNESSE, M.J., ROSENBERG, M.S. & JENNIONS, M.D. (2013) Recovering missing or partial data from
studies: a survey of conversions and imputations for meta-analysis. In *Handbook of Meta-*
analysis in Ecology and Evolution pp. 195–206. Princeton University Press, Princeton.

LARSEN, O.N. (2020) To Shout or to Whisper? Strategies for Encoding Public and Private Information
in Sound Signals. In *Coding Strategies in Vertebrate Acoustic Communication* (eds T. AUBIN &
N. MATHEVON), pp. 11–44. Springer International Publishing, Cham.

*LIJTMAER, D.A. & TUBARO, P.L. (2007) A reversed pattern of association between song dialects and
habitat in the rufous-collared sparrow. *The Condor* **109**, 658–667.

LONING, H., GRIFFITH, S.C. & NAGUIB, M. (2022) Zebra finch song is a very short-range signal in the wild:
evidence from an integrated approach. *Behavioral Ecology* **33**, 37–46.

LÜDECKE, D. (2019) esc: Effect Size Computation for Meta Analysis. Zenodo.
https://zenodo.org/record/1249217 [accessed 31 October 2023].

MARTEN, K. & MARLER, P.M. (1977) Sound transmission and its significance for animal vocalization. I.
Temperate habitats. *Behavioral Ecology and Sociobiology* **2**, 271–290.

MARTEN, K., QUINE, D. & MARLER, P. (1977) Sound transmission and its significance for animal
vocalization: II. Tropical forest habitats. *Behavioral Ecology and Sociobiology* **2**, 291–302.

*MASON, N.A., SHULTZ, A.J. & BURNS, K.J. (2014) Elaborate visual and acoustic signals evolve
independently in a large, phenotypically diverse radiation of songbirds. *Proceedings of the*
Royal Society B: Biological Sciences **281**, 20140967.

MCLEAN, M.J., BISHOP, P.J. & NAKAGAWA, S. (2012) Male quality, signal reliability and female choice:
assessing the expectations of inter-sexual selection: Assessing the expectations of inter-
sexual selection. *Journal of Evolutionary Biology* **25**, 1513–1520.

*MEAUX, E., PEABOTUWAGE, I., MAMMIDES, C., MALYKHINA, K., QUAN, R.-C. & GOODALE, E. (2021)
Behavioural variables influence contact call rate more than characteristics of the vegetation
in a group-living passerine species. *Behavioural Processes* **185**, 104345.

*MEJÍAS, M.A., RONCAL, J., IMFELD, T.S., BOISEN, S. & WILSON, D.R. (2020) Relationships of song structure
to phylogenetic history, habitat, and morphology in the vireos, greenlets, and allies
(Passeriformes: Vireonidae). *Evolution* **74**, 2494–2511.

MENDOZA-HENAO, A.M., ZAMUDIO, K.R., GUAYASAMIN, J.M., ESCALONA, M. & PARRA-OLEA, G. (2023)
Environment rather than character displacement explains call evolution in glassfrogs.
Evolution **77**, 355–369.

MENEZES, J.C.T. & SANTOS, E.S.A. (2020) Habitat structure drives the evolution of aerial displays in
birds. *Journal of Animal Ecology* **89**, 482–493.

MICHONNEAU, F., BROWN, J.W. & WINTER, D.J. (2016) rotl: an R package to interact with the Open Tree
of Life data. *Methods in Ecology and Evolution* **7**, 1476–1481.

MIKULA, P., VALCU, M., BRUMM, H., BULLA, M., FORSTMEIER, W., PETRUSKOVÁ, T., KEMPENAERS, B. &
ALBRECHT, T. (2021) A global analysis of song frequency in passerines provides no support for
the acoustic adaptation hypothesis but suggests a role for sexual selection. *Ecology Letters*
24, 477–486.

MOHER, D., LIBERATI, A., TETZLAFF, J. & ALTMAN, D.G. (2009) Preferred Reporting Items for Systematic
Reviews and Meta-Analyses: The PRISMA Statement. *Journal of Clinical Epidemiology* **62**,
1006–1012.

MORTON, E.S. (1975) Ecological sources of selection on avian sounds. *The American Naturalist* **109**,
17–34.

NAGUIB, M. (2003) Reverberation of rapid and slow trills: Implications for signal adaptations to long-
range communication. *The Journal of the Acoustical Society of America* **113**, 1749–1756.

*NAGUIB, M., HAMMERSCHMIDT, K. & WIRTH, J. (2001) Microgeographic variation, habitat effects and
individual signature cues in calls of chiffchaffs *Phylloscopus collybita canarensis*. *Ethology*
107, 341–355.

NAKAGAWA, S., LAGISZ, M., JENNIONS, M.D., KORICHEVA, J., NOBLE, D.W.A., PARKER, T.H., SÁNCHEZ-TÓJAR, A.,
YANG, Y. & O'DEA, R.E. (2022) Methods for testing publication bias in ecological and
evolutionary meta-analyses. *Methods in Ecology and Evolution* **13**, 4–21.

NAKAGAWA, S., LAGISZ, M., O'DEA, R.E., POTTIER, P., RUTKOWSKA, J., SENIOR, A.M., YANG, Y. & NOBLE, D.W.A.
(2023a) orchaRd 2.0: An R package for visualising meta-analyses with orchard plots. *Methods
in Ecology and Evolution* **14**, 2003–2010.

NAKAGAWA, S., NOBLE, D.W.A., SENIOR, A.M. & LAGISZ, M. (2017) Meta-evaluation of meta-analysis: ten
appraisal questions for biologists. *BMC Biology* **15**, 18.

NAKAGAWA, S. & SANTOS, E.S.A. (2012) Methodological issues and advances in biological meta-analysis.
Evolutionary Ecology **26**, 1253–1274.

NAKAGAWA, S. & SCHIELZETH, H. (2013) A general and simple method for obtaining R^2 from generalized
linear mixed-effects models. *Methods in Ecology and Evolution* **4**, 133–142.

NAKAGAWA, S., YANG, Y., MACARTNEY, E.L., SPAKE, R. & LAGISZ, M. (2023b) Quantitative evidence
synthesis: a practical guide on meta-analysis, meta-regression, and publication bias tests for
environmental sciences. *Environmental Evidence* **12**, 8.

*NASCIMENTO, V.D.D. (2014) A evolução do canto em aves : integrando morfologia, filogenia e
ambiente. PhD Thesis, Universidade Federal do Paraná, Curitiba, Brazil.

NEMETH, E., DABELSTEEN, T., PEDERSEN, S.B. & WINKLER, H. (2006) Rainforests as concert halls for birds:
Are reverberations improving sound transmission of long song elements? *The Journal of the
Acoustical Society of America* **119**, 620–626.

NICHOLLS, J.A. & GOLDIZEN, A.W. (2006) Habitat type and density influence vocal signal design in satin
bowerbirds: Habitat-related call variation in bowerbirds. *Journal of Animal Ecology* **75**, 549–
558.

O'DEA, R.E., LAGISZ, M., JENNIONS, M.D., KORICHEVA, J., NOBLE, D.W.A., PARKER, T.H., GUREVITCH, J., PAGE,
M.J., STEWART, G., MOHER, D. & NAKAGAWA, S. (2021) Preferred reporting items for systematic
reviews and meta-analyses in ecology and evolutionary biology: a PRISMA extension.
Biological Reviews **96**, 1695–1722.

OPHIR, A.G., SCHRADER, S.B. & GILLOOLY, J.F. (2010) Energetic cost of calling: general constraints and
species-specific differences. *Journal of Evolutionary Biology* **23**, 1564–1569.

*PANG-CHING, J.M., PAXTON, K.L., PAXTON, E.H., PACK, A.A. & HART, P.J. (2018) The effect of isolation,
fragmentation, and population bottlenecks on song structure of a Hawaiian honeycreeper.
Ecology and Evolution **8**, 2076–2087.

PARADIS, E., CLAUDE, J. & STRIMMER, K. (2004) APE: Analyses of Phylogenetics and Evolution in R
language. *Bioinformatics* **20**, 289–290.

PEARSE, W.D., MORALES-CASTILLA, I., JAMES, L.S., FARRELL, M., BOIVIN, F. & DAVIES, T.J. (2018) Global
macroevolution and macroecology of passerine song. *Evolution* **72**, 944–960.

*PERLA, B.S. & SLOBODCHIKOFF, C.N. (2002) Habitat structure and alarm call dialects in Gunnison's
prairie dog (*Cynomys gunnisoni*). *Behavioral Ecology* **13**, 844–850.

*PETERS, G., BAUM, L., PETERS, M.K. & TONKIN-LEYHAUSEN, B. (2009) Spectral characteristics of intense
mew calls in cat species of the genus *Felis* (Mammalia: Carnivora: Felidae). *Journal of
Ethology* **27**, 221–237.

*PETERS, G. & PETERS, M.K. (2010) Long-distance call evolution in the Felidae: effects of body weight,
habitat, and phylogeny. *Biological Journal of the Linnean Society* **101**, 487–500.

*PICK, J.L., NAKAGAWA, S. & NOBLE, D.W.A. (2019) Reproducible, flexible and high-throughput data
extraction from primary literature: The METADIGITISE R package. *Methods in Ecology and
Evolution* **10**, 426–431.

R CORE TEAM (2022) R: A language and environment for statistical computing. R Foundation for
Statistical Computing, Vienna, Austria. http://www.R-project.org/.

*RHEINDT, F.E., GRAFE, T.U. & ABOUHEIF, E. (2004) Rapidly evolving traits and the comparative method:
how important is testing for phylogenetic signal? *Evolutionary Ecology Research* **6**, 377–396.

RIVERA, M., EDWARDS, J.A., HAUBER, M.E. & WOOLLEY, S.M.N. (2023) Machine learning and statistical
classification of birdsong link vocal acoustic features with phylogeny. *Scientific Reports* **13**,
7076.

ROTHSTEIN, S.I. & FLEISCHER, R.C. (1987) Vocal Dialects and Their Possible Relation to Honest Status
Signalling in the Brown-Headed Cowbird. *The Condor* **89**, 1–23.

RSTUDIO TEAM (2021) RStudio: Integrated Development Environment for R. RStudio, PBC, Boston, MA.
http://www.rstudio.com/.

RYAN, M.J., TUTTLE, M.D. & TAFT, L.K. (1981) The costs and benefits of frog chorusing behavior.
Behavioral Ecology and Sociobiology **8**, 273–278.

*RYAN, M.J. & WILCZYNSKI, W. (1991) Evolution of intraspecific variation in the advertisement call of a
cricket frog (*Acris crepitans*, Hylidae). *Biological Journal of the Linnean Society* **44**, 249–271.

*SAUNDERS, J. & SLOTOW, R. (2004) The evolution of song structure in southern African birds: an
assessment of the acoustic adaptation hypothesis. *Ostrich* **75**, 147–155.

*SEBASTIÁN-GONZÁLEZ, E., VAN AARDT, J., SACCA, K., BARBOSA, J.M., KELBE, D. & HART, P.J. (2018) Testing
the acoustic adaptation hypothesis with native and introduced birds in Hawaiian forests.
Journal of Ornithology **159**, 827–838.

SEDDON, N. (2005) Ecological adaptation and species recognition drives vocal evolution in Neotropical
suboscine birds. *Evolution* **59**, 200–215.

SENIOR, A.M., GRUEBER, C.E., KAMIYA, T., LAGISZ, M., O'DWYER, K., SANTOS, E.S.A. & NAKAGAWA, S. (2016)
Heterogeneity in ecological and evolutionary meta-analyses: its magnitude and implications.
Ecology **97**, 3293–3299.

*SHY, E. & MORTON, E.S. (1986) Adaptation of amplitude structure of songs to propagation in field
habitat in song sparrows. *Ethology* **72**, 177–184.

*SIMPSON, R.K. & MCGRAW, K.J. (2018) Multiple signaling in a variable environment: expression of
song and color traits as a function of ambient sound and light. *Biotropica* **50**, 531–540.

SLABBEKOORN, H. & PEET, M. (2003) Birds sing at a higher pitch in urban noise. *Nature* **424**, 267–267.

SLABBEKOORN, H. & SMITH, T.B. (2002) Habitat-dependent song divergence in the little greenbul: an
analysis of environmental selection pressures on acoustic signals. *Evolution* **56**, 1849–1858.

*SMITH, J.I. & YU, H.-T. (1992) The association between vocal characteristics and habitat type in
Taiwanese passerines. *Zoological science* **9**, 659–664.

*SMITH, T.B., MILÁ, B., GRETHER, G.F., SLABBEKOORN, H., SEPIL, I., BUERMANN, W., SAATCHI, S. & POLLINGER,
J.P. (2008) Evolutionary consequences of human disturbance in a rainforest bird species
from Central Africa. *Molecular Ecology* **17**, 58–71.

*SORJONEN, J. (1986) Factors affecting the structure of song and the singing behaviour of some
northern European passerine birds. *Behaviour* **98**, 286–304.

STANIEWICZ, A., SOKOŁOWSKA, E., MUSZYŃSKA, A. & BUDKA, M. (2023) Competition for acoustic space in a
temperate-forest bird community. *Behavioral Ecology* **34**, 1043–1054.

STERNE, J.A.C., SUTTON, A.J., IOANNIDIS, J.P.A., TERRIN, N., JONES, D.R., LAU, J., CARPENTER, J., RUCKER, G.,
HARBORD, R.M., SCHMID, C.H., TETZLAFF, J., DEEKS, J.J., PETERS, J., MACASKILL, P., SCHWARZER, G., ET
AL. (2011) Recommendations for examining and interpreting funnel plot asymmetry in meta-
analyses of randomised controlled trials. *BMJ* **343**, d4002.

SUEUR, J., KRAUSE, B. & FARINA, A. (2019) Climate Change Is Breaking Earth's Beat. *Trends in Ecology &*
Evolution **34**, 971–973.

*TIETZE, D.T., MARTENS, J., FISCHER, B.S., SUN, Y., KLUSSMANN-KOLB, A. & PÄCKERT, M. (2015) Evolution of
leaf warbler songs (Aves: Phylloscopidae). *Ecology and Evolution* **5**, 781–798.

TONINI, J.F.R., PROVETE, D.B., MACIEL, N.M., MORAIS, A.R., GOUTTE, S., TOLEDO, L.F. & PYRON, R.A. (2020)
Allometric escape from acoustic constraints is rare for frog calls. *Ecology and Evolution* **10**,
3686–3695.

*TSAO, C.-M. (2016) Call Attributes of Moltrecht's Treefrog *Rhacophorus moltrechti* on Different
Calling Sites. MSc Thesis, National Sun Yat-sen University, Taiwan.

*TUBARO, P.L. & LIJTMAER, D.A. (2006) Environmental correlates of song structure in forest grosbeaks
and saltators. *The Condor* **108**, 120–129.

*Tᴜʙᴀʀᴏ, P.L. & Mᴀʜʟᴇʀ, B. (1998) Acoustic Frequencies and Body Mass in New World Doves. *The*
Condor **100**, 54–61.

*Tᴜʙᴀʀᴏ, P.L. & Sᴇɢᴜʀᴀ, E.T. (1994) Dialect differences in the song of *Zonotrichia capensis* in the
southern Pampas: a test of the acoustic adaptation hypothesis. *The Condor* **96**, 1084–1088.

*Tᴜʙᴀʀᴏ, P.L., Sᴇɢᴜʀᴀ, E.T. & Hᴀɴᴅғᴏʀᴅ, P. (1993) Geographic Variation in the Song of the Rufous-
Collared Sparrow in Eastern Argentina. *The Condor* **95**, 588–595.

Tʏᴀᴄᴋ, P.L. (2020) A taxonomy for vocal learning. *Philosophical Transactions of the Royal Society B:*
Biological Sciences **375**, 20180406.

Vɪᴇᴄʜᴛʙᴀᴜᴇʀ, W. (2010) Conducting Meta-Analyses in *R* with the **metafor** Package. *Journal of*
Statistical Software **36**, 1–48.

Vɪᴇᴄʜᴛʙᴀᴜᴇʀ, W. (2023) Confidence intervals for the amount of heterogeneity accounted for in meta-
regression models. Ghent, Belgium.
https://www.wvbauer.com/lib/exe/fetch.php/talks:2023_viechtbauer_eam_ci_for_r2.pdf.

ᴅᴇ Vɪʟʟᴇᴍᴇʀᴇᴜɪʟ, P. & Nᴀᴋᴀɢᴀᴡᴀ, S. (2014) General Quantitative Genetic Methods for Comparative
Biology. In *Modern Phylogenetic Comparative Methods and Their Application in Evolutionary*
Biology (ed L.Z. Gᴀʀᴀᴍsᴢᴇɢɪ), pp. 287–303. Springer Berlin Heidelberg, Berlin, Heidelberg.

*Wᴀᴀs, J.R. (1988) Song pitch – habitat relationships in white-throated sparrows: cracks in acoustic
windows? *Canadian Journal of Zoology* **66**, 2578–2581.

*Wᴀssᴇʀᴍᴀɴ, F.E. (1979) The relationship between habitat and song in the White-throated Sparrow.
The Condor **81**, 424.

Wᴇʟᴋʟɪɴ, J.F., Bʀᴀɴᴄʜ, C.L., Pɪᴛᴇʀᴀ, A.M., Sᴏɴɴᴇɴʙᴇʀɢ, B.R., Bᴇɴᴇᴅɪᴄᴛ, L.M., Hᴇɪɴᴇɴ, V.K., Kᴏᴢʟᴏᴠsᴋʏ, D.Y.
& Pʀᴀᴠᴏsᴜᴅᴏᴠ, V.V. (2023) Females pair with males larger than themselves in a socially
monogamous songbird. *Animal Behaviour* **199**, 35–49.

Wɪᴄᴋʜᴀᴍ, H. (2016) *ggplot2: elegant graphics for data analysis*Second edition. Springer, Cham.

Wɪᴄᴋʜᴀᴍ, H., Fʀᴀɴᴄ̧ᴏɪs, R., Hᴇɴʀʏ, L. & Mᴜ̈ʟʟᴇʀ, K. (2022) dplyr: A Grammar of Data Manipulation.
https://CRAN.R-project.org/package=dplyr.

*Wɪʟᴇʏ, R.H. (1991) Associations of song properties with habitats for territorial oscine birds of
eastern North America. *The American Naturalist* **138**, 973–993.

Wɪʟᴇʏ, R.H. & Rɪᴄʜᴀʀᴅs, D.G. (1978) Physical constraints on acoustic communication in the
atmosphere: Implications for the evolution of animal vocalizations. *Behavioral Ecology and*
Sociobiology **3**, 69–94.

Wɪʟᴇʏ, R.H. & Rɪᴄʜᴀʀᴅs, D.G. (1982) Adaptations for Acoustic Communication in Birds: Sound
Transmission and Signal Detection. In *Acoustic Communication in Birds* pp. 131–181.
Elsevier.

*Wɪʟʟɪᴀᴍs, J.M. & Sʟᴀᴛᴇʀ, P.J.B. (1993) Does Chaffinch *Fringilla coelebs* song vary with the habitat in
which it is sung? *Ibis* **135**, 202–208.

WINKLER, D.W., BILLERMAN, S.M. & LOVETTE, I.J. (2020) Cardinals and Allies (Cardinalidae). In *Birds of*
the World (eds S.M. BILLERMAN, B.K. KEENEY, P.G. RODEWALD & T.S. SCHULENBERG), p. Cornell Lab
of Ornithology.

*WOOD, C.M., SCHMIDT, S.M. & PEERY, M.Z. (2019) Spatiotemporal Patterns of the California Spotted
Owl's Territorial Vocalizations. *Western Birds* **50**, 232–242.

YU, G., SMITH, D.K., ZHU, H., GUAN, Y. & LAM, T.T. (2017) GGTREE : an R package for visualization and
annotation of phylogenetic trees with their covariates and other associated data. *Methods in*
Ecology and Evolution **8**, 28–36.

ZHAO, L., SANTOS, J.C., WANG, J., RAN, J., TANG, Y. & CUI, J. (2021) Noise constrains the evolution of call
frequency contours in flowing water frogs: a comparative analysis in two clades. *Frontiers in*
Zoology **18**, 37.

*ZIMMERMAN, B.L. (1983) A comparison of structural features of calls of open and forest habitat frog
species in the central Amazon. *Herpetologica* **39**, 235–246.

1195

X. SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

Text S1. Differences between the pre-registered methods and the ones used are explained in this document.

Table S1. Definition of each acoustic variable and the general predictions of the acoustic adaptation hypothesis for a closed habitat in comparison to an open habitat.

Table S2. Source data table of effect sizes included in the meta-analysis.

Table S3. Number of studies and effect sizes for each moderator category included in this meta-analysis for primary studies of within- and among-species comparisons.

Table S4. Studies and effect sizes included in a previous meta-analysis testing the acoustic adaptation hypothesis in birds by Boncoraglio & Saino (2007) and effect sizes included in our study converted to Fisher's z. Inconsistencies between this and our study were divided in six different categories: (1) effect size differs more than 0.10; (2) effect size is similar but differs in sign; (3) effect size could not be retrieved for our study; (4) study excluded from our meta-analysis based on our selection criteria; (5) different methodological approach: authors might have pooled the data and we did not; (6) different methodological approach: we used raw values.

Table S5. Summary statistics of moderator models of within- and among-species comparisons using restricted maximum likelihood approach. Moderators include habitat classification (categorical versus continuous), habitat contrast of categorical classifications (0 versus 1), and body size correction (no versus yes). For comparisons among species, moderators also include phylogeny correction (no versus yes). The first category mentioned in parentheses is used as the reference level for the model results. Phylogenetically controlled multilevel meta-analytic single predictor models are shown with Omnibus tests (Q_M, Wald-type chi-square test) and pseudo-R^2.

Table S6. Global tests using the Bayesian approach. Results of intercept-only general linear-mixed

effects models are shown for the entire dataset (global model) and subsets with respect to animal

class (Aves, Mammalia or Amphibia). For among-species comparisons of birds, the results are also

shown for a subset of studies that accounted for phylogenetic non-independence ('Aves (phylogeny

correction)'). Table shows number of effect sizes (k), number of species or taxonomic groups (N),

estimates of *r* with 95% confidence intervals in parentheses and heterogeneity I^2 (variance arising

from phylogenetic affinities, taxon, study identity, and observation). Model estimates are shown as

posterior modes with 95% Highest Posterior Density (HPD) intervals in brackets.

Table S7. Test of moderators explaining for within- and among-species comparisons using the

Bayesian approach. Moderators include habitat classification (categorical versus continuous), habitat

contrast of categorical classifications (0 versus 1), and body size correction (no versus yes). For

comparisons among species, moderators also include phylogeny correction (no versus yes). The first

category mentioned in parentheses is used as the reference level for the model results. Model

estimates (i.e., estimated difference between groups) are shown as posterior modes with 95%

Highest Posterior Density (HPD) intervals obtained from phylogenetically controlled general linear-

mixed effects models. The variance explained by the entire model and by the fixed factors is given as

the conditional R^2 and the marginal R^2, respectively, with 95% HPD intervals in brackets.

Table S8. Sensitivity analysis examining the robustness of the overall effect size (r) to the assumption

of different ρ values (within-study correlation) used to construct the variance-covariance metrics

that account for non-independence of sampling variances. Table shows estimates of r with 95%

confidence intervals in parentheses.

Table S9. Global effect sizes (Fisher's z) for the acoustic adaptation hypothesis obtained from the

restricted maximum likelihood approach. Results of intercept-only General Linear-Mixed Effects

Models are shown for the entire dataset (global model) and for subsets with respect to animal class

(Aves). For among-species comparisons of birds, the results are shown for a subset of studies that

accounted for phylogenetic non-independence (Aves (phylogeny correction)). Table shows number

of effect sizes (k), number of species or taxa (N), and estimates of Fisher's z and the 95% confidence

intervals in parentheses.

Fig. S1. Phylogeny used to account for phylogenetic non-independence in statistical analysis of

within-species comparisons. Doughnut charts show the proportion of sample sizes (i.e., number of

Pearson correlation coefficients) and the number of species.

Fig. S2. Phylogeny used to account for phylogenetic non-independence in analysis of among-species

comparisons. Doughnut charts show the proportion of sample sizes (i.e., number of Pearson

correlation coefficients) and the number of taxa.

Fig. S3. Density plots showing posterior distributions obtained from Bayesian models for the

differences in Pearson correlation coefficients (r) among animal classes obtained from general linear

mixed-effects models (see Methods). Positive values indicate support for the AAH.

Fig. S4. Orchard plots for within-species comparisons showing individual effect sizes and mean effect

size obtained from REML models for each acoustic variable. Number of effect sizes and number of

taxa (k = number of effect sizes (number of taxa)) are shown on the left for each variable. Bold

horizontal bars are 95% confidence intervals (CIs) and thin black lines are precision intervals. Positive

Pearson correlation coefficients (r) indicate support for the acoustic adaptation hypothesis. Solid

vertical line indicates the estimated global effect size obtained from the REML model with random

factors only, and the grey area denotes the 95% CIs (r = -0.049, 95% CIs: -0.202 to 0.104). Circle size

reflects effect size precision (log(1/SE)).

Fig. S5. Orchard plots for among-species comparisons showing individual effect sizes and mean

effect size from REML models for each acoustic variable. Number of effect sizes and number of taxa

(k = number of effect sizes (number of taxa)) are shown on the left for each variable. Bold horizontal

bars are 95% confidence intervals (CIs) and thin black lines are precision intervals. Positive Pearson

correlation coefficients (r) indicate support for the acoustic adaptation hypothesis. Solid vertical line

indicates the estimated global effect size obtained from the REML model with random factors only,

and the grey area denotes the 95% CIs (r = 0.111, 95% CIs: -0.031 to 0.254). Circle size reflects effect

size precision (log(1/SE)).

Fig. S6. Orchard plots for comparisons within bird species showing individual effect sizes and mean

effect size obtained from REML models for each acoustic variable. Number of effect sizes and

number of taxa (k = number of effect sizes (number of taxa)) are shown on the left for each variable.

Bold horizontal bars are 95% confidence intervals (CIs) and thin black lines are precision intervals.

Positive Pearson correlation coefficients (r) indicate support for the acoustic adaptation hypothesis.

Solid vertical line indicates the estimated global effect size obtained from the REML model with

random factors only, and the grey area denotes the 95% CIs (r = 0.075, 95% CIs: 0.027 to 0.123).

Circle size reflects effect size precision (log(1/SE)).

Fig. S7. Orchard plots for comparisons among bird species showing individual effect sizes and mean

effect size from REML models for each acoustic variable. Number of effect sizes and number of taxa

(k = number of effect sizes (number of taxa)) are shown on the left for each variable. Bold horizontal

bars are 95% confidence intervals (CIs) and thin black lines are precision intervals. Positive Pearson

correlation coefficients (r) indicate support for the acoustic adaptation hypothesis. Solid vertical line

indicates the estimated global effect size obtained from the REML model with random factors only

and the grey area denotes the 95% CIs (r = 0.150, 95% CIs: 0.024 to 0.276). Circle size reflects effect

size precision (log(1/SE)).

Fig. S8. Orchard plots for comparisons that accounted for phylogenetic non-independence

(phylogeny correction) among bird species showing individual effect sizes and mean effect size from

REML models for each acoustic variable. Number of effect sizes and number of taxa (k = number of

effect sizes (number of taxa)) are shown on the left for each variable. Bold horizontal bars are 95%

confidence intervals (CIs) and thin black lines are precision intervals. Positive Pearson correlation

coefficients (r) indicate support for the acoustic adaptation hypothesis. Solid vertical line indicates

the estimated global effect size from the REML model including only phylogeny corrected studies

with random factors only, and the grey area denotes the 95% CIs (r = 0.036, 95% CIs: -0.062 to

0.134). Circle size reflects effect size precision (log(1/SE)).

Fig. S9. Bar and density plots for within- (left) and among-species (right) comparisons showing the

distribution of effect sizes (Pearson correlation coefficient (r)). Plots are shown for entire dataset (A

and B) and for a subset of studies focusing on birds (C and D). Positive Pearson correlation

coefficients (r) indicate support for the acoustic adaptation hypothesis. Solid vertical black lines

indicate the estimated global effect size obtained from the REML model with random factors only for

global models (A: r = -0.049; B: r = 0.111) and for the ones including only comparisons of birds (C: r =

0.075; D: r = 0.150).

Fig. S10. Caterpillar plots of leave-one-out analyses for within- (left) and among-species (right)

comparisons based on datasets with one study left out at a time from model fitting. Plots are shown

for entire dataset (A and B) and for a subset of studies focusing on birds (C and D). Estimated global

effect sizes are ordered by mean estimate, and with horizontal bars showing 95% confidence

intervals (CIs). Positive Pearson correlation coefficients (r) indicate support for the acoustic

adaptation hypothesis. Solid vertical black lines indicate the estimated global effect size obtained

from the REML model with random factors only, and the grey area denotes the 95% CIs for global

models (A: r = -0.049, 95% CIs: -0.202 to 0.104; B: r = 0.111, 95% CIs: -0.031 to 0.254) and for the

ones including only comparisons of birds (C: r = 0.075, 95% CIs: 0.027 to 0.123; D: r = 0.150, 95% CIs:

0.024 to 0.276).

Fig. S11. Plots for the assessment of publication bias for within- (left) and among-species (right)

comparisons. Regressions (A and B) show the relationship between effect size (Fisher's z) and its

standard error testing whether studies with smaller sample sizes and lower precision are more likely

to be published when reporting larger effect sizes. Solid lines show the fit of the multivariate linear

mixed-effects model with year and all the moderators as fixed effects and study identifier, effect size

identifier, taxon name, and phylogeny as random terms (within-species comparisons: slope = -0.887,

95% CI = [-1.766, -0.008], t-value = -1.990, P-value = 0.048; among-species comparisons: slope = -

0.206, 95% CI = [-1.382, 0.970], t-value = -0.346, P-value = 0.730). Circle size reflects precision of

effect sizes (log(1/SE)). Funnel plots for raw values (C and D) and for meta-analytic residuals (E and F)

were obtained from multivariate linear mixed-effects model with all the moderators as fixed effects

and study identifier, effect size identifier, taxon name, and phylogeny as random terms. Dashed lines

indicate the estimated global effect size. Bold and thin black solid lines denote the expected 95% and

99% confidence limits, respectively, purely due to sampling heterogeneity. Asymmetries along the

global effect size may reflect publication biases. Time-lag bias analysis (G and H) showing the

relationship (linear regression) between effect size (Pearson correlation coefficient, r) and the year

of publication of studies that did (and did not) account for phylogenetic non-independence (within-

species comparisons: slope ± SE = -0.001 ± 0.002, t-value = -0.545, P-value = 0.586; among-species

comparisons: slope[yes] ± SE = 0.002 ± 0.004, t-value = 0.514, P-value = 0.609, slope[no] ± SE = -

0.005 ± 0.003, t-value = -1.625, P-value = 0.108). Circle size reflects effect size precision (1/SE).

1334

1335 **Fig. 1**. Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) diagram of

1336 the protocol for screening, including and excluding studies for this meta-analysis and number of

1337 studies identified at each stage. OATD, Open Access Theses and Dissertations.

1338

1339

Fig. 2. Caterpillar plot for (A) within- and (B) among-species comparisons. Effect sizes are displayed

by taxonomic class, ordered by mean estimate, and with horizontal bars showing 95% confidence

intervals (CIs). Symbols of the point estimate illustrate the acoustic variable measured. Positive

Pearson correlation coefficients (*r*) indicate support for the acoustic adaptation hypothesis. Solid

vertical black lines indicate the estimated global effect size obtained from the REML model with

random factors only, and the grey area denotes the 95% CIs (A: *r* = -0.049, 95% CIs: -0.202 to 0.104;

B: *r* = 0.111, 95% CIs: -0.031 to 0.254).

1347

Fig. 3. Orchard plots for (A) within- and (B) among-species comparisons showing effect sizes

(Pearson correlation coefficients, *r*) and number of taxa (k = total number of effect sizes (total

number of taxa) and mean effect size from REML models. Solid horizontal bars are 95% confidence

intervals (CIs) of global effect sizes obtained for each taxon. Solid vertical black lines indicate the

estimated global effect size from the REML model with the random factors only, and grey bands

denote the 95% CIs (A: *r* = -0.049, 95% CIs: -0.202 to 0.104; B: *r* = 0.111, 95% CIs: -0.031 to 0.254).

Circle size reflects precision of effect sizes (log(1/SE)).

Table 1. Global effect sizes for the acoustic adaptation hypothesis obtained from the restricted maximum likelihood approach. Results of intercept-only

General Linear-Mixed Effects Models are shown for the entire dataset (global model) and for subsets with respect to animal class (Aves, Mammalia or

Amphibia). For among-species comparisons of birds, the results are shown for a subset of studies that accounted for phylogenetic non-independence (Aves

(phylogeny correction)). Table shows number of effect sizes (k), number of species or taxonomic groups (N), estimates of r with 95% confidence limits in

parentheses and heterogeneity I^2 (total, variance arising from phylogenetic affinities, taxon, study identity, and observation).

Type of study	Taxonomic range	k	N	r		z-value	P-value	I^2 Total	I^2 Phylogeny	I^2 Taxon	I^2 Study	I^2 Observation
Within species	Global model	219	37	-0.049	(-0.202,0.104)	-0.622	0.534	0.955	0.138	0.000	0.280	0.537
	Aves (all)	189	32	0.075	(0.027,0.123)	3.057	**0.002**	0.937	0.000	0.000	0.000	0.937
	Mammalia	16	3	-0.201	(-0.844,0.441)	-0.614	0.539	0.979	0.000	0.886	0.000	0.092
	Amphibia	14	2	-0.482	(-1.096,0.132)	-1.537	0.124	0.892	0.272	0.272	0.272	0.076
Among species	Global model	152	20	0.111	(-0.031,0.254)	1.532	0.126	0.950	0.055	0.025	0.642	0.228
	Aves (all)	129	14	0.150	(0.024,0.276)	2.331	**0.020**	0.945	0.000	0.020	0.692	0.233
	Aves (phylogeny correction)	67	11	0.036	(-0.062,0.134)	0.724	0.469	0.820	0.000	0.385	0.000	0.435
	Mammalia	11	3	-0.275	(-0.649,0.100)	-1.435	0.151	0.972	0.611	0.000	0.328	0.033
	Amphibia	12	3	0.232	(-0.109,0.574)	1.334	0.182	0.865	0.474	0.000	0.000	0.390

1360