Differential functions of the dorsal and intermediate regions of the hippocampus for optimal goal-directed navigation in VR space

Hyeri Hwang[1], Seung-Woo Jin[2], Inah Lee[1]*

[1]Department of Brain and Cognitive Sciences

Seoul National University

Gwanak-ro 1, Gwanak-gu

Seoul, Korea

08826

[2]Department of Psychiatry and Behavioral Sciences

University of Washington

1959 N.E. Pacific St, Seattle

WA, 98195, USA

*Corresponding author:

E-mail: inahlee@snu.ac.kr

Phone: +82-2-880-8013

Abstract

Goal-directed navigation requires the hippocampus to process spatial information in a value-

dependent manner, but its underlying mechanism needs to be better understood. Here, we

investigated whether the dorsal (dHP) and intermediate (iHP) regions of the hippocampus

differentially function in processing place and its associated value information. Rats were

trained in a place-preference task involving reward zones with different values in a visually

rich VR environment where two-dimensional navigation was possible. Rats learned to use

distal visual scenes effectively to navigate to the reward zone associated with a higher reward.

Inactivation of both dHP and iHP with muscimol altered the efficiency and precision of

wayfinding behavior, but iHP inactivation induced more severe damage, including impaired

place preference. Our findings suggest that the iHP is more critical for value-dependent

navigation toward higher-value goal locations.

Introduction

It has long been suggested that the hippocampus is the neural substrate of a 'cognitive map' – a map-like representation of the spatial environment that allows flexible spatial navigation (O'Keefe and Nadel, 1978). The cognitive map is also needed for remembering important events in space. Animals in the natural environment often navigate to achieve goals, such as finding food or avoiding predators, and this goal-directed navigation involves remembering places and their associated values. It has been reported that the receptive fields of place cells in the hippocampus tend to accumulate near a goal location or shift towards it (Hollup et al., 2001; Kennedy and Shapiro, 2009; Dupret et al., 2010). One could argue that the hippocampus must process task- or goal-relevant information, including the value of a place, to achieve the goal. However, the specific hippocampal processes involved in integrating the two types of representation – place and value – towards goal-oriented behavior are still largely unknown.

Such integration may occur along the dorsoventral axis of the hippocampus. Previous anatomical studies (Krettek and Price, 1977; Swanson et al., 1978; Pikkarainen et al., 1999; Tao et al., 2021) suggest that the hippocampus can be divided along its dorsoventral axis into dorsal (dHP), intermediate (iHP), and ventral (vHP) hippocampal subregions based on different anatomical characteristics. The dHP is connected with brain regions that process visuospatial information, including the retrosplenial cortex and the caudomedial entorhinal cortex (van Groen and Wyss, 2003; Dolorfo and Amaral, 1998); it also communicates with the iHP via bidirectional extrinsic connections but exhibits limited connections with the vHP (Tao et al., 2021; Swanson et al., 1978). The iHP receives heavy projections from valence-related areas, such as the amygdala and ventral tegmental area (VTA) – subcortical inputs that are less prominent in the dHP (Pikkarainen et al., 1999; Felix-Ortiz and Tye, 2014; Gasbarri et al., 1994). The vHP also has connections with the iHP and value-representing areas like the amygdala, but it does not project heavily to the dHP (Tao et al., 2021; Swanson et al., 1978; Pikkarainen et al., 1999; Krettek and Price, 1977). Notably, compared with the dHP, the iHP and vHP have much heavier connections with the medial prefrontal cortex (mPFC), contributing to goal-directed action control (Hoover and Vertes, 2007; Liu and

Carter, 2018). Additionally, the three subregions along the dorsoventral axis display different

gene expression patterns that corroborate the anatomical delineations (Dong et al., 2009;

Bienkowski et al., 2018). Overall, the iHP subregion of the hippocampus appears to be

ideally suited to integrating information from the dHP and vHP.

Surprisingly, beyond the recognition of anatomical divisions, the available literature

on the functional differentiation of subregions along the dorsoventral axis of the hippocampus,

particularly in the context of value representation, is somewhat inconsistent. Specifically,

there is physiological evidence that the size of a place field becomes larger as recordings of

place cells move from the dHP to the vHP (Jung et al., 1994; Maurer et al., 2005; Kjelstrup et

al., 2008; Royer et al., 2010). Thus, it has been thought that the dHP is more specialized for

fine-grained spatial representation than the iHP and vHP. However, when it comes to the

neural representation of value information of a place, results are mixed. Several studies have

reported that place fields recorded in the dHP respond to internal states and motivational

significance based on their accumulation near behaviorally significant locations (e.g., reward

locations or escape platforms; Hollup et al., 2001; Kennedy and Shapiro, 2009; Dupret et al.,

2010), or to the reward per se (Gauthier and Tank, 2018). In contrast, others have reported

that dHP place cells do not alter their activity according to a change in reward or reward

location and thus do not represent value information (Duvelle et al., 2019; Jin and Lee, 2021;

Speakman and O'Keefe, 1990).

Furthermore, although the iHP and vHP have mainly been studied in the context of

fear and anxiety, several studies have also reported spatial representation and value-related

signals in these subregions. Specifically, prior studies reported that rats with a dysfunctional

dHP retained normal goal-directed, target-searching behavior if the iHP and vHP were intact

(Moser et al., 1995; de Hoz et al., 2003). Moreover, lesions in the iHP have been shown to

impair rapid place learning in the water maze task (Bast et al., 2009). Our laboratory also

reported that place cells in the iHP, but not the dHP, instantly respond to a change in spatial

value and overrepresent high-value locations (Jin and Lee, 2021).

Based on existing experimental evidence, we hypothesize that the iHP is the primary

locus for associating spatial representation with value information, distinguishing it from the

dHP and vHP. In the current study, we investigated the differential functions of the dHP and

iHP in goal-directed spatial navigation by monitoring behavioral changes after

pharmacological inactivation of either of the two regions as rats performed a place-preference

task in a two-dimensional (2D) virtual reality (VR) environment. In this experimental

paradigm, rats learned to navigate toward one of two hidden goal locations associated with

different reward amounts. Whereas inactivation of the dHP mainly affected the precision of

wayfinding, iHP inactivation impaired value-dependent navigation more severely by

affecting place preference.

Results

Well-trained rats align themselves toward the high-value zone before departure in the

place-preference task

We established a VR version of a place-preference task (**Figure 1A**) in which rats could

navigate a 2D environment by rolling a spherical treadmill with their body locations fixed,

allowing them to run at the apex of the treadmill. Body-fixed rats (n = 8) were trained to

explore a virtual circular arena surrounded by multiple distal visual landmarks (houses, rocks,

mountains, and trees) (**Figure 1B**). Rats were always started at the center of the arena, and

the arena contained two unmarked reward zones – a high-value zone and a low-value zone –

each associated with different amounts of honey water (6:1 ratio between high- and low-

value zones). A trial started with the rat facing one of six start directions – north (N),

northeast (NE), southeast (SE), south (S), southwest (SW), and northwest (NW) – determined

pseudorandomly to guarantee equal numbers of trials in all directions. In the example trial

shown in **Figure 1C**, the rat was heading in the NE direction at the start location ('Trial

Start'), then turned to the left side to run toward the W goal zone ('Navigation'). Once the rat

arrived at one of the reward zones, the synchronization between the spherical treadmill

movement and the virtual environment stopped, and multiple drops of honey water were

delivered via the licking port ('Reward'). Then, during an inter-trial interval (ITI), the LCD

screens turned gray, and the rat was required to remain still for 5 seconds to initiate the

subsequent trial. The pre-surgical training session consisted of 60 trials, which were reduced

to 40 during post-surgical training.

Pre-surgical training began after shaping in the VR environment. On average, it took

13 days for rats to reach pre-surgical criteria, namely, to complete 60 trials and visit the high-

value goal zone in more than 75% of completed trials (see Methods for the detailed

performance criteria). Well-trained rats exhibited two common behaviors during the pre-

surgical training. First, although it was not required in the task, they learned to rotate the

spherical treadmill counterclockwise to move around in the virtual environment (presumably

to perform energy-efficient navigation). To rule out the potential effect of hardware bias or

any particular aspect of peripheral landscape to make rats turn only to one side, we measured

the direction of the first body-turn in each trial on the last day of shaping and the first day of

the main task (i.e., before rats learned the reward zones). There was no significant difference

between the clockwise and counterclockwise turns (p=0.46 for shaping, p=0.76 for main task;

Wilcoxon signed-rank test), indicating that the stereotypical pattern of counterclockwise

body-turn appeared only after the rats learned the reward locations.

Second, once a trial started, the animal rotated the treadmill immediately to align its

starting direction with the visual scene associated with the high-value reward zone. After

setting the starting direction, the rat started to run on the spherical treadmill, moving the

treadmill forward to navigate directly toward the reward zone. All eight rats displayed this

strategy during the later learning phase but not during the earlier learning stage, suggesting

that the start-scene–alignment strategy was learned during training. Because the initial

rotational scene alignment before departure was an essential component of the task and this

behavior was not readily detectable with position-based analysis, we based most of our

behavioral analysis on the directional information defined by the allocentric reference frame

of the virtual environment (**Figure 2A**). Because we did not measure the rat's head direction

in the current study, the allocentric directional information represented the angular position of

a particular scene in the virtual environment displayed in the center of the screen.

To establish the direction in which the rat departed the starting point at the center

after scene alignment, we first defined a *departure circle* – a virtual circle (~20 cm in

diameter) in the VR environment at the center of the arena (**Figure 2A**). In the example trial

shown in **Figure 2A**, the rat faced the NE direction (315°) at the trial start but immediately

turned its body to the NW direction upon starting and ran straight toward the high-value zone

after that. Since the initial scene rotations at the start point cannot be visualized in the

position-based graph, we made a *scene rotation plot* that visualizes the rotational movement

traces in the virtual environment. The scene rotation plot covers the period from the start of

the trial to when the rat leaves the departure circle (**Figure 2B**).

On the first day of training for the task ('Novice' stage), the rat produced almost no

rotation of the VR environment until he exited the departure circle, indicating that the animal

ran straight in the initially set start direction without adjusting the scene orientation. As a

result, rats missed the target reward zones in most trials (**Figure 2B** and **2C**). However, by

the last day of training ('Expert' stage), there were noticeable rotational shifts in all

directional traces (i.e., counterclockwise rotations) that converged on the high-value reward

zone (**Figure 2B** and **2D**). This was the case for all trials except those in which the initial

start direction almost matched the orientation of the high-value reward zone (i.e., 225° or

NW). Furthermore, the average travel distance and latency for each start direction declined

from the novice to the expert stage, suggesting that the rats navigated more efficiently toward

reward zones in the later stage of learning by pre-adjusting their starting scene direction at the

trial start (**Figure 2C** and **2D**).

Overall, the marked differences in orienting behaviors between early and late

learning stages suggest that rats could discriminate the high-value reward zone from the low-

value zone in our VR environment and show that they preferred visiting the high-value

reward zone over the low-value zone. It also indicates that rats could explore the VR

environment using allocentric visual cues to find the critical scenes associated with the high-

value zone before leaving the starting point (i.e., departure circle).

Departing orientation and perimeter-crossing direction provide a measure of

navigational efficiency and precision, respectively

To analyze behavioral changes during learning in more detail, we analyzed various learning-

related parameters at different stages of pre-surgical training. For this, we focused on days in

which rats visited the high-value zone on more than 75% of trials for two consecutive days –

the performance criterion for completion of pre-surgical training. These two consecutive days

(post-learning days) were grouped and averaged for each rat as the post-learning group

('POST' in **Figure 3A-ii**) and compared with the two consecutive days immediately

preceding the post-learning days (pre-learning days; 'PRE' in **Figure 3A-ii**).

We first measured the departing direction when crossing the departure circle

(Departing Direction [DD]; **Figure 3A-i**). As indicated in **Figure 2**, well-trained rats rotated

the VR environment to place the target VR scene (i.e., high-value reward zone scene) ahead

before departure. Therefore, alignment of the departure direction with the high-value zone at

the beginning of navigation indicated that the rat remembered the scenes associated with the

high-value zone. For example, the distribution of pre-learning days departure directions was

widely distributed without any directional bias; as such, its mean vector was small (**Figure**

3A-ii). On the other hand, the departure directions of post-learning sessions mostly converged

on the direction aligned with the high-value zone, resulting in a larger mean vector length

compared with that in the pre-learning session. The distributions of averaged departure

directions for all rats significantly differed between 'PRE' and 'POST' ($p < 0.001$, Kuiper's

test), verifying that departure direction is a valid index of the acquisition of the high-value

zone and efficient navigation. To investigate how accurately rats oriented themselves directly

to the high-value zone before leaving the start point, we also calculated the average deviation

angle of departure directions (DD-deviation) – the angle between the departure direction and

the high-value zone ($180°$, measured at the center of the zone) – for each rat (**Figure 3A-iii**).

A comparison between pre- and post-learning sessions showed that the DD-deviation

significantly declined after learning. This implies that well-trained rats aligned their bodies

more efficiently to directly navigate to the high-value zone ($p < 0.01$, Wilcoxon signed-rank

test).

Rats adjusted their navigational routes further, even after exiting the departure circle,

to navigate more accurately and straight to the goal, avoiding the wall surrounding the arena.

Such fine spatial tuning (i.e., navigation precision), measured as the decrease in DD-deviation,

only appeared after the rats learned the high-value reward location. To quantify navigation

precision, we measured the perimeter-crossing direction (PCD; **Figure 3B-i**), defined as the

angle at which the rat first touched the unmarked circular boundary along the arena's

perimeter, which shares the inner boundaries of the reward zones (green dashed lines in

Figure 3B-i). During pre-learning, the PCD was randomly distributed along the perimeter

('PRE' in **Figure 3B-ii**). On the other hand, in most post-learning stage trials, rats crossed the

unmarked peripheral boundaries only in the vicinity of the high-value zone ('POST' in

Figure 3B-ii). Since rats usually turned counterclockwise during navigation, the convergence

of crossings near the northern edge of the high-value zone indicates that they took a shortcut

– the most efficient route – to enter the goal zone. The PCD distributions were significantly

different between pre- and post-learning stages ($p < 0.001$; Kuiper's test) (**Figure 3B-ii**). The

deviation angle between the PCD and the high-value zone center also significantly decreased

with learning (**Figure 3B-iii**), indicating that the navigation of rats to the goal became more

accurate.

Additionally, to investigate whether the rats used a certain landmark as a beacon to

find the reward zones, we conducted the landmark omission test as a part of control

experiments. Here, one of the landmarks was omitted, and the landmark to be made disappear

was pseudorandomly manipulated on a trial-by-trial basis. The omission of one landmark,

regardless of its identity, did not cause a specific behavioral change in finding the reward

zones, suggesting that the rats were not relying on a single visual landmark when finding the

reward zones. The result can be reported anecdotally only because of an insufficient sample

size (n=3), not permitting any meaningful statistical testing.

Navigation is impaired by inactivation of either the dHP or iHP, but only iHP

inactivation affects place-preference behavior

To dissociate the roles of the dHP and iHP, we inactivated either the dHP or iHP in an

individual animal using muscimol (MUS), a GABA-A receptor agonist, before the rat

performed the place-preference task. To allow within-subject comparisons in performance,

we bilaterally implanted two pairs of cannulas – one targeting the dHP and the other targeting

the iHP – in the same rat after it successfully reached pre-surgical training criteria (**Figure**

4A). After 1 week of recovery from surgery, rats were retrained to regain a level of

performance similar to that in the pre-surgical training period (**Figure 4B**), after which the

drug injection schedule was started.

We divided rats into two drug-injection groups (n = 4 rats/group) to counterbalance

the injection order between the dHP and iHP. Rats in one group received drug infusion into

the dHP first, whereas rats in the other group were injected into the iHP first. For all rats,

phosphate-buffered saline (PBS) was initially injected in both regions as a vehicle control.

For analytical purposes, we first ensured no statistical difference in performance between the

two PBS sessions (dPBS and iPBS; see below) and then averaged them into a single PBS

session to increase statistical power. During the PBS session, rats tended to take the most

efficient path to the high-value zone, as they had done during pre-surgical training (**Figure**

5A). They aligned the VR scene at the start with the high-value zone for all start directions

and then ran directly toward the goal zone. Notably, once the start scene alignment was

complete, rats usually moved quickly and straight without slowing in the middle. Also, their

navigation paths led them directly toward the center of the goal zone. During subsequent

dHP-inactivation sessions, rats appeared less accurate, bumping into the arena wall in many

trials (dMUS in **Figure 5A**), but most of these wall bumps occurred in the vicinity of the

high-value zone, and rats quickly compensated for their error by turning their bodies to target

the reward zone correctly after wall bumping. In contrast, in iHP-inactivation sessions, the

trajectories were largely disorganized, and the wall-bumping locations were no longer limited

to the vicinity of the high-value zone. In some trials, rats moved largely randomly (as shown

in 860-17-24 in **Figure 5A)** and appeared to visit the low-value zone significantly more than

during PBS or dMUS sessions.

To quantitatively analyze these observations, we compared the proportions of visits

to the high-value zone among drug conditions (**Figure 5B**), finding a significant difference in

the percentage of correct target visits among drug conditions ($F_{(2,14)} = 10.56$, $p < 0.01$, one-

way repeated-measures ANOVA; $p = 0.25$ for dPBS vs. iPBS, Wilcoxon signed-rank test). A

post hoc analysis revealed a significant decrease in the iMUS session compared to the PBS

session ($p < 0.05$, Bonferroni-corrected post hoc test). In contrast, no significant differences

were found in other conditions, although there was a decreasing trend in the iMUS compared

to PBS ($p = 0.2$ for PBS vs. dMUS; $p = 0.1$ in dMUS vs. iMUS). These results indicate that

dHP-inactivated rats show a strong preference for the high-value zone, as they did in control

sessions, but that the performance of iHP-inactivated rats was impaired in our place-

preference task, as reflected in their significantly more frequent visits to the low-value zone

compared with controls.

It is unlikely that these differences stemmed from generic sensorimotor impairment

as a result of MUS infusion because running speed remained unchanged across drug

conditions ($F_{(2,14)} = 0.99$, $p = 0.37$, one-way repeated-measures ANOVA; $p = 0.95$ for dPBS

vs. iPBS, Wilcoxon signed-rank test) (**Figure 5C**). Furthermore, rats remained motivated

throughout the testing session, as evidenced by the absence of a significant difference in the

number of trials across drug groups ($F_{(1.16, 8.13)} = 1.34$, $p = 0.29$, one-way repeated-measures

ANOVA with Greenhouse-Geisser correction; $p = 1.0$ for dPBS vs. iPBS, Wilcoxon signed-

rank test; data not shown), although there was an increase in the session duration ($F_{(2,14)} = $

6.46, $p < 0.05$, one-way repeated-measures ANOVA; $p = 0.27$ for dPBS vs. iPBS, Wilcoxon

signed-rank test; data not shown) during MUS sessions (dMUS and iMUS) compared with

the PBS session (p-values < 0.05, Bonferroni-corrected post hoc test). This increase in

session duration was attributable to arena wall bumping events, which usually entailed a

recovery period before rats left the peripheral boundaries and moved again toward the goal.

These observations indicate that inactivation of the iHP significantly impairs the rat's ability

to effectively navigate to the higher-value reward zone in a VR environment without

affecting goal-directedness or locomotor activity.

To determine how effectively rats traveled to the goal in each condition, we also

quantified the errors made in each condition by assessing the number of perimeter crossings

(**Figure 5D**). To avoid duplicate assessments, we only counted an event as a perimeter

crossing when the rat crossed the perimeter boundary from inside to outside. Rats tended to

make more errors in dMUS sessions compared with controls, and errors were even more

prevalent in iMUS sessions ($F_{(2,14)} = 18.59$, $p < 0.001$, one-way repeated-measures ANOVA;

$p = 0.39$ for dPBS vs. iPBS, Wilcoxon signed-rank test; $p < 0.01$ for PBS vs. dMUS, $p < 0.01$

in PBS vs. iMUS; $p < 0.05$ for dMUS vs. iMUS, Bonferroni-corrected post hoc test). During

PBS sessions, navigation was mostly precise, resulting in just one perimeter crossing. In the

dMUS sessions, precision declined, but the rats were relatively successful in finding the high-

value zone, with most trials being associated with a slightly increased number of perimeter

crossings. In contrast, rats in the iMUS sessions failed to find the high-value zone. They

seemed undirected, exhibiting a significantly increased number of perimeter crossings

compared with the other two sessions. Taken together, these results indicate that iHP

inactivation more severely damages normal goal-directed navigational patterns than dHP

inactivation in our place-preference task.

The iHP causes more damage to value-dependent spatial navigation than the dHP,

which is important for navigational precision

To further differentiate among conditions, we examined departure direction and PCD –

indices of the effectiveness and precision of navigation, respectively (**Figure 3**). We first

investigated the distribution of departure directions in all trials for all rats and calculated the

resultant mean vector (**Figure 6A**). Note that dPBS and iPBS sessions were separately

illustrated here for better visualization of changes in behavioral pattern for each subregion.

Whereas departure directions for both PBS sessions (dPBS and iPBS) were distributed

relatively narrowly towards the high-value zone, those for dMUS sessions were more widely

distributed, and their peak pointed away from the reward zone. In the case of the iHP-

inactivation session, some departure directions were even pointed towards the opposite side

of the target goal zone (i.e., the low-value zone). Thus, the mean vectors from PBS sessions

were relatively longer than those from MUS sessions. The mean vectors for PBS sessions

also stayed within the range of the high-value zone, whereas those for MUS sessions pointed

either toward the edge of the reward zone (dMUS) or the outside of the reward zone (iMUS).

We next quantitatively confirmed these observations, comparing the mean direction

for each drug condition to determine how inactivation affected the accuracy of the body

alignment of rats at departure (**Figure 6A**). A Watson-Williams test indicated that the mean

angles of departure directions in all four drug conditions (dPBS, iPBS, dMUS, and iMUS) for

all rats significantly differed from each other ($F_{(3,1253)} = 7.78$, $p < 0.001$). Post hoc pairwise

comparisons showed that inactivation of either the dHP or iHP significantly altered departure

directions compared with the PBS condition ($p < 0.05$ for dPBS vs. dMUS; $p < 0.001$ for

iPBS vs. iMUS; $p = 0.66$ for dPBS vs. iPBS; Watson-Williams test). Moreover, the mean

departure directions of dMUS and iMUS sessions were displaced from the center of the high-

value zone compared with those of PBS sessions (i.e., dPBS and iPBS), suggesting that the

rats did not accurately align themselves to the target reward zone at the time of departure.

The mean vector of the iMUS session also appeared smaller than that of the other conditions,

indicating a less concentrated distribution of departure directions with iHP inactivation.

Unfortunately, it was not possible to perform a statistical comparison between dMUS and

iMUS because the departure directions for the iMUS session were too dispersed to yield a

mean vector with a sufficient length to compare directions between the two conditions

(averaged mean vector length of dMUS and iMUS sessions < 0.45; Berens, 2009).

The mean vector lengths for departure directions were also significantly different

among drug conditions ($F_{(2,14)} = 12.64$, $p < 0.001$, one-way repeated-measures ANOVA; $p =$

0.55 for dPBS vs. iPBS, Wilcoxon signed-rank test) (**Figure 6B**), with a post hoc analysis

showing a significant difference between the iMUS session and both PBS ($p < 0.01$) and

dMUS ($p = 0.05$) sessions; however, no significant difference was found between PBS and

dMUS sessions ($p = 0.24$, Bonferroni-corrected post hoc test). The profound performance

deficits in the iHP-inactivated condition were also confirmed by examining the DD-deviation

from the target direction, defined as the center of the high-value zone (**Figure 6C**).

Specifically, we found that DD-deviations were significantly different among drug conditions

($F_{(2,14)} = 13.37$, $p < 0.001$, one-way repeated-measures ANOVA; $p = 0.38$ for dPBS vs. iPBS,

Wilcoxon signed-rank test), with a Bonferroni-corrected post hoc test revealing a significant

increase in DD-deviation in the iMUS session compared with both the PBS session ($p < 0.01$)

and the dMUS session ($p < 0.05$). Again, no significant difference was found between PBS

and dMUS sessions ($p = 0.19$). These results demonstrate that disruption of the dHP does not

significantly affect the ability of rats to orient themselves effectively at departure to target the

high-value reward zone. In contrast, inactivation of the iHP across all trials caused rats to

depart the starting location without strategically aligning to the scene and consequently

failing to hit the target zone effectively and directly.

Next, we ran similar analyses for the PCD (**Figure 7**), also investigating the PCD

distribution and its mean vector for each drug condition (**Figure 7A**). PCD distributions

appeared similar to those for departure direction; the PCD distributions of PBS sessions were

narrowly contained within the high-value reward zone, whereas those of MUS sessions were

more dispersed and misaligned with the reward zone. Again, the PCD distribution of the

iMUS session showed some occurrences near the low-value zone. An examination of the

resulting mean vectors using a Watson-Williams test revealed a significant difference in

mean PCD angle in all sessions except for the comparison between the two PBS sessions

($F_{(3,1253)}$ = 16.22, p < 0.001; p = 0.08 for dPBS vs. iPBS). The mean PCD angle of the dMUS

session was shifted toward the upper end of the high-value zone (p < 0.001 for dPBS vs.

dMUS), whereas that of the iMUS session was outside of the reward zone (p < 0.001 for

iPBS vs. iMUS). Notably, iHP inactivation resulted in more severe errors in finding the high-

value zone than dHP inactivation (p < 0.01 for dMUS vs. iMUS). Interestingly, with iHP

inactivation, several PCDs were found near the low-value zone, an outcome that rarely

occurred in other conditions. Considering the decreased percentage of high-value zone visits

(**Figure 5**), some of these trials ended with the rat visiting the low-value zone, suggesting an

impaired ability of the animal to perform goal-directed navigation strategically.

The PCD mean vector length was largest in the PBS condition, shortest in the iMUS

condition, and intermediate in the dMUS condition ($F_{(2,14)}$ = 15.67, p < 0.001, one-way

repeated-measures ANOVA; p = 0.55 for dPBS vs. iPBS, Wilcoxon signed-rank test) (**Figure**

7B). Unlike the mean vector length for departure direction, the PCD mean vector length

differed between PBS and dMUS sessions, suggesting that wayfinding behavior is explicitly

disrupted by dHP inactivation, albeit to a lesser extent compared with iHP inactivation (p <

0.05 for PBS vs. dMUS; p < 0.01 for PBS vs. iMUS; p = 0.06 for dMUS vs. iMUS,

Bonferroni-corrected post hoc test).

On the other hand, the PCD deviation angle from the center of the high-value zone

increased in inverse order: smallest in the PBS condition and largest in the iMUS condition

($F_{(2,14)}$ = 17.24, p < 0.001, one-way repeated-measures ANOVA; p = 0.55 for dPBS vs. iPBS,

Wilcoxon signed-rank test) (**Figure 7C**). Similar to the PCD mean vector length data, the

significant increase in deviation angle after dHP inactivation indicates that dHP-inactivated

rats failed to achieve fine spatial tuning toward the high-value zone compared with controls

(p < 0.05 for PBS vs. dMUS, Bonferroni-corrected post hoc test). iHP inactivation also

resulted in less accurate navigation, including perimeter crossings – effects that were more

severe than those caused by dHP inactivation (p < 0.01 for PBS vs. iMUS; p = 0.06 for

dMUS vs. iMUS, Bonferroni-corrected post hoc test).

Overall, results based on the PCD measure revealed that dHP-inactivated rats showed

decreased precision in arriving at the goal, as reflected in the significant deviation of their

PCD from the high-value zone. The PCD distribution was also not as narrow as under control

conditions. Notably, deficits in navigation performance were even more severe in rats with

iHP inactivation, and their performance impairment was qualitatively different from that

observed with dHP inactivation in terms of both efficiency and precision of navigation. Again,

these results suggest that, while the dHP is essential for accurate wayfinding, the iHP is

crucial for value-dependent navigation to the higher-reward location.

Hippocampal inactivation does not impair cue-guided navigation or goal-directedness

After the drug-injection stage, we trained five of the same rats used in the main task in a

visual cue-guided navigation task to verify whether MUS inactivation of the hippocampus

resulted in deficits in goal-directed navigation in general (**Figure 8A-i**). We used the same

circular arena from the main task but removed all allocentric visual landmarks. The rat was

started from a fixed location near the periphery of the arena. As the trial started, a spherical

visual landmark with a checkered pattern flickered on either the left or right side

(pseudorandomized across trials) of the rat's starting position, serving as a beacon. When the

rat arrived at the landmark area (see Methods), the connection between treadmill movement

and the virtual environment stopped, and honey water was provided as a reward. The rewards

provided by left and right reward zones were the same in terms of both quality and quantity.

In this version of the navigation task, the rat's navigation was simply guided by the

visual beacon, a type of task that the literature suggests is not hippocampus-dependent

(Morris et al., 1986; Packard et al., 1989). Rats learned the task rapidly. Specifically, it took 3

days on average for rats to reach the criterion of completing 40 trials with an excess travel

distance of less than 0.1 meters (see Methods). Moreover, examining their trajectories

suggested that rats had no problem moving toward the visual landmark, whether it appeared

on the left or right of the starting location (**Figure 8A-ii**). Rats arrived at the reward zone

directly in most trials ('Direct hit') but bumped into the arena wall in some trials. Given the

presence of a strong visual landmark, which served as a beacon, trials in which the rat

bumped the arena walls were considered failed trials (**Figure 8A-ii**).

Finally, we applied the same drug injection schedule for the main task after the rats

reached the abovementioned criterion. A one-way repeated-measures ANOVA revealed that

the proportion of direct-hit trials did not significantly differ across drug conditions, indicating

no significant change in navigation precision when the goal was marked by the visual beacon

($F_{(2,8)} = 1.60$, p = 0.26; p = 0.50 for dPBS vs. iPBS, Wilcoxon signed-rank test) (**Figure 8B**).

These results also imply that no generic sensory-motor or motivational deficits were involved.

Collectively, these observations confirm that MUS injections in the hippocampus do not alter

the ability of the rat to move around freely in the VR environment in a goal-directed fashion

when the hippocampus is not necessary for the task.

Discussion

In the current study, we inactivated the dorsal or intermediate hippocampal region in rats

performing a place-preference task in VR space to investigate the functional differentiation

along the dorsoventral hippocampal axis during goal-directed navigation. Inactivation of the

intermediate region, but not the dorsal region, of the hippocampus produced a marked

reduction in the rat's ability to conduct strategic goal-directed navigation in the virtual space

without affecting goal-directedness or locomotor ability. We further examined navigational

quality by measuring the precision of scene alignment upon departure and by assessing the

efficiency (i.e., directness) of travel to the target goal zone (i.e., higher-value zone) without

bumping into walls on the arena boundaries. We found that dHP-inactivated rats were

modestly, but significantly, impaired, not only in precisely targeting the goal at the time of

departure but also in effectively traveling to the goal zone, compared with controls.

Importantly, however, the ability of these dHP-inactivated rats to head toward the higher-

value zone in the VR environment was unimpaired. In contrast, iHP-inactivated rats were

severely impaired in the initial targeting of the goal zone at the time of departure and traveled

somewhat aimlessly in the VR environment compared with both controls and dHP-

inactivated rats. Our findings suggest that the dHP is essential for finding the most effective

travel path for precise spatial navigation and that the iHP is necessary for navigating the

space in a value-dependent manner to achieve goals.

472 Rats use allocentric visual scenes and landmarks to target the goal zone and adjust their

473 paths accordingly during navigation in the VR environment

In the current paradigm, rats rotated the spherical treadmill counterclockwise immediately

after the trial started at the VR arena's center, presumably to find the visual scene to guide

them directly toward the goal zone (i.e., high-value zone) upon departure. This initial

orientation of departure – or departure direction (DD) – seems critical in our task, as

evidenced by the fact that, during training, rats that miscalculated the departure direction

usually bumped into the wall and had to reorient themselves at various positions within the

environment. Once the rats learned the task, they oriented themselves before leaving the start

point by rotating the visual environment until they found the goal-associated visual scenes

and then ran straight toward the goal zone. These behavioral characteristics suggest that our

task is heavily dependent on the rat's ability to use the allocentric reference frame of the

visual environment.

Prior studies suggest that, in an environment where the directional information comes

largely from allocentric visual cues, the spiking activity of place cells is significantly

modulated by directional visual cues, a finding that holds in both real and virtual

environments (Acharya et al., 2016; Ravassard et al., 2013; Aronov and Tank, 2015). In one

of these studies (Acharya et al.), directional modulation of place cells was observed even

during random foraging in the absence of a goal-directed memory task. Notably, this was also

true for spatial view cells in nonhuman primates (Rolls and O'Mara, 1995; Georges-Francois

et al., 1999). Although we did not record place cells in our study, hippocampal place cells

could be predicted to exhibit directional firing patterns associated with the visual scenes

along the periphery of the current VR environment. Because rats in the current study rotated

the environment until they found the target-matching scene without leaving the center starting

point, our VR task may be an ideal behavioral paradigm for examining the directional firing

of place cells in future studies.

499 Inactivating the dHP impairs navigational precision but does not affect place preference

500 based on differential reward values

Our working model posits that the dHP represents a fine-scaled spatial map of an

environment, in this case, a VR environment, that allows an animal to map its location

precisely and choose the most efficient travel routes. Our experimental results support this

model, demonstrating that dHP-inactivated rats deviated slightly, but significantly, from the

ideal target heading at the time of departure (measured by departure direction), resulting in

crossing the area boundary near the target goal zone. Nonetheless, it is important to note that

dHP-inactivated rats in our study oriented themselves normally in the direction of the high-

value reward zone at the time of departure, suggesting that the value-coding cognitive map

and its use were intact and able to spatially guide the rats to the high-reward area in the

absence of a functioning dHP. We argue that such intact place preference performance with

reasonable spatial navigation ability is supported by the iHP (presumably in connection with

the vHP) in dHP-inactivated rats.

Whether the dHP represents value signals remains a matter of controversy.

According to previous studies, place fields of the dHP seem to translocate to or accumulate

near the location with motivational significance (e.g., reward zone), and where the strategic

importance is higher (e.g., choice point in the T-maze) (Hollup et al., 2001; Lee et al., 2006;

Kennedy and Shapiro, 2009; Dupret et al., 2010; Ainge et al., 2011; Valenti et al., 2018). For

instance, the overrepresentation of the escape platform in a water maze – a location of high

motivational significance – was observed in the neural firing patterns of place cells in the

hippocampus (Hollup et al., 2001). In addition, Lee et al. reported that dHP place fields

gradually translocate toward the goal arm of a continuous T-maze (Lee et al., 2006), and

Dupret and colleagues suggested a goal-directed reorganization of hippocampal place fields

based on an experimental paradigm in which reward locations were changed daily (Dupret et

al., 2010). Such accumulation of spatial firing is not restricted to the goal location, as place

fields recorded from the dHP were reported to be unevenly distributed near the start box and

the choice point of a T-maze (Ainge et al., 2011). One potential explanation for the

discrepancy between our study and studies that reported apparent valence-dependent signals

in the dHP could be that the dHP processes motivational and strategic significance (from the

perspective of task demand), which is not always the same as the reward. Significance might

include task demand, such as a change between random and directed search of reward

(Markus et al., 1995), or a change in a significant environment stimulus from which the goal

location needs to be calculated (Gothard et al., 1996). However, none of these were changed
in our experimental paradigm, which might explain why dHP inactivation did not affect
place-preference behavior.

Another possibility is that the dHP responds only to a more radical change in value,
such as the presence or absence of reward, but not to different amounts of the same reward.
Indeed, hippocampal neuronal activity does not show an explicit response to reward value in
rats trained to visit arms of a plus-maze in descending order of reward amount (Tabuchi et al.,
2002). Moreover, when the reward is unexpectedly altered to a less preferred one, thus
decreasing motivational significance, place cells in the dHP remain mostly unchanged (Jin
and Lee, 2021). These results suggest that the dHP is not crucial to maintaining value
preference, a finding in line with the observed absence of an effect on place preference after
dHP inactivation in our study. A recent study in which mice were trained to associate a
particular odor with an appetitive outcome and distinguish it from the non-rewarded odor
suggested that the dHP is responsible for stimulus identity, not saliency (Biane et al., 2023).
This might be another possible interpretation of our dHP results since we used the same
honey water reward for both reward zones.

The iHP may contain a value-associated cognitive map with reasonable spatial
resolution for value-based navigation

iHP-inactivated rats showed poor goal-directed navigation, characterized by misalignment of
their departing orientation with the goal zone and arrival points that were often far removed
from the goal zone compared with the same rats under both control and dHP-inactivated
conditions. Particularly, rats changed their heading directions during the navigation when
they were not confident with the location of the higher reward, resulting in a less efficient
route to the goal location. Rats showing this type of behavior tended to hit the perimeter of
the arena first before correcting their routes. Therefore, when considered together with DD,
our PCD measure could tell that the rats not hitting the goal zone directly after departure were
impaired in orienting themselves to the target zone accurately from the start, not in
maintaining the correct heading direction to the goal zone at the start location.

Although there is still a possibility that the levels of expression of GABA-A

receptors might be different along the longitudinal axis of the hippocampus, these results

support our working model that the iHP is critical for representing a value-associated

cognitive map of the environment. iHP-inactivated rats, presumably unable to utilize such a

value-representing map, could not strategically plan and organize their behaviors to target the

high-value area in the current study. Consequently, the fine-grained spatial map present in the

dHP may be of little use without the guidance of the value-associated map in the iHP,

accounting for the poor navigational performance of iHP-inactivated rats. The value-

associated cognitive map in the iHP may still show reasonable spatial specificity, as

evidenced by the larger, but still specifically located, place fields in the iHP compared with

the dHP (Jin and Lee, 2021).

The involvement of the iHP in spatial value association has been reported or

implicated in several studies. For example, Bast and colleagues reported that rapid place

learning is disrupted by removing the iHP and vHP, even when the dHP remains undamaged

(Bast et al., 2009). On the other hand, if the iHP is spared but the dHP and vHP are removed

by lesioning, rats in a water maze test quickly learn a new platform location normally.

Moreover, a change in reward value induced an immediate global remapping response and a

greater overrepresentation of the reward zone with a higher value in iHP neurons (Jin and Lee,

2021). Another recent study by Jarzebowski et al. focused on how hippocampal place cells

change their firing patterns during the learning process for several sets of changing reward

locations (Jarzebowski et al., 2022). The results from this study suggest that, in the iHP, the

same place cells persistently fire across different reward locations, thus tracking the changes

in reward locations.

Anatomically, the iHP is in an ideal position to represent associations between a

space and its value by intrahippocampal connections from both the dHP and vHP (Tao et al.,

2021; Swanson et al., 1978). Importantly, the vHP is known to receive much heavier

projections from value-processing subcortical areas, such as the amygdala and VTA,

compared with the upper two-thirds of the hippocampus (Krettek and Price, 1977; Swanson

et al., 1978; Pikkarainen et al., 1999; Felix-Ortiz and Tye, 2014; Gasbarri et al., 1994). Thus,

although the iHP also receives afferent projections from these areas, it is highly likely that the

vHP plays a crucial role in the value-related representation of the iHP. Notably, both the

amygdala and VTA are known to be involved in processing palatability information (Tye and

Janak, 2007; Fontanini et al., 2009; Chen et al., 2020), and the amygdala has a subpopulation

of neurons dedicated to encoding positive values (Kim et al., 2016; Beyeler et al., 2016).

These anatomical studies support our working model of the iHP in integrating place-value

information.

It is worth noting that the iHP sends direct projections to the mPFC, which is thought

to be involved in behavioral control and action (Hoover and Vertes, 2007; Liu and Carter,

2018). Our experimental paradigm required rats to choose and navigate towards one of two

reward zones with different values, a task structure that must demand active cognitive control,

presumably by the mPFC in collaboration with the hippocampus. It is also possible that

inactivation of the iHP prevents the transfer of the dHP's spatial information to the mPFC via

the iHP, which may explain why iHP inactivation produces severe deficits in goal-directed

navigation in the current task. Based on these findings, we propose a working model in which

the iHP associates spatial value information with the cognitive map of the dHP and sends

value-associated spatial information to the mPFC, which translates the space-value–

integrated representation into action (Bast, 2011; Bast et al., 2009).

Limitations

We tested the differential functions of the hippocampal subregions in the long axis, dHP and

iHP, by inactivating each subregion during goal-directed navigation. The subregional

inactivation allowed us to compare the differences in navigational patterns directly between

the drug conditions within subjects. However, our study includes only behavioral results and

further mechanistic explanations as to the processes underlying the behavioral deficits require

physiological investigations at the cellular level. Neurophysiological recordings during VR

task performance could answer, for example, the questions such as whether the value-

associated map in the iHP is built upon the map inherited from the dHP or it is independently

developed in the iHP. Also, although our observations and behavioral data strongly suggest

that rats rely on allocentric visual scenes in the VR environment instead of a single or limited

set of landmarks, it is still difficult to prove experimentally whether rats used the cognitive

map of the virtual arena to find the high-value zone or they had an alternative strategy to find

the goal.

Methods

Subjects

Eight male Long-Evans rats (8 weeks old) were housed individually under a 12-hour

light/dark cycle in a temperature- and humidity-controlled environment. Rats were food-

restricted to maintain ~80% of their free-feeding weight, but water was provided ad libitum.

The experimental protocol (SNU-200504-3-1) complied with the guidelines of the

Institutional Animal Care and Use Committee of Seoul National University. Based on our

prior studies (Park et al., 2017; Yoo and Lee, 2017; Lee et al., 2014), the sample size of our

study was set to the least number to achieve the necessary statistical power in the current

within-subject study design for ethical commitments and practical considerations (i.e.,

relatively long training periods).

2D VR system

We established our own VR environment consisting of a circular arena surrounded by

multiple landmarks using a game engine (Unreal Engine [UE] 4.14.3; Epic Games, Inc., USA)

(**Figure 1A** and **1B**). The VR environment was presented via five adjacent LCD monitors

covering 270 degrees of the visual field. Rats were body-restrained at the top of a spherical

treadmill, and a silicone-coated Styrofoam ball with 400mm diameter was placed on multiple

ball bearings. Rats could move their heads freely; body jackets were used to anchor their

positions, limiting their body movements to a 120-degree range. As rats rolled the treadmill,

their movement was recorded by three rotary encoders (DBS60E-BGFJD1024; Sick, Inc.,

Germany) attached to the treadmill surface. The signal from the encoders was then sent to the

computer and synchronized with the movement in the virtual environment via an Arduino

interface board (Arduino Leonardo; Arduino, Italy) and MATLAB R2021a (MathWorks,

USA). A licking port was placed in front of the rats and moved in association with their body

movement. It was maintained in a retracted position but was extended toward the snout by a

linear motor (L16-R; Actuonix Motion Devices, Canada); an infrared sensor (FD-S32;

Panasonic Industry, Japan) detected rats' tongues to record licking behavior. When rats

arrived at either reward zone, the solenoid valve (VA212-3N; Aonetech, Republic of Korea),

controlled by the UE via the Arduino interface (Arduino UNO; Arduino, Italy), dispensed

honey water as a reward. The amount of honey water dispensed for high-value and low-value

zones was maintained at a ratio (in drops) of 12:2, with 12 μL per drop.

Behavioral paradigm

After several days of handling, rats were moved to the VR apparatus and trained to roll the

treadmill to navigate the virtual environment ('Shaping'; **Figure 4B**). In this session, rats had

to reach a flickering checkerboard-shaped sphere randomly spawned on a circular arena (1.6

meters in diameter) to obtain a honeywater reward. After rats had completed more than 60

trials on two consecutive days, they were assumed to have adapted to navigating freely. They

were moved to pre-surgical training ('Pre-training') – a 2D VR version of the place-

preference task. For the pre-training session, rats were required to find hidden reward zones

using the surrounding scene, including various landmarks, such as houses, mountains, and

arches, on the same circular arena from the shaping session. The start position was located at

a fixed point in the center of the arena, and reward zones were located at the east and west

sides of the circular platform; reward zones were positioned at a slight distance from the

arena wall to prevent rats from employing a thigmotaxis strategy. Therefore, the shortest path

length between the start position and the reward zone was 0.62 meters. A trial started with a

heading in one of six start directions, pseudorandomly chosen, and ended when the rat arrived

at either reward zone. Pre-surgical training criteria were defined by the number of trials (60

trials in 40 minutes), high-value zone visit percentage (>75%), and average excess travel

distance (<0.6 meters). If a rat successfully achieved training criteria 2 days in a row, it

received cannula implantation surgery.

After the surgery, rats were allowed 1 week of recovery ('Recovery') and then were moved to

post-surgical training ('Post-training'). During post-training, rats were tested on the same

place-preference task until they achieved the same criteria as pre-surgical training, except that

the trial number was reduced to 40 and the average excess travel distance was reduced to less

than 1 meter. This point marked the beginning of the drug injection stage; four rats received

their initial injection in the dHP, and the other four rats were injected first in the iHP to

counterbalance the injection order ('Place Preference Task').

Object-guided navigation task

After completing drug injections, we trained five of the eight rats from the main task for an

object-guided navigation task to investigate whether drug infusion caused any motor- or

motivation-related impairments ('Probe'; **Figure 4B**). Note the smaller sample size in the

object-guided navigation task. This was because the task was later added to the study design.

In this task, the rat simply had to find and navigate toward a flickering object; because there

was no need for the rat to use the surrounding scene to locate the reward, this probe test was

hippocampus-independent. For the probe test, the rat started from the south of the arena;

concurrently, a flickering checkerboard-shaped sphere appeared on either the left or right side

of the screen. When the rat arrived at the reward zone (i.e., a 0.4m-radius circle surrounding

the object), the visual stimulus stopped, a honeywater reward was given, and the trial ended.

No landmarks surrounded the circular arena to distinguish the environment from that in the

main task. The criterion for training included the completion of 40 trials with less than 0.1

meter of mean excess travel distance, calculated as the shortest path length between the start

location and the reward zone; a time limit of 60 seconds was also imposed. The drug infusion

schedule from the place-preference task was then repeated, at which point rats were sacrificed

for histological procedures.

Surgery

After rats reached pre-surgical training criteria, they were implanted with four commercial

cannulae (P1 Technologies, USA), bilaterally targeting the dHP and the iHP, enabling within-

subject comparisons in performance between dHP inactivation (dMUS) and iHP inactivation

(iMUS) conditions (**Figure 4A**). Animals were first anesthetized with an intraperitoneal

injection of sodium pentobarbital (Nembutal, 65 mg/kg), then their heads were fixed in a

stereotaxic frame (Kopf Instruments, USA). Isoflurane (0.5-2% mixed with 100% oxygen)

was used to maintain anesthesia throughout the surgery. The cannula tips targeted

approximately the upper blades of the dentate gyrus of both regions (AP -3.8 mm, ML ±2.6

mm, DV -2.7 mm for the dHP; AP -6.0 mm, ML ±5.6 mm, DV -3.2 mm with a 10-degree tilt

for the iHP) to inactivate each subregion effectively. The cannula, consisting of a 26-gauge

guide cannula coupled with a 33-gauge dummy cannula, was fixed to the target location by

several skull screws and bone cement. After the surgery, ibuprofen syrup was orally

administered for pain relief, and the animal was kept in an intensive care unit overnight.

Drug infusion

For drug injection, the rat was first anesthetized with isoflurane. Then, 0.3 to 0.5 μl of either

phosphate-buffered saline (PBS) or muscimol (MUS; 1mg/ml, dissolved in saline) was

infused into each hemisphere via a 33-gauge injection cannula at an injection speed of 0.167

μl/min, based on our previous study (Lee et al., 2014; Kim et al., 2012). The injection

cannula and dummy cannula extended 1 mm below the tip of the guide cannula. The injection

cannula was left in place for 1 minute after completing the drug infusion to ensure stable

diffusion of the drug. Then, it was slowly removed from the guide cannula and replaced by

the dummy cannula. The rat was kept in a clean cage to recover from anesthesia completely

and monitored for side effects for 20 minutes, then was moved to the VR apparatus for

behavioral testing. If the rat showed any side effect, particularly sluggishness or aggression,

we reduced the drug injection amount in the rat by 0.1 μl until we found the dosage with

which there was no visible side effect. As a result, five of the rats received 0.4 μl, two

received 0.3 μl, and one received 0.5 μl.

Histology

After completing the probe test, animals were sacrificed by inhalation of an overdose of CO_2.

Rats were then transcardially perfused, first with PBS, administered with a syringe, and then

with a 4% v/v formaldehyde solution, delivered using a commercial pump (Masterflex Easy-

Load II Pump; Cole-Parmer, USA). The brain was extracted and placed in a 4% v/v

formaldehyde-30% sucrose solution at 4℃ until it sank to the bottom of the container. After

gelatin embedding, the brain was sectioned at 40 µm using a microtome (HM430; Thermo-

Fisher Scientific, USA), and sections were mounted on subbed slide glasses for Nissl staining.

Statistical analysis

Data were statistically analyzed using custom programs written in MATLAB R2021a

(MathWorks, USA), Prism 9 (GraphPad, USA), and SPSS (IBM, USA). Statistical

significance was determined using the Wilcoxon signed-rank test and one-way repeated-

measures analysis of variance (RM ANOVA) followed by a Bonferroni post hoc test.

Although most of our statistics were based on the non-parametric tests for the relatively small

sample size (n=8), we used the parametric RM ANOVA for comparing three groups (i.e.,

PBS, dMUS, and iMUS) because it is the most commonly known and widely used statistical

test in such comparison. However, we also performed statistical test with the alternatives for

reference, and the statistical significances were not changed with any of the results. For

directional analysis, Kuiper's test and Watson-Williams test were used. However, the latter

test was considered inapplicable for the mean angle when the average mean vector length

between two samples was less than 0.45 (Berens, 2009). The significance level was set at

$\alpha = 0.05$, and all error bars indicate the standard error of the mean (SEM).

Competing Interests

The authors have no conflicts of interest to declare.

Acknowledgments

This work was supported by the National Research Foundation of Korea

(2019R1A2C2088799, 2021R1A4A2001803, 2022M3E5E8017723) and the Global Ph.D.

Fellowship program (2019H1A2A1073456). We thank Heesoo Oh for his assistance in

behavioral training.

Data Availability

The data used in this study are deposited on GitHub at

https://github.com/hhwang28/Behavioral-Data.

References

Acharya L, Aghajan ZM, Vuong C, Moore JJ, Mehta MR (2016) **Causal influence of visual**
cues on hippocampal directional selectivity *Cell* **164**: 197–207.

Aronov D, Tank DW (2014) **Engagement of neural circuits underlying 2D spatial**
navigation in a rodent virtual reality system *Neuron* **84**: 442–456.

Bast T (2011) **The hippocampal learning-behavior translation and the functional**
significance of hippocampal dysfunction in schizophrenia *Current Opinion in*
Neurobiology **21**: 492–501.

Bast T, Wilson IA, Witter MP, Morris RGM (2009) **From rapid place learning to**
behavioral performance: A key role for the intermediate hippocampus *PLoS Biology*
7: e1000089.

Berens P (2009) **CircStat: A MATLAB toolbox for circular statistics** *Journal of Statistical*
Software **31**: 1–21.

Beyeler A, Namburi P, Glober GF, Simonnet C, Calhoon GG, Conyers GF, Luck R, Wildes
CP, Tye KM (2016) **Divergent routing of positive and negative information from the**
amygdala during memory retrieval *Neuron* **90**: 348–361.

Biane JS, Ladow MA, Stefanini F, Boddu SP, Fan A, Hassan S, Dundar N, Apodaca-
Montano DL, Zhou LZ, Fayner V, Woods NI, Kheirbek MA (2023) **Neural dynamics**
underlying associative learning in the dorsal and ventral hippocampus *Nature*
Neuroscience **26**: 798–809.

Bienkowski MS, Bowman I, Song MY, Gou L, Ard T, Cotter K, Zhu M, Benavidez NL,
Yamashita S, Abu-Jaber J, Azam S, Lo D, Foster NN, Hintiryan H, Dong H-W (2018)
Integration of gene expression and brain-wide connectivity reveals the multiscale
organization of mouse hippocampal networks *Nature Neuroscience* **21**: 1628–1643.

Chen L, Lu Y-P, Chen H-Y, Huang S-N, Guo Y-R, Zhang J-Y, Li Q-X, Luo C-Y, Lin S-W,
Chen Z-N, Hu L-H, Wang W-X, Li H-Y, Cai P, Yu C-X (2020) **Ventral tegmental area**
GABAergic neurons induce anxiety-like behaviors and promote palatable food
intake *Neuropharmacology* **173**: 108114.

De Hoz L, Knox J, Morris RGM (2003) **Longitudinal axis of the hippocampus: Both**
septal and temporal poles of the hippocampus support water maze spatial learning
depending on the training protocol *Hippocampus* **13**: 587–603.

Dolorfo CL, Amaral DG (1998) **Entorhinal cortex of the rat: Organization of intrinsic**
connections *Journal of Comparative Neurology* **398**: 49-82.

Dong H-W, Swanson LW, Chen L, Fanselow MS, Toga AW (2009) **Genomic–anatomic**
evidence for distinct functional domains in hippocampal field CA1 *Proceedings of*
the National Academy of Sciences **106**: 11794–11799.

Dupret D, O'Neill J, Pleydell-Bouverie B, Csicsvari J (2010) **The reorganization and**
reactivation of hippocampal maps predict spatial memory performance *Nature*
Neuroscience **13**: 995–1002.

Duvelle É, Grieves RM, Hok V, Poucet B, Arleo A, Jeffery K, Save E (2019) **Insensitivity**
of place cells to the value of spatial goals in a two-choice flexible navigation task *The*
Journal of Neuroscience **39**: 2522-2541.

Felix-Ortiz AC and Tye KM (2014) **Amygdala inputs to the ventral hippocampus**
bidirectionally modulate social behavior *The Journal of Neuroscience* **34**: 586-595.

Fontanini A, Grossman SE, Figueroa JA, Katz DB (2009) **Distinct subtypes of basolateral**
amygdala taste neurons reflect palatability and reward *The Journal of Neuroscience*,
29: 2486–2495.

Gasbarri A, Packard MG, Campana E, Pacitti C (1994) **Anterograde and retrograde**
tracing of projections from the ventral tegmental area to the hippocampal
formation in the rat *Brain Research Bulletin* **33**: 445–452.

Gauthier JL, Tank DW (2018) **A dedicated population for reward coding in the**
hippocampus *Neuron* **99**: 179-193.e7.

Georges-Francois P, Rolls ET, Robertson RG (1999) **Spatial view cells in the primate**
hippocampus: Allocentric view not head direction or eye position or place *Cerebral*
Cortex **9**: 197–212.

Gothard K, Skaggs W, Moore K, McNaughton B (1996) **Binding of hippocampal CA1**
neural activity to multiple reference frames in a landmark-based navigation task
The Journal of Neuroscience **16**: 823–835.

Hollup SA, Molden S, Donnett JG, Moser M-B, Moser EI (2001) **Accumulation of**

hippocampal place fields at the goal location in an annular watermaze task *The*

Journal of Neuroscience **21**: 1635–1644.

Hoover WB, Vertes RP (2007) **Anatomical analysis of afferent projections to the medial**

prefrontal cortex in the rat *Brain Structure and Function* **212**: 149–179.

Jarzebowski P, Hay YA, Grewe BF, Paulsen O (2022) **Different encoding of reward**

location in dorsal and intermediate hippocampus *Current Biology* **32**: 834-841.e5.

Jin S-W, Lee I (2021) **Differential encoding of place value between the dorsal and**

intermediate hippocampus *Current Biology* **31**: 3053-3072.e5.

Jung M, Wiener S, McNaughton B (1994) **Comparison of spatial firing characteristics of**

units in dorsal and ventral hippocampus of the rat *The Journal of Neuroscience*, **14**:

7347–7356.

Kennedy PJ, Shapiro ML (2009) **Motivational states activate distinct hippocampal**

representations to guide goal-directed behaviors *Proceedings of the National*

Academy of Sciences **106**: 10805–10810.

Kim J, Pignatelli M, Xu S, Itohara S, Tonegawa S (2016) **Antagonistic negative and**

positive neurons of the basolateral amygdala *Nature Neuroscience* **19**: 1636–1646.

Kjelstrup KB, Solstad T, Brun VH, Hafting T, Leutgeb S, Witter MP, Moser EI, Moser M-B

(2008) **Finite scale of spatial representation in the hippocampus** *Science* **321**: 140–

143.

Krettek JE, Price JL (1977) **Projections from the amygdaloid complex and adjacent**

olfactory structures to the entorhinal cortex and to the subiculum in the rat and cat

Journal of Comparative Neurology **172**: 723–752.

Lee I, Griffin AL, Zilli EA, Eichenbaum H, Hasselmo ME (2006) **Gradual translocation of**

spatial correlates of neuronal firing in the hippocampus toward prospective reward

locations *Neuron* **51**: 639–650.

Lee KJ, Park SB, Lee I (2014) **Elemental or contextual? It depends: individual difference**

in the hippocampal dependence of associative learning for a simple sensory stimulus

Frontiers in Behavioral Neuroscience **8**: 217.

Liu X, Carter AG (2018) **Ventral hippocampal inputs preferentially drive corticocortical**
neurons in the infralimbic prefrontal cortex *The Journal of Neuroscience* **38**: 7351–
7363.

Markus E, Qin Y, Leonard B, Skaggs W, McNaughton B, Barnes C (1995) **Interactions**
between location and task affect the spatial and directional firing of hippocampal
neurons *The Journal of Neuroscience* **15**: 7079–7094.

Maurer AP, VanRhoads SR, Sutherland GR, Lipa P, McNaughton B (2005) **Self-motion and**
the origin of differential spatial scaling along the septo-temporal axis of the
hippocampus *Hippocampus* **15**: 841-852.

Morris RGM, Hagan JJ, Rawlins JNP (1986) **Allocentric spatial learning by**
hippocampectomised rats: A further test of the "spatial mapping" and "working
memory" theories of hippocampal function *The Quarterly Journal of Experimental*
Psychology **38**: 365-395.

Moser MB, Moser EI, Forrest E, Andersen P, Morris RG (1995) **Spatial learning with a**
minislab in the dorsal hippocampus *Proceedings of the National Academy of Sciences*
92: 9697–9701.

O'Keefe J, Dostrovsky J (1971) **The hippocampus as a spatial map. Preliminary evidence**
from unit activity in the freely-moving rat *Brain Research* **34**: 171–175.

O'Keefe J, Nadel L (1978) *The hippocampus as a cognitive map*. Oxford: Clarendon.

Packard MG, Hirsh R, White NM (1989) **Differential effects of fornix and caudate nucleus**
lesions on two radial maze tasks: evidence for multiple memory systems *The Journal*
of Neuroscience **9**: 1465-1472.

Park EH, Ahn JR, Lee I (2017) **Interactions between stimulus and response types are**
more strongly represented in the entorhinal cortex than in its upstream regions in
rats *eLife* **6**: e32657.

Pikkarainen M, Rönkkö S, Savander V, Insausti R, Pitkänen A (1999) **Projections from the**
lateral, basal, and accessory basal nuclei of the amygdala to the hippocampal
formation in rat *Journal of Comparative Neurology* **403**: 229–260.

Ravassard P, Kees A, Willers B, Ho D, Aharoni D, Cushman J, Aghajan ZM, Mehta MR
(2013) **Multisensory control of hippocampal spatiotemporal selectivity** *Science* **340**:
1342–1346.

Rolls ET, O'Mara SM (1995) **View☐responsive neurons in the primate hippocampal**
complex *Hippocampus* **5**: 409–424.

Royer S, Sirota A, Patel J, Buzsaki G (2010) **Distinct representations and theta dynamics**
in dorsal and ventral hippocampus *The Journal of Neuroscience* **30**: 1777–1787.

Speakman A, O'Keefe J (1990) **Hippocampal complex spike cells do not change their**
place fields if the goal is moved within a cue controlled environment *European*
Journal of Neuroscience **2**: 544–555.

Swanson LW, Wyss JM, Cowan WM (1978) **An autoradiographic study of the**
organization of intrahippocampal association pathways in the rat *Journal of*
Comparative Neurology **181**: 681–715.

Tao S, Wang Y, Peng J, Zhao Y, He X, Yu X, Liu Q, Jin S, Xu F (2021) **Whole-brain**
mapping the direct inputs of dorsal and ventral CA1 projection neurons *Frontiers in*
Neural Circuits **15**: 643230.

Tye KM, Janak PH (2007) **Amygdala neurons differentially encode motivation and**
reinforcement *The Journal of Neuroscience* **27**: 3937–3945.

Valenti O, Mikus N, Klausberger T (2018) **The cognitive nuances of surprising events:**
Exposure to unexpected stimuli elicits firing variations in neurons of the dorsal
CA1 hippocampus *Brain Structure and Function* **223**: 3183–3211.

Van Groen T, Wyss JM (2003) **Connections of the retrosplenial granular b cortex in the**
rat *Journal of Comparative Neurology* **463**: 249-263.

Yoo SW and Lee I (2017) **Functional double dissociation within the entorhinal cortex for**
visual scene-dependent choice behavior *eLife* **6**: e21543.

Figure Legends

Figure 1. Place-preference task in a 2D VR environment. (A) 2D VR setup. (B) Bird's-

eye view of the virtual environment. Various landmarks surrounded a circular arena, and a

fixed start location ('St') was at the center. Reward zones are illustrated with white dashed

lines for visualization purposes. (C) Place-preference task paradigm. A trial started with one

of six pseudorandomly chosen start directions ('Trial Start'). In this example, the rat started

the trial facing the northeast (NE) direction, highlighted in green. Subsequent navigation is

illustrated here with the associated scene ('Navigation'). A dot on the gray trajectory

indicates the rat's current location and the black arrow describes the head direction. When the

rat arrived at a reward zone, honey water was delivered within 8 seconds, with the visual

scene frozen ('Reward'). Finally, a gray screen appeared, denoting an inter-trial interval; if

the rat remained still (<5 cm/s) for 5 seconds, the subsequent trial began ('ITI').

Figure 2. Common body-turning behavior of rats after learning. (A) The reference frame

of the virtual environment. The six start directions are illustrated with the red high-value zone

(180°) and blue low-value zone (0°). On the right, the departure circle (DC) is denoted with a

purple dashed line, and the start direction is marked with a black arrowhead and a green

arrow. (B) Overall changes in scene direction over the normalized distance between the start

location and the DC (left). Each colored line indicates the median change of scene direction

in trials with each start location, and red and blue arrowheads mark high- and low-value zone

centers, respectively. The 0°-to-360° range was repeated in the ordinate of the plot to capture

rotational movements in opposite directions (positive and negative directions for clockwise

and counterclockwise rotations, respectively). The gray lines on the right show the rat's

trajectory within the DC. These examples were excerpted from the first and last days of pre-

training of a single rat. The numbers after 'Novice' and 'Expert' indicate the rat and session

number of the example. (C) Individual examples of scene directions and trajectories in the

novice session. Scene direction change for each direction is drawn separately (top) for

individual trials. The black arrowhead indicates that specific start direction. Trajectories

within the DC (middle) and the whole arena (bottom) are also illustrated according to the

indicated color code. Mean travel distance in meters and latency in seconds are shown below

the VR arena trajectory. (D) Same as (C), but for the expert session.

Figure 3. Learning index for efficient navigation during pre-surgical training. (A)

Changes in departing direction (DD) with learning. (i) Schematic of DD (purple dot), with the

departure circle shown as a dashed line. (ii) Distribution of DDs in pre- and post-learning

sessions from all rats (rose plots). Gray denotes the pre-learning session, whereas purple

indicates the post-learning session. Mean vectors are illustrated as arrows with the same color

scheme, and their lengths are indicated at the upper right side of the plot. (iii) Schematic of

the DD-deviation angle (angle between the high-value zone center and the DD) and

comparisons of DD-deviation angles between pre- and post-learning sessions. Each dot

represents data from one rat. (B) Same as (A), but for perimeter-crossing direction (PCD;

green dot). The perimeter is drawn as a green dashed circle. **$p < 0.01$.

Figure 4. Cannula implantation locations and schedules for training and drug injection.

(A) Cannula positions marked. (i) Example of bilaterally implanted cannula tracks in Nissl-

stained sections in the dHP and iHP. (ii) Tip locations illustrated in the atlas, with different

colors for individual rats (n = 8). (B) Training schedule. Rats were divided into two groups

(n = 4/group) to counterbalance the injection order for the main task and probe test.

Figure 5. Changes in navigational pattern with each drug condition. (A) Sample

trajectories in each drug condition. Black arrowheads indicate the start direction and the gray

line shows the trajectory for each trial. Numbers above each trajectory indicate the

identification numbers for rat, session, and trial. (B) Mean high-value zone visit percentage

for each drug condition. Gray, green, and orange each indicate PBS, dMUS, and iMUS

sessions, respectively. (C) Average running speed. (D) Number of perimeter crossings. For

the PBS session, dPBS and iPBS sessions were first tested for significant differences between

sessions; if they were not different, they were averaged to one PBS session for analysis

purposes. *$p < 0.05$.

Figure 6. dHP and iHP inactivation differentially affect efficient goal-directed

navigation. (A) Grouped comparison of DD in each drug condition. (i) Distributions of DDs

in each drug condition (rose plots) and a comparison of their mean directions. Gray plots,

PBS sessions; green plots, dHP inactivation; orange plots, iHP inactivation. Red and blue arcs

indicate high- and low-value zones, respectively. Statistically significant differences in mean

vectors, illustrated as arrows, are indicated with asterisks. The mean directions of all four

conditions were first compared together (Watson-Williams test); a post hoc pairwise

comparison was subsequently applied if the average mean vector length of the two sessions

was greater than 0.45. The number on the upper right side of the plot shows the length of the

mean vector. (B, C) Changes in mean vector length (B) and deviation angles from the high-

value zone center (C) of the DD in each drug session. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$.

1000

1001   **Figure 7. Precision of goal-directed navigation is more severely impaired with iHP**

1002   **inactivation.** (A-C) Same as Figure 6, except showing PCD. *$p < 0.05$, **$p < 0.01$,

1003   ***$p < 0.001$.

1004

Figure 8. Goal-directedness and navigational capacity are unaffected by drug infusion.

(A) Object-guided navigation task as a probe test. (i) A flickering object appeared on either

the left ('Left trial') or right ('Right trial') side of the screen. The start location is marked

with a yellow dot, with a white arrow indicating the start direction, which remained the same

for both trial types. (ii) Example of trajectories in one session. Blue and red lines represent

trajectories from left and right trials that directly arrived at the reward zones, whereas gray

lines indicate failed trials. Green dashed lines denote reward zones. (B) Comparison of the

proportion of each drug condition's direct hit trials (both left and right).

Figure 1

A

B

Low-Value Zone 💧 x 1

High-Value Zone 💧 x 6

St

C

Trial Start

Navigation

Reward

ITI

0.5m

Six start directions

Honey water for 8s

<5cm/s for 5s

Figure 2

Figure 3

A

(i) Departure Direction (DD)

(ii) PRE 0.17   POST 0.55

(iii) 180° 164° 344°
164° = DD Deviation Angle

DD Deviation Angle (°)
PRE POST
**

B

(i) Perimeter-Crossing Direction (PCD)

(ii) PRE 0.26   POST 0.77

(iii) 255° 180° 75°
75° = PCD Deviation Angle

PCD Deviation Angle (°)
PRE POST
**

Figure 4

A

(i)

dHP
AP -3.8mm

iHP
AP -6.0mm

(ii)

- r735
- r739
- r741
- r768
- r772
- r773
- r835
- r860

B

		Surgery				Place Preference Task	Object-Guided Navigation	

Shaping | Pre-training | ▼ Recovery | Post-training | | Place Preference Task | Object-Guided Navigation | Histology

Injection   Injection

Drug Injection Schedule

Group 1 (n=4)	dPBS	dMUS	iPBS	iMUS

| Group 2 (n=4) | iPBS | iMUS | dPBS | dMUS |

Figure 5

Figure 6

A

PBS   MUS

dHP   0.67   **   0.50
        dPBS   dMUS

n.s.   DD
            p<0.001

iHP   0.59   ***   0.31
        iPBS   iMUS

B

**

n.s.   *

DD Mean Vector Length

1.0
0.8
0.6
0.4
0.2
0.0
     PBS   dMUS   iMUS

C

**

n.s.   *

DD Deviation Angle (°)

     PBS   dMUS   iMUS

Figure 7

Figure 8

A

(i)

Object-Guided Navigation Task

(ii)

- - - **Reward Zone**
— **Left trials (Direct hit)**
— **Right trials (Direct hit)**
— **Failed trials**

B