The GPI sidechain of *Toxoplasma gondii* prevents parasite pathogenesis

Julia A Alvarez[1,2], Elisabet Gas-Pascual[3], Sahil Malhi[1], Ferdinand Ngale Njume[1], Juan C Sánchez-Arcila[1], Hanke van der Wel[3], Yanlin Zhao[4], Gabriella Ceron[1], Jasmine Posada[1], Scott P Souza[1,2], George S Yap[4], Christopher M West[3], Kirk D C Jensen[1,5]
#Corresponding author: kjensen5@ucmerced.edu

[1]Department of Molecular and Cell Biology, School of Natural Sciences, University of California, Merced, California, United States of America
[2]Quantitative and Systems Biology Graduate Program, University of California, Merced, California, United States of America
[3]Department of Biochemistry and Molecular Biology, Center for Tropical and Emerging Global Diseases, and Complex Carbohydrate Research Center, University of Georgia, Athens, Georgia, United States of America
[4]Department of Medicine and Center for Immunity and Inflammation, New Jersey Medical School, Rutgers University, Newark, New Jersey, United States of America
[5]Health Sciences Research Institute, University of California, Merced, California, United States of America

RUNNING TITLE
Role of the GPI sidechain in *T. gondii* pathogenesis

ABBREVIATIONS
GPI, Glycosylphosphatidylinositol; MFI, mean fluorescence intensity

ABSTRACT

Glycosylphosphatidylinositols (GPIs) are highly conserved anchors for eukaryotic cell surface proteins. The apicomplexan parasite, *Toxoplasma gondii*, is a widespread intracellular parasite of warm-blooded animals whose plasma membrane is covered with GPI-anchored proteins, and free GPIs called GIPLs. While the glycan portion is conserved, species differ in sidechains added to the triple mannose core. The functional significance of the Glcα1,4GalNAcβ1- sidechain reported in *Toxoplasma gondii* has remained largely unknown without an understanding of its biosynthesis. Here we identify and disrupt two glycosyltransferase genes and confirm their respective roles by serology and mass spectrometry. Parasites lacking the sidechain on account of deletion of the first glycosyltransferase, PIGJ, exhibit increased virulence during primary and secondary infections, suggesting it is an important pathogenesis factor. Cytokine responses, antibody recognition of GPI-anchored SAGs, and complement binding to PIGJ mutants are intact. In contrast, the scavenger receptor CD36 shows enhanced binding to PIGJ mutants, potentially explaining a subtle tropism for macrophages detected early in infection. Galectin-3, which bind GIPLs, exhibits a slight enhancement of binding to PIGJ mutants, and the protection of galectin-3 knockout mice from lethality suggests that *Δpigj* parasite virulence in this context is sidechain dependent. Parasite numbers are not affected by *Δpigj* early in the infection in wildtype mice, suggesting a breakdown of tolerance. However, increased tissue cysts in the brains of mice infected

47 with *Δpigj* parasites indicate an advantage over wildtype strains. Thus, the GPI
48 sidechain of *T. gondii* plays a crucial and diverse role in regulating disease outcome in
49 the infected host.
50
Summary

The functional significance of sidechain modifications to the GPI anchor is yet to be
determined because the glycosyltransferases responsible for these modifications have
not been identified. Here we present identification and characterization of both *T. gondii*
GPI sidechain-modifying glycosyltransferases. Removal of the glycosyltransferase that
adds the first GalNAc to the sidechain results in parasites without a sidechain on the
GPI, and increased parasite virulence. Loss of the second glycosyltransferase results in
a sidechain with GalNAc alone, and no glucose added, and has negligible effect on
parasite virulence. This indicates GPI sidechains as fundamental to host-parasite
interactions.

61
Introduction

Protozoan parasites are widespread and cause prominent diseases including
malaria, leishmaniasis, chagas, and toxoplasmosis. One notable feature of protozoans
is their extensive decoration of glycosylinositolphospholipids (GIPL) and GPI-anchored
proteins (GPI-AP). The GPI was first discovered and characterized in trypanosomes,
the protozoan parasite that causes African sleeping sickness in humans (1). Since its
discovery, it has been shown to have a conserved core structure across the eukaryotic
kingdom; EtNP-6Manα1-2Manα1-6Manα1-4GlcN1-6*myo*-inositol-phospholipid (where
EtNP, Man, and GlcN are ethanolamine phosphate, mannose, and glucosamine,
respectively) (1,2). Apicomplexan parasites use GPI-APs to assist attachment to and
cellular invasion of host cells (3) and as such, the core GPI synthetic pathway enzymes
are each essential gene products required for *Toxoplasma gondii* (*T. gondii)* and
Plasmodium sp. survival and intracellular infection (4,5).

In the case of *T. gondii*, a widespread apicomplexan parasite of warm-blooded
animals and humans, cellular attachment and invasion are mediated by a large family of
T. gondii GPI-APs called surface antigen glycoprotein (SAG)-related super family (SRS)
(6,7). It is known that these dominant SAG antigens are targeted by *T. gondii*-specific
antibodies following infection (8), but also play important roles in virulence. For
example, SAG1, the dominant antigen in the parasite's lytic stage has been shown to
promote small intestinal ileitis in mice (9), and promotes parasite survival *in vivo* (10)
and in activated macrophages, by an unknown mechanism (11). Another GPI-AP of *T.
gondii*, SRS35 (also known as p18 or SAG4), has been shown to promote mouse
macrophage invasion and virulence of the parasite (12). In contrast, overexpression of
the GPI-AP SRS29C (p35) quells *T. gondii* virulence and promotes mouse survival to an
otherwise lethal infection (6).

While many GPI-APs have been considered for their role in parasite virulence,
the GPI that anchors them has long been shown to be immunogenic in *T. gondii* and
other parasites. For instance, *T. cruzi* GIPLs are potent activators of TLR2 (13).
Similarly, GIPLs of *P. falciparum* are thought to be major pathogenesis factors, and are
shown to activate TLR2 and TLR4 (14), which in turn causes lethal inflammatory
responses (15). The GIPL of *T. gondii* activates TLR2 and TLR4 (16). *T. gondii* and

Plasmodium sp. GIPLs are also robustly targeted by antibodies during the early stages
of infection in both humans and in mice (17–25). Collectively, these data demonstrate
that parasitic GPIs illicit robust innate and adaptive immune responses during infection.
Whereas the core GPI structure is conserved across the eukaryotic kingdom,
species differ in how their GPI is modified through the attachment of additional sugars
and other moieties to the mannose core, called sidechain modifications (26–28) (**Fig
S1**). The significance of these sidechains and why species possess structural diversity
in their GPI modifications is unclear because the responsible GPI glycosyltransferases
(GT) have remained unidentified in most species. However in humans, the
glycosyltransferase PGAP4 was recently determined to be the GT responsible for
adding D-GalNAc to the first mannose of the human GPI, the first subunit of the
sidechain (29). PGAP4 is widely conserved among eukaryotic species that are similarly
substituted, including *Caenorhabditis elegans*. In the case of *T. gondii,* the GPI
sidechain occurs on the same first mannose in two glycoforms: one consisting of a
single identically linked D-GalNAc, and the other in which the GalNAc is extended by an
α4-linked D-Glc (17). Interestingly, however, PGAP4 is not conserved in *T. gondii*
despite sharing the same GalNAc branch at the first mannose (**Fig S1**) (29). When the
PGAP4 homolog was deleted in mice the deficiency resulted in elevated blood alkaline
phosphatase levels, impaired bone formation, decreased locomotion, impaired memory,
and more enhanced vulnerability to prion diseases (30). These results point to GPI
sidechains having far-reaching and important roles in mammals.
In the context of eukaryotic microbes, even less is known regarding the functional
significance of the GPI sidechain. For example, in *Plasmodium* and fungi there is a
fourth mannose added as a sidechain to the third mannose of the core GPI by the
SMP3 glycosyltransferase (**Fig S1**). Removal of *SMP3* in *Candida albicans* interferes
with fungal viability *in vitro* (31). In *P. falciparum* which has a fourth mannose GPI
sidechain, but lacks a homolog to *SMP3*, its GT *PIGB* that is responsible for adding the
third mannose to the GPI core can add the fourth mannose when complemented into *C.
albican Δsmp3* mutants. Whether *P. falciparum PIGB* encodes the GT responsible for
adding the mannose GPI sidechain is uncertain because *Δpigb* parasites fail to grow *in
vitro* (32). Multiple GPI sidechain GT's have been identified in *Trypanosoma brucei* (*T.
brucei*) parasites which exhibit multiple elaborately branched sidechains emanating from
the second mannose of its GPI core. The *T. brucei* GT *TbGT8*, which adds the first
branch point of one of the sidechain branches, was shown to be nonessential in
bloodstream- and procyclic-form stages of the parasite (33). *TbGT3* is expressed in the
bloodstream and procyclic stages of *T. brucei*, and mutants remained infectious to
tsetse flies, though the Gal addition to the non-reducing-terminal GlcNAc residue of the
GPI sidechain was lost (34). Loss of *TbGT10* resulted in a minor fitness cost evidenced
by reduced growth *in vitro*, likely a consequence of impaired carbohydrate synthesis,
but *TbGT10* was responsible for adding all β6 linkages to the GPI sidechain, resulting in
10 less sugars in *TbGT10* null parasites (35). Although removal of the aforementioned
T. brucei GT's reduces some of the GPI sidechain branching complexity, none of these
GT mutants lack a GPI sidechain, suggesting functional redundancy may exist between
different GPI glycoforms to support *T. brucei* fitness. Whether expression of specific
glycoforms of the GPI are required for *in vivo* fitness of eukaryotic microbes is largely
unknown.

While there have been several eukaryotic GPI sidechain modifying
glycosyltransferases identified to date, none have addressed whether lacking a GPI
sidechain entirely or whether specific glycoforms are required for microbial
pathogenicity. Therefore, we sought to address these questions in *T. gondii* and test
whether specific GPI glycoforms impact parasite virulence and immune recognition.
Here we identify the two sidechain-modifying glycosyltransferases in *T. gondii* and
report that loss of the GPI sidechain promotes virulence. Whereas antibody recognition
of parasite GPI-APs and inflammatory cytokines appear intact to GPI sidechain null
parasites, we present indirect evidence for galectin-3 and the scavenger receptor,
CD36, in mediating certain phenotypes associated with sidechain deficiency in *T. gondii*
strains.
150
MATERIALS & METHODS
Parasite strains and cell lines
Human foreskin fibroblasts (HFFs) monolayers were grown in DMEM GlutaMAX^{TM} (4.5
g/L D-glucose) (Life Technologies #10-566-024) supplemented with 2 mM L-glutamine,
20% heat inactivated (HI) fetal bovine serum (FBS) (Omega Scientific), 1% penicillin-
streptomycin (Life Technologies, #15140122), and 0.2% gentamycin (Life Technologies
#15-710-072). Mouse Embryonic Fibroblasts (MEFs) were grown in DMEM GlutaMAX^{TM}
(4.5 g/L D-glucose) supplemented with 10% HI FBS, 20mM HEPES, 1% penicillin-
streptomycin, and 0.2% gentamycin. *Toxoplasma gondii* strains were passaged in HFFs
in "Toxo medium" (4.5 g/L D-glucose, L-glutamine in DMEM GlutaMAX^{TM} supplemented
with 1% HI FBS and 1% penicillin-streptomycin). The following clonal strains were used
(clonal types are indicated in parentheses): RH *Δku80 Δhxgprt* (type I), RH *GFP:cLUC*
(type I) (clone 1-1), RH *Δku80 Δhxgprt Δpigj::HXGPRT* (type I) (clone C12), RH *Δku80*
Δhxgprt Δpige::DHFR-TS (type I) (36), GT1 (type I), GT1 *Δpigj:DHFR-TS* (type I) (clone
B8), GT1 *Δpige::DHFR-TS* (type I) (clone C11), CEP *hxgprt-* (type III), CEP *hxgprt-*
GFP:cLUC (type III), CEP *hxgprt- Δpigj:hxgprt GFP:cLUC* (type III) (clone D10), CEP
hxgprt- Δpigj::HXGPRT PIGJ-HA$_{3x}$:DHFR-TS GFP:cLUC (type III) (clone C2), CEP
hxgprt- Δpige::hxgprt GFP:cLUC (type III) (clone D4). All strains, oligos and plasmids
used for the generation of strains in this study can be found in **Table S1**.
Generation of gene edited *T. gondii* strains
Cas-9 and single guide RNA (gRNA) expression plasmid, pSS013 (gift from Jeroen
Saeij, University of California, Davis) was designed to target *Tg_207750* (Uniport
S8FC68)*, PIGJ*. For *Tg_266320* (Uniprot S7VW57)*, PIGE* a modified dual-guide pU6-
universal plasmid targeting two different exons within the gene was previously
generated (36). Selectable markers were produced by PCR amplification. An amplicon
of the pyrimethamine selectable cassette (*DHFR-TS)* was amplified from the pLoxp-
DHFR-TS-mCherry plasmid (Addgene plasmid #70147) containing similar homology
arms to the targeted protospacer sequences*. The amplicon of *HXGPRT* was generated
from the plasmid pTKO-att (a gift from Jeroen Saeij, University of California, Davis) with
homology arms flanking the targeted protospacer sequences of the targeted gene. The
expression Cas9/gRNA plasmid and amplicon were co-transfected via electroporation at
a 5:1 ratio. Transfectants were selected and cloned in medium containing
pyrimethamine (1 µM) or 50 µg/mL of MPA (Axxora) and 50 µg/mL of xanthine (Alfa
Aesar) and screened for the disruption the locus by diagnostic PCR.

Generation of the *PIGJ* complementation construct and complementation strain

The pLIC-HA-DHFR-TS (gift from Jeroen Saeij, University of California, Davis) plasmid was treated simultaneously with PacI (NEB #R0547S) and AvrII (NEB #R0174S) restriction endonucleases to allow for insertion of *TgVEG_PIGJ* by homology directed ligation. The full-length coding region with introns spliced out of *PIGJ* was amplified using the Q5 high fidelity polymerase (NEB, M0491) from a CEP cDNA library preparation using primers designed to contain 19 bp homology with the 3' end of the *PIGJ* promoter sequence and 25 bp homology with end of the digested pLIC-HA plasmid that contains and provides a 3x HA tag, respectively. The CEP cDNA library was prepared from CEP total RNA preparation using the High-Capacity cDNA Reverse Transcription Kit (ThermoFisher, cat# 4368814) according to manufacturer's instructions. For amplification of the *PIGJ* promoter region, a 1000 bp non-coding region upstream of the start ATG of *PIGJ* was amplified from CEP genomic DNA with Q5 high fidelity polymerase according to manufacturer's protocol. The forward primer contained a 19 bp homology sequence to the end of the digested pLIC-HA plasmid and the reverse shared a 22 bp homology sequence with the 5' end of the amplified PIGJ coding-cDNA sequence described above. The amplified promoter and coding *PIGJ* DNA sequence were first purified from 1% agarose gels using the ZymocleanTM Gel DNA recovery kit (Zymo Research, #D4007) and assembled in frame into the digested pLIC-HA -DHFR-TS plasmid in frame with the 3x HA tag using the NEBuilder HiFi DNA Assembly cloning Kit (NEB, cat# E5520) according to the manufacturer's instructions. The CEP *hxgprt- Δpigj::HXGPRT GFP:cLUC* strain was transfected with the linearized pLIC-HA-DHFR-TS PIGJ-HA3x plasmid, grown in pyrimethamine selection medium and cloned by limiting dilution to generate the *CEP Δpigj::PIGJ-HA$_{3x}$* complementation strain.

The HA tag was visualized by fluorescent microscopy. 8×10^5 HFFs were plated on coverslips in 24-well plates overnight before being infected with 8×10^4 parasites overnight. Cells were fixed with 3% formaldehyde and permeabilized with blocking buffer (1X PBS with 10% normal goat serum, and 0.01% saponin) and stained with rat anti-HA (1:50) (3F10, Sigma). Secondary goat anti-rat AF594 (Thermo Fisher cat#A11007) antibodies were used at (1:3000), and DAPI (1:10000), for detection and visualized using a fluorescent microscope (Nikon Eclipse T*i*-5).

Mice

C57BL/6J (B6), A/J, and *Lgals3-/- (*B6.Cg-*Lgals3^{tm1Poi}/J)* mice were purchased from Jackson Laboratories. *Tlr2/4* double knockout B6 mice (B6.129P2-*Tlr4^{tm1Aki}* B6.129P2-*Tlr2^{tm1Aki}*) were a generous gift from Dr. Greg Barton (UC Berkeley). Mice were maintained under specific pathogen free conditions at UC Merced. Female mice of 6-10 weeks of age were used for experiments unless otherwise stated.

Ethics statement

Mouse work was performed in accordance with the National Institutes of Health Guide to the Care and Use of Laboratory Animals. Mouse protocols used have been reviewed and approved by UC Merced's Committee on Institutional Animal Care and Use Committee (IACUC, #AUP-20-0015). UC Merced has an Animal Welfare Assurance filed with OLAW (A4561-01), is registered with USDA (93-R-0518), and the UC Merced Animal Care Program is AAALAC accredited (001318).

Parasite infections and serotyping

Parasite injections were prepared by scraping T-25 flasks containing vacuolated HFFs
and sequential syringe lysis first through a 25G needle followed by a 27G needle. The
parasites were spun at 34 x g for 5 minutes to remove debris and the supernatant was
transferred, followed by a spin at 611 x g and washing with sterile 1X PBS (Life
Technologies, #10-010-049). For primary infections, mice were infected intraperitoneally
(i.p.) with 104 tachyzoites of type III CEP *hxgprt-* parental and mutant strains. Parasite
viability of the inoculum was determined by a plaque assay. In brief, 100 or 300
tachyzoites were plated in HFF monolayers grown in a 24-well plate and 4-6 days later
were counted by microscopy (4x objective) (Nikon Eclipse Ti-5).
At 30 to 35 days after primary infection, 50µl of blood was harvested and
collected in Eppendorf tubes containing 5µL 0.5 M EDTA and placed on ice. Blood was
pelleted at 9391 x g for 5 minutes, and blood plasma was collected from the
supernatant and stored at -80°C. To evaluate the seropositivity of the mice, HFFs were
grown on coverslips and infected with green fluorescent protein (GFP)-expressing RH
(1-1) overnight. 18 hrs later, cells were fixed with 3% formaldehyde in PBS,
permeabilized with a permeabilization solution (3% bovine serum albumin fraction V
(Omega, FB-11), 2% normal goat serum (Omega, NG-11) 0.2 M Triton X-100, 0.01%
sodium azide), incubated with a 1:100 dilution of collected blood plasma for 2 hrs at
room temperature, washed with 1X PBS, and detected with Alexa Fluor 594-labeled
goat secondary antibodies specific for mouse IgG (Thermo Fisher, cat#A11032) in
permeabilization solution. Seropositive parasites were observed by
immunofluorescence microscopy.
For secondary infections, seropositive mice were infected i.p. with 5×10^4
tachyzoites of type I RH *Δku80* parental and mutant strains. Parasite viability of the
inoculum was determined by plaque assay described above.

Cyst enumeration using Dolichos-FITC

To quantify brain cysts, brains were dissected and placed in 10mLs of 1X PBS on ice.
Brains were homogenized using a 10mL syringe with a 18G needle by extrusion several
times through the needle. The homogenate was spun at 611 x g for 7 minutes and then
resuspended in 1mL of 1X PBS. 100µL of the brain homogenate was fixed in ice cold
methanol at a 1:10 dilution for 5 minutes, spun in a microcentrifuge at 5200 x g for 5
minutes and washed once and resuspended in 1mL 1X PBS. The fixed homogenate
was stained with a 1:150 dilution of FITC-conjugated Dolichos biflorus agglutinin (Vector
Laboratories cat# FL-1031) in 1X PBS and slowly rotated at 4°C overnight. Samples
were washed twice with 1X PBS and resuspended in 1mL of 1X PBS. 50µL of stained
homogenate was plated in a 96 well plate and four replicates of each sample were
enumerated for cysts with an inverted fluorescence microscope using a 20x objective
(Nikon Eclipse T*i*-5).

Superinfections

Mice that survived 35 days of secondary infection were euthanized and brain
homogenates were generated and resuspended in 1mL of 1X PBS as described above.
100µL of the homeogenate was cultured in HFFs and monitored for parasite growth.
Once parasite growth was established, the media was switched to selection media
containing MPA/xanthine for secondary infections to select against the primary infecting
CEP *hxgprt-* strain, and for *HXGPRT* expressing secondary infection strain. Parasites
that grew in selection media were also confirmed to be type I strains through restriction

fragment length polymorphism analysis. In brief, isolated DNA from MPA/xanthine selected parasites was PCR amplified using GRA6 primers and the purified amplicon was fragmented with MseI (NEB cat# R0525M). Products were run on a gel to determine parasite type I or III based on unique fragmentation sizes indictive of the genotype.

SDS-PAGE and immunoblotting for parasite lysate antigen

To generate parasite lysate antigens *T. gondii* was cultured in HFF monolayers and expanded to approximately 2×10^8 parasites in a T175 flask. Parasites were syringe-lysed, washed with sterile 1X PBS, and pelleted at 611 x g for 7 minutes. The pelleted parasites were lysed with Lameli Buffer (0.0625M Tris Base, 0.07M SDS, 10% glycerol, 5% β-mercaptoethanol) and centrifuged at 14,000 x g for 20 minutes to remove large insoluble debris. The supernatant was aliquoted and stored at -80°C. Parasite lysates were separated via SDS-PAGE in 4-20% acrylamide hand cast gels before transfer to PVDF membrane. Membranes were blocked with 10% fortified bovine milk (Raley's) dissolved in Tris-Buffered Saline with 0.1% Tween (TBS-T 0.1%) for 1-2 hrs at room temperature or overnight at 4°C. Blots were then probed with *T. gondii* GPI anchor glycoform-specific antibody clones T3 3F12 (BEI) which binds GalNAc GIPL, or T5 4E10 (a gift from Jean Francois Dubremetz U. Montpelier) which binds GalNAc + Glc GIPL in blocking buffer for 72 hrs at 4°C. Membranes were washed with TBS-T 0.1% three times and incubated for one hr at room temperature with goat anti-mouse Ig horseradish peroxidase (HRP)-conjugated antibodies (Southern Biotech, anti-IgM secondary 1:1000 (cat# 1020-05), anti-IgG 1:7500 (cat# 1030-05)). Membranes were then washed with TBS-T 0.1% three times and developed with Immobilon Forte Western HRP Substrate (Millipore, WBLUF0500). All blots were imaged via chemiluminescence on a ChemiDoc Touch (cat#12003153, Bio-Rad). Image Lab 6.1 software (Bio-Rad) was used for analysis of bands.

NanoLC-MS and MALDI-TOF-MS analysis of GIPLs and GPI-anchored protein glycans released from parasites

Parasite pellets corresponding to approximately 3×10^8 parasites were harvested as described in 2.2.6 but from 15 T175 flasks and thoroughly rinsed in ice-cold PBS (Corning cat # 21-040 CV), before being pelleted and stored at 80°C. Control, uninfected HFF monolayers were harvested in 1 mM EDTA in PBS and washed 3 times in ice-cold PBS by centrifugation (8 minutes, 2000 x *g*, 4°C). Parasite and host cell pellets were then delipidated, and the protein pellet was extracted with butan-1-ol saturated water to enrich for GPI-anchored proteins, as described (36). GIPL fractions were obtained from the clarified lipid extracts generated during initial pellet delipidation. Fatty acids (FA) were removed from both GPI-anchored proteins and GIPLs by incubation in 0.5 M NH_4OH (in water for GPI-anchored proteins or after bath sonication in 70% ethanol for GIPLs) under rotation for 6 hrs at 4°C. The glycan core was then released by hydrofluoric acid treatment, C18 cleaned up, re-N-acetylated, permethylated, and dried as described (36).

For nLC-MS, permethylated glycans were redissolved in MeOH and 10μl of were mixed with 90μl 0.1% FA in water. 10μl was injected into a PepMap Acclaim analytical C18 (75 μm, 15 cm, 2 μm pore size) column maintained at 60°C in an Ultimate 3000 RSLC coupled to a Q-Exactive-Plus mass spectrometer (Thermo Fisher Scientific). Column was equilibrated for 10 minutes at 97.5% LC-MS Buffer A (0.1% FA in water)

and ramped up to 35% LC-MS Buffer B (80% (v/v) over 2 minutes. Glycan separation was achieved using a linear gradient from 35% to 70% Buffer B over 150 minutes at a flow rate of 300 nl/minutes. The effluent was introduced into the mass spectrometer by nanospray ionization in positive mode via a stainless-steel emitter with spray voltage set to 1.9 kV and capillary temperature set at 275°C. The MS method consisted of a survey Full MS scan at 70,000 resolution in positive ion mode, followed by MS(2) fragmentation of the top 10 most intense peaks using HCD at 40% collision energy and an isolation window of 2 m/z. Dynamic exclusion was set to exclude ions for fragmentation for 30 sec. All the data were processed manually using the Xcalibur 2.0 software package.

For MALDI-TOF-MS, permethylated samples were redissolved in MeOH and 0.5µl aliquots were mixed 0.5µl MeOH saturated with either 2,5-dihydroxybenzoic acid (DHB) or α-cyano-4-hydroxycinnamic acid (CHCA) on the MALDI plate and allowed to air dry. Samples were then analyzed on an ABI 4700 MALDI-TOF-MS operated in positive ion reflectron mode for m/z range 500-5000.

Flow cytometry of peritoneal exudate cells after infection

Mice were infected i.p. with 10^6 tachyzoites, and 3 hrs post injection mice were euthanized and peritoneal cavity lavaged with 4mL of 1X PBS and 3 mL of air. The peritoneal exudate fluid was passed through a 70 µm cell strainer, cells were pelleted for staining. Samples were blocked for 30 minutes in FACS buffer (2% FBS 1X PBS) containing Fc Block anti-CD16/32 (2.4G2, BD Biosciences) (1:100 dilution), 5% normal hamster serum, and 5% normal rat serum (Jackson ImmunoResearch). After blocking, cells were stained at a 1:100 dilution for 30 minutes with the following antibodies: CD11b BUV395 (M1/70, BD), Gr.1 PE (RB6-8C5, BioLegend), F4.80 BV421 (BM8, Thermo Fisher), MHC II APC (AF6-120.2, Thermo Fisher), B220 APC-Cy7 (RA3-6B2, BioLegend), CD19 BUV785 (6D5, BioLegend), NK 1.1 PE Cy7 (PK136, Thermo Fisher), CD3ε BV510 (17A2, BD) and Propidium Iodine (PI) (1:1000). After incubation, cells were washed three times in FACS buffer and resuspended in FACS buffer. Samples were run on a ZE5 flow cytometer (Bio Rad) and analyzed using FlowJo software version 10.

Flow cytometry for parasite antigen expression

Syringe lysed parasites were plated in a micro-titer plate at $8x10^5$ parasites per well and incubated with one of the following antibodies at (1:100) to detect antigen expression: mouse anti-SAG1 (T4 IE5, BEI), mouse anti-SAG3 (T4 1F12, BEI), mouse anti-p35 (T4 3F12, BEI). After 20 minutes incubation on ice, parasites were washed twice with FACS buffer and stained with (1:100) goat anti-mouse IgG-APC (BioLegend, cat# Poly4053). After incubation, cells were washed and resuspended in FACS buffer as described above. Samples were run on a ZE5 flow cytometer and analyzed using FlowJo software version 10.

Serum antibody parasite binding assay

For serum reactivity analysis, syringe-lysed GFP-expressing strains were fixed in 3% formaldehyde for 20 minutes, washed twice in PBS, and plated in 96-well micro-titer plates at $4x10^5$ parasites/well. The parasites were then incubated with serum from chronically infected mice, in serum concentrations ranging from 10^{-2} to 10^{-6} diluted in FACS buffer, for 20 minutes at 37°C. Parasites were then washed with FACS buffer and placed on ice for incubation with anti-isotype detection antibodies depending on application: anti-IgG3-BV421 (R40-82, BD Bioscience), anti-IgM-PE/Cy7 (RMM-1,

BioLegend), anti-IgG1-APC (RMG1-1, BioLegend), anti-IgG2b-PE (RMG2b-1,
BioLegend), anti-IgG2a-PerCP/Cy5.5 (RMG2a-62, BioLegend). Parasites were washed
and resuspended in FACS buffer as described before. Flow cytometry was performed
on an LSR analyzer (BD) and analyzed using FlowJo software version 10. The MFI was
computed by gating on forward and side scatter characteristics consistent with parasites
and further gated for GFP+ if the parasite expressed GFP using FlowJo.

Complement C3b binding assay
For analysis of C3b binding, parasites were syringe lysed, washed, and resuspended to
a concertation of 1×10^7 parasites per mL in Hank's Balancing Salt Solution Buffer
(HBSS) (ThermoFisher, cat#14175095). Parasites were plated in a 96-well plate at 10^6
parasites per well with 10% blood plasma from naive mice and incubated at 37°C for 30
minutes. After incubation, complement activation was stopped by washing with cold 1X
PBS. Parasites were spun down at 611 x g for 3 minutes and washed 3 times using cold
1X PBS. Parasites were then fixed using 3% formaldehyde for 10 minutes and then
washed using cold 1X PBS. Parasites were then stained with mouse anti-C3b (10C7)
(Invitrogen MA1-70054) (1:200) for 20 minutes. After primary stain, parasites were
washed with cold 1X PBS and stained with anti-mouse IgG1 APC (RMG1-1, BioLegend)
(1:100) for 20 minutes on ice. After incubation, parasites were washed 3 times and
resuspended in 1X PBS for analysis. Samples were run on a ZE5 machine and data
analyzed using FlowJo version 10.3. MFI was determined as described above.

Galectin-3 binding assay
Recombinant Gal-3 (R&D Systems, Cat# 1197-GA-050) was pre-diluted in phosphate-
buffered saline (PBS) containing 10% bovine serum albumin (BSA) and 14 mM β-
mercaptoethanol. Parasites were syringe lysed, washed, and 8×10^6 parasites were
incubated with either 10 or 5 µg of recombinant Gal-3 for one hour at 4°C in FACS
buffer. Parasites were washed to remove unbound Gal-3 and incubated with a PE-
conjugated anti-Gal-3 antibody (eBioM3/38 (M3/38), Invitrogen) at a 1:100 dilution in
FACS buffer for 20 minutes at 4°C. After two further washes, the samples were
analyzed using a ZE5 Cell Analyzer (Bio-Rad) and data processed as described above.

CD36-Fc binding assay
10^7 parasites were incubated with 0.5 µg recombinant human IgG Fc (R&D Systems,
cat# 110-HG) or recombinant CD36-Fc (R&D Systems, #2519-CD) in 50µl binding
buffer (0.14 M NaCl, 2.5mM $CaCl_2$, 0.01M HEPES (pH7.4), 3% BSA) for 1 hr at 15°C.
Parasites were washed once to remove unbound recombinant protein and the parasites
were then incubated with mouse anti-SAG1-AF405 (TP3cc, Novus Biologicals) and
donkey anti-human IgG-Daylight 550 (Thermo Fisher, SA5-10127). The stained
parasites were acquired by Attune NxT flow cytometry. The CD36-Fc MFI was
computed by gating on GFP+/SAG1+ parasites using FlowJo.

Opsonization and invasion assays
Bone marrow-derived macrophages (BMDMs) were generated in L929 conditioned
medium from female C57BL/6J mice as previously described (37). BMDMs were plated
a day in advance on a coverslip lined 24-well plate at 8 $\times10^5$ cells in 1mL per well, in
"BMDM media" supplemented with 10% L929 (20% HI FBS, 1% penicillin-streptomycin,
1% non-essential amino acids (ThermoFisher, cat#11-140-050), 1% sodium pyruvate).
The day of the experiment, freshly lysed parasites were washed and resuspended for
an MOI of 0.5 in Toxo media warmed to 37°C. In a 96-well plate, serum dilutions were

made in Toxo media that was pre-warmed to 37°C. Parasites were added to the diluted
serum and incubated at 37°C for 20 minutes before being added to the BMDM coverslip
wells with fresh Toxo medium. The BMDM plates were spun at 304 x g for 5 minutes to
synchronize parasite invasion between wells and incubated at 37°C for 40 minutes to
allow for phagocytosis or invasion. After 40 minutes, media was replaced with 300µl of
3% formaldehyde in 1X PBS, fixed for 20 minutes, washed three times with 1X PBS,
blocked and permeabilized with "Blocking Buffer" for 1 hr (1X PBS with 10% normal
goat serum, and 0.01% saponin). Cells were stained with rabbit anti-GRA7 (1:5000) (gift
from John Boothroyd, Stanford) and rat anti-LAMP1 (1D4B, BD Biosciences) (1:500)
diluted in blocking buffer for 1 hr or overnight at 4°C. After primary antibody incubation,
wells were washed three times with 1X PBS before secondary antibody incubation.
Secondary antibodies were goat anti-rabbit AF647 (1:3000) (Thermo Fisher,
cat#A21245) goat anti-rat AF594 (1:1000) (Thermo Fisher cat#A11007), and DAPI
(1:10000), and if parasites were not GFP-expressing, goat anti-Toxo conjugated FITC
(1:400) (ViroStat, cat#0283) was also used. Fluorescence microscopy was performed
(Nikon Eclipse Ti-5) and images were captured at 60x and blinded for analysis. Images
were processed (Nikon Elements) and 80-100 GFP+ events were quantified for parasite
association with Lamp1+ or for containment within a ring-like GRA7+ parasitophorous
vacuole (PV); fraction opsonization was calculated as (Lamp1+GFP+ counts /
Lamp1+GFP+ plus GRA7 PV+GFP+ total counts).
For invasion assays BMDMs were plated on coverslips in 24-well plates
overnight as described above and infected the next day at an MOI of 0.25 parasites.
Plates were spun at 304 x g to synchronize invasion and incubated at 37°C for 20
minutes before washing once in 1X PBS and fixing in 3% formaldehyde. Cells were not
permeabilized. Wells were washed and stained with mouse anti-SAG1 (1:5000) for 1 hr.
Cells were washed, and secondary antibodies goat anti-mouse AF594 (1:3000) and
DAPI (1:1000) were used for 1 hr. Fluorescence microscopy was performed as
described above, with 100 GFP+ parasites counted per slide and quantified for parasite
for SAG1+ staining. Invaded parasites were determined to be GFP+ SAG1-. Fraction of
invasion was calculated as (SAG1-GFP+ counts / total GFP+ counts).

Neutralization assay

MEFs were plated a day in advance in a 96-well plate at $5x10^5$ cells per well in MEF
media. The day of the experiment, freshly lysed parasites were washed and
resuspended to a concentration of $1.6x10^7$ in in Toxo media warmed to 37°C. In a 96
well plate, blood plasma 1:10 dilution was made in Toxo media pre-warmed to 37°C and
incubated at 37°C in 10% serum for 20 minutes before being added to the MEFs, such
that in the absence of serum the MOI is approximately 0.5. The plate was spun at 304 x
g for 3 minutes to synchronize parasite invasion and incubated at 37°C for 2 hrs.
Supernatant was carefully removed, and 20µl of trypsin was added to each well and
incubated at 37°C for 5 minutes to dislodge cells from the plate. Cells were harvested
with 100µl of cold 1X PBS, transferred to a FACS plate and washed 3 times with FACS
buffer to remove trypsin. Cells were resuspended in 1:1000 PI in FACS buffer, and then
analyzed via flow cytometry for GFP+ PI- MEFs to indicate invasion. Neutralization was
defined as a ratio: the percentage of PI- cells infected with parasites incubated with
serum divided by the percentage of PI- cells infected with parasites without serum
incubation. For non-GFP+ expressing parasites, cells were fixed after trypsinization in

3% formaldehyde, washed, and blocked for 1 hr before staining with anti-Toxo-FITC
(ViroStat, cat#0283). Neutralization ratios were determined as described, but PI was not
included in the discrimination. Samples were run on a ZE5 machine and data analyzed
using FlowJo software version 10.

Parasite lytic cycle
For plaque sizes, HFFs were plated in 24-well plates and allowed to grow to confluency
before adding 100 or 300 syringe-lysed parasites and allowed to grow for 5 days.
Plaque sizes were measured (Nikon Elements) under 4x objective (Nikon Eclipse T*i*-5).

For parasites per vacuole assays, 8×10^5 MEFs were plated on coverslips in 24-
well plates and allowed to adhere before infecting with 10^5 parasites. Plates were spun
at 475 x g to synchronize invasion and incubated for 15 minutes at 37°C. After 15
minutes cells were washed twice to remove any unattached parasites. Fresh Toxo
media was added, and plates incubated at 37°C for 16 hours before being fixed,
blocked with 0.1% saponin blocking buffer (as described above), and stained for 1 hr
with rabbit anti-GRA7 (1:3000). Cells were washed and incubated with secondary goat
anti-rabbit AF 594 (1:3000) (Thermo Fisher, cat#A11037) and DAPI (1:10,000) for 1 hr.
100 GFP+ vacuoles per slide were quantified for number of parasites residing in GRA7+
vacuoles under 100X objective (Nikon Eclipse T*i*-5).

For attachment time course assays, HFFs were plated in 24-well plates and
allowed to grow to confluency. 200 syringe lysed parasites were added per well and
plates were spun at 475 x g to synchronize invasion and incubated for 30, 60, or 90
minutes before washing 5 times in 1X PBS to remove any unattached parasites. Fresh
Toxo media was added and plates were incubated at 37°C for 5 days before counting
plaques under the microscope. Plaque numbers were normalized to those obtained
from wells that did not undergo any washes.

Quantitative PCR for cytokine expression and parasite burden
Cells were collected from mouse peritoneal lavages as described above and spleens.
Cells were filtered through a 70 µm filter and centrifuged 611 x g for 7 minutes to pellet
and washed with sterile 1X PBS before pelleting again. Samples were removed of red
blood cells through treatment of ACK lysis buffer (0.5 M NH_4Cl, 0.01 M $KHCO_3$, .0001 M
EDTA) for 5 minutes before being washed with 1X PBS. Cells were stored in RNA-later
(ThermoFisher, cat#AM7021) in -80°C until RNA isolation. After thawing samples,
aliquots were removed and diluted with 1X PBS and centrifuged to pellet cells and
remove RNA-later. RNA was isolated using the RNEasy Mini Kit (QIAGEN, cat#74134)
and cDNA was synthesized using the Invitrogen High-Capacity cDNA Reverse
Transcription Kit (ThermoFisher, cat#4368814). Quantitative PCR was performed on
synthesized cDNA using iTaq Universal SYBR Green Supermix (BioRad, #172511).
Normalization of all samples was calculated in comparison to *Actb* expression levels.
Fold change in cytokines were determined through the $2^{-\Delta\Delta Ct}$ method.

For parasite burden determination by quantitative PCR, peritoneal exudate and
spleen cells, liver and lung tissues were harvested, and appropriate amounts were
digested according to manufacturer's protocols using the QIAGEN DNeasy Blood &
Tissue Kit (#69504). Brains were isolated and DNA purified as follows. In brief, whole
brains were harvested in 5 mL of 1X PBS and homogenized through an 18G needle of a
10mL syringe and passed up and down through the needle 10 times before being spun
down at 611 x g for 7 minutes to pellet. 100 µL of brain homogenates were processed

using the QIAGEN DNeasy Blood & Tissue kit according to the manufacturer's
directions. Standard curves were made from known numbers of isolated parasite DNA
ranging from 10^1 to 10^6 parasites per well. Isolated DNA samples were amplified
targeting the *Toxoplasma*-specific B1 gene, and the standard curve was used to
quantify parasite number in each sample per µg of isolated DNA.

Statistics
Statistical analyses were performed with GraphPad Prism 8 software. Statistical
significance was defined as $P < 0.05$. P values between two groups were calculated
using paired or non-paired two-tailed t-tests. One or two-way ANOVA was used for
comparison across more than two groups with Tukey or Dunnett correction, as
recommended by Prism. Survival curve significance was calculated using log-rank
Mantel-Cox testing.

RESULTS

***Tg_207750* and *Tg_266320* are predicted to be the *T. gondii* GPI sidechain-**
modifying glycosyltransferases, PIGJ and PIGE, respectively.
T. gondii strains are known to differ in virulence in various intermediate hosts
(38). In laboratory mice, type III strains are referred to as avirulent because they require
a dose of over 10^5 parasites to achieve 50% lethality (38,39). Meanwhile, type I strains
are highly virulent, and one parasite is enough to be 100% lethal in naïve mice (40,41).
However, when C57BL/6J mice given a primary infection with the avirulent type III strain
CEP and allowed to progress to a chronic infection, they survive secondary infections
(i.e., "challenge") with high dose type I strain RH parasites due to protective
immunological memory responses that are generated following the initial parasite
exposure (42–44). In contrast, when mice infected with CEP are challenged with the
type I strain GT1 they succumb (43). RH and GT1 are highly similar type I strains and
the gene *Tg_207750* was noted to be six times more highly expressed in RH compared
to GT1 (45). Based on sequence analysis, *Tg_207750* was predicted to be a
glycosyltransferase that adds an acetylated hexose (HexNAc) to a core mannose of
complex N-glycans (45). Moreover, *Tg_207750* was one of two *T. gondii* βHexNAc
transferase-like GTs we proposed to be responsible for adding the GalNAc to the GPI of
T. gondii (36). Thus, we hypothesize that *Tg_207750* is partially responsible for
differences in secondary infection virulence between RH and GT1 type I strains and the
GPI GalNAc transferase we name here as "PIGJ" (**Fig 1A**).
Informatic analysis revealed *Tg_207750* is predicted to be a type 2
transmembrane protein (46) with a C-terminal glycosyltransferase domain with greatest
similarity to CAZy family GT17 (36) (**Fig 1B**). The occurrence of a non-synonymous
SNP within the GT17 CAZy-like domain was also noted between RH and GT1.
Reciprocal BLASTp studies indicate that PIGJ is coccidian specific and not found in
Plasmodium or *Cryptosporidium spp.*, or Piroplasm apicomplexans (not shown). The
catalytic DxD domain is separated from the transmembrane domain by a stretch of
about 170 aa, which in other GTs comprises a stem-like region that may mediate
protein interactions or appropriately orient the catalytic domain. The 143-aa cytoplasmic
region is predicted to be disordered based on amino acid composition. AlphaFold
predictions confirm the potential of the GT17 region to fold as a GT-A superfamily

domain, but do not confidently predict a structure for the stem-like or cytoplasmic
regions. Now referred to as PIGJ, *Tg_207750* evolved independently of PGAP4, the GT
that catalyzes the same reaction in animals (30). PGAP4 is a CAZy GT109 family
member that bears no recognizable homology with CAZy GT17 sequences and has a
distinct architecture that includes a transmembrane hairpin inserted within the canonical
catalytic domain. Thus, formation of the same GalNAcb1,4Man linkage in coccidian
apicomplexans and in animal hosts is the result of convergent evolution.
The gene *Tg_266320* was previously predicted to encode the GPI αGlcT that
mediates addition of the terminal Glc to GalNAc to complete the disaccharide sidechain
(36) (**Fig 1A**). *Tg_266320* is also predicted to be a type 2 transmembrane protein (**Fig.
1B**). Now referred to as "PIGE", *Tg_266320* has greatest similarity to CAZy GT32
glycosyltransferases (36). BLASTp searches suggest that this protein is likewise
restricted to coccidian apicomplexans (not shown). The most similar GT found is
lactosylceramide 4-alpha-galactosyltransferase (a.k.a. globotriaosylceramide or Gb3
synthase), which attaches αGal to the 4-position of a β-linked Gal acceptor across many
eukaryotes (47). Thus, the linkage formed is the same, but the donor and acceptor
sugars are distinct, which is an evolutionary variation common among CAZy GT32 GTs.
GT32 GTs utilize sugar nucleotide rather than Dol-P-sugar donors, as previously
documented for the enzyme that assembles the Glcα1,4-linkage in the *Toxoplasma* GPI
(48). Though PIGE and Gb3 synthase are both type 2 membrane proteins that modify
glycolipids, PIGE is distinctive with its ~385 aa long stem-like region and a ~160 aa
poorly conserved, likely disordered cytoplasmic region, whereas Gb3 synthase is only
360 aa overall with a negligible cytoplasmic region. Given evidence for multiprotein
complexes in the GPI biosynthetic pathways of other organisms (49), the cytoplasmic
and stem-like regions of PIGJ and PIGE might integrate them into complexes for orderly
and efficient sequential GPI-assembly in the rER (50). This hypothesis is consistent with
the lack of KDEL-like rER retentions signals at their C-termini.
580
Western blot analysis implicates *Tg_207750* as PIGJ
To address their roles in the GPI sidechain synthetic pathway, each predicted GT
gene was targeted for disruption by CRISPR-Cas9 and repair with a drug selectable
marker (**Fig S2 S3**). Mutants of each gene were made in both type I (RH, GT1) and type
III (CEP) genetic backgrounds (**Table S1**), and integration of the selectable markers
was confirmed with diagnostic primers and PCR (**Fig S2 S3**). Using glycoform-specific
antibodies that bind the protein-free GPI lipid (GIPL) sidechains (17), western blot
analysis revealed these antibodies lose recognition of both sidechain glycoforms
derived from the GT1 *Δ207750* and CEP *Δ207750* strains (**Fig 1C**). This indicates that
the enzyme encoded by *Tg_207750* is likely responsible for GalNAc addition to the
mannose core of the GPI. Western blot analysis of GT1 *Δ266320* and CEP *Δ266320*
mutants retained GIPL recognition by the T3 3F12 GalNAc GIPL glycoform specific
antibody, and in the case of GT1 *Δ266320* increased detection was routinely observed.
In contrast, recognition of the GalNAc + Glc glycoform by the T5 4E10 antibody was
unimpeded against GIPL from GT1 *Δ266320* and CEP *Δ266320* mutants (**Fig 1C**). A
technical note of importance, while the T5 4E10 antibody has preferential reactivity to
GalNAc + Glc compared to GalNAc GIPLs of *T. gondii* (17), it can detect GalNAc

sidechain bearing GIPLs of human intermediate GPIs (29), suggesting significant cross-
reactivity to antigen by this antibody clone.
To further explore the recognition potential by these antibodies, we decided to
monitor their ability to bind intact parasites by flow cytometry. Like that of our western
blot analysis, T3 3F12 and T5 4E10 staining of the GT1 *Δ207750* mutant was
significantly reduced compared to the parental parasite strain. However, the staining
was not entirely abrogated (**Fig S4**). Regarding the GT1 *Δ266320* strain, T3 3F12
recognition of the mutant parasite is enhanced suggesting increased presence of the
GalNAc bearing GPI sidechains. In contrast, T5 4E10 recognition of GT1 *Δ266320* is
only marginally reduced, calling into question whether this GT is responsible for adding
the terminal Glc to the GPI sidechain, at least using this antibody clone which has
significant cross-reactivity to GPI antigens (17,29). Finally, while *Tg_207750* is more
highly expressed in RH compared to GT1 type I strains (45,51), we do not detect GIPL
in the RH strain via western blot using these antibodies (**Fig 1C**), nor do we detect the
disaccharide sidechain by mass spectrometry of GPI anchors isolated from this strain
(36). Consistent with these observations, reduced T5 4E10 and T3 3F12 binding of the
RH strain relative to the GT1 strain is observed by flow cytometry, with only slight
reductions in T3 3F12 staining of the RH *Δ207750* compared to the parental RH strain
is seen (**Fig S4**).

Structural analysis confirms GPI sidechain glycosyltransferases of *T. gondii*.
Given the low detection of GIPL in the RH background by GIPL-specific antibody
and mass spec, the GT1 strain was subjected to GPI sidechain structural analysis via
mass spectrometry (**Fig 2**). A previous study developed a new method to isolate and
analyze the glycan component of GPI anchors using mass spectrometry (36). The
method first enriched for GPI-anchored proteins using butan-1-ol extraction of material
delipidated by extraction with $CHCl_3$ and MeOH. The fraction was saponified to remove
potential fatty acids (FAs), and the glycan core was released by phosphodiester bond
cleavage with hydrofluoric acid (HF). Finally, the glycan was N-acetylated and
permethylated, and analyzed initially by MALDI-TOF-MS. The most abundant ion from
the parental GT1 strain corresponded to a glycan with the composition of four hexose
residues, two HexNAcs, and one hexitol, or H4N2Ino (**Fig 2A**). This composition
corresponds to the previously described structure consisting of three Man, one Glc, one
GalNAc, and one inositol, as shown in Figure 1A. In addition, less abundant ions were
found that correspond to H3N2Ino and H3N1Ino, suggesting the presence of isoforms
with a monosaccharide side arm, or none at all. Ions whose composition corresponded
to multimers of hexose, H3, H4 and H5, were also detected.
Since parasites are grown in HFFs, we assessed the potential contribution of
HFFs to these ions, since mammals also express GPI anchor glycans with a
disaccharide side chain on the α4-linked (first) Man residue (29). Preliminary analysis of
N-glycans, performed as described in Gas-Pascual et al. (2019), showed the presence
of N-glycans from HFFs as well as tachyzoites (data not shown), such that >25% of the
sample was potentially of HFF origin. Therefore, a GPI-anchor fraction from HFFs was
examined. As shown in Figure 2D, this sample contained a series of hexose oligomers
whose abundance decreases with length, which potentially represent breakdown
products of glycogen, as well as an ion corresponding to H3N1Ino. Previous analysis of

a preparation of spontaneously lysed extracellular tachyzoites that were unlikely to be
contaminated by host cells, and that were washed extensively to minimize serum
contaminants, yielded no detectable hexosamers (36), consistent with the low level of
starch accumulation at the tachyzoite stage. However, the ratio of H3N1Ino to Hex4 was
small in HFFs compared to the large ratio in the GT1 sample (**Fig 2A, D**). If the amount
of H4 is used as a measure of contamination of HFFs to the parasite to sample, it is
evident that the great majority of the GPI glycans originated from the parasites.

To verify these interpretations, the GT1 GPI preparation was also examined by
MS analysis in an Orbitrap mass spectrometer after separation by C18 nano-HPLC. As
shown in Figure 3, the elution profile of all ions (base peak chromatogram) from the
GT1 strain included a prominent peak eluting at 49.04 min (**Fig 3A**). This peak
coincided with the extracted ion chromatogram analysis (EIC) of H4N2Ino in the lower
trace of panel A. A second peak eluting at about 33.77 min corresponded to the elution
positions of H3N1Ino and H3N2Ino, which were not well resolved by this method. These
ions eluted as a mixture of primarily singly charged H^+ or NH_4^+ ions, and doubly charged
ions in either $2H^+$ or H^+/NH_4^+ states, as shown in the right-hand panels. These ions
matched the expected *m*/*z* values with 1-2 orders of magnitude higher accuracy
compared to the MALDI-TOF-MS method. Furthermore, collision-based decomposition
(HCD MS-2) revealed a series of daughter ions that confirmed that it consisted of
various combinations of monosaccharides including non-reducing end Hex residues
(**Fig S5A**) that are completely consistent with structure shown in Figure 1A. As
expected, all 3 GPI isoforms were detected, but at a ratio of 1.0:0.25:0.13 for
H4N2Ino:H3N2Ino:H3N1Ino, respectively (**Fig 3A**), consistent with the MALDI-TOF-MS
data (**Fig 2A**). Most of the remaining ions in the base peak chromatogram matched the
elution positions of the H3, H4 and H5 hexosamers, as described for the HFF GPI
sample (**Fig 3D**). Note that these isomers elute in pairs with the same *m*/*z* value,
indicating that they represent a mixture of reducing end α- and β-anomers with distinct
elution times. While all three GPI isoforms were also detected in the HFF samples, they
were of very low relative abundance and H3N1Ino rather than H4N2Ino was the most
abundant ion. Taken altogether, the results confirm the presence of the previously
described Glc-GalNAc- side chain of parasite GPI anchors, and good evidence for
isomers with only GalNAc or no side chain at all.

To confirm the predicted role of PIGE as the GPI Glc transferase, the GPI
fraction from *Δpige* parasites was analyzed as above. Strikingly, only the H3N2Ino and
H3N1Ino isoforms were detected, along with substantial levels of H4 and H5 (**Fig 2B**).
Using the more sensitive nLC-MS method, the full-length H4N2Ino isoform was
detected but at 0.005% of the level of H3N2Ino (**Fig 3B**), which is likely a contaminant
from the HFF cells. To confirm the predicted role of PIGJ as the GPI GalNAc
transferase, the GPI fraction from *Δpigj* parasites was similarly analyzed. In this case,
essentially only the H3N1Ino glycan was detected together with substantial H3 and H4
glycans (**Fig 3C**). The trace levels of H4N2Ino and H3N2Ino glycans detected were
readily explained as carryover from the host HFF cells as indicated by the abundant H3
and H4 glycans. Both mutant forms of the GPI-anchor fragmented to yield expected
glycan products (**Fig S5B, C**).

T. gondii GIPLs have been described to possess the same Glc-GalNAc- sidearm
on the same α4-linked core Man residue. To confirm the findings described above for

GPI-anchors, the method for isolating GPI-glycans was applied to the organic fraction
obtained from the initial delipidation step used to isolate GPI-anchored proteins, as
described in Material and Methods. Using the same MALDI-TOF-MS method as above
though with a different matrix, a series of ions was observed that suggested the
presence of H4N2Ino, H3N2Ino, H3N1Ino, and H4 (**Fig S6A**). However, the m/z values
for the parasite glycans were 36 units smaller than expected, whereas H4 had the
expected value. In comparison, a parallel analysis using a different matrix yielded the
expected m/z for the most abundant ion (**Fig S6A inset**) albeit also substantial -36 ion.
In addition, ions at -14 units were detected at low abundance, suggesting slightly
incomplete methylation. To examine whether these mass defects were an artifact of
MALDI-ionization, the sample was also analyzed by nLC-MS. Significantly, only ions
corresponding to the expected values were detected (**Fig S7A**), and ions with mass
defects of -36 and -14 were insignificant. Therefore, these defects, which were also
present at trace levels in the GPI-anchor analysis (**Fig 2B, C**) but not originally noted,
were evidently artifacts of the MALDI ionization method. The ratio of
H4N2Ino:H3N2INo:H3N1Ino (1.0:0.33:0.17) was like that of the GPI isoform ratios
observed in the MALDI-TOF-MS method.
Analysis of the GIPL fractions from *Δpige* and *Δpigj* strains yielded the same
findings as from the GPI-anchor studies. Using either MALDI-TOF-MS (**Fig S6B**) or
nLC-MS (**Fig S7B**), the *Δpige* sample contained primarily the monosaccharide side
chain with negligible disaccharide side chain presence. Similarly, the *Δpigj* sample
lacked the side chain altogether (**Figs S6C, S7C**) with the miniscule levels of side chain
isoforms attributable to host cell contamination (**Figs S6D, S7C**). Thus, the same
glycosyltransferases act to generate the Glc-GalNAc- side-arm in both GPI-APs and
GIPLs, and a significant fraction of glycans contain a monosaccharide or no side-arm in
the wildtype GT1 strain as well. In summary, mass spec results confirm the predicted
role of both glycosyltransferases. *Δ266320*, the gene hereon referred to as PIGE, lacks
the GalNAc + Glc glycoform, and *Δ207750*, the gene hereon referred to as PIGJ,
completely lacks both glycoforms of the sidechain (GalNAc, GalNAc + Glc) in the GT1
background.
720
PIGJ mutants have increased primary and secondary infection virulence.
With the knowledge that PIGJ and PIGE are the *T. gondii* GPI sidechain GTs, the
opportunity presented itself to test the role of distinct GPI glycoforms in microbial
virulence. To this end both primary and secondary infections were performed using the
PIGJ and PIGE mutant strains. Importantly, primary infections with the CEP *Δpigj* strain
resulted in 60% mortality compared to 0% mortality following wildtype infections (**Fig
4A**). In contrast, mice infected with CEP *Δpige* survived primary infections (**Fig 4A**).
Secondary infections were performed in mice chronically infected with the type III strain,
and as previously reported (43,44), these mice survive type I RH challenge due to the
formation of protective immunity (**Fig 4B**). Importantly, RH *Δpigj* but not RH *Δpige*
mutants caused lethal secondary infections. Chronically infected C57BL/6J mice were
also challenged with type I GT1 mutants and succumbed similarly to all three strains:
GT1, GT1 *Δpigj,* and GT1 *Δpige* (**Fig S8**). Given C57BL6/J mice are susceptible to GT1
challenges (43), it is difficult to observe virulence increases in this context. To explore
this further, the virulence phenotypes of the GPI sidechain mutants were screened in

A/J mice, which are genetically resistant to primary infections with low virulence type II
strains (52) and secondary type I GT1 infections (53). As observed in C57BL/6 mice,
CEP *Δpigj* caused lethality during primary infections in A/J mice (**Fig 4C**). During
secondary infections with the RH and GT1 GPI sidechain mutants, however, A/J mice
were still resistant (**Fig 4D**). Hence, the *Δpigj* deletion produces a virulence phenotype
that overcomes the genetic basis for resistance to primary but not secondary infections
in A/J mice. Collectively, these data suggest that the GPI sidechain of *T. gondii* prevents
lethal outcomes in its hosts, and that the virulence phenotype is less impacted by the
terminal Glc residue transferred by PIGE. Furthermore, the data is consistent with the
original hypothesis that differential *Tg_207750* gene expression contributes to the type I
strain differences in secondary infection virulence, at least in the susceptible C57BL/6
genetic background.
748
Complementation of *PIGJ* partially rescues the survival phenotype.
Given the most consistent phenotype surrounding the GPI sidechain occurred
with the CEP *Δpigj* strain, a PIGJ complementation strain was generated in this
background and efforts were focused on primary infection virulence. The full-length
coding region with introns spliced out of *TgVEG_207750* was amplified from cDNA and
fused with 1000 bps of the gene's promoter, followed by delivery as a transgene into
CEP *Δpigj* parasites using a pLIC-DHFR-3xHA plasmid which adds a C-terminal
hemagglutinin tag to the gene of interest (54). Western blot analysis shows the C2
complementation clone, CEP *Δpigj* + *PIGJ*$_{3xHA}$, has its GPI sidechain glycoforms
restored albeit to levels lower than wildtype (**Fig 5A**), and the presence of the HA tag is
detectable by fluorescence microscopy (**Fig 5B**). Both diffuse and puncta distribution of
PIGJ$_{3xHA}$ are seen throughout the cell of the parasite with no distinctive localization
pattern to determine its subcellular localization. Survival, weight loss and cyst formation
following primary infection with the C2 complementation strain were assessed. While
complete rescue of the avirulence phenotype was not attained following C2 infections, a
significant increase in mouse survival and reduced weight loss occurred during primary
infections (**Fig 5C**), indicating a partial complementation of the virulence phenotype is
achieved. When brain cyst burdens were analyzed in the survivors of the primary
infections, significantly more brain cysts were detected in the mice infected with CEP
Δpigj compared to wildtype strains (**Fig 5D**). However, the complementation strain was
not significantly different compared to the WT or *Δpigj* mutant strains (**Fig 5D**).
To assess whether fitness defects may explain the *in vivo* phenotypes, lytic cycle
analysis was performed. Consistent with genome wide fitness studies (4) (**Fig 1A**),
deletion of PIGJ did not impede parasite growth in HFFs or MEFs, nor attachment to
host cells (**Fig S9**). It was noted CEP *Δpigj* had significantly larger plaque areas than
wildtype CEP, and this gain in plaque size was retained in the complementation strain,
suggesting that either the level of complementation of PIGJ was insufficient or the
background of the *Δpigj* strain gained a growth advantage during cloning irrespective of
the *PIGJ* gene (**Fig S9**). Regardless, differences in *in vitro* growth rates between strains
do not correlate with their pathogenesis phenotypes. In summary, virulence inversely
correlates with the expression of *PIGJ* and implicates the GPI sidechain of *T. gondii* as
a factor that inhibits pathogenesis.
781

Early parasite burden and cytokine responses are unchanged following CEP Δ*pigj* primary infections.

The early weight loss and death during primary infections pointed to a possible immune defect responsible for the increased virulence of the PIGJ mutants. Therefore, mice were infected with either the wildtype CEP or CEP Δ*pigj* strains and multiple organs were assessed for parasite burden as well as cytokine responses in the peritoneal cavity (**Fig 6**). *T. gondii* elicits a robust TH1 response and IFNγ is essential for survival, therefore gene expression of major cytokines known to be involved in TH1 immunity (*Il6, Il12b, Ifng, Cxcl10)* and its regulation (*Il10*) were measured. As GIPL induces robust TLR2/4-dependent TNFα production in macrophages (16), this cytokine was measured (*Tnfa*), as well as type I IFN (*Ifna, Ifnb1*) and type I IFN induced genes (*Isg15, Mx1*) due to the protective role this pathway plays during primary *T. gondii* infection (55). While there were slight increases in some cytokine response to the PIGJ mutant compared to wildtype infections, no significant differences were detected (**Fig 6A, C, E**). Moreover, parasite burden in the peritoneum, spleen, lung, and liver appeared the same between mutant and parental strain as measured by qPCR (**Fig 6B, D, F**), or by flowcytometry for GFP+ parasites in the peritoneum (**Fig S10**). Therefore, host susceptibility to primary infection with the PIGJ mutant is not related to overt dysregulation in the cytokine response or parasite burden during acute infection.

Antibody reactivity, function and expression of GPI-anchored SAGs remain largely intact to PIGJ mutants.

We wondered whether the loss of the sidechain influenced regulation of GPI anchored protein expression and/or antibody recognition of GPI-APs, and if this correlated with parasite virulence. Some reports suggest GPI-sidechains may alter the protein conformation of the GPI-AP (56), and the amount of time a GPI-AP spends folded over on the plasma membrane (57), both of which could potentially impact antibody recognition of GPI-APs. To this end, the surface expression of major GPI anchored surface proteins SAG1, SAG3, and p35 on wildtype and Δ*pigj* strains was measured with monoclonal antibodies by flow cytometry. For both RH and CEP strains, deletion of PIGJ did not prevent surface expression nor detection of the aforementioned SAGs (**Fig 7A**).

Parasite-specific antibody derived from the sera of mice infected with *T. gondii* allows a global assessment for any defects that polyclonal antibody recognition might have against a multitude of surface antigens regulated by the GPI sidechain. To this end, parasites were incubated with serum from chronically infected mice, and fluorescently labeled anti-isotype secondary antibodies were used to detect isotypes of parasite-bound serum antibodies as previously described (53). Again, no differences were observed for IgM, IgG3, IgG1, Ig2a/c and IgG2b reactivity to wildtype and PIGJ mutants over a range of serum dilutions (**Fig 7B**) (not shown). Complement component 3 (C3) binds the surface of *T. gondii* and is required for host resistance to infection (58). Incubating parasites with naïve serum as a source of complement and detection with an anti-C3 antibody revealed no difference in C3 recognition between WT and PIGJ mutant strains (**Fig S11**).

To screen for any functional defects of immune or naïve sera due to the loss of the GPI sidechain, opsonization and neuralization assays were performed. For

opsonization, parasites were incubated with sera before being added to bone marrow
derived macrophages (BMDMs) and allowed to either invade or be phagocytosed. Via
microscopy, successful invasion, and formation of the parasitophorous vacuole is
marked by GRA7+, while LAMP-1 marks the phagolysosome and when associated with
parasites, indicates opsonization (59). No significant differences in opsonization were
detected between parental and GPI mutant strains, although the CEP *Δpigj* strain
trended slightly higher compared to the parental CEP strain (**Fig 7C**). To assay
neutralization, serum coated parasites were incubated with mouse embryonic
fibroblasts (MEFs) and assessed for relative invasion by flow cytometry, and no
differences in antibody neutralization were observed between strains (**Fig 7D**).
Therefore, effector functions of immune and naïve sera are largely intact to GPI
sidechain null *T. gondii* parasites.
840
The terminal glucose is required for immune serum IgM recognition of GIPL.
Given the terminal Glc but not the GalNAc+Glc GPI sidechains is recognized by
serum from humans latently infected with *T. gondii* (21), GIPL reactivity of serum from
chronically infected mice was assessed against the GPI mutants by western analysis.
Sera from both C57BL/6J (**Fig 8A**) and A/J mice were used (**Fig 8B**). First, GIPL
reactivity, as observed by a band at 4-5 kDa when probed with anti-mouse IgM, is more
robust using A/J mouse serum, but faint recognition can be seen with the C57BL/6J
serum as well. Second, and consistent with the flow cytometry data, antibody reactivity
to parasite antigens 18 kDa and larger did not change between the wildtype and
knockout parasites. Third, RH GIPL is not detected by poly clonal immune sera, which
is again consistent with previous mass-spectrometry and western blot analysis.
Importantly, loss of IgM GIPL reactivity to *Δpige* and *Δpigj* was clearly observed for both
GT1 and CEP mutants (**Fig 8A B**). Therefore, the GalNAc + Glc glycoform of the GPI
sidechain is required for serum antibody recognition of GIPL, as previously inferred
(17,21). However, given the drop in serum IgM reactivity to both virulent (*Δpigj*) and
avirulent (*Δpige*) mutants, we reason this phenotype is unrelated to primary and
secondary infection virulence in mice.
858
Non-significant macrophage tropism but enhanced CD36 binding of PIGJ
mutants.
To test whether cellular tropism of GFP+ expressing parasites may explain
difference in host susceptibility early in infection, mice were infected with 10^6 parasites
and 3 hours later peritoneal exudate cells were analyzed by flow cytometry as
previously described (60). Invasion of PECs was estimated by analyzing FSC and SSC
parameters of GFP+ events to differentiate by size (**Fig 9A**). Given similar GFP signal
(**Fig 9B**), no differences in cell associated and intracellular (FSC^{int-hi} SSC^{int-hi}) vs. non-
cell associated and extracellular (FSC^{lo} SSC^{lo}) GFP+ parasites were observed between
strains (**Fig 9C**). Cell death, as inferred by PI staining, was also similar between
parasites, suggesting similar death kinetics of parasite infected peritoneal cells (**Fig 9D**).
The cell associated GFP+ events were further analyzed for markers to identify
infected cell types as a means of assessing parasite tropism. Gr-1^{hi} Cd11b+ neutrophils
had no significant differences in parasite infection (**Fig 9E**). Regarding macrophages,
there was a higher frequency of bone marrow-derived macrophages (MHCII^{hi} F4/80^{lo}

"BMDM" cells) infected with CEP *Δpigj* compared to WT strains, indicating a tropism for
BMDMs (**Fig 9F**). Infection frequencies of BMDMs with the complementation strain
mirrored that of the WT strain and was reduced in relation to *Δpigj*. A similar trend was
observed for MHCII^{hi}, F4/80^{hi} yolk-sack macrophages (YSM) (**Fig 9G**). However, due to
the biological variability, differences in BMDM or YSM tropism between strains did not
reach statistical significance. To test whether this subtle tropism was related to
enhanced invasion of macrophages, invasion assays were performed using *in vitro*
differentiated BMDMs. When BMDMs were infected for 20 minutes and
intracellular/extracellular parasites were quantified, no difference in cellular invasion
was observed between the parental and *Δpigj* strains (**Fig 9H**).
CD36 is a scavenger receptor expressed in many tissues and immune cells that
plays a role in fatty acid uptake, angiogenesis and phagocytosis (61). Moreover, *T.*
gondii tropism for macrophages is mediated in part by CD36 (60). Therefore, parasites
were incubated with recombinant CD36, and binding was measured via flow cytometry.
CD36 binding was found to be significantly increased to CEP *Δpigj* compared to
wildtype CEP strains (**Fig 9I J**). Whether macrophage tropism is inhibited by PIGJ,
presumably through the addition of the GPI sidechain, and whether strain-dependent
differences in macrophage association are mediated by CD36 is unknown.
892
Virulence of *Δpigj* strains is galectin-3 and sex dependent, but not TLR-2 nor -4
dependent.
Since TLR2 and TLR4 (16) and galectin-3 recognize *T. gondii* GIPL (50), we
hypothesized that TLR and/or galectin-3 signaling would mediate differences in the
PIGJ-dependent virulence. First, *Tlr2/Tlr4 -/-* double knockout mice were used to study
survival. However, the double knockout mice similarly succumbed to CEP *Δpigj* while
surviving to CEP (**Fig 10A**), suggesting that differences in the GPI sidechain do not
impact host-parasite interactions mediated by these PRRs. In contrast, mice lacking the
gene for galectin-3, *Lgals3 -/-*, survived both to CEP and to CEP *Δpigj* infections (**Fig**
10B). This indicates that the *Δpigj* virulence observed in wildtype mice is dependent on
galectin-3. The CEP *Δpigj* infected mice still had significant weight loss, but this
occurred later in the infection (**Fig 10C**). Despite the weight loss, the mice survived, and
did not have significant differences in brain cyst burdens (**Fig 10D**). When recombinant
galectin-3 binding was measured on whole parasites via flow cytometry, an increase in
galectin-3 binding to CEP *Δpigj* parasites was observed (**Fig 10E F**). It is unknown
whether host factors like CD36 and galectin-3, unimpeded by a sidechain, preferentially
bind the GPI mannose core to regulate the pathogenesis of *Δpigj* strains. Regardless,
when mice lack galectin-3, *Δpigj* virulence is disrupted and the *Lgals3 -/-* mice are
protected.
One final and surprising observation was that male C57BL/6J mice infected in
these studies were resistant to both primary and secondary infections with *Δpigj*
mutants (**Fig S12**). These mice sustained their weight similarly to the wildtype infected
mice (**Fig S12B D**) and had similar survival (**Fig S12A C**). There are sex differences in
a variety of immune responses, which have been reviewed (62). Specifically in *T.*
gondii, females have been shown to be more susceptible to acute infections compared
to males, with different cytokine kinetics, and the females were found to have higher

brain cyst burdens (63). While the sex basis for this phenotype was not explored further,
this observation may help unravel the mechanism behind *Δpigj* virulence.
921
922　**DISCUSSION**
923
This study aimed to address the following questions and gaps in the field's
knowledge: 1) What are the GPI sidechain-modifying enzymes of *T. gondii*, and 2) Does
the sidechain of *T. gondii* play a role in parasite virulence? In our attempts to answer
these questions we made several critical observations. Firstly, we identified the two
glycosyltransferases responsible for GPI sidechain modification in *T. gondii*,
Tg_207750, named PIGJ, which adds the GalNAc to the mannose of the core
backbone, and *Tg_266320*, named PIGE, which adds the glucose to the GalNAc
sidechain. We confirmed their activity using mass spectrometry analysis of the GPI in
the knockout parasites, which clarified ambiguities using monoclonal antibodies which
apparently have some cross-reactivity to unknown epitopes. In addition to the
characterization of these glycosyltransferases, we explored their role in parasite
virulence in both primary and secondary infections which revealed that PIGJ mutants
are virulent in both settings. This study is a first analysis for the role of the GPI
sidechain for any microbe in pathogenesis.
Removal of the core GPI synthesizing enzymes is lethal in mammals and in
eukaryotic microbes. However, it remained an outstanding question whether complete
removal of the GPI sidechain would have fitness bearing impacts in parasites. The
genes identified here do not have deleterious impacts on the lytic cycle, and the fitness
scores for PIGJ and PIGE align with this supposition (**Fig 1A**). In performing reciprocal
blasts using *T. gondii* PIGJ and PIGE, we find evidence that homologous enzymes exist
in coccidian species, but not in more distantly related apicomplexan species such as
Plasmodium and *Cryptosporidium* (not shown). Coccidia parasites are orally acquired
parasites of warm-blooded animals that have an intra-epithelial stage in their
development, but many form tissue cysts, which require parasite dissemination
following infection. Whether all coccidia parasites express similar GalNAc + Glc bearing
sidechains is unknown but would be expected from the identification of PIGJ and PIGE
homologs across this subclass of parasites. It is tempting to speculate that the GPI
sidechain, with a structure similar to that of mammals (**Fig S1**), use this similarity as the
result of convergent evolution, to mimic a fundamental biological process mediated by
this glycoform and required in these environments.
In our search for PIGJ and PIGE functions, we characterized and knocked out
the enzymes responsible for GPI sidechain synthesis in both type I (GT1, RH) and a
type III (CEP) strains of *T. gondii*. Parasites lacking PIGJ, but not those lacking PIGE,
exhibit increased virulence in primary and secondary infections, demonstrating the GPI
sidechain modulates parasite virulence by preventing lethality in its host. To elucidate
the mechanism for increased virulence of the PIGJ mutant we characterized a variety of
immune parameters and parasite burden following infection. Our findings indicate early
parasite burden in the peritoneal cavity, liver, and lung are similar, yet the PIGJ mutants
can evade clearance and establish higher cyst burdens in the brain. Cytokine
responses, antibody reactivity, opsonization, neutralization, and parasite surface
antigen expression were all similar between strains that express a GPI sidechain and

those that do not. We speculate that a breakdown of infectious tolerance occurs early
during infection when the parasite lacks a GPI sidechain. Whether enhanced tissue
damage results following infection with GPI sidechain null parasites, is unknown.
To further explore where the pathogenesis might take place, a subtle cell tropism
for BMDMs was observed in the peritoneal cavity 3 hrs after infection, and CD36
binding was found to be increased with CEP *Δpigj*. We also found that *Lgals3* -/- mice
were protected against primary infections with CEP *Δpigj*, indicating a fundamental role
of galectin-3 in the susceptibility of wildtype mice to CEP *Δpigj* infections. Galectin-3 is
one of the few known host factors that bind *T. gondii* GIPL (50), and recombinant
galectin-3 preferentially bound CEP *Δpigj* strains. Galectin-3 was previously shown to
be required for macrophage TNFα production in response to *T. gondii*, and it was
proposed that galectin-3 might act as a co-receptor that presents the GPI to TLR2 (50).
Our data of PIGJ infections in double knockout mice of *Tlr2/4* -/- indicated that TLR2/4
signaling was not involved in the virulence differences. It is likely there are other cellular
functions mediated by galectin-3 binding to GIPL that do not involve TLR2/4, however,
the exact mechanisms are unknown. Previous reports have shown that following *T.*
gondii infection, galectin-3 is upregulated in peripheral tissues during acute infection of
T. gondii and were susceptible to intraperitoneal infections with type II strains of *T.*
gondii (64). Interestingly, *Lgals3* -/- mice had higher Th1 responses with increased IL-
12p40 and IFNγ detected in the sera, as well as from cultured splenocytes after
infection (64). It is possible the enhanced Th1 response underpins the resistance of
Lgals3 -/- mice to type III strains but renders them susceptible to more virulent type II
strains. Regardless, further mechanistic insight is required to draw correlations between
galectin-3 and the GPI sidechain, and to place this interaction as central to the disease
outcome of PIGJ mutants.
Of further note on the CD36 pathway, it is possible that mice infected with *Δpigj*
strains are more able to phagocytose parasites via CD36, allowing for enhanced
infection and escape via the phagosome cellular-invasion route reported for less virulent
parasite strains (59). *Cd36* -/- susceptibility to *T. gondii* infections was suggested to
correlate with a breakdown in tissue homeostasis following infection (60). This was
revealed by monitoring serum indicators of tissue stress including GDF-15 and FGF21.
GDF-15 induces anorexia by binding to the GFRAL receptor expressed by neurons of
the hindbrain (65), while GFG21 is a hormone that controls energy homeostasis and
adiposity (66) and is often induced in multiple tissues by nutrient starvation and
endoplasmic reticulum stress (67). Given the PIGJ mutants appear normal with respect
to parasite burden and the disease outcome is sex linked, we suspect the GPI sidechain
promotes tissue homeostasis during infection by potentially controlling mechanisms of
tolerance via CD36 interactions and/or other host pathways yet to be identified.
In our initial approach to confirm the loss of the Glc and GalNAc sidechain
additions, we utilized glycoform specific antibodies T3 3F12 and T5 4E10. While
complete removal of the sidechain in the *Δpigj* mutants was clearly observed through
western blotting, the removal of the glucose addition was not. The T5 4E10 antibody,
though it is described to be specific for the GalNAc + Glc glycoform, has considerable
cross-reactivity to the GalNAc only glycoform, as seen in previous studies utilizing it
(17,29). The T3 3F12 antibody clone is an IgG3 isotype, and the T5 4E10 clone is an
IgM isotype. It is interesting to note, B-1 cells preferentially produce antibodies of IgM

and IgG3 isotypes. B-1 cells are responsible for natural self-reactive IgM antibodies
and are known to contribute to T-independent response, most of which are against non-
protein antigens (68). We originally looked into the GPI as a non-protein antigen after
we found that B-1 cells are important for immunity to secondary infections of T. gondii
(53). With respect to the IgM response, we did note that IgM recognition of GIPL was
highly sensitive to the terminal glucose (**Fig 8**), as previously observed using chemically
synthesized GPI and serum from latently infected humans (21). Whether the antibody
response to GIPL is from B-1 cells is unknown. Regardless, the lack of antibody
reactivity to GIPLs of the PIGE and PIGJ mutants, cannot account for why these strains
differ in primary and secondary infection virulence.
The analysis of the type I RH strain presents additional insights into the
significance of the sidechain and its biosynthesis. Previous reports have indicated
detection of RH GIPL and its sidechain (17). However, a recent report using mass
spectrometry of RH GPI-AP revealed only the linear mannose core of GPI with no
sidechain (36). A potential cause for this could be the lab adaptation of this strain. This
strain was initially passaged in mice and then in tissue culture for over 60 years before
ever being frozen down. Due to this, the RH strain has become lab adapted, and
acquired some unique characteristics (45). For instance, RH has lost its ability to form
orally infectious tissue cysts, grows much more rapidly than other strains, and has
increased extracellular viability (69,70). We have discovered that the type I RH strain
has lost its ability to evade immunological memory responses that are generated
following vaccination or natural infection, which we demonstrated to occur with other
type I strains like GT1 (43,53). It is likely that for whatever reason, RH has lost its
expression of the GPI sidechain through this laboratory evolution. However, it was an
initial report that the gene *Tg_207750* was almost 6 times more highly expressed in RH
compared to GT1 that led us to investigate it as a GPI sidechain GT candidate (45).
Overexpression may play a role in the loss of the sidechain, and a second possible
explanation is the previously reported non-synonymous SNP that generates a L620R
substitution uniquely in this strain relative to all other available sequences (**Fig 1B**) (not
shown). This position is conserved as a hydrophobic core of this region of the enzyme
that is involved in coordinating a PO_4 group of the UDP-GalNAc substrate. Whether this
acquired mutation in RH is inactivating to PIGJ, which in turn leads this strain to
compensate by overexpressing the gene is unknown. Interestingly, full deletion of the
gene increased virulence in secondary infections, suggesting an additional role for PIGJ
that might be mediated by other regions of the protein. It is possible that PIGJ, like most
other enzymes in the GPI processing pathway (49), belongs to a multiprotein complex
that depends in part on the presence of the PIGJ protein, and that its absence disturbs
some other aspect of GPI anchor assembly such as selectivity or efficiency.
In conclusion, this study identified and characterized both GPI sidechain
modifying glycosyltransferases of *T. gondii*. In addition, for the first time in any microbe,
the effects of the complete loss of the sidechain were studied for its role in
pathogenicity. The results presented here indicate a fundamental role of the presence of
the sidechain in host survival, and that when the sidechain of the GPI is lost, so is host
resistance to *T. gondii*. Although most of the major surface antigens (SAGs) of *T. gondii*
are GPI-anchored proteins and the GPI is known to be targeted by antibody responses
in a variety of protozoan infections (19,26), our findings here would suggest that for *T.*

gondii antigens, the sidechain presence does not impact antibody reactivity to GPI-
anchored proteins. However, whatever other humoral functions are at play regarding
host recognition of GIPL, they are wildly disrupted when the sidechain is lost. We
suggest both a fundamental, and yet diverse role for the GPI sidechain in microbial
pathogenesis. Based on our data, we propose a model in which enhanced CD36 and
galectin-3 binding of PIGJ mutants lead to pathology, perhaps through differences in
cellular tropism and unknown inflammatory mediators yet to be defined (**Fig S13**).
1064

ACKNOWLEDGEMENTS
Jean Françios-Dubremetz (U Montpellier) for the T5 4E10 ascites. Boris Striepen (U
Penn) for technical support and guidance. John Boothroyd (Stanford) for the rabbit
GRA7 polyclonal antibody. Gregory Barton (UC Berkeley) for the *Tlr2/4* -/- mice. Jeroen
Saeij (UC Davis) for plasmids. Dave Gravano for assistance with flow cytometry (UC
Merced, SCIF). The following monoclonal antibodies (produced in vitro) were obtained
through BEI Resources, NIAID, NIH: Anti-*Toxoplasma gondii*
Glycosylphosphatidylinositol Anchor, Clone T3 3F12, NR-50253; anti-*T. gondii* Surface
Antigen 1, Clone T4 1E5, NR-50255, anti-*T. gondii* Surface Antigen 3, Clone T4 1F12,
NR-50257, anti-*T. gondii* Surface Antigen P35, Clone T4 3F12, NR-50259. This work
was supported by NIH grants 1R01 AI137126 and 1R21 AI176186 to KDCJ; 1R21
AI123161 to CW; R21 AI171670 to GY. JP was supported by the supplement 3R01
AI137126-04S1, and GC was supported by a UC LEADS fellowship.

1078
FIGURE LEGENDS
Figure 1: Glycoform specific antibodies implicate *Tg_207750* as the GalNAc GPI
sidechain modifying enzyme, PIGJ.
A) Schematic of the T. gondii GPI with known and putative glycosyltransferases
involved in the GPI synthetic pathway. Transferases show their gene ID, gene name,
transferase type, fitness score (the more negative the fitness score being more lethal to
the parasite when inactivated) (4), and the linkage they form. The two novel
glycosyltransferases identified and characterized in this study are labeled in color and
named 'PIGJ' and 'PIGE' responsible for the GalNAc and Glc additions, respectively. B)
Gene models for *Tg_207750* (PIGJ) and *Tg_266320* (PIGE) from ToxoDB were
translated and subjected to BLASTp and other homology searches to predict the
evolution and function of the proteins. Regions of predicted function are represented:
TM transmembrane, putative stem region, DxD catalytic site, N-sequon potential
glycosylation sites and a non-synonymous SNP at position 620 in PIGJ between GT1
(L) and RH (R) type I strains. C) Representative western blots confirming the loss of the
sidechain in *Δ207750* mutants through sidechain specific antibody reactivity to the low
molecular weight (<10 kDa) GIPL antigen (Clones: T3 3F12 binds to the GalNAc
glycoform and T5 4E10 prefers the GalNAc + Glc glycoform of GIPL). GIPL from CEP
and GT1 *Δ266320* mutants are recognized by both glycoform-specific antibodies. GIPL
from RH was not detected by T3 3F12 nor T5 4E10. Representative of 3-5 experiments.
1099
1100 **Figure 2: MALDI-TOF-MS analysis of GPI preparations from GPI-anchored**
1101 **proteins reveal the identities of PIGE and PIGJ as GPI sidechain**
1102 **glycosylransferases.**

Glycans isolated from GPI-anchored proteins from tachyzoite stage parasites were N-acetylated, permethylated, mixed with CHCA matrix and analyzed in reflectron and positive ion mode. The m/z range of 800-1700 is shown. Ions that correspond to singly charged (z=1) sodiated GPI glycans are labeled in green, with the observed monoisotopic m/z value (in dark red), and the predicted m/z values for the assignment (in black). As an example, the composition assignment for an ion possessing three hexoses (known to be mannoses), one HexNAc (known to be N-acetylglucosamine), and one hexitol (known to be inositol) is represented as H3N1Ino. Ions that correspond to Hex(n) oligomers are labeled in red. A) Parental GT1 strain. B) *Δpige*. C) *Δpigj*. D) Host cells (HFFs).

Figure 3: nLC-MS analysis of GPI-anchor preparations confirm the GPI sidechain transferase activity of PIGJ and PIGE in *T. gondii*.

Isolated GPI permethylated glycans described in Figure 2 were reanalyzed by nLC separation on a C18 column and hyphenated analysis in an Orbitrap mass spectrometer in positive ion mode. The left column of panels shows base peak chromatograms (all ions m/z 500-2000) and extracted ion chromatograms (EIC) for each of the indicated targets (m/z ranges are specified in panels A and D). The ratio of ion intensities over the EIC m/z ranges shown for H3N1Ino, H3N2Ino, and H4N2Ino are shown relative to the most abundant ion. The most abundant ions are labeled in the base peak chromatogram. The right column of panels shows representative mass spectra of the two most abundant ions eluting with the target and used for quantitating relative levels. Levels of hexosamers were not quantitated. Observed m/z values are in dark red, expected values are in black, and the charge state is as indicated. A) Parental GT1 strain. B) *Δpige*. C) *Δpigj*. D) Host cells (HFFs).

Figure 4: Deletion of the GPI sidechain glycosyltransferase PIGJ but not PIGE results in increased parasite virulence.

A) C57BL/6J (B6) mice were given primary infections of 10^4 parasites (i.p.) of the indicated variants of the avirulent type III CEP *hxgprt- GFP:cLuc* strains. Cumulative survival from 2-4 experiments is plotted (mice; n = 20 CEP, 19 CEP *Δpigj*, n = 11 CEP *Δpige*). B) B6 mice given a primary infection with CEP *hxgprt-* and 35 days later (i.e. "chronically infected") were given a secondary infection with 5×10^4 parasites of the indicated variants of the type I RH strain and tracked for survival. Cumulative survival from 4 experiments is plotted (mice; n = 9 RH, n = 12 RH *Δpigj*, n = 6 RH *Δpige*). C) A/J mice were given primary infections of 10^4 parasites of either CEP or CEP *Δpigj* and monitored for survival for 30 days. Cumulative survival from 2 experiments is plotted (mice; n = 20 CEP, n = 20 CEP *Δpigj*,) D) After A/J mice were given a primary infection with the avirulent type III strain CEP, 35 days later were given a secondary infection with 5×10^4 parasites of the indicated type I strains and tracked for survival. A single experiment was performed (mice; n = 5 RH, n = 3 RH *Δpigj*, n = 2 GT1, n = 5 GT1 *Δpigj*, n = 10 GT1 *Δpige*). For survival analysis, significance was determined by Log-rank (Mantel-Cox) test, **** p <0.001, *** p<0.01.

Figure 5: Complementation of PIGJ into *Δpigj* partially restores avirulence of the type III CEP strain.

A) Western blot analysis using glycoform specific antibodies for the two GPI sidechains demonstrates GIPL reactivity is restored to the complementation strain. B) Fluorescence microscopy showing HA-tagged PIGJ localization in CEP Δ*pigj* + *PIGJ$_{3XHA}$* parasites. The complementation strain expresses GFP, as all CEP GPI mutants were derived from a CEP *hxgprt- GFP:cLuc* parental strain. C) Mice were injected i.p. with 10^4 parasites of CEP, CEP Δ*pigj*, or CEP Δ*pigj* + *PIGJ* strains and monitored for survival for 30 days. Weight loss after primary infection was normalized to the initial weight (=1), the average +/- SD of each cohort is shown. Cumulative results from 5 experiments are plotted (mice; n = 4 CEP, n = 14 CEP Δ*pigj*, n = 36 CEP Δ*pigj* + *PIGJ*). D) Brain cysts were numerated from survivors of the indicated. Each dot represents an individual mouse, and the cumulative average +SD from 2 experiments is plotted. Statistical significance for weight loss was calculated by one-way ANOVA with multiple comparisons and a Dunnett correction (D14 and D21 * p<0.05 CEP vs. CEP Δ*pigj*, D28 *** p<0.001 CEP vs. CEP Δ*pigj*). Statistical significance for brain cysts was calculated by a one-way ANOVA with multiple comparisons and a Tukey correction; * p<0.05, n.s. non-significant.

Figure 6: Early parasite burden and cytokine responses are similar between PIGJ mutant and wildtype strains.
A) Day 3 and C) day 5 after infection with the indicated parasite strains, peritoneal cavity exudate cells and E) splenocytes were harvested and RNA isolated. cDNA was synthesized from RNA and qPCR was performed to measure gene expression levels, which were normalized to uninfected mouse levels (naive=1). Cumulative results from 2-5 experiments are plotted. Each dot is the result of an individual mouse. B) Day 3 and D) 5 after infection, DNA was isolated from the peritoneal exudate cells and qPCR was performed using the *Toxoplasma* specific B1 primers to measure parasite burden relative to the DNA in each sample. Cumulative results of 2-3 experiments is plotted, with each dot representing an individual mouse. F) 5 days after infection, spleen, liver, and lungs were harvested, DNA was isolated, and qPCR was performed using the *Toxoplasma* specific B1 primers to measure parasite burden between infections. Each dot represents the result of an individual mouse and cumulative results from 2 experiments are plotted. Statistics were performed with an unpaired t-tests, for which no condition revealed a significant difference between CEP and CEP Δ*pigj* infections.

Figure 7: Surface expression of GPI-anchored SAGs, antibody recognition and functions against PIGJ mutants are intact.
A) Parasite surface expression of p35, SAG3, and SAG1. Fixed parasites were incubated with primary antibodies against the respective surface antigen, and secondary fluorescent anti-isotype antibodies were used to measure via flow cytometry. Representative histograms of 4 different experiments shown. CEP and CEP Δ*pigj* (top row); RH and RH Δ*pigj* (bottom row). B) Fixed parasites were incubated with serum from CEP chronically infected C57BL/6J mice, and antibodies bound to parasites were detected with fluorescent anti-isotype antibodies. Representative histograms of 2-3 experiments displaying parasite-specific antibody reactivity to the indicated CEP strains at 10^{-2} serum dilution (top row), or RH strains (bottom row). C) Parasites were incubated with 1% serum from CEP chronically infected C57BL/6J mice ("chronic serum") or naïve

mice for 20 minutes before allowing to invade or be phagocytosed for 40 minutes.
Opsonization was calculated as LAMP1+/ total LAMP1+ or GRA7+ for each parasite
observed. Each dot represents the ratio obtained after counting 100 parasites by
fluorescence microscopy for an individual serum, and samples were blinded. Plotted is
the average ratio +SD; no serum controls were also assessed. D) Parasites were
incubated with 10% chronic serum from C57BL/6J mice for 20 minutes before allowing
to invade MEFs for 2 hours. MEFs were measured for GFP+ (CEP strains) or by
intracellular staining with FITC-labeled parasite-specific antibodies (RH strains)
indicating parasite invasion and normalized to infections in the absence of serum. Each
dot represents the result from an individual serum. Statistics for opsonization was
calculated with one-way ANOVA with Tukey correction and neutralization were
calculated with unpaired t-tests; ns, non-significant
1207
Figure 8: Sera IgM recognition of GIPL requires the terminal glucose.
Parasite lysate was analyzed via western blot. Blots were probed with serum from CEP
chronically infected A) C57BL/6J (B6) or B) A/J mice and anti-mouse IgM-HRP was
used to detect IgM reactivity to parasite lysate antigens of the indicated genotypes. Red
boxes and arrows indicate the region where the low molecular weight 4-5 kDa GIPL
antigen migrates.
1214
Figure 9: Enhanced CD36 binding and slight macrophage tropism 3 hours after
infection.
Mice (C57BL/6J) were injected i.p. with 10^6 parasites of either CEP, CEP *Δpigj* or CEP
Δpigj + PIGJ and PECs were analyzed 3 hours later by flow cytometry. A) Gating
strategy for B-G. B) Total GFP+ signal in the peritoneal exudate; GFP frequency of total
events. C) GFP+ events were separated by size using FSC and SSC parameters to
determine extracellular or "non cell associated" and intracellular or "cell associated"
GFP+ parasites. D) Frequency PI+ cells of the indicated GFP+ category. E) Frequency
of cell associated GFP+ events that were GR-1^{hi} Cd11b+ neutrophils. F) Frequency of
cell associated GFP+ events that were MHCII^{hi} F4/80^{lo} bone marrow-derived
macrophages (BMDM), or G) MHCII^{lo} F4/80^{hi} yolk sack-derived macrophages (YSM).
Plotted is cumulative of 7 experiments, each dot represents a single mouse (mice; n =
12 wt, n = 12 *Δpigj*, n = 8 *Δpigj +PIGJ*). Statistics performed were one-way ANOVAs
with multiple comparisons and a Tukey correction; none were significant, values are
indicated. H) *In vitro* differentiated BMDMs were plated on coverslips overnight and
infected with 100 and 300 of GFP+ parasites per well, and parasites were allowed 20
minutes to invade cells before being fixed and stained for fluorescence microscopy. No
permeabilization was used, and SAG1 staining was performed. Intracellular parasites
were not stained with SAG1 but were GFP+, while extracellular parasites are
SAG1+GFP+. The fraction of intracellular parasites is plotted, each dot is a
representative of one experiment, n = 3 experiments. I) Parasites were incubated for 1
hr with recombinant CD36-Fc and CD36 binding was measured with anti-human-IgG-
Daylight-550 via flow cytometry. Representative histogram, and J) average +SD MFI of
CD36 binding from 4 experiments is plotted.
1239
Figure 10: CEP *Δpigj* virulence is galectin-3 dependent.

A) *Tlr2/4 -/-* mice were given primary infections of 10^4 parasites of CEP or CEP *Δpigj*
and monitored for survival for 30 days. Plotted are cumulative results from 2
experiments (mice; n = 5 CEP, n = 6 CEP *Δpigj*). B) *Lgals3 -/-* mice were given primary
infections of 10^4 parasites of CEP or CEP *Δpigj* and monitored for survival for 30 days.
Plotted are the cumulative results from 4 experiments (mice; n = 15 CEP, n = 23 CEP
Δpigj). C) Plotted is the average +/-SD fraction of initial weight for each cohort analyzed
in B. D) Brain cysts were quantified from surviving mice in B. Shown is 2 experiments,
(mice; n = 2 CEP, n = 5 CEP *Δpigj*). For survival analysis, significance was determined
by Log-rank (Mantel-Cox) test, * p <0.05. For weight loss, significance was determined
by one-way ANOVA test, * p <0.05. For brain cysts unpaired t-tests were performed, ns
= non-significant. E) Whole parasites were incubated with 10 ug/mL of recombinant
mouse galectin-3 and galectin-3 binding was detected with anti-Gal-3-PE by flow
cytometry. Representative histogram of galectin-3 binding; N-1 control is the staining in
the absence of galectin-3. F) MFIs of galectin-3 parasite binding from 9 experiments are
plotted. MFI values are connected by experiment and statistical analysis was performed
with a paired, one-tailed t-test. Normalized MFIs values (CEP=1) are also plotted, and
statistical analysis was performed with an unpaired, one-tailed t-test; * p < 0.05.
1258
SUPPLEMENTAL FIGURE LEGENDS
Figure S1. GPI sidechains differ between eukaryotic species.
Schematic of the diversity of GPI sidechains between species (*T. gondii*, *P. falciparum*,
T. brucei procyclic stage, *H. sapiens*). Note the conserved mannose backbone with
species variability of sidechain modifications. Not all GPI glycoforms for each species
are represented.
1265
1266 **Figure S2: Approach and PCR confirmation of *Tg_207750* disruption.**
1267 Schematic of the CRISPR Cas9 targeted disruption of the *Tg_207750* (*PIGJ*) locus and
1268 insertion of the selectable marker *HXGPRT* in A, or *DHFR-TS* in B. C-E) PCR
1269 confirmation of disruption of the targeted locus and insertion of the selectable marker
1270 within the cut site using the primers indicated in panels A or B.
1271
Figure S3: Approach and PCR confirmation of *Tg_266320* disruption.
Schematic of the CRISPR Cas9 targeted disruption of the T*GGT1_266320* (*PIGE*) locus
and insertion of the selectable marker *HXGPRT* in A, or *DHFR-TS* in B. C-D) PCR
confirmation of disruption of the target locus and insertion of the selectable marker
within the cut site using primers in A or B. In the case of CEP *Δ266320*, the *HXGPRT*
selectible marker inserted into the Cas9 cut site in exon one in reverse orientation, but
not the second site in exon 3. In the case of GT1 *Δ266320*, the *DHFR-TS* selectible
marker inseted in reverse orietnation in the Cas9 cut site in exon 1, while the cut site in
exon 3 was repaired with a concatemerized DHFR-TS as prposed based upon the
diagnostic PCR.
1282
Figure S4: Decreased T3 3F12 and T5 4E10 recognition of GT1 *Δpigj* mutants by
flow cytometry.
A) Fixed parasites of the indicated type I GT1 and RH parasite strains were stained with
GPIL glycoform specific antibodies T3 3F12 (mouse IgG3) and T5 4E10 (mouse IgM).

Secondary fluorescent anti-isotype antibodies were used to detect T3 3F12 and T5
4E10 binding. SAG3 positive parasites were analyzed to distinguish parasites from
debris. Data is representative of 3-8 experiments. B) Representative histograms from
the flow cytometry analysis comparing the parental with the mutant strains of the data
shown in A. In addition, a no staining control is plotted for comparison (grey histogram).
1292
Figure S5. Fragmentation analysis of GPI-anchor glycans.
Isolated primary GPI ions were selected in MS(1) as described in Figure 3 and
subjected to collision-based MS(2) fragmentation during the nLC-MS run. Residual
parent ions are labeled at the high m/z end, and decomposition products at lower m/z
values. Green arrows trace sequential fragmentation pathways. A) H4N2 glycan from
strain GT1. B) H3N2 glycan from GT1 *Δpige*. C) H3N1 from GT1-*Δpigj*.
1299
Figure S6: MALDI-TOF-MS analysis of GIPL preparations is consistent with PIGE
and PIGJ as being the GPI sidechain glycosyl transferases in *T. gondii*.
Glycans isolated from GPIL samples from tachyzoite stage parasites were N-acetylated
and permethylated, and mixed with DHB matrix and analyzed as in Figure 2. A)
Parental GT1 strain. The inset shows a sample prepared in CHCA rather than DHB
matrix. See text for the basis of assignment of ions that differ by a m/z defect of -36. B)
Δpige. C) *Δpigj*. D) Host cells (HFFs).
1307
Figure S7: nLC-MS analysis of GIPL preparations is consistent with PIGE and
PIGJ as being the GPI sidechain glycosyl transferases in *T. gondii*.
Isolated GIPL fractions described in Figure S6 were reanalyzed by nLC separation on a
C18 column and hyphenated analysis in an Orbitrap mass spectrometer in positive ion
mode. The left-hand column of panels shows base peak chromatograms (all ions m/z
500-2000) and extracted ion chromatograms (EIC) for each of the indicated targets (m/z
ranges given in panels A and D). The ratio of ion intensities over the EIC m/z ranges
shown for H3N1Ino, H3N2Ino, and H4N2Ino are shown relative to the most abundant
ion. The most abundant ions are labeled in the base peak chromatogram. The right-
hand column of panels shows representative mass spectra of the one or two most
abundant ions eluting with the target and used for quantitating relative levels. Levels of
hexosamers were not quantitated. Observed m/z values are in dark red, expected
values are in black, and the charge state is as indicated. A) Parental GT1 strain. B)
Δpige. C) *Δpigj*. D) Host cells (HFFs).
1322
Figure S8: C57BL/6J mice succumb to secondary infections with the type I GT1
GPI sidechain mutants similarly to the parental GT1 strain.
C57BL/6J (B6) mice given a primary infection with the avirulent type III strain CEP, 35
days later were given a secondary infection with $5x10^5$ parasites of the type I strain
GT1, GT1 *Δpigj* or GT1 *Δpige* strains and tracked for survival. Plotted is the result from
one experiment (mice; n = 3 wt GT1, n=5 GT1 *Δpigj*, n=1 GT1 *Δpige*).
1329
Figure S9: PIGJ mutants have no apparent fitness defects *in vitro*.
A) HFF cells in 24 well plates were infected with either CEP, CEP *Δpigj*, or CEP *Δpigj* +
PIGJ and allowed to grow for 5 days. On day 5 plaque sizes were measured and

plotted. Cumulative data from 4 experiments is shown, each dot represents a single
plaque area.
B) MEFs were infected for 16 hours before being fixed and stained for GRA7 to mark
the PV. Parasites per vacuole were quantified, counting 100 vacuoles per experiment,
and fractions are plotted for each experiment. Cumulative data from 4 experiments is
plotted, each dot is fraction obtained from an individual experiment. C) HFFs were
infected with 200 parasites per well and allowed various timepoints to attach before
being washed extensively and cultured for 5 days before quantifying plaque numbers,
which were normalized to the plaque counts from wells that were not washed (=1).
Cumulative average +/-SD from 4 experiments is plotted. For panels A-B, statistics
performed were one-way ANOVAs with multiple comparisons, and a Holm-Sidak's
correction; **** $p < 0.0001$ (GFP vs KO), *** $p < 0.001$ (GFP vs C2); ns, not significant.
For panel C, t-tests yielded non-significant values.
1346
Figure S10: Parasite burden is similar between wildtype and PIGJ mutant strains
in peritoneal exudate cells as assessed by flow cytometry.
3 and 5 days after infection with the indicated GFP expressing parasite strains,
peritoneal cavity exudate cells (PerC) were harvested and PI negative cells were
analyzed by flow cytometry. Average frequency (+SD) of infected GFP+ PerC among
total PI- cells is shown for each infection. Cumulative results from 2-3 experiments is
plotted, with each dot representing an individual mouse. Statistics were performed with
an unpaired t-test and values indicated, no condition revealed a significant difference
between CEP and CEP *Δpigj* infections.
1356
Figure S11: Complement C3b binding is unimpaired to PIGJ mutants.
Live parasites were incubated with serum from naïve mice and C3b binding was
measured with anti-C3b-APC. A) Representative histogram from 4 experiments each
with different naïve serums is shown. Staining controls include use of heat inactivated
serum and N-1, which is the staining in the absence of serum. C) Average +SD MFI of
C3b binding, each dot is the value of a different serum.
1363
Figure S12: Male C57BL/6J mice are protected against *Δpigj* primary and
secondary infections.
A-B) Male C57BL/6J (B6) mice were given primary infections i.p. with 10^4 parasites of
either CEP or CEP *Δpigj* and monitored for survival in A, and weight loss in B, for 30
days. Cumulative results from 2 experiments are plotted (mice; n = 6 wt, n = 8 *Δpigj*). C-
D) Male B6 mice were first given a primary infection of 10^4 CEP parasites, and after 35
days were given a challenge infection of either RH or RH *Δpigj* and monitored for
survival in C, and weight loss in D. Cumulative results from 2 experiments are plotted
(mice; n = 2 wt, n = 4 *Δpigj*). For survival analyses, significance was determined by Log-
rank (Mantel-Cox) test, and for weight loss, significance was determined by a one-way
ANOVA test with multiple comparisons with Dunnett correction, none of which were
found significant.
1376
Figure S13: Working model of *Δpigj* pathogenicity.

With the loss of sidechain in *Δpigj* mutants we have found an increase in CD36 binding,
an increase in BMDM tropism, and an increase in galectin-3 binding. We hypothesize
that there are other inflammatory signals possibly leading to tissue damage that
ultimately are responsible for the death of the hosts, and the cause for increased
pathogenicity in sidechain-null mutants.
1383
1384 **SUPPLEMENTAL TABLE LEGEND**
1385 **Table S1. Parasite strains, oligos and plasmids.**
1386 Parasite strains, oligos and plasmids used in this study are described and their origins
1387 indicated.
1388
1389

REFERENCES

1. Ferguson MAJ, Homans SW, Dwek RA, Rademacher TW. Glycosyl-phosphatidylinositol moiety that anchors Trypanosoma brucei variant surface glycoprotein to the membrane. Science (80-). 1988;239(4841):753–9.
2. Nosjean O, Briolay A, Roux B. Mammalian GPI proteins: sorting, membrane residence and functions. Biochim Biophys Acta - Rev Biomembr. 1997;1331(2):153–86.
3. Paing MM, Tolia NH. Multimeric Assembly of Host-Pathogen Adhesion Complexes Involved in Apicomplexan Invasion. PLoS Pathog. 2014;10(6).
4. Sidik SM, Huet D, Ganesan SM, Huynh MH, Wang T, Nasamu AS, et al. A Genome-wide CRISPR Screen in Toxoplasma Identifies Essential Apicomplexan Genes. Cell. 2016;166(6):1423-1435.e12.
5. Basagoudanavar SH, Feng X, Krishnegowda G, Muthusamy A, Gowda DC. Plasmodium falciparum GPI mannosyltransferase-III has novel signature sequence and is functional. Biochem Biophys Res Commun. 2007;364(4):748–54.
6. Wasmuth JD, Pszenny V, Haile S, Jansen EM, Gast AT, Sher A, et al. Integrated bioinformatic and targeted deletion analyses of the SRS gene superfamily identify SRS29C as a negative regulator of toxoplasma virulence. MBio. 2012;3(6).
7. Lekutis C, Ferguson DJP, Grigg ME, Camps M, Boothroyd JC. Surface antigens of Toxoplasma gondii: Variations on a theme. Int J Parasitol. 2001;31(12):1285–92.
8. Manger ID, Hehl AB, Boothroyd JC. The surface of Toxoplasma tachyzoites is dominated by a family of glycosylphosphatidylinositol-anchored antigens related to SAG1. Infect Immun. 1998;66(5):2237–44.
9. Rachinel N, Buzoni-Gatel D, Dutta C, Mennechet FJD, Luangsay S, Minns LA, et al. The Induction of Acute Ileitis by a Single Microbial Antigen of Toxoplasma gondii . J Immunol. 2004;173(4):2725–35.
10. Tachibana Y, Hashizaki E, Sasai M, Yamamoto M. Host genetics highlights IFN-γ-dependent Toxoplasma genes encoding secreted and non-secreted virulence factors in in vivo CRISPR screens. Cell Rep. 2023;42(6):112592.
11. Krishnamurthy S, Maru P, Wang Y, Bitew MA, Mukhopadhyay D, Yamaryo-botté Y. RESEARCH ARTICLE CRISPR Screens Identify Toxoplasma Genes That Determine. 2023;(March):1–20.
12. Hamie M, Tawil N, El Hajj R, Najm R, Moodad S, Hleihel R, et al. P18 (SRS35/TgSAG4) Plays a Role in the Invasion and Virulence of Toxoplasma gondii. Front Immunol. 2021;12(June):1–16.
13. Campos MAS, Almeida IC, Takeuchi O, Akira S, Valente EP, Procópio DO, et al. Activation of Toll-Like Receptor-2 by Glycosylphosphatidylinositol Anchors from a Protozoan Parasite. J Immunol. 2001;167(1):416–23.
14. Krishnegowda G, Hajjar AM, Zhu J, Douglass EJ, Uematsu S, Akira S, et al. Induction of proinflammatory responses in macrophages by the glycosylphosphatidylinositols of Plasmodium falciparum: Cell signaling receptors, glycosylphosphatidylinositol (GPI) structural requirement, and regulation of GPI activity. J Biol Chem. 2005;280(9):8606–16.
15. Schofield L, Hewittt MC, Evans K, Slomos MA, Seeberger PH. Synthetic GPI as a

candidate anti-toxic vaccine in a model of malaria. Nature. 2002;418(6899):785–
9.

16.   Debierre-Grockiego F, Campos MA, Azzouz N, Schmidt J, Bieker U, Resende
MG, et al. Activation of TLR2 and TLR4 by glycosylphosphatidylinositols derived
from Toxoplasma gondii. J Immunol. 2007;179(2):1129–37.

17.   Striepen B, Zinecker CF, Damm JBL, Melgers PAT, Gerwig GJ, Koolen M, et al.
Molecular structure of the "low molecular weight antigen" of Toxoplasma gondii: A
glucose α1-4 N-acetylgalactosamine makes free glycosyl-phosphatidylinositols
highly immunogenic. J Mol Biol. 1997;266(4):797–813.

18.   Tomavo S, Couvreur G, Leriche MA, Sadak A, Achbarou A, Fortier B, et al.
Immunolocalization and characterization of the low molecular weight antigen (4-5
kDa) of Toxoplasma gondii that elicits an early IgM response upon primary
infection. Parasitology. 1994;108(2):139–45.

19.   Naik RS, Branch OH, Woods AS, Vijaykumar M, Perkins DJ, Nahlen BL, et al.
Glycosylphosphatidylinositol Anchors of Plasmodium falciparum. J Exp Med.
2000;192(11):1563–76.

20.   Giraldo M, Cannizzaro H, Ferguson MAJ, Almeida IC, Gazzinelli RT. Fractionation
of membrane components from tachyzoite forms of Toxoplasma gondii:
Differential recognition by immunoglobulin M (IgM) and IgG present in sera from
patients with acute or chronic toxoplasmosis. J Clin Microbiol. 2000;38(4):1453–
60.

21.   Götze S, Azzouz N, Tsai YH, Groß U, Reinhardt A, Anish C, et al. Diagnosis of
toxoplasmosis using a synthetic glycosylphosphatidyl-inositol glycan. Angew
Chemie - Int Ed. 2014;53(50):13701–5.

22.   Naik RS, Krishnegowda G, Ockenhouse CF, Gowda DC. Naturally elicited
antibodies to glycosylphosphatidylinositols (GPIs) of Plasmodium falciparum
require intact GPI structures for binding and are directed primarily against the
conserved glycan moiety. Infect Immun. 2006;74(2):1412–5.

23.   Gilson PR, Nebl T, Vukcevic D, Moritz RL, Sargeant T, Speed TP, et al.
Identification and stoichiometry of glycosylphosphatidylinositol-anchored
membrane proteins of the human malaria parasite Plasmodium falciparum. Mol
Cell Proteomics. 2006;5(7):1286–99.

24.   França CT, Li Wai Suen CSN, Carmagnac A, Lin E, Kiniboro B, Siba P, et al. IgG
antibodies to synthetic GPI are biomarkers of immune-status to both Plasmodium
falciparum and Plasmodium vivax malaria in young children. Malar J.
2017;16(1):1–10.

25.   Sharma SD, Mullenax J, Araujo FG, Erlich HA, Remington JS. Western Blot
analysis of the antigens of Toxoplasma gondii recognized by human IgM and IgG
antibodies. J Immunol. 1983;131(2):977–83.

26.   Ropert C, Gazzinelli RT. Signaling of immune system cells by
glycosylphosphatidylinositol (GPI) anchor and related structures derived from
parasitic protozoa. Curr Opin Microbiol. 2000;3(4):395–403.

27.   Debierre-Grockiego F, Schwarz RT. Immunological reactions in response to
apicomplexan glycosylphosphatidylinositols. Glycobiology. 2010 Jul 1;20(7):801–
11.

28.   Ferguson MAJ. The structure, biosynthesis and functions of

glycosylphosphatidylinositol anchors, and the contributions of trypanosome research. J Cell Sci. 1999;112(17):2799–809.

29. Hirata T, Mishra SK, Nakamura S, Saito K, Motooka D, Takada Y, et al. Identification of a Golgi GPI-N-acetylgalactosamine transferase with tandem transmembrane regions in the catalytic domain. Nat Commun. 2018;9(1).

30. Hirata T, Kobayashi A, Furuse T, Yamada I, Tamura M, Tomita H, et al. Loss of the N-acetylgalactosamine side chain of the GPI-anchor impairs bone formation and brain functions and accelerates the prion disease pathology. J Biol Chem. 2022;298(3):101720.

31. Grimme SJ, Colussi PA, Taron CH, Orlean P. Deficiencies in the essential Smp3 mannosyl-transferase block glycosylphosphatidylinositol assembly and lead to defects in growth and cell wall biogenesis in Candida albicans. Microbiology. 2004;150(10):3115–28.

32. Cortes LK, Scarcelli JJ, Taron CH. Complementation of essential yeast GPI mannosyltransferase mutations suggests a novel specificity for certain Trypanosoma and Plasmodium pigB proteins. PLoS One. 2014;9(1):1–9.

33. Izquierdo L, Nakanishi M, Mehlert A, Machray G, Barton GJ, Ferguson MAJ. Identification of a glycosylphosphatidylinositol anchor-modifying β1-3 N-acetylglucosaminyl transferase in Trypanosoma brucei. Mol Microbiol. 2009;71(2):478–91.

34. Izquierdo L, Acosta-Serrano A, Mehlert A, Ferguson MA. Identification of a glycosylphosphatidylinositol anchor-modifying β1-3 galactosyltransferase in Trypanosoma brucei. Glycobiology. 2015;25(4):438–47.

35. Duncan SM, Nagar R, Damerow M, Yashunsky D V., Buzzi B, Nikolaev A V., et al. A Trypanosoma brucei β3 glycosyltransferase superfamily gene encodes a β1-6 GlcNAc-transferase mediating N-glycan and GPI anchor modification. J Biol Chem. 2021;297(4):14–6.

36. Gas-Pascual E, Ichikawa HT, Sheikh MO, Serji MI, Deng B, Mandalasi M, et al. CRISPR/Cas9 and glycomics tools for Toxoplasma glycobiology. J Biol Chem. 2019;294(4):1104–25.

37. Kongsomboonvech AK, Rodriguez F, Diep AL, Justice BM, Castallanos BE, Camejo A, et al. Naïve CD8 T cell IFNγ responses to a vacuolar antigen are regulated by an inflammasome-independent NLRP3 pathway and Toxoplasma gondii ROP5. Vol. 16, PLoS Pathogens. 2020. 1–35 p.

38. Mukhopadhyay D, Arranz-Solís D, Saeij JPJ. Influence of the Host and Parasite Strain on the Immune Response During Toxoplasma Infection. Front Cell Infect Microbiol. 2020;10(October).

39. Howe DK, Sibley LD. Toxoplasma gondii Comprises Three Clonal Lineages: Correlation of Parasite Genotype with Human Disease. J Infect Dis. 1995;172(6):1561–6.

40. Pfefferkorn ER and LCP. Toxoplasma gondii: Isolation and Preliminary Characterization of Temperature-Sensitive Mutants. Experimen. 1976;376(39):365–76.

41. Sibley LD, Boothroyd JC. Virulent strains of Toxoplasma gondii comprise a single clonal lineage. Nature. 1992;359(6390):82–5.

42. Gigley JP, Bhadra R, Khan IA. CD8 T cells and toxoplasma gondii: A new

paradigm. J Parasitol Res. 2011;2011.

43.   Jensen KDC, Grotenbreg GM, Frickel E-M, Julien L, Cordeiro C, Camejo A, et al.
Toxoplasma gondii Superinfection and Virulence during Secondary Infection
Correlate with the Exact ROP5/ROP18 Allelic Combination. MBio. 2015;6(2):1–
15.

44.   Splitt SD, Souza SP, Valentine KM, Castellanos BE, Curd AB, Hoyer KK, et al.
PD-L1, TIM-3, and CTLA-4 Blockade Fails To Promote Resistance to Secondary
Infection with Virulent Strains of Toxoplasma gondii . Infect Immun. 2018;86(9):1–
16.

45.   Yang N, Farrell A, Niedelman W, Melo M, Lu D, Julien L, et al. Genetic basis for
phenotypic differences between different Toxoplasma gondii type I strains. BMC
Genomics [Internet]. 2013;14(1):1. Available from: BMC Genomics

46.   Hallgren J, Tsirigos KD, Damgaard Pedersen M, Juan J, Armenteros A, Marcatili
P, et al. DeepTMHMM predicts alpha and beta transmembrane proteins using
deep neural networks. bioRxiv [Internet]. 2022;2022.04.08.487609. Available
from:
https://www.biorxiv.org/content/10.1101/2022.04.08.487609v1%0Ahttps://www.bi
orxiv.org/content/10.1101/2022.04.08.487609v1.abstract

47.   Keusch JJ, Manzella SM, Nyame KA, Cummings RD, Baenziger JU. Cloning of
GB3 synthase, the key enzyme in globo-series glycosphingolipid synthesis,
predicts a family of α1,4-glycosyltransferases conserved in plants, insects, and
mammals. J Biol Chem. 2000;275(33):25315–21.

48.   Striepen B, Dubremetz JF, Schwarz RT. Glucosylation of
glycosylphosphatidylinositol membrane anchors: Identification of uridine
diphosphate-glucose as the direct donor for side chain modification in toxoplasma
gondii using carbohydrate analogues. Biochemistry. 1999;38(5):1478–87.

49.   Komath S, Fujita M, Hart G. Essentials of Glycobiology. In: 4th ed. Cold Spring
Harbor (NY): Cold Spring Harbor Laboratory Press; 2022.

50.   Debierre-Grockiego F, Niehus S, Coddeville B, Elass E, Poirier F, Weingart R, et
al. Binding of Toxoplasma gondii glycosylphosphatidylinositols to galectin-3 is
required for their recognition by macrophages. J Biol Chem. 2010;285(43):32744–
50.

51.   Minot S, Melo MB, Li F, Lu D, Niedelman W, Levine SS, et al. Admixture and
recombination among Toxoplasma gondii lineages explain global genome
diversity. Proc Natl Acad Sci U S A. 2012;109(33):13458–63.

52.   McLeod R, Eisenhauer P, Mack D, Brown C, Filice G, Spitalny G. Immune
responses associated with early survival after peroral infection with Toxoplasma
gondii. J Immunol. 1989;142(9):3247–55.

53.   Souza SP, Splitt SD, Sanchez-Arcila JC, Alvarez JA, Wilson JN, Wizzard S, et al.
Genetic mapping reveals Nfkbid as a central regulator of humoral immunity to
Toxoplasma gondii. Vol. 17, PLoS Pathogens. 2021. 1–33 p.

54.   Huynh M-H, Carruthers VB. Tagging of Endogenous Genes in a Toxoplasma
gondii Strain Lacking Ku80 . Eukaryot Cell. 2009;8(4):530–9.

55.   Matta SK, Olias P, Huang Z, Wang Q, Park E, Yokoyama WM, et al. Toxoplasma
gondii effector TgIST blocks type I interferon signaling to promote infection. Proc
Natl Acad Sci U S A. 2019;116(35):17480–91.

56. Bate C, Nolan W, Williams A. Sialic acid on the glycosylphosphatidylinositol
anchor regulates PrP-mediated cell signaling and prion formation. J Biol Chem.
2016;291(1):160–70.
57. Banerjee P, Silva DV, Lipowsky R, Santer M. The importance of side branches of
glycosylphosphatidylinositol anchors: a molecular dynamics perspective.
Glycobiology. 2022;32(11):933–48.
58. Sikorski PM, Commodaro AG, Grigg ME. Toxoplasma gondii Recruits Factor H
and C4b-Binding Protein to Mediate Resistance to Serum Killing and Promote
Parasite Persistence in vivo. Front Immunol. 2020;10(January):1–16.
59. Zhao Y, Marple AH, Ferguson DJP, Bzik DJ, Yap GS. Avirulent strains of
Toxoplasma gondii infect macrophages by active invasion from the phagosome.
Proc Natl Acad Sci U S A. 2014;111(17):6437–42.
60. Zhao Y, Reyes J, Rovira-Diaz E, Fox BA, Bzik DJ, Yap GS. Cutting Edge: CD36
Mediates Phagocyte Tropism and Avirulence of Toxoplasma gondii . J Immunol.
2021;207(6):1507–12.
61. Silverstein RL, Febbraio M. CD36, a Scavenger Receptor Involved in Immunity,
Metabolism, Angiogogenesis, and Behavior. Sci Signal. 2010;2(72):1–16.
62. Klein SL, Flanagan KL. Sex differences in immune responses. Nat Rev Immunol.
2016;16(10):626–38.
63. Roberts CW, Cruickshank SM, Alexander J. Sex-determined resistance to
Toxoplasma gondii is associated with temporal differences in cytokine production.
Infect Immun. 1995;63(7):2549–55.
64. Bernardes ES, Silva NM, Ruas LP, Mineo JR, Loyola AM, Hsu DK, et al.
Toxoplasma gondii infection reveals a novel regulatory role for galectin-3 in the
interface of innate and adaptive immunity. Am J Pathol. 2006;168(6):1910–20.
65. Hsu JY, Crawley S, Chen M, Ayupova DA, Lindhout DA, Higbee J, et al. Non-
homeostatic body weight regulation through a brainstem-restricted receptor for
GDF15. Nature. 2017;550(7675):255–9.
66. BonDurant LD, Ameka M, Naber MC, Markan KR, Idiga SO, Acevedo MR, et al.
FGF21 Regulates Metabolism Through Adipose-Dependent and -Independent
Mechanisms. Cell Metab. 2017;25(4):935-944.e4.
67. Tezze C, Romanello V, Sandri M. FGF21 as modulator of metabolism in health
and disease. Front Physiol. 2019;10(MAR):1–9.
68. Baumgarth N. A Hard(y) Look at B-1 Cell Development and Function. J Immunol.
2017;199(10).
69. Villard O, Candolfi E, Ferguson DJP, Marcellin L, Kien T. Loss of oral infectivity of
tissue cysts of Toxoplasma gondii RH strain to outbred Swiss Webster mice. Int J
Parasitol. 1997;27(12):1555–9.
70. Khan A, Behnke MS, Dunay IR, White MW, David Sibley L. Phenotypic and gene
expression changes among clonal type i strains of toxoplasma gondit. Eukaryot
Cell. 2009;8(12):1828–36.
1615
1616
1617

A GPI of *Toxoplasma gondii*

TGGT1_321660
PIGB; DPM α2ManT
Fitness -3.27

TGGT1_270220
PIGV; DPM α6ManT
Fitness -3.05

TGGT1_310100
PIGM; DPM α4ManT
Fitness -4.66

TGGT1_241860
PIGA; PI α6GlcNAcT
Fitness -5.14

TGGT1_266320
PIGE; α4GlcT
Fitness -0.99

TGGT1_207750
PIGJ; β4GalNAcT
Fitness -0.47

P Phosphate
In Inositol
Glucosamine
Mannose
GalNAc
Glucose

B

TGGT1_207750 PIGJ Uniprot S8FC68 851 aa

NH₂ — intracellular | TM | stem? | CAZy GT17-like
N-sequon DxD L620R N-sequon
100 a.a.

TGGT1_266320 PIGE Uniprot S7VW57 1097 aa

NH₂ — intracellular | TM | stem? | CAZy GT32-like
DxD

C

Primary: anti-GalNAc (T3 3F12)
Secondary: anti-IgG

CEP | CEP Δ207750 | CEP Δ266320 | GT1 | GT1 Δ266320 | GT1 Δ207750 | RH | RH Δ207750

4-5 kDa

Primary: anti-GalNAc + Glc (T5 4E10)
Secondary: anti-IgM

CEP | CEP Δ207750 | CEP Δ266320 | GT1 | GT1 Δ207750 | GT1 Δ266320 | RH | RH Δ207750

4-5 kDa

MALDI-TOF-MS analysis of GPI anchor preparations

nLC-MS analysis of GPI anchor preparations

A GIPL Western Blot

T3 3F12   4-5 kDa

T5 4E10   4-5 kDa

CEP   CEP Δpigj   CEP Δpigj + PIGJ

B CEP *GFP:cLuc Δpigj + PIGJ*$_{3xHA}$

DAPI   GFP   HA

C B6 Primary Infection

Weight Loss

- CEP
- CEP Δpigj
- CEP Δpigj +PIGJ

D Brain Cysts